

Contribution of liquid, NAT and ice particles to chlorine activation and ozone depletion during Antarctic winter and spring

O. Kirner¹, R. Müller², R. Ruhnke³, and H. Fischer³

¹Karlsruhe Institute of Technology, Steinbuch Centre for Computing (SCC), Karlsruhe, Germany

²Institute of Energy and Climate Research – Stratosphere (IEK-7), Research Centre Jülich GmbH, Jülich, Germany

³Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research (IMK), Karlsruhe, Germany

Abstract. Heterogeneous reactions in the Antarctic stratosphere are the cause of chlorine activation and ozone depletion, but the relative roles of different types of PSCs in chlorine activation is an open question. We use multi-year simulations of the chemistry-climate model EMAC to investigate the impact that the various types of PSCs have on Antarctic chlorine activation and ozone loss.

One standard and three sensitivity EMAC simulations have been performed. In all simulations a Newtonian relaxation technique using the ERA-Interim reanalysis was applied to simulate realistic synoptic conditions. In the three sensitivity simulations, we only changed the heterogeneous chemistry on PSC particles by switching on and off the chemistry on liquid, NAT and ice particles. The results of these simulations show that the significance of heterogeneous reactions on NAT and ice particles for chlorine activation and ozone depletion in Antarctic winter and spring is small in comparison to the significance of heterogeneous reactions on liquid particles. In comparison to liquid particles, is subordinate regarding chlorine activation and ozone depletion in Antarctic winter and spring. The heterogeneous chemistry on liquid particles is sufficient to activate at least 90% of the chlorine reservoir species. Liquid particles alone are sufficient to activate almost all of the available chlorine, with the exception of the upper PSC regions between 10 and 30 hPa, where temporarily ice particles show a relevant contribution. Shortly after the first PSC occurrence, NAT particles contribute a small fraction to chlorine activation. With the exception of the upper PSC regions between 10 and 30 hPa where temporarily the ice particles have a relevant contribution to the chlorine activation and during the initial PSC occurrence with short NAT contributions the liquid particles alone are sufficient to activate almost all of the available chlorine.

In the model simulations heterogeneous chemistry on

liquid particles is responsible for more than 90% of the ozone depletion in Antarctic spring in the model simulations. In high southern latitudes heterogeneous chemistry on ice particles causes only up to 5 DU of additional ozone depletion in the column and heterogeneous chemistry on NAT particles less than 0.5 DU. Only up to 5 DU of column ozone in high southern latitudes is depleted by chlorine activation due to additional heterogeneous chemistry on ice particles and less than 0.5 DU due to additional heterogeneous chemistry on NAT particles.

The simulated HNO₃, ClO and O₃ results agree closely with observations from the Microwave Limb Sounder onboard NASA's Aura satellite.

1 Introduction

Polar Stratospheric Clouds (PSCs) consist of supercooled ternary solution (STS, type 1b PSC), nitric acid trihydrate (NAT, type 1a PSC) and ice particles (type 2 PSC). An accurate representation of PSCs and cold binary solutions of the sulphuric acid background/water (binary) aerosol is essential for the correct simulation of chlorine activation and polar ozone depletion in chemistry-climate-models (CCMs). The liquid (cold binary aerosols and STS particles) and solid particles (NAT and ice particles) allow heterogeneous reactions to proceed, which cause the activation of chlorine reservoirs (Solomon et al., 1986) and the production of chlorine radicals leading to ozone destruction. In addition, NAT particles cause and for the stratospheric denitrification of the stratosphere (through NAT particles), resulting in a delay in the deactivation of this active chlorine in polar spring (e.g., Toon et al., 1986; Lowe and MacKenzie, 2008).

Here we revisit the question of the importance of different types of PSC and stratospheric aerosol for heterogeneous chlorine activation and ozone depletion in Antarctic winter and spring using To consider the relevance of PSCs

~~for polar chlorine activation, a new algorithm based on an efficient growth of NAT particles, developed by van den Broek et al. (2004), has been implemented into the submodel PSC (Kirner et al., 2011) of the chemistry-climate model ECHAM5/MESSy for Atmospheric Chemistry (EMAC, Jöckel et al., 2006). to improve the representation of the PSCs in this model. The submodel PSC and the new NAT parameterisation are described in Sect. 3 and in Kirner et al. (2011).~~

Over the last years a discussion developed about the **different** contribution of cold binary aerosols, STS, NAT and ice particles to chlorine activation. Drdla and Müller (2012) have raised the question of the necessity of NAT and ice particles for reaching the full polar stratospheric chlorine activation. ~~They also argue that assuming only liquid particles or even liquid binary particles only leads to a minor reduction of the simulated chlorine activation and ozone loss. They further argue that even the heterogeneous reactivity on cold binary aerosol is sufficient to explain the major fraction of polar stratospheric chlorine activation. In contrast, Solomon et al. (2014) have argued that in sunlit air in the polar stratosphere, deactivation, i.e. the reformation of HCl and ClONO₂, increases the rate at which heterogeneous reactions must proceed to keep pace with deactivation.~~

Wegner et al. (2012) examined the importance of the cold binary aerosols for chlorine activation in the Arctic winters 2004/2005 and 2009/2010 with the help of in-situ observations. They concluded that the surface area provided by PSCs did not significantly impact chlorine activation rates in these both winters. Wohltmann et al. (2013) showed, based on simulations of the Arctic winter 2009/2010 with the Lagrangian model ATLAS, that liquid aerosols alone allow to explain the observed mixing ratios of active chlorine and ozone from MLS, ACE-FTS and in-situ measurements.

However, in these Arctic studies the contribution of ice particles to chlorine activation was not investigated and there is still the question if liquid particles are also sufficient to activate the complete inorganic chlorine during Antarctic winters. Therefore, we performed a standard EMAC simulation from 2000 to 2012 and three additional sensitivity simulations from 2005 to 2012 with different assumptions for heterogeneous chemistry on stratospheric aerosol and PSC particles. By comparing the results of the EMAC standard and sensitivity simulations it is possible to determine the contribution of liquid particles (in EMAC the cold binary aerosols and STS particles are not distinguished, see Kirner et al., 2011), NAT particles and ice particles to chlorine activation in polar winter and **to determine the** corresponding ozone depletion in polar spring.

For comparison of our model results with observations we choose the Earth Observing System (EOS) Microwave Limb Sounder (MLS) (Waters et al., 2006) onboard NASA's Aura satellite.

2 The EMAC model

The ECHAM/MESSy Atmospheric Chemistry (EMAC) model is a numerical chemistry and climate simulation system that includes submodels describing tropospheric and middle atmosphere processes and their interaction with oceans, land and human influences (Jöckel et al., 2006). It uses the Modular Earth Submodel System (MESSy) to link multi-institutional ~~numerical-computer~~ codes (Jöckel et al., 2005). The core atmospheric model is the 5th generation European Centre Hamburg general circulation model (ECHAM5, Roeckner et al., 2006). For the present study we applied EMAC (ECHAM5 version 5.3.01, MESSy version 1.10) in the T42L39MA-resolution, i.e. with a spherical truncation of T42 (corresponding to a quadratic Gaussian grid of ~~approx. approximately~~ 2.8° by 2.8° in latitude and longitude) with 39 vertical hybrid pressure levels up to 0.01 hPa (~~approx. approximately~~ 80 km) and with a time step of 10 min. ~~The approximate vertical resolution in the stratosphere is between 1.5 and 2.5 km decreasing with altitude.~~

Besides the submodel MECCA1 (Sander et al., 2005) for the calculation of gas-phase chemistry and the submodel PSC for the simulation of polar stratospheric clouds (Kirner et al., 2011) ~~among others~~ the following submodels have been used in our study: OFFLEM for offline emissions of trace gases and aerosols, TNUDGE for tracer nudging (both Kerkweg et al., 2006), ~~DRYDEP for dry deposition of trace gases and aerosols (Kerkweg et al., 2006b), SEDI for the sedimentation of aerosol particles (Kerkweg et al., 2006a), JVAL for the calculation of photolysis rates (Landgraf and Crutzen, 1998), SCAV for the scavenging and liquid phase chemistry in cloud and precipitation (Tost et al., 2006a), CONVECT for the parameterisation of convection (Tost et al., 2006), LNOX for the source of NO_x produced by lightning (Tost et al., 2007b), PTRAC for additional prognostic tracers (Jöckel et al., 2008), CVTRANS for convective tracer transport, TROPOP for diagnosing the tropopause and boundary layer height, H2O for stratospheric water vapour, RAD4ALL for the radiation calculation, and HETCHEM for calculating reaction rate coefficients of heterogeneous reactions on aerosols (see Jöckel et al., 2006, and references therein), and CLOUD for calculating the cloud cover as well as cloud microphysics including precipitation (Tost et al., 2007a).~~

Reaction rate coefficients for gas phase reactions and absorption cross sections for photolysis are taken **mainly** from Atkinson et al. (2007) and Sander et al. (2011).

A Newtonian relaxation technique for the prognostic variables temperature, vorticity, divergence and the surface pressure is implemented in EMAC to reproduce realistic synoptic conditions. We applied this nudging technique in our simulations above the boundary layer and below 1 hPa with nudging strengths as recommended in van Aalst (2005) using the ERA-Interim reanalysis (Dee et al., 2011) from the European Centre for Medium-range Weather Forecasts (ECMWF).

3 The submodel PSC

The submodel PSC is originally based on the “Mainz Photochemical Box Model” (Crutzen et al., 1992; Müller et al., 1994; Grooß, 1996; Meilinger, 2000) and was improved and extended by Buchholz (2005) and Kirner (2008). A comprehensive description of the submodel PSC is given by Kirner et al. (2011). The submodel PSC simulates the number densities, mean radii and surface areas of liquid sulphuric acid aerosols, supercooled ternary solutions (STS), nitric acid trihydrate (NAT) particles and ice particles. Different sedimentation schemes for these particles are included (the trapezoid scheme is used in our simulations (Buchholz, 2005)). Together with the submodels HETCHEM and MECCA1 the reaction rate coefficients and reaction rates of all heterogeneous reactions are calculated, which are important for the ozone depletion in polar spring. Heterogeneous reaction rates and their temperature dependencies on NAT were calculated according to the parameterisation of Carslaw et al. (1995) based on the measurements of Hanson and Ravishankara (1993). The heterogeneous reaction rate coefficients on liquid particles are taken from Hanson and Ravishankara (1994) and Hanson et al. (1994), the uptake coefficients and reaction probabilities for ice particles from Sander et al. (2011).

Heterogeneous reaction rates and their temperature dependencies on NAT were calculated according to the parameterisation of Carslaw and Peter (1997) based on the measurements of Hanson and Ravishankara (1993). Carslaw and Peter (1997) also developed a parameterisation for heterogeneous reactions on NAT based on the measurements of Abbatt and Molina (1992). Using the latter parameterisation results in substantially lower (by a factor of more than 50) heterogeneous reactivity on NAT and in a significantly reduced chemical ozone loss (Carslaw et al., 1997); therefore this latter parameterisation was not considered here.

The heterogeneous reaction rate coefficients on liquid particles are taken from Hanson and Ravishankara (1994) and Hanson et al. (1994). An alternative parameterisation for reactions on liquid particles was reported by Shi et al. (2001), which results however in a heterogeneous reactivity very close to the one employed here (e.g., Wegner et al., 2012).

The heterogeneous reactivity of ice particles is much larger than that of other types of PSCs because of the much larger surface area of ice PSCs; the uptake coefficients and reaction probabilities for ice particles are taken from Sander et al. (2011).

The formation of STS ($\text{HNO}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$) in the model is based on Carslaw et al. (1995) and takes place through uptake of nitric acid (HNO_3) by liquid sulphuric acid (binary) aerosols ($\text{H}_2\text{SO}_4\text{-H}_2\text{O}$) (described in detail in Kirner et al. (2011)).

The formation of ice particles in the model is based on the thermodynamic approach of Marti and Mauersberger (1993) with:

$$\text{H}_2\text{O}(\text{ice}) = \frac{(e_{\text{H}_2\text{O}(\text{total})} - E_{\text{H}_2\text{O}}^{\text{ice}})}{p} \quad (1)$$

The ratio of ice phase to dry air ($\text{H}_2\text{O}(\text{ice})$ in mol mol^{-1}) depends on the difference of the total partial pressure of H_2O ($e_{\text{H}_2\text{O}(\text{total})}$ in Pa) and the saturation vapour pressure of H_2O over ice particles ($E_{\text{H}_2\text{O}}^{\text{ice}}$ in Pa). For the ice nucleation 50 % H_2O supersaturation is assumed to be necessary (Tabazadeh et al., 1997; Peter and Grooß, 2012).

With the help of $\text{H}_2\text{O}(\text{ice})$ the total mass of ice particles (m_{ICE}) as well as their total volume (V_{ICE} in m^3) is calculated. The total number density (N_{ICE}) and the mean radius (r_{ICE}) of ice particles is calculated with the help of a minimum ice radius (r_{min}) and a maximum number density of ice particles (N_{max}). In our simulations we assumed for $r_{\text{min}} = 1 \times 10^{-5}$ cm and for $N_{\text{max}} = 0.042 \text{ cm}^{-3}$ (Höpfner et al., 2006).

With N_{ICE} and r_{ICE} it is possible to calculate the surface area density of ice (A_{ICE}) important for the calculation of the heterogeneous reaction rates on ice particles.

Besides the parameterisations for the formation of liquid and ice particles, there are two parameterisations for the formation of NAT particles included in the PSC submodel. It is possible to choose either a “thermodynamical NAT parameterisation” assuming instantaneous thermodynamical equilibrium (Hanson and Mauersberger, 1988) or the new “kinetic growth NAT parameterisation” for NAT particles based on the efficient growth and sedimentation algorithm of Carslaw et al. (2002) and van den Broek et al. (2004) implemented into the submodel by Kirner et al. (2011). In both cases, NAT formation takes place below the NAT existence temperature (T_{NAT} , depending on the pressure and on the partial pressures of HNO_3 and H_2O) with the assumption of a necessary super cooling of 3 K (Schlager and Arnold, 1990; Dye et al., 1992).

We used the kinetic growth NAT parameterisation for this study. Hence, also homogeneous NAT nucleation is possible in our simulations.

The time-related growth of the NAT particles in the kinetic growth NAT parameterisation is based on the calculation of the surface growth factor (G in $\text{m}^2 \text{s}^{-1}$):

$$G = \frac{D_{\text{HNO}_3}^* M_{\text{NAT}}}{\rho_{\text{NAT}} R_{\text{gas}} T} (e_{\text{HNO}_3} - E_{\text{HNO}_3}^{\text{NAT}}) \quad (2)$$

G is depending on the temperature (T in K), the difference between the partial HNO_3 vapour pressure (e_{HNO_3} in Pa) and the saturation vapour pressure of HNO_3 over NAT ($E_{\text{HNO}_3}^{\text{NAT}}$ in Pa), as well as on the diffusion coefficient of HNO_3 in air ($D_{\text{HNO}_3}^*$ in $\text{m}^2 \text{s}^{-1}$). For negative G , the particle size decreases through evaporation.

With the help of this surface growth factor it is possible to simulate the growth of NAT particles. For integration of this growth concept in EMAC, which is an Eulerian model,

it is necessary to split the NAT particles into different size bins. In the kinetic growth NAT parameterisation of the sub-model PSC consequently a separation into eight size bins is implemented (see Table 1). The minimum and maximum NAT radii in the different size bins, as well the maxima of the NAT number densities are assumed based on Fahey et al. (2001).

The total number density of NAT (N_{NAT}) is calculated as the sum over all eight size bins. The mean NAT radius (r_{NAT}) is calculated from the NAT radii of each size bin, weighted with the corresponding NAT number density. With N_{NAT} and r_{NAT} it is possible to calculate the surface area density of NAT (A_{NAT}) important for the calculation of the heterogeneous reaction rates on NAT particles.

A comprehensive description of the submodel PSC is given by Kirner et al. (2011).

4 Performed simulations

Four different EMAC simulations were performed using the submodel PSC: a standard simulation (*Standard*) from 2000 to 2012 with heterogeneous chemistry on liquid, NAT and ice particles and three additional sensitivity simulations from 2005 to 2012. In the three sensitivity simulations, we only changed the heterogeneous chemistry on PSC particles by switching on and off the chemistry on liquid, NAT and ice particles. We performed one sensitivity simulation without heterogeneous chemistry (*NoHet*), one sensitivity simulation with heterogeneous chemistry only on liquid particles (*Liquid*) and one sensitivity simulation with heterogeneous chemistry only on liquid and NAT particles, but excluding heterogeneous chemistry on ice particles (*LiquidNAT*) (see also Table 2).

The assumptions on the formation and distribution of the PSC particles are the same for the four simulations. Also the denitrification caused by the sedimentation of NAT particles is simulated in all simulations consistently. Only the assumptions regarding the heterogeneous chemistry on liquid, NAT and ice particles are different.

Through the assumptions made here in the sensitivity simulations we only test the impact of the heterogeneous reactivity of the different PSC particle types. An alternative scenario would be to make assumptions on the formation of PSC (especially NAT) and test the combined effect of heterogeneous chemistry and particle setting.

To simulate realistic synoptic conditions we apply in all four EMAC simulations a Newtonian relaxation technique of the prognostic variables temperature, vorticity, divergence and the surface pressure is used. We applied this nudging technique above the boundary layer and below 1 hPa with nudging strengths as recommended in van Aalst (2005) using the ERA-Interim reanalysis (Dee et al., 2011) from the European Centre for Medium-range Weather Forecasts (ECMWF).

The chemistry of the standard simulation is initialized from a previously completed climatological simulation, the meteorological conditions from ERA-Interim reanalysis. For initialization of the three sensitivity simulations the rerun files of the standard simulation are used. Boundary conditions for greenhouse gases are in all four EMAC simulations from the IPCC-A1B scenario (IPCC, 2007). Boundary conditions for ozone depleting substances (CFCs and halons) are from the WMO-Ab scenario (WMO, 2007). We increased the CH_3Br boundary condition by a constant value of 5 pptv (WMO, 2007), because very short lived bromine species are not included in our simulations.

The simulation includes a comprehensive chemistry setup with 104 gas phase species, 234 gas phase reactions, 67 photolysis reactions, and 11 heterogeneous reactions on liquid aerosols, NAT- and ice particles (Jöckel et al., 2006).

5 The MLS observations

The Earth Observing System (EOS) Microwave Limb Sounder (MLS) (Waters et al., 2006) onboard NASA's Aura satellite launched on 15 July 2004. It observes in a sun-synchronous orbit (98° inclination) temperature and trace gas profiles between 82°S and 82°N by scanning the Earth's limb and measuring the microwave emission. The MLS vertical resolution depends on the considered species and usually ranges between 3-4 km in troposphere/stratosphere and 8 km in the mesosphere (Livesey et al., 2011).

In this study we are using MLS Level-2 data Version 3.3 of HNO_3 , ClO , and O_3 Level 2 data. In the stratosphere the precision of the HNO_3 profile is about 0.7 ppbv with a vertical resolution between 3 to 5 km (Sect. 3.11, Livesey et al., 2011). The precision of ClO is between 0.1 and 0.3 ppbv with a vertical resolution between 3 to 5 km. (Santee et al., 2008). In the uppermost troposphere and stratosphere the precision of MLS O_3 data is between 0.04 and 0.5 ppm with a vertical resolution of about 3 km (Froidevaux et al., 2008).

For comparison with the EMAC results we analysed the MLS observations of the year 2007 (Figs. 1, 4, and 6).

6 Mean development of liquid, NAT and ice particles in EMAC and simulated denitrification

The formation and distribution of PSCs are simulated in EMAC very reasonably. For example, in Fig. 1 (first to third row) the simulated denitrification in 46 hPa due to NAT particles from the standard simulation is presented. After the temperature has dropped below 192 K, and thus the necessary temperature threshold is reached for NAT formation, the first NAT particles are formed on 15 May 2004/12 May 2007 over Antarctica. Due to the continued cooling from 19 to 31 May 2004/12 May 2007 to 31 May 2007 the number density of NAT increases up to 240 m^{-3} $2.4 \times 10^{-4}\text{ cm}^{-3}$ and, simultaneously, the NAT- PSCs are expanding into a larger

area. ~~In HNO₃ the~~ The first hint of denitrification due to sedimentation of NAT is visible on ~~19 May 2004~~ ~~19 May 2007~~. Ongoing NAT formation and sedimentation of NAT particles leads ~~in EMAC~~ to reduced HNO₃ in the area in which NAT particles are present.

The MLS measurements of HNO₃ in 46 hPa (Fig. 1, bottom row) agree closely with the EMAC simulations. The MLS HNO₃ values are in general higher than simulated in EMAC, but the first hint of denitrification occurs about at the same time as in the standard simulation and the time development of HNO₃ is similar.

Besides ~~the~~In addition to NAT particles, also ice particles and liquid particles ~~in form of cold binary solutions (H₂SO₄-H₂O) and supercooled ternary solutions (HNO₃-H₂SO₄-H₂O)~~ are calculated with the submodel PSC. In Fig. 2 the time development of liquid particles, NAT and ice particles is shown ~~in form of~~ as an average over the latitudes from 80° S to 90° S and over the five years from 2005 to 2009. ~~The illustrated results~~The results shown in Fig. 2 are valid for the standard and for all three sensitivity simulations, i.e. the formation and development of stratospheric aerosol and PSCs are ~~the same~~ identical for all simulations. ~~In average~~NAT is ~~visible from mid-May until October~~ in the EMAC simulations ~~visible from mid-May until October in the area of lower temperatures~~. The relatively slow increase of the number density ~~from 0 to the average maximum of over~~ to values of more than ~~200 m⁻³ × 10⁻⁴ cm⁻³~~ by end of May is caused by ~~different first NAT appearances~~differences in the timing of the first NAT appearances in the individual years of 2005 to 2009. The decrease of NAT number densities between June to September is caused by the sedimentation of larger NAT particles and the related denitrification. Ice particles exist mainly from June to September with mean maxima of number densities of over ~~40 000 m⁻³ = 0.04 cm⁻³~~. In comparison to liquid and NAT particles they have the largest surface area densities with values up to 38 μm² cm⁻³. The liquid particles have the second largest surface area densities with values up to 6.8 μm² cm⁻³, the NAT particles the smallest with maxima of 0.11 μm² cm⁻³.

7 Contribution of NAT, ice and liquid particles to stratospheric chlorine activation and ozone depletion

The contribution of heterogeneous reactions on liquid, NAT and ice particles to chlorine activation and the corresponding ozone depletion in polar spring is determined by comparing the standard and the sensitivity simulations. ~~By comparing the standard and the sensitivity simulations it is possible to determine the contribution of the heterogeneous reactivity of liquid, NAT and ice particles to polar chlorine activation and corresponding ozone depletion in polar spring. In this way it is possible to assess~~ The impact of the liquid particles is assessed by ~~subtraction~~subtracting the results of the simula-

tion *NoHet* from the results of the simulation *Liquid*. The absolute contribution of additional NAT chemistry is calculated by ~~subtraction~~subtracting the results of the simulation *Liquid* from the simulation *LiquidNAT*, and the contribution of additional heterogeneous chemistry on ice particles by ~~subtraction~~subtracting the results of the simulation *LiquidNAT* from the simulation *Standard*.

In Figs. 3 to 7 we examine the chlorine activation and ozone depletion of the different EMAC simulations and compare the results of the standard simulation with MLS observations. For a better statistics we averaged the simulated ClO_x and O₃ results from EMAC over the southern high latitudes from 80° S to 90° S and over the five year time period from 2005 to 2009 (Figs. 3 and 5) and the ozone columns as zonal mean over the same five years (Fig. 7). For the comparison of the EMAC results with MLS observations the year 2007 is chosen (Figs. 4 and 6).

In Fig. 3 (top-left), the evolution ~~the averaged development~~ of ClO_x in the standard simulation ~~and the contributions of liquid, NAT and ice particles are~~ is illustrated shown. ~~The contributions of liquid, NAT and ice particles are shown in Fig. 3 (top right and two bottom panels).~~ It is obvious that the simulated ClO_x development with maximum values of about 2.9 ppbv is mainly caused ~~due to~~by the heterogeneous chemistry on liquid particles with contributions up to 2.6 ppbv. There is almost no extra chlorine activation due to the additional heterogeneous chemistry on NAT particles with a contribution below 0.1 ppbv and only a slight additional chlorine activation due to heterogeneous chemistry on ice particles. ~~The relevant contribution of the ice particles is mainly~~Ice particles only show a relevant contribution in the upper PSC region in which only few liquid particles are simulated (see Fig. 2). A maximum of about 0.33 ppbv ~~extra~~additional ClO_x is present in the region between 10 and 30 hPa from end of June until mid-September due to additional heterogeneous chemistry on ice.

Liquid particles contribute more than 90% to chlorine activation for most of the time in Antarctic winter and spring in the performed EMAC simulations. ~~The contribution of liquid particles to chlorine activation is for most of the time over 90%.~~ The contribution is almost 100% from ~~Between~~ June and to August in the region between 30 hPa and 100 hPa ~~the contribution is almost 100%.~~ The heterogeneous chemistry of additional ice particles is ~~Only~~ only relevant in the upper PSC region, as mentioned above, ~~the heterogeneous chemistry of additional ice particles is relevant~~ with maximum contributions in the range of 10% to 15% of the total chlorine activation. The NAT particles show only a relevant contribution to chlorine activation ~~Only~~ during the initial PSC occurrence ~~have the NAT particles a relevant contribution to chlorine activation in Mid-May in the region between 20 hPa and 50 hPa, with contributions of up to 10%.~~ in Mid-May in the region between 20 hPa and 50 hPa.

The MLS measurements of ClO in 79.5° S in Antarctic winter and spring (Fig. 4) compare well with the ClO de-

velopment of the EMAC standard simulation. In EMAC and MLS increased ClO mixing ratios are visible between mid-August and end of September in the middle and upper stratosphere. In the middle stratosphere maximum values over 1 ppbv are reached in early September both in the measurements and in the simulation. In the upper stratosphere the simulated ClO development is also close to the MLS observations.

The evolution of ozone development resulting from the chlorine activation in the different EMAC simulations is shown in Fig. 5. The ozone depletion is highest from mid-September to late October forming, resulting in the formation of the ozone hole (e.g., Groöb et al., 2011). The largest impact contribution to this ozone depletion comes from the chlorine activated on liquid particles with values an ozone depletion of more than 2500 ppbv. In contrast in the upper PSC region only values up to 170 ppbv during the maximum ozone depletion can be attributed to the simulation of additional heterogeneous chemistry on ice particles. In contrast, the contribution that can be attributed to additional chemistry on ice particles reaches values of only 170 ppbv in the upper PSC region between 10 to 30 hPa. A contribution of NAT particles is visible in the region between 20 to 40 hPa in September and October, but the values are close to zero.

The comparison of the EMAC ozone evolution with MLS in 79.5° S in Antarctic winter and spring (Fig. 6) is very reasonable. The observed ozone hole is represented in EMAC, even if the O₃ decrease is slightly smaller than in MLS. In general, the O₃ mixing ratios are slightly higher in EMAC than in the observations, but the time evolution of O₃ agree closely with MLS.

The total effect on ozone chemistry of applying heterogeneous chemistry on different PSC particles is illustrated in form of the time development of ozone columns in the Southern Hemisphere in Fig. 7. Fig. 7 shows the total effect that heterogeneous chemistry on different PSC particles has on the time development of the ozone column. Liquid particles are mainly responsible for the ozone depletion noticeable as a decrease in ozone columns by more than 130 DU in early October in high southern latitudes. The ozone column shows a decrease of more than 130 DU by early October in high southern latitudes. Chlorine activation on liquid particles is mainly responsible for this decrease. The contribution of liquid particles to the simulated ozone depletion during September to October in the southern high latitudes the contribution of liquid particles to the simulated ozone depletion is at least 95 % during September to October at southern high latitudes.

The contribution of additional heterogeneous chemistry on ice particles to Antarctic ozone depletion only reaches up to 5 DU and is always smaller less than 5 %. The contribution of additional heterogeneous chemistry on NAT particles is less than 0.5 DU and thus almost negligible. On the other hand, the NAT particles are essential for the denitrification in

polar spring and thus have an additional indirect influence on the formation of the ozone hole.

8 Conclusions

The significance of heterogeneous reactions on NAT and ice particles with respect to chlorine activation and ozone depletion in Antarctic winter and spring is small compared to reactions on liquid particles in standard and sensitivity our simulations performed with the EMAC model. The simulated heterogeneous chemistry on liquid particles is sufficient to activate at least about 90 % of the chlorine.

With the exception of the upper PSC regions between 10 and 30 hPa, where temporarily the ice particles have relevant contribution to chlorine activation and during the initial PSC occurrence with NAT contributions, the liquid particles are sufficient to activate almost all of the available chlorine reservoirs alone. Liquid particles alone are sufficient to activate almost all of the available chlorine, with the exception of the upper PSC regions between 10 and 30 hPa, where temporarily ice particles show a relevant contribution when few other types of PSCs are present. Shortly after the first PSC occurrence, NAT particles contribute a very small fraction to chlorine activation.

In the model simulations, heterogeneous chemistry on liquid particles is mainly responsible for more than 90 % of the ozone depletion in polar spring in our EMAC simulations. In high southern latitudes heterogeneous chemistry on ice particles causes only up to 5 DU of additional ozone depletion in the column and heterogeneous chemistry on NAT particles less than 0.5 DU. Only up to 5 DU of column ozone in high southern latitudes is depleted by chlorine activation due to additional heterogeneous chemistry on ice particles and less than 0.5 DU due to additional heterogeneous chemistry on NAT particles.

It might be surprising that the enormous increase of heterogeneous reactivity provided by the frequent occurrence of ice PSCs has rather little impact on chlorine activation in the simulations presented here. However, a stronger heterogeneous reactivity does not necessarily translate into stronger chlorine activation. Frequently, chlorine activation through heterogeneous chemistry is rate limited by the gas phase formation of reaction partners for heterogeneous reactions (e.g., Müller et al., 1994). This is in particular true for the time period after the initial activation step when HCl and ClONO₂ are titrated against each other on the first occurrence of heterogeneous reactivity when temperatures sink below the threshold for chlorine activation. If under such circumstances, likely in polar night, ice particles form days or weeks after the initial activation step, they will have very little impact on chlorine activation.

The MLS measurements of HNO₃, ClO and O₃ in Antarctic winter and spring of 2007 are close to the EMAC simulations. The observed time evolution of these three substances

is represented very reasonably in EMAC, even if the HNO_3 mixing ratios slightly smaller and O_3 values slightly higher than in MLS.

An open question that remains is the influence of our assumed NAT number densities and size distribution based on Fahey et al. (2001). In other studies (e.g., Northway et al., 2002; Voigt et al., 2005; Pitts et al., 2011) different NAT number densities and particle size distributions are observed. But the current estimates of NAT number densities are rather low and in the range of Fahey et al. (2001). In comparison to the surface area densities of liquid particles and ice particles the NAT surface area densities are low in all studies. Therefore, the influence of the assumed NAT number densities and size distribution should be small.

Here in this study we have confirmed the dominant role of liquid particles to polar chlorine activation for Antarctic conditions. This finding should be taken into account in studies of future polar ozone development when the stratospheric halogen loading declines. By confirming the major contribution of liquid particles to chlorine activation some of the latest published findings could be essential for the future study of the stratospheric polar ozone development. Hence, regarding the future polar ozone development evolution it will on the one hand be crucial how strong the future cooling in the lower and middle stratosphere will be in polar winters and springs and how strong this will be affecting the occurrence and heterogeneous reactivity of PSCs (e.g., Butchart et al., 2010; Bohlinger et al., 2014). On the other hand also the future development increase of the background stratospheric aerosol, as discussed in by Solomon et al. (2011), can also lead to an altered distribution of liquid particles with implications for the polar chlorine activation.

Acknowledgements. We acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Karlsruhe Institute of Technology. We thank the MLS team for providing us their data. This MLS data were acquired as part of the activities of NASA's Science Mission Directorate, and are archived and distributed by the Goddard Earth Sciences (GES) Data and Information Services Center (DISC).

References

- Abbatt, J.P.D. and Molina, M.J.: Heterogeneous interactions of ClONO_2 and HCl on nitric acid trihydrate at 202 K, *J. Phys. Chem.*, 96, 7674-7679, doi:10.1021/j100198a036, 1992.
- Atkinson, R., Baulch, D.L., Cox, R.A., Crowley, J.N., Hampson, R.F., Hynes, R.G., Jenkin, M.E., Rossi, M.J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III gas phase reactions of inorganic halogens, *Atmos. Chem. Phys.*, 7, 981-1191, doi:10.5194/acp-7-981-2007, 2007.
- Bohlinger, P., Sinnhuber, B.-M., Ruhnke, R., and Kirner, O.: Radiative and dynamical contributions to past and future Arctic stratospheric temperature trends, *Atmos. Chem. Phys.*, 14, 1679-1688, doi:10.5194/acp-14-1679-2014, 2014.
- Buchholz, J.: Simulations of physics and chemistry of polar stratospheric clouds with a general circulation model, Ph.D. thesis, Johannes Gutenberg University Mainz, 2005.
- Butchart, N., Cionni, I., Eyring, V., Shepherd, T.G., Waugh, D.W., Akiyoshi, H., Austin, J., Brühl, C., Chipperfield, M.P., Cordero, E., Dameris, M., Deckert, R., Dhomse, S., Frith, S.M., Garcia, R.R., Gettelman, A., Giorgetta, M.A., Kinnison, D.E., Li, F., Mancini, E., McLandress, C., Pawson, S., Pitari, G., Plummer, D.A., Rozanov, E., Sassi, F., Scinocca, J.F., Shibata, K., and Tian, W.: Chemistry-climate model simulations of 21st century stratospheric climate and circulation changes, *J. Climate*, 23, 5349-5374, doi:10.1175/2010JCLI3404.1, 2010.
- Carslaw, K.S., Luo B., and Peter T.: An analytic expression for the composition of aqueous HNO_3 - H_2SO_4 stratospheric aerosols including gas phase removal of HNO_3 , *Geophys. Res. Lett.*, 22, 14, 1877-1880, doi:10.1029/95GL01668, 1995.
- Carslaw, K.S. and Peter, T.: Uncertainties in reactive uptake coefficients for solid stratospheric particles - 1. Surface chemistry, *Geophys. Res. Lett.*, 24, 1743-1746, 1997.
- Carslaw, K.S., Peter, T., and Müller, R.: Uncertainties in Reactive Uptake Coefficients for Solid Stratospheric Particles - 2. Effect on Ozone Depletion, *Geophys. Res. Lett.*, 24, 1747-1750, 1997.
- Carslaw, K.S., Kettleborough, J.A., Northway, M.J., Davies, S., Gao, R.S., Fahey, D.W., Baumgardner, D.G., Chipperfield, M.P., and Kleinböhl, A.: A vortex-scale simulation of the growth and sedimentation of large nitric acid hydrate particles, *J. Geophys. Res.*, 107, 8300, doi:10.1029/2001JD000467, 2002.
- Crutzen, P.J., Müller, R., Brühl, C., and Peter, T.: On the potential importance of the gas-phase reaction $\text{CH}_3\text{O}_2 + \text{ClO} \rightarrow \text{ClOO} + \text{CH}_3\text{O}$ and the heterogeneous reaction $\text{HOCl} + \text{HCl} \rightarrow \text{H}_2\text{O} + \text{Cl}_2$ in ozone hole chemistry, *Geophysical Research Letters*, 19, 1113-1116, doi:10.1029/92GL01172, 1992.
- Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A.P., Monge-Sanz, B.M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.*, 137, 553-597, doi: 10.1002/qj.828, 2011.
- Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the polar stratosphere, *Ann. Geophys.*, 30, 1055-1073, doi:10.5194/angeo-30-1055-2012, 2012.
- Dye, J.E., Baumgardner, D., Gandrud, B.W., Kawa, S.A., Kelly, K.K., Lowenstein, M., Ferry, G.V., Chan, K.R., and Gary, B.L.: Particle size distributions in Arctic polar stratospheric clouds, growth and freezing of sulphuric acid droplets, and implications for cloud formation. *J. Geophys. Res.* 97, 8015-8034, doi:10.1029/91JD02740, 1992.
- Fahey, D.W., Gao, R.S., Carslaw, K.S., Kettleborough, J., Popp, P.J., Northway, M.J., Holecek, J.C., Ciciora, S.C., McLaughlin, R.J., Thompson, T.L., Winkler, R.H., Baumgardner, D.G., Gandrud, B., Wennberg, P.O., Dhaniyala, S., McKinney, K., Peter, T., Salawitch, R.J., Bui, T.P., Elkins, J.W., Webster, C.R., Atlas, E.L., Jost, H., Wilson, J.C., Herman, R.L., Kleinböhl, A., and von König, M.: The Detection of Large HNO_3 -Containing Parti-

- cles in the Winter Arctic Stratosphere, *Science*, 291, 1026-1031, doi:10.1126/science.1057265, 2001
- Froidevaux, L., Jiang, Y.B., Lambert, A., Livesey, N.J., Read, W.G., Waters, J.W., Browell, E.V., Hair, J.W., Avery, M.A., McGee, T.J., Twigg, L.W., Sumnicht, G.K., Jucks, K.W., Margitan, J.J., Sen, B., Stachnik, R.A., Toon, G.C., Bernath, P.F., Boone, C.D., Walker, K.A., Filipiak, M.J., Harwood, R.S., Fuller, R.A., Manney, G.L., Schwartz, M.J., Daffer, W.H., Drouin, B.J., Cofield, R.E., Cuddy, D.T., Jarnot, R.F., Knosp, B.W., Perner, V.S., Snyder, W.V., Stek, P.C., Thurstans, R.P., and Wagner, P.A.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *J. Geophys. Res.*, 113, D15S20, doi:10.1029/2007JD008771, 2008.
- Groß, J.-U.: Modelling of Stratospheric Chemistry based on HALOE/UARS Satellite Data, Ph.D. thesis, Universität Mainz, 1996.
- Groß, J.-U., Brauttsch, K., Pommrich, R., Solomon, S., and Müller, R.: Stratospheric ozone chemistry in the Antarctic: what determines the lowest ozone values reached and their recovery?, *Atmos. Chem. Phys.*, 11, 12217-12226, doi:10.5194/acp-11-12217-2011, 2011.
- Hanson, D.R., and Mauersberger K.: Laboratory studies of the nitric acid trihydrate: implications for the south polar stratosphere, *Geophys. Res. Lett.*, 15, 855-858, doi:10.1029/GL015i008p00855, 1988.
- Hanson, D.R., and Ravishankara, A.R.: Reaction of ClONO₂ with HCl on NAT, NAD, and frozen sulfuric acid and hydrolysis of N₂O₅ and ClONO₂ on frozen sulfuric acid, *J. Geophys. Res.*, 96, 22931-22936, doi:10.1029/93JD01929, 1993.
- Hanson, D.R., and Ravishankara, A.R.: Reactive uptake of ClONO₂ onto sulfuric acid due to reaction with HCl and H₂O, *J. Phys. Chem.*, 98, 5728-5735, doi:10.1021/j100073a026, 1994.
- Hanson, D.R., Ravishankara, A.R., and Solomon, S.: Heterogeneous reactions in sulfuric acid aerosols: A framework for model calculations, *J. Geophys. Res.*, 99, 3615-3629, doi:10.1029/93JD02932, 1994.
- Höpfner, M., Luo, B.P., Massoli, P., Cairo, F., Spang, R., Snels, M., Di Donfrancesco, G., Stiller, G., von Clarmann, T., Fischer, H., and Biermann, U.: Spectroscopic evidence for NAT, STS, and ice in MIPAS infrared limb emission measurements of polar stratospheric clouds, *Atmos. Chem. Phys.*, 6, 1201-1219, doi:10.5194/acp-6-1201-2006, 2006
- IPCC (Intergovernmental Panel on Climate Change): Climate Change 2007: Synthesis Report. A Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Core Writing Team, Pachauri, R. K., and Reisinger, A., IPCC, Geneva, Switzerland, 2007.
- Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy)-a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433-444, doi:10.5194/acp-5-433-2005, 2005.
- Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M.G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmos. Chem. Phys.*, 6, 5067-5104, doi:10.5194/acp-6-5067-2006, 2006.
- Jöckel, P., Kerkweg, A., Buchholz-Dietsch, J., Tost, H., Sander, R., and Pozzer, A.: Technical note: Coupling of chemical processes with the Modular Earth Submodel System (MESSy) submodel TRACER, *Atmos. Chem. Phys.*, 8, 1677-1687, doi:10.5194/acp-8-1677-2008, 2008.
- Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical note: An implementation of the dry removal processes DRY-DEPosition and sedimentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617-4632, doi:10.5194/acp-6-4617-2006, 2006a.
- Kerkweg, A., Sander, R., Tost, H., and Jöckel, P.: Technical note: Implementation of prescribed (OFFLEM), calculated (ON-LEM), and pseudo-emissions (TNUDGE) of chemical species in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 3603-3609, doi:10.5194/acp-6-3603-2006, 2006.
- Livesey, N.J., Read, W.G., Froidevaux, L., Lambert, A., Manney, G.L., Pumphrey, H.C., Santee, M.L., Schwartz, M.J., Wang, S., Cofield, R.E., Cuddy, D.T., Fuller, R.A., Jarnot, R.F., Jiang, J.H., Knosp, B.W., Stek, P.C., Wagner, P.A., and Wu, D.L.: Earth Observing System (EOS) Aura Microwave Limb Sounder (MLS) Version 3.3 Level 2 data quality and description document, available at: <http://mls.jpl.nasa.gov/data/datadocs.php> (last access: 06 November 2014), 2011.
- Kirner, O.: Prozessstudien der stratosphärischen Chemie und Dynamik mit Hilfe des Chemie-Klima-Modells ECHAM5/MESSy1, Ph.D. thesis, Universität Karlsruhe, 2008.
- Kirner, O., Ruhnke, R., Buchholz-Dietsch, J., Jöckel, P., Brühl, C., and Steil, B.: Simulation of polar stratospheric clouds in the chemistry-climate-model EMAC via the submodel PSC, *Geosci. Model Dev.*, 4, 169-182, doi:10.5194/gmd-4-169-2011, 2011.
- Landgraf, J., and Crutzen, P.J.: An efficient method for online calculations of photolysis and heating rates, *J. Atmos. Sci.*, 55, 863-878, doi:10.1175/1520-0469(1998)055<0863:AEMFOC>2.0.CO;2, 1998.
- Lowe, D., and MacKenzie, A.R.: Polar stratospheric cloud microphysics and chemistry, *J. Atmos. Sol.-Terr. Phys.*, 70, 13-40, doi:10.1016/j.jastp.2007.09.011, 2008.
- Marti, J., and Mauersberger, K.: A survey and new measurements of ice vapor pressure at temperatures between 170 and 250 K, *Geophys. Res. Lett.*, 20, 363-366, doi:10.1029/93GL00105, 1993.
- Meilinger, S.K.: Heterogeneous Chemistry in the Tropopause Region: Impact of Aircraft Emissions, Ph.D. thesis, Swiss Federal Institute of Technology (ETH), Zürich, 2000.
- Müller, R., Peter, T., Crutzen, P.J., Oelhaf, H., Adrian, G.P., Von Clarmann, T., Wegner, A., Schmidt, U., and Lary, D.: Chlorine chemistry and the potential for ozone depletion in the arctic stratosphere in the winter of 1991/92, *Geophys. Res. Lett.*, 21, 14271430, doi:10.1029/94GL00465, 1994.
- Northway, M.J., Gao, R.S., Popp, P.J., Holecek, J.C., Fahey, D.W., Carslaw, K.S., Tolbert, M.A., Lait, L.R., Dhaniyala, S., Flagan, R.C., Wennberg, P.O., Mahoney, M.J., Herman, R.L., Toon, G.C., and Bui, T.P.: An analysis of large HNO₃-containing particles sampled in the Arctic stratosphere during the winter of 1999/2000, *J. Geophys. Res.*, 107(D20), 8298, doi:10.1029/2001JD001079, 2002.
- Peter, T. and Groß, J.-U.: Polar Stratospheric Clouds and Sulfate Aerosol Particles: Microphysics, Denitrification and Heteroge-

- neous Chemistry, in: *Stratospheric Ozone Depletion and Climate Change*, edited by Müller, R., pp. 108–144, Royal Society of Chemistry, doi:10.1039/9781849733182-00108, ISBN: 978-1-84973-002-0, 2012.
- 830 Pitts, M.C., Poole, L.R., Dörnbrack, A., and Thomason, L.W.: 890
The 2009/2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, doi:10.5194/acp-11-2161-2011, 2011.
- 835 Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Koernblueh, L., Manzini, E., Schlese, U., and Schulzweida, 895
U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Climate*, 19, 3771–3791, doi:10.1175/JCLI3824.1, 2006.
- 840 Sander, R., Kerkweg, A., Jöckel, P., and Lelieveld, J.: Technical note: The new comprehensive atmospheric chemistry module 900
MECCA, *Atmos. Chem. Phys.*, 5, 445–450, doi:10.5194/acp-5-445-2005, 2005.
- 845 Sander, S.P., Abbatt, J., Barker, J.R., Burkholder, J.B., Friedl, R.R., Golden, D.M., Huie, R.E., Kolb, C.E. Kurylo, M.J., Moortgat, G.K., Orkin, V.L. and Wine, P.H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 17, JPL Publication 10-6, Jet Propulsion Laboratory, Pasadena, 2011.
- 850 Santee, M. L., MacKenzie, I.A., Manney, G.L., Chipperfield, M.P., Bernath, P.F., Walker, K.A., Boone, C.D., Froidevaux, L., 910
Livesey, N.J., and Waters, J.W.: A study of stratospheric chlorine partitioning based on new satellite measurements and modeling, *J. Geophys. Res.*, 113, D12307, doi:10.1029/2007JD009057, doi:10.1029/2007JD009057, 2008.
- 855 Schlager, H., and Arnold, F.: Measurement of stratospheric gaseous 915
nitric acid in the Winter arctic vortex using a novel rocket-borne mass spectrometer method, *Geophys. Res. Lett.*, 17, 433–436, doi:10.1029/GL017i004p00433, 1990.
- 860 Shi, Q., Jayne, J.T., Kolb, C.E., Worsnop, D.R., and Davidovits, P.: Kinetic model for reaction of ClONO₂ with H₂O and HCl and 920
HOCl with HCl in sulfuric acid solutions, *J. Geophys. Res.*, 106, 24,259–24,274, doi:10.1029/2000JD000181, 2001.
- 865 Solomon, S., Garcia, R.R., Rowland, F.S., and Wuebbles, D.J.: On the depletion of Antarctic ozone, *Nature*, 321, 755–758, doi:10.1038/321755a0, 1986. 925
- Solomon, S., Daniel, J.S., Neely, R.R., Vernier, J.P., Dutton, E.G., and Thomason, L.W.: The Persistently Variable Background Stratospheric Aerosol Layer and Global Climate Change, *Science*, 333, 6044, 866–870, doi:10.1126/science.1206027, 2011.
- 870 Solomon, S., Haskins, J., Ivy, D. J., and Min, F.: Fundamental differences between Arctic and Antarctic ozone depletion, *Proc. Natl. Acad. Sci.*, 111, 6220–6225, doi:10.1073/pnas.1319307111, 2014. 930
- 875 Tabazadeh, A., Toon, O.B., and Jensen, E.J.: Formation and implications of ice particle nucleation in the stratosphere, *Geophys. Res. Lett.*, 24, 2007–2010, doi:10.1029/97GL01883, 1997. 935
- Toon, O.B., Hamill, P., Turco, R.P., and Pinto, J.: Condensation of HNO₃ and HCl in the winter polar stratospheres, *Geophys. Res. Lett.*, 13, 1284–1287, doi:10.1029/GL013i012p01284, 1996. 880
- ~~Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive scavenging submodel for global atmospheric chemistry modelling, *Atmos. Chem. Phys.*, 6, 565–574, doi:10.5194/acp-6-565-2006, 2006a.~~
- 885 Tost, H., Jöckel, P., and Lelieveld, J.: Influence of different convection parameterisations in a GCM, *Atmos. Chem. Phys.*, 6, 5475–5493, doi:10.5194/acp-6-5475-2006, 2006.
- ~~Tost, H., Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., and Lelieveld, J.: Global cloud and precipitation chemistry and wet deposition: tropospheric model simulations with ECHAM5/MESSy1, *Atmos. Chem. Phys.*, 7, 2733–2757, doi:10.5194/acp-7-2733-2007, 2007a.~~
- ~~Tost, H., Jöckel, P., and Lelieveld, J.: Lightning and convection parameterisations—uncertainties in global modelling, *Atmos. Chem. Phys.*, 7, 4553–4568, doi:10.5194/acp-7-4553-2007, 2007b.~~
- Van Aalst, M.K.: Dynamics and Transport in the Stratosphere - simulations with a general circulation model, Ph.D. thesis, Institute for Marine and Atmospheric Research Utrecht, Netherlands, 2005.
- Van den Broek, M.M.P., Williams, J.E., and Bregman, A.: Implementing growth and sedimentation of NAT particles in a global Eulerian model, *Atmos. Chem. Phys.*, 4, 1869–1883, doi:10.5194/acp-4-1869-2004, 2004.
- Voigt, C., Schlager, H., Luo, B.P., Dörnbrack, A., Roiger, A., Stock, P., Curtius, J., Vössing, H., Borrmann, S., Davies, S., Konopka, P., Schiller, C., Shur, G., and Peter, T.: Nitric Acid Trihydrate (NAT) formation at low NAT supersaturation in Polar Stratospheric Clouds (PSCs), *Atmos. Chem. Phys.*, 5, 1371–1380, doi:10.5194/acp-5-1371-2005, 2005.
- Waters, J.W., Froidevaux, L., Harwood, R.S., Jarnot, R.F., Pickert, H.M., Read, W.G., Siegel, P.H., Cofield, R.E., Filipiak, M.J., Flower, D.A., Holden, J.R., Lau, G.K., Livesey, N.J., Manney, G.L., Pumphrey, H.C., Santee, M.L., Wu, D.L., Cuddy, D.T., Lay, R.R., Loo, M.S., Perun, V.S., Schwartz, M.J., Stek, P.C., Thurstans, R.P., Boyles, M.A., Chandra, K.M., Chavez, M.C., Chen, G.S., Chudasama, B.V., Dodge, R., Fuller, R.A., Girard, M.A., Jiang, J.H., Jiang, Y.B., Knosp, B.W., LaBelle, R.C., Lam, J.C., Lee, K.A., Miller, D., Oswald, J.E., Patel, N.C., Pukala, D.M., Quintero, O., Scaff, D.M., Van Snyder, W., Tope, M.C., Wagner, P.A., and Walch, M.J.: The Earth Observing System Microwave Limb Sounder (EOS MLS) on the Aura satellite, *IEEE T. Geosci. Remote*, 44, 1075–1092, doi:10.1109/TGRS.2006.873771, 2006.
- Wegner, T., Groöb, J.-U., von Hobe, M., Stroth, F., Sumińska-Ebersoldt, O., Volk, C.M., Hösen, E., Mitev, V., Shur, G., and Müller, R.: Heterogeneous chlorine activation on stratospheric aerosols and clouds in the Arctic polar vortex, *Atmos. Chem. Phys.*, 12, 11095–11106, doi:10.5194/acp-12-11095-2012, 2012.
- Wohlmann, I., Wegner, T., Müller, R., Lehmann, R., Rex, M., Manney, G.L., Santee, M.L., Bernath, P., Sumiska-Ebersoldt, O., Stroth, F., von Hobe, M., Volk, C.M., Hösen, E., Ravagnani, F., Ulanovsky, A., and Yushkov, V.: Uncertainties in modelling heterogeneous chemistry and Arctic ozone depletion in the winter 2009/2010, *Atmos. Chem. Phys.*, 13, 3909–3929, doi:10.5194/acp-13-3909-2013, 2013.
- WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and Monitoring Project Report No. 50, Geneva, Switzerland, 2007.

Table 1. The eight size bins in submodel PSC used for the kinetic growth NAT parameterisation.

| Size bin | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|--|------|------|------|------|------|------|------|------|
| Minimum NAT radius (μm) | 0.0 | 0.2 | 1.0 | 2.0 | 6.0 | 9.0 | 12.0 | 16.0 |
| Maximum NAT radius (μm) | 0.2 | 1.0 | 2.0 | 6.0 | 9.0 | 12.0 | 16.0 | 20.0 |
| Mean NAT radius ($r_{\text{NAT}_{\text{bin}}}$ in μm) | 0.1 | 0.6 | 1.5 | 4.0 | 7.5 | 10.5 | 14.0 | 18.0 |
| Maximum of NAT number density ($N_{\text{NAT}_{\text{bin}}}$ in 10^{-5} particles cm^{-3}) | 3.28 | 3.28 | 3.28 | 3.28 | 3.28 | 3.28 | 1.64 | 1.64 |

Table 2. The four performed EMAC simulations with different assumptions for heterogeneous chemistry.

| Simulations | Time period | Heterogeneous chemistry |
|------------------|-------------|----------------------------------|
| <i>Standard</i> | 2000–2012 | on liquid, NAT and ice particles |
| <i>NoHet</i> | 2005–2012 | none |
| <i>Liquid</i> | 2005–2012 | on liquid particles |
| <i>LiquidNAT</i> | 2005–2012 | on liquid and NAT particles |




Fig. 1. Denitrification in the standard simulation and from MLS – time series of simulated temperature in K (top row) and the simulated distribution of N_{NAT} in cm^{-3} (second row) and HNO_3 in ppbv from EMAC (third row) and HNO_3 in ppbv from MLS (bottom row) for southern polar latitudes on 46 hPa approximately 21.6 km for the dates 12 May 2007, 19 May 2007, and 31 May 2007. In the temperature plots the isothermes of 195 K (white) and 192 K (light blue) are marked.




Fig. 2. Stratospheric temperature, liquid and solid particles – time series averaged from 80° to 90° S and over the years 2005 to 2009 for temperature in K (top left), number densities of NAT (N_{NAT}) and ice particles (N_{ICE}) in cm^{-3} (top center and right), as well as surface area densities of liquid particles (A_{LIQ}), NAT (A_{NAT}) and ice particles (A_{ICE}) in $\mu\text{m}^2 \text{cm}^{-3}$ (bottom left to right). **The results shown are valid for the standard and for all three sensitivity simulations.**




Fig. 3. Chlorine activation – time series of ClO_x in ppbv in the standard simulation during Antarctic winter and spring ~~in~~ **averaged over** 80° to 90° S ~~averaged over~~ and the years 2005 to 2009 (top left) and contribution of liquid particles (top right), additional contribution by activation on NAT particles (bottom left) and additional contribution by activation on ice particles (bottom right). ClO_x is defined as the sum of $\text{ClO} + \text{OClO} + \text{HOCl} + 2\text{Cl}_2 + 2\text{Cl}_2\text{O}_2$. **Note that the color bars have different units.**




Fig. 4. Comparison of ClO from EMAC with MLS observations – time series of ClO in ppbv from MLS (left) and from the standard EMAC simulation (right) during Antarctic winter and spring of 2007 in 79.5° S.




Fig. 5. Ozone depletion – time series of O₃ in ppbv in the standard EMAC simulation during Antarctic winter and spring averaged over 80° to 90° S averaged over and the years 2005 to 2009 (top left) and contribution of liquid particles (top right), additional NAT particles (bottom left) and additional ice particles (bottom right). Note that the color bars have different units.




Fig. 6. Comparison of O₃ from EMAC with MLS observations – time series of O₃ in ppbv from MLS (left) and from the standard EMAC simulation (right) during Antarctic winter and spring of 2007 in 79.5° S.




Fig. 7. Ozone columns – development of ozone column in DU in the standard simulation during Antarctic winter and spring in the Southern Hemisphere averaged over the years 2005 to 2009 (top left) and contribution of liquid particles (top right), additional NAT particles (bottom left) and additional ice particles (bottom right) to ozone depletion. Note that the color bars have different units.