



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

A pathway analysis of global aerosol processes

N. A. J. Schutgens and P. Stier

Department of Physics, University of Oxford, Parks Road, OX1 3PU, UK

Received: 15 May 2014 – Accepted: 17 May 2014 – Published: 10 June 2014

Correspondence to: N. A. J. Schutgens (schutgens@physics.ox.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We present a detailed budget of the changes in atmospheric aerosol mass and numbers due to various processes: emission, nucleation, coagulation, H₂SO₄ condensation and in-cloud production, ageing and deposition. The budget is created from monthly-averaged tracer tendencies calculated by the global aerosol model ECHAM5.5-HAM2 and allows us to investigate process contributions at various length- and time-scales. As a result, we show in unprecedented detail what processes drive the evolution of aerosol. In particular, we show that the processes that affect aerosol masses are quite different from those affecting aerosol numbers. Condensation of H₂SO₄ gas onto pre-existing particles is an important process, dominating the growth of small particles in the nucleation mode to the Aitken mode and the ageing of hydrophobic matter. Together with in-cloud production of H₂SO₄, it significantly contributes to (and often dominates) the mass burden (and hence composition) of the hydrophilic Aitken and accumulation mode particles. Particle growth itself is the leading source of number densities in the hydrophilic Aitken and accumulation modes, with their hydrophobic counterparts contributing (even locally) relatively little. As expected, the coarse mode is dominated by primary emissions and mostly decoupled from the smaller modes. Our analysis also suggests that coagulation serves mainly as a loss process for number densities and that, relative to other processes, it is a rather unimportant contributor to composition changes of aerosol. The analysis is extended with sensitivity studies where the impact of a lower model resolution or pre-industrial emissions is shown to be small. We discuss the use of the current budget for model simplification, prioritisation of model improvements, identification of potential structural model errors and model evaluation against observations.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

1 Introduction

In the past decennia, the importance of atmospheric aerosol to the Earth's climate and biosphere has become increasingly clear. Aerosols change the global radiation budget both directly (Angstrom, 1962) and indirectly (Twomey, 1974; Albrecht, 1989). They affect the temperature structure of the atmosphere (Hansen et al., 1997; Lohmann and Feichter, 2005) and may have consequences for the hydrological cycle (Lohmann and Feichter, 1997). Dust aerosols transport nutrients for the biosphere over long distances (Vink and Measures, 2001; McTainsh and Strong, 2007; Maher et al., 2010; Lequy et al., 2012). Finally, anthropogenic aerosols can pose health hazards for humans (Dockery et al., 1993; Brunekreef and Holgate, 2002; Ezzati et al., 2002; Smith et al., 2009; Beelen et al., 2013).

To increase our understanding of the role played by aerosols, increasingly sophisticated models have been built over the past 10–20 years (Ghan and Schwartz, 2007). The early models represented only a single aerosol species like sulfate or dust, e.g. Langner and Rodhe (1991), Feichter et al. (1996), Tegen and Lacis (1996), Roelofs et al. (1998), Lohmann et al. (1999), and Rasch et al. (2000). Following that, models started to simulate a combination of aerosol species that were externally mixed, (Take-mura et al., 2000; Chin et al., 2002). These days several models exist that allow time-varying sizes and species compositions of aerosol particles, e.g. Easter et al. (2004), Liao et al. (2004), Liu et al. (2005), Stier et al. (2005), Spracklen et al. (2005), Bauer et al. (2008), Seland et al. (2008), Yu and Luo (2009), and Aan de Brugh et al. (2011). This increase in sophistication allowed the inclusion of more processes: from emission, transport and deposition for the earlier models to new particle formation, coagulation and gaseous condensation in the latter. That added detail in simulation comes at a cost in the form of CPU resources.

It is not clear to what extent the increased detail has led to better representation of the physical aerosol system. AERCOM is a community effort to intercompare global aerosol models and evaluate them against observations (Kinne et al., 2006; Schulz

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the simulation, whether there is a hierarchy in process importance and how a process operates (again, the model is treated as a black box).

Pathway analysis is a methodology to assess how aerosols are processed in a model and which processes are important. It offers a budget of the model tendencies that affect all tracers that are used to represent the aerosols. The basic idea is simple and an extension of the emission/deposition budgets that many models routinely produce. Because pathway analysis offers a relative budget of the individual processes as they are represented in a model, it offers a systematic approach for the reduction of the complexity of (global) aerosol models, thereby allowing us to build faster and leaner aerosol modules for use in climate simulations or data-assimilation experiments. Likewise, it helps identify processes that may contain important structural errors and helps prioritise model improvements.

In this paper, we present a pathway analysis for ECHAM5.5-HAM2 (Stier et al., 2005; Zhang et al., 2012) with the M7 microphysics module (Vignati et al., 2004). The challenge in this pathway analysis is to visualise the aerosol processes acting on 25 tracers (18 for masses and 7 for numbers) through 121 tendencies (17 for emissions, 25 for depositions and 79 for remaining processes like nucleation and coagulation). The M7 microphysics module in ECHAM-HAM is also implemented in regional aerosol models like COSMO and HIRLAM, and is rather similar to the GLOMAP module in the global aerosol model HadGEM-UKCA.

In this paper, we start in Sect. 2 with an explanation of the ECHAM5.5-HAM2 model and in particular the processes that govern aerosol evolution. Following, we will describe our methodology (Sect. 3). Next, we will briefly discuss a year-long simulation of ECHAM5.5-HAM2 with a special focus on the contribution from the different aerosol modes (Sect. 4.1). In Sect. 4.2, the global lifetimes of aerosol species, aerosol modes and aerosol species per mode will be discussed, while Sect. 4.3 examines the global transfer of mass and numbers among the aerosol modes. A detailed analysis of processes per aerosol mode is given in Sect. 4.4. The sensitivity of this analysis to model

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

assumptions like grid resolution and emission dataset is considered in Sect. 4.5. We will summarise our work and its results in Sect. 5.

2 The ECHAM-HAM model

The global aerosol model ECHAM-HAM consists of an aerosol module HAM (Stier et al., 2005, 2007) coupled to an atmospheric general circulation model ECHAM (Roeckner et al., 2003, 2006). ECHAM-HAM has been used to study non-linearities in aerosol response due to emission changes (Stier et al., 2006a), aerosol effects in a transient climate (Stier et al., 2006b), aerosol activation and cloud-processing (Roelofs et al., 2006), aerosol indirect effects (Lohmann et al., 2007), the impact of pollution mitigation on climate forcing (Kloster et al., 2008), the impact of volcanic eruptions on climate (Niemeier et al., 2009; Timmreck et al., 2010), the impact of aerosol nucleation on radiative forcing (Makkonen et al., 2009; Kazil et al., 2010), dimming and brightening of surface radiation due to aerosols (Folini and Wild, 2011), climate forcing due to secondary organic aerosols (O'Donnell et al., 2011) and aerosol indirect effects due to shipping emissions (Peters et al., 2012) to name but a few studies.

The general circulation model ECHAM was developed at the Max Planck Institute for Meteorology and evolved from the model at the European Centre for Medium-Range Weather Forecasting. It solves the prognostic equations for vorticity, divergence, surface pressure and temperature using spherical harmonics with triangular truncation. Tracers like water vapour, liquid and ice hydrometeors, various trace gases as well as aerosols are advected with a flux-form semi-Lagrangian transport scheme (Lin and Rood, 1996) on a Gaussian grid. ECHAM can be nudged to meteorological reanalysis fields. More details can be found in Roeckner et al. (2003, 2006).

The aerosol module HAM calculates the global evolution of five aerosol species: sulfate (SO_4), particulate organic matter (POM), black carbon (BC), sea salt (SS) and dust (DU). These species are the constituents of both internally and externally mixed aerosol particles whose size distribution is represented by 7 uni-modal log-normal dis-

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tributions called modes. These 7 modes describe four size classes (nucleation, Aitken, accumulation and coarse) and two hygroscopic classes (hydrophobic and hydrophilic). Most of these modes contain time- and space-varying mixtures of aerosol species, see Table 1. To predict aerosol evolution, ECHAM-HAM (without explicit SOA, see later) uses 25 tracers: 7 for number mixing ratios and 18 for mass mixing ratios. H₂SO₄ nucleation and condensation, coagulation and ageing are handled by the M7 sub-module by Vignati et al. (2004). The processes described in M7 cause redistribution of aerosol mass and numbers among the modes. Part of HAM is a sulphur cycle model (Feichter et al., 1996) that predicts the evolution of dimethyl sulphide (DMS), sulphur dioxide (SO₂) and gaseous sulfate (SO₄⁻) using monthly mean fields of the oxidants OH, H₂O₂, NO₂ and O₃, calculated off-line by the MOZART chemical transport model (Horowitz et al., 2003). The aerosol in HAM is affected by the meteorology calculated by ECHAM, and in turn provides feedback to ECHAM as it affects atmospheric radiative transfer and cloud microphysics.

Over time, various improvements have been made to ECHAM-HAM and currently a distinction is made between the initial version HAM 1 (Stier et al., 2005) and the newer version HAM 2 (Zhang et al., 2012). While using the same modal structure (Table 1), HAM 2 added new parametrisations for nucleation, sea salt and dust emissions, a water uptake scheme based on κ -Köhler theory and an explicit scheme for secondary organic aerosol (SOA) formation. For a detailed overview of the differences between HAM 1 and HAM 2, see Zhang et al. (2012) who also define the default choices for an ECHAM5.5-HAM2 experiment.

In this paper, we will use ECHAM5.5-HAM2, nudged to ERA-interim meteorological reanalysis. To reduce the complexity of the analysis, we chose not to use the default option in HAM 2 to use explicit SOA modelling and instead use the implicit scheme from HAM 1 where scaled biogenic emissions are assumed to directly condense into pre-existing aerosol (SOA mass formation).

2.1 Aerosol processes in ECHAM-HAM

An overview of all possible processes is shown in Table 2. It lists the physical or chemical process, the code used in many of this paper's figures, the explicit pathways and the species involved. Pathways are indicated by right-pointing arrows, with the numbers on either side indicating the involved modes (1: nucleation, 2: hydrophilic Aitken, 3: hydrophilic accumulation, 4: hydrophilic coarse, 5: hydrophobic Aitken, 6: hydrophobic accumulation, 7: hydrophobic coarse). On the left of the arrow, the mode that loses mass ("m") or numbers ("n"), and on the right of the arrow the mode that gains mass or numbers are shown. Some processes instantaneously lead to ageing of particles. As an example take condensation of H_2SO_4 onto hydrophobic accumulation dust particles. This involves the SO_4 species, which condenses on the hydrophobic accumulation mode which has no mass tracer for sulfate. Instead, the condensed sulfate and a portion of the dust particles are immediately moved to the hydrophilic accumulation mode.

2.2 Primary emissions

The emissions of sea salt and dust are based on diagnosed 10 m wind speeds from the model. Sea salt uses the emission parametrisations by Monahan et al. (1986) and Smith and Harrison (1998) as suggested by Guelle et al. (2001). The species is emitted into the hydrophilic accumulation and coarse modes only. Dust emission is based on the scheme by Tegen (2002), as modified by Stier et al. (2005). Dust is emitted into the hydrophobic accumulation and coarse modes and is hence assumed to be hydrophobic initially.

The emissions of sulfate, particulate organic matter and black carbon are based on inventories (Stier et al., 2005; Zhang et al., 2012). It is assumed that 2.5 % of emitted SO_2 (itself based on an inventory) is emitted as sulfate aerosol with prescribed size, depending on the source. Sulfate from volcanoes is injected at pre-specified heights in the atmosphere into the hydrophilic Aitken and accumulation modes. Sulfate from

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

biomass burning is likewise injected at pre-specified heights in the atmosphere into the hydrophilic Aitken and accumulation modes. Anthropogenic emissions of sulfate, on the other hand, are distributed over the hydrophilic Aitken, accumulation and coarse modes depending on the exact source.

Organic carbon emissions distribute 65 % of their mass in the hydrophilic modes and 35 % in the hydrophobic mode. Most sources of organic matter (including wild fires) cause emission into the Aitken mode. The exception is the treatment of biogenic emissions where half of the hydrophilic biogenic material condenses onto Aitken particles and the other half on accumulation particles.

Black carbon, irrespective of source, is emitted into the hydrophobic Aitken mode.

2.3 Removal processes

Wet and dry deposition as well as sedimentation (together called: removal processes) are all modelled in ECHAM-HAM. For details, we refer to Stier et al. (2005) and Zhang et al. (2012). Both rain-out and below-cloud scavenging of aerosol by water is considered and precipitation can re-evaporate. The standard version of ECHAM-HAM does not explicitly model aerosol in hydrometeors (but see Hoose et al., 2008). Dry deposition is based on a surface resistance model and sedimentation is calculated using Stokes theory. Sedimentation is not considered for the smallest particles (nucleation and Aitken). Removal processes assume that all particles within the mode have the same (wet) mode radius when calculating efficiencies and will remove particles accordingly.

2.4 Nucleation of H₂SO₄ gas

Binary nucleation of the H₂SO₄-H₂O system (Seinfeld and Pandis, 2006, Ch. 11) converts sulfate gas into sulfate aerosol. In ECHAM-HAM, two parametrisation schemes are used to calculate the formation of new nucleation mode sulfate aerosol based on environmental conditions. Binary homogenous nucleation is parametrized according to

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kazil and Lovejoy (2007) and is the main nucleation mechanism yielding large particle numbers at the top of the troposphere. Boundary layer nucleation events over forest areas is modelled according to cluster activation theory (Kulmala et al., 2006) and parametrized as in Riipinen et al. (2007).

While the parametrisation by Kazil and Lovejoy (2007) is based on theory and numerous laboratory experiments, the parametrisation by Riipinen et al. (2007) is based on observations at two (European) sites only. See also Kazil et al. (2010) for an evaluation of nucleation processes in ECHAM-HAM.

2.5 Condensation of H₂SO₄ gas

Sulfate gas condenses easily onto aerosol and cloud particles (Seinfeld and Pandis, 2006, Ch. 12 and 13). In ECHAM-HAM, a distinction is made between the condensation of sulfate in cloudy and cloud-free parts of a grid box. In the cloud-free parts, condensation of sulfate gas onto aerosol particles is calculated using the mass balance equation in Fuchs (1964), assuming that in each mode all aerosol particles have the same size (mode radius). The accommodation coefficients (“sticking” coefficients) are 1.0 for hydrophilic and 0.3 for hydrophobic modes. In the cloudy part of the gridbox, it is assumed that all available sulfate gas condenses onto the cloud particles (which are much larger than the aerosol and present a much larger surface). However, the in-cloud condensed sulfate gas is then assumed to redistribute over the aerosol in the grid box. Since HAM2 does not specifically model aerosol in cloud particles, the cloud-condensed H₂SO₄ is allocated to the largest hydrophilic mode available. If no such mode exists in a grid box, the largest hydrophobic mode available is used instead.

Condensation of H₂SO₄ onto aerosol is calculated at the mode radii only (size distribution of the modes are ignored). In practice, cloud-condensed H₂SO₄ is never allocated to any of the hydrophobic modes.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.6 In-cloud production of H₂SO₄ aerosol

A third process for the production of sulfate aerosol species is aqueous-phase oxidation of dissolved SO₂, S(IV), into dissolved SO₄, S(VI), (Seinfeld and Pandis, 2006, Ch. 7). In HAM, SO₂ dissolves in clouds into S(IV) products which are subsequently oxidized by O₃ and H₂O₂ into SO₄ (Feichter et al., 1996). The resulting sulfate aerosol species is allocated to the hydrophilic accumulation and coarse modes (distributed according to available number densities). Oxidation rates are pH-dependent. If no hydrophilic accumulation and coarse mode aerosol exists in a grid box, new hydrophilic coarse aerosol is created, although this rarely happens. In fact, almost all in-cloud produced H₂SO₄ is allocated to pre-existing hydrophilic accumulation particles.

2.7 Coagulation

In ECHAM-HAM, coagulation is modelled by considering Brownian coagulation processes for the smaller and/or more numerous modes. Coagulation results in a decrease of number densities for intra-modal coagulation (i.e. nucleation particles coagulating with their own kind) or a decrease of number density in one mode and a mass transfer to another (larger or more hydrophilic) mode for inter-modal coagulation. The model for coagulation is based on the work by Fuchs (1964) and represents the change in number density N as

$$\frac{\partial N_i}{\partial t} = 4\pi N_i \sum_j \alpha_{ij}(D_i + D_j)(r_i + r_j)\beta_{ij}N_j \quad (1)$$

where i represents the mode whose particles are captured by particles in mode j . D is the diffusivity and r the radius of the particles of a certain mode. For inter-modal coagulation $\alpha_{ij} = 1$, but for intra-modal coagulation it is $\alpha_{ij} = \frac{1}{2}$. Fuchs introduced the factor β_{ij} that corrects the Brownian diffusion of a smaller particle close to a larger particle. The coagulation efficiency is 1 for all modes.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Not all possible coagulation processes are considered in ECHAM-HAM. For a list of those represented, see Table 2. In particular, intra-modal coagulation of the hydrophobic accumulation and coarse modes and the hydrophilic coarse mode are ignored, the latter in contrast to what is mentioned in Vignati et al. (2004).

The solution of Eq. (1) for one model time-step is the analytical solution assuming that all $N_{j \neq i}$ are constant. Our model for coagulation assumes all particles of the same mode have an identical radius (being the mode radius of the number size distribution). For typical aerosol sizes, intra-modal coagulation increases as size decreases, while inter-modal coagulation is more effective when the coagulating particles are more disparate in size (Seinfeld and Pandis, 2006, Sect. 13.3.1). As a consequence, the use of a single size for all particles in a mode may have an impact on coagulation. In addition, no consideration has been given to the impact of shear flow, turbulence, gravitational settling or electrical forces on coagulation. Finally, the theory by Fuchs assumes spherical particles which may not be entirely appropriate for black carbon or dust particles.

2.8 Ageing of hydrophobic aerosol

Hydrophobic aerosol becomes hydrophilic through the collection of hydrophilic species like sulfate or carbons. In HAM, the collection of hydrophilic sulfate onto hydrophobic carbon and dust aerosols is modelled explicitly. Hydrophilic carbons by themselves do not cause ageing (we also do not consider chemical ageing). The hydrophilic species are collected in the hydrophobic modes through either condensation of sulfate gas or coagulation with hydrophilic aerosol. This collection instantaneously leads to a transfer of the hydrophilic species and part of the hydrophobic aerosol to a hydrophilic mode. The following pathways exist:

- Condensation of sulfate gas onto particles of the hydrophobic modes and subsequent transfer to hydrophilic modes of the same size class.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Coagulation of smaller hydrophilic particles with the hydrophobic modes and subsequent transfer to the hydrophilic mode of the same size class as the hydrophobic mode.
- Coagulation of particles of the hydrophobic Aitken mode with hydrophilic Aitken and accumulation modes.

The resulting mass transfer for the latter pathway (note there will be no number transfer) is calculated using the coagulation equation. The mass and numbers transfer for the first two pathways are calculated by assuming the SO_4 accumulated in one time-step coats as many hydrophobic mode particles as possible with a single layer of sulfate. Hence this calculation uses total accumulated SO_4 mass and the radius of the hydrophobic mode. Only for a steady-state will this provide an accurate estimate of the tendency.

The ageing processes represented in ECHAM-HAM are shown in Table 2 in the columns on the right.

2.9 Redistribution among hydrophilic modes

Each mode has a spatio-temporally varying mode radius that should remain within its allotted size range (see Table 1). As a result of aforementioned processes like condensation and production of H_2SO_4 or intermodal coagulation, particle sizes of hydrophilic modes may grow. When the particles becomes too large, part of the aerosol in such a hydrophilic mode will be transferred to the next larger hydrophilic mode. Conceptually, part of the assumed size distribution of the mode will extend beyond an allowed maximum size and will be moved to the next larger mode. This process is sometimes called “mode-merging” but “redistribution” is a more accurate description.

3 Methodology

In this paper, process refers to a physical or chemical event that modifies aerosol mass and number densities. The processes that we will consider are: nucleation of H₂SO₄, emission of primary aerosol, condensation of H₂SO₄ onto pre-existing aerosol, in-cloud production of liquid phase H₂SO₄, coagulation of aerosol, ageing of aerosol as well as wet and dry deposition and sedimentation. In the model, the effect of these processes is represented by tendencies. A single process can be described by several tendencies, e.g. the condensation of H₂SO₄ is handled in the model through 7 mass tendencies (one for each mode). All tendencies that modify any single aerosol mode are called the pathways into and out of that mode. Note that these tendencies do *not* include tendencies due to tracer transport but are purely due to physical and chemical processes in the atmospheric column.

Conceptually, the mass density in a grid box of the model is updated as follows:

$$m(t + \delta t) = m(t) + \delta m_{\text{emission}} + \delta m_{\text{nucleation}} + \delta m_{\text{condensation}} + \dots - \delta m_{\text{deposition}} - \dots, \quad (2)$$

where the tendencies $\delta m \geq 0$. For our analysis, model tendencies for aerosol mass and number densities due to the various processes at each time-step were stored and averaged over a month, e.g.:

$$\Delta m_{\text{emission}} = \frac{1}{T} \int_0^T \delta m_{\text{emission}} dt \quad (3)$$

Tendencies can be distinguished into gain (e.g. emission) and loss (e.g. deposition) tendencies, depending on whether they increase or decrease the tracer they affect. The sum of all gain tendencies will be called the total gain tendency. Fractional tendencies are calculated by dividing the tendency due to one (gain or loss) process by the total

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(gain or loss) tendency, e.g.:

$$f_{\text{emission}} = \frac{\Delta m_{\text{emission}}}{\sum_{\text{all gain processes}} \Delta m} \quad (4)$$

These fractional tendencies ($0 \leq f \leq 1$) provide a lot of information on which processes are dominant and which are not. Although fractional tendencies can be calculated for each model grid box, it will make more sense to calculate them for an atmospheric column, as a zonal average or for a region (see Fig. 1 for a definition of the regions used in this paper).

In this paper, we will analyse three different experiments, all conducted with ECHAM5.5-HAM2. First, a base-line experiment with present day emissions at a grid resolution of T63L31 (this indicates a triangular truncation of the spherical harmonics at zonal wave number 63, giving grid boxes of approximately $2^\circ \times 2^\circ$ or ~ 210 km at the equator, and 31 atmospheric levels). Secondly, two sensitivity studies where either the model resolution (T31L19) or the emissions (pre-industrial) were changed. The base-line experiment has a setup identical to the baseline experiment in Zhang et al. (2012), with the exception of SOA for which we use the older (implicit) scheme, where biogenic emissions are assumed to condense immediately onto pre-existing aerosol.

4 Analysis

4.1 Contributions per mode to model observables

Before analysing the processes in ECHAM5.5-HAM2, we want to understand their consequence in terms of the contributions that the different modes give to essential aerosol fields like AOT¹, PM_{2.5}², CCN³ and the distribution of black carbon mass, see Figs. 2

¹Aerosol Optical Thickness.

²Mass of Particulate Matter with sizes less than 2.5 μm .

³Cloud Condensation Nuclei.

and 3. First, we note that the global distributions of these fields are as expected; there is a notable land-ocean contrast and values tend to be highest near known sources of aerosol, either man-made or natural.

We see that in all cases that the hydrophilic accumulation mode has a significant and often dominating contribution. AOT is determined by the hydrophilic accumulation mode and the hydrophobic and hydrophilic coarse modes (note the absence of any significant contribution by the hydrophobic accumulation mode). Over polluted regions, hydrophilic accumulation mode dominates, while over the Sahara and the Saharan outflow coarse particles dominate. Over ocean, it is typically a mix of hydrophilic accumulation and coarse modes that defines AOT. In contrast, the coarse modes dominate PM_{2.5} everywhere although sizeable contributions from the hydrophilic accumulation mode exist locally. Note that over land, the hydrophobic coarse mode has usually a significant contribution. CCN at supersaturation $S = 1\%$, on the other hand are dominated by hydrophilic particles of smaller size classes: Aitken and accumulation modes. Over the remote oceans, the Aitken mode actually dominates CCN, although there is no primary emission of Aitken size particles there. At low $S = 0.1\%$ values (not shown), CCN is dominated by the hydrophilic accumulation mode. Finally, most black carbon is present in the hydrophilic accumulation mode although a sizeable mass fraction is carried by the hydrophobic Aitken mode (into which it is emitted).

4.2 Lifetimes and loss rates

In this section, we will analyse typical time-scales in our aerosol simulations. In particular, the lifetimes (Seinfeld and Pandis, 2006, Ch. 2) of aerosol species and modes will be considered,

$$\tau = \frac{B}{I} = \frac{B}{R} \quad (5)$$

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where B is the global burden of a species or mode, I its global introduction (through e.g. emissions, nucleation or coagulation) and R its global removal (through e.g. depositions or coagulation).

The definition of life-time presupposes that steady state conditions hold, at least when averaging over suitably long time-scales. We checked that the sum of all yearly-averaged tendencies for each species per mode indeed resulted in a near zero value (some discrepancies are expected). Remaining discrepancies are below 1 % for each mode.

Table 3 shows life-times for all three experiments and all species and modes involved. For the base-line experiment, we see that the species life-times are reasonable and generally agree with other studies (Textor et al., 2006). New are the separate life-times for modes and species within a mode.

We see that the nucleation mode has a short life-time of less than half a day. The life-time for numbers is even much shorter because intra-coagulation (an important process for the nucleation mode as we will later see) affects number densities but not mass densities.

The Aitken mode also has relatively short life-times, especially for POM and BC in the hydrophilic mode. These species are quickly processed into the hydrophilic accumulation mode as we will later see. The life-time of SO_4 is substantially larger than that of POM and BC, the result of a different vertical distribution. The carbons are mostly found in the boundary layer and lower troposphere, but the sulfate mostly in the free troposphere. The number life-time of hydrophilic Aitken is much larger than its mass life-time. This is again the result of the vertical distribution, where many small Aitken (sulfate-dominated) particles are found high up in the atmosphere while the hydrophilic Aitken mass is more concentrated in the lower troposphere.

As expected, the hydrophilic accumulation has the largest life-time. This is after all where aerosol accumulates. Life-times for the SO_4 , POM and BC species are consequently determined by their life-times in the hydrophilic accumulation mode. However, this is not the case for the dust and seasalt species that have the majority of their emis-

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sions in the coarse mode. All lifetimes in the hydrophilic coarse mode are smaller than in the hydrophilic accumulation mode.

For the T31L19 experiment, we see that life-times agree more-or-less with the baseline experiment. Largest differences are found for the large hydrophobic dust modes (accumulation and coarse), sea salt in the hydrophilic accumulation mode and sulfate in the hydrophilic coarse mode. Lifetimes of the nucleation and Aitken modes that are strongly affected by coagulation are surprisingly robust.

When using the pre-industrial emissions, what stands out is the sensitivity of those life-times that are influenced by the condensation of H_2SO_4 gas, which itself is the result of conversion of SO_2 . We can expect ageing to happen slower in the pre-industrial climate than in the present day (see also Stier et al., 2006a). Indeed, lifetimes for the hydrophobic modes is up to a factor 2 larger for pre-industrial emissions. In contrast, the life-times of natural aerosol in the hydrophilic accumulation and coarse modes (dust and sea salt) hardly change.

4.3 Global pathways through HAM

In this section, we analyse the global pathways of aerosol mass and number by studying the net mass and numbers transfer among the modes.

The top panel in Fig. 4 shows an annual and global average of net mass tendencies between modes and between the modes and the rest of the physical environment. Mass enters the aerosol system through the nucleation and Aitken modes as well as the hydrophobic accumulation and coarse modes. Mass leaves the system through the hydrophilic accumulation and coarse modes. Most mass enters through the hydrophobic coarse mode, followed by the hydrophilic and hydrophobic Aitken mode. This net mass flow hides the fact that e.g. a lot of seasalt is emitted and deposited in both the hydrophilic accumulation and coarse modes (for more details see Sect. 4.4). In contrast to the Aitken and coarse modes, most mass in the hydrophilic accumulation mode has first been processed in other modes.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The bottom panel in Fig. 4 shows a similar graphic but now for an annual and global average of net number tendencies. The pink arrows denote numbers that are lost due to coagulation when smaller particles impact upon larger ones. The picture is very different with most numbers entering through the nucleation mode and the large majority being “captured” by existing larger particles. Only a small fraction of nucleation particles eventually grows into hydrophilic Aitken particles but it is a very significant amount compared to other sources. Similarly, most hydrophilic accumulation mode particles result from the growth of hydrophilic Aitken particles. In contrast, the hydrophilic coarse mode seems almost isolated from the smaller modes.

4.4 Spatial distribution of processes per mode

A more detailed analysis of the pathways per mode will now be presented. We will discuss global distributions of each mode and its species composition as well as the processes that increase or reduce mass and numbers of that mode. Starting point for each discussion will be a plot as in Fig. 6.

This plot consists of six panels. The top-left panel shows the global distribution of the mass burdens. The top-right panel shows global fractional tendencies. The remaining four panels show global maps of the gain and loss tendencies for mass and numbers. In each of these panels, pie-charts are used to denote the relative contributions of either species (top-left panel) or processes (all other panels). Excepting the top-right panel, the radii of the pie-charts show the burden or total tendency using a logarithmic scaling as indicated in the legend at the bottom of each panel. Number burdens correlate fairly well with masses and are not shown. Although our baseline ECHAM5.5-HAM2 simulation was done at T63L31, results have been averaged over 64 grid-boxes for graphical clarity. Note that the top-left global map uses a different colour coding (see Table 3) than the other panels (see Table 2).

4.4.1 The nucleation mode

The global distribution (Fig. 6a) of this mode shows a marked latitudinal variation but is otherwise quite homogeneous (using a linear scale for mass densities reveals higher masses near SO_2 sources). The nucleation mode is created through nucleation and subsequent condensation of SO_4 gas molecules and depleted through capture by larger particles or redistribution to the hydrophilic Aitken mode (Fig. 6b). Intra-modal coagulation is an important process due to the large number densities in the Upper Troposphere/Lower Stratosphere (see also Fig. ??). In Sect. 4.2, we pointed out the significant difference in lifetime estimates derived from either mass or number tendencies. This discrepancy is explained by intra-modal coagulation that depletes numbers but leaves masses intact. Particle growth due to condensation will eventually lead to redistribution (Fig. 6c and d), (the absolute mass tendencies due to condensation and redistribution are correlated). Finally, we note that coagulation of nucleation mode particles with hydrophilic coarse mode or the hydrophobic modes represent negligible loss terms. Neither are wet and dry deposition important for the nucleation mode as the majority of particles are so high up in the atmosphere.

Figure ?? reveals that nucleation occurs mainly near the top of the troposphere. Below the layer, mass gain due to condensation dominates. Within the layer, coagulation is responsible for most mass loss, while below it redistribution to the hydrophilic Aitken mode is more important.

Regionally, nucleation and condensation show a very distinct annual cycle due to solar insolation (and the resulting temperature profile in the atmosphere) as shown in Fig. 7 for East Asia. Even so, the fractional tendencies do not vary much, neither for the gain nor for the loss tendencies. This is seen for all regions (see Fig. 1) we have examined although the relative contribution of condensation and nucleation will differ (see also Fig: 6c).

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.4.2 The hydrophobic Aitken mode

The global distribution of the hydrophobic Aitken mode (Fig. 8a) is very distinct and closely related to its emission sources. The hydrophobic Aitken mode is created through primary emission of organic matter and black carbon (Fig. 8b). Actually, mass burdens and mass column tendencies have very similar distributions, a consequence of the short life-times (Table 3) of less than a day. The hydrophobic Aitken mode is more or less depleted where it is created (Fig. 8d and e), with conversion into the hydrophilic mode being the major process. This conversion (ageing) is accomplished through 4 pathways: condensation of H_2SO_4 (the dominant process) and coagulation of hydrophobic Aitken with either the nucleation or the hydrophilic Aitken or accumulation modes. Although condensation of H_2SO_4 adds only a little mass to the hydrophobic Aitken mode (Fig. 8b), it causes a great loss of both mass and numbers. Only a single layer of H_2SO_4 is needed in our model to coat hydrophobic Aitken particles and turn them hydrophilic. Coagulation accounts for a bit less than 25 % of the mass and number loss tendencies on a global scale (Fig. 8b), slightly more than the combined deposition processes. Intra-coagulation is a relatively unimportant process.

There is a marked vertical separation of the ageing processes (Fig. 9): coagulation with nucleation particles is important in the nucleation layer identified previously, and the upper free troposphere, while condensation matters most in the lower troposphere and boundary layer.

The typical ageing time-scale varies greatly with location as a global map shows (Fig. 10). Far away from sulfate sources (Amazon, Africa), ageing time-scales are long (~ 2 days) but near known sites of anthropogenic SO_2 sources (e.g. Eastern Europe, India, East Asia) those time-scales are short (~ 0.3 days).

4.4.3 The hydrophilic Aitken mode

The global distribution of hydrophilic Aitken mass shown in Fig. 11a is fairly homogeneous, although its columnar composition varies. The hydrophilic Aitken mode receives

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a lot of mass and numbers from other modes, in contrast to the previous two modes discussed. A variety of processes (Fig. 11b) drive its evolution: mass derives from H_2SO_4 condensation, ageing of hydrophobic Aitken and primary emission of particulate organic matter in fairly equal measures, while numbers come mostly from redistribution from the nucleation mode. The mass tendencies (Fig. 11c and d) show a distinct ocean-land contrast. Over the oceans, H_2SO_4 condensation matters most but over land primary emission and ageing matter more. Most organic mass derives from hydrophobic Aitken particles aged through H_2SO_4 condensation (32 %) and from primary emissions (64 %), while most sulfate mass is the result of H_2SO_4 condensation (85 %, with only 1.5 % due to aged hydrophobic Aitken particles). Black carbon derives entirely from aged hydrophobic Aitken particles.

In contrast, mass loss tendencies are dominated by redistribution to the hydrophilic accumulation mode (Fig. 11b), with deposition being more important over ocean and at higher latitudes (Fig. 11d). Number loss tendencies are dominated by coagulation (mostly intra-coagulation) and again deposition (mostly over oceans and at higher latitudes). Note that redistribution to the hydrophilic accumulation mode is an important loss tendency for numbers near primary organic sources. The main process driving particle growth appears to be primary carbon emission and ageing of hydrophobic Aitken particles. Both processes introduce relatively large particles into the hydrophilic Aitken mode that will quickly be moved over to the hydrophilic accumulation mode. In contrast, at altitudes where there is a large influx of nucleation (sulfate) particles into the Aitken mode, there is little loss to the accumulation mode. Another way to put it is this: in the hydrophilic Aitken mode, particles that consist predominantly of sulfate tend to be smaller than particles that predominantly consist of carbons. This affects the redistribution tendency and leads to a significant difference in life-times (see Table 3) in the hydrophilic Aitken mode for organic matter and black carbon (0.2–0.4 days) and for sulfate (3.5 days). These life-times in turn explain why sulfate is so prominent in the global mass distribution (Fig. 11a) while the carbons are more obvious in the mass gain

tendencies (Fig. 11c), see also Eq. (5). It should be noted that wet and dry deposition of both sulfate and carbon aerosols in this mode is relatively unimportant.

The seasonal cycle of the fractional tendencies for the hydrophilic Aitken mode is shown for the Savanna region in Fig. 12. In the Savanna, there is a pronounced seasonal cycle of biomass burning emission (also seen in the hydrophobic Aitken mode) that drives the mass tendencies. The fractional mass tendencies are nevertheless rather constant: as emissions increase, the other processes scale up or down (note there is a small increase in the contributions from processes other than primary emission or ageing). The number tendencies show quite different behaviour as they are to a large extent dependent on the evolution of the nucleation mode. A similar behaviour (strong seasonal cycle in the mass tendencies, but fairly constant fractional mass tendencies) can be seen for the Amazon (not shown).

In contrast, East-Asia shows a seasonal cycle where both absolute and fractional mass tendencies vary. Here the cycle is mostly driven by the condensation of H_2SO_4 (the emission varies as well but not as much). The fractional mass tendency for H_2SO_4 condensation shows a similar cycle as the total mass tendency. The same seasonal cycle in H_2SO_4 condensation can be seen for the nucleation, accumulation and coarse modes. The growth of nucleation particles due to condensation results in increased redistribution to hydrophilic Aitken, causing a very similar cycle for the number tendencies as for the mass tendencies in hydrophilic Aitken. Similar behaviour can be seen over Europe (not shown).

4.4.4 The hydrophobic accumulation mode

The hydrophobic accumulation mode is due to the emission of dust particles (Fig. 14b). As a consequence, it is found mainly near deserts (Fig. 14a), although low levels of hydrophobic dust can be found throughout the atmosphere. As with the hydrophobic Aitken mode, mass tendencies and distributions are closely linked (Fig. 14a, c and d), although the dust lifetime is longer than that of hydrophobic Aitken, Table 3, and outflows are more conspicuous.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The main loss process for hydrophobic accumulation is ageing (conversion to hydrophilic material) through either H_2SO_4 condensation (the dominant process) or coagulation with smaller hydrophilic modes (nucleation and Aitken). Regionally, strong differences may exist: over a significant part of Eur-Asia, wet deposition is actually the dominant loss pathway for hydrophobic accumulation. Note that significant ageing happens in the outflows over ocean, west of North-Africa and west of Australia.

The hydrophobic accumulation mode contributes relatively little to overall mass of dust in the air. A more detailed description of what happens to hydrophobic dust will be postponed until the discussion of the hydrophobic coarse mode.

4.4.5 The hydrophilic accumulation mode

Of the hydrophilic modes discussed so far, the accumulation mode has the least homogeneous global distribution (Fig. 15a). This is a direct consequence of the multiple pathways that add mass and numbers to it (Fig. 15b, c and e). Primary emissions (mostly sea salt), condensation and in-cloud production of H_2SO_4 as well as redistribution from the hydrophilic Aitken mode all cause significant mass gains. These processes show very obvious regional patterns (Fig. 15c): seasalt emission dominates in the southern oceans, redistribution from hydrophilic Aitken contributes near carbon sources, and on the continents of the industrial Northern Hemisphere condensation and in-cloud production of H_2SO_4 often dominates.

As with the hydrophilic Aitken mode, there is a distinct difference between mass and numbers gain tendencies: almost all numbers derive from the hydrophilic Aitken mode, with some contribution from primary emissions. In contrast, loss tendencies are due to wet deposition mainly.

Surprisingly, even near dust sources, aged dust is not the main contributor to the hydrophilic accumulation mode. This can be seen in both the accumulation density and tendency maps.

The variety of sources for hydrophilic accumulation matter also leads to a colourful picture of the vertical distribution of species compositions, see Fig. 16. Almost every-

where sulfate is the major ingredient (mass-wise) of aerosol, except in the lower atmosphere over the southern oceans where sea salt dominates. With very few exceptions, hydrophilic accumulation aerosol is always a mixture with at least ~ 25% of its mass different from the main constituent.

Different seasonal cycles are present in regional hydrophilic accumulation aerosol, see Figs. 18–19. Over East-Asia, we see the same cycle in fractional mass tendency for H_2SO_4 condensation as we saw for the hydrophilic Aitken mode (Fig. 13). Note that total mass tendency does not change much and so any increase in H_2SO_4 condensation is met by a reduction in in-cloud H_2SO_4 production. Over the Savanna, the hydrophilic accumulation tendencies are clearly driven by emission of carbons into the Aitken mode (Fig. 12). Unsurprisingly, redistribution is the major gain term for both mass and numbers. Note that the numbers loss process due to intra-coagulation shows a strong seasonal cycle in trend with the gain of mass and numbers. Finally, we have a look at the southern oceans (Fig. 19). Although absolute gain tendencies are rather constant, there is an obvious cycle in the fractional tendencies. This cycle is again related to sulfate: mass tendencies are driven by in-cloud production while number densities are driven by condensation onto and subsequent redistribution of hydrophilic Aitken particles.

4.4.6 The hydrophobic coarse mode

Like the hydrophobic accumulation mode, the hydrophobic coarse, results from the emission of dust. The coarse particles are of course bigger than the accumulation particles, and significantly more mass although less numbers can be found in this mode than in the accumulation mode. Yet the overall description of densities and tendencies is very similar (Fig. 20).

Ageing time-scales for the hydrophobic coarse mode can be seen in Fig. 21. These time-scales can differ by a factor of ~ 10, from 0.8 days to 7 days. Along the main dust belt (North Africa, Arabian Peninsula, Asia) in particular, time-scales decrease with longitude as progressively more industrially polluted areas are encountered.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.4.7 The hydrophilic coarse mode

The hydrophilic coarse mode consists composition-wise mostly of seasalt and dust (Fig. 22a). The seasalt is due to primary emissions while the dust is aged hydrophobic coarse dust (Fig. 22c). Even though various pathways exist that bring in aerosol from smaller modes (growth of hydrophilic accumulation, coagulation) or through other processes (condensation and in-cloud production of H_2SO_4), these processes hardly matter at all (Fig. 22b). The evolution of the coarse modes is to a large extent decoupled from the other modes.

Dust outflows over ocean are not very apparent in the hydrophilic accumulation mode (Fig. 15a) but very obvious in the hydrophilic coarse mode (Fig. 22a). More-over, comparing Figs. 20 and 22, it appears that in the Asian outflow most dust is hydrophilic while near Africa and Australia the outflow really is an external mixture of hydrophobic and hydrophilic coarse dust. Asian dust sources are more inland and the dust travels over heavily industrialised regions before flowing out over the ocean, so there is plenty of time for dust to age. Ageing times over East-Asia are much reduced (Fig. 21).

There are no pathways for the hydrophilic coarse mode loss other than wet and dry deposition and sedimentation, but due to the size of the involved particles and their vertical distribution all three deposition processes play a part in removing this mode from the air.

As the coarse modes are decoupled from the other modes, their seasonal cycle can be quite different. We show the cycle over East-Asia as an example, see Fig. 23. Seasalt emission is obviously the most important source of hydrophilic coarse over East-Asia. The second major source of hydrophilic coarse matter is aged dust. Note that redistribution from hydrophilic accumulation particles into the coarse mode only contributes when the absolute tendencies are low.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.5 Sensitivity studies

The final part of this analysis concerns the sensitivity of our results to the assumptions in using a grid resolution of T63L31 and present-day emissions. Grid resolution is known to affect e.g. dust emissions, nucleation events and removal processes (note that in ECHAM-HAM, the dust emission scheme has been tuned to yield very similar results independent of resolution). Pre-industrial primary emissions of aerosol were less and in particular reduced emission of SO₂ gas will affect sulfate nucleation and condensation. Especially coagulation, which depends on a product of two number densities, may be sensitive to changes in either resolution or emissions. However, as we will show, the pathway analysis (that is, the fractional tendencies) is (are) not fundamentally affected.

Figure 24 shows regional fractional tendencies for East Asia (see Fig. 1) of all processes for both masses and numbers at two different grid resolutions, the baseline T63L31 and sensitivity T31L19 experiments. Since these data all lie closely to the $y = x$ line, we conclude that grid resolution has no significant impact on pathways, even though there is a little bit of scatter. Other regions show similar results (not shown). Note that the resolutions used in this paper, which are typical for climate models with interactive aerosol, are not able to resolve typical black carbon plumes as measured by the HIPPO campaigns (Weigum et al., 2012).

Figure 25 shows regional fractional tendencies of all processes for both masses and numbers for either pre-industrial (PI) or present-day (PD) emission scenarios. Again, we show results for the East-Asia region (see Fig. 1) where large differences in emission exist. Consequently, looking at the left panels for mass and number gains, we see that primarily emissions (grey shades) and ageing through sulfate condensation (brown) have changed. In particular, the emission of organic matter and black carbon into the hydrophobic Aitken changes significantly in its importance for numbers gain. Looking at the right panels for mass and number losses, we see that mostly wet deposition (light grey) of the hydrophobic modes and ageing through sulfate con-

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



denensation (brown) have changed. Under present day conditions, ageing due to sulfate condensation of hydrophobic modes has become more important and this is balanced by a decreased importance of deposition as a loss process of the same hydrophobic modes. It's not that the deposition processes become less effective but rather ageing speeds up, resulting in the shorter life-time of the hydrophobic modes that we already discussed in Sect. 4.2.

Regardless of these changes due to emissions the overall hierarchy of processes remains unaltered. In particular, most data points in Fig. 25 remain within the band $y = x \pm 0.1$ showing there are no substantial changes in pathways. That is not to say that the absolute tendencies hardly change. In fact, PI tendencies for e.g. sulfate condensation are substantially reduced from PD tendencies, but as the aerosol system adjusts, fractional tendencies change much less. This is shown in Fig. 26, where ratios of PI to PD tendencies are plotted against ratios of PI to PD fractional tendencies, again for East Asia. We see, in the top left panel, that the absolute tendencies of sulfate condensation (light purple) on the nucleation mode or sulfate production (dark purple) for the hydrophilic accumulation mode change substantially (3–10 ×) but their fractional tendencies change by no more than 20%. Similarly, in the bottom left and top right panel, the absolute tendencies for redistribution (red) are changed by a factor 2–4 ×, while their fractional tendencies change with less than 10%. The obvious exception to this are the tendencies for ageing due to sulfate condensation (brown) in all panels, that show substantial changes in both absolute and fractional tendencies. Similar conclusions can be drawn for other regions (not shown).

5 Conclusions

We present a pathway analysis of aerosol processes in the Earth's atmosphere. This pathway analysis is a budget of mass and number tendencies due to primary emissions, nucleation, H₂SO₄ condensation and in-cloud production, coagulation, ageing and wet and dry deposition as well as sedimentation. This budget is constructed

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from monthly averaged model tendencies due to these processes as modelled by ECHAM5.5-HAM2. To our knowledge, this is the first time such a detailed budget for global aerosol processes has been constructed.

The model ECHAM-HAM represents the aerosol species sulfate, organic matter, black carbon, sea salt and dust with 7 distinct modes of four size classes (nucleation, Aitken, accumulation and coarse) and two hygroscopic classes (hydrophobic and hydrophilic). Each mode is presented by a number of tracers: one for its number density and from one to five for the mass densities of its constituent species. Aerosol in ECHAM-HAM can thus be both externally and internally mixed. The M7 microphysics module in ECHAM-HAM is also implemented in regional aerosol models like COSMO and HIRLAM, and is rather similar to the GLOMAP module in the global aerosol model HadGEM-UKCA.

The pathway analysis gives an unprecedented view of how global aerosol is processed. First, aerosol mass and numbers are influenced by quite different processes. A striking example is the hydrophilic accumulation mode. Over most of the globe, its aerosol mass results from condensation and in-cloud production of H_2SO_4 or primary emission of sea-salt, while its numbers are determined by growth of hydrophilic Aitken particles. Next, our analysis shows there is a rather stable hierarchy of importance among the processes, with some processes contributing significantly and others hardly to the aerosol evolution. Among the significant processes, the dominant process differs substantially with region, altitude and season. Our analysis maps out these variations. The coarse modes are mostly governed by primary emissions of seasalt and dust. The evolution of these modes is very distinct and almost entirely decoupled from that of the smaller nucleation, Aitken and accumulation modes. The latter modes are also influenced by primary emissions (of sulfate and carbons) but H_2SO_4 condensation, in-cloud production of H_2SO_4 and coagulation often play more substantial roles. Coagulation is only important among the smaller and more numerous modes, where it serves as a loss term (coagulation barely adds mass to the larger modes). Particle growth is shown to be an important process for introducing large numbers of parti-

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

cles into the Aitken and accumulation mode. The analysis also shows how the ageing timescale of hydrophobic organic matter and black carbon or hydrophobic dust differs by an order of magnitude across the globe, even when only considering major source regions. Finally, we present species compositions of each mode resulting from the
5
aforementioned processes. It shows in particular the dominant contribution of sulfate to hydrophilic accumulation mode aerosol on the Northern Hemisphere.

Interestingly, the hierarchy of importance for the aerosol processes is relatively unaffected by what are normally considered major changes in numerical experimental setup. We have conducted experiments with coarser grid resolutions or pre-industrial emissions that lead to changes in the overall simulation (e.g. see Table 3), while leaving the hierarchy intact. Whenever we saw a change, it usually concerned two major
10
processes that exchanged their relative importance. Most notably, when using pre-industrial emissions, ageing of hydrophobic aerosols slows down and wet deposition becomes a more important pathway for removal of hydrophobic aerosol. Changes in
15
grid resolution (from T63L31 to T31L19) have less effect on process hierarchy than changing the emissions.

A pathway analysis may be used in several ways. First of all, it provides a detailed narrative of how aerosol is processed. Secondly, it offers ideas for the reduction of complexity in aerosol models. Current state-of-the-art aerosol modules require a significant
20
part of the overall resources required for global (climate) simulations. Thirdly, a pathway analysis helps to prioritise model improvement as a clear hierarchy in importance of the processes was found. Even though the processes are modelled with various degrees of abstraction, the magnitude of their effect should be reasonable. Fourthly, by identifying where and when what processes dominate, a process-based evaluation
25
of a model against relevant observations should become easier. Finally, and related to the last two points, a pathway analysis offers suggestions as to what kind of structural error may affect aerosol simulations.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Future work will include a significant reduction of complexity of M7, the aerosol microphysics routine in ECHAM-HAM. Currently we are extending our pathway analysis to other models to explore the robustness of our findings.

Acknowledgements. This work was supported by the National Environmental Research Council grants nr NE/G006148/1 (project AEROS) and NE/J024252/1 (project GASSP). Computational resources were made available by Deutsches Klimarechenzentrum (DKRZ) through support from the Bundesministerium für Bildung und Forschung (BMBF). The first author likes to thank Sebastian Rast (DKRZ) for his kind help in running ECHAM-HAM on Blizzard. The figures in this paper were prepared using David W. Fanning's coyote library for IDL.

References

- Aan de Brugh, J. M. J., Schaap, M., Vignati, E., Dentener, F., Kahnert, M., Sofiev, M., Huijnen, V., and Krol, M. C.: The European aerosol budget in 2006, *Atmos. Chem. Phys.*, 11, 1117–1139, doi:10.5194/acp-11-1117-2011, 2011. 15047
- Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227–1230, 1989. 15047
- Angstrom, B. A.: Atmospheric turbidity, global illumination and planetary albedo of the earth, *Tellus*, XIV, 435–450, 1962. 15047
- Bauer, S. E., Wright, D. L., Koch, D., Lewis, E. R., McGraw, R., Chang, L.-S., Schwartz, S. E., and Ruedy, R.: MATRIX (Multiconfiguration Aerosol TRacker of mIXing state): an aerosol microphysical module for global atmospheric models, *Atmos. Chem. Phys.*, 8, 6003–6035, doi:10.5194/acp-8-6003-2008, 2008. 15047
- Beelen, R., Raaschou-Nielsen, O., Stafoggia, M., Andersen, Z. J., Weinmayr, G., Hoffmann, B., Wolf, K., Samoli, E., Fischer, P., Nieuwenhuijsen, M., Vineis, P., Xun, W. W., Katsouyanni, K., Dimakopoulou, K., Oudin, A., Forsberg, B., Modig, L., Havulinna, A. S., Lanki, T., Turunen, A., Oftedal, B., Nystad, W., Nafstad, P., De Faire, U., Pedersen, N. L., Östenson, C.-G., Fratiglioni, L., Penell, J., Korek, M., Pershagen, G., Eriksen, K. T., Overvad, K., Ellermann, T., Eeftens, M., Peeters, P. H., Meliefste, K., Wang, M., Bueno-de Mesquita, B., Sugiri, D., Krämer, U., Heinrich, J., de Hoogh, K., Key, T., Peters, A., Hampel, R., Concin, H., Nagel, G., Ineichen, A., Schaffner, E., Probst-Hensch, N., Künzli, N., Schindler, C., Schikowski, T.,

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Adam, M., Phuleria, H., Vilier, A., Clavel-Chapelon, F., Declercq, C., Grioni, S., Krogh, V., Tsai, M.-Y., Ricceri, F., Sacerdote, C., Galassi, C., Migliore, E., Ranzi, A., Cesaroni, G., Badaloni, C., Forastiere, F., Tamayo, I., Amiano, P.: Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the multicentre ESCAPE project, *Lancet*, 6736, 1–11, doi:10.1016/S0140-6736(13)62158-3, 2013. 15047
- 5 Brunekreef, B. and Holgate, S. T.: Air pollution and health, *Lancet*, 360, 1233–1242, doi:10.1016/S0140-6736(02)11274-8, 2002. 15047
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric aerosol optical thickness from the GOCART model and comparisons with satellite and sun photometer measurements, *J. Atmos. Sci.*, 59, 461–483, 2002. 15047
- 10 Dockery, D., Pope, A., Xu, X., Spengler, J., Ware, J., Fay, M., Ferris, B., and Speizer, F.: An association between air pollution and mortality in six U.S. cities, *New Engl. J. Med.*, 329, 1753–1759, 1993. 15047
- 15 Easter, R. C., Ghan, S. J., Zhang, Y., Saylor, R. D., Chapman, E. G., Laulainen, N. S., Abdul-razzak, H., Leung, L. R., Bian, X., and Zaveri, R. A.: MIRAGE: model description and evaluation of aerosols and trace gases, *J. Geophys. Res.*, 109, D20210, doi:10.1029/2004JD004571, 2004. 15047
- Ezzati, M., Lopez, A. D., Rodgers, A., Vander Hoorn, S., and Murray, C. J. L.: Selected major risk factors and global and regional burden of disease, *Lancet*, 360, 1347–1360, doi:10.1016/S0140-6736(02)11403-6, 2002. 15047
- 20 Feichter, J., Kjellstrom, E., Rodhe, H., Dentener, F., Lelieveld, J., and Roelofs, G.-J.: Simulation of the tropospheric sulfur cycle in a global climate model, *Atmos. Environ.*, 30, 1693–1707, 1996. 15047, 15051, 15055
- 25 Folini, D. and Wild, M.: Aerosol emissions and dimming/brightening in Europe: sensitivity studies with ECHAM5-HAM, *J. Geophys. Res.*, 116, D21104, doi:10.1029/2011JD016227, 2011. 15050
- Fuchs, N. A.: *The Mechanics of Aerosols*, Pergamon Press, Oxford, 1964. 15054, 15055
- Ghan, S. J. and Schwartz, S. E.: Aerosol properties and processes: a path from field and laboratory measurements to global climate models, *B. Am. Meteorol. Soc.*, 88, 1059–1083, doi:10.1175/BAMS-88-7-1059, 2007. 15047
- 30 Ghan, S. J., Liu, X., Easter, R. C., Zaveri, R., Rasch, P. J., Yoon, J.-H., and Eaton, B.: Toward a minimal representation of aerosols in climate models: comparative decomposi-

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion of aerosol direct, semidirect, and indirect radiative forcing, *J. Climate*, 25, 6461–6476, doi:10.1175/JCLI-D-11-00650.1, 2012. 15048

Guelle, W., Schulz, M., Balkanski, Y., and Dentener, F.: Influence of the source formulation on modeling the atmospheric global distribution of sea salt aerosol, *J. Geophys. Res.*, 106, 27509, doi:10.1029/2001JD900249, 2001. 15052

Hansen, J., Sato, M., and Ruedy, R.: Radiative forcing and climate response, *J. Geophys. Res.*, 102, 6831–6864, 1997. 15047

Hoose, C., Lohmann, U., Bennartz, R., Croft, B., and Lesins, G.: Global simulations of aerosol processing in clouds, *Atmos. Chem. Phys.*, 8, 6939–6963, doi:10.5194/acp-8-6939-2008, 2008. 15053

Horowitz, L. W., Walters, S., Mauzerall, D. L., Emmons, L. K., Rasch, P. J., Granier, C., Tie, X., Lamarque, J.-F., Schultz, M. G., Tyndall, G. S., Orlando, J. J., and Brasseur, G. P.: A global simulation of tropospheric ozone and related tracers: description and evaluation of MOZART, version 2, *J. Geophys. Res.*, 108, 4784, doi:10.1029/2002JD002853, 2003. 15051

Kazil, J. and Lovejoy, E. R.: A semi-analytical method for calculating rates of new sulfate aerosol formation from the gas phase, *Atmos. Chem. Phys.*, 7, 3447–3459, doi:10.5194/acp-7-3447-2007, 2007. 15054

Kazil, J., Stier, P., Zhang, K., Quaas, J., Kinne, S., O'Donnell, D., Rast, S., Esch, M., Ferrachat, S., Lohmann, U., and Feichter, J.: Aerosol nucleation and its role for clouds and Earth's radiative forcing in the aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 10, 10733–10752, doi:10.5194/acp-10-10733-2010, 2010. 15050, 15054

Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Bernsten, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment – optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, doi:10.5194/acp-6-1815-2006, 2006. 15047

Kloster, S., Dentener, F., Feichter, J., Raes, F., van Aardenne, J., Roeckner, E., Lohmann, U., Stier, P., and Swart, R.: Influence of future air pollution mitigation strategies on total aerosol radiative forcing, *Atmos. Chem. Phys.*, 8, 6405–6437, doi:10.5194/acp-8-6405-2008, 2008. 15050

A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Koffi, B., Schulz, M., Bréon, F.-M., Griesfeller, J., Winker, D., Balkanski, Y., Bauer, S., Berntsen, T., Chin, M., Collins, W. D., Dentener, F., Diehl, T., Easter, R., Ghan, S., Ginoux, P., Gong, S., Horowitz, L. W., Iversen, T., Kirkevåg, A., Koch, D., Krol, M., Myhre, G., Stier, P., and Takemura, T.: Application of the CALIOP layer product to evaluate the vertical distribution of aerosols estimated by global models: AeroCom phase I results, *J. Geophys. Res.*, 117, D10201, doi:10.1029/2011JD016858, 2012. 15048
- Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Cluster activation theory as an explanation of the linear dependence between formation rate of 3nm particles and sulphuric acid concentration, *Atmos. Chem. Phys.*, 6, 787–793, doi:10.5194/acp-6-787-2006, 2006. 15054
- Langner, J. and Rodhe, H.: A global three-dimensional model of the tropospheric sulfur cycle, *J. Atmos. Chem.*, 13, 225–263, 1991. 15047
- Lee, L. A., Carslaw, K. S., Pringle, K. J., and Mann, G. W.: Mapping the uncertainty in global CCN using emulation, *Atmos. Chem. Phys.*, 12, 9739–9751, doi:10.5194/acp-12-9739-2012, 2012. 15048
- Lee, L. A., Pringle, K. J., Reddington, C. L., Mann, G. W., Stier, P., Spracklen, D. V., Pierce, J. R., and Carslaw, K. S.: The magnitude and causes of uncertainty in global model simulations of cloud condensation nuclei, *Atmos. Chem. Phys.*, 13, 8879–8914, doi:10.5194/acp-13-8879-2013, 2013. 15048
- Lequy, E., Conil, S., and Turpault, M.-P.: Impacts of Aeolian dust deposition on European forest sustainability: a review, *Forest Ecol. Manag.*, 267, 240–252, doi:10.1016/j.foreco.2011.12.005, 2012. 15047
- Liao, H., Seinfeld, J. H., Adams, P. J., and Mickley, L. J.: Global radiative forcing of coupled tropospheric ozone and aerosols in a unified general circulation model, *J. Geophys. Res.*, 109, D16207, doi:10.1029/2003JD004456, 2004. 15047
- Lin, S.-J. and Rood, R. B.: Multidimensional flux-form semi-Lagrangian transport schemes, *Mon. Weather Rev.*, 124, 2046–2070, 1996. 15050
- Liu, X., Penner, J. E., and Herzog, M.: Global modeling of aerosol dynamics: model description, evaluation, and interactions between sulfate and nonsulfate aerosols, *J. Geophys. Res.*, 110, D18206, doi:10.1029/2004JD005674, 2005. 15047
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*, 5, 709–739, doi:10.5194/gmd-5-709-2012, 2012. 15048

Lohmann, U. and Feichter, J.: Impact of sulfate aerosols on albedo and lifetime of clouds: a sensitivity study with the ECHAM4 GCM, *J. Geophys. Res.*, 102, 13685–13700, 1997. 15047

Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715–737, doi:10.5194/acp-5-715-2005, 2005. 15047

Lohmann, U., Salzen, K. V., Mcfarlane, N., and Leighton, H. G.: Tropospheric sulfur cycle in the Canadian general circulation model, *J. Geophys. Res.*, 104, 833–858, 1999. 15047

Lohmann, U., Stier, P., Hoose, C., Ferrachat, S., Kloster, S., Roeckner, E., and Zhang, J.: Cloud microphysics and aerosol indirect effects in the global climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 7, 3425–3446, doi:10.5194/acp-7-3425-2007, 2007. 15050

Maher, B., Prospero, J., Mackie, D., Gaiero, D., Hesse, P., and Balkanski, Y.: Global connections between aeolian dust, climate and ocean biogeochemistry at the present day and at the last glacial maximum, *Earth-Sci. Rev.*, 99, 61–97, doi:10.1016/j.earscirev.2009.12.001, 2010. 15047

Makkonen, R., Asmi, A., Korhonen, H., Kokkola, H., Järvenoja, S., Räisänen, P., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., Järvinen, H., Lohmann, U., Bennartz, R., Feichter, J., and Kulmala, M.: Sensitivity of aerosol concentrations and cloud properties to nucleation and secondary organic distribution in ECHAM5-HAM global circulation model, *Atmos. Chem. Phys.*, 9, 1747–1766, doi:10.5194/acp-9-1747-2009, 2009. 15050

McTainsh, G. and Strong, C.: The role of aeolian dust in ecosystems, *Geomorphology*, 89, 39–54, doi:10.1016/j.geomorph.2006.07.028, 2007. 15047

Monahan, E., Spiel, D., and Davidson, K. L.: A Model of marine aerosol generation via whitecaps and wave disruption, in: *Oceanic Whitecaps*, edited by: Monahan, E. and Niocaill, G. M., D. Reidel, the Netherlands, 167–174, 1986. 15052

Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase

A budget for aerosol processesN. A. J. Schutgens and
P. Stier

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

II simulations, *Atmos. Chem. Phys.*, 13, 1853–1877, doi:10.5194/acp-13-1853-2013, 2013. 15048

Niemeier, U., Timmreck, C., Graf, H.-F., Kinne, S., Rast, S., and Self, S.: Initial fate of fine ash and sulfur from large volcanic eruptions, *Atmos. Chem. Phys.*, 9, 9043–9057, doi:10.5194/acp-9-9043-2009, 2009. 15050

O'Donnell, D., Tsigaridis, K., and Feichter, J.: Estimating the direct and indirect effects of secondary organic aerosols using ECHAM5-HAM, *Atmos. Chem. Phys.*, 11, 8635–8659, doi:10.5194/acp-11-8635-2011, 2011. 15050

Peters, K., Stier, P., Quaas, J., and Graßl, H.: Aerosol indirect effects from shipping emissions: sensitivity studies with the global aerosol-climate model ECHAM-HAM, *Atmos. Chem. Phys.*, 12, 5985–6007, doi:10.5194/acp-12-5985-2012, 2012. 15050

Rasch, P. J., Barth, M. C., Kiehl, J. T., Schwartz, E., Benkovitz, C. M., and America, N.: A description of the global sulfur cycle and its controlling processes in the National Center for Atmospheric Research Community Climate Model, Version 3, *J. Geophys. Res.*, 105, 1367–1385, 2000. 15047

Riipinen, I., Sihto, S.-L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinilä, K., Kerminen, V.-M., Laaksonen, A., and Lehtinen, K. E. J.: Connections between atmospheric sulphuric acid and new particle formation during QUEST III–IV campaigns in Heidelberg and Hyytiälä, *Atmos. Chem. Phys.*, 7, 1899–1914, doi:10.5194/acp-7-1899-2007, 2007. 15054

Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblüeh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The Atmospheric General Circulation Model ECHAM5, Part I: Model Description, *Tech. Rep. 349*, Max Planck Institute for Meteorology, Hamburg, 2003. 15050

Roeckner, E., Rokoř, R. B., Sch, M. E., Iorgetta, M. G., and Aagemann, S. H.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Climate*, 19, 3771–3791, 2006. 15050

Roelofs, G.-J., Lelieveld, J., and Ganzeveld, L.: Simulation of global sulfate distribution and the influence on effective cloud drop radii with a coupled photochemistry sulfur cycle model, *Tellus B*, 50, 224–242, doi:10.1034/j.1600-0889.1998.t01-2-00002.x, 1998. 15047

Roelofs, G. J., Stier, P., Feichter, J., Vignati, E., and Wilson, J.: Aerosol activation and cloud processing in the global aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 6, 2389–2399, doi:10.5194/acp-6-2389-2006, 2006. 15050

A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- 5 Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty, *Atmos. Chem. Phys.*, 13, 2423–2434, doi:10.5194/acp-13-2423-2013, 2013. 15048
- 10 Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006. 15047
- 15 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons, Hoboken, New Jersey, 2006. 15053, 15054, 15055, 15056, 15060
- Seland, O. Y., Iversen, T., Kirkevåg, A., and Storelvmo, T.: Aerosol-climate interactions in the CAM-Oslo atmospheric GCM and investigation of associated basic shortcomings, *Tellus A*, 60, 459–491, doi:10.1111/j.1600-0870.2008.00318.x, 2008. 15047
- 20 Smith, K. R., Jerrett, M., Anderson, H. R., Burnett, R. T., Stone, V., Derwent, R., Atkinson, R. W., Cohen, A., Shonkoff, S. B., Krewski, D., Pope, C. A., Thun, M. J., and Thurston, G.: Public health benefits of strategies to reduce greenhouse-gas emissions: health implications of short-lived greenhouse pollutants, *Lancet*, 374, 2091–103, doi:10.1016/S0140-6736(09)61716-5, 2009. 15047
- 25 Smith, M. H. and Harrison, N. M.: The sea spray generation function, *J. Aerosol Sci.*, 29, S189–S190, 1998. 15052
- Spracklen, D. V., Pringle, K. J., Carslaw, K. S., Chipperfield, M. P., and Mann, G. W.: A global off-line model of size-resolved aerosol microphysics: I. Model development and prediction of aerosol properties, *Atmos. Chem. Phys.*, 5, 2227–2252, doi:10.5194/acp-5-2227-2005, 2005. 15047
- 30 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, doi:10.5194/acp-5-1125-2005, 2005. 15047, 15049, 15050, 15051, 15052, 15053

A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stier, P., Feichter, J., Kloster, S., Vignati, E., and Wilson, J.: Emission-induced nonlinearities in the global aerosol system: results from the ECHAM5-HAM aerosol-climate model, *J. Climate*, 19, 3845–3862, 2006a. 15050, 15062

Stier, P., Feichter, J., Roeckner, E., Kloster, S., and Esch, M.: The evolution of the global aerosol system in a transient climate simulation from 1860 to 2100, *Atmos. Chem. Phys.*, 6, 3059–3076, doi:10.5194/acp-6-3059-2006, 2006b. 15050

Stier, P., Seinfeld, J. H., Kinne, S., and Boucher, O.: Aerosol absorption and radiative forcing, *Atmos. Chem. Phys.*, 7, 5237–5261, doi:10.5194/acp-7-5237-2007, 2007. 15050

Stier, P., Schutgens, N. A. J., Bellouin, N., Bian, H., Boucher, O., Chin, M., Ghan, S., Huneus, N., Kinne, S., Lin, G., Ma, X., Myhre, G., Penner, J. E., Randles, C. A., Samset, B., Schulz, M., Takemura, T., Yu, F., Yu, H., and Zhou, C.: Host model uncertainties in aerosol radiative forcing estimates: results from the AeroCom Prescribed intercomparison study, *Atmos. Chem. Phys.*, 13, 3245–3270, doi:10.5194/acp-13-3245-2013, 2013. 15048

Takemura, T., Okamoto, H., Maruyama, Y., Numaguti, A., Higurashi, A., and Nakajima, T.: Global three-dimensional simulation of aerosol optical thickness distribution of various origins, *J. Geophys. Res.*, 105, 17853, doi:10.1029/2000JD900265, 2000. 15047

Tegen, I.: Impact of vegetation and preferential source areas on global dust aerosol: results from a model study, *J. Geophys. Res.*, 107, 4576, doi:10.1029/2001JD000963, 2002. 15052

Tegen, I. and Lacis, A.: Modeling of particle size distribution and its influence on the radiative properties of mineral dust aerosol, *J. Geophys. Res.*, 101, 19237–19244, 1996. 15047

Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, 6, 1777–1813, doi:10.5194/acp-6-1777-2006, 2006. 15048, 15061

Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Feichter, J., Fillmore, D., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I. S. A., Iversen, T., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, M. S., Seland, Ø., Stier, P., Takemura, T.,

A budget for aerosol processesN. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Tie, X.: The effect of harmonized emissions on aerosol properties in global models – an AeroCom experiment, *Atmos. Chem. Phys.*, 7, 4489–4501, doi:10.5194/acp-7-4489-2007, 2007. 15048

5 Timmreck, C., Graf, H.-F., Lorenz, S. J., Niemeier, U., Zanchettin, D., Matei, D., Jungclaus, J. H., and Crowley, T. J.: Aerosol size confines climate response to volcanic super-eruptions, *Geophys. Res. Lett.*, 37, L24705, doi:10.1029/2010GL045464, 2010. 15050

Twomey, S.: Pollution and the planetary albedo, *Atmos. Environ.*, 8, 1251–1256, 1974. 15047

10 Vignati, E., Wilson, J., and Stier, P.: M7: an efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res.*, 109, D22202, doi:10.1029/2003JD004485, 2004. 15049, 15051, 15056

Vink, S. and Measures, C.: The role of dust deposition in determining surface water distributions of Al and Fe in the South West Atlantic, *Deep-Sea Res. Pt. II*, 48, 2787–2809, doi:10.1016/S0967-0645(01)00018-2, 2001. 15047

15 Weigum, N. M., Stier, P., Schwarz, J. P., Fahey, D. W., and Spackman, J. R.: Scales of variability of black carbon plumes over the Pacific Ocean, *Geophys. Res. Lett.*, 39, L15804, doi:10.1029/2012GL052127, 2012. 15071

Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691–7710, doi:10.5194/acp-9-7691-2009, 2009. 15047

20 Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J., Wan, H., Rast, S., and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity to improvements in process representations, *Atmos. Chem. Phys.*, 12, 8911–8949, doi:10.5194/acp-12-8911-2012, 2012. 15049, 15051, 15052, 15053, 15059

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Table 1. ECHAM-HAM aerosol mode structure. Mode radii are constrained to the given size ranges. Identification numbers label the modes and are used in figures throughout the paper. Species composition differs by mode. Only the hydrophilic modes experience wet growth due to humidity.

Size class	[μm]	Hydrophilic ID	Hydrophilic species	Hydrophobic ID	Hydrophobic Species
nucleation	$r \leq 0.005$	1	SO ₄		
Aitken	$0.005 \leq r \leq 0.05$	2	SO ₄ , POM, BC	5	POM, BC
accumulation	$0.05 \leq r \leq 0.5$	3	SO ₄ , POM, BC, SS, DU	6	DU
coarse	$r \geq 0.5$	4	SO ₄ , POM, BC, SS, DU	7	DU

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 2. Aerosol processes in ECHAM-HAM. The columns present from left to right: the process; the identification code used in various figures; the exact pathway, using numbers to identify mode and letters to identify number (n) or mass (m) tendencies; the species involved. In the right two columns, specific ageing processes are detailed. In ECHAM-HAM they are the instantaneous result of some process mentioned on the left.

Process	Code	Pathway	Species	aging		
				Code	Pathway	Species
Emission	SO4	→ 2n,m	SO ₄			
Emission	SO4	→ 3n,m	SO ₄			
Emission	SO4	→ 4n,m	SO ₄			
Emission	POM	→ 2n,m	POM			
Emission	POM	→ 3m	POM			
Emission	POM	→ 5n,m	POM			
Emission	SS	→ 3n,m	SS			
Emission	SS	→ 4n,m	SS			
Emission	BC	→ 5n,m	BC			
Emission	DU	→ 6n,m	DU			
Emission	DU	→ 7n,m	DU			
Nucleation	NCL	→ 1n,m	SO ₄			
Condensation	CND	→ 1m	SO ₄			
Condensation	CND	→ 2m	SO ₄			
Condensation	CND	→ 3m	SO ₄			
Condensation	CND	→ 4m	SO ₄			
Condensation	CND	→ 5	SO ₄			
Condensation	CND	→ 6	SO ₄			
Condensation	CND	→ 7	SO ₄			
In-cloud production	PRD	→ 3m	SO ₄			
In-cloud production	PRD	→ 4n,m	SO ₄			
Coagulation	11	1n →				
Coagulation	12	1n,m → 2m	SO ₄			
Coagulation	13	1n,m → 3m	SO ₄			
Coagulation	14	1n,m → 4m	SO ₄			
Coagulation	15	1n,m → 5	SO ₄			
Coagulation	16	1n,m → 6	SO ₄			
Coagulation	17	1n,m → 7	SO ₄			
Coagulation	22	2n →				
Coagulation	23	2n,m → 3m	SO ₄ , POM, BC			
Coagulation	24	2n,m → 4m	SO ₄ , POM, BC			
Coagulation	26	2n,m → 6	SO ₄ , POM, BC			
Coagulation	27	2n,m → 7	SO ₄ , POM, BC			
Coagulation	33	3n →				
Coagulation	55	5n →				
Coagulation	52	5n,m → 2m	POM, BC			
Coagulation	53	5n,m → 3m	POM, BC			
Redistribution	RDB	1n,m → 2n,m	SO ₄			
Redistribution	RDB	2n,m → 3n,m	SO ₄ , POM, BC			
Redistribution	RDB	3n,m → 4n,m	SO ₄ , POM, BC, DU, SS			
Wet deposition	WET	all →	SO ₄ , POM, BC, DU, SS			
Dry deposition	DRY	all →	SO ₄ , POM, BC, DU, SS			
Sedimentation	SED	all* →	SO ₄ , POM, BC, DU, SS			
				CAG 5n,m → 2n,m SO ₄ , POM, BC CAG 6n,m → 3n,m SO ₄ , DU CAG 7n,m → 4n,m SO ₄ , DU 152 5n,m → 2n,m SO ₄ , POM, BC 163 6n,m → 3n,m SO ₄ , DU 174 7n,m → 4n,m SO ₄ , DU 263 6n,m → 3n,m SO ₄ , POM, BC, DU 274 7n,m → 4n,m SO ₄ , POM, BC, DU		
				effectively an ageing process		
				effectively an ageing process		

*: all, except nucleation mode

A budget for aerosol processes

N. A. J. Schutgens and P. Stier

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




A budget for aerosol processes

 N. A. J. Schutgens and
P. Stier

Table 3. Life-times [days] of all modes and species in ECHAM5.5-HAM2.

Mode	Species	base-line		T31L19		pre-industrial	
		masses	numbers	masses	numbers	masses	numbers
	■ SO ₄	4.6		5.5		5.3	
	■ POM	6.0		5.0		5.7	
	■ BC	6.1		6.0		7.8	
	■ SS	0.6		0.5		0.6	
	■ DU	4.8		4.2		5.0	
1: nucleation	SO ₄	0.5	0.2	0.4	0.2	0.6	0.2
2: Aitken (hydrophilic)		1.0	4.7	1.1	4.9	1.8	5.6
	SO ₄	3.5		3.3		4.4	
	POM	0.2		0.2		0.4	
	BC	0.4		0.5		0.8	
3: accumulation (hydrophilic)		3.9	6.2	4.0	6.7	3.5	6.8
	SO ₄	4.4		5.1		4.9	
	POM	6.0		5.1		5.4	
	BC	6.0		6.0		7.4	
	SS	1.7		1.3		1.7	
	DU	7.6		6.7		7.8	
4: coarse (hydrophilic)		0.9	3.4	0.8	3.1	0.8	3.2
	SO ₄	2.7		4.0		2.8	
	POM	2.0		2.1		1.9	
	BC	1.9		2.4		2.9	
	SS	0.6		0.5		0.6	
	DU	5.1		5.0		5.2	
5: Aitken (hydrophobic)		0.8	0.7	0.7	0.5	1.6	1.5
	POM	0.9		0.7		1.6	
	BC	0.7		0.6		1.5	
6: accumulation (hydrophobic)	DU	2.2	2.2	1.7	1.8	3.3	3.4
7: coarse (hydrophobic)	DU	2.3	3.3	1.7	2.7	3.0	4.6

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A budget for aerosol processesN. A. J. Schutgens and
P. Stier**Figure 1.** Definition of the regions that will be used for part of the analysis in this paper.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 2. Contributions by different modes to AOT and CCN for the baseline experiment. The pie-chart colours show contribution by mode (see legend below lowest panel), the pie-chart's size the overall magnitude (legend at the bottom of each panel). From top to bottom: AOT at 550 nm (linear scale); column-integrated CCN at $S = 1\%$ (logarithmic scale).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 3. Contributions by different modes to surface $\text{PM}_{2.5}$ and black carbon mass burden for the baseline experiment. The pie-chart colours show contribution by mode (see legend below lowest panel), the pie-chart's size the overall magnitude (legend at the bottom of each panel). From top to bottom: surface $\text{PM}_{2.5}$ (logarithmic scale); black carbon mass burden (logarithmic scale).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


 N. A. J. Schutgens and
P. Stier


Figure 4. Globally and annually averaged net mass and number tendencies (arrows) between the 7 modes (dots) in ECHAM5.5-HAM2. The vertical arrows at the top and bottom of the figure represent H₂SO₄ nucleation, condensation and production, aerosol primary emission and deposition processes. The arrows between the modes represent coagulation, ageing and redistribution processes. Pink arrows are number loss tendencies for coagulation processes. The width of the arrow scales logarithmically with the magnitude of the net tendency. In addition, numerical values for the annual and global averages are given.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A budget for aerosol processes


 N. A. J. Schutgens and
P. Stier


Figure 5. Global distribution of annual burdens (a) and tendencies (b–f) of the nucleation mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels a, c–f). Panel (b) shows a global average of the tendencies.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 6. Zonal distribution of annual densities (**a**) and tendencies (**b** and **c**) of the nucleation mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates the zonally-averaged density or tendency (logarithmic scale as shown in black at the top of **a**, **b** and **c**).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier




Figure 7. Seasonal cycle of regional tendencies of the nucleation mode over East-Asia (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 8. Global distribution of annual burdens **(a)** and tendencies **(b–f)** of the hydrophobic Aitken mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels **a**, **c–f**). Panel **(b)** shows a global average of the tendencies.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 9. Zonal distribution of annual densities (a) and tendencies (b) of the hydrophobic Aitken mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates the zonally-averaged density or tendency (logarithmic scale as shown in black at the top of the panels a and b).

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier




Figure 10. Ageing time-scales of the hydrophobic Aitken mode. This time-scale is defined as $N/(\partial N/\partial t)$, with N the number density and $\partial N/\partial t$ its time-derivative *due to ageing processes only*. The colours indicate processes (Table 2), the radius of the pie indicates ageing time-scale (inverse linear scale).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 11. Global distribution of annual burdens **(a)** and tendencies **(b–f)** of the hydrophilic Aitken mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels **a**, **c–f**). Panel **(b)** shows a global average of the tendencies.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and P. Stier




Figure 12. Seasonal cycle of regional tendencies of the hydrophilic Aitken mode over the Savanna (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and P. Stier




Figure 13. Seasonal cycle of regional tendencies of the hydrophilic Aitken mode over East-Asia (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 14. Global distribution of annual burdens **(a)** and tendencies **(b–f)** of the hydrophobic accumulation mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels **a**, **c–f**). Panel **(b)** shows a global average of the tendencies.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 15. Global distribution of annual burdens (a) and tendencies (b–f) of the hydrophilic accumulation mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels a, c–f). Panel (b) shows a global average of the tendencies.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier




Figure 16. Zonal distribution of annual densities of the hydrophilic accumulation mode. The colours indicate species (defined in Table 3). The pie-chart shows fractional contributions, while the radius of the pie indicates the zonally-averaged density (logarithmic scale as shown in black at the top).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A budget for aerosol processes

N. A. J. Schutgens and P. Stier




Figure 17. Seasonal cycle of regional tendencies of the hydrophilic accumulation mode over the Savanna (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and P. Stier




Figure 18. Seasonal cycle of regional tendencies of the hydrophilic accumulation mode over East-Asia (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier




Figure 19. Seasonal cycle of regional tendencies of the hydrophilic accumulation mode over the southern ocean (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes


N. A. J. Schutgens and
P. Stier

Figure 20. Global distribution of annual burdens (a) and tendencies (b–f) of the hydrophobic coarse mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels a, c–f). Panel (b) shows a global average of the tendencies.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 21. Ageing time-scales of the hydrophobic coarse mode. This time-scale is defined as $N/(\partial N/\partial t)$, with N the number density and $\partial N/\partial t$ its time-derivative *due to ageing processes only*. The colours indicate processes (Table 2), the radius of the pie indicates ageing time-scale (inverse linear scale).

A budget for aerosol processes

N. A. J. Schutgens and P. Stier




Figure 22. Global distribution of annual burdens (a) and tendencies (b–f) of the hydrophilic coarse mode. The colours indicate either species (defined in Table 3) or processes (defined in Table 2). The pie-chart shows fractional contributions, while the radius of the pie indicates total column burden or total column tendency (logarithmic scale as shown in black at the bottom of panels a, c–f). Panel (b) shows a global average of the tendencies.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



A budget for aerosol processes

N. A. J. Schutgens and P. Stier




Figure 23. Seasonal cycle of regional tendencies of the hydrophilic coarse mode over East-Asia (see Fig. 1). The coloured bars show fractional tendencies (left axis) for the processes defined in Table 2. The solid line (right axis) shows the total tendency.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A budget for aerosol processes

N. A. J. Schutgens and
P. Stier

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 24. Fractional tendencies for the T31L19 low-resolution experiment vs. the baseline T63L31 high-resolution experiment. Colours indicate processes (Table 2), numbers indicate modes (Table 3). Results are shown for East Asia.

A budget for aerosol processes

N. A. J. Schutgens and
P. Stier




Figure 25. Fractional tendencies for the Pre-Industrial experiment vs. the Present Day experiment. Colours indicate processes (Table 2), numbers indicate modes (Table 3). Results are shown for East Asia.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion









Figure 26. Pre-industrial (PI) to Present Day (PD) ratios of absolute tendencies against PI to PD ratios of fractional tendencies. Colours indicate processes (Table 2), numbers indicate modes (Table 3). The size of the symbol indicates fractional tendency in the PD experiment, with the largest symbols having a fractional tendency of about 1 (linear scale). The slanted lines represent $y = 2^{-n}x$, with $n = 0, 1, 2, 3$. Results are shown for East Asia.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

