



**A case study over
China**

A.-M. Sundström et al.

On the use of satellite remote sensing based approach for determining aerosol direct radiative effect over land: a case study over China

A.-M. Sundström¹, A. Arola², P. Kolmonen³, Y. Xue⁴, G. de Leeuw^{1,3}, and M. Kulmala¹

¹Department of Physics, University of Helsinki, Helsinki, Finland

²Finnish Meteorological Institute, Kuopio, Finland

³Finnish Meteorological Institute, Helsinki, Finland

⁴Institute of Remote Sensing and Digital Earth, Chinese Academy of Sciences, Beijing, China

Received: 28 March 2014 – Accepted: 21 May 2014 – Published: 10 June 2014

Correspondence to: A.-M. Sundström (anu-maija.sundstrom@helsinki.fi)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

A satellite-based approach to derive the aerosol direct short wave (SW) radiative effect (ADRE) was studied in an environment with highly variable aerosol conditions over Eastern China from March to October 2009. The method is based on using coincident SW Top of the Atmosphere (TOA) fluxes from the Clouds and the Earth's Radiant Energy System (CERES) and aerosol optical depths (AODs) from the Moderate Resolution Imaging Spectroradiometer (MODIS). The estimate for instantaneous clear sky ADRE is obtained by establishing linear regression between CERES fluxes and MODIS AODs. Even though the approach has been used in a number of studies, less focus has been paid to the method itself. In this study the main goals were first to study the method in more detail as well as its applicability over Eastern China, and second to derive a satellite-based estimate of ADRE over the study area. Before the linear fitting, CERES fluxes were normalized to a fixed solar zenith angle, Earth–Sun distance and atmospheric water vapour content to reduce the noise in the flux observations that was not related to aerosols. The satellite based clear sky estimates for median instantaneous and diurnally averaged ADRE over the study area were -8.8 W m^{-2} , and -5.1 W m^{-2} , respectively. Over heavily industrialized areas the cooling at TOA was two to more than three times the median value, and associated with high AODs (> 0.5). Especially during the summer months positive ADREs were observed locally over dark surfaces. This was most probably a method artifact related to systematic change of aerosol type, subvisual cloud contamination or both. The key question in the satellite-based approach is the accuracy of the estimated aerosol-free TOA flux ($F_{0,\text{TOA}}$). Comparison with simulated $F_{0,\text{TOA}}$ showed that both the satellite method and the model produced qualitatively similar spatial patterns, but absolute values differed. In 58 % of the cases the satellite based $F_{0,\text{TOA}}$ was within $\pm 10 \text{ W m}^{-2}$ of the modeled value. Over bright surfaces the satellite-based method tend to produce lower $F_{0,\text{TOA}}$ than the model.

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Aerosols affect the Earth's climate directly by scattering and absorbing solar and infrared radiation, and indirectly by acting as a cloud condensation or ice nuclei and thus modifying the lifetime and radiative properties of clouds. The aerosol direct radiative effect (ADRE) describes the change of energy in the Earth's radiation field due to the scattering and absorption by aerosols. At the Top of the Atmosphere (TOA) ADRE is defined as the difference between the outgoing short wave (SW) solar flux without ($F_{0,TOA}$) and with aerosols present ($F_{aer,TOA}$) in the atmosphere. ADRE is considered as the combined radiative effect of anthropogenic and natural aerosols whereas the aerosol radiative forcing refers to the radiative effect of anthropogenic aerosols only and requires the separation between natural and anthropogenic aerosol components (e.g. Heald et al., 2013). Negative values of ADRE (or forcing) correspond to increased outgoing radiation and planetary cooling, whereas positive values correspond to decreased outgoing radiation at TOA and increased atmospheric warming. Several studies conclude that globally the clear sky ADRE and total forcing are negative (e.g. Haywood and Boucher, 2000; Jacobson, 2001a; Bellouin et al., 2005; Loeb and Manalo-Smith, 2005; Schulz et al., 2006; Quaas et al., 2008; Bellouin et al., 2008; Garcia et al., 2012; Myhre et al., 2013), but the estimates of the magnitude vary. This is mainly due to the large spatial and temporal variation of the aerosol concentration and mass as well as their relatively short lifetime in the atmosphere. Also, different methods and models used to estimate the radiative effect provide different results. The recent IPCC report (2013) summarizes the estimates of the global direct aerosol radiative effect at TOA for clear sky to range mainly from about -0.1 to -0.8 W m^{-2} , observation (satellite) based methods giving somewhat more negative estimates than the models. It is noted that locally the values of the direct aerosol radiative effect can differ significantly from the global estimates. Positive values of ADRE at TOA can be observed when, e.g., highly absorbing aerosols are transported over bright surfaces such as desert or snow. Also the direct radiative forcing of some anthropogenic components such as black carbon

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



have been estimated to be positive (e.g. Jacobson, 2001b; Schulz et al., 2006; Myhre et al., 2013).

During the past decade an increasing number of observation-based studies of ADRE and aerosol forcing have been carried out where remote sensing from space plays an important role. E.g. Yu et al. (2004); Bellouin et al. (2005) and Yu et al. (2006) showed global estimates of the aerosol radiative effect and forcing by combining remote sensing observations from Moderate Imaging Spectroradiometer (MODIS) and radiative transfer calculations, and more recently Thomas et al. (2013) used the Advanced Along Track Scanning Radiometer (AATSR) based global aerosol optical depth (AOD) dataset GlobAEROSOL with a radiative transfer model. In these studies the remote sensing observations have been used as an input to the radiative transfer model to define the TOA SW fluxes with aerosols. Another remote sensing-based approach to determine the SW ADRE at TOA is to use the AOD from satellite observations with coincident broadband flux observations from instrument such as the Clouds and the Earth's Radiant Energy System (CERES), which provides TOA fluxes in three broadband channels. The advantage is that the outgoing SW flux with aerosols at TOA is directly obtained from CERES measurements, so there is no need to use models to estimate the aerosol properties to infer ADRE (Loeb and Manalo-Smith, 2005). The challenge on the other hand is to obtain an estimate for the outgoing SW flux at TOA without aerosols from the measurement data, since aerosols are always present in the atmosphere. The aerosol-free flux is derived by establishing a linear relationship between coincident observations of SW fluxes and AODs, and then extrapolating to $AOD = 0$. It is noted that the resulting ADRE is an instantaneous value, i.e. representative only for the approximate time (and solar zenith angle, SZA) of the satellite overpass. Radiative transfer codes are needed to model the diurnal variation of the SZA, and to expand the obtained satellite observation-based ADRE to represent the 24 h monthly averages (Remer and Kaufman, 2006) comparable with other estimates of ADRE values presented in the literature. This satellite-based approach has been previously used over ocean e.g. by Loeb and Manalo-Smith (2005); Zhao et al. (2008)

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Christopher (2011), and over land e.g. by Patadia et al. (2008a, b); Sena et al. (2013) and most recently by Feng and Christopher (2013) (over land and ocean).

The coincident satellite observations of broad band fluxes and AOD provide an unique way to derive an observational based estimate for ADRE. However, even though this approach has been used in several studies, less attention has been paid on the method itself, which gives the motivation for this study. From a climate and Earth's radiative balance point of view the highly variable aerosol conditions over East Asia make the region important and interesting. For this study a part of Eastern of China (20–45° N, 110–125° E, Fig. 1) was selected as the region of interest, hosting one of the fastest growing economies in the world with rapidly expanding anthropogenic activity, industry, urbanization, and large-scale agriculture. Along with deriving the ADRE for 8 months from March to October 2009, the key questions in this study were focused on the method, e.g. how the satellite-based aerosol-free flux compares to the modeled flux, is the method working better with certain aerosol loading or surface, or could some method related parameter be used to identify possible method artifacts. The initial approach and threshold values were adapted from previous work of Sena et al. (2013) and Patadia et al. (2008a) with the difference that in this study also positive values of the ADRE are allowed. In the first part of the study a normalization of the CERES TOA fluxes is introduced in order to decrease the noise in flux observations not related to aerosols. In the second part of the study the ADRE is defined, and the aerosol-free flux values obtained from the satellite method are compared to the modeled values. In addition, dependencies between different linear fitting related parameters and, e.g., surface albedo are studied. Cases of positive ADRE are also investigated in some detail, and possible reasons for the positive sign (warming) are discussed.

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Remote sensing data

2.1 CERES SSF data

The CERES instruments onboard the EOS Terra, Aqua, and on Suomi NPP platforms provide radiance measurements in the shortwave (0.3–5.0 μm), infrared window (8.0–12.0 μm) and total (8.0–200 μm) broadband channels. The measured radiances are converted to TOA-fluxes using angular distribution models (e.g. Loeb and Kato, 2002). In this study data from the Terra platform were used from March to October 2009. The equatorial overpass time for Terra is about 10:30 LT. The CERES Single Scanner Footprint (SSF) dataset combines the CERES broadband observations with spatially and temporally coincident aerosol and cloud observations from the higher-resolution imager MODIS on the same satellite platform, and meteorological information provided by the Global Modeling and Assimilation Office (GMAO). The CERES-SSF footprint nadir resolution is about 20 km and includes point-spread-function weighted averages of 10 km MODIS collection 5 aerosol and cloud parameters. In this work only data over land were considered.

2.2 MODIS land data

The MODIS Land Cover Type Climate Modeling Grid product (MCD12C1) was used to identify inland water bodies within the study area. Based on the data, observations over rivers and lakes were removed from the dataset to reduce the surface inhomogeneity within the grid cells. For radiative transfer simulations monthly means of the SW broad band black-sky albedo were calculated using the MODIS MCD43C3 albedo product. The albedo files contain 16 days of combined level 3 data from Aqua and Terra (e.g. Gao et al., 2005; Zhang et al., 2010). The monthly mean albedo was defined as a weighted average including all the data where the 16 days measurement period overlapped with the month of interest.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Reference data

3.1 AERONET inversion data and aerosol type

AERONET (Aerosol Robotic Network, <http://aeronet.gsfc.nasa.gov>, Holben et al., 1998) is a global ground-based monitoring network of aerosol optical, microphysical, and radiative properties, providing observations that are available in a public domain. AERONET uses Cimel sunphotometers to measure AOD at 340, 380, 440, 500, 675, 870 and 1020 nm, but also provides retrievals of other aerosol parameters, such as the complex refractive index and single scattering albedo (SSA). Figure 1 shows the AERONET stations within the study area that provided data during 2009 and were used in this work. To get a picture of the prevailing aerosol types within the study area, level 2.0 (cloud-screened and quality assured) SSA was analyzed, and when level 2.0 data was not available level 1.5 (cloud-screened) SSA with certain criteria were used: from level 1.5 data only those observations were included in the analysis where all other quality criteria for lev 2.0 were met except the AOD threshold (Arola et al., 2013).

Figure 2 shows the monthly variation from March to October of SSA (lev 2.0) at 440 nm and the difference between SSA at 440 nm and the average of near infrared (NIR; 675, 870, and 1020 nm) channels in Beijing AERONET station. During spring lower SSA and negative difference of the 440 nm and NIR-channel SSA indicated the presence of dust type aerosol, whereas high SSA during summer and slightly positive difference between spectral SSA indicated urban-type pollution (Dubovik et al., 2002). Similar but somewhat weaker seasonal variation was observed in XiangHe, Xinglong, and Taihu stations. For Hong Kong and Dalanzagad stations lev. 1.5 data were used because lev. 2.0 data did not have enough data points for analysis. In Hong Kong SSA did not have clear seasonal variation, and showed throughout the year similar values than in Beijing during summer. In Dalanzagdad the variation of SSA (lev. 1.5 data) was relatively large, but the SSA medians indicated slightly more absorbing aerosol type than in other stations. Overall, based on the AERONET data it can be estimated that in the major part of the study area the SSA (at 440 nm) varies mainly around 0.9, in

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the southern part mainly around 0.95 and in the remote desert areas in the north west around 0.85.

3.2 Simulated TOA-fluxes and critical albedo

The magnitude and sign (cooling/warming) of ADRE at TOA with certain solar zenith angle is a combined result of aerosol type, loading, and brightness of the underlying surface. To get a priori estimate of what kind of ADRE pattern could be expected over the study area, a number of simulations of TOA-fluxes were carried out with LibRadtran radiative transfer code (Mayer and Kylling, 2005) using different scenarios about aerosol type, loading (AOD) and surface. Figure 3 shows the mean SW black-sky albedo in the study area calculated from the MODIS MCD43C3 albedo product, and simulated SW fluxes at TOA as a function of SW broad band surface albedo with varying AOD for two types of aerosols. For both of the aerosol types a critical albedo can be seen, where the TOA fluxes for increasing AOD are lower than for the aerosol-free case, resulting in positive values of ADRE. For highly scattering aerosols ($SSA = 0.97$) the critical albedo was about 0.42, whereas for more absorbing aerosols ($SSA = 0.80$) the critical albedo was much lower, about 0.14. The MODIS albedo map shows that in major part of the study area the black-sky SW albedo varied between 0.1 and 0.15, whereas over the highly populated and industrialized areas, e.g. the one extending from Beijing towards south west, the surface was somewhat brighter with an albedo about 0.17. The Gobi desert in the north and north west was significantly brighter than other areas (SW albedo ~ 0.25), which makes the area more susceptible to positive values of ADRE even if the aerosols were not highly absorbing.

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Method

4.1 Deriving the instantaneous SW ADRE from the satellite observations

The instantaneous SW ADRE at the top of the atmosphere for a given location and time is defined as

$$ADRE(lat, lon, \theta) = F_{0,TOA}(lat, lon, \theta) - F_{aer,TOA}(lat, lon, \theta) \quad (1)$$

where $F_{0,TOA}$ and $F_{aer,TOA}$ are the upward shortwave fluxes at TOA without and with aerosols, θ is the solar zenith angle, lat and lon are the location coordinates. While the CERES flux observations represent the instantaneous values of $F_{aer,TOA}$, the estimate for $F_{0,TOA}$ is obtained by establishing a linear regression between coincident CERES fluxes and MODIS AODs, and extrapolating the regression line to AOD = 0. In each 0.5° grid cell the cloud-free flux and AOD observations were collected over one month as in, e.g., Patadia et al. (2008b) and Sena et al. (2013). Similarly, to avoid very large pixel sizes, the solar and viewing zenith angles were restricted to less than 60°, and the AOD values were limited to 2.0 to avoid possible nonlinearities that could appear with extreme aerosol loadings. To filter out cloudy or partly cloudy pixels, the clear area percent coverage parameter at subpixel resolution was used, which is based on the MODIS 250 m resolution cloud mask. All CERES pixels where the clear area was less than 99.9 % were removed as in Sena et al. (2013). For a successful regression it was required that the number of coincident flux-AOD observations in a grid cell was at least 10 per month, and that the absolute value of correlation coefficient was 0.2 or larger. It is noted that in contrast to Patadia et al. (2008b) and Sena et al. (2013), also negative correlations between the AOD and SW TOA fluxes (i.e. positive ADRE) were allowed in this study.

The essential assumption in this method is that the effective aerosol type does not change considerably over a month, and SW flux changes are mainly related to the change in aerosol loading. This assumption does not hold if some aerosol episode such as biomass burning or dust outbreak occur, but overall the ADRE obtained from

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over
China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fitting should be representative for the monthly mean conditions at the time of the satellite overpass. Furthermore, the observed $F_{\text{aer,TOA}}$ not only depends on aerosols, but is also affected by changes in SZA, surface albedo, atmospheric water vapour content and the day-of-year (DOY, i.e. Sun–Earth distance). The SZA has a seasonal and diurnal variation reaching maximum values at sunrise and sunset, and minimum at noon. ADRE values depend on the SZA so that for larger values of the SZA, the ADRE at TOA becomes more negative. Hence, with large zenith angles cooling is stronger, and could be further enhanced with increasing aerosol loading. The amount of atmospheric water vapour is also subject to substantial seasonal variation within the study area. The climate in Eastern China is characterized by dry winters, and humid, rainy summers. The rainy season starts in May when the monsoon spreads gradually from the south east onto the Chinese mainland, and lasts until about September. With the increasing humidity ADRE at TOA becomes less negative. To account for these variations that are not related to aerosols, and to minimize the noise, the observed CERES fluxes $F_{\text{aer,TOA}}$ were normalized to fixed values of SZA, DOY, and water vapour content before establishing the linear fitting against AOD:

$$F_{\text{TOA,norm.}}^{\text{CER}} = F_{\text{aer,TOA}}^{\text{CER}} \cdot \frac{F_{\text{TOA}}^{\text{mod}}(\text{norm.SZA}) \cdot F_{\text{TOA}}^{\text{mod}}(\text{norm.WV}) \cdot F_{\text{TOA}}^{\text{mod}}(\text{norm.DOY})}{F_{\text{TOA}}^{\text{mod}}(\text{obs.SZA, WV, DOY})^3} \quad (2)$$

where the superscript “CER” refers to a CERES observation and “mod” to a modeled TOA flux. The subscript “norm” refers to the fixed values of SZA, water vapor content (WV) and DOY, respectively, to which the observed CERES fluxes are normalized. The fixed values for SZA and water vapor content were the monthly means in each grid cell, calculated from those SSF-file observations that were included in the linear fitting. The fixed DOY was the 15th day of the month, and $F_{\text{TOA}}^{\text{mod}}(\text{obs.SZA, WV, DOY})$ was the modeled CERES observation. In each modeled flux the input values for SZA, WV content, DOY, surface albedo, and AOD were taken from the CERES SSF file except in those cases when a fixed value was used for one of the three first parameters. In the reference flux simulations scattering aerosol type (single scattering albedo SSA =

0.97 at 550 nm) was assumed. Before deciding the aerosol type used in the simulated reference fluxes, a number of tests with more absorbing aerosol types were carried out to evaluate the sensitivity of the normalization procedure to the changing aerosol type. The tests indicated only small differences between the reference flux values associated with more absorbing aerosol types, and the effect on the normalization and linear fitting against AOD was minimal.

When quantifying ADRE, the normalization was carried out in only those grid cells where the number of proper observations was at least 10, and hence a successful linear fitting was possible. After the normalization of the CERES fluxes, the linear regression with the MODIS AOD was established, and instantaneous $F_{0,TOA}$ and ADRE were defined for those cells where the absolute value of correlation between the normalized fluxes and AOD was 0.2 or greater. The normalization according to the surface albedo was not included in this study, and hence the possible variations in surface albedo could still cause scatter in $F_{TOA,norm.}^{CER}$ values. Figure 4 illustrates the effect of normalization of the CERES fluxes on the linear regression in one grid cell (40.0° N, 117.0° E, October 2009). Overall, when considering the whole study period, the normalization increased the positive correlation between the CERES fluxes and the MODIS AODs, and decreased the root-mean-square error (RMSE) in the majority of the cases. It is noted that here RMSE indicates the goodness of the fit between CERES flux and MODIS AOD, and thus should not be interpreted as an uncertainty of $F_{0,TOA}$. Figure 5 illustrates the absolute values of the normalization corrections made to the observed CERES fluxes. As is seen, the largest corrections in units of $W m^{-2}$ are due to the normalization to fixed SZA, which also had a major contribution to the improved correlations between AOD and TOA fluxes. Normalization to the mean water vapor content was important especially during seasons and at locations when the water vapor content had large variation, but overall the distribution of absolute corrections was not as wide as for fixed SZA. The correction of observed fluxes for fixed DOY was mainly between -1.0 and $1.0 W m^{-2}$.

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ADRE obtained from the fitting is an instantaneous value, representative for the time of the satellite overpass time (SZA), i.e. about 10:30 LT. To estimate the 24 h mean ADRE and account for the diurnal variation of SZA, the instantaneous ADRE needs to be scaled with the help of modeled ADRE as in (Remer and Kaufman, 2006):

$$ADRE_{24h}^{Sat.}(lat, AOD, WV, \alpha) = ADRE_{Inst.}^{Sat.}(SZA, AOD, WV, \alpha) \cdot \frac{ADRE_{24h}^{mod}(lat, AOD, WV, \alpha)}{F_{inst.}^{mod}(SZA, AOD, WV, \alpha)} \quad (3)$$

where “mod” refers to modeled fluxes, and “ α ” to surface albedo. The scaling coefficient for 24 h ADRE was defined for each grid cell and the diurnal variation of SZA was calculated with 2 h time resolution. Since the diurnal variation of AOD and water vapour are not known, the parameters were assumed to be constant in the model simulations, and in each grid cell the monthly mean values of AOD, water vapor content and surface albedo were calculated from the CERES SSF files.

5 Results and discussion

5.1 Aerosol direct radiative effect

The monthly SW ADREs at TOA estimated using the satellite-based method were derived for eight months from March to October 2009. The best data coverage was obtained for October, when ADRE was successfully derived for 58 % of the land grid cells but for July ADRE was obtained for only 16 % of the land grid cells, which is mainly due to lack of cloud-free observations during the humid summer months. The seasonal medians of the instantaneous and diurnally averaged ADRE as well as the median AODs in the fitting are shown in Fig. 6. It is noted that the median AODs were calculated only for those grid cells, and of those AOD observations that were included in the linear fitting against CERES fluxes. Hence the seasonal AOD distributions did not e.g. include

observations where $\text{AOD} \geq 2.0$. The median instantaneous ADRE over the whole measurement period was -8.8 W m^{-2} , and the 24 h ADRE was -5.1 W m^{-2} . The instantaneous and diurnally averaged ADREs were negative over most of the study area, and the strongest cooling at TOA was often associated with large AOD, e.g. during spring and autumn over the area extending from south of Tianjin to Handan and further to Chengdu. Another example is Shanghai, where the cooling at TOA was stronger than over the surrounding areas. Figure 7 shows all the instantaneous ADREs derived in this study as a function of AOD, illustrating the connection of aerosol loading and cooling at TOA. However, some exceptions were also found. Some of the highest AODs were observed during spring over area south west from Wuhan Fig. 6c, but both the instantaneous and 24 h ADREs showed less negative, or even positive values over that area.

The spatial pattern of the ADRE during the summer was somewhat different than during the two other seasons. Strongest cooling effect was observed over the Jiangsu province north from Shanghai, where the instantaneous ADRE values were twice as negative as during spring or autumn. This was most probably related to crop burning in maize and wheat fields (Huang et al., 2012). On a monthly basis the other notable change during the summer was the more frequent appearance of pixels with positive ADRE, also over areas where the warming effect was not expected. The number of positive ADRE cases increased already during May, and then decreased by September. The case of positive ADRE is discussed in more detail in the following section.

5.2 Cases of positive ADRE

In the case of positive ADRE (i.e. warming at TOA), the correlation between normalized CERES fluxes and MODIS AODs is negative, i.e. the flux values decrease with increasing AOD. As shown in Sect. 3.2, positive values of ADRE were mainly expected over the desert, but were unexpectedly observed also over other areas, especially during the summer months. In some of the pixels initially weak positive ADRE changed to negative when the normalization procedure was applied, indicating that the positive



A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



effect was an artefact most probably related to variation of SZA or water vapor. However, even after the normalization a number of pixels with positive ADRE still existed over relatively dark surfaces. In fact, only 11 % of the positive ADRE cases were observed over bright surface (SW albedo ≥ 0.2). Since detailed information on aerosol types in the study area for specific days and locations was not available, the possibility of extremely absorbing aerosols causing some of the warming effect could not be definitely ruled out. One of the locations with positive ADRE was an area north west from Shanghai (around 33°N , 116.5°E), in May showing one of the highest negative correlation between the CERES fluxes and MODIS AODs outside the desert area. That area was selected as a test case for more detailed study.

Figure 8 illustrates the linear fitting after the normalization for the 33°N , 116.5°E grid cell in May 2009. The correlation coefficient between the normalized CERES fluxes and MODIS AOD was -0.51 , resulting in an instantaneous ADRE of 20.83 W m^{-2} . The aerosol-free TOA flux estimate obtained from fitting was 198.2 W m^{-2} , which is about 18 W m^{-2} higher than the modeled value for $F_{0,\text{TOA}}$ for the same grid cell. According to the simulations, a brighter surface with an albedo of about 0.175 would be required to produce $F_{0,\text{TOA}}$ similar to the fitting. However, in that grid cell the surface was darker, the broadband SW albedo was about 0.14. In May 2009 the particular grid cell consisted of observations from seven different days having altogether 33 coincident TOA flux-AOD observations. Figure 8 shows that when AOD was between 0.4 and 0.6, the normalized flux values varied a lot; from values of about 160 W m^{-2} to 190 W m^{-2} . For example, on 6 and 7 May the aerosol loading was about the same (AOD ~ 0.43), but the normalized flux values on these days differed by about 20 W m^{-2} . One explanation for such a large change in the flux values could be a significant change in aerosol type. According to the radiative transfer simulations, if the AOD would be about 0.4 the SSA should decrease from over 0.95 to about 0.85 to cause such a large difference in TOA flux values. This would imply that the assumption in the method of non-systematic changes in aerosol type was not valid in this case. On the other hand, this could be also related to cirrus contamination, which would mean that both AOD and TOA-flux observations were

A case study over
China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



affected. The study by e.g. Sun et al. (2011) showed that globally up to about 50 % of MODIS-derived clear sky scenes could actually be covered by invisibly thin cirrus clouds. Also, the climatology of all cirrus type clouds (not only subvisual) over China by Chen and Liu (2005) shows that the occurrence is highest during summer months.

Since there are no additional data available about aerosol types or cirrus, either of the possible explanations could be confirmed. However, the reference simulations of $F_{0,TOA}$ indicate that the satellite-based $F_{0,TOA}$ is too large, and the positive ADRE over the region was most probably a method artifact.

5.3 Comparison of $F_{0,TOA}$ with radiative transfer model

The advantage of the satellite based approach for defining ADRE is that $F_{aer,TOA}$ is directly obtained from CERES measurements, and models are not needed to estimate the aerosol or surface properties. Hence to assess the satellite method's ability to produce ADRE, the key question is how well this method can produce $F_{0,TOA}$. Since there is no validation (measurement) data for $F_{0,TOA}$, the LibRadtran radiative transfer model was used to create a dataset for a comparison. It is noted that the comparison between the satellite and model based $F_{0,TOA}$ does not necessarily indicate which method is giving more correct results but it illustrates in which cases the two methods produced similar estimates and when not. The $F_{0,TOA}$ was modeled for each eight months in the same $0.5^\circ \times 0.5^\circ$ grid that was used in the satellite-based method as explained in Sect. 3. $F_{0,TOA}$ was modeled in each month for only those cells where the conditions for a successful linear fitting were met. The monthly mean surface albedo was calculated from the MODIS land product as explained in Sect. 2.2, whereas the input for SZA and water vapour content were the monthly means calculated from the CERES SSF file using only those observations that were included in the fitting. The DOY was set to 15th day of the month.

The results show that the satellite fitting method and the radiative transfer simulations produce qualitatively similar $F_{0,TOA}$ spatial pattern. Figure 9 shows an example of $F_{0,TOA}$ from both methods for March 2009. The satellite-based approach caught the

A case study over
China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



main features similar to the modeled aerosol-free fluxes even some differences in the absolute $F_{0,TOA}$ values can be observed. Considering all pixels for which ADRE was determined between March and October 2009, in 31 % of the cases the satellite observations based method produced $F_{0,TOA}$ that was within 5 W m^{-2} , and in 58 % within 10 W m^{-2} from the modeled values (Fig. 10). In 19 % of the cases the satellite based $F_{0,TOA}$ was more than 10 W m^{-2} lower than the model, and in 23 % cases more than 10 W m^{-2} higher than the model. During the summer months the relative number of the extreme difference between satellite-based method and modeled $F_{0,TOA}$ was largest. During the whole eight month period there were only few areas where the satellite-based approach was producing systematically lower or higher $F_{0,TOA}$ than the model. In most of the pixels the the satellite-model $F_{0,TOA}$ difference changed from month to month.

The next step was to investigate whether the difference of aerosol-free flux between satellite and model ($\Delta F_{0,TOA}$) depends on some fitting related parameter such as the correlation coefficient between AODs and normalized CERES fluxes or RMSE, or vice versa, could some of these parameters predict e.g. extremely large $\Delta F_{0,TOA}$. Figure 11 shows density plots of $\Delta F_{0,TOA}$ (satellite-model estimate) as a function of correlation coefficient, RMSE, dynamic AOD range and number of observations obtained in a grid cell in one month. With high positive correlation ($R > 0.5$) $\Delta F_{0,TOA}$ was more likely within $\pm 10 \text{ W m}^{-2}$ but large differences in $F_{0,TOA}$ could still appear between the two methods, mainly satellite-based $F_{0,TOA}$ being lower than the modeled value. Results did not show any clear critical value of correlation which would indicate smaller differences between the satellite-based method and simulations. On the other hand, when the correlation was negative, it seems that in most of the cases the satellite based method produces larger $F_{0,TOA}$ than the model. Also with broad dynamic AOD range (monthly max.–min. AOD) or a high number of observations per month the count of extreme $\Delta F_{0,TOA}$ are less. Large dynamic AOD range was most often also associated with high positive correlation between AODs and TOA-fluxes and lower RMSE (not shown). On the other

hand, 78 % of the pixels having negative correlation between fluxes and AODs were associated with dynamic AOD range less than 0.4.

One of the major parameters affecting the CERES flux observations is the surface brightness, and hence the surface SW albedo could potentially have a large effect on the satellite-based approach for deriving ADRE. Even though the albedo varied considerably (between 0.1 and 0.3) within the study area, the relative variation (standard deviation/mean) of the SW albedo within the grid cells was typically small, about 2–5 %, when calculated using the MODIS SW black-sky albedo. Figure 12 shows $\Delta F_{0,TOA}$ as a function of grid cell mean MODIS SW black-sky albedo. Especially over bright surfaces the satellite-based method tend to produce lower estimate for $F_{0,TOA}$ than the model. This might be partly related to very small dynamic AOD range that was often observed over bright desert areas, but it is also noted that MODIS can have challenges of retrieving AOD over bright surfaces (Levy et al., 2010).

6 Conclusions

This study examined a satellite-based approach for determining the aerosol direct radiative effect (ADRE) over Eastern China (20–45° N, 100–125° E) from March to October 2009. In addition to the derived ADRE estimates, the method itself was investigated in detail. As ADRE at TOA is determined as the difference between SW fluxes without and with aerosols, the key challenge of this observation-based method is how well it can produce an estimate for the flux without aerosols. Because the aerosol-free flux can not be measured, the satellite-based approach uses coincident observations of SW broad band fluxes from CERES and AODs from MODIS to determine the monthly aerosol-free flux at TOA by establishing a linear regression between the two parameters. It is assumed that the changes in flux values are related to changes in aerosol loading, when the aerosol-free flux can be obtained as the y-intercept of the regression line. The satellite based ADRE estimate is an instantaneous value, representative only at the time of the satellite overpass (in this study at about 10:30 LT). Radiative transfer

calculations are needed to expand the estimate for diurnally averaged 24 h ADRE. The approach in this work was similar to that presented in Sena et al. (2013) and Pata-dia et al. (2008a) with the difference that also positive values of ADRE (i.e. negative correlation between TOA fluxes and AOD) were allowed.

The CERES flux observations not only depend on changes in aerosol loading, but were also affected by the variation of solar zenith angle (SZA), water vapor (WV), Sun–Earth distance (DOY), and surface albedo. Since the linear regression was performed using coincident observations from one month, the variation of these parameters also caused noise in the observed CERES fluxes. In this work a normalization procedure to fixed SZA, WV and DOY to the CERES fluxes was applied before the actual linear fitting. The results show that the normalization decreased the non-aerosol induced noise in the flux observations, and in the majority of the cases increased the positive correlation with AOD and decreased RMSE. In some cases the normalization also changed ADRE from weakly positive to negative, and hence possibly removed some method artifacts.

As expected, the ADRE over the study area was overall negative, and the strong cooling at TOA was often associated with high AOD. The instantaneous median ADRE of the study area was -8.8 W m^{-2} , and the diurnally averaged ADRE was -5.1 W m^{-2} , but locally a large variation around these medians were observed. Over heavily industrialized and populated areas the ADRE could be more than three times the median values, which indicated that within the study area the antropogenic aerosol emissions have large contribution to the cooling of the atmosphere. The obtained satellite-based ADRE estimates are in line with values found in other studies using different methods (e.g. Thomas et al., 2013). Locally some positive ADREs, especially during the summer moths, were observed outside desert areas where surface is darker, and the aerosols should have been strongly absorbing to produce a warming effect at TOA. In fact 78 % of all positive ADRE cases were observed over darker surfaces than desert. The majority of these cases were most probably method artifacts related to some systematic change in aerosol type, subvisual cloud contamination or both. In some locations this

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



conclusion was also supported by the high values of satellite-based $F_{0,TOA}$. However, on the whole, the pixels with positive ADREs were different, and some additional in situ data, e.g. aerosol single scattering albedos would have been needed to define case by case whether the warming effect was real or not.

One of the key questions in this study was how well the satellite method produces the aerosol-free flux. For comparison the aerosol-free fluxes for each month were also modeled using the LibRadtran radiative transfer code and MODIS broad band albedos. Results show that both methods produced qualitatively similar spatial patterns of $F_{0,TOA}$, but the absolute values differed somewhat. In 58 % of all cases the $F_{0,TOA}$ difference between the satellite method and the model was within $\pm 10 \text{ W m}^{-2}$. Extreme differences were often associated with low dynamic AOD range (< 0.4), and/or grid cells with less than 20 observations per month. The results also showed that over bright surfaces the satellite-based method tend to produce lower $F_{0,TOA}$ than model.

The satellite-based approach for determining ADRE offers a valuable observation-based comparison for the model simulated estimates of aerosol direct radiative effects. This study shows that this method can be applied over areas having large variations in aerosol load and surface properties, but attention should be paid especially to cases of positive ADRE. Over bright surfaces positive ADRE at TOA is physically justified, but over darker surfaces strongly absorbing aerosols are required to produce warming at TOA. One good indication of a method artifact which also applies in cases of negative ADRE, could be the satellite-based value of $F_{0,TOA}$. Comparison of satellite-based $F_{0,TOA}$ to e.g. simulated thresholds of $F_{0,TOA}$ could help to identify the apparent artifact cases.

Acknowledgements. This work has been supported by Academy of Finland Centre of Excellence program (project no. 272041), Marco Polo (EU FP7 SPACE Grant, agreement no 606953), MEGAPOLI (EU FP7 FP7-ENV-2007.1.1.2.1), EUCAARI (EU FP6, contract no. 036833-2), and Aerosol-cci (ESA-ESRIN project AO/1-6207/09/I-LG). The CERES SSF-data was obtained from the Atmospheric Science Data Center at the NASA Langley Research Cen-

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ter, and the MODIS surface data products were obtained through the online Data Pool at the NASA Land Processes Distributed Active Archive Center (LP DAAC).

References

- Arola, A., Eck, T. F., Huttunen, J., Lehtinen, K. E. J., Lindfors, A. V., Myhre, G., Smirnov, A., Tripathi, S. N., and Yu, H.: Influence of observed diurnal cycles of aerosol optical depth on aerosol direct radiative effect, *Atmos. Chem. Phys.*, 13, 7895–7901, doi:10.5194/acp-13-7895-2013, 2013. 15119
- Bellouin, N., Boucher, O., Haywood, J., and Reddy, M.: Global estimate of aerosol direct radiative forcing from satellite measurements, *Nature*, 438, 1138–1141, 2005. 15115, 15116
- Bellouin, N., Jones, A., Haywood, J., and Christopher, S.: Updated estimate of aerosol direct radiative forcing from satellite observations and comparison against the Hadley Centre climate model, *J. Geophys. Res.*, 113, D10205, doi:10.1029/2007JD009385, 2008. 15115
- Chen, B. and Liu, X.: Seasonal migration of cirrus clouds over the Asian Monsoon regions and the Tibetan Plateau measured from MODIS/Terra, *Geophys. Res. Lett.*, 32, L01804, doi:10.1029/2004GL020868, 2005. 15127
- Christopher, S. A.: Satellite remote sensing methods for estimating clear sky shortwave top of atmosphere fluxes used for aerosol studies over the global oceans, *Remote Sens. Environ.*, 115, 3002–3006, 2011. 15117
- Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanre, D., and Slutsker, I.: Variability of absorption and optical properties of key aerosol types observed in worldwide locations, *J. Atmos. Sci.*, 59, 590–608, 2002. 15119
- Feng, N. and Christopher, S.: Satellite and surface-based remote sensing of Southeast Asian aerosols and their radiative effects, *Atmos. Res.*, 122, 544–554, 2013. 15117
- Gao, F., Schaaf, C., Strahler, A., Roesch, A., Lucht, W., and Dickinson, R.: MODIS bidirectional reflectance distribution function and albedo Climate Modeling Grid products and the variability of albedo for major global vegetation types, *J. Geophys. Res.*, 110, D01104, doi:10.1029/2004JD005190, 2005. 15118
- García, O. E., Díaz, J. P., Expósito, F. J., Díaz, A. M., Dubovik, O., Derimian, Y., Dubuisson, P., and Roger, J.-C.: Shortwave radiative forcing and efficiency of key aerosol types us-

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing AERONET data, Atmos. Chem. Phys., 12, 5129–5145, doi:10.5194/acp-12-5129-2012, 2012. 15115

Haywood, J. and Boucher, O.: Estimated of the direct and indirect radiative forcing due to Tropospheric aerosols: a review, Rev. Geophys., 38, 513–543, 2000. 15115

5 Heald, C. L., Ridley, D. A., Kroll, J. H., Barrett, S. R. H., Cady-Pereira, K. E., Alvarado, M. J., and Holmes, C. D.: Beyond direct radiative forcing: the case for characterizing the direct radiative effect of aerosols, Atmos. Chem. Phys. Discuss., 13, 32925–32961, doi:10.5194/acpd-13-32925-2013, 2013. 15115

10 Holben, B., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET – a federated instrument network and data archive for aerosol characterization, Remote Sens. Environ., 66, 1–16, 1998. 15119

Huang, X., Li, M., Li, J., and Song, Y.: A high-resolution emission inventory of crop burning in fields in China based on MODIS thermal anomalies/fire products, Atmos. Environ., 50, 9–15, 2012. 15125

15 Jacobson, M.: Global direct radiative forcing due to multicomponent anthropogenic and natural aerosols, J. Geophys. Res., 106, 1551–1568, 2001a. 15115

Jacobson, M.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols, Nature, 409, 695–697, doi:10.1038/35055518, 2001b. 15116

20 Levy, R. C., Remer, L. A., Kleidman, R. G., Mattoo, S., Ichoku, C., Kahn, R., and Eck, T. F.: Global evaluation of the Collection 5 MODIS dark-target aerosol products over land, Atmos. Chem. Phys., 10, 10399–10420, doi:10.5194/acp-10-10399-2010, 2010. 15129

Loeb, N. G. and Kato, S.: Top-of-atmosphere direct radiative effect of aerosols over the tropical oceans from the Clouds and the Earth's Radiant Energy System (CERES) satellite instrument, J. Climate, 15, 1474–1484, 2002. 15118

25 Loeb, N. G. and Manalo-Smith, N.: Top-of-atmosphere direct radiative effect of aerosols over global oceans from merged CERES and MODIS observations, J. Climate, 18, 3506–3526, 2005. 15115, 15116

Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations – description and examples of use, Atmos. Chem. Phys., 5, 1855–1877, doi:10.5194/acp-5-1855-2005, 2005. 15120

30 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D.,

A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over
China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys.*, 13, 1853–1877, doi:10.5194/acp-13-1853-2013, 2013. 15115, 15116

Patadia, F., Gupta, P., and Christopher, S. A.: First observational estimates of global clear sky shortwave aerosol direct radiative effect over land, *Geophys. Res. Lett.*, 35, L04810, doi:10.1029/2008JD010754, 2008a. 15117, 15130

Patadia, F., Gupta, P., Christopher, S. A., and Reid, J. S.: A Multisensor satellite-based assessment of biomass burning aerosol radiative impact over Amazonia, *J. Geophys. Res.*, 113, D12214, doi:10.1029/2007JD009486, 2008b. 15117, 15121

Quaas, J., Boucher, O., Belloiun, N., and Kinne, S.: Satellite-based estimate of the direct and indirect aerosol climate forcing, *J. Geophys. Res.*, 113, D05204, doi:10.1029/2007JD008962, 2008. 15115

Remer, L. A. and Kaufman, Y. J.: Aerosol direct radiative effect at the top of the atmosphere over cloud free ocean derived from four years of MODIS data, *Atmos. Chem. Phys.*, 6, 237–253, doi:10.5194/acp-6-237-2006, 2006. 15116, 15124

Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006. 15115, 15116

Sena, E. T., Artaxo, P., and Correia, A. L.: Spatial variability of the direct radiative forcing of biomass burning aerosols and the effects of land use change in Amazonia, *Atmos. Chem. Phys.*, 13, 1261–1275, doi:10.5194/acp-13-1261-2013, 2013. 15117, 15121, 15130

Sun, W., Videen, G., Kato, S., Lin, B., Lukashin, C., and Hu, Y.: A study of subvisual clouds and their radiation effect with a synergy of CERES, MODIS, CALIPSO, and AIRS data, *J. Geophys. Res.*, 116, D22207, doi:10.1029/2011JD016422, 2011. 15127

Thomas, G. E., Chalmers, N., Harris, B., Grainger, R. G., and Highwood, E. J.: Regional and monthly and clear-sky aerosol direct radiative effect (and forcing) derived from

the GlobAEROSOL-AATSR satellite aerosol product, Atmos. Chem. Phys., 13, 393–410, doi:10.5194/acp-13-393-2013, 2013. 15116, 15130

Yu, H., Dickinson, R. E., Chin, M., Kaufman, Y., Zhou, M., Zhou, L., Tian, Y., Dubovik, O., and Holben, B. N.: Direct radiative effect of aerosols as determined from a combination of MODIS retrievals and GOCART simulations, J. Geophys. Res., 109, D03206, doi:10.1029/2003JD003914, 2004. 15116

Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. A., Anderson, T. L., Balkanski, Y., Bellouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T., and Zhou, M.: A review of measurement-based assessments of the aerosol direct radiative effect and forcing, Atmos. Chem. Phys., 6, 613–666, doi:10.5194/acp-6-613-2006, 2006. 15116

Zhang, X., Liang, S., Wang, K., Li, L., and Gui, S.: Analysis of global land surface short-wave broadband albedo from multiple data sources, IEEE J. Sel. Top. Appl., 3, 296–305, doi:10.1109/JSTARS.2010.2049342, 2010. 15118

Zhao, T., Yu, H., Laszlo, I., Chin, M., and Conant, W.: Derivation of component aerosol direct radiative forcing at the top of the atmosphere for clear-sky oceans, J. Quant. Spectrosc. Ra., 109, 1162–1186, 2008. 15116

**A case study over
China**

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over
China

A.-M. Sundström et al.




Figure 1. Map of the study area with AERONET stations that provided data during 2009. AERONET stations are 1 = Beijing, 2 = XiangHe, 3 = Xinglong, 4 = Dalanzadgad, 5 = Lanzhou city, 6 = Taihu, 7 = Hong Kong Polytech., and 8 = Hong Kong Hok Tsui.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A case study over
China**

A.-M. Sundström et al.




Figure 2. Monthly variation of the SSA at 440 nm (left) and the difference between SSA at 440 nm and average of Near Infrared Channels (NIR, 670, 870, and 1020 nm) (right) at the Beijing AERONET station. The red central marks in each box indicate the median values while the upper and lower edges of the boxes indicate 75th and 25th percentiles. The whiskers are showing extreme values and the outliers are marked with “+”.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion



(a)



(b)



(c)

Figure 3. Monthly mean (September 2009) of the shortwave black-sky broadband albedo obtained from the MODIS MCD43C3 data **(a)**, and radiative transfer simulations of TOA SW fluxes as a function of surface albedo and varying aerosol loading for highly scattering **(b)** and absorbing **(c)** aerosol type.

A case study over China

A.-M. Sundström et al.




Figure 4. An example of the linear regression between CERES TOA fluxes and MODIS AODs without (left panel) and with (right panel) the normalization of the observed fluxes. The data is from a grid cell that covers the XiangHe AERONET station. The observations are made during October 2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over China

A.-M. Sundström et al.




Figure 5. Distribution of the absolute corrections made to the observed CERES fluxes due to the normalization to fixed solar zenith angle, atmospheric water vapour content and day of year. The distribution includes all CERES observations within the study area between March and October 2009 that have been used in the fitting. White areas denote pixels where enough data has not been available for a successful linear fitting for any of the months.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion






(a) Instantaneous ADRE median Mar.-May 2009. (b) 24 h ADRE median Mar.-May 2009. (c) AOD median Mar.-May 2009.



(d) Instantaneous ADRE median Jun.-Aug. 2009. (e) 24 h ADRE median Jun.-Aug. 2009. (f) AOD median Jun.-Aug. 2009.



(g) Instantaneous ADRE median Sep.-Oct. 2009. (h) 24 h ADRE median Sep.-Oct. 2009. (i) AOD median Sep.-Oct. 2009.

Figure 6. Seasonal medians of the instantaneous (left) and 24 h averaged (middle) ADRE over the study area as well as the corresponding seasonal medians of AODs which have been used in the linear fitting against CERES fluxes. The 24 h averaged ADRE accounts only for the SZA diurnal variation, whereas other parameters such as AOD and water vapor content were assumed constant in the diurnal averaging.

A case study over China

A.-M. Sundström et al.




Figure 7. Aerosol radiative effect as a function of AOD. The AOD values denote the center of each 0.1 unit bin. Red central marks denote the median values of the ADRE whereas the upper and lower edges of the box denote the 75th and 25th percentiles. Whiskers show extreme values and outliers are denoted by individual “+” marks.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A case study over China

A.-M. Sundström et al.




Figure 8. Linear regression in lat = 33.0° N , lon = 116.5° E grid cell in May 2009 illustrating the negative correlation between normalized fluxes and AOD obtained outside the desert area. The MODIS black-sky SW albedo for the grid cell was 0.14.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over China

A.-M. Sundström et al.




Figure 9. The $F_{0,TOA}$ obtained with satellite-based method and radiative transfer simulations for March 2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A case study over
China**

A.-M. Sundström et al.




Figure 10. Difference of $F_{0,TOA}$ obtained from satellite-based fitting method and radiative transfer model simulations including all pixels within the study area from March to October 2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A case study over
China

A.-M. Sundström et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 11. The difference between satellite-based and modeled $F_{0,TOA}$ as a function of **(a)** correlation coefficient between CERES fluxes and MODIS AODs, **(b)** RMSE of the linear fitting, **(c)** dynamic AOD range (the difference between grid cell monthly max and min AOD values), and **(d)** number of observations in a grid cell/month.

**A case study over
China**

A.-M. Sundström et al.




Figure 12. The difference between satellite-based and modeled $F_{0,TOA}$ as a function of the MODIS black-sky SW albedo.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

