

1 **Erythemal ultraviolet irradiation trends in the Iberian**  
2 **Peninsula from 1950 to 2011**

3

4 **R. Román<sup>1</sup>, J. Bilbao<sup>1</sup>, and A. de Miguel<sup>1</sup>**

5 [1]{University of Valladolid, Laboratorio de Atmósfera y Energía, Department of Applied  
6 Physics, Valladolid, Spain. }

7 Correspondence to: R. Román (robertor@fa1.uva.es)

8

9 **Abstract**

10 Erythemal ultraviolet (UVER) irradiation was reconstructed at nine Spanish locations, with  
11 series starting around 1950 in five of these places. Each series was checked by applying  
12 homogeneity tests in order to discard non-homogeneous series. Available series were used to  
13 create an averaged Iberian Peninsula UVER series. Results indicate that annual UVER  
14 irradiation in the Iberian Peninsula increased by  $155 \text{ Jm}^{-2}$  (6.5%) between 1950 and 2011 due  
15 to the decrease observed in atmospheric ozone rather than changes in aerosol and clouds. By  
16 contrast, annual UVER irradiation increased by  $135 \text{ Jm}^{-2}$  (5.6%) between 1985 and 2011,  
17 mainly due to changes in aerosol and clouds.

18

19

20

21

22

23

24

25

## 1 **1 Introduction**

2 Among other effects, ultraviolet (UV) radiation, which is a part of total solar shortwave (SW)  
3 radiation, produces harmful effects on human skin, such as erythema (sunburn) induction  
4 (UNEP, 2003). Nevertheless, UV radiation exposure is also beneficial for the health, for  
5 example, by contributing towards human Vitamin D synthesis (Webb, 2006). The  
6 effectiveness of UV radiation in producing erythema on human skin is usually quantified by  
7 the erythemal action spectrum (McKinlay and Diffey, 1987), and the UV radiation weighted  
8 by this spectrum is erythemal ultraviolet (UVER) radiation.

9 The damage caused to human skin by UVER radiation is cumulative and is proportional to  
10 exposure time (WHO, 1995). It is therefore important to know both present-day as well as  
11 past UVER radiation levels in order to estimate future epidemiological data related to diseases  
12 caused by sun exposure. However, it was not until the 1970s and 1980s that the first long-  
13 term UVER measurement databases appeared (Scotto et al., 1988; Blumthaler and Ambach,  
14 1990; Correl et al., 1992; Borkowski, 2000). In fact, the longest UVER series in Spain started  
15 in 1995. In order to obtain UVER data previous to the use of instrumentation, several authors  
16 have reconstructed UVER series using other available records like SW radiation, sunshine  
17 duration, and cloud cover (Lindfors et al., 2003, 2007; den Outer et al., 2005, 2010; Rieder et  
18 al., 2008; Walker, 2010; Antón et al., 2011a; Bilbao et al., 2011 among others).

19 UVER radiation is sensitive to factors such as ozone, clouds, and aerosol particles in the  
20 atmosphere. Over the last few decades, the presence of these factors in the atmosphere has  
21 changed, and might have affected past UVER levels.

22 Solar radiation reaching the Earth's surface ranges mainly between 305 nm and 2800 nm, its  
23 wavelength integration being known as SW radiation, and it plays a key role in the Earth's  
24 energy balance and radiation budget. SW radiation decreased between 1950 and the mid  
25 1980s in the Northern Hemisphere, a phenomenon known as "global dimming" (Stanhill and  
26 Cohen, 2001). SW radiation began to increase in the mid 1980s in the Northern Hemisphere, a  
27 phenomenon known as "global brightening" (Wild et al., 2005). Dimming were caused  
28 because aerosol loads increased in the Northern Hemisphere between 1950 and the mid 1980s  
29 absorbing and scattering (aerosol direct effect) more radiation, although after the mid-1980s  
30 the aerosol load started to decrease (Wild, 2009, 2012). Changes in aerosols can modify the  
31 microphysical properties of clouds (aerosol indirect effect), since aerosols act as condensation  
32 nuclei, which contributes to enhance the dimming and brightening phenomena. The

1 mentioned aerosol and cloud changes might cause variations in the amount of UVER  
2 radiation reaching Earth. Dimming and brightening phenomena were also observed in the  
3 Iberian Peninsula by Sánchez-Lorenzo et al. (2007, 2013a, b) and Mateos et al. (2013, 2014).

4 The atmospheric total ozone column (TOC) evidenced major depletion after the late 1970s up  
5 to the mid 1990s due to strong atmospheric emission of halogen gases between the 1960s and  
6 1980s (WMO, 2011). TOC evolution in the Iberian Peninsula was studied in depth by Román  
7 et al. (2014d) in the dimming and brightening periods, with usually negative but not  
8 significant trends being reported in both periods, and a statistically significant trend of -  
9  $0.7\%dc^{-1}$  (percent per decade units) in the annual TOC between 1950 and 2011. Antón and  
10 Mateos (2013) showed that these TOC changes have no significant influence on SW radiation  
11 (e.g., the SW radiative effects due to TOC variations between late 1970s and 2012 were  
12 smaller than  $0.2 Wm^{-2}$  over the Iberian Peninsula). Nevertheless, these TOC changes might  
13 prove extremely relevant for UVER evolution over the last few decades, marking the  
14 difference between the SW and UVER trends in the past.

15 Many authors have found statistically significant positive trends over the last decades for  
16 UVER radiation at different places in the following European countries: Austria, Czech  
17 Republic, Finland, Greece, Germany, the Netherlands, Norway, Sweden, and Switzerland  
18 (Lindfors et al., 2003; 2007; Rieder et al., 2008; Walker, 2010; den Outer et al., 2010;  
19 Krzyscin et al., 2011). From these papers, the increase in UVER radiation values was mainly  
20 attributed to ozone depletion since the late 1970s; some of them also suggested that the  
21 UVER increase was partially caused by a diminishing of cloudiness or aerosols.

22 UVER radiation presents high levels in the Iberian Peninsula due to the great values of the  
23 elevation solar, making it an interesting area to study the evolution of UVER radiation.  
24 However, in the Iberian Peninsula, UVER radiation has only been reconstructed at Valladolid,  
25 since 1991 (Bilbao et al., 2011), and at Badajoz and Cáceres (only in the summer months)  
26 since 1950 (Antón et al., 2011a), a significant rise in UVER radiation levels having been  
27 reported at the three sites.

28 Recently Román et al. (2014d) obtained the TOC trends from 1950 to 2011 at the Iberian  
29 Peninsula; and Román et al. (2014a) the SW and temperature trends for the same period and  
30 region, but not the UVER trends due to the lack of UVER data. As a first step, Román et al.  
31 2014b simulated the UVER irradiance under cloudless conditions at the Iberian Peninsula  
32 with a radiative transfer model, using as inputs monthly climatological values of aerosols,

1 water vapour, etc. They also characterized the uncertainty of these simulations caused by the  
2 monthly variability (standard deviation of the used monthly climatological values) and  
3 uncertainty of the inputs. These cloudless simulations are useful to reconstruct UVER  
4 radiation under all conditions (e.g., Lindfors et al., 2003, 2007; den Outer et al., 2005;  
5 Walker, 2010; Bilbao et al., 2011).

6 Therefore, in this framework, the main objectives of the present paper are: 1) to develop a  
7 new UVER reconstruction model and optimise another one (both models based on cloudless  
8 UVER simulations); 2) to reconstruct UVER irradiation series since 1950 using the developed  
9 models with data at certain Spanish locations over the Iberian Peninsula; and 3) to analyse  
10 their evolution and trends of the reconstructed dataset. A novel issue of this paper is the  
11 analysis of the annual and seasonal (not only summer) UVER trends at the Iberian Peninsula  
12 from 1950, since these trends were not yet calculated in previous studies. Another novel issue  
13 is the quantification of the role of aerosols and clouds (both together), and ozone in the  
14 changes of UVER during the last six decades.

15 The paper is structured as follows: Sect. 2 shows the relevant information concerning the  
16 locations, the instrumentation, and the explanation of all the data used. The methods used to  
17 obtain the reconstructed UVER series are developed and explained in detail in Sect. 3. Sect. 4  
18 presents the main results for the evolution and trends of UVER. The factors not taken into  
19 account in the work are mentioned in Sect. 5. Finally, Sect. 6 summarises the main results and  
20 conclusions.

21

## 22 **2 Place, instrumentation, and data**

### 23 **2.1 Places and instrumentation**

24 All data used in this paper were taken at nine Spanish radiometric stations located in the  
25 Iberian Peninsula. These locations are marked in Fig. 1 and their coordinates are also shown  
26 in Table 1. The Iberian Peninsula is well covered by these locations. One of these stations is  
27 controlled by the University of Valladolid and is located in the village of “Villalba de los  
28 Alcores” (de Miguel et al. 2012). The rest are controlled by the Spanish Meteorological  
29 Agency (Moreta et al., 2013). Hourly UVER and SW irradiance were measured at all these  
30 stations, although sunshine duration and temperature were not measured at the Villalba  
31 station. Therefore, the sunshine duration and meteorological variables, as with temperature,

1 measured at the Valladolid Airport AEMet station were thus considered the same as at the  
2 Villalba station, since the two stations are located just a few kilometres away from each other.

3 Hourly UVER irradiance was recorded at the nine locations using UVB-1 pyranometers  
4 (Yankee Environmental Systems Inc.). These pyranometers were periodically calibrated by a  
5 two-step method (Vilaplana et al., 2009), which provides a combined uncertainty (68%  
6 confidence) of between 5.4% and 8.0% for the measured hourly UVER data (Hülßen and  
7 Gröbner, 2007). This uncertainty was considered the maximum (8.0%) in this work. The  
8 oldest UVER data recorded in Spain date from 1<sup>st</sup> November 1995 and continue up to the  
9 present day at the Madrid station.

10 Hourly SW irradiance was initially measured at each location using a CM6B pyranometer  
11 (Kipp & Zonen), whose spectral response ranges from 305 nm to 2,800 nm. The expanded  
12 uncertainty (95% confidence) of the hourly SW recorded by this pyranometer is 8%, and was  
13 the expanded uncertainty assumed by all available hourly SW measurements even when more  
14 recent records were taken using improved pyranometers (displaying less uncertainty). The  
15 longest SW data in Spain date from 11<sup>th</sup> July 1973 and continue up to the present day at the  
16 Madrid station.

17 In general, it was not possible to obtain information on the instruments used for sunshine  
18 duration records at the various stations, although most were probably Campbell-Stokes  
19 heliographs (Sánchez-Lorenzo et al., 2007). This heliograph comprises a spherical lens which  
20 concentrates direct radiation from the sun onto a dark paper card, which is burned when direct  
21 radiation exceeds a certain threshold. The combined uncertainty of the sunshine duration  
22 records was assumed to be 15 minutes (0.25 h). The daily sunshine fraction (F) is the ratio of  
23 the measured sunshine duration to the same sunshine duration under cloudless conditions  
24 ( $SunDu_{cl}$ ). This variable was calculated by the following equation (Iqbal, 1983):

$$25 \quad SunDu_{cl} = \frac{24}{\pi} \arccos\left(\frac{\cos(\theta_s) - \sin(\delta)\sin(\phi)}{\cos(\delta)\cos(\phi)}\right) \quad (1)$$

26 where  $SunDu_{cl}$  is in hours,  $\delta$  is the solar declination,  $\Phi$  is the location latitude, and  $\theta_s$  is the  
27 solar zenith angle (SZA) at sunset and sunrise (equal to 87° in this case).  $SunDu_{cl}$  was  
28 calculated between the solar zenith angle of 87° near sunrise and the SZA equal to 87° near  
29 sunset, since direct solar radiation usually is not enough to burn the dark paper card even  
30 under cloud free conditions for a SZA below 87°. The oldest available F data in Spain date

1 from 1<sup>st</sup> January 1920 and continue up to the present day at the Madrid station. F data have  
2 been available at certain locations (A Coruña, Madrid, San Sebastián, Tortosa, and Villalba)  
3 since the 1950s.

4 The instruments used to take all the mentioned measurements were well calibrated on a  
5 regular basis, following World Meteorological Organization (WMO) recommendations  
6 (Webb et al., 2006; WMO, 2008) for instrument maintenance, and involved: bubble levelling  
7 of the instruments, cleaning domes, monitoring and replacing desiccant, etc. In addition, a  
8 high-quality control of UVER, SW, and F data was applied to all available data in order to  
9 reject possible spurious and outlier data. Daily UVER and SW irradiation data were obtained  
10 integrating the hourly values each day.

## 11 **2.2 Additional data**

12 Other atmospheric variables were also used in this work: aerosol optical depth (AOD),  
13 Angström exponent, aerosol single scattering albedo (SSA), water vapour column (w), TOC  
14 and surface albedo. The daily data of these variables (except TOC) were used to calculate a  
15 monthly climatology in order to be used as input in a radiative transfer model to calculate  
16 UVER and SW irradiance under cloudless conditions (See Sect. 3.1). These daily data are  
17 described in this section, and the monthly climatology values were obtained from them.

18 Daily AOD at 433 nm and 670 nm (“MISR-Terra Prod.ver.31: MIL3DAE.004” product) from  
19 the MISR instrument (Multi-angle Imaging SpectroRadiometer) were obtained at each  
20 location from 2000 to 2012. The Angström Exponent was directly calculated using the AOD  
21 values at the two wavelengths. The daily w (“MODIS-Terra Ver. 5.1: MOD08\_D3.051”  
22 product) from MODIS (MODerate resolution Imaging Spectroradiometer) was also obtained  
23 at each location between 2000 and 2012. AOD and w data were downloaded from the  
24 GIOVANNI application (GES-DISC Interactive Online Visualization ANd aNalysis  
25 Infrastructure; <http://disc.sci.gsfc.nasa.gov/giovanni>) as an averaged 0.2°x0.2° square centred  
26 at each location (Acker and Leptoukh, 2007). The SSA at 354 nm and 500 nm retrieved from  
27 the OMI (Ozone Monitoring Instrument) instrument between 2004 and 2011 were also  
28 obtained for all locations as overpass files available at AVDC (Aura Validation Data Center).  
29 These data are the same as those used by Román et al. (2014c), who calculated the uncertainty  
30 of some of these products in the Iberian Peninsula. The combined uncertainty of AOD at 433  
31 nm and 670 nm is 0.074 and 0.054, respectively. The combined uncertainty of the Angström

1 Exponent is below 0.5 when AOD at 433 nm is above 0.25, except for high Angström  
2 Exponent values. The combined uncertainty of the water vapour column is between 0.38 cm  
3 ( $w=0.5$  cm) and 0.52 cm ( $w=3$  cm).

4 A daily TOC series for 1950 to 2011 was available for each location. These series comprised  
5 different databases: ground-based ozone; ozone retrieved from TOMS (Total Ozone Mapping  
6 Spectrometer) instrument on board Nimbus-7, Meteor-3, and Earth-probe satellites; TOC  
7 from OMI; retrieved TOC from GOME (Global Ozone Monitoring Experiment) and GOME-  
8 2 instruments on board ERS-2 and MetOp-A satellites; and reconstructed ozone data from the  
9 COST-726 project (Krzyscin, 2008; [www.cost726.org](http://www.cost726.org)). The construction of these TOC series  
10 was explained by Román et al. (2014d) who, by means of an intercomparison with ground  
11 measurements, calculated that the combined uncertainty of the daily TOC values of these  
12 series was around 10 Dobson Unit (DU).

13 Surface albedo data between 2000 and 2011 were obtained each eight days at seven  
14 wavelength ranges (459-479 nm, 545-565 nm, 620-670 nm, 841-876 nm, 1230-1250 nm,  
15 1628-1652 nm, and 2105-2155 nm) from the MCD43A3 product of MODIS instruments  
16 (Schaaf et al., 2002). In addition, daily surface albedo at 360 nm between 1957 and 2002 was  
17 obtained from the COST-726 project database as an interpolation at each location of the  
18 available data grid (Schwander et al., 1999; Tanskanen, 2004). More information concerning  
19 the albedo and ozone data used in this work is available in Román et al. (2014d).

20

### 21 **3 Reconstructed UVER series**

#### 22 **3.1 Simulations under cloudless conditions**

23 Global, diffuse and direct horizontal UVER and SW irradiance were simulated under  
24 cloudless conditions using a radiative transfer model (UVSPEC/libRadtran) from 1950 to  
25 2011 each hour for all locations shown in Fig. 1. UVSPEC is the main tool of the libRadtran  
26 (version 1.7 in this work) software package developed by Mayer and Kylling (2005). For  
27 UVER simulations, irradiance was calculated each 1 nm from 280 nm to 400 nm under cloud-  
28 free conditions using the “cdisor” solver with six streams (Buras et al., 2011) and the  
29 “SUSIM SL2” extraterrestrial spectrum (Van Hoosier et al., 1988), these obtained values then  
30 being weighted with the erythemal action spectrum. For SW simulations, the model was run  
31 under cloudless conditions using the “twostr” solver (Kylling et al., 1995), the extraterrestrial

1 spectrum from Kurucz (1992), and the pseudo-spectral k-distribution “SBDART” from  
2 Ricchiazzi et al. (1998). Irradiance was calculated from 305 nm to 800 nm in 2 nm intervals,  
3 from 800 nm to 1,600 nm in 5 nm intervals, and from 1,600 to 2,800 nm in 10 nm intervals,  
4 these spectral values then being spectrally integrated to obtain SW irradiance. An hourly  
5 (UVER or SW) irradiance value was simulated at a fixed SZA given by the averaged cosine  
6 of the SZA over the hour.

7 The UVSPEC model was run using standard vertical profiles. A mid-latitude summer  
8 atmosphere with spring-summer aerosol profiles was used as input for the months from May  
9 to October, with a mid-latitude winter atmosphere with fall-winter aerosol profiles being  
10 selected for the other months (Anderson et al., 1987; Shettle, 1989). These vertical profiles  
11 were rescaled with monthly climatological tables of water vapour, AOD at 443 nm, Angström  
12 Exponent, and SSA (at 354 nm for UVER simulations and at 500 nm for SW ones). These  
13 climatology tables (one per location and variable) comprised 12 monthly averaged (using all  
14 available data) values for each variable, said climatological tables being available in Román et  
15 al. (2014c). The daily TOC at each location was included in the inputs, changing the value for  
16 each location on each day. Finally, the surface albedo values were also monthly averaged for  
17 each wavelength (twelve monthly values at each location for each available wavelength),  
18 these monthly values being linearly interpolated to obtain the monthly surface albedo at all  
19 the other wavelengths to be then used as input in the radiative transfer code (it is well detailed  
20 in Román et al., 2014d).

21 Both the combined and the expanded uncertainty of all simulations were calculated using the  
22 results obtained by Román et al. (2014b), who calculated the maximum variations in  
23 simulated UVER and SW irradiance caused by the uncertainty of the inputs. Simulated hourly  
24 SW and UVER values were also compared with global SW and UVER irradiance  
25 measurements under cloudless conditions by Román et al. (2014b). It was found there was  
26 better agreement for low SZA values, and that the differences between simulations and  
27 measurements were in agreement within the uncertainty. Daily UVER and SW irradiation  
28 were calculated for each day at each location, and the uncertainties of these daily values were  
29 also calculated. Román et al. (2014b) compared these simulations with measured irradiation  
30 under cloudless conditions, with better agreement being found for the spring and summer  
31 months. For all months and locations together, a mean bias error (MBE) of -0.1% and a root



1 mean square error (RMSE) of 3.6% for the SW case, and an MBE of 2.9% and an RMSE of  
2 7.7% for the UVER case, were also reported.

### 3 3.2 Reconstruction models

#### 4 3.2.1 Cloud Modification Factor

5 The cloud modification factor (CMF) is defined as the ratio between measured radiation and  
6 simulated radiation under cloudless conditions:

$$7 \quad CMF_R = \frac{R_{me}}{R_{cl}} \quad (2)$$

8 where R can be SW or UVER, the “me” index indicating the measured R value, and the “cl”  
9 index being for simulated R radiation under cloudless conditions. CMF can be calculated for  
10 hourly irradiance or daily irradiation values. CMF is a useful variable to quantify cloudiness,  
11 CMF near 1 indicating cloudless conditions, CMF close to 0 being for high overcast  
12 cloudiness, and CMF above 1 indicating enhancement effects (Sabburg and Parisi, 2006;  
13 Sabburg and Calbó, 2009; Piedehierro et al., 2014). CMF for SW ( $CMF_{SW}$ ) is usually  
14 different for UVER ( $CMF_{UVER}$ ). The relationships between  $CMF_{UVER}$  and other variables, like  
15  $CMF_{SW}$ , are an important research topic since they prove useful for reconstructing UVER  
16 data. If a relationship between  $CMF_{UVER}$  and  $CMF_{SW}$  is given by a function f:

$$17 \quad CMF_{UVER} = f(CMF_{SW}) \quad (3)$$

18 then UVER radiation can be obtained when the measured SW radiation, UVER and SW  
19 simulations under cloudless conditions, and the f function are known:

$$20 \quad UVER = f(CMF_{SW}) UVER_{cl} \quad (4)$$

21 where UVER is the calculated UVER radiation, and  $UVER_{cl}$  is the simulated UVER under  
22 cloudless conditions. The main goal of this section is to obtain relationships such as the one  
23 given by Eq. (3) in order to reconstruct UVER data as a function of other variables as in Eq.  
24 (4).

#### 25 3.2.2 Model based on hourly SW irradiance measurements

26 Some UVER reconstruction models are based on different measured variables (Calbó et al.,  
27 2005), with several UVER reconstruction models being based on SW radiation

1 measurements, which were normally measured before UVER (Bodeker and McKenzie, 1996;  
2 Kaurola et al., 2000; Matthijsen et al., 2000; den Outer et al., 2000, 2005; Lindfors et al.,  
3 2007; Walker, 2010; Antón et al., 2011b; Bilbao et al., 2011). The data used to obtain a model  
4 based on SW were the measured values when both SW and UVER irradiance measurements  
5 were available. A total of 294,047 pairs of data (SW and UVER) were available for SZA  
6 below 85° taking all locations into account. Hourly CMF was calculated for hourly SW and  
7 UVER irradiance with the mentioned available data. The reconstruction model proposed in  
8 this paper was based on the model developed by Bilbao et al. (2011), suggesting the next  
9 relationship:

$$10 \quad CMF_{UVER} = CMF_{SW}^{c+d \cos(SZA)} \quad (5)$$

11 where  $c$  and  $d$  are two parameters which can be calculated by a least square fit. Bilbao et al.  
12 (2011) calculated  $c$  and  $d$  using all data measured at one location and did not take uncertainty  
13 into account in the variables of Eq. (5). It motivates this paper to improve the reconstruction  
14 method, hence a novelty in this paper is that  $c$  and  $d$  are calculated considering a similar  
15 number of data for different cloudiness and SZA conditions, and taking into account the  
16 uncertainty in these data, which optimised the mentioned method. To this end, a dataset of  
17  $CMF_{UVER}$ - $CMF_{SW}$  pairs of data was selected in this paper.

18 All available data pairs of  $CMF_{UVER}$ - $CMF_{SW}$  from all locations were divided into data groups  
19 taking into account the cosine of SZA. Eighty-seven data groups were obtained separated by  
20 intervals of the cosine of SZA, from 0.095 to 0.965 in 0.01 steps. Each data group was  
21 divided into a further 15 subgroups considering the  $CMF_{SW}$  value in the intervals from 0 to  
22 1.5 in 0.1 bins. A total of 1305 subgroups were available. Fifty pairs of  $CMF_{UVER}$ - $CMF_{SW}$   
23 data were chosen randomly for each subgroup. Some groups with high  $CMF_{SW}$  values had  
24 fewer than 50 data. Finally, 49,777 data pairs were randomly selected. This method of  
25 choosing data ensures that the number of data selected is similar and is representative for any  
26 cloudiness and SZA condition.

27 The combined uncertainty of each selected data was known; hence the  $c$  and  $d$  parameters of  
28 Eq. (5) were calculated by the weighted least square method using the 49,777 chosen data.  
29 The weight in the fit was the inverse of the square combined uncertainty. Uncertainty in SZA  
30 was considered null. The result of the fit gave a  $c$  value equal to 0.6106 with a combined  
31 uncertainty of 0.0014, and a  $d$  value equal to 0.358 with a combined uncertainty of 0.002.

1 These calculated values are different to those obtained by Bilbao et al. (2011). Once the c and  
 2 d values were calculated, the f function of Eq. (4) can be expressed as:

$$3 \quad f(CMF_{SW}) = CMF_{SW}^{0.6106 \pm 0.0014 + (0.358 \pm 0.002) \cos(SZA)} \quad (6)$$

4 UVER irradiance can be reconstructed using Eq. (6) in Eq. (4) taking into account cloudless  
 5 simulations and measured SW irradiance. This model for reconstructing UVER irradiance  
 6 was called ‘‘M-SW’’.

### 7 **3.2.3 Model based on daily sunshine fraction measurements**

8 Another variable used to reconstruct UVER radiation is sunshine fraction (Lindfors et al.,  
 9 2003), due to the availability of longer series than SW; it has the advantage to reconstruct  
 10 older data. The sunshine fraction represents the fraction of day when sun was not blocked by  
 11 clouds. Therefore, by way of an initial approach, daily direct (on horizontal) UVER  
 12 irradiation ( $UVER^{dir}$ ) can be expressed as:

$$13 \quad UVER^{dir} = UVER_{cl}^{dir} F \quad (7)$$

14 where  $UVER_{cl}^{dir}$  is the daily direct UVER irradiation simulated under cloudless conditions on  
 15 horizontal surface. Assuming Eq. (7), daily global UVER irradiation can thus be expressed as  
 16 the sum of the direct (on horizontal surface) and diffuse ( $UVER^{dif}$ ) components:

$$17 \quad UVER = UVER_{cl}^{dir} F + UVER^{dif} \quad (8)$$

18 As a second approximation, daily diffuse UVER irradiation can be assumed as a function (g),  
 19 which depends on the daily diffuse UVER under cloudless conditions ( $UVER_{cl}^{dif}$ ) and on F:

$$20 \quad UVER^{dif} = g(UVER_{cl}^{dif}, F) \quad (9)$$

21 Eq. (9) in Eq. (8) gives:

$$22 \quad UVER = UVER_{cl}^{dir} F + g(UVER_{cl}^{dif}, F) \quad (10)$$

23 Global UVER irradiation can be obtained through F measurements and cloudless simulations  
 24 using Eq. (10) once the g function is known. The g function can be calculated by F and UVER  
 25 measurements:

$$26 \quad g = \frac{UVER - UVER_{cl}^{dir} F}{UVER_{cl}^{dif}} \quad (11)$$

1 In order to calculate the  $g$  values a new dataset was chosen made up of daily UVER  
 2 irradiation and  $F$  measurements. Days without SW, UVER or  $F$  measurements were discarded  
 3 in the new dataset. Available data (all locations) were then divided by the season: winter  
 4 (January, February, and December), spring (March, April, and May), summer (June, July, and  
 5 August) and autumn (September, October, and November). 10% of the available data for each  
 6 season were randomly selected and used to obtain the  $g$  function. Figure 2 shows the values  
 7 of  $g$  calculated by Eq. (11) as a function of  $UVER_{cl}^{dif}$  for four  $F$  intervals. The  $g$  values  
 8 increase linearly with cloudless diffuse UVER irradiation, showing a correlation coefficient  
 9 ( $r$ ) above 0.87 for all  $F$  intervals. Moreover, the dependence of  $g$  on  $UVER_{cl}^{dif}$  varies with  $F$ ,  
 10 and shows a higher slope when  $F$  increases. The y-intercept of the fits is close to zero, then,  
 11 assuming that  $UVER^{dif}$  should be equal to zero when  $UVER_{cl}^{dif}$  was zero, the  $g$  function could  
 12 be expressed as:  $g = B(F) UVER_{cl}^{dif}$  (12)

13 where  $B$  is a parameter which depend on  $F$ .  $B$  was calculated through a least square fit using  
 14 the same data as in Fig. 2, but for the following  $F$  intervals:  $F=0$ ,  $0 < F < 0.2$ ,  $0.2 \leq F < 0.4$ ,  
 15  $0.4 \leq F < 0.6$ ,  $0.6 \leq F < 0.8$ ,  $0.8 \leq F < 1$ ,  $F \geq 1$ . These intervals were chosen in order to have more  
 16 resolution of  $F$  for the model development; while the intervals of Fig. 2 were wider because  
 17 they were enough to find the kind of dependence on  $F$  of  $g$  function versus  $UVER_{cl}^{dif}$ . The  
 18 values obtained are shown in Fig. 3 with their combined uncertainty as a function of  $F$ . This  
 19  $B$  parameter increases with  $F$ , and is close to 1 when  $F$  tends to 1. Uncertainty is around 0.2  
 20 except for  $F$  equal to 1, when it strongly increases up to 0.55 (caused in part by the high  
 21 uncertainty in  $UVER_{cl}^{dir}$ , which is not reduced multiplying by  $F$ ). The  $B$  value can be obtained  
 22 by interpolating the values of Fig. 3 for an  $F$  measurement. This interpolated value of  $B$  can  
 23 be used to retrieve global UVER irradiation taking into account Eq. (12) in Eq. (10):

$$24 \quad UVER = UVER_{cl}^{dir} F + B(F) UVER_{cl}^{dif} \quad (13)$$

25 The method for retrieving global UVER irradiation finding a  $B$  value by an  $F$  measurement  
 26 interpolation in Fig. 3 and using this  $B$  value with the measured  $F$  and cloudless simulations  
 27 in Eq. (13) was called “M-F” in this paper.

### 28 **3.2.4 Models vs. measurements**

29 Hourly UVER irradiance was reconstructed with model M-SW using the available data which  
 30 were not used to develop the model M-SW. Twenty-five UVER data (three days) were  
 31 rejected applying a new quality control. A total of 220,105 pairs of measured ( $UVER_{me}$ ) and

1 reconstructed ( $UVER_{mo}$ ) UVER irradiance data were available for each model considering  
2  $SZA < 80^\circ$ . These data pairs were used to quantify the agreement between UVER reconstructed  
3 by model and measured UVER irradiance. The combined and expanded uncertainty of all  
4 reconstructed values were calculated deriving the equations of the model and taking into  
5 account the uncertainty in the used measurements and cloudless simulations. Table 2 shows  
6 the MBE and the RMSE obtained in the comparison between the reconstructed and measured  
7 hourly UVER values. The model M-SW slightly overestimates the measurements showing a  
8 MBE of 1.6% ( $0.7 \text{ mWm}^{-2}$ ); the RMSE was 15.8% ( $6.2 \text{ mWm}^{-2}$ ). A depth analysis (not  
9 shown) indicated that model M-SW fit better within the measurements for  $SZA < 55^\circ$   
10 ( $RMSE < 10\%$ ); the MBE is close to 0% for  $SZA < 75^\circ$ ; the best RMSE and MBE values  
11 appears for  $CMF_{SW}$  values between 0.2 and 1.1.

12 In order to quantify the agreement between the models and the daily UVER measurements, all  
13 data not used to develop models M-F (90% of data per season) were selected. The  
14 reconstructed UVER irradiance these days was integrated to obtain the reconstructed daily  
15 UVER irradiation by model M-SW. Daily irradiance was also reconstructed by model M-F.  
16 The available number of pairs of data for reconstructed and measured daily UVER irradiation  
17 was 21,349 as can be seen in Table 2. Table 2 also shows that model M-SW fit better (with a  
18 RMSE of 8.4%) within measurements than model M-F (RMSE higher than 20%); moreover,  
19 model M-F overestimates the daily measurements showing a MBE of 5.1%. RMSE and MBE  
20 were also calculated for different F intervals (not shown) and model M-SW presents RMSE  
21 values lower than 10% and MBE closest to 0% for all F intervals; model M-F show a good  
22 agreement with measurements for F higher than 0.5 but RMSE and MBE increases when F  
23 decreases.

24 All available data were used to obtain daily UVER irradiation, and these values were monthly  
25 averaged (at least 25 days per month) for each location and year. The same monthly averages  
26 were reconstructed with the models. The MBE and RMSE obtained using the monthly  
27 averages of reconstructed and measured UVER irradiation are shown in Table 2. These values  
28 are similar for both models, being the MBE around 2% and RMSE around below 6.5%. The  
29 number of data for the models is different due to the lack of F data in some periods at Madrid  
30 in the 2000s decade. Monthly averages were used to obtain the annual averaged values  
31 (always with all 12 months available) of UVER irradiation at each location. Annual  
32 reconstructed and measured data were compared also in Table 2 showing the statistical

1 estimators of the comparison. Both models present a good agreement with measurements  
2 being RMSE lower than 3% and MBE below 1% in absolute value. 95% of the reconstructed  
3 annual UVER data with model M-SW shows a difference below 5% with the measurements.  
4 Finally, the differences between measured and reconstructed values are in agreement within  
5 uncertainty for both models and for all temporal resolutions (not shown). Model M-F  
6 measurements are not the best for reconstructing daily UVER irradiation although it  
7 evidences great agreement with monthly and annual averaged measurements. A depth  
8 comparison not shown in this paper about these models can be found in Román (2014).

### 9 **3.3 Final reconstructed series**

10 Hourly UVER irradiance was reconstructed at each location, when hourly SW records were  
11 available, using the method referred to as M-SW. Daily reconstructed UVER irradiation was  
12 obtained by integrating the hourly reconstructed values. Table 1 (M-SW column) shows the  
13 number of daily UVER irradiation data reconstructed by the model based on SW records.  
14 This number is around 10,000 (27 years) for A Coruña, Cáceres, and Murcia, and is over  
15 13,000 (36 years) for Madrid. Villalba shows the lowest number of reconstructed UVER data  
16 by this model due to the scant number of SW records at this location. When SW records were  
17 not available, UVER irradiation was reconstructed using the method M-F and based on F  
18 measurements. Table 1 (M-F column) shows the number of daily UVER irradiation data  
19 reconstructed by the model based on F records. This number varies with the location, and is  
20 higher for Villalba, San Sebastián, and Tortosa. The reconstructed daily UVER irradiation  
21 series were completed with daily measured UVER irradiation when SW and F records were  
22 not available. The number of measured data used to form the UVER series is shown in Table  
23 1 (Measured data column). This number of data is low compared to the other reconstructed  
24 data, is less than 20 at all locations, and is zero in Murcia and Villalba. Finally, a long-term  
25 daily UVER irradiation series was obtained at each location using models and measurements.  
26 The total number of data of these series and the year of the first UVER irradiation value are  
27 shown in Table 1. A Coruña, Madrid, San Sebastián, and Villalba show UVER series with  
28 more than 20,000 data, all commencing in the 1950s. The lowest number of daily UVER data  
29 is around 10,000 for Cáceres and Murcia, whose UVER series started in the mid 1980s. The  
30 combined uncertainty of all daily UVER irradiation values of the obtained series was also  
31 calculated taking into account the uncertainty in the cloudless simulations and measured  
32 values.

## 1 4 Results and discussion

### 2 4.1 Anomalies and homogeneity

3 Monthly averages of daily UVER irradiation were calculated using the available reconstructed  
4 series taking into account at least 20 daily UVER data per month, year, and location. The  
5 monthly series obtained were deseasonalized calculating their monthly anomalies considering  
6 the reference period from 1985 to 2011. UVER anomaly (A) in month “m” and year “y” is  
7 thus calculated as:

$$8 \quad A_{m,y} = UVER_{m,y} - \frac{1}{N} \sum_{y'=1985}^{2011} UVER_{m,y'} \quad (14)$$

9 where N is the number of data used in the sum of Eq. (14). Monthly UVER irradiation  
10 anomalies were calculated for all months at all locations. The monthly anomalies of UVER  
11 irradiation at the nine locations were averaged, obtaining a new monthly series of anomalies  
12 representative of the Iberian Peninsula. This was then called the “Iberian Peninsula” series.  
13 Annual UVER anomalies were calculated averaging the monthly anomalies when at least six  
14 monthly data were available for each year. Seasonal anomalies were also calculated when at  
15 least two monthly data were available in winter (December, January, and February), spring  
16 (March, April, and May), summer (June, July, and August), and autumn (September, October,  
17 and November). Winter anomalies were calculated with the January and February anomalies  
18 for a specific year, together with the December anomaly of the previous year.

19 Homogeneity of all these averaged daily UVER irradiation anomaly series was tested in a  
20 similar way to the TOC and SW series analyzed by Román et al. (2014a, d). The homogeneity  
21 tests are mainly applied in order to know whether the series are valid for trend studies or, on  
22 the contrary, they are not valid due to instrumentation/measurement problems; additionally,  
23 the homogeneity tests can provide information about break points in some climatic trends. In  
24 this case, the null hypothesis assumes that a temporal series is homogenous, and this  
25 hypothesis was verified using the Standard Normal Homogeneity Test (SNHT), the Pettitt  
26 test, the Buishand test, and the Von Neumann ratio (Wijngaard et al., 2003). Hakuba et al.  
27 (2013) assumed that a temporal series is non-homogeneous when the null hypothesis is  
28 rejected with a confidence of 99% by at least three of the four mentioned tests. The four tests  
29 were directly carried out on the annual UVER series. Eight (all except Madrid and Murcia) of  
30 the ten (nine locations plus the averaged Iberian Peninsula) annual series show non-

1 homogeneity around the mid 1980s, which could be caused by a climate change in UVER  
2 from the dimming to brightening periods. Homogeneity analysis was thus performed for the  
3 same annual UVER series for the 1950-1984 and 1985-2011 periods. The first period  
4 evidences inhomogeneities in San Sebastián and A Coruña, and the second period is free of  
5 inhomogeneities. The same result was obtained by Román et al. (2014c) for the annual SW  
6 irradiation series.

7 The monthly series was also subjected to homogeneity analysis by applying the four tests to  
8 the synthetic reference series generated with UVER data from the other locations  
9 (Alexandersson and Moberg, 1997; Sánchez-Lorenzo et al., 2013c). No monthly UVER series  
10 shows inhomogeneities for the 1985-2011 period, and only one month shows inhomogeneity  
11 in Madrid and San Sebastián in the 1950-1984 period. These results thus indicate that all the  
12 UVER anomaly series can be considered homogeneous, or at least not inhomogeneous  
13 enough to change the UVER series values.

#### 14 **4.2 UVER temporal evolution**

15 Figure 4 shows the annual UVER daily irradiation anomaly series for the nine locations and  
16 the averaged Iberian Peninsula series. Anomalies are shown with their combined uncertainty.  
17 Figure 4 panels present a 21-year Gaussian low-pass filter to reduce noise in the evolution.  
18 Moreover, the linear trends calculated by the least square method are plotted for the 1950-  
19 2011, 1950-1984, and 1985-2011 periods. Madrid shows a strong peak increasing at the  
20 beginning of 1970s while A Coruña exhibits a negative peak around 1980. These peaks look  
21 strange, but the homogeneity tests do not consider it as inhomogeneities, and they also  
22 appeared in the sunshine fraction and SW irradiation series at the same places (Román et al.,  
23 2014a). In addition, a relative peak in the daily maximum temperature series appeared in the  
24 dates periods in the same places (not shown), which evidences that the UVER peaks were real  
25 but does not explain the cause of them. All annual UVER anomalies of Fig. 4 display an  
26 increase for the periods 1950-2011, and 1985-2011, except in A Coruña. By contrast, in the  
27 period 1950-1984, the annual UVER series which contains most data shows a slight decrease.  
28 San Sebastián exhibits the clearest change in UVER in the 1980s, which explains the break  
29 point detected in this location with the homogeneity tests. Román et al. (2014a) and Román et  
30 al. (2014d) found that SW irradiation and TOC decreased until mid 1980s (the dimming  
31 period) at the same locations. These results suggest that UVER in the dimming period  
32 experienced two opposite effects: UVER tended to decrease due mainly to the increase of



1 aerosols (as in SW irradiation), and, at the same time, UVER tended to increase due to ozone  
2 depletion. The two effects offset one another, leading to a not significant change in UVER  
3 irradiation over the dimming period.

4 An example of the balanced effects between the impact of aerosol increase and ozone  
5 depletion can be seen after a major volcanic eruption during which vast amounts of gaseous  
6 compounds can be shot into the stratosphere. These are precursors for the atmospheric  
7 formation of sulphate aerosol particles which in turn provide surfaces for heterogeneous  
8 processes on polar stratospheric clouds in the lower stratosphere, enhancing ozone depletion  
9 (Peter, 1997; Solomon, 1999; Rieder et al., 2013). In sum, aerosol load increases and ozone  
10 decreases after a violent volcanic eruption. The volcanic eruptions at El Chichón (México)  
11 and Pinatubo (Philippines) in 1982 and 1991, respectively, are highlighted in Fig. 4. Román et  
12 al. (2014a) found that the years after these eruptions, a reduction in SW irradiation due to the  
13 increase in sulphate aerosol particles is apparent. However, by contrast, UVER irradiation  
14 shows little increase in most of the panels in Fig. 4. The UVER increase is caused by the  
15 strong decrease in TOC after volcanic eruptions detected at these locations by Román et al.  
16 (2014d), which leads to an increase in UVER more than a UVER decrease caused by  
17 aerosols.

### 18 **4.3 UVER trends: dimming and brightening periods**

19 Figure 4 shows the annual UVER evolution and trends in qualitative but not quantitative  
20 terms. In order to quantify them, the temporal trends of the monthly, seasonal, and annual  
21 UVER anomaly series were assumed to be the trends obtained by the Theil-Sen (TS)  
22 estimator and its 95% confidence interval (95CI). The statistical significance of each  
23 calculated trend was evaluated by the non-parametric Mann-Kendall test, considering three  
24 types of trends: with a confidence of 99% ( $p < 0.01$ ), with a confidence of 95% but not 99%  
25 ( $p < 0.05$ ), or non-significant at least at 95% confidence ( $p \geq 0.05$ ). All these estimators were  
26 calculated following the methods of Gilbert (1987). If the Mann-Kendall test considered a  
27 trend as statistically significant with at least 95% confidence, this trend was then assumed to  
28 be just significant. A trend was only calculated when a series comprised more than 10 data.  
29  $TS_{O_3}$  is the TS trend calculated with the same UVER irradiation series but simulated under  
30 cloudless conditions. The  $TS_{O_3}$  value gives the UVER trend caused by changes in TOC  
31 because aerosol and clouds changes are not included in cloudless simulations.  $TS_{ac}$  is defined

1 as TS minus  $TS_{O_3}$ , and represents the UVER trend brought about by changes in aerosol and  
2 clouds (both together).

3 The trends (and their significance and 95CI) of monthly, seasonal, and annual UVER  
4 irradiation series were calculated for all locations at three periods: 1950-1984 (considered as  
5 the dimming period), 1985-2011 (considered as the brightening period), and 1950-2011  
6 (whole period). Fig. 5 shows all these values following the methodology used by Walker  
7 (2010), who plotted the trends as points which type depends on the trend significance and he  
8 also added the 95% confidence intervals as error bars. The significant seasonal and annual  
9 trends are also shown in Table 3, which also shows  $TS_{O_3}$  and  $TS_{ac}$  trends. The most  
10 representative trends of the dimming period are those obtained for the San Sebastián, Madrid,  
11 Villalba, and Iberian Peninsula series, since they have fewer missing data. The mentioned  
12 series show no significant trends in the dimming period except San Sebastián, which  
13 evidences a significant trend of  $-211 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $-7.7\%\text{dc}^{-1}$ ) in May, and an annual trend of  $-48$   
14  $\text{Jm}^{-2}\text{dc}^{-1}$  ( $-2.8\%\text{dc}^{-1}$ ). The negative annual trend found in San Sebastián is mainly caused by  
15 changes in aerosol and clouds radiative effects rather than ozone effects since  $TS_{ac}$  is much  
16 higher than  $TS_{O_3}$ . Román et al. (2014a) found many more significant trends in SW irradiation  
17 for the same locations during the dimming period, underpinning the key role played by ozone  
18 decrease in UVER trends during the dimming period.

19 The brightening period (1985-2011) has the advantage that all UVER series are complete.  
20 UVER trends are mainly significant in summer and in the annual series. All series, except for  
21 A Coruña and Madrid, present significant trends in summer, and are  $109 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $2.5\%\text{dc}^{-1}$ )  
22 for the Iberian Peninsula series. The annual trend of the Iberian Peninsula series is  $50 \text{ Jm}^{-2}\text{dc}^{-1}$   
23 ( $2.1\%\text{dc}^{-1}$ ). The  $TS_{ac}$  and  $TS_{O_3}$  values in Table 3 reveal that UVER irradiation increased in the  
24 brightening period due to a reduction in aerosols and clouds radiative effects and in ozone  
25 effects. The ozone changes represent approximately two thirds of the trend caused by aerosol  
26 and clouds changes.

27 As regards the 1950-2011 period, the most interesting series are San Sebastián, Madrid,  
28 Villalba, and the Iberian Peninsula series for the same reason as during the dimming period.  
29 All annual UVER series trends, except Madrid, are significant at 99% (95% for Murcia), this  
30 trend being  $25 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $1.1\%\text{dc}^{-1}$ ) in the Iberian Peninsula, which indicates an increase of  $155$   
31  $\text{Jm}^{-2}$  ( $6.5\%$ ) in annual UVER irradiation over the last 62 years in the Iberian Peninsula. March  
32 presents positive and significant trends for the four most complete series, ranging from  $31 \text{ Jm}^{-2}$

1  $^2\text{dc}^{-1}$  (San Sebastián:  $2.2\%\text{dc}^{-1}$ ) to  $74 \text{ Jm}^{-2}\text{dc}^{-1}$  (Villalba:  $4.1\%\text{dc}^{-1}$ ). June and July exhibit the  
2 highest UVER trends, all of them proving significant except Madrid in July. The UVER trend  
3 in the Iberian Peninsula is  $83 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $1.9\%\text{dc}^{-1}$ ) in June and  $47 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $1.0\%\text{dc}^{-1}$ ) in July.  
4 As regards seasonal trends, all are significant in spring and summer except for spring in  
5 Madrid. The trend in the Iberian Peninsula series is  $32 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $1.2\%\text{dc}^{-1}$ ) in spring and  $54$   
6  $\text{Jm}^{-2}\text{dc}^{-1}$  ( $1.2\%\text{dc}^{-1}$ ) in summer. UVER trends in the 1950-2011 period are mainly caused by  
7 changes in TOC because these trends are similar to the obtained values of  $\text{TS}_{\text{o3}}$ ,  $\text{TS}_{\text{ac}}$  being  
8 around zero. This result indicates that aerosol and clouds presence increasing during dimming  
9 and decreasing during brightening are well compensated over the 1950-2011 period.

#### 10 **4.4 UVER trends: comparison with literature.**

11 In the previous section, UVER trends were calculated for three specific periods. Other  
12 authors, however, have calculated UVER trends for other periods in the literature. UVER  
13 trends were thus recalculated with the same UVER anomaly series but for other periods in  
14 this section in order to compare them with the results of other works.

15 Lindfors et al. (2003) reconstructed UVER irradiation at Sodankylä (Finland) between 1950  
16 and 1999 also using sunshine duration records, and reported two significant trends for this  
17 period:  $3.9\%\text{dc}^{-1}$  in March and  $-3.3\%\text{dc}^{-1}$  in July. Slightly higher significant trends were found  
18 in March for Madrid ( $5.3\%\text{dc}^{-1}$ ), Villalba ( $6.6\%\text{dc}^{-1}$ ), and the Iberian Peninsula ( $4.4\%\text{dc}^{-1}$ ) in  
19 the same period. These series do not present statistically significant trends in July for the  
20 1950-1999 period.

21 Josefsson (2006) analyzed the measured UVER at Norrköping (Sweden) between 1983 and  
22 2003, finding significant trends in UVER irradiation in spring ( $7.8\%\text{dc}^{-1}$ ), autumn ( $8.2\%\text{dc}^{-1}$ ),  
23 and in the annual series ( $5.2\%\text{dc}^{-1}$ ). Significant although lower trends were also detected in  
24 the same period in the series analyzed in this paper. The Iberian Peninsula series showed a  
25 significant trend of  $4.4\%\text{dc}^{-1}$  in spring and of  $2.7 \%\text{dc}^{-1}$  for the annual series.

26 Lindfors et al. (2007) reconstructed UVER irradiation using SW records from 1983 to 2005 in  
27 Bergen (Norway), Jokionen (Finland), Norrköping and Sodankylä, and reported a significant  
28 increase in annual UVER at Sodankylä ( $4.1\%\text{dc}^{-1}$ ). For the same period, trends in the annual  
29 UVER series at Cáceres, Murcia, and the Iberian Peninsula were similar and also significant,  
30 being  $3.2\%\text{dc}^{-1}$ ,  $2.7\%\text{dc}^{-1}$ , and  $2.7\%\text{dc}^{-1}$ , respectively.

1 den Outer et al. (2010) obtained UVER irradiation between 1980 and 2006 using different  
2 reconstruction models at eight European locations: Sodankylä, Jokionen, Norrköping,  
3 Potsdam (Germany), Lindenberg (Germany), Bilthoven (the Netherlands), Hradec Kralove  
4 (Czech Republic), and Thessaloniki (Greece). The annual UVER trends obtained by den  
5 Outer et al. (2010) range between  $2.8\%dc^{-1}$  and  $5.8\%dc^{-1}$ . The trends in the annual UVER  
6 series of this paper are all significant for the same period except for Murcia, and range  
7 between  $1.8\%dc^{-1}$  (Madrid) and  $5.3\%dc^{-1}$  (San Sebastián), with the Iberian Peninsula trend  
8 being  $3.2\%dc^{-1}$ . These results are similar to those obtained in literature.

9 Walker (2010) analyzed the trends of reconstructed UVER irradiation between 1981 and 2007  
10 at four Swiss locations: Davos, Payerne, Locarno, and Jungfrauoch, with the UVER  
11 irradiation trend proving significant in March and June for all four locations during this  
12 period. Spring and summer months present the highest number of significant trends in the  
13 UVER series of this work for the 1981-2007 period. Annual trends in Switzerland were  
14 similar to Spain (eight of the ten are significant) with values varying between  $2\%dc^{-1}$  and  
15  $4\%dc^{-1}$  from 1981 to 2007 in both countries, with  $3.0\%dc^{-1}$  being the trend in the annual  
16 Iberian Peninsula series.

17 Krzyscin et al. (2011) studied the UVER radiation observed at Belsk (Poland) between 1976  
18 and 2008, and found an annual trend of  $5.6\%dc^{-1}$ . In the same period, the trend in the annual  
19 Iberian Peninsula series was  $2.8\%dc^{-1}$ , half that of Belsk.

20 As regards UVER trends in the Iberian Peninsula obtained by other authors, three papers are  
21 well known. Bilbao et al. (2011) reconstructed UVER at Valladolid from 1991 to 2010 and  
22 found significant trends in summer ( $3.5\%dc^{-1}$ ) and autumn ( $4.1\%dc^{-1}$ ), similar to those  
23 obtained for the 1991-2010 period at Valladolid with the reconstructed series used in this  
24 paper:  $3.6\%dc^{-1}$  (summer) and  $5.9\%dc^{-1}$  (autumn). Antón et al. (2011a) reconstructed UVER  
25 irradiance at solar noon in summer from 1950 to 2000 at Badajoz (Spain) and Cáceres, and  
26 obtained a trend of  $4.9\%dc^{-1}$  for the 1979-2000 period at Cáceres. The significant UVER  
27 irradiation trend at Cáceres during the same period was  $5.2\%dc^{-1}$  using data from the present  
28 work, a similar value to that obtained by Antón et al. (2011a). This means that UVER  
29 irradiance trend at solar noon was similar to daily UVER irradiation trend at Cáceres. Finally,  
30 Ialongo et al. (2011) calculated UVER irradiation from 1979 to 2010 over the whole world  
31 using satellite images and found a UVER trend in the Iberian Peninsula of around  $2.5\%dc^{-1}$  in

1 March and October. The annual Iberian Peninsula series of the present work shows a similar  
2 trend ( $2.8\%dc^{-1}$ ) for the same period.

3 A comparison between the results obtained and those reported in the literature reveals that  
4 UVER changes in Europe have been similar, at least over the last few decades.

#### 5 **4.5 Effect of UVER uncertainty on trends**

6 The trends obtained were calculated without considering uncertainty in the UVER irradiation  
7 values, although uncertainty might influence the value and significance of trends. The effect  
8 of uncertainty on trends was studied following the method used by Román et al. (2014a, d):

9 For one specific series of UVER anomalies  $A$ , with  $N$  values  $A_i$ , and  $\sigma(A_i)$  being the  
10 uncertainty of the  $i$ -value of  $A$ , a normally distributed (centred on zero with a standard  
11 deviation of  $\sigma(A_i)$ ) random number ( $b_i$ ) is generated for each  $i$ -value. A normal distribution  
12 with a standard deviation of  $\sigma(A_i)$  is selected since it implies that the probability of finding a  
13 value  $A_i+b_i$  in the interval  $[A_i -\sigma(A_i), A_i +\sigma(A_i)]$  is about 68%, a probability that increases to  
14 95% when the interval is  $[A_i -2\sigma(A_i), A_i +2\sigma(A_i)]$ . Once the  $N$  values of  $b_i$  for all values of  $A$   
15 are obtained, a new synthetic series (SS) is formed as the sum of the original  $A$  series and the  
16 random numbers, the  $i$ -value of the SS series being equal to  $A_i+b_i$ . The SS series is physically  
17 valid since it is inside the uncertainty of the  $A$  series.

18 10,000 synthetic SS series were randomly obtained for each series analyzed in the previous  
19 section following the methodology described, and their trends and significance were  
20 calculated. The percentage of the 10,000 series whose trend is statistically significant at 95%  
21 ( $P(p<0.05)$ ), and at 99% ( $P(p<0.01)$ ), was calculated. Only the series with a value of  
22  $P(p<0.05)$  and  $P(p<0.01)$  above 95% and 99% were considered statistically significant at 95%  
23 and 99%, respectively. 55% (65 out of 119, considering the 10 locations and the 12 monthly +  
24 4 seasonal + 1 annual) of the significant trends (in 1950-2011, 1950-1984, and 1985-2011  
25 periods) with at least 95% confidence are considered significant (95% confidence) using the  
26 criterion based on uncertainties. All series not considered significant in the previous section  
27 are not significant with the uncertainty-based criterion.

28 The  $P(p<0.05)$  and  $P(p<0.01)$  values of the seasonal and annual UVER series considered  
29 statistically significant with at least 95% confidence taking into account the uncertainty are  
30 shown in Table 4. High  $P(p<0.05)$  values do not imply high  $P(p<0.01)$  values. The median  
31 and standard deviation of the 10,000 values of TS calculated from all synthetic series are also

1 included in Table 4. These trends are similar to the trends in Table 3, the differences between  
2 them being below  $5 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $0.1\%\text{dc}^{-1}$ ) in most cases, although this difference reaches  $18 \text{ Jm}^{-2}\text{dc}^{-1}$  ( $0.4\%\text{dc}^{-1}$ ) for Ciudad Real in summer in the 1985-2011 period. These differences are  
3 always below the standard deviation of the trend given in Table 4 (except for Ciudad Real in  
4 summer in the 1985-2011 period).  
5

6

## 7 **5 Factors not taken into account**

8 The results of this paper were obtained using reconstructed data series by models. However,  
9 the paper is not without certain limitations. Changes in aerosol optical depth, surface albedo  
10 or water vapour column were not considered, these variables being used in the radiative  
11 transfer model as monthly climatology.

12 The lack of AOD data before 2000 led to use a climatological table which does not contain  
13 the aerosol changes in the 1950-2011 period. However the aerosol effect is included the SW  
14 and F measurements (like clouds) and, as a first approximation, the reconstruction models  
15 transfer this effect to the UVER radiation. This approximation is not valid for the case of  
16 water vapour because it affects SW and F but not UVER.

17 If the water vapour column had increased, cloudless SW irradiation would have decreased  
18 and, therefore, reconstructed UVER should be higher. Daily water vapour column trends were  
19 calculated between 1957 and 2002 in the Iberian Peninsula using the daily ERA-40 data  
20 (Uppala et al., 2005; Lindfors et al., 2007). These trends (not shown) indicate a slight water  
21 vapour decrease in recent decades, which did not always prove significant, but which might  
22 contribute to reducing the UVER trends obtained.

23 Trends in surface albedo at 360 nm (Sect. 2.2) from 1958 to 2002 were calculated (not  
24 shown), revealing that, apart from a slight decrease in winter months, surface albedo has  
25 suffered no significant changes in recent decades. This result indicates that the UVER trend  
26 obtained in winter might be slightly lower due to changes in albedo, but should not affect the  
27 remaining months.

28 A further factor to take into account should be the uncertainty of the data used (caused in part  
29 by the monthly variability of the radiative transfer inputs), since certain trends cannot be  
30 considered significant when uncertainty is taken into account, as can be seen in Sect. 4.5.  
31 Moreover, the averaged Iberian Peninsula anomaly series was calculated using nine locations,

1 with only four or five locations having data available for the years prior to 1970. This number  
2 of locations might not be sufficient to obtain a representative averaged result for the Iberian  
3 Peninsula. However, by way of an initial approximation, the number of locations was  
4 considered representative since when using the same locations, Román et al. (2014a) obtained  
5 similar results in SW irradiation to Sánchez-Lorenzo et al. (2013a), who used more locations  
6 to obtain an averaged Iberian Peninsula series.

7

## 8 **6 Conclusions**

9 UVER irradiation in the Iberian Peninsula increased by  $2.1\%dc^{-1}$  (annual) and  $2.5\%dc^{-1}$   
10 (summer) for the 1985-2011 period, and  $1.1\%dc^{-1}$  (annual) and  $1.3\%dc^{-1}$  (summer) for the  
11 1950-2011 period. The amount of ozone in the atmosphere is returning to pre-1980 levels due  
12 to the reduction in halogen gases subsequent to the Montreal Protocol. This decrease supports  
13 the belief that increased UVER radiation over the past 27 years has in large part been due to a  
14 reduction in the release of aerosols into the atmosphere, also reducing the cloud presence  
15 (brightening). However, increased UVER was mainly caused by ozone depletion during the  
16 1950-2011 period. A significant UVER trend can become non-significant when uncertainty in  
17 the reconstructed data is taken into account.

## 18 **Acknowledgements**

19 The authors gratefully acknowledge the financial support extended by the Spanish Ministry of  
20 Science and Innovation for project CGL2011-25363. The authors also thank the staff at the  
21 AEMet for the data used and for their effort in establishing and maintaining the stations. The  
22 authors gratefully thank the OMI, TOMS, GOME, GOME-2, MODIS, and MISR teams for  
23 the satellite data used in this study, as well as the staff of the COST-726 project for the  
24 reconstructed ozone data. The NASA GES DISC and the AVDC are also acknowledged for  
25 the GIOVANNI application and the satellite overpass files. Roberto Román would like to  
26 thank the University of Valladolid for its support through the PIF-UVa grant.

27

28

29

30

## 1 **References**

- 2 Acker, J. G. and Leptoukh, G.: Online Analysis Enhances Use of NASA Earth Science Data,  
3 Eos, Trans. AGU, 88, 2 14-17, 2007.
- 4 Alexandersson, H. and Moberg, A.: Homogenization of Swedish temperature data. Part I:  
5 Homogeneity test for linear trends, Int. J. Climatol., 17, 25–34, 1997.
- 6 Anderson, G., Clough, S., Kneizys, F., Chetwynd, J., and Shettle, E.: AFGL atmospheric  
7 constituent profiles (0-120 km), Tech. Rep. AFGL-TR-86-0110, Air Force Geophys. Lab.,  
8 Hanscom Air Force Base, Bedford, Mass., 1986.
- 9 Antón, M., Serrano, A., Cancillo, M. L., García, J. A., and Madronich, S.: Application of an  
10 analytical formula for UV Index reconstructions for two locations in Southwestern Spain,  
11 Tellus, 63B, 1052–1058, 2011a.
- 12 Antón, M., Serrano, A., Cancillo, M. L., García, J.A., and Madronich, S.: Empirical  
13 evaluation of a simple analytical formula for the ultraviolet index, Photochem. Photobiol., 87,  
14 478–482, 2011b.
- 15 Antón, M., and Mateos, D: Shortwave radiative forcing due to long-term changes of total  
16 ozone column over the Iberian Peninsula, Atmos. Environ., 81, 532-537, doi:  
17 10.1016/j.atmosenv.2013.09.047, 2013.
- 18 Bilbao, J., Román, R., de Miguel, A., and Mateos, D.: Long-term solar erythemal UV  
19 irradiance data reconstruction in Spain using a semiempirical method, J. Geophys. Res., 116,  
20 D22211, 2011.
- 21 Blumthaler, M. and Ambach, W.:\_Indication of increasing solar ultraviolet-B radiation flux in  
22 alpine region, Science, 248, 206-208, 1990.
- 23 Bodeker, G. E. and McKenzie, R.L.: An algorithm for inferring surface UV irradiance  
24 including cloud effects, J. Appl. Meteorol., 35, 10, 1860–1877, 1996.
- 25 Borkowski, J. L.: Homogenization of the Belsk UV-B series (1976-1997) and trend analysis,  
26 J. Geophys. Res., 105, D4, 4873–4878, 2000.
- 27 Buras, R., Dowling, T., and Emde, C.: New secondary-scattering correction in DISORT with  
28 increased efficiency for forward scattering, J. Quant. Spectrosc. Ra., 112, 2028–2034,  
29 doi:10.1016/j.jqsrt.2011.03.019, 2011.



- 1 Calbó, J., Pagès, D., and González, J. A.: Empirical studies of cloud effects on UV radiation:  
2 A review, *Rev. Geophys.*, 43, 2, 1-28, 2005.
- 3 Correl, D. L., Clark, O., Goldberg, B., Goodrich, V. R., Hayes Jr., D. R., Klein, W. H., and  
4 Schecher, W. D.: Spectral ultraviolet-B radiation at the Earth's surface: Long-term variations  
5 at 39°N, 77°W, *J. Geophys. Res.*, 97, 7579-7591, 1992.
- 6 de Miguel, A., Bilbao, J., Román, R., and Mateos, D.: Measurements and attenuation of  
7 erythemal radiation in Central Spain, *Int. J. Climatol.*, 32, 929–940, 2012.
- 8 den Outer, P. N., Slaper, H., Matthijsen, J., Reinen, H. A. J. M., and Tax, R.: Variability of  
9 ground-level ultraviolet: Model and measurement, *Radiat. Prot. Dosim.*, 91, 105-110, 2000.
- 10 den Outer, P. N., Slaper, H., and Tax, R. B.: UV radiation in the Netherlands: Assessing long-  
11 term variability and trends in relation to ozone and clouds, *J. Geophys. Res.*, 110, D02203,  
12 2005.
- 13 den Outer, P. N., Slaper, H., Kaurola, J., Lindfors, A., Kazantzidis, A., Bais, A. F., Feister, U.,  
14 Junk, J., Janouch, M., and Josefsson, W.: Reconstructing of erythemal ultraviolet radiation  
15 levels in Europe for the past 4 decades, *J. Geophys. Res.*, 115, D10102, 2010.
- 16 Gilbert, R. O.: Statistical methods for environmental pollution monitoring, Van Nostrand  
17 Company, Hoboken, N. J., 320 pp, 1987.
- 18 Hakuba, M. Z., Sánchez-Lorenzo, A., Folini, D., and Wild, M.: Testing the homogeneity of  
19 short-term surface solar radiation series in Europe, *AIP Conf. Proc.*, 1531, 700, 2013.
- 20 Hülsen, G. and Gröbner, J.: Characterization and calibration of ultraviolet broadband  
21 radiometers measuring erythemal weighted irradiance, *Appl. Opt.*, 26, 23, 5877-5886, 2007.
- 22 Ialongo, I., Arola, A., Kujanpää, J., and Tamminen, J.: Use of satellite erythemal UV products  
23 in analysing the global UV changes, *Atmos. Chem. Phys.*, 11, 9649–9658, 2011.
- 24 Iqbal, M.: An introduction to solar radiation, Academic Press, 0-12-373750-8, 1983.
- 25 Josefsson, W.: UV-radiation 1983-2003 measured at Norrköping, Sweden, *Theor. Appl.*  
26 *Climatol.*, 83, 59-76, 2006.
- 27 Kaurola, J., Taalas, P., Koskela, T., Borkowski, J., and Josefsson, W.: Long-term variations of  
28 UV-B doses at three stations in northern Europe, *J. Geophys. Res.*, 105, D16, 20813–20820,  
29 2000.

1 Krzyscin, J. W.: Statistical reconstruction of daily total ozone over Europe 1950 to 2004, *J.*  
2 *Geophys. Res.*, 113, D07112, 2008.

3 Krzyscin, J. W., Sobolewski, P. S., Jarosławski, J., Podgórski, J., and Rajewska-Wiech, B.:  
4 Erythemal UV observations at Belsk, Poland, in the period 1976–2008: data homogenization,  
5 climatology, and trends, *Acta Geophys.*, 59, 155–182, 2011.

6 Kurucz, R.: Synthetic infrared spectra, Proceedings of the 154th Symposium of the  
7 International Astronomical Union (IAU), 1992.

8 Kylling, A., Stamnes, K., and Tsay, S. C.: A reliable and efficient two-stream algorithm for  
9 spherical radiative transfer: Documentation of accuracy in realistic layered media, *J. Atmos.*  
10 *Chem*, 21, 115-150, 1995.

11 Lindfors, A., Arola, A., Kaurola, J., Arola, A., Taalas, P., and Svenoe, T.: Long-term  
12 erythemal UV doses at Sodankylä estimated using total ozone, sunshine duration, and snow  
13 depth, *J. Geophys. Res.*, 108 (D16), 4518, 2003.

14 Lindfors, A., Kaurola, J., Arola, A., Koskela, T., Lakkala, K., Josefson, W., Olseth, J. A., and  
15 Johnsen, B.: A method for reconstruction of past UV radiation based on radiative transfer  
16 modeling: Applied to four stations in northern Europe, *J. Geophys. Res.*, 112, D23201, 2007.

17 Mateos, D., Antón, M., Sanchez-Lorenzo, A., Calbó, J., and Wild, M.: Long-term changes in  
18 the radiative effects of aerosols and clouds in a mid-latitude region (1985–2010), *Global and*  
19 *Planetary Change*, 111, 288–295, doi: 10.1016/j.gloplacha.2013.10.004, 2013.

20 Mateos, D., Sanchez-Lorenzo, A., Antón, M., Cachorro, V. E., Calbó, J., Costa, M. J., Torres,  
21 B., and Wild, M.: Quantifying the respective roles of aerosols and clouds in the strong  
22 brightening since the early 2000s over the Iberian Peninsula, *J. Geophys. Res.*, 119, doi:  
23 10.1002/2014JD022076, 2014.

24 Matthijsen, J., Slaper, H., Reinen, H. A. J. M., and Velders, G. J. M.: Reduction of solar UV  
25 by clouds: A comparison between satellite-derived cloud effects and ground-based radiation  
26 measurements, *J. Geophys. Res.*, 105, D4, 5069-5080, 2000.

27 Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative  
28 transfer calculations – description and examples of use, *Atmos. Chem. Phys.*, 5, 1855–1877,  
29 2005.

1 McKinlay, A. F. and Diffey, B. L.: A reference action spectrum for ultraviolet induced  
2 erythema in human skin, *Commission Internationale de l' Eclairage (CIE)*, 6, 17–22, 1987.

3 Moreta J. R., García, R., Martín, L., Montero, J., San Atanasio, J. M., Hernández, J. L., Díaz,  
4 A., Vicente, R., C.R.N., and López, M.: AEMet contribution to the WMO/GAW programme,  
5 GAW 2013 Symposium, WMO Secretariat, Geneva, 2013.

6 Peter, T.: Microphysics and heterogeneous chemistry of polar stratospheric clouds, *Annu.*  
7 *Rev. Phys. Chem.*, 48, 785–822, 1997.

8 Piedehierro, A. A., Antón, M., Cazorla, A., Alados-Arboledas, L., Olmo, F. J.: Evaluation of  
9 enhancement events of total solar irradiance during cloudy conditions at Granada  
10 (Southeastern Spain), *Atmos. Res.*, 135:136, 1-7, 2014.

11 Ricchiazzi, P., Yang, S., Gautier, C., and Sowle, D.: SBDART: A research and Teaching  
12 software tool for plane-parallel radiative transfer in the Earth's atmosphere, *Bulletin of the*  
13 *American Meteorological Society*, 79, 2101–2114, 1998.

14 Rieder, H. E., Holawe, F., Simic, S., Blumthaler, M., Krzyscin, J. W., Wagner, J. E.,  
15 Schmalwieser, A. W., and Weihs, P.: Reconstruction of erythemal UV-doses for two stations  
16 in Austria: a comparison between alpine and urban regions, *Atmos. Chem. Phys.*, 8, 6309–  
17 6323, 2008.

18 Rieder, H. E., Frossard, L., Ribatet, M., Staehelin, J., Maeder, J. A., Di Rocco, S., Davison, A.  
19 C., Peter, T., Weihs, P., and Holawe, F.: On the relationship between total ozone and  
20 atmospheric dynamics and chemistry at mid-latitudes – Part 2: The effects of the El  
21 Niño/Southern Oscillation, volcanic eruptions and contributions of atmospheric dynamics and  
22 chemistry to long-term total ozone changes, *Atmos. Chem. Phys.*, 13, 165–179, 2013.

23 Román, R.: Reconstrucción y análisis de la radiación ultravioleta eritemática en la Península  
24 Ibérica desde 1950, PhD Thesis, Dep. of Appl. Phys., Univ. of Valladolid, Valladolid, Spain,  
25 2014.

26 Román, R., Bilbao, J., and de Miguel, A.: Reconstruction of six decades of daily total solar  
27 shortwave irradiation in the Iberian Peninsula using sunshine duration records, *Atmos.*  
28 *Environ.*, 99, 41–50, 2014a.

1 Román, R., Bilbao, J., and de Miguel, A.: Solar radiation simulations in the Iberian Peninsula:  
2 Accuracy and sensitivity to uncertainties in inputs of a radiative transfer model, *J. Quant.*  
3 *Spectrosc. Ra.*, 145, 95–109, doi: [10.1016/j.jqsrt.2014.04.028](https://doi.org/10.1016/j.jqsrt.2014.04.028), 2014b.

4 Román, R., Bilbao, J., and de Miguel, A.: Uncertainty and variability in satellite-based water  
5 vapor column, aerosol optical depth and Angström Exponent, and its effect on radiative  
6 transfer simulations in the Iberian Peninsula, *Atmos. Environ.*, 89, 556-569, 2014c.

7 Román, R., Bilbao, J., and de Miguel, A.: Uncertainty of different atmospheric ozone  
8 retrievals and its effect on temporal trends and on a radiative transfer simulations in the  
9 Iberian Peninsula, *J. Geophys. Res. Atmos.*, 119, published online,  
10 doi:10.1002/2013JD021260., 2014d.

11 Sabburg, J. and Parisi, A. V.: Spectral dependency of cloud enhanced UV irradiance, *Atmos.*  
12 *Res.* 81, 206–214, 2006.

13 Sabburg, J., and Calbó, J.: Five years of cloud enhanced surface UV radiation measurements  
14 at two sites (in the Northern and Southern Hemispheres), *Atmos. Res.*, 93, 4, 902-912, 2009.

15 Sánchez-Lorenzo, A., Brunetti, M., Calbó, J., and Martin-Vide, J.: Recent spatial and  
16 temporal variability and trends of sunshine duration over the Iberian Peninsula from a  
17 homogenized data set, *J. Geophys. Res.*, 112, D20115, 2007.

18 Sánchez-Lorenzo, A., Calbó, J., and Wild, M.: Global and diffuse solar radiation in Spain:  
19 Building a homogeneous dataset and assessing their trends, *Global Planet. Change*, 100, 343-  
20 352, 2013a.

21 Sánchez-Lorenzo, A., Wild, M., Guijarro, J. A., Brunetti, M., Bartok, B., Mystakidis, S.,  
22 Hakuba, M., and Müller, G.: Reassessment and update of the trends in the surface solar  
23 radiation over Europe by means of homogenized series from the GEBA, European  
24 Geophysical Union (EGU) General Assembly 2013, 2013b.

25 Sánchez-Lorenzo, A., Wild, M., and Trentmann, J.: Validation and stability assessment of the  
26 monthly mean CM SAF surface solar radiation dataset over Europe against a homogenized  
27 surface dataset (1983-2005), *Rem. Sens. Environ.*, 134 355-366, 2013c.

28 Schaaf, C. B., et al.: First Operational BRDF, Albedo and Nadir Reflectance Products from  
29 MODIS, *Remote Sens. Environ.*, 83, 135-148, 2002.

1 Schwander, H., Mayer, B., Ruggaber, A., Albold, A., Seckmeyer G., and Koepke, P.: Method  
2 to determine snow albedo values in the UV for radiative transfer modelling, *Appl. Optics*, 38,  
3 18, 3869–3875, 1999.

4 Scotto, J., Cotton, G., Urbach, F., Berger, D., and Fears, T.: Biologically effective ultraviolet  
5 radiation: Surface measurement in the United States, 1974 to 1985, *Science*, 239, 762–764,  
6 1988.

7 Shettle, E.: Models of aerosols, clouds and precipitation for atmospheric propagation studies,  
8 Atmospheric propagation in the uv, visible, ir and mm-region and related system aspects, 454  
9 in AGARD Conference Proceedings, 1989.

10 Solomon, S.: Stratospheric ozone depletion: a review of concepts and history, *Rev. Geophys.*,  
11 37, 275–316, 1999.

12 Stanhill, G. and Cohen, S.: Global dimming: A review of the evidence for a widespread and  
13 significant reduction in global radiation, *Agric. For. Meteorol.*, 107, 255-278, 2001.

14 Tanskanen, A.: Lambertian Surface Albedo Climatology at 360 nm from TOMS Data Using  
15 Moving Time-Window Technique, Proceedings of the XX Quadrennial Ozone Symposium,  
16 2004.

17 UNEP (United Nations Environment Programme): UNEP assessment reports: Environmental  
18 effects of ozone depletion and its interactions with climate change: 2002 assessment,  
19 *Photochem. Photobiol. Sci.*, 2, 1-72, 2003.

20 Uppala, S., et al.: The ERA-40 re-analysis, *Q. J. R. Meteorol. Soc.*, 131, 2961-3012, 2005.

21 Van Hoosier, M. E., Bartoe, J.-D. F., Brueckner, G. E., and Prinz, D. K.: Absolute solar  
22 spectral irradiance 120 nm - 400 nm (Results from the Solar Ultraviolet Spectral Irradiance  
23 Monitor - SUSIM- Experiment on board Spacelab 2), *Astro. Lett. and Communications*, 27,  
24 163-168, 1988.

25 Vilaplana, J. M., Cachorro, V. E., Sorribas, M., Luccini, E., de Frutos, A. M., Berjón, A., and  
26 de la Morena, B.: Modified calibration procedures for a yankee environmental system UVB-1  
27 biometer based on spectral measurements with a brewer spectrophotometer, *Photochem.*  
28 *Photobiol.*, 82, 508–514, 2009.

29 Walker, D.: Cloud effects on erythemal UV radiation in a complex topography, PhD Thesis,  
30 *Veröffentlichungen der MeteoSchweiz*, 86, 106 p, ISSN: 1422-1381, 2010.

1 Webb, A. R.: Who, what, where and when-influences on cutaneous vitamin D synthesis, *Prog.*  
2 *Biophys. Mol. Biol.*, 92, 17–25, 2006.

3 Webb, A., Gröbner, J., and Blumthaler, M.: A Practical Guide to Operating Broadband  
4 Instruments Measuring Erythemally Weighted Irradiance, COST726, 22595, 92-898-0032-1,  
5 2006.

6 WHO (World Health Organization): Protection against exposure to ultraviolet radiation,  
7 Technical Report WHO/EHG 17, 1995.

8 WHO (World Health Organization): Global Solar UV Index: A Practical Guide, 28 pp., ISBN  
9 92-4-159007-6, Geneva, Switzerland, 2002.

10 Wijngaard, J. B., Klein-Tank, A. M. G., and Können, G. P.: Homogeneity of 20th century  
11 European daily temperature and precipitation series, *Int. J. Climatol.*, 23, 679-692, 2003.

12 Wild, M.: Global dimming and brightening: A review, *J. Geophys. Res.*, 114, D00D16, 2009.

13 Wild, M.: Enlightening global dimming and brightening, *Bull. Amer. Meteor. Soc.*, 93, 27–  
14 37, 2012.

15 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis,  
16 A., Russak, V., and Tsvetkov, A.: From dimming to brightening: Decadal changes in solar  
17 radiation at Earth’s surface, *Science*, 308, 847-850, 2005.

18 WMO (World Meteorological Organization): Guide to meteorological instruments and  
19 methods of observation, 7th Edn. WMO Publication 8, Geneva, Switzerland, 2008.

20 WMO (World Meteorological Organization): WMO Scientific Assessment of Ozone  
21 Depletion: 2010, Global Ozone Research and Monitoring Project, Report No 52, World  
22 Meteorological Organization, Geneva, Switzerland, 2011.

23  
24  
25  
26  
27  
28

1 **Tables**

2 Table 1: Coordinates of the AEMet stations used, and number of data of the final  
3 reconstructed series which were filled with M-SW model, M-F model or with measurements.  
4 The total number of data of each final reconstructed series and the year when it began are  
5 included.

Location	Latitude	Longitude	Altitude (m.s.l)	M-SW data	M-F data	Measured data	Total data	First year
Ciudad Real	38°59'21''N	3°55'13''W	628	5717	9300	6	15023	1970
San Sebastián	43°18'23''N	2°02'28''W	251	7428	15029	9	22466	1950
A Coruña	43°21'57''N	8°25'17''W	58	9600	11388	2	20990	1951
Madrid	40°27'06''N	3°43'27''W	664	13208	9158	7	22373	1950
Cáceres	39°28'17''N	6°20'20''W	394	9517	1054	7	10578	1983
Murcia	38°00'07''N	1°10'15''W	61	10035	101	0	10136	1984
Tortosa	40°49'14''N	0°29'29''E	44	5081	12476	15	17572	1954
Valladolid	41°39'00''N	4°46'00''W	735	6813	7139	16	13968	1973
Villalba	41°48'50''N	4°55'48''W	840	3712	18180	0	21892	1951

6

7

8

9

10

11

12

13

14

15

1 Table 2: Statistical estimators calculated with N pairs of measured and reconstructed UVER  
 2 radiation for M-SW and M-F models and for different temporal resolutions. The hourly values  
 3 were obtained using data with SZA below 80°.

Resolution	Model	N	MBE (%)	RMSE (%)
Hourly	M-SW	220105	1.6	15.8
Daily	M-SW	21349	1.5	8.4
Daily	M-F	21349	5.1	24.7
Monthly	M-SW	845	1.8	5.2
Monthly	M-F	783	2.3	6.4
Annual	M-SW	38	0.3	2.9
Annual	M-F	35	-0.6	2.6

4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19



1 Table 3: Statistically significant UVER irradiation trends with a confidence of 99% (95%  
2 marked with an asterisk) and their 95% confidence interval, at different seasons and locations  
3 for the 1950-2011, 1950-1984, and 1985-2011 periods.  $TS_{O_3}$  and  $TS_{ac}$  are also included. N is  
4 the number of data used.


Location	Period	Season	N	TS ( $Jm^{-2}dc^{-1}$ )	TS (% $dc^{-1}$ )	$TS_{O_3}$ ( $Jm^{-2}dc^{-1}$ )	$TS_{ac}$ ( $Jm^{-2}dc^{-1}$ )	95CI ( $Jm^{-2}dc^{-1}$ )
Ciudad Real	1950-2011	Spring	41	78	2.59	64	14	(15:140)
Ciudad Real	1950-2011	Summer	41	90	1.88	46	44	(26:153)
Ciudad Real	1950-2011	Annual	41	49	1.86	30	19	(19:76)
Ciudad Real	1985-2011	Summer	27	207	4.31	61	146	(52:310)
Ciudad Real	1985-2011	Annual	27	71	2.67	28	43	(23:123)
San Sebastián	1950-2011	Spring	62	42	1.98	42	0	(9:75)
San Sebastián	1950-2011	Summer	61	59	1.81	59	0	(23:92)
San Sebastián*	1950-2011	Autumn	62	13	0.96	4	9	(-2:27)
San Sebastián	1950-2011	Annual	62	28	1.56	29	-1	(10:46)
San Sebastián*	1950-1984	Winter	35	-15	-2.97	2	-17	(-27:1)
San Sebastián	1950-1984	Spring	35	-109	-5.40	-33	-76	(-173:-31)
San Sebastián	1950-1984	Annual	35	-48	-2.76	-5	-43	(-79:-15)
San Sebastián*	1985-2011	Spring	27	100	4.46	64	36	(-1:202)
San Sebastián	1985-2011	Summer	27	164	4.85	88	76	(58:269)
San Sebastián*	1985-2011	Annual	27	68	3.68	45	23	(3:114)
A Coruña*	1950-2011	Spring	57	34	1.47	30	4	(1:71)
A Coruña	1950-2011	Summer	56	67	1.85	41	26	(27:111)
A Coruña	1950-2011	Annual	58	28	1.41	20	8	(11:42)
A Coruña*	1950-1984	Spring	30	-86	-3.93	-21	-65	(-164:9)
A Coruña*	1950-1984	Annual	31	-34	-1.82	-1	-33	(-76:5)
Madrid*	1950-2011	Summer	61	33	0.71	40	-7	(2:61)
Cáceres*	1950-2011	Spring	29	94	3.04	46	48	(11:206)
Cáceres	1950-2011	Summer	29	173	3.56	66	107	(56:275)
Cáceres	1950-2011	Annual	29	87	3.29	29	58	(41:125)
Cáceres	1985-2011	Summer	27	163	3.35	54	109	(23:295)
Cáceres	1985-2011	Annual	27	82	3.05	24	58	(32:120)
Murcia	1950-2011	Summer	28	138	3.04	2	136	(31:218)
Murcia*	1950-2011	Annual	28	44	1.71	-6	50	(-11:96)
Murcia	1985-2011	Summer	27	137	3.03	-6	143	(29:226)
Tortosa	1950-2011	Spring	48	48	1.78	51	-3	(10:79)
Tortosa	1950-2011	Summer	48	63	1.50	54	9	(29:99)
Tortosa	1950-2011	Annual	48	34	1.48	31	3	(15:54)
Tortosa	1985-2011	Summer	27	111	2.61	66	45	(23:205)
Valladolid	1950-2011	Winter	38	24	3.62	13	11	(5:45)
Valladolid	1950-2011	Spring	38	103	3.70	78	25	(43:174)
Valladolid	1950-2011	Summer	38	139	3.08	68	71	(53:193)
Valladolid	1950-2011	Annual	38	68	2.84	40	28	(36:97)
Valladolid*	1985-2011	Summer	27	137	2.98	65	72	(-11:280)
Valladolid*	1985-2011	Annual	27	56	2.28	30	26	(2:111)
Villalba	1950-2011	Winter	61	17	2.52	13	4	(6:29)
Villalba	1950-2011	Spring	61	47	1.70	47	0	(10:85)
Villalba	1950-2011	Summer	60	53	1.16	50	3	(19:84)
Villalba	1950-2011	Annual	61	30	1.25	30	0	(13:47)
Villalba*	1985-2011	Summer	27	130	2.79	63	67	(6:257)
Villalba*	1985-2011	Autumn	27	83	4.94	-1	84	(6:128)
Villalba*	1985-2011	Annual	27	63	2.54	31	32	(2:125)
Iberian Peninsula	1950-2011	Spring	62	33	1.19	38	-5	(4:62)
Iberian Peninsula	1950-2011	Summer	62	54	1.25	46	8	(26:78)
Iberian Peninsula	1950-2011	Annual	62	25	1.05	24	1	(12:38)
Iberian Peninsula	1985-2011	Summer	27	109	2.47	46	63	(24:206)
Iberian Peninsula*	1985-2011	Annual	27	50	2.09	19	31	(6:91)

1 Table 4: Statistically significant UVER irradiation trends considered as the median of 10,000  
2 trends (standard deviation in parenthesis), with a confidence of 99% (95% marked with an  
3 asterisk), and the P(p<0.05) and P(p<0.01) values at various locations and seasons for the  
4 1950-2011, 1950-1984, and 1985-2011 periods.

Location	Period	Season	TS (Jm <sup>-2</sup> dc <sup>-1</sup> )	TS (%dc <sup>-1</sup> )	P(p<0.05) (%)	P(p<0.01) (%)
Ciudad Real*	1950-2011	Spring	78 (5.9)	2.59 (0.19)	99.98	73.97
Ciudad Real*	1950-2011	Summer	88 (8.8)	1.84 (0.19)	99.99	91.96
Ciudad Real	1950-2011	Annual	48 (3.2)	1.81 (0.12)	100	99.98
Ciudad Real*	1985-2011	Summer	189 (17.9)	3.92 (0.37)	99.99	91.40
Ciudad Real*	1985-2011	Annual	73 (5.7)	2.75 (0.21)	100	98.52
San Sebastián*	1950-2011	Spring	41 (3.0)	1.95 (0.14)	99.98	57.53
San Sebastián	1950-2011	Summer	60 (4.5)	1.83 (0.14)	100	99.97
San Sebastián	1950-2011	Annual	28 (1.5)	1.54 (0.08)	100	100
San Sebastián*	1950-1984	Spring	-106 (9.2)	-5.24 (0.46)	100	99.01
San Sebastián*	1950-1984	Annual	-46 (4.4)	-2.63 (0.25)	100	95.18
San Sebastián	1985-2011	Summer	161 (12.7)	4.76 (0.38)	100	99.46
San Sebastián*	1985-2011	Annual	67 (4.4)	3.56 (0.24)	99.19	21.76
A Coruña	1950-2011	Summer	68 (4.3)	1.86 (0.12)	100	100
A Coruña	1950-2011	Annual	27 (1.4)	1.39 (0.07)	100	100
Cáceres*	1950-2011	Spring	95 (8.3)	3.06 (0.27)	100	30.58
Cáceres	1950-2011	Summer	177 (12.1)	3.65 (0.25)	100	99.96
Cáceres	1950-2011	Annual	88 (3.5)	3.28 (0.13)	100	100
Cáceres*	1985-2011	Summer	168 (14.0)	3.45 (0.29)	99.97	83.57
Cáceres	1985-2011	Annual	79 (4.4)	2.96 (0.16)	100	100
Murcia*	1950-2011	Summer	136 (10.2)	3.00 (0.23)	100	94.61
Murcia*	1985-2011	Summer	135 (10.9)	2.99 (0.24)	99.99	78.76
Tortosa*	1950-2011	Spring	46 (4.1)	1.72 (0.16)	99.51	51.72
Tortosa	1950-2011	Summer	65 (6.0)	1.56 (0.14)	100	99.91
Tortosa	1950-2011	Annual	34 (2.1)	1.48 (0.09)	100	100
Valladolid*	1950-2011	Winter	24 (2.2)	3.62 (0.34)	99.93	69.07
Valladolid	1950-2011	Spring	103 (7.2)	3.71 (0.26)	100	99.98
Valladolid	1950-2011	Summer	130 (10.3)	2.87 (0.23)	100	99.94
Valladolid	1950-2011	Annual	68 (3.2)	2.83 (0.13)	100	100
Villalba	1950-2011	Winter	17 (1.4)	2.57 (0.20)	100	99.78
Villalba*	1950-2011	Spring	47 (4.2)	1.68 (0.15)	99.80	62.46
Villalba*	1950-2011	Summer	55 (7.1)	1.20 (0.16)	99.79	91.98
Villalba	1950-2011	Annual	31 (2.3)	1.26 (0.10)	100	99.99
Iberian Peninsula*	1950-2011	Spring	33 (1.9)	1.22 (0.07)	100	41.45
Iberian Peninsula	1950-2011	Summer	54 (2.7)	1.25 (0.06)	100	100
Iberian Peninsula	1950-2011	Annual	25 (0.9)	1.05 (0.04)	100	100
Iberian Peninsula*	1985-2011	Summer	111 (6.1)	2.53 (0.14)	100	94.36
Iberian Peninsula*	1985-2011	Annual	50 (2.1)	2.06 (0.09)	100	36.93

5  
6  
7  
8  
9

1 **Figures**



2

3 Figure 1: Distribution of selected Spanish stations located in the Iberian Peninsula.

4

5

6

7

8

9

10

11

12

13


14

15

16

17

18



1

2 Figure 2: the  $g$  function against the simulated diffuse UVER under cloudless conditions for  
 3 four sunshine fraction (F) intervals. A linear fit calculated with the N data, and its equation  
 4 and correlation coefficient (r) are included in each panel.

5

6


7

8

9

10

11



1

2 Figure 3: Parameters a (left), b (middle), and B (right) calculated for different sunshine  
 3 fraction (F) intervals. The error bar represents the combined uncertainty.

4

5

6

7

8

9

10

11

12

13

14

15


16

17

18

19

20



1

2 Figure 4: Evolution of the annual UVER irradiation anomalies and their uncertainties for ten  
 3 series. The red line corresponds to a linear fit between 1950 and 2011, and green lines to  
 4 linear fits in the 1950-1984 and 1985-2011 periods. The solid black line is a 21-year Gaussian  
 5 low-pass filter, and the years 1982 and 1991 are marked with a dashed black line.

6

7


8

9

10

11

12



1  
 2 Figure 5: UVER trends for different months, seasons, and for annual and three periods. The  
 3 error bars are the 95% confidence interval and their colour represents the location of the  
 4 legend. The green circles represent statistically significant trends with 99% confidence  
 5 ( $p < 0.01$ ), yellow squares represent statistically significant trends with 95% confidence  
 6 ( $p < 0.05$ ), and red triangles represent non-statistically significant trends with at least 95%  
 7 confidence.

8