



Climate-relevant  
physical properties  
of molecular  
constituents

M. A. Upshur et al.

# Climate-relevant physical properties of molecular constituents relevant for isoprene-derived secondary organic aerosol material

M. A. Upshur<sup>1</sup>, B. F. Strick<sup>1</sup>, V. F. McNeill<sup>2</sup>, R. J. Thomson<sup>1</sup>, and F. M. Geiger<sup>1</sup>

<sup>1</sup>Department of Chemistry, Northwestern University, Evanston, IL 60202, USA

<sup>2</sup>Department of Chemical Engineering, Columbia University, New York, NY 10027

Received: 30 April 2014 – Accepted: 15 June 2014 – Published: 27 June 2014

Correspondence to: V. F. McNeill (vfm2103@columbia.edu), R. J. Thomson (r-thomson@northwestern.edu), and F. M. Geiger (geigerf@chem.northwestern.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Secondary organic aerosol (SOA) particles, formed from gas-phase biogenic volatile organic compounds (BVOCs), contribute large uncertainties to the radiative forcing that is associated with aerosols in the climate system. Reactive uptake of surface-active organic oxidation products of BVOCs at the gas–aerosol interface can potentially decrease the overall aerosol surface tension and therefore influence their propensity to act as cloud condensation nuclei (CCN). Here, we synthesize and measure some climate-relevant physical properties of SOA particle constituents consisting of the isoprene oxidation products  $\alpha$ -,  $\delta$ -, and *cis*- and *trans*- $\beta$ -IEPOX (isoprene epoxide), as well as *syn*- and *anti*-2-methyltetraol. Following viscosity measurements, we use octanol-water partition coefficients to quantify the relative hydrophobicity of the oxidation products while dynamic surface tension measurements indicate that aqueous solutions of  $\alpha$ - and *trans*- $\beta$ -IEPOX exhibit significant surface tension depression. We hypothesize that the surface activity of these compounds may enhance aerosol CCN activity, and that *trans*- $\beta$ -IEPOX may be highly relevant for surface chemistry of aerosol particles relative to other IEPOX isomers.

## 1 Introduction

Secondary organic aerosol (SOA) particles make up a substantial fraction of tropospheric aerosol and are known to lead to negative radiative forcing (Kanakidou et al., 2005; Carlton et al., 2009; Williams et al., 2011), yet, their formation ranks among the least understood processes in the atmosphere (Kanakidou et al., 2005; Hallquist et al., 2009; Goldstein and Galbally, 2007; Galbally et al., 2007; Riipinen et al., 2011). Many studies (Carlton et al., 2009; Hallquist et al., 2009; Kroll and Seinfeld, 2008; Lin et al., 2012; Worton et al., 2013; Kourtev et al., 2014; Ehn et al., 2014; Bates et al., 2014; Claeys et al., 2004a) support the idea that the gas-phase oxidation products of biogenic volatile compounds can either (a) partition to existing particles due to reduced volatil-

### Climate-relevant physical properties of molecular constituents

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ity compared to the parent compounds or (b) dissolve in aerosol or cloud water and participate in aqueous phase reactions to form low-volatility material. Surface tension is expected to be of particular importance for SOA formation and growth as it involves processes occurring at the interface between the SOA particle phase and the gas phase (Wang and Wexler, 2013). Moreover, atmospheric particles, once formed, can contain thousands of organic compounds or surfactants that can decrease the surface tension and thereby change aerosol particle properties such as cloud droplet formation, reactivity, and ice nucleation (Schwier et al., 2013; McNeill et al., 2014). Specifically, it has been reported that organic surfactants can influence the propensity of atmospheric aerosol particles to act as cloud condensation nuclei (CCN) by depressing the surface tension at the moment of activation (Facchini et al., 1999; Kiss et al., 2005; Shulman et al., 1996; Asa-Awuku et al., 2008; Facchini et al., 2000; Hitzenberger, 2002; Novakov and Penner, 1993; Salma et al., 2006; Sareen et al., 2013). Lower surface tension values result in decreases in the water vapor supersaturation required for cloud droplet activation, depending on ionic content, pH, temperature, and meteorological conditions. McNeill and coworkers recently showed that volatile surfactant species such as methylglyoxal and acetaldehyde can suppress surface tension at the gas–aerosol interface beyond predictions based on bulk surface tension measurements, leading to significantly enhanced CCN activity (Sareen et al., 2013). Chemical reactions at the surface and in the bulk of the particle between aerosol components may also influence overall surface tension and thus impact the dependence of CCN activity on the presence of surfactants. Yet, surface tension effects of many compounds relevant for SOA particle formation remain largely uncharacterized (McNeill et al., 2014; Woo et al., 2013).

Given their importance for global SOA particle formation (Carlton et al., 2009; Paulot et al., 2009; Claeys et al., 2004b), we report here surface tension values, measured using pendant drop tensiometry, of suspended drops of deionized water and 1.0 M ammonium sulfate solutions containing 0.1 to 30 mM concentrations of synthetically prepared isoprene-derived SOA particle constituents. Specifically, we studied the iso-



**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the limited solubility of the 2-methyltetraol compounds in octanol, stock solutions (~ 45 mM) of the tetraol compounds were prepared in deionized water. Equal volumes of stock solution and 1-octanol presaturated with water were mixed in three separate 15 mL polypropylene conical tubes. In all cases, phases of the solvent systems were mutually saturated by shaking for ~ 24 h on a mechanical shaker at room temperature. The three mixtures for each compound were subsequently centrifuged for 5 min at 3000 rpm to ensure complete phase separation. Three aliquots of the octanol phase were taken to determine the concentration of the IEPOX compounds (1–4), the tetraols (5, 6) and the alkene diol (7) compound.

The concentration of the compounds from the octanol phase for the epoxides and tetraols were determined using an Agilent 5973 gas chromatograph-mass spectrometer with a FFAP column (length 30 m, inner diameter 0.25 mm, film thickness 0.25  $\mu\text{m}$ ) and a quadrupole analyzer and EI ionization. The injector and detector temperatures were 260  $^{\circ}\text{C}$  and 250  $^{\circ}\text{C}$  respectively. For the alkene diol and IEPOX compounds, the oven had an initial temperature of 40  $^{\circ}\text{C}$  and a final temperature of 200  $^{\circ}\text{C}$  with a ramp rate of 15  $^{\circ}\text{C min}^{-1}$ . For the tetraol compounds, the oven had an initial temperature of 150  $^{\circ}\text{C}$  and a final temperature of 220  $^{\circ}\text{C}$  with a ramp rate of 30  $^{\circ}\text{C min}^{-1}$ . The gas flow rate was 1.0  $\text{mL min}^{-1}$ .

For IEPOX and alkene diol compounds, the quantity of the compound present at equilibrium in the aqueous phase was calculated from difference between the quantity of the compound originally introduced and the quantity in the octanol phase determined using the mass balance technique.

### 2.3 Viscosity studies

All viscosities were measured using solutions of 0.325  $\text{g mL}^{-1}$  of the compound of interest and 0.1625  $\text{g mL}^{-1}$   $(\text{NH}_4)_2\text{SO}_4$ . Viscosity measurements are relative to a control solution (0.1625  $\text{g mL}^{-1}$   $(\text{NH}_4)_2\text{SO}_4$  in deionized  $\text{H}_2\text{O}$ ) and were determined using a technique similar to a Cannon–Fenske viscometer, by measuring the time taken for





## Climate-relevant physical properties of molecular constituents

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sible hydroxyl groups of the epoxides due to the placement of the methyl group and possibly indicate that  $\alpha$ -IEPOX (4) would be the isomer most likely to partition into the organic-rich phase of particles. Replacement of the epoxide group in  $\alpha$ -IEPOX (4) with a simple alkene (7) shifted the  $\log(K_{ow})$  upward by about 0.4 units. This demonstrates that removal of the polar epoxide group significantly increases hydrophobicity.

The exact  $K_{ow}$  values of the two tetraol diastereomers (5 and 6) could not be determined, possibly due to their very limited solubility in octanol. GC traces of the octanol fraction in tetraol partitioning experiments showed that concentrations of the tetraols in the octanol fractions were below the detection limit. This indicates that the  $\log(K_{ow})$  values for tetraol compounds 5 and 6 would be much more negative than the values found for the IEPOX compounds.

Relative viscosities are listed in Fig. 1. The substances tested are all viscous liquids from room temperature down to  $-40^\circ\text{C}$ . The epoxides (1–3) have a viscosity similar to glycerol ( $1.98 \pm 0.03$ ), whereas the 2-methyltetraols (5, 6) are slightly more viscous and almost gelatinous.

### 3.2 Dynamic surface tension measurements

Based on the relevance of surface tension measurements in the prediction of new particle formation and aerosol CCN properties, the effect of concentration on surface tension over time was measured for the four epoxide isomers (1–4) and the two tetraol diastereomers (5, 6) in  $\text{dH}_2\text{O}$  and in  $1.0\text{ M }(\text{NH}_4)_2\text{SO}_4$ . As shown in Fig. 2, results in  $\text{dH}_2\text{O}$  showed that the  $\alpha$ -IEPOX (4) is by far the most surface active of the epoxide compounds. At the highest concentration measured (30 mM), interfacial tension for  $\alpha$ -IEPOX (4) was lowered by 5% at  $t = 0\text{ s}$  and decreased an additional 14% over the course of 10 min relative to  $\text{dH}_2\text{O}$ . While some of this decrease may be due to evaporation, the majority of the effect is most likely due to the migration of  $\alpha$ -IEPOX (4) to the surface of the droplet. Based on partitioning coefficients,  $\alpha$ -IEPOX (4) is the most hydrophobic of the epoxides and therefore would be more likely to partition from the bulk of the aqueous droplet to the surface.





(known as the dry diameter) as well as the surface tension of the droplet can alter its propensity to act as a cloud condensation nucleus. Köhler theory describes cloud droplet activation and growth from soluble particles (Köhler, 1936; Seinfeld and Pandis, 2008). The Köhler curve is given by:

$$s = \frac{A}{D_p} - \frac{B}{D_p^3} \quad (2)$$

with

$$A = \frac{4M_w\sigma}{RT\rho_w} \quad \text{and} \quad B = \frac{6n_sM_w}{\pi\rho_w} \quad (3)$$

where  $s$  is the supersaturation,  $D_p$  is the diameter of the aqueous droplet,  $M_w$  is the molecular weight of water and  $\rho_w$  its density,  $R$  is the gas constant,  $T$  is temperature,  $\sigma$  is surface tension, and  $n_s$  is the number of moles of solute. A decrease in surface tension due to the presence of surfactants would therefore decrease parameter  $A$  and result in increased CCN activation. If the bulk solute content of the particle remains constant, the effect of organic surfactants on equilibrium CCN activity can be assumed to be purely surface tension based. This assumption is valid based on the fact that gas-phase isoprene oxidation products will be continuously taken up at the gas-aerosol interface as they are consumed in heterogeneous reactions within the bulk and at the surface of the aerosol (Sareen et al., 2010). Using this assumption, the critical supersaturation for particles of a given size can be described as follows:

$$s_c^* = \left(\frac{\sigma}{\sigma_w}\right)^{3/2} s_c \quad (4)$$

Here,  $s_c^*$  is the critical supersaturation,  $\sigma_w$  and  $\sigma$  are the surface tension of water and the particle, respectively, and  $s_c$  is the critical supersaturation of particle with the surface tension of water ( $72.8 \text{ mN m}^{-1}$ ) (Engelhart et al., 2008). For all IEPOX and 2-methyltetraol compounds, dynamic surface tension measurements were fit to exponential curves in order to determine the equilibrium surface tension at  $t = \infty$  (Table 1). The

Climate-relevant  
physical properties  
of molecular  
constituents

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



equilibrium surface tension was used in Eq. (4) to calculate the critical supersaturation ratio ( $s_c^*/s_c$ ), which are listed in Table 2.

While there is some uncertainty regarding the in-particle concentrations of IEPOX and its reaction products, we can make reasonable estimates of these values based on field and modeling studies. Chan et al. (2010) reported up to  $24 \text{ ng m}^{-3}$  of IEPOX in Yorkville, GA during the 2008 August Mini-Intensive Gas and Aerosol Study (AMIGAS). During that period, they also measured  $33.4 \text{ microgram m}^{-3}$  of  $\text{PM}_{2.5}$ . Therefore, the observed IEPOX loading corresponds to an in-particle concentration of  $\sim 7 \text{ mM}$ , assuming  $1.2 \text{ g cm}^{-3}$  for the particle density. The McNeill group's coupled gas-aqueous aerosol chemistry model, Gas Aerosol Model for Mechanism Analysis (GAMMA McNeill et al., 2012), has been updated to include the latest aqueous phase IEPOX chemistry and physical parameters (Nguyen et al., 2013). GAMMA 1.4 simulations predict up to  $12 \text{ mM}$  in-particle IEPOX concentrations in a rural scenario. Therefore, we take here  $7.5 \text{ mM}$  as an example of an atmospherically relevant IEPOX in-particle concentration and find that  $\alpha$ -IEPOX (4) exhibited the largest decrease in  $s_c^*/s_c$  (9%), with an even larger decrease observed (23%) in  $1.0 \text{ M } (\text{NH}_4)_2\text{SO}_4$ . *trans*- $\beta$ -IEPOX (1) was also observed to lower surface tension and therefore is also expected to lead to decreased supersaturation ratios and enhanced CCN activity. The potential of *trans*- $\beta$ -IEPOX to enhance CCN activity is particularly significant based on recent studies demonstrating that *trans*- $\beta$ -IEPOX is the most abundantly produced isomer relative to other IEPOX isomers during isoprene oxidation (Bates et al., 2014). At  $10 \text{ mM}$ ,  $s_c^*/s_c$  for *trans*- $\beta$ -IEPOX (1) is predicted to decrease by 8% in  $\text{dH}_2\text{O}$  and in  $(\text{NH}_4)_2\text{SO}_4$ . Surface tension depression, and therefore the predicted impact on CCN activity, was less significant for *cis*- $\beta$ -IEPOX (2),  $\delta$ -IEPOX (3) and the 2-methyltetraols (5, 6). On a per-mole basis, surface tension depression by *trans*- $\beta$ -IEPOX is similar to that observed for methylglyoxal in bulk solutions (Sareen et al., 2010). The Henry's Law constant for IEPOX is several orders of magnitude higher than that of methylglyoxal (Nguyen et al., 2014), leading to a greater potential for suppression of aerosol surface tension by these species via bulk effects. That being said, as demonstrated by Sareen et al. (2013), bulk absorption

## Climate-relevant physical properties of molecular constituents

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of surface-active gases is apparently not a requirement for surface tension depression and enhanced CCN activity. Our results set the stage for future investigations of the effects of *trans*- $\beta$ -IEPOX on the CCN activity of aqueous aerosols.

## 5 Conclusions

In conclusion, we report dynamic surface tension measurements, using pendant drop tensiometry, of synthetically prepared isoprene-derived SOA particle constituents. Specifically, we studied the isoprene oxidation products  $\alpha$ -,  $\delta$ - and *cis*- and *trans*- $\beta$ -isoprene epoxide (IEPOX) (1–4) and *syn*- and *anti*-2-methyltetraol (5, 6) compounds. In addition, we experimentally determined octanol-water partitioning coefficients ( $K_{ow}$ ) and viscosities of these compounds. Results these experiments revealed that  $\alpha$ -IEPOX (4) is the most hydrophobic and surface active of the compounds studied here, however the hydrophobicity of these compounds did not coincide with surface activity for all compounds. Calculation of supersaturation ratios from surface tension values demonstrated that *trans*- $\beta$ -IEPOX (1) lowers supersaturation ratios significantly while the largest decrease in supersaturation ratios was calculated for  $\alpha$ -IEPOX (4). Other compounds measured, *cis*- $\beta$ -IEPOX (2),  $\delta$ -IEPOX (3), and the 2-methyltetraols (5, 6), demonstrated less significant surface activity and therefore minimal decreases in supersaturation ratios at higher concentrations.

The enhanced surface activity of *trans*- $\beta$ -IEPOX (1) and its potential to significantly decrease supersaturation ratios is particularly important based on its correlation with recent sum frequency generation (SFG) spectroscopy studies towards the identification of molecular constituents on the surfaces of isoprene-derived SOA particles (Ebben et al., 2014). This study identified *trans*- $\beta$ -IEPOX (1) as the closest match to the SFG spectra of isoprene-derived SOA surfaces, which coupled with surface tension experiments presented here, strongly indicates that *trans*- $\beta$ -IEPOX (1) may be present in higher concentrations at the surface of aerosol particles relative to other IEPOX isomers. This conclusion is also supported by the study by Wennberg and coworkers

### Climate-relevant physical properties of molecular constituents

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where *trans*- $\beta$ -IEPOX (1) was found to be produced in higher yields relative to other IEPOX isomers during isoprene oxidation by hydroxyl radicals. Reactive uptake of IEPOX compounds into aerosol particles by acid-catalyzed epoxide ring opening can also lead to formation of organosulfate and organonitrate derivatives (Worton et al., 2013; Surratt et al., 2007, 2010) so future studies will involve synthesizing these derivatives and analyzing their surface activity and other atmospherically relevant properties.

**The Supplement related to this article is available online at doi:10.5194/acpd-14-17197-2014-supplement.**

*Acknowledgements.* MAU gratefully acknowledges support from a NASA Earth and Space Science Fellowship. This work was supported by the Initiative for Sustainability and Energy at Northwestern (ISEN) and the National Science Foundation (NSF) Environmental Chemical Sciences Program in the Division of Chemistry under Grant No. 1212692.

## References

- Adamson, A. W. and Gast, A. P.: Physical Chemistry of Surfaces, 6th edn., John Wiley & Sons Inc., New York, 1997. 17202
- Asa-Awuku, A., Sullivan, A. P., Hennigan, C. J., Weber, R. J., and Nenes, A.: Investigation of molar volume and surfactant characteristics of water-soluble organic compounds in biomass burning aerosol, *Atmos. Chem. Phys.*, 8, 799–812, doi:10.5194/acp-8-799-2008, 2008. 17199
- Bates, K. H., Crounse, J. D., St Clair, J. M., Bennett, N. B., Nguyen, T. B., Seinfeld, J. H., Stoltz, B. M., and Wennberg, P. O.: *J. Phys. Chem. A*, 118, 1237–1246, 2014. 17198, 17200, 17208
- Broekhuizen, K., Kumar, P. P., and Abbatt, J. P. D.: *Geophys. Res. Lett.*, 31, 107–112, 2004. 17206

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmos. Chem. Phys.*, 9, 4987–5005, doi:10.5194/acp-9-4987-2009, 2009. 17198, 17199
- Chan, M. N., Surratt, J. D., Claeys, M., Edgerton, E. S., Tanner, R. L., Shaw, S. L., Zheng, M., Knipping, E. M., Eddingsaas, N. C., Wennberg, P. O., and Seinfeld, J. H.: *Environ. Sci. Technol.*, 44, 4590–4596, 2010. 17208
- 5 Claeys, M., Wang, W., Ion, A. C., Kourtchev, I., Gelencsér, A., and Maenhaut, W.: *Atmos. Environ.*, 38, 4093–4098, 2004a. 17198
- Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: *Science*, 303, 1173–1176, 2004b. 17199
- 10 Corrigan, C. E. and Novakov, T.: *Atmos. Environ.*, 33, 2661–2668, 1999. 17206
- Cruz, C. N. and Pandis, S. N.: A study of the ability of pure secondary organic aerosol to act a cloud condensation nuclei, *Atmos. Environ.*, 31, 2205–2214, 1997. 17206
- 15 Cruz, C. N. and Pandis, S. N.: The effect of organic coatings on the cloud condensation nuclei activation of inorganic atmospheric aerosol, *J. Geophys. Res.*, 103, 13111–13123, 1998. 17206
- Drozd, G. T. and McNeill, V. F.: Organic matrix effects on the formation of light-absorbing compounds from alpha-dicarbonyls in aqueous salt solution, *Environ. Sci. Process. Impacts*, 16, 741–747, 2014. 17202
- 20 Ebben, C. J., Strick, B. F., Upshur, M. A., Chase, H. M., Achtyl, J. L., Thomson, R. J., and Geiger, F. M.: Towards the identification of molecular constituents associated with the surfaces of isoprene-derived secondary organic aerosol (SOA) particles, *Atmos. Chem. Phys.*, 14, 2303–2314, doi:10.5194/acp-14-2303-2014, 2014. 17200, 17209
- 25 Ehn, M., Thornton, J. A., Kleist, E., Sipila, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I. H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurten, T., Nielsen, L. B., Jorgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petaja, T., Wahner, A., Kerminen, V. M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, 2014. 17198
- 30

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Ekström, S., Nozière, B., and Hansson, H.-C.: The Cloud Condensation Nuclei (CCN) properties of 2-methyltetrols and C3–C6 polyols from osmolality and surface tension measurements, *Atmos. Chem. Phys.*, 9, 973–980, doi:10.5194/acp-9-973-2009, 2009. 17206
- Engelhart, G. J., Asa-Awuku, A., Nenes, A., and Pandis, S. N.: CCN activity and droplet growth kinetics of fresh and aged monoterpene secondary organic aerosol, *Atmos. Chem. Phys.*, 8, 3937–3949, doi:10.5194/acp-8-3937-2008, 2008. 17207
- Facchini, M. C., Mircea, M., Fuzzi, S., and Charlson, R. J.: Cloud albedo enhancement by surface-active organic solutes in growing droplets, *Nature*, 401, 257–259, 1999. 17199
- Facchini, M. C., Decesari, S., Mircea, M., Fuzzi, S., and Loglio, G.: Surface tension of atmospheric wet aerosol and cloud/fog droplets in relation to their organic carbon content and chemical composition, *Atmos. Environ.*, 34, 4853–4857, 2000. 17199, 17206
- Finizio, A., Mackay, D., Bidleman, T., and Harner, T.: Octanol-air partition coefficient as a predictor of partitioning of semi-volatile organic chemicals to aerosols, *Atmos. Environ.*, 31, 2289–2296, 1997. 17203
- Galbally, I. E., Lawson, S. J., Weeks, I. A., Bentley, S. T., Gillett, R. W., Meyer, M., and Goldstein, A. H.: Volatile organic compounds in marine air at Cape Grim, Australia, *Environ. Chem.*, 4, 178–182, 2007. 17198
- Goldstein, A. H. and Galbally, I. E.: Volatile organic compounds in marine air at Cape Grim, Australia, *Environ. Chem.*, 4, 178–182, 2007. 17198
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009. 17198
- Henning, S., Rosenørn, T., D’Anna, B., Gola, A. A., Svenningsson, B., and Bilde, M.: Cloud droplet activation and surface tension of mixtures of slightly soluble organics and inorganic salt, *Atmos. Chem. Phys.*, 5, 575–582, doi:10.5194/acp-5-575-2005, 2005. 17206
- Hitzenberger, R.: Surface tension of Rax cloud water and its relation to the concentration of organic material, *J. Geophys. Res.*, 107, D244752, doi:10.1029/2002JD002506, 2002. 17199
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkan-

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- ski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005. 17198
- Keene, W. C.: Closure evaluation of size-resolved aerosol pH in the New England coastal atmosphere during summer, *J. Geophys. Res.*, 109, D23307, doi:10.1029/2004JD004801, 2004. 17202
- Kiss, G., Tombacz, E., and Hansson, H. C.: Surface tension effects of humiclike substances in the aqueous extract of tropospheric fine aerosol, *J. Atmos. Chem.*, 50, 279–294, 2005. 17199
- Klopffer, W., Rippen, G., and Frische, R.: Physicochemical Properties as Useful Tools for Predicting the Environmental Fate of Organic Chemicals, *Ecotox. Environ. Safe.*, 6, 294–301, 1982. 17203
- Kohler, H.: The nucleus in the growth of hygroscopic droplets., *T. Faraday Soc.*, 32, 1152–1161, 1936. 17207
- Kourtchev, I., Fuller, S. J., Giorio, C., Healy, R. M., Wilson, E., O'Connor, I., Wenger, J. C., McLeod, M., Aalto, J., Ruuskanen, T. M., Maenhaut, W., Jones, R., Venables, D. S., Sodeau, J. R., Kulmala, M., and Kalberer, M.: Molecular composition of biogenic secondary organic aerosols using ultrahigh-resolution mass spectrometry: comparing laboratory and field studies, *Atmos. Chem. Phys.*, 14, 2155–2167, doi:10.5194/acp-14-2155-2014, 2014. 17198
- Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593–3624, 2008. 17198
- Leo, A., Hansch, C., and Elkins, D.: Partition coefficients and their uses. *Chem. Rev.*, 71, 525–616, 1971. 17203
- Li, Z., Williams, A. L., and Rood, M. J.: Influence of soluble surfactant properties on the activation of aerosol particles containing inorganic solute, *J. Atmos. Sci.*, 55, 1859–1866, 1998. 17205
- Li, Z., Schwier, A. N., Sareen, N., and McNeill, V. F.: Reactive processing of formaldehyde and acetaldehyde in aqueous aerosol mimics: surface tension depression and secondary organic products, *Atmos. Chem. Phys.*, 11, 11617–11629, doi:10.5194/acp-11-11617-2011, 2011. 17205
- Lin, Y. H., Zhang, Z. F., Docherty, K. S., Zhang, H. F., Budisulistiorini, S. H., Rubitschun, C. L., Shaw, S. L., Knipping, E. M., Edgerton, E. S., Kleindienst, T. E., Gold, A., and Surratt, J. D.:

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Isoprene epoxydiols as precursors to secondary organic aerosol formation: acid-catalyzed reactive uptake studies with authentic compounds, *Environ. Sci. Technol.*, 46, 250–258, 2012. 17198

Liu, P. S. K., Leaitch, W. R., Banic, C. M., Li, S. M., Ngo, D., and Megaw, W. J.: Aerosol observations at Chebogue Point during the 1993 North Atlantic Regional Experiment: Relationships among cloud condensation nuclei, size distribution, and chemistry, *J. Geophys. Res.*, 101, 28971–28990, 1996. 17206

Matijevic, E. and Pethica, B. A.: The properties of ionized monolayers. Part 1. Sodium dodecyl sulphate at the air/water interface, *Trans. Faraday Soc.*, 54, 1383–1389, 1958. 17205

McNeill, V. F., Woo, J. L., Kim, D. D., Schwier, A. N., Wannell, N. J., Sumner, A. J., and Barakat, J. M.: Aqueous-Phase Secondary Organic Aerosol and Organosulfate Formation in Atmospheric Aerosols: A Modeling Study, *Environ. Sci. Technol.*, 46, 8075–8081, 2012. 17208

McNeill, V. F., Sareen, N., and Schwier, A. N.: Surface-active organics in atmospheric aerosols, *Top. Curr. Chem.*, 339, 201–259, 2014. 17199

Meylan, W. M. and Howard, P. H.: Estimating octanol-air partition coefficients with octanol-water partition coefficients and Henry's law constants, *Chemosphere*, 61, 640–644, 2005. 17203

Nguyen, Q. T., Skov, H., Sørensen, L. L., Jensen, B. J., Grube, A. G., Massling, A., Glasius, M., and Nøjgaard, J. K.: Source apportionment of particles at Station Nord, North East Greenland during 2008–2010 using COPREM and PMF analysis, *Atmos. Chem. Phys.*, 13, 35–49, doi:10.5194/acp-13-35-2013, 2013. 17208

Nguyen, T. B., Coggon, M. M., Bates, K. H., Zhang, X., Schwantes, R. H., Schilling, K. A., Loza, C. L., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Organic aerosol formation from the reactive uptake of isoprene epoxydiols (IEPOX) onto non-acidified inorganic seeds, *Atmos. Chem. Phys.*, 14, 3497–3510, doi:10.5194/acp-14-3497-2014, 2014. 17208

Novakov, T. and Penner, J. E.: Large contribution of organic aerosols to cloudcondensation – nuclei concentrations, *Nature*, 365, 823–826, 1993. 17199

Noziere, B., Baduel, C., and Jaffrezo, J. L.: The dynamic surface tension of atmospheric aerosol surfactants reveals new aspects of cloud activation, *Nat. Commun.*, 5, 3335–3342, 2014. 17202

Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kurten, A., St Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325, 730–733, 2009. 17199

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Pradeep Kumar, P., Broekhuizen, K., and Abbatt, J. P. D.: Organic acids as cloud condensation nuclei: Laboratory studies of highly soluble and insoluble species, *Atmos. Chem. Phys.*, 3, 509–520, doi:10.5194/acp-3-509-2003, 2003. 17206
- 5 Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., and Sherman, D. E.: The Effects of Low Molecular Weight Dicarboxylic Acids on Cloud Formation, *J. Phys. Chem. A*, 105, 11240–11248, 2001. 17206
- Raymond, T. M.: Cloud activation of single-component organic aerosol particles, *J. Geophys. Res.*, 107, 4787–4795, 2002. 17206
- Raymond, T. M.: Formation of cloud droplets by multicomponent organic particles, *J. Geophys. Res.*, 108, 4469–4477, 2003. 17206
- 10 Riipinen, I., Pierce, J. R., Yli-Juuti, T., Nieminen, T., Häkkinen, S., Ehn, M., Junninen, H., Lehtipalo, K., Petäjä, T., Slowik, J., Chang, R., Shantz, N. C., Abbatt, J., Leaitch, W. R., Kerminen, V.-M., Worsnop, D. R., Pandis, S. N., Donahue, N. M., and Kulmala, M.: Organic condensation: a vital link connecting aerosol formation to cloud condensation nuclei (CCN) concentrations, *Atmos. Chem. Phys.*, 11, 3865–3878, doi:10.5194/acp-11-3865-2011, 2011. 17198
- Surface tension of atmospheric humic-like substances in connection with relaxation, dilution, and solution pH, *J. Geophys. Res.*, 111, 205–212, 2006. 17199
- Sareen, N., Schwier, A. N., Shapiro, E. L., Mitroo, D., and McNeill, V. F.: Secondary organic material formed by methylglyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 10, 997–1016, doi:10.5194/acp-10-997-2010, 2010. 17205, 17207, 17208
- 20 Sareen, N., Schwier, A. N., Latham, T. L., Nenes, A., and McNeill, V. F.: Surfactants from the gas phase may promote cloud droplet formation, *Proc. Natl. Acad. Sci. USA*, 110, 2723–2728, 2013. 17199, 17206, 17208
- Schwier, A., Mitroo, D., and McNeill, V. F.: Surface tension depression by low-solubility organic material in aqueous aerosol mimics, *Atmos. Environ.*, 54, 490–495, 2012. 17205
- Schwier, A. N., Viglione, G. A., Li, Z., and Faye McNeill, V.: Modeling the surface tension of complex, reactive organic–inorganic mixtures, *Atmos. Chem. Phys.*, 13, 10721–10732, doi:10.5194/acp-13-10721-2013, 2013. 17199
- 30 Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley, New York, 1998. 17207

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Shulman, M. L., Jacobson, M. C., Carlson, R. J., Synovec, R. E., and Young, T. E.: Dissolution behavior and surface tension effects of organic compounds in nucleating cloud droplets, *Geophys. Res. Lett.*, 23, 277–280, 1996. 17199

Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L.,  
5 Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for organosulfates in secondary organic aerosol, *Environ. Sci. Technol.*, 41, 517–527, 2007. 17210

Surratt, J. D., Chan, A. W., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P.,  
10 Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene. *Proc. Natl. Acad. Sci. USA*, 107, 6640–6645, 2010. 17210

Wang, J. and Wexler, A. S.: Adsorption of organic molecules may explain growth of newly nucleated clusters and new particle formation, *Geophys. Res. Lett.*, 40, 2834–2838, 2013. 17199

15 Williams, J., Crowley, J., Fischer, H., Harder, H., Martinez, M., Petäjä, T., Rinne, J., Bäck, J., Boy, M., Dal Maso, M., Hakala, J., Kajos, M., Keronen, P., Rantala, P., Aalto, J., Aaltonen, H., Paatero, J., Vesala, T., Hakola, H., Levula, J., Pohja, T., Herrmann, F., Auld, J., Mesarchaki, E., Song, W., Yassaa, N., Nölscher, A., Johnson, A. M., Custer, T., Sinha, V., Thieser, J., Pouvesle, N., Taraborrelli, D., Tang, M. J., Bozem, H., Hosaynali-Beygi, Z.,  
20 Axinte, R., Oswald, R., Novelli, A., Kubistin, D., Hens, K., Javed, U., Trawny, K., Breitenberger, C., Hidalgo, P. J., Ebben, C. J., Geiger, F. M., Corrigan, A. L., Russell, L. M., Ouwersloot, H. G., Vilà-Guerau de Arellano, J., Ganzeveld, L., Vogel, A., Beck, M., Bayerle, A., Kampf, C. J., Bertelmann, M., Köllner, F., Hoffmann, T., Valverde, J., González, D., Riekkola, M.-L., Kulmala, M., and Lelieveld, J.: The summertime Boreal forest field measurement intensive (HUMPPA-COPEC-2010): an overview of meteorological and chemical  
25 influences, *Atmos. Chem. Phys.*, 11, 10599–10618, doi:10.5194/acp-11-10599-2011, 2011. 17198

Woo, J. L., Kim, D. D., Schwier, A. N., Li, R., and McNeill, V. F.: Aqueous aerosol SOA formation: impact on aerosol physical properties, *Faraday Discuss.*, 165, 357–367, 2013. 17199

30 Worton, D. R., Surratt, J. D., Lafranchi, B. W., Chan, A. W., Zhao, Y., Weber, R. J., Park, J. H., Gilman, J. B., de Gouw, J., Park, C., Schade, G., Beaver, M., Clair, J. M., Crouse, J., Wennberg, P., Wolfe, G. M., Harrold, S., Thornton, J. A., Farmer, D. K., Docherty, K. S., Cubison, M. J., Jimenez, J. L., Frossard, A. A., Russell, L. M., Kristensen, K., Glasius, M.,

**Climate-relevant  
physical properties  
of molecular  
constituents**

M. A. Upshur et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mao, J., Ren, X., Brune, W., Browne, E. C., Pusede, S. E., Cohen, R. C., Seinfeld, J. H., and Goldstein, A. H.: Observational insights into aerosol formation from isoprene, *Environ. Sci. Technol.*, 47, 11403–11413, 2013. 17198, 17210

5 Yuan, Y., Renbaum-Wolff, L., Carreras-Sospedra, M., Hanna, S. J., Hiranuma, N., Kamal, S., Smith, M. L., Zhang, X., Weber, R. J., Shilling, J. E., Dabdub, D., Martin, S. T., and Bertram, A. K.: Images reveal that atmospheric particles can undergo liquid-liquid phase separations, *Proc. Natl. Acad. Sci. USA*, 109, 13188–13193, 2012. 17203

10 Zhang, Q., Jimenez, J. L., Worsnop, D. R., and Canagaratna, M.: A case study of urban particle acidity and its influence on secondary organic aerosol, *Environ. Sci. Technol.*, 41, 3213–3219, 2007. 17202



## Climate-relevant physical properties of molecular constituents

M. A. Upshur et al.

**Table 2.** Supersaturation ratios for IEPOX (1–4) and 2-methyltetraols (5, 6)<sup>a, b</sup>.


Concentration (mM)						
	1	2	3	4	5	6
0	1.0 (1.02)	1.00 (1.0)	1.00 (1.04)	1.0 (1.04)	0.99 (1.05)	0.99 (1.04)
0.1	1.0 (1.04)	1.00 (1.03)	1.0 (1.04)	0.98 (1.0)	0.99 (1.0)	0.99 (1.04)
1	0.98 (1.02)	0.99 (1.03)	0.99 (1.0)	0.9 (1.00)	1.0 (1.0)	0.97 (1.04)
5	0.97 (1.02)	0.99 (1.03)	0.98 (1.04)	0.96 (0.86)	0.98 (1.01)	0.98 (1.03)
7.5	0.92 (1.01)	0.98 (1.03)	0.98 (1.03)	0.90 (0.80)	0.97 (1.00)	0.97 (1.03)
10	0.90 (0.96)	1.0 (1.02)	0.96 (1.03)	0.86 (0.78)	1.0 (1.00)	0.97 (1.02)
20	0.84 (0.97)	0.9 (1.00)	0.93 (1.01)	0.80 (0.66)	0.90 (0.98)	0.78 (0.95)
30	0.78 (0.86)	0.9 (0.93)	0.9 (1.0)	0.74 (0.60)	0.85 (0.96)	0.77 (0.94)

<sup>a</sup> Supersaturation ratios ( $s_c^*/s_c$ ) for solutions of IEPOX (1–4) and 2-methyltetraols (5, 6) in  $\text{dH}_2\text{O}$ .

<sup>b</sup> Values in parentheses denote supersaturation ratios ( $s_c^*/s_c$ ) in 1 M  $(\text{NH}_4)_2\text{SO}_4$ . Errors in ratios varied from 0.01–0.2 for IEPOX 1–3 and *syn*-2-methyltetraol 5, 0.01–0.1 for IEPOX 4, and 0.01–0.09 for *anti*-2-methyltetraol 6.

Climate-relevant  
physical properties  
of molecular  
constituents

M. A. Upshur et al.




a. Viscosities are reported as a time dependent ratio relative to a control solution of  $(\text{NH}_4)_2\text{SO}_4$  in deionized  $\text{H}_2\text{O}$  b.  $\alpha$ -IEPOX (4) was insoluble in  $\text{dH}_2\text{O}$  at higher concentrations used for viscosity experiments c. 2-methyltetraols (5,6) had concentrations in octanol below the detection limit d. Viscosity for alkene diol (7) was not measured due to its increased hydrophobicity relative to  $\alpha$ -IEPOX (4) observed during partitioning experiments.

**Figure 1.** Octanol–water partition coefficients and viscosity values for epoxides 1–4 and tetraols 5 and 6.

Climate-relevant  
physical properties  
of molecular  
constituents

M. A. Upshur et al.




**Figure 2.** Dynamic surface tension measurements for IEPOX compounds in dH<sub>2</sub>O. **(A)** *trans*- $\beta$ -IEPOX (1), **(B)** *cis*- $\beta$ -IEPOX (2), **(C)**  $\delta$ -IEPOX (3), **(D)**  $\alpha$ -IEPOX (4).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Climate-relevant  
physical properties  
of molecular  
constituents


M. A. Upshur et al.



**Figure 3.** Dynamic surface tension measurements for IEPOX compounds in 1 M  $(\text{NH}_4)_2\text{SO}_4$ . **(A)** *trans*- $\beta$ -IEPOX (1), **(B)** *cis*- $\beta$ -IEPOX (2), **(C)**  $\delta$ -IEPOX (3), **(D)**  $\alpha$ -IEPOX (4).

Climate-relevant  
physical properties  
of molecular  
constituents

M. A. Upshur et al.



**Figure 4.** Dynamic surface tension measurements for 2-methyltetraol compounds in dH<sub>2</sub>O (**A** and **B**) and 1 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (**C** and **D**). (**A**, **C**) *syn*-2-methyltetraol (**5**) (**B**, **D**) *anti*-2-methyltetraol (**6**).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)