



Long-term variability  
of dust events in  
Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.

# Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova<sup>1,2</sup>, O. Arnalds<sup>1</sup>, and H. Olafsson<sup>2,3,4</sup>

<sup>1</sup>Agricultural University of Iceland, Hvanneyri, Borgarnes, Iceland

<sup>2</sup>University of Iceland, Department of Physics, Reykjavík, Iceland

<sup>3</sup>Icelandic Meteorological Office, Reykjavík, Iceland

<sup>4</sup>Bergen School of Meteorology, Geophysical Institute, University of Bergen, Bergen, Norway

Received: 26 May 2014 – Accepted: 18 June 2014 – Published: 27 June 2014

Correspondence to: P. Dagsson-Waldhauserova (pavla@lbhi.is)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Long-term frequency of atmospheric dust observations was investigated for the southern part of Iceland and merged with results obtained from the Northeast Iceland (Dagsson-Waldhauserova et al., 2013). In total, over 34 dust days per year on average occurred in Iceland based on conventionally used synoptic codes for dust. Including codes 04–06 into the criteria for dust observations, the frequency was 135 dust days annually. The Sea Level Pressure (SLP) oscillation controlled whether dust events occurred in NE (16.4 dust days annually) or in southern part of Iceland (about 18 dust days annually). The most dust-frequent decade in S Iceland was the 1960s while the most frequent decade in NE Iceland was the 2000s. A total of 32 severe dust storms (visibility < 500 m) was observed in Iceland with the highest frequency during the 2000s in S Iceland. The Arctic dust events (NE Iceland) were typically warm and during summer/autumn (May–September) while the Sub-Arctic dust events (S Iceland) were mainly cold and during winter/spring (March–May). About half of dust events in S Iceland occurred in winter or at sub-zero temperatures. A good correlation was found between PM<sub>10</sub> concentrations and visibility during dust observations at the stations Vik and Storhofdi. This study shows that Iceland is among the dustiest areas of the world and dust is emitted the year-round.

## 1 Introduction

Frequency of dust episodes is monitored around many of the major desert areas of the world. Detailed and long-term studies on wind erosion variability can potentially explain the climatological and environmental changes in past. Periodical dust occurrences can affect ecosystem fertility and spatial and temporal distribution of animal and vegetation species similarly to climate variations (Fields et al., 2010). Oceanic ecosystems receive high amounts of nutrient rich dust spread over large areas where deserts occur near the sea (Arnalds et al., 2014). The long-term dust variability studies based on the

### Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

meteorological observations present up to 90 years old records from North America, Africa, Asia and Australia (N'TchayiMbourou et al., 1997; Qian et al., 2002; Natsagdorj et al., 2003; Ekström et al., 2004; Jamalizadeh et al., 2008; Steenburgh et al., 2012). Engelstaedter et al. (2003) reported high dust activity at many weather stations  
5 located in high-latitude regions. Cold climate regions are represented by long-term dust frequency in Northeast Iceland (Dagsson-Waldhauserova et al., 2013). Dust emission intensity and deposition rates in active glacial environment have been found very high, in some cases far exceeding those in lower latitudes (Bullard, 2013). Ganopolski et al. (2009) calculated glaciogenic dust deposition  $> 50 \text{ g m}^{-2} \text{ yr}^{-1}$  at the last glacial  
10 maximum with highest rates over the north-western Europe. Recently, the highest deposition rates of glaciogenic dust  $> 500 \text{ g m}^{-2} \text{ yr}^{-1}$  are reported from Iceland (Arnalds, 2010, see also Bullard, 2013).

Dust events in Arctic/Sub-Arctic region have been observed in Alaska (Nickling, 1978; Crusius et al., 2011), Greenland (Bullard, 2013), Svalbard (Dornbrack et al.,  
15 2010) and Iceland (Arnalds, 2010; Prospero et al., 2012; Thorarinsdottir and Arnalds, 2012). Arctic coastal zones are considered as the windiest regions on Earth (Eldridge, 1980). Strong winds in Iceland are causing some of the most extreme wind erosion events recorded on Earth (Arnalds et al., 2013).

The highest dust emissions in Arctic regions are associated with summer and early  
20 autumn (Nickling, 1978; Bullard, 2013; Dagsson-Waldhauserova et al., 2013). Dust concentrations in Sub-Arctic regions peak in spring (April–June, Prospero et al., 2012). Cold and winter periods are, however, of higher glaciogenic dust deposition than warm periods (Ganopolski et al., 2009). Dust events are frequent during dry years (Steenburgh et al., 2012; Dagsson-Waldhauserova et al., 2013), but suspended dust  
25 has also been observed during high precipitation and low wind conditions (Dagsson-Waldhauserova et al., 2014).

Iceland is an important source of volcanic sediments that are subjected to intense aeolian activity (Arnalds, 2010; Prospero et al., 2012; Thorarinsdottir and Arnalds, 2012; Arnalds et al., 2013) and is likely the largest glaciogenic dust source area in

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the Arctic/Sub-Arctic region. Total emissions of dust from Icelandic dust sources are of the range 30 to 40 million tons annually with 5–14 million tons deposited annually over the Atlantic and Arctic Oceans (Arnalds et al., 2014). Seven major dust plume sources have been identified (Arnalds, 2010). These sources are all in vicinity of glaciers. The most active glacial flood plain, Dyngjúsandur, covers an area of about 270 km<sup>2</sup> with up to 10 m thick sediments and is the main source for dust events in NE Iceland and towards Arctic (Dagsson-Waldhauserova et al., 2013). The major dust sources in South Iceland are Skeidararsandur, Myrdalssandur, Mælifellssandur, Landeyjasandur resulting in dust events south towards Europe during northerly winds, but alternatively towards Reykjavik and North America during easterly winds. The Hagavatn plume area is the source for frequent dust events towards Reykjavik and North America (the ocean southwest of Iceland). Glaciogenic dust from the Mælifellssandur area contains fine sharp-tipped shards with bubbles and 80 % of the particulate matter is volcanic glass rich in heavy metals (Dagsson-Waldhauserova et al., 2014). Such physical properties of the particles allow rapid suspension of moist particles within only a few hours after rains. In situ measurements from other dust plume areas are not available.

Dust suspension is related to reduced visibility. Wang et al. (2008) found a good correlation between PM<sub>10</sub> concentrations and visibility during dust observation. The visibility-dust formula can be used for dust concentration estimations where no aerosol mass concentration measurements are conducted (Dagsson-Waldhauserova et al., 2013). The relationship between dust concentration and visibility has not been investigated in Iceland.

The main objectives of this study were to explore the long-term (63 years) frequency of dust events in Iceland. Emphasis was given on determining the climatology and character of Arctic and Sub-Arctic dust events. In addition, the relationship between available dust concentrations and visibility during dust observation was investigated and the frequency of dust events placed in an international perspective.

## 2 Methods

### 2.1 Meteorological data and PM measurements

A network of 30 weather stations (15 in S Iceland, 8 in NE Iceland, and 7 in NW Iceland) operated by the Icelandic Meteorological Office was chosen for the study (Fig. 1). Table 1 shows the duration of station operation with majority of stations in operation since 1949. The data consist of conventional meteorological parameters such as wind velocity, wind direction, temperature and visibility, accompanied by synoptic codes of present weather. Present weather refers to atmospheric phenomena occurring at the time of observation, or which has occurred preceding the time of observation (IMO, 1981). The synoptic codes (ww) for present weather which refer to dust observation are 7–9, and 30–35. In addition, codes 4–6 are considered, but only if the codes for primary or secondary past weather (ww1, ww2) are 3 for blowing soil, dust, sand and dust storm (IMO, 1981; Dagsson-Waldhauserova et al., 2013). Weather observations were made 3–8 times a day.

Meteorological observations (synoptic codes for dust including 04–06 and visibility) were evaluated with available particulate matter (PM) mass concentrations data provided by the Environmental Agency of Iceland (EAI). The PM<sub>10</sub> data were obtained from the permanent station in Reykjavik (Grensasvegur, since 1996) and temporary stations in Vík and Kirkjubæjarklaustur (2010–2011). The Reykjavik station is equipped with Thermo EMS Andersen FH 62 I-R instrument, the Kirkjubæjarklaustur station with the Grimm EDM 365 and Thermo 5014 measured concentrations in Vík. Distance between the meteorological and EAI stations in Reykjavik and Kirkjubæjarklaustur is about one kilometer and several kilometers in Vík. Data set of dust concentrations (1997–2002, 2010) from the High-volume Filter Aerosol Sampler in Vestmannaeyjar (Westmann Islands) was used for evaluation of the dust codes and visibility at the Stórhofdi station (Prospero et al., 2012). Daily dust concentrations were correlated with the minimum visibility during dust observations during the preceding 24 h.

### Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Most of the conventional dust studies do not include synoptic codes 04–06 for “Visibility reduced by volcanic ashes”, “Dust haze” and “Widespread dust in suspension in the air” into the criteria for dust observation (Dagsson-Waldhauserova et al., 2013). Comparing these codes with available dust concentration measurements showed that  $PM_{10}$  concentration  $> 41 \mu g m^{-3}$  (about a double mean concentration) was exceeded in about 80 % of the 04–06 code cases. We have not included these codes in this long-term study except that ww1 or ww2 was 3.

## 2.2 Analysis

The initial dataset was built from the occurrence of “dust observation” made at one or more weather stations. Long-term dust activity was expressed in dust days. A “dust day” was defined as a day when at least one station recorded at least one dust observation. About 29 % of the observations did not include information on the present weather and they were excluded from the dataset.

Dust concentration measurements can be compared to the weather observations at few stations in Iceland and for a short time period. For the stations where  $PM_{10}$  measurements were available, we applied a power regression to determine the relationship between dust concentrations and visibility during dust codes including 04–06 (methods detailed in Wang et al., 2008). Visibility during dust observation was used to classify the severity of dust events in past (Dagsson-Waldhauserova et al., 2013).

## 3 Results

### 3.1 Frequency, spatial and temporal variability in dust production

A mean of 34.4 dust days per year was observed in Iceland during the period 1949–2011. An annual mean of 16.4 dust days (total of 1033 days) was recorded in NE Iceland (Dagsson-Waldhauserova et al. 2013) and about 17.9 dust days (total of 1153 days) occurred annually in southern parts of Iceland in 1949–2011. Figure 2 shows that

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the most dust active decade in Iceland was the 1960s while the 1980s were the lowest in number of dust days. For the southern part of Iceland, the highest frequency of dust events was in the 1950s–1960s, whereas the 2000s was the most frequent decade in the NE Iceland. The Grimsstadir station (NE) is the dustiest weather observation location in Iceland with > 12 dust days annually. The following dusty stations with > 3 dust days annually are represented in Table 2: Hofn (S), Vatnsskardsholar (S), Egilsstadir (NE), and Hella (S). The stations with highest dust frequency in southern part of Iceland are described in Fig. 2 (NE stations published in Dagsson-Waldhauserova et al. 2013a). The stations Hofn and Vatnsskardsholar reported highest number of dust days in the 1950s–1960s, the station Hella observed highest dust period in the 1960s–1970s and a new station in Hjarðarland (established in 1990) was the most active in the 2000s. Dust events were less severe in the 2000s than in the 1950s–1990s reflected by increased visibility during dust observations. Mean visibility during dust observations in S Iceland was 23.3 km indicating more severe dust events in S than in the NE Iceland or that weather stations are closer to major dust sources. Including codes 04–06 into the criteria for dust observation, the annual mean dust-day frequency was 135 dust days with 101 dust days observed in S Iceland and 34 dust days in NE Iceland.

### 3.1.1 Annual and seasonal dust day variability

An annual number of dust days in 1949–2011 is depicted in Fig. 3. The dustiest years were 1955, 1966 and 2010, when over 55 dust days occurred annually. The least dusty period was 1987–1990 with 11–15 dust days annually. Dust events occurred more frequently in southern part of Iceland than in NE Iceland in 1949–1954, 1962–1975, 1978–1981, and 2009–2011. The NE dust events were observed more often in 1955–1961, 1976–1977, 1982–1986, and 1992–2008 (except 1994, 2003). There is clear trend of having either the south or the north more active at a time. The years with relatively severe dust events (and annual visibility during dust observations < 15 km) were 1949, 1966, 1975, 1996, and 1998.

**Long-term variability  
of dust events in  
Iceland (1949–2011)**P. Dagsson-  
Waldhauserova  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The seasonal distribution of dust days in southern part of Iceland showed that about 47% of dust events occurred in winter (November–March) or during sub-zero temperatures. Dust days, as shown in Fig. 4, were most often in May (18% of dust days), April (13%) and March (11%). The lowest occurrence of dust days (< 6%) was in January, December, August and September. Contrarily, dust events in NE Iceland occurred mainly in summer and early autumn (May–September, Dagsson-Waldhauserova et al., 2013).

## 3.2 Climatology of dust events

### 3.2.1 Long-term trends in meteorological parameters of dust events (DE)

The mean DE temperature in southern part of Iceland was 3 °C with minimum 1.4 °C in the 1960s and maximum 5 °C in the 2000s (Fig. 5a). There was a great variability in DE temperatures, especially during the most active dust decade, the 1960s. The DE were the coldest in NE Iceland during the 1960s as well, but the warmest DE period was the 1950s (Dagsson-Waldhauserova et al., 2013). The mean DE temperature in the NE was significantly higher than in S Iceland, about 10.5 °C.

Dust observations in S Iceland reported high mean DE wind velocity of 13.6 m s<sup>-1</sup>, where the maximum mean of 15.6 m s<sup>-1</sup> was during the 1980s and the minimum of 11.9 m s<sup>-1</sup> during the 2000s (Fig. 5b). Extreme DE winds exceeding 30 m s<sup>-1</sup> occurred mainly in the 1960s and the 1970s. The mean DE wind velocity in NE Iceland was 10.3 m s<sup>-1</sup> with the maximum of 11.9 m s<sup>-1</sup> during the 2000s and the minimum of 8.6 m s<sup>-1</sup> in the 1980s (Dagsson-Waldhauserova et al., 2013).

The most common wind direction during dust events in S Iceland was N–NE, mainly reported from the stations Höfn, Hella, Vatnsskardsholar, Kirkjubaejarklaustur, Storföldi, Eyrarbakki, Vik, Thingvellir, Hjarðarland, Keflavik, and Reykjavik (Fig. 6). Dust events were often observed from the wind direction ENE (Haell, Vatnsskardsholar), E–ESE (Storföldi, Vatnsskardsholar, Thingvellir, Reykjavik, Keflavik), NW–



NNW (Höfn), and W–WNW (Vatnsskardsholar). The DE wind directions in NE Iceland were predominantly SW–S and SSE–SE (Dagsson-Waldhauserova et al., 2013).

### 3.2.2 Seasonal patterns in meteorological parameters of dust events

Seasonal variability in temperature and wind velocity during dust events in S Iceland is depicted in Fig. 7. The DE mean temperatures in October–May period are several degrees lower than the long-term monthly temperatures (higher in June–August period). Generally, the DE temperature in S Iceland was about 1.7°C lower than the long-term mean. Contrarily, the DE temperatures in NE Iceland were about 3°C higher than monthly long-term temperatures (Dagsson-Waldhauserova et al., 2013).

The DE wind velocities were significantly higher (5–11 m s<sup>-1</sup>) than long-term monthly wind velocities (Fig. 7b). The highest DE winds in S Iceland were from December to April while the lowest DE winds occurred in summer (June–September). This corresponds to the long-term monthly wind velocity trends. The mean DE wind velocity was 7.7 m s<sup>-1</sup> higher than long-term mean wind velocity. The difference is most pronounced during the winter months. In NE Iceland, the DE winds were about 4–7 m s<sup>-1</sup> higher than long-term means with maxima in May and September–October (Dagsson-Waldhauserova et al., 2013). Generally, the DE winds were about 3 m s<sup>-1</sup> lower in NE than S Iceland.

### 3.2.3 Dust event classification and meteorology

Reported dust events were of different severity. Where no atmospheric dust measurements are available, visibility during dust observation is used to estimate the dust event severity. Table 2 describes the dust event classes based on the visibility ranges. The most frequent were dust observations of “Suspended” and “Moderate suspended dust” (NE 73 %; S 59 %) with visibility 10–70 km, “Severe” and “Moderate haze” (NE 24 %; S 32 %) with visibility 1–10 km, and “Severe” and “Moderate dust storm” (NE 3 %; S 5 %)

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with visibility < 1 km. There were 32 “Severe Dust Storms” (visibility < 500 m) observed in Iceland (14 in NE mostly in the 1950s, 18 in S mostly in the 2000s).

The DE wind velocity increased with the DE severity, but the DE temperature decreased with the DE severity, except for “Moderate dust storm” recorded mostly at the Vik station in S Iceland. The parameters show that dust events in southern part of Iceland were observed as more severe than in NE Iceland.

Most of the dust classes in S Iceland occurred in April and May. Severe dust storms were most frequent in March and January at Vik, Hella, Kirkjubæjarklaustur, Hæll, Eyrarbakki and Vatnsskardsholar stations. The station Vik located only about 10 km from the Myrdalssandur dust source reported the mean DE visibility of 2 km indicating very severe dust events. Following stations with the lowest mean DE visibility were Raufarhofn (NE, 15 km), Höfn (18.3 km), Kirkjubæjarklaustur (20.1 km), Storhofdi (20.4 km), and Hella (21.1 km). The highest mean DE velocity was measured at the most windy station Storhofdi ( $22.6 \text{ m s}^{-1}$ ) while the lowest mean DE winds were at the station Thingvellir. Thingvellir recorded also the highest mean DE temperature ( $8.5^\circ\text{C}$ ) in S Iceland. The lowest DE temperatures were in Höfn ( $-2.3^\circ\text{C}$ ) located downwind Vatnajökull glacier.

About 18% of dust events in S Iceland were observed at more stations in the same time (two stations: 12.5%, three stations: 3.4%, four or more stations: 1.5%). Dust co-observations were mostly in Kirkjubæjarklaustur and Höfn, Kirkjubæjarklaustur and Vatnsskardsholar, and Kirkjubæjarklaustur with Hella. The Reykjavik station observed dust together with Hella or Thingvellir.

### 3.3 Relationship between $\text{PM}_{10}$ concentrations and visibility

Hourly  $\text{PM}_{10}$  concentrations were compared with corresponding visibility data during dust observations at available stations. Good correlation ( $R^2 = 0.73$ ) and considerable correlation ( $R^2 = 0.48$ ) were found between dust concentration and visibility by power function fitting at the stations in Vik and Vatnsskardsholar (Fig. 8a and b). Weak relationship between  $\text{PM}_{10}$  concentrations and visibility during dust codes ( $R^2 < 0.3$ ) was

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



found at the stations Reykjavik and Kirkjubæjarklaustur. Figure 8c shows visibility of all available dust codes plotted against corresponding  $PM_{10}$  concentrations together at all stations. Power function analysis resulted in moderate correlation ( $R^2 = 0.37$ ,  $p < 0.01$ ). Daily dust concentrations from the High-volume Filter Aerosol Sampler at Storhofdi during 1997–2002 and 2010 were well correlated with the 24 h minimum visibility ( $R^2 = 0.71$ , Fig. 8d).

## 4 Discussion

An annual mean of 34 dust days recorded in Iceland is comparable to dust studies from the active parts of China (35 dust days  $yr^{-1}$ , Qian et al., 2002), Mongolia (40 dust days  $yr^{-1}$ , Natsagdorj et al., 2003), and Iran (Jamalizadeh et al., 2008). The synoptic coding protocols can, however, contribute up to 15 % underestimation of annual dust day number (O’Loingsigh et al., 2010). Moreover, synoptic codes 04–06 showed a good agreement with increased  $PM_{10}$  concentrations. Including these codes into the criteria for dust observation, the annual mean dust-day frequency would be fourfold higher than applying conventionally used dust codes. This results in a total of 135 dust days per year on average for Iceland with 101 dust days observed in S Iceland and 34 dust days in NE Iceland. Such frequency can be found in parts of Australia and Africa (Ekström et al., 2004; N’TchayiMbourou et al., 1997). High numbers of dust observations presented here reflect previous studies showing high dust deposition rates in Iceland (Arnalds, 2010; Prospero et al., 2012; Thorarinsdottir and Arnalds, 2012; Bullard, 2013; Arnalds et al., 2013, 2014) and places the country among the important dust production areas of the world. Iceland is likely the most largest and active high-latitude cold dust source.

Trends in global dust emissions show high dust frequency during the 1950–1960s and low frequency during 1980s in the USA, Australia and China as well as in Iceland (Steenburgh et al., 2012; Ekström et al., 2004; Qian et al., 2002). The 2000s were reported as the most active decade in Iran and in NE Iceland (Jamalizadeh et al., 2008).

Dust periods retrieved from the ice-cores data during GISP2 project in Greenland correlate with the NE Iceland dust frequency 1950–1990 (Donarummo et al., 2002).

Generally, the period 1950–1965 was warm and dry in Iceland resulting in frequent dust suspension (Hanna et al., 2004). For the NE Iceland, the dustiest year 1955 with 37 dust days, coincides with one of the warmest and driest years in NE Iceland (Hanna et al., 2004). For the southern part of Iceland, the most frequent and severe dust event period was during 1965–1968. It was a period of below-average precipitation reported at stations Reykjavik, Stykkisholmur and Vestmannaeyjar (Hanna et al., 2004) while the 1965 was the driest year in SW Iceland for the past 100 years. The 20th century warm period in Iceland (1920s–1965) ended very abruptly in 1965 with about 1 °C drop in mean annual temperature (Hanna et al., 2004). The most exceptional year was, however, the year 1966 with 40 dust days reported in S Iceland. Not only was October 1966 reported as the driest October in Icelandic history, but also February 1966 in Reykjavik. Together with extremely strong maximum winds of more than 40 m s<sup>-1</sup>, the meteorological conditions in February 1966 caused at least 11 days of extremely severe dust storms. Local newspaper reported several large roofs removed from the houses, ships tore away from the harbors and planes turned around (Morgunblaðið, 1966).

The seventies were cold with high precipitation, but strong winds were often observed in S Iceland bringing the dust into suspension. The 1980s and 1990s were cold and with high precipitation in S Iceland while the 1990s were warm in the NE (Hanna et al., 2004). High frequency of dust events in NE Iceland during the 2000s was associated with dry and warm Junes. High number of dust days in S Iceland in 2010 was often because of resuspension of volcanic ash from the Eyjafjallajökull eruption during very frequent northerly winds (Petersen et al., 2012). The annual differences in dust event frequency do not correspond to trends of the global climate drivers such as the North Atlantic Oscillation (NAO), the Arctic Oscillation or prevailing ocean currents (Dagsson-Waldhauserova et al., 2013). The main driver is likely an orthogonal pattern to NAO, the dipole of Sea Level Pressure (SLP) oscillation oriented east-west (Dagsson-Waldhauserova et al., 2013).

Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The position of the Icelandic low determines whether dust plumes travel in a north-east or southerly direction. Higher frequency and severity of DE (low visibility and high wind speeds) in S Iceland than in NE Iceland is likely due to the close proximity of the S stations to the dust sources as well as higher number of the stations in the South (Fig. 1). The Grimsstadir station is > 100 km from the Dyngjusandur source while the southerly stations are in range of tens of km from the sources. Dust deposition rates and DE severity decrease exponentially with distance from the source (Arnalds et al., 2014). The local dust sources in S Iceland are also affected by milder oceanic climate during the winter while the NE highland dust sources are covered by snow for much of the winter. The dustiest weather station, Grimsstadir, is located downwind from the most active glacial plain in Iceland, Dyngjusandur, N of the Vatnajokull glacier. The most active stations are equally distributed around the areas with very high dust deposition (Arnalds, 2010) from the central NE, SE, S to SW Iceland. The land reclamation activities from the 1950s and 1970s (Crofts, 2011) result in decreased dust activity at the stations Hella and Höfn (Fig. 2).

The seasonal distribution of dust events in Iceland shows that the high dust period is from March to October. The NE dust events are typically warm, occurring during summer/autumn (May–September) while the S dust events are mainly cold, occurring during winter/spring (March–May). This is related to the SLP pattern which controls the warm southerly winds in NE Iceland as well as the cold northerly winds in S Iceland (Bjornsson and Jonsson, 2003). The S dust events are, however, more equally distributed during the year. The winter season is related to mild temperatures and high winds in S Iceland. Relatively high mean dust concentrations were measured during winter (January–March) at station Storhofdi (Prospero et al., 2012). The winter cold dust storms were frequently observed also in Mongolia (Natsagdorj et al., 2003). The highest number of dust storms occurred in March–May while the mean March–April temperatures were sub-zero. May is the driest and dustiest month in Iceland while June and September are the driest months only in NE Iceland (Hanna et al., 2004; Dagsson-Waldhauserova et al., 2013). Nevertheless, dust events can be observed also during

high precipitation seasons < 4 h after the rain (Dagsson-Waldhauserova et al., 2014). This agrees that even the highest precipitation year such 1972 can be of relatively high dust frequency.

5 Visibility during dust observations is an important indicator of dust event severity. To estimate the empirical relationship between visibility and dust concentration in Iceland, we compared available PM<sub>10</sub> concentrations with visibility based on methods in Wang et al. (2008). We found moderate correlation ( $R^2 = 0.37$ ,  $p < 0.01$ ) between dust concentrations and visibility which was likely caused due to several factors: (i) visibility was observed manually and only the prevailing visibility ( $\phi > 180^\circ$ ) recorded, (ii) generally  
10 low number of measurements, (iii) the stations were located in different distance of each other, (iv) time resolution between the dust and weather measurements, and v) station Reykjavik with majority of the measurements was influenced by anthropogenic aerosols. More observations are therefore needed.

15 Some of the most severe dust events in Iceland during 2007–2011 were captured by the Moderate Resolution Imaging Spectroradiometer (MODIS) flying on NASA's Terra satellite. Several dust plumes, shown on a visible wavelength, exceeded 1000 km travelling towards Europe, North America and Arctic. It was calculated that dust events caused deposition over 370 000 km<sup>2</sup> oceanic area around Iceland carrying 6–14 million tons of dust (Arnalds et al., 2014). The majority of dust, containing high amounts  
20 of bioavailable iron, is deposited in early spring in southern parts of Iceland. Oceanic biochemical cycles and productivity might therefore be strongly affected by local aeolian processes. We also emphasize here that considerably high dust event frequency and long-range transport of Icelandic dust may affect the environment and climate on macro scale. Icelandic dust aerosol should be included in climate projections as well  
25 as in the European and Arctic air pollution studies.

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 5 Conclusions

This study of long-term dust observations in Iceland showed that dust-day frequency in cold high-latitude areas can be comparable to the major desert areas in the world. It was found that dust events often occurred during winter and at sub-zero temperatures.

5 Observed dust events were more severe in southern part of Iceland than in NE Iceland, most likely because of close proximity of the southerly weather stations to major dust sources. The highest frequency of dust events was during the 1960s in S Iceland while most of dust events in NE Iceland occurred during the 2000s. The highest number of severe dust storms (visibility < 500 m) was observed in southern part of Iceland during  
10 the 2000s. Synoptic codes for dust were in good agreement with available dust concentration measurements; codes 04–06 should be considered in dust studies. There was a moderate correlation found between available PM<sub>10</sub> concentrations and visibility during the dust observations in Iceland. More synchronised dust and weather measurements are therefore needed. Iceland can be considered as the largest and most  
15 active desert and dust source at the boundary of the Arctic and Sub-Arctic region.

*Acknowledgements.* The work was supported by the Eimskip Fund of The University of Iceland and by the Nordic Centre of Excellence for Cryosphere-Atmosphere Interactions in a Changing Arctic Climate (CRAICC). We would like to thank Joseph Prospero from the University of Miami, USA, and Thorsteinn Johannsson from the Environment Agency of Iceland for providing the PM  
20 data for the dust measurements.

## References

- Arnalds, O.: Dust sources and deposition of aeolian materials in Iceland, *Icel. Agr. Sci.*, 23, 3–21, 2010.
- 25 Arnalds, O., Thorarinsdottir, E. F., Thorsson, J., Dagsson-Waldhauserova, P., and Agustsdottir, A. M.: An extreme wind erosion event of the fresh Eyjafjallajökull volcanic ash, *Nature Sci. Rep.*, 3, 1257, doi:10.1038/srep01257, 2013.

### Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Arnalds, O., Olafsson, H., and Dagsson-Waldhauserova, P.: Quantification of iron-rich volcanogenic dust emissions and deposition over ocean from Icelandic dust sources, *Biogeosciences Discuss.*, 11, 5941–5967, doi:10.5194/bgd-11-5941-2014, 2014.
- Bjornsson, H. and Jonsson, T.: Climate and climatic variability at Lake Mývatn, *Aquat. Ecol.*, 38, 129–144, 2003.
- Bullard, J. E.: Contemporary glacial inputs to the dust cycle, *Earth Surf. Proc. Land.*, 38, 71–89, 2013.
- Crofts, R. (Ed.): *Healing the Land*, Soil Conservation Service of Iceland, Reykjavik, Iceland, 2011.
- Crusius, J., Schroth, A. W., Gasso, S., Moy, C. M., Levy, R. C., and Gatica, M.: Glacial flour dust storms in the Gulf of Alaska: hydrologic and meteorological controls and their importance as a source of bioavailable iron, *Geophys. Res. Lett.*, 38, L06602, doi:10.1029/2010GL046573, 2011.
- Dagsson-Waldhauserova, P., Arnalds, O., and Olafsson, H.: Long-term frequency and characteristics of dust storm events in Northeast Iceland (1949–2011), *Atmos. Environ.*, 77, 117–127, 2013.
- Dagsson-Waldhauserova, P., Arnalds, O., Olafsson, H., Skrabalova, L., Sigurdardottir, G. M., Branis, M., Hladil, J., Skala, R., Navratil, T., Chadimova, L., von Lowis of Menar, S., Thorsteinsson, T., Carlsen, H. K., and Jonsdottir, I.: Physical properties of suspended dust during moist and low-wind conditions in Iceland, *Icel. Agr. Sci.*, 27, 2014.
- Dörnbrack, A., Stachlewska, I. S., Ritter, C., and Neuber, R.: Aerosol distribution around Svalbard during intense easterly winds, *Atmos. Chem. Phys.*, 10, 1473–1490, doi:10.5194/acp-10-1473-2010, 2010.
- Donarummo, J. J., Ram, M., and Stolz, M. R.: Sun/dust correlations and volcanic interference, *Geophys. Res. Lett.*, 29, 1361, doi:10.1029/2002gl014858, 2002.
- Eikström, M., McTainsh, G. H., and Chappell, A.: Australian dust storms: temporal trends and relationships with synoptic pressure distributions (1960–99), *Int. J. Climatol.*, 24, 1581–1599, 2004.
- Eldridge, F. R. (Ed.): *Wind Machines*, edited by: Van Nostrand, R., New York, USA, 1980.
- Engelstaedter, S., Kohfeld, K. E., Tegen, I., and Harrison, S. P.: Controls of dust emissions by vegetation and topographic depressions: an evaluation using dust storm frequency data, *Geophys. Res. Lett.*, 30, 1294, doi:10.1029/2002GL016471, 2003.



**Long-term variability  
of dust events in  
Iceland (1949–2011)**

P. Dagsson-  
Waldhauserova  
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fields, J. P., Belnap, J., Breshears, D. D., Neff, J. C., Okin, G. S., Whicker, J. J., Painter, T. H., Ravi, S., Reheis, M. C., and Reynolds, R. L.: The ecology of dust, *Front. Ecol. Environ.*, 8, 423–430, 2010.

Ganopolski, A., Calov, R., and Claussen, M.: Simulation of the last glacial cycle with a coupled climate ice-sheet model of intermediate complexity, *Clim. Past*, 6, 229–244, doi:10.5194/cp-6-229-2010, 2010.

Hanna, E., Jonsson, T., and Box, J. E.: An analysis of Icelandic climate since the nineteenth century, *Int. J. Climatol.*, 24, 1193–1210, 2004.

IMO: Reglur um veðurskeyti og veðurathuganir, Weather Observer Handbook, The Icelandic Meteorological Office, Reykjavik, Iceland, 85 pp., 1981.

Jamalizadeh, M. R., Moghaddamnia, A., Piri, J., Arbabi, V., Homayounifar, M., and Shahryari, A.: Dust Storm Prediction Using ANNs Technique (A Case Study: Zabol City), *Proc. World Aca. Sci. Eng. Tech.*, 33, 529–537, 2008.

Morgunblaðið: Þök fjúka af húsum, skip slitna upp, Roofs blow off houses, boats tore away, 1, available at: [http://timarit.is/view\\_page\\_init.jsp?issId=113054&pageld=1372641&lang=is&q=1966](http://timarit.is/view_page_init.jsp?issId=113054&pageld=1372641&lang=is&q=1966) (last access: 4 April 2014), 1966.

N'Tchayi Mbourou, G., Berrand, J. J., and Nicholson, S. E.: The diurnal and seasonal cycles of wind-borne dust over Africa north of the Equator, *J. Appl. Meteorol.*, 36, 868–82, 1997.

Natsagdorj, L., Jugder, D., and Chung, Y. S.: Analysis of dust storms observed in Mongolia during 1937–1999, *Atmos. Environ.*, 37, 1401–1411, 1978, 2003.

Nickling, W. G.: Eolian sediment transport during dust storms: Slims River Valley, Yukon Territory, *Can. J. Earth Sci.*, 15, 1069–1084.

O'Loingsigh, T., Mc Tainsh, G., Tapper, N., and Shinkfield, P.: Lost in code: a critical analysis of using meteorological data for wind erosion monitoring, *Aeolian Res.*, 2, 49–57, 2010.

Petersen, G. N., Björnsson, H., and Arason, T.: The impact of the atmosphere on the Eyjafjallajökull 2010 eruption plume, *Geophys. Res. Lett.*, 117, D00U07, doi:10.1029/2011JD016762, 2012.

Prospero, J. M., Bullard, J. E., and Hodgkins, R.: High-latitude dust over the North Atlantic: Inputs from Icelandic proglacial dust storms, *Science*, 335, 1078, 2012.

Qian, W. H., Quan, L., and Shi, S.: Variations of the dust storm in China and its climatic control, *J. Climate*, 15, 1216–1229, 2002.

Steenburgh, W. J., Massey, J. D., and Painter, T. H.: Episodic dust events of Utah's Wasatch front and adjoining region, *J. Clim. Appl. Meteorol.*, 51, 1654–1669, 2012.

Thorarinsdottir, E. F. and Arnalds, O.: Wind erosion of volcanic materials in the Hekla area, South Iceland, *Aeol. Res.*, 4, 39–50, 2012.

Wang, Y. Q., Zhang, X. Y., Gong, S. L., Zhou, C. H., Hu, X. Q., Liu, H. L., Niu, T., and Yang, Y. Q.: Surface observation of sand and dust storm in East Asia and its application in CUACE/Dust, *Atmos. Chem. Phys.*, 8, 545–553, doi:10.5194/acp-8-545-2008, 2008.

5

**Long-term variability  
of dust events in  
Iceland (1949–2011)**

P. Dagsson-  
Waldhauserova  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Weather stations in Iceland reporting synoptic observations. Observation period, number of dust observations, dust days and dust days per year are included. Stations are listed in descending order from the highest number of dust days.

Station	Observation period	Dust days	Dust observations	Dust day yr <sup>-1</sup>
<b>Grimsstadir</b>	1949–2011	791	1685	12.6
<b>Hofn</b>	1949–2011	243	575	3.9
<b>Vatnsskardsholar</b>	1949–2011	234	408	3.7
<b>Egilsstadir</b>	1949–1998	192	386	3.8
<b>Hella</b>	1958–2005	179	368	3.7
<b>Kirkjubaejarklaustur</b>	1931–2011	158	274	2
<b>Storhofdi</b>	1949–2011	118	204	1.9
<b>Haell</b>	1949–2011	94	132	1.5
<b>Hveravellir</b>	1965–2004	91	124	2.3
<b>Eyrbakki</b>	1957–2011	80	120	1.5
Vik	1961–2011	76	96	1.5
Keflavik	1952–2011	68	96	1.1
Vopnafjordur	1961–2011	64	83	1.3
Thingvellir	1949–1984	56	81	1.6
Reykjavik	1949–2011	41	70	0.7
Raufarhofn	1949–2011	41	61	0.7
Hjardarland	1990–2011	38	56	1.7
Sidumuli	1949–2011	30	37	0.5
Akureyri	1949–2011	26	26	0.4
Galtarviti	1953–1994	15	16	0.4
<b>Stadarholl</b>	1961–2011	12	15	0.2
<b>Stykkisholmur</b>	1949–2011	9	13	0.1
<b>Reykholar</b>	1961–2004	8	9	0.2
<b>Kollaleira</b>	1976–2007	5	7	0.2
<b>Blonduos</b>	1949–2003	5	6	0.1
<b>Natabu</b>	1949–2004	3	4	0.1
<b>Blafeldur</b>	1998–2011	2	2	0.1
<b>Bergstadir</b>	1978–2011	2	2	0.1
<b>Hornbjargsviti</b>	1949–2004	1	1	0.02
<b>Reykir i Hrutafj.</b>	1997–2011	1	1	0.1

**Long-term variability of dust events in Iceland (1949–2011)**

P. Dagsson-Waldhauserova et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.


**Table 2.** Dust event classification based on visibility criteria. Frequency of dust events, mean wind velocity, mean temperature, and annual number of dust days of each dust class are included. S represents southern part and NE northeastern part of Iceland.

Dust event class	Visibility (km)	Frequency (%)		Wind velocity ( $\text{m s}^{-1}$ )		Temperature ( $^{\circ}\text{C}$ )		Number of dust days $\text{yr}^{-1}$	
		S	NE	S	NE	S	NE	S	NE
Severe dust storm	$\leq 0.5$	1.2	< 1	15.7	16.2	-1.7	8.4	0.3	0.2
Moderate dust storm	0.5–1.0	3.5	2	13.6	14.9	4.1	9.4	1.1	0.5
Severe haze	1.0–5.0	14	10	15.0	13.0	1.1	10.6	3.0	2
Moderate haze	5.0–10.0	17	13	14.7	11.3	1.7	10.9	4.1	3
Suspended dust	10.0–30.0	42	46	13.5	9.9	3.0	10.6	10	10
Moderate susp. dust	30.0–70.0	16	27	11.7	10.2	3.7	10.0	6	7

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.




**Figure 1.** A map showing the locations of weather stations in Northeast and central Iceland (large black circles) and stations in northwestern and southern part of Iceland (small circles). Base map from the Agricultural University of Iceland Erosion Database (Soil Erosion in Iceland).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Long-term variability of dust events in Iceland (1949–2011)


P. Dagsson-  
Waldhauserova  
et al.



**Figure 2.** Total number of dust days, all stations combined to the left (blue bars for southern and northwestern part of Iceland, brown bars for Northeast Iceland). Individual stations in South Iceland sorted by decades to the right. Lines represent mean visibility (blue for S, brown for NE Iceland).

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.




**Figure 3.** Number of dust days (blue bars for southern and northwestern part of Iceland, brown bars for Northeast Iceland) and 3-year moving averages of dust day frequency (red for NE, light blue for S Iceland).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.



**Figure 4.** Number of dust days per month (bars) and monthly means of dust visibility (line) in southern part of Iceland in 1949–2011.




**Long-term variability  
of dust events in  
Iceland (1949–2011)**P. Dagsson-  
Waldhauserova  
et al.

**Figure 5.** Temperature (A) and wind velocity (B) for dust events in southern part of Iceland in 1949–2011. The boxes demarcate the range in which half the data can be found. The red lines represent the mean and the circles the median.

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-  
Waldhauserova  
et al.




**Figure 6.** Wind directions (WD) during dust events in southern part of Iceland in 1949–2011. Weather stations that observed mainly WD 0–18° – Höfn, Eyrarbakki, Kirkjubæjarklaustur, Storhofdi, Thingvellir; WD 18–36° – Höfn, Vatnsskardsholar, Hjardarland, Reykjavik, Keflavik; WD 36–54° – Hella, Vatnsskardsholar, Vik; WD 54–72° – Haell, Vatnsskardsholar; WD 90–108° – Storhofdi, Vatnsskardsholar; WD 270–306° – Vatnsskardsholar; and WD 306–342° – Höfn.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Long-term variability of dust events in Iceland (1949–2011)


P. Dagsson-Waldhauserova et al.



**Figure 7.** Monthly mean values (solid lines) of temperature **(A)** and wind velocity **(B)** during dust events in S Iceland in 1949–2011. Dashed lines represent the total mean values in 1949–2011.

## Long-term variability of dust events in Iceland (1949–2011)

P. Dagsson-Waldhauserova et al.



**Figure 8.** Hourly  $PM_{10}$  concentrations with corresponding visibility at stations: **(A)** – VÍk, **(B)** – Vatnsskardsholar, and **(C)** – all stations (Reykjavik, Vik, Vatnsskardsholar, and Kirkjubæjarklaustur). **(D)** represents daily  $PM_{10}$  concentrations concentrations from the High-volume Filter Aerosol Sampler with corresponding minimum 24 h visibility.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)