Fast transport from Southeast Asia boundary layer sources to Northern Europe: rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon anticyclone

B. Vogel¹, G. Günther¹, R. Müller¹, J.-U. Grooß¹, P. Hoor², M. Krämer¹, S. Müller², A. Zahn³, and M. Riese¹

55

¹Forschungszentrum Jülich, Institute of Energy and Climate Research – Stratosphere (IEK-7), Jülich, Germany ²Institute for Atmospheric Physics, University of Mainz, Mainz, Germany

³Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

Correspondence to: B. Vogel (b.vogel@fz-juelich.de)

Abstract. Enhanced tropospheric trace gases such as CO, CH_4 , and H_2O and reduced stratospheric O_3 were measured in situ in the lowermost stratosphere over Northern Europe on 26 September 2012 during the TACTS aircraft campaign.

- ⁵ The measurements indicate that these air masses clearly differ from the stratospheric background. The calculation of 40 day backward trajectories with the trajectory module of the CLaMS model shows that these air masses are affected by the Asian monsoon anticyclone. Some air masses origi-
- nate from the boundary layer in Southeast Asia/West Pacific 40 and are rapidly lifted (1–2 days) within a typhoon up to the outer edge the Asian monsoon anticyclone. Afterwards the air parcels are entrained by the anticyclonic circulation of the Asian monsoon. The subsequent long-range transport (8–14)
- ¹⁵ days) of enhanced water vapour and pollutants to the lower-⁴⁵ most stratosphere in Northern Europe is driven by eastward transport of tropospheric air from the Asian monsoon anticyclone caused by an eddy shedding event. We find that the combination of rapid uplift by a typhoon and eastward eddy
- shedding from the Asian monsoon anticyclone is a novel fast 50 transport pathway that may carry boundary emissions from Southeast Asia/West Pacific within approximately 5 weeks to the lowermost stratosphere in Northern Europe.

25 1 Introduction

30

One of the most pronounced circulation patterns in the the upper troposphere and lower stratosphere (UTLS) during boreal summer is the Asian summer monsoon circulation. It consists of a large-scale anticyclone in the UTLS extending from Asia to the Middle East from early June until end of September. The Asian monsoon anticyclone extends into the lowermost stratosphere (e.g., Randel and Park, 2006), where the tropopause above the monsoon is higher than in the extratropics by about 50 K (e.g., Dunkerton, 1995; Highwood and Hoskins, 1998). The anticyclone is flanked by an equatorial easterly jet to the south and by the subtropical westerly jet to the north and is characterised by low values of Potential Vorticity (PV) (e.g., Randel and Park, 2006).

In general, the Asian monsoon circulation provides an effective pathway for water vapour (Ploeger et al., 2013) and pollutants to the lower stratosphere of the Northern Hemisphere. Water vapour (H_2O) is the most important greenhouse gas and moistening of the stratosphere is an important driver of climate change (e.g., Forster and Shine, 1999, 2002; Shindell, 2001; Smith et al., 2001; Vogel et al., 2012). In particular, even small changes of H₂O in the UTLS have an impact on surface climate (e.g., Solomon et al., 2010; Riese et al., 2012). In spite of its radiative importance changes and trends in UTLS water vapour are only poorly quantified (e.g., Hurst et al., 2011; Kunz et al., 2013). Furthermore, increasing amounts of the greenhouse gas methane (CH_4) enhance stratospheric water vapour concentrations by methane oxidation (e.g., Röckmann et al., 2004; Riese et al., 2006; Rohs et al., 2006). Methane emissions by rice paddies in India and Southeast Asia (e.g., Khalil et al., 1998; Huang et al., 2004) may provide an important contribution to stratospheric CH₄ in this context. Enhanced water vapour concentrations in the UTLS affect not only the radiative balance, but have also the potential to affect stratospheric chemistry (e.g., Kirk-Davidoff et al., 1999; Dvortsov and Solomon, 2001; Vogel et al., 2011a). An increasing population and growing industries in Asia along with climate change have the potential to enhance future entry concentrations of water vapour and pollutants into the stratosphere transported by the Asian mon-

soon anticyclone. Therefore it is important to identify trans-120 port pathways to the global stratosphere to allow potential environmental risks to be assessed.

The Asian summer monsoon is associated with strong upward transport of tropospheric source gases by deep convec-

- ⁷⁰ tion which is confined by the strong anticyclonic circulation ¹²⁵ (e.g., Li et al., 2005; Randel and Park, 2006; Park et al., 2007, 2008, 2009). Hence, in satellite measurements relatively large concentrations of tropospheric trace gases such as water vapour (H_2O) (Rosenlof et al., 1997; Jackson et al.,
- ⁷⁵ 1998), carbon monoxide (CO) (Li et al., 2005; Park et al., ¹³⁰ 2008), and methane (CH₄) (Park et al., 2004; Xiong et al., 2009) are found in the Asian monsoon anticyclone, isolated by a transport barrier from the surrounding air. Conversely stratospheric trace gases such as O_3 show low concentra-
- tions in the anticyclone (Randel and Park, 2006; Liu et al., 135 2009; Konopka et al., 2010). However, the impact of different boundary layer sources on the chemical composition of the air in the Asian monsoon anticyclone (e.g., Li et al., 2005; Park et al., 2009; Chen et al., 2012; Bergman et al., 2013)
- and the mechanisms for transport into the lowermost strato- 140 sphere (Dethof et al., 1999; Park et al., 2009; Randel et al., 2010; Bourassa et al., 2012) are subject of current debate.

Trajectory calculations suggest air mass contributions in the Asian monsoon anticyclone from boundary sources orig-

- inating in India/Southeast Asia (including the Bay of Bengal) ¹⁴⁵
 and the Tibetan Plateau (Chen et al., 2012; Bergman et al., 2013). Chen et al. (2012) found the main contribution to air at tropopause height from the tropical Western Pacific region and the South China Seas, while Bergman et al. (2013) found
- ⁹⁵ that contributions from the Tibetan Plateau are most important at 100 hPa. In the Asian monsoon, findings by Chen et al. (2012) indicate that timescales of transport from the boundary layer to the tropopause region by deep convection overshooting are about 1–2 days, while several weeks by large
- scale ascent. Further, there is evidence that emissions on the 155 eastern side of the anticyclone (northeast India and southwest China) are lifted upward and trapped in the Asian monsoon anticyclone (Li et al., 2005).

There is evidence that the mean upward transport at the eastern/southeastern side of the Asian monsoon anticyclone 160 is a gateway of tropospheric air to the stratosphere by direct convective injection (Rosenlof et al., 1997; Park et al., 2007, 2008; Chen et al., 2012). However, the impact of this effect on the composition of the stratosphere has not been isolated

- from the influence of the transport in the deep tropics, en- 165 trained into the upward Brewer–Dobson circulation (Gettelman et al., 2004; Bannister et al., 2004), which is the primary transport pathway of air from the troposphere to the stratosphere (Holton et al., 1995).
- ¹¹⁵ One of the possible pathways for long-range transport of ¹⁷⁰ air masses from the Asian monsoon anticyclone to the extratropical lowermost stratosphere are smaller anticyclones

breaking off a few times each summer from the main anticyclone characterised by low PV values (Hsu and Plumb, 2001; Popovic and Plumb, 2001; Garny and Randel, 2013). This process is referred to as "eddy shedding". Westward transport of monsoon air masses by eddy shedding from the Asian monsoon anticyclone seems to be a common phenomenon, in contrast to eastward migrating anticyclones that occur less frequently (Popovic and Plumb, 2001; Garny and Randel, 2013). Garny and Randel (2013) found that westward propagation often appears after periods of strong convective forcing. Hsu and Plumb (2001) inferred from shallow-water calculations that, if the anticyclone is sufficiently asymmetric, the elongated anticyclone becomes unstable and westward eddy shedding occurs. In contrast, eastward eddy shedding is ascribed to the interaction of the monsoon anticyclone with an eastward moving mid-latitude synoptic-scale tropospheric cyclone (Dethof et al., 1999).

Eddy shedding events have the potential to carry air with tropospheric trace gases such as water vapour or pollutants from the Asian monsoon anticyclone to mid and high latitudes of the Northern Hemisphere (Dethof et al., 1999; Garny and Randel, 2013). These air masses can be transported into the extratropical lower stratosphere where they are eventually mixed irreversibly with the surrounding stratospheric air (Dethof et al., 1999; Garny and Randel, 2013) and thus affect the chemical and radiative balance of the extra-tropical UTLS.

In this paper, we show that fast transport from boundary emissions from Southeast Asia has the potential to affect the chemical composition of the lowermost stratosphere over Northern Europe. We focus on the question how transport pathways and time scales from surface emissions from Southeast Asia are influenced by the Asian monsoon circulation and its interaction with tropospheric weather systems.

We use in situ measurements obtained during the TACTS (Transport and Composition in the Upper Troposphere and Lowermost Stratosphere) aircraft campaign 2012 and Lagrangian backward trajectories calculated with the trajectory module of the Chemical Lagrangian Model of the Stratosphere (CLaMS) (e.g., McKenna et al., 2002a,b; Konopka et al., 2010, and references therein). Pure trajectory models represent the advective transport by the resolved flow using three-dimensional winds from meteorological data sets and are therefore useful for investigating of the origin of air masses, in particular in the region of the Asian monsoon anticyclone (Dethof et al., 1999; Chen et al., 2012; Bergman et al., 2013). The paper is organised as follows: Sect. 2 describes the measurements and Sect. 3 the trajectory calculations. In Sect. 4, the interaction of tropospheric weather systems, in particular of typhoons, with the Asian monsoon anticyclone and the resulting impact on the behaviour of the backward trajectories is presented. Our findings are discussed in Sect. 5 followed by a short summary and conclusions in Sect. 6.

2 Measurements of enhanced tropospheric trace gases in the stratosphere over Northern Europe

225 We use in situ measurements obtained during the TACTS aircraft campaign in August and September 2012. The TACTS 175 campaign was designed to study Transport and Composition in the Upper Troposphere and Lowermost Stratosphere. It was conducted jointly with the ESMVal (Earth System Model Validation) measurement campaign. Both TACTS and 230 ESMVal were performed using the German High Altitude 180 and LOng Range Research Aircraft HALO, a Gulfstream V. Here, we focus on the last research flight performed during TACTS/ESMVal on 26 September 2012 based out of Special Airport Oberpfaffenhofen (near Munich, 48° N 11° E), 235 Germany. The flightpath is shown in Fig. 1 in conjunction 185 with PV at 375 K potential temperature. PV is based on ERA-Interim reanalysis data (Dee et al., 2011) (see Sect. 3). The aircraft was flown roughly to the west to the British Isles, then to the north to the Atlantic Ocean, thereafter to the east towards Scandinavia. A closed loop flight track (hexagon)²⁴⁰ 190 was conducted over Norway to test the capability of the newly developed Gimballed Limb Observer for Radiance Imaging of the Atmosphere (GLORIA) to sound the atmosphere tomographically (Kaufmann et al., 2014; Riese et al., 2014). Thereafter, a flight pattern with a steep descent to the ²⁴⁵ 195 lower troposphere and directly afterwards a steep ascent back to the stratosphere (referred to as "dive") was flown inside the hexagon. The research flight ended in Oberpfaffenhofen. In this work, measurements of the following in situ instru-250 ments are used: 200 - The measurements of CO and CH₄ during with the TRIHOP TACTS/ESMVal were made instrument, which is an updated version of the threechannel tunable diode laser instrument for atmospheric 255 research (TRISTAR), which has been used during the 205 SPURT¹ project (Hoor et al., 2004; Engel et al., 2006). During the flight on 26 September 2012, TRIHOP achieves a precision of 0.9 ppbv for CO and 10 ppbv for $\rm CH_4$ for 1.5 s integration time. From the standard devi-260 210

ation of the in-situ calibrations, which were performed every 20 minutes, we estimate a reproducibility of the calibrations of 2.3 ppbv for CO and 15 ppbv for CH_4 without any corrections applied. The reproducibility can be regarded as an upper limit for the drift of the 265 measurements between the calibrations.

215

220

- The water measurements on board the HALO aircraft were obtained by the Fast In-situ Stratospheric Hygrometer (FISH) which is based on the Lyman- α photofragment fluorescence technique. Instrument and calibration procedure are described in Zöger et al. (1999). The FISH inlet is mounted forward-facing thus measuring total water, i.e. the sum of gas-phase water and water in ice particles. The procedure to correct the data for oversampling of ice particles in the forward-facing inlet is discussed in Schiller et al. (2008).

- A light-weight (14.5 kg) instrument (named FAIRO) for measuring ozone (O₃) with high accuracy (2%) and high time resolution (10 Hz) was developed for the use on board HALO. It combines a dual-beam UV photometer with an UV-LED as light source and a dry chemiluminescence detector (Zahn et al., 2012). The performance of FAIRO was excellent during all 13 flights during TACTS/ESMVAL.
- Potential temperature is deduced from the Basic Halo Measurement and Sensor System (BAHAMAS) which yield basic meteorological and avionic data of HALO.

Figure 2 shows potential temperature (top, black dots) at the measurement location, CO and O₃ (middle), and CH₄ and H_2O (bottom) along the flightpath of the flight on 26 September 2012. In the first half of the flight at altitudes between 370 K and 380 K potential temperature, enhanced concentrations of tropospheric trace gases CO, CH_4 , and H_2O were measured in the lowermost stratosphere and also reduced values of the stratospheric trace gas O_3 . In Fig. 2, the time period from 09:05 UTC (Universal Time Coordinated) to 10:17 UTC with enhanced tropospheric trace gases and reduced ozone is shaded in grey. Strong gradients are found in tracer measurements (at the beginning and the end of this time period) at the same level of potential temperature indicating that air masses were sampled with strongly different origin compared to the background air probed in this part of the flight. The region showing the enhanced tropospheric trace gases is referred to as "region of interest". The region of interest is located at the northwestern flank of the flight path over the Atlantic Ocean as shown in Fig. 1 (shown as white line in the flightpath). The potential vorticity (PV) at 375 K potential temperature indicates that in the region of interest and in the hexagon lower values of PV (7.3-8.0 PVU, yellow in Fig. 1) occur than in the stratospheric background, corroborating that these air masses have a different origin, that is more tropospheric, compared to the background. This is also evident in the Fig. 2 where, within the flight part conducted within the hexagon shortly before the dive ($\approx 13:00$ UTC), enhanced CO, CH₄, and H₂O and reduced O₃ were measured simultaneously. Between 08:05 UTC and 08:23 UTC a tropospheric signal is also evident in the observations (see Fig. 2) measured over the British Isles (Fig. 1). However, in the region of interest the measured signatures are most pronounced and therefore we focus here on that part of the flight.

3 Results of backward trajectory calculations

To study the origin of air in the region of interest (highlighted in grey in Fig. 2) 40 day backward trajectories are calculated

¹SPURenstofftransport in der Tropopausenregion (SPURT)

along the flightpath using wind data from the ERA-Interim

- reanalysis (Dee et al., 2011) (with a horizontal resolution $_{330}$ of $1^{\circ} \times 1^{\circ}$) provided by the European Centre for Medium-Range Weather Forecasts (ECMWF). For trajectory calculations, the trajectory module of the Chemical Lagrangian Model of the Stratosphere (CLaMS) (McKenna et al., 2002b;
- ²⁸⁰ Konopka et al., 2012, and references therein) is used. Tra-³³⁵ jectory calculations with CLaMS can be performed in both the diabatic mode and the kinematic mode, i.e. using potential temperature or pressure, respectively, as the vertical coordinate (Ploeger et al., 2010). Here, the diabatic approach
 ²⁸⁵ applied using the diabatic heating rate (with contributions ³⁴⁰
- from radiative heating including the effects of clouds, latent heat release, mixing and diffusion) as vertical velocity including the daily cycle (more details see Ploeger et al., 2010). The transport is simulated with vertical velocities from the
- diabatic heating rate in the UTLS (< 300 hPa) and using 345 a pressure based hybrid vertical coordinate in the troposphere (> 300 hPa) (Pommrich et al., 2014).

40 day backward trajectories starting at the observation along the flight path (every 10 s) on 26 September 2012 and ending at the origin of the air masses in the past (17 Au- $_{350}$ gust 2012) are calculated (in total 3582 trajectories). The potential temperature at the air mass origin (Θ_{org}) gives information about the vertical transport of the air mass along the trajectory. In Fig. 2 (top), the potential temperature at the

air mass origin (Θ_{org}) is shown as red dots. Most Θ_{org} val-355 ues lie above or at the same level of potential temperature as during the flight, consistent with the descent of air masses in the lower stratosphere of the Northern Hemisphere. However, in the region of interest (432 trajectories) some of the

³⁰⁵ air masses originate at much lower levels of potential tem- ³⁶⁰ perature, namely between 295 K and 360 K.

In Fig. 3, trajectories from two different Θ_{org} intervals, 295–320 K (left) and 320–360 K (right), are shown colourcoded by potential temperature (top) and by days reversed from 26 September 2012 (middle). Further, the geographi-365

from 26 September 2012 (middle). Further, the geographi-36 cal air mass origin (bottom, red dots in embedded map) and potential temperature vs. time along the 40 day backward trajectories (bottom) colour-coded by latitude are shown.

All air parcels with Θ_{org} below 360 K are affected by air masses originating from the Asian monsoon anticyclone ³⁷⁰ (Fig. 3, top). The air parcels are separated from the Asian monsoon circulation in the region over Japan, East China, and Southeast Siberia approximately 8–14 days before they were sampled during the flight on 26 September 2012 (Fig. 3, middle). 375

Air masses originating at much lower levels of potential temperature, below 320 K ($\Theta_{org} < 320$ K), are lifted between 35 and 40 days before the flight over the West Pacific (Fig. 3, left middle). Afterwards, they move first anti-clockwise at the edge of the anticyclone (at ≈ 365 K), then turn over Japan ³⁸⁰

before they become entrained into the Asian monsoon circulation and then move further around the Asian monsoon (left middle). For Θ_{org} between 320 K and 360 K, some of

325

the air masses intrude directly into the anticyclone and move clockwise round the Asian monsoon at the edge of the anticyclonic circulation. Other trajectories perform first a loop over the West Pacific ocean (right middle).

Some of the air parcels with Θ_{org} lower than 320 K show a very rapid uplift between 23 and 25 August with a maximum ascent rate of 41 K day⁻¹ (= 523 hPa day⁻¹) and originate in Southeast Asia (left bottom). Air parcels with Θ_{org} between 320 K and 360 K are characterised by rapid uplift up to 13 K day⁻¹ (= 139 hPa day⁻¹) between 18 and 24 August and originate in the West Pacific (northern or western of Philippines) or from northern of India (right bottom). An overview about maximum and mean vertical velocities is given in Table 1.

In the following we discuss also transport pathways of air masses originate in higher levels of potential temperature, namely in intervals for Θ_{org} between 360–370 K, 370–380 K, and 380–420 K.

Most air masses originating at Θ_{org} values between 360 K and 370 K experienced a moderately rapid uplift with mean values of about $2 \,\text{K}\,\text{day}^{-1}$ (= $22 \,\text{hPa}\,\text{day}^{-1}$) roughly in the region of the Asian monsoon anticyclone. Most of these air masses originate in North Africa, South Asia, and in the West Pacific (see Fig. 4 (top)). The trajectories are separated from the anticyclone approximately 8–14 days before the flight on 26 September 2012, and therefore transport air masses from inside the Asian monsoon anticyclone to Northern Europe (shown in Fig. 1 left in the electronic supplement of this paper).

At levels above, for Θ_{org} between 370 K and 380 K, the trajectories are affected by both the Asian monsoon anticyclone and the subtropical westerly jet. The moderate uplift along the trajectories is 1 K day^{-1} (= 16 hPa day^{-1}) in the mean. The majority of air masses originate in regions around the Asian monsoon anticyclone and in Central America (see Fig. 4 (middle)). In the latter case, the trajectories are most likely affected by the North American monsoon (shown in Fig. 1 right in the electronic supplement of this paper).

Most air masses that originate at Θ_{org} between 380 K and 420 K are dominated by the global circulation patterns with a descent of air masses in the Northern Hemisphere lower stratosphere. These trajectories do not circulate around the Asian monsoon anticyclone (except for two trajectories) and the air mass origins are spread out in the entire Northern Hemisphere, excluding regions affected by the Asian monsoon anticyclone such as South Asia or parts of North Africa (see Fig. 4 (bottom) and Fig. 2 in the electronic supplement of this paper). Our findings inferred from backward trajectory calculations are summarised in Table 1.

Our trajectory calculations suggest that air masses from altitudes, which are potentially strongly affected by boundary emissions from Southeast Asia and the West Pacific, are rapidly uplifted and are transported within approximately 5 weeks to Northern Europe. The air mass origins are not found in surface regions located closed to the core of the 440

480

Asian monsoon such as North India, South India or East China. This suggests that boundary emissions from these regions may need a longer time period for upward transport 435 within the Asian monsoon anticyclone to reach the lower-

most stratosphere over Northern Europe. However, we can-

not rule out that a strong uplift in convection over India and Tibet is not well enough resolved in ERA-Interim reanalysis data.

385

In this work, we focus on the region of interest (cf. Sect. 2), however also in other parts of the flight enhanced tropospheric signals in tracers and reduced ozone are measured

- between 08:05 UTC and 08:23 UTC and along the hexagon before the dive. The latter signal is also caused by trans-445 port of air masses from lower levels of potential temperature as evident in 40 day backward trajectories (Fig. 3, top). The tropospheric signal between 08:05 UTC and 08:23 UTC
- is not present in 40 day backward trajectories, but is evident
 in 60 day backward trajectories as discussed in Sect. 5.3.

4 Interaction between the Asian monsoon anticyclone and other tropospheric weather systems 455

To understand the behaviour of the backward trajectories presented in the previous section, the meteorology of the Asian monsoon area will be analysed based on ERA-Interim reanalysis data (Dee et al., 2011). In August 2012, the Asian₄₆₀ monsoon anticyclone was well established over Southern Asia and the Middle East. The Asian monsoon anticyclone

⁴¹⁰ is characterised by low PV values, indicating that it consists mainly of air masses of tropospheric origin. To the north the Asian monsoon anticyclone is connected to the subtrop-465 ical jet, to the south the equatorial westward flow is adjacent. The Asian monsoon anticyclone extends over a height range from 400 kPa or 240 K unword well into the lowerment.

415 range from 400 hPa or 340 K upward well into the lowermost stratosphere.

During August and September 2012 several tropical cy- $_{470}$ clones had impact on meteorological conditions in the vicinity of the Asian monsoon anticyclone. Further, during

⁴²⁰ September 2012 the subtropical jet over Asia was disturbed by strong Rossby waves triggered by low pressure systems travelling with the Arctic jet. The interaction between these ⁴⁷⁵ disturbances and the Asian monsoon anticyclone is discussed in the next sections.

425 4.1 Very rapid uplift in tropical cyclones in August 2012

In the northwestern Pacific region, tropical cyclones (typhoons) are observed at all times of the year, with storm activity peaking in late northern summer (e.g., Emanuel, 2003). In August and September 2012, several major typhoons of the category 4 or 5 according to the Saffir–Simpson hurriBetween 18 and 20 August 2012, the typhoon Tembin (named Igme by the Philippine Atmospheric, Geophysical and Astronomical Services Administration PAGASA) grew from a tropical depression to a category 4 typhoon southeast of Taiwan². Only a few days later the tropical typhoon Bolaven (PAGASA name Julian) formed east of the Philippines and moved northwest towards Taiwan. Bolaven was also classified as a class 4 typhoon³. After having made landfall twice in Taiwan, Tembin started to interact with Bolaven on 25 August, forcing Bolaven to move further to the north, where it made landfall in Korea on 28 August. Two days later, Tembin made landfall in Korea as well.

Figure 5 (top) shows the geographical position of the typhoons Tembin and Bolaven on 24 August 2012. Tembin was located over Taiwan, Bolaven further to the east over the Philippine Sea, being much stronger than Tembin at that time. The position of Bolaven coincides in time and space with the very rapid uplift of air masses evident in the trajectory calculations for Θ_{org} between 295 K and 320 K on 23/24 August 2012 (see Fig. 3, bottom).

Figure 5 (bottom) shows that both typhoons lifted air masses rapidly from the lower troposphere up to the tropopause region. The maximum updraft in a typhoon occurs at the eyewall, a ring of very deep convective clouds extending from the outer edge of the eye outward another 20–50 km (Emanuel, 2003). Maximum vertical uplift up to 300 hPa within 6 h (0.014 hPas⁻¹) is found at around 135° E at the eastern flank of Bolaven. This very rapid vertical transport is consistent with upward motion calculated along the 40 day backward trajectories of up to 276 hPa/6 h (see Table 1).

Bolaven moved northward during the next 4 days until it made landfall in Korea. This motion is also visible in the 40 day backward trajectories for Θ_{org} between 295 K and 310 K. Along the trajectories the air parcels move first cyclonically around the typhoon, before they move northward and enter the outer edge of the anticyclonic flow of the Asian monsoon anticyclone over Korea at ≈ 370 K (see Fig. 3 left middle panel).

For our trajectory calculations ERA-Interim reanalysis data are used. In ERA-Interim reanalysis data changes are implemented to improve deep and mid-level convection compared to previous reanalysis data (Dee et al., 2011), however small-scale rapid uplift in convective cores is not included. To show that the uplift in typhoons as represented in the ERA-Interim reanalysis data is strong enough to transport air parcels from the boundary layer to the Asian monsoon anticyclone as found in our backward trajectory calculations, we calculated forward trajectories starting in

the category 4 or 5 according to the Saffir–Simpson hurricane wind scale (SSHWS) occurred in the northwestern Pacific region.

²see e.g., http://www.nasa.gov/mission_pages/hurricanes/ archives/2012/h2012_Tembin.html

³see e.g., http://www.nasa.gov/mission_pages/hurricanes/ archives/2012/h2012_Bolaven.html

520

525

the boundary layer around the position the typhoon Bolaven ⁵³⁵ on 23 August 2012 12:00 UTC (in a window of 15°N-25°N,

- ⁴⁸⁵ 128°E-140°E and below ≈ 310 K potential temperature). The trajectories are started at a horizontal grid of $0.5^{\circ} \times 0.5^{\circ}$ and a vertical grid of 1 K. Within 48 hours (23-25 August 2012 12:00 UTC), 10% of these trajectories experience ⁵⁴⁰ a strong uplift greater than 50 K potential temperature as
- 490 shown in Fig 6. This very rapid uplift is also found on other days (22-26 August 2012) for both Bolaven and Tembin (not shown here). These calculations demonstrate that trajectories based on ERA-Interim reanalysis data show very 545 rapid uplift and can reach the lower stratosphere, in spite
- of the fact that ERA-Interim reanalysis data do not resolve small-scale rapid uplift in convective cores. Therefore, the ERA-Interim reanalysis data are well suited to identify transport pathways from the boundary layer in Asia to the 550 lowermost stratosphere over Northern Europe. We expect
 that in data sets including in addition the small-scale rapid
- uplift in convective cores, the very rapid uplift of air masses in tropical cyclones should be even more pronounced.

555

575

- In summary, our trajectory calculations show that typhoons in the northern West Pacific have the potential to very rapidly uplift air masses from the sea surface up to altitudes of the Asian monsoon anticyclonic circulation within 1 to 2 days. Balloon measurements of water vapour and 560 ozone in the Asian monsoon anticyclone launched in Kunming (China) in August 2009 show likewise a very rapid up-
- lift of boundary sources associated with the typhoon Morakot and afterwards injection of these air masses in the Asian monsoon anticyclone (Munchak et al., 2010). These mea-565 surements support the results of our trajectory calculations
- for September 2012 showing that rapid uplift by typhoons in the West Pacific may transport air parcels from the boundary layer directly in the outer edge of the Asian monsoon anticyclone.

4.2 Asian monsoon anticyclone and eddy shedding in September 2012

In August 2012, the Asian monsoon anticyclone was well established over Southern Asia and the Middle East. During both August and September the Asian monsoon anticyclone was influenced by strong Rossby waves and by several tropical cyclones.

A time sequence of PV fields derived from ERA-Interim reanalysis data at 370 K potential temperature over Asia from 9 until 20 September 2012 is shown in Fig. 7 (colours) highlighting the fine-scale structure of PV in the anticyclone and

the eastward propagation of filaments with low PV values at 585 its northeastern flank (showing PV values < 5 PVU). In addition, the horizontal velocity (white arrows) characterising spatial and temporal evolution of the flow in the Asian monsoon anticyclone are shown.

On 9 September 2012, the Asian monsoon anticyclone was located over South Asia and Northeast Africa (Fig. 7). At the northeastern flank of the anticyclone filaments with PV values lower than 5 PVU evolved between 10-14 September. The filament with low PV was extruded eastward along the subtropical westerly jet. A separation of the filament from the Asian monsoon circulation occurred and a small anticyclonic system developed during 15-16 September 2012. Thereafter, the secondary small anticyclone was elongated, moved westward and merged with another newly formed filament on the eastern flank of the Asian monsoon with low PV values close to Japan on 17/18 September. On 20 September, again a secondary anticyclone, larger than the secondary anticyclone from 15/16 September, was separated from the Asian monsoon anticyclone and carried air with low PV to the Pacific Ocean. Finally, the low PV air moved further eastwards along the subtropical jet across the extratropical tropopause to Northern Europe and dissipated slowly with time in the Northern Hemisphere lowermost stratosphere.

In September 2012, the subtropical jet was disturbed by strong Rossby waves triggered by low pressure systems travelling with the Arctic jet. The interaction between these disturbances and the subtropical jet can lead to baroclinic instabilities and eddy shedding (e. g. Orlanski and Sheldon, 1995) a process of peeling off secondary anticyclonic structures occurring a few times each summer (Dethof et al., 1999; Hsu and Plumb, 2001; Popovic and Plumb, 2001) (cf. Sect. 1). Figure 7 shows that both the Asian monsoon anticyclone and the secondary eastward propagating anticyclones are characterised by low PV values, indicating that it consists mainly of air masses of tropospheric origin. Both the Asian monsoon anticyclone and the secondary eastward propagating anticyclone are still a tropospheric feature. The transport of air masses into the lowermost stratosphere occurs afterwards. In contrast, tropospheric intrusions transport directly quasi-isentropically air masses from the tropical tropopause layer (TTL) into the extra-tropical lowermost stratosphere, where the air masses are mixed irreversibly (e.g., Vogel et al., 2011b). This can be caused by Rossby wave breaking events along the subtropical jet (e.g., Homeyer et al., 2011).

Fig. 7 shows in addition the temporal movement of the super typhoon Sanba (PAGASA name Karen) the strongest typhoon (class 5) during 2012 in the western Pacific region⁴. It made landfall on 17 September in South Korea (see Fig. 7) coincident with the process of peeling off secondary anticyclonic structures of the Asian monsoon anticyclone. Whether typhoon Sanba had an impact on the eddy shedding event of the Asian monsoon anticyclone in September 2012, and whether this in a more general feature of typhoons, is an interesting open question, which requires further investigations.

⁴see e.g., http://www.nasa.gov/mission_pages/hurricanes/ archives/2012/h2012_Sanba.html

The general behaviour of 40 day backward trajectories shown in Fig. 1 of the supplementary material and in Fig. 3 (main body of the paper) can be explained by the temporal evolution of the Asian monsoon anticyclone. The period 640 between 10 and 20 September 2012 is very active in terms

between 10 and 20 September 2012 is very active in terms of release of filaments and eddy shedding events with two pronounced events on 14 and 20 September on the northeast flank of the Asian monsoon anticyclone (see Fig. 7) causing the separation of the trajectories from the flow around the 645 core of the anticyclone. Nearly all trajectories originating below 380 K show this separation between 10 and 20 September 2012.

In summary, our trajectory calculations show that typhoons in the northern west Pacific have the potential to very 650

rapidly lift air masses from the sea surface to the edge of the Asian monsoon anticyclone within 1 to 2 days. In combination with the monsoon anticyclone and eastward eddy shedding events, this constitutes a new rapid transport pathway of surface emissions form the West Pacific to Northern 655

Europe. In our study, the minimum transport times from surface to the lowermost stratosphere over Northern Europe are approximately 5 weeks. Moistening of the stratosphere is an important driver of climate change, therefore the role of this new rapid transport pathway in moistening of the extratrop- 660

ical lower stratosphere during summer needs further investigation.

Moreover, in the last 30 years a poleward migration occurred of the latitude where tropical cyclones achieve their lifetime-maximum intensity (Kossin et al., 2014). If this environmental change continues, more tropical cyclones 665

migrate polewards to the northeastern flank of the Asian monsoon anticyclone. Therefore, the frequency of occurrence of the direct injections of polluted wet boundary layer air masses by very rapid uplift in typhoons into the Asian monsoon anticyclone could increase.

670

675

615

5 Discussion

5.1 Limitations of trajectory calculations

In general, trajectory calculations have limitations due to trajectory dispersion by errors through interpolation of the wind data to the air parcel positions at a specific time. Over the timescales in question, also mixing can be relevant. 680 Backward trajectory calculations as performed in our study can diagnose the origin of air parcels measured during the research flight on 26 September 2012 over Northern Europe. To confirm whether the pathways identified in backward tra-

jectories are robust, forward trajectories calculations starting 685 at the origin of the air parcels are useful. To corroborate our finding that trajectories starting in Pacific typhoons, circulat-695 ing around the Asian monsoon anticyclone, released by eddy

shedding events and are finally transported eastwards by the

subtropical jet can reach the region of interest over Northern Europe during the TACTS measurements on 26 September 2012, we calculated forward trajectories starting within the typhoon Bolaven and Tembin between 22 and 26 August 2012 at noon. These calculations show that trajectories starting on 22 to 26 August from typhoon Tembin do not reach the region of interest. However trajectories starting at typhoon Bolaven on 22, 23, 24 and 25 August (12:00 UTC) reach the region of interest. Fig. 8 shows trajectories started at the typhoon Bolaven on 23 August at noon (see Sect. 4.1). Shown are only trajectories that reach the region of interest during the flight on 26 September 2012 12:00 UTC (at locations between 365 K and 385 K, between 55°N and 70°N, and between 20°W and 0°W). These calculations confirm that air masses from the typhoon Bolaven have the potential to reach the region of interest within ≈ 5 weeks. Our calculations show that trajectories based on wind fields from ERA-Interim reanalysis data experience rapid uplift in tropical cyclones and can reach the flightpath of the TACTS flight on 26 September 2012 in the lower stratosphere over Northern Europe. The coincidence of forward and backward trajectory analysis demonstrates that the transport pathways identified in our study are unaffected by limitations due to trajectory dispersion caused by interpolation processes. Further, the typhoon Bolaven could be identified as source of air masses from low levels of potential temperature.

5.2 Chemical contribution of air masses with different origins

All air parcels measured in the region of interest (except at the very beginning and end) are chemically affected by the troposphere, however only 5% of the backward trajectories originate in the troposphere ($\Theta_{\rm org} < 360 \, {\rm K}$) and 34% in the Asian monsoon anticyclone or UTLS (360 K $< \Theta_{\rm org} <$ 380 K). The trajectories describe only advective transport, but in the real atmosphere in addition mixing processes occur. Therefore the region of interest is a mixture of air masses from the troposphere and the stratosphere which is reflected in different origins of the backward trajectories. To infer the relative contribution of the 2% air parcels lifted within the typhoon to the composition of the measured air parcels in the region of interest also mixing processes have to be taken into account. This is not possible with a pure trajectory study, here only the relative contributions of the trajectory origins can be determined.

Deduced from the number of trajectories originating in the lower troposphere or in the Asian monsoon anticyclone (5 % + 12 %), we conclude that in the region of interest a small amount air (17 %) with high concentrations of CO, CH₄, and H₂O is mixed with a huge amount of unpolluted dry stratospheric air (61 %). Filaments or eddy shedding events (see Fig. 7) transport polluted wet tropospheric air from the Asian monsoon into the lowermost stratosphere

- over Northern Europe. We assume that within these fila-690 ments or small anticyclones the tropospheric amount of trace gases is frozen in the core and only at the edge of 745 the filaments/anticyclones mixing with stratospheric air occurred, therefore the tropospheric characteristic of these
- air masses can be transported over a large distance from 695 Asia to Europe before the filaments dissipate slowly in the lowermost stratosphere. This kind of mixing process is 750 described in more detail e. g. in Vogel et al. (2011b) in a case study analyzing the transport of a topospheric intrusions from the tropical tropopause layer (TTL) into the lowermost 700

stratosphere over mid and high latitudes.

5.3 Sixty days backward trajectories

A non-negligible fraction of the parcels (12%) in the region of interest (cf. Sect. 2) is also transported from the Asian 760 705 monsoon region towards Northern Europe. In order to assess the geographical origin of these air parcels in the boundary layer, we calculated backward trajectories further beyond the 40 days limit. For that purpose, 60 days backward trajectories were calculated, a common trajectory length to study trans-765 710

port processes in the Asian monsoon region (ranging from a couple of weeks to a few months (e. g. Chen et al., 2012; Bergman et al., 2013).

For the region of interest, the origins of 60 days backward trajectories coming from low levels of potential tem-770 715 perature, from the $\Theta_{\rm org}$ interval 285 K–320 K, are clustered in Southeast Asia, West Pacific Ocean, Bay of Bengal, and the Arabian Sea. 53 % of these trajectories from $\Theta_{\rm org}$ values between 285 K and 320 K are from locations between

10°S, 40°N, 60°E, and 160°E. The origins of other trajecto-720 ries are distributed in several different geographical regions (see Fig. 3 in the electronic supplement of this paper).

The origins of trajectories from higher levels of potential temperature, from the $\Theta_{\rm org}$ interval 320 K–360 K, are concentrated around the Tibetan Plateau, Northeast China, 725 Mongolia, Southeast Asia, and the Pacific Ocean. 55,% of these trajectories from $\Theta_{\rm org}$ values between 320 K and 360 K are from locations between 0°N, 50°N, 60°E, and $^{^{780}}$ 160°E. Also here, the origins of other trajectories are distributed globally (see Fig. 3 in the electronic supplement 730

of this paper). Calculations of 60 day backward trajectories confirm that the origins of the air masses with enhanced values of tropospheric trace gases such as $\rm CH_4,\, CO,\, and^{785}$ H₂O found in the region of interest during the TACTS flight on 26 September 2012 are mainly due to boundary source 735 emissions in Asia.

In Sect. 3, it is shown that in addition to the region of inter-⁷⁹⁰ est (cf. Sect. 2), in an other part of the flight enhanced tropospheric signals in tracers and reduced ozone are measured be-740 tween 08:05 UTC and 08:23 UTC. However, in 40 day backward trajectories no evidence is found that these air masses

are from lower levels of potential temperature. But calculations of 60 day backward trajectories show that also in this part of the flight, as well as in the region of interest, some of the trajectories originate at low levels of potential temperature between 295 K and 320 K. Fig. 9 shows that these air parcels are also from Southeast Asia and experience very rapid uplift on 2-3 August 2012 southeast of the Philippines. The reason for the strong upward transport is again a typhoon, namely typhoon 'Haikui'⁵.

The results from 60 day backward trajectories demonstrate that very rapid uplift in typhoons in combination with the Asian monsoon anticyclone is not a single case of long-range transport, but rather a fast transport pathway from boundary layer in Asia to the lowermost stratosphere over Northern Europe that occurs more frequently. How frequently this pathway is expected to take place is an interesting open question. During the typhoon season 2012⁶ 11 typhoons occurred between June and September. But also tropical depressions and tropical storms have the potential for very rapid uplift of boundary layer emissions. The impact of each of theses events in combination with the Asian monsoon anticyclone has to be analysed. We expect that the impact of this pathway is smaller compared to other (slower) transports from the tropical troposphere into the extratropical lower stratosphere. However, the impact of this rapid transport pathway is clearly evident in the TACTS measurements. The impact of this transport pathway also depends on the particular species considered. The shorter lived the species, the more important the fast transport pathway will be provided that the lifetime of the species is longer than the transport time.

Conclusions 6

755

On 26 September 2012, a filament with enhanced tropospheric trace gases such as CH₄, CO, and H₂O and reduced stratospheric O₃ was measured over Northern Europe in the extratropical lowermost stratosphere during a flight on board the German Research Aircraft HALO during the TACTS campaign. Trajectory calculations show that these air masses originate in Southeast Asia and are affected by the Asian monsoon anticyclone. Based on our analysis of these observations and the associated trajectory calculations, we propose a new rapid transport pathway of approximately 5 weeks from the boundary layer sources in Southeast Asia to the location of the measurement in the lowermost stratosphere over Northern Europe.

Our findings show that air originating in the polluted boundary layer in South East Asia can be very rapidly lifted

⁵see e.g., http://www.nasa.gov/mission_pages/hurricanes/archives/ 2012/h2012_Damrey.html

⁶see e. g.:http://en.wikipedia.org/wiki/2012_Pacific_typhoon_season; season summary

within 1–2 days by Pacific typhoons into air masses circulating around the Asian monsoon anticyclone (at ≈ 365 K potential temperature). Similar timescales are found for deep convective overshooting in this region (Chen et al., 2012)

- or in convective transport in the West African tropics (e.g., 850 Fierli et al., 2011). Therefore, our findings confirm evidence from previous studies that at the eastern side of the anticyclone tropospheric air masses are lifted and trapped in the anticyclone (Li et al., 2005) or can be injected by convec-
- tion into the stratosphere (Rosenlof et al., 1997; Park et al., 855 2007, 2008; Chen et al., 2012). Moreover, boundary emissions from India or China lifted within the Asian monsoon anticyclone are not found in 40 day backward trajectories, but in 60 day backward trajectories suggesting that transport form houndary emissions form India or China form India end China theorem.
- from boundary emissions from India and China through the 860
 Asian monsoon anticyclone is comparatively slow.

However, we cannot rule out that a strong uplift in convection over India and Tibet is not well enough resolved in ERA-Interim reanalysis data in contrast to convection in tropical cyclones that are much larger than most convective 865

cells. This might lead to a certain underestimation of upward transport in smaller scale convective systems compared to upward transport in typhoons.

810

- In Fig. 10, selected 40 day backward trajectories representing characteristic pathways with different vertical velocities 870 (cf. Table 1) are shown as functions of potential temperature and longitude summarising our results. The figure illustrates different transport pathways of air masses with different origins to the location of the measurement over North-
- ern Europe. The observed air masses in the region of interest 875 are a mixture of air masses with distinct origins in different fractions (cf. Table 1). The combination of very rapid uplift by a typhoon and eastward eddy shedding from the Asian monsoon anticyclone yield fast transport (≈ 5 weeks) from
- Southeast Asia boundary layer sources to Northern Europe 880 (2% of all trajectories, red). Some of the 40 day backward trajectories originate in the West Pacific troposphere (3%, blue) or within the Asian monsoon anticyclone over South Asia/North Africa (12%, green). Air masses that originate
- in the UTLS at altitudes between 370 K and 380 K poten-885 tial temperature are mainly from the edge of the Asian monsoon anticyclone (22 %, yellow). Most air parcels originate at higher altitudes between 380 K and 420 K. These air parcels come from the entire Northern Hemisphere, with the exception of the region of the Asian monsoon anticyclone (61 %,
- grey).

Further, our calculations show that after the injection into ⁸⁹⁰ the anticyclonic circulation, the air masses circulate clock-wise, in an upward spiral, at the edge of the Asian monsoon

- anticyclone around the core of the anticyclone and do not enter the core of the anticyclone itself. In addition to the upward transport within the Asian monsoon connecting surface air with enhanced pollution to the lower stratosphere (e.g., Park et al., 2009; Randel et al., 2010; Chen et al., 2012; Bergman
- et al., 2013), the injection into the outer edge of the anticy-895

clone and subsequent clockwise circulation around the core of the anticyclone is an additional pathway of air mass transport from the Asian monsoon anticyclone to the lowermost stratosphere. During the Nabro volcanic eruption in 2011, direct injections in the upper troposphere at the southwest flank of the Asian monsoon anticyclone (over Northwest Africa) have been observed by satellite observations by the Michelson Interferometer for Passive Atmospheric Sounding (MI-PAS) (Griessbach et al., 2013). The MIPAS observations also show clockwise transport of air masses injected directly at the southwest flank in the outer edge of the Asian monsoon anticyclone. This support our findings about the clockwise transport of air masses injected directly at the east flank by typhoons around the core of the Asian monsoon anticyclone.

Our analysis shows that eastward eddy shedding events occurred in mid September 2012. The subtropical jet was disturbed by strong Rossby waves triggered by low pressure systems travelling with the Arctic jet. The interaction between these disturbances and the subtropical jet led to the peeling off of secondary anticyclonic structures from the Asian monsoon anticyclone.

The backward trajectories show that the transport time of water vapour and pollutants from the Asian monsoon anticyclone to Northern Europe caused by eddy shedding events and transport along the subtropical jet is about 8-14 days in the case study discussed here. Thereafter, the trajectories travel further northeastwards along the subtropical jet to the Pacific Ocean and finally to Northern Europe. In our study, moisture and pollution are transported very fast within up to 5 weeks from the lower troposphere in Southeast Asia to the lowermost stratosphere over Northern Europe caused by the combination of very rapid uplift by a typhoon and eastward eddy shedding from the Asian monsoon anticyclone. This rapid transport pathway into the lowermost stratosphere is for short-lived substances (e.g. bromine-containing very shortlived source gases) of particular importance. Eddy shedding events have the potential for long-range transport of enhanced concentrations of boundary layer sources such as water vapour and pollutants from Southeast Asia to mid and high latitudes of the Northern Hemisphere lowermost stratosphere.

Supplementary material related to this article is available online at: http://\@journalurl/\@pvol/\@ fpage/\@pyear/\@journalnameshortlower-\@pvol-\ @fpage-\@pyear-supplement.pdf.

Acknowledgements. We sincerely thank Andreas Engel and Harald Bönisch (University Frankfurt) for coordinating the TACTS campaign and the HALO crew and pilots for conducting the regarded flight. The authors gratefully acknowledge Paul Konopka and Felix Plöger (both at Research Centre Jülich) for helpful discussions on the Asian monsoon circulation and Laura Pan (NCAR, Boulder) for discussions on the balloon observations in China associated with the strong uplift in typhoons. We thank the European Centre of Medium-Range Weather Forcasts (ECMWF) for providing 955

meteorological analyses and the ERA-Interim reanalysis data. Our 900 activities, flight planning and forecasts were partly founded by the German Science Foundation (Deutsche Forschungsgemeinschaft, DFG) under the DFG project LASSO (HALO-SPP 1294/GR 3786). The authors thank three anonymous reviewers for their very helpful 960 comments. 905

The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

References 910

references

- Bannister, R. N., O'Neill, A., Gregory, A. R., and Nissen, K. M.: The role of the south-east Asian monsoon and other seasonal features in creating the "tape-recorder" signal in the Unified Model, Q. J. Roy. Meteor. Soc., 130, 1531-1554, 2004. 975
- 915 Bergman, J. W., Fierli, F., Jensen, E. J., Honomichl, S., and Pan, L. L.: Boundary layer sources for the regional contributions Asian anticyclone: to а verconduit, Geophys. Res., 118, tical J. 2560-2575, doi:http://dx.doi.org/10.1002/jgrd.5014210.1002/jgrd.50142, 920 980 2013.
- Bourassa, A. E., Robock, A., Randel, W. J., Deshler, T., Rieger, L. A., Lloyd, N. D., Llewellyn, E. J. T., and Degenstein, D. A.: Large volcanic aerosol load in the stratosphere
- linked to Asian monsoon transport, Science, 337, 78-81, 985 925 2012
 - Chen, B., Xu, X. D., Yang, S., and Zhao, T. L.: Climatological perspectives of air transport from atmospheric boundary
- layer to tropopause layer over Asian monsoon regions dur-990 930 ing boreal summer inferred from Lagrangian approach, Atmos. Chem. Phys., 12, 5827-5839, doi:http://dx.doi.org/10.5194/acp-12-5827-201210.5194/acp-12-5827-2012, 2012.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Bal-995 935 samo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Koehler, M.,
- Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Mor7000 940 crette, J. J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Q. J. Roy. Meteor. Soc., 137, 553-597,
- doi:http://dx.doi.org/10.1002/qj.82810.1002/qj.828, 2011. 945 1005 Dethof, A., O'Neill, A., Slingo, J. M., and Smit, H. G. J.: A mechanism for moistening the lower stratosphere involving the Asian summer monsoon, Q. J. Roy. Meteor. Soc., 556, 1079-1106, 1999.
- Dunkerton, T. J.: Evidence of meridional motion in the summer 950 lower stratosphere adjacent to monsoon regions, J. Geophys.

Res., 100, 16675-16688, 1995.

965

970

- Dvortsov, V. L. and Solomon, S.: Response of the stratospheric temperatures and ozone to past and future increases in stratospheric humidity, J. Geophys. Res., 106, 7505-7514, 2001.
- Emanuel, K.: Tropical cyclons, Annu. Rev. Earth Pl. Sc., 3, 75-104, 2003.
- Engel, A., Bönisch, H., Brunner, D., Fischer, H., Franke, H., Günther, G., Gurk, C., Hegglin, M., Hoor, P., Königstedt, R., Krebsbach, M., Maser, R., Parchatka, U., Peter, T., Schell, D., Schiller, C., Schmidt, U., Spelten, N., Szabo, T., Weers, U., Wernli, H., Wetter, T., and Wirth, V.: Highly resolved observations of trace gases in the lowermost stratosphere and upper troposphere from the Spurt project: an overview, Atmos. Chem. Phys., 6, 283-301, doi:http://dx.doi.org/10.5194/acp-6-283-200610.5194/acp-6-283-2006, 2006.
- Fierli, F., Orlandi, E., Law, K. S., Cagnazzo, C., Cairo, F., Schiller, C., Borrmann, S., Di Donfrancesco, G., Ravegnani, F., and Volk, C. M.: Impact of deep convection in the tropical tropopause layer in West Africa: in-situ observations and mesoscale modelling, Atmos. Chem. Phys., 11, 201-214, doi:http://dx.doi.org/10.5194/acp-11-201-201110.5194/acp-11-201-2011, 2011.
- Forster, P. and Shine, K. P.: Stratospheric water vapour change as possible contributor to observed stratospheric cooling, Geophys. Res. Lett., 26, 3309-3312, doi:http://dx.doi.org/10.1029/1999GL01048710.1029/1999GL010487, 1999.
- Forster, P. and Shine, K. P.: Assessing the climate impact of trends in stratospheric water vapor, Geophys. Res. Lett., 29, 10-1-10-4 doi:http://dx.doi.org/10.1029/2001GL01390910.1029/2001GL013909, 2002.

Garny, H. and Randel, W. J .: Dynamic variability of the Asian monsoon anticyclone observed in potential vorticity and correlations with tracer distributions, J. Geophys. Res., 118, 13421-13433,

- doi:http://dx.doi.org/10.1126/science.121937110.1126/science.1219371,doi:http://dx.doi.org/10.1002/2013JD02090810.1002/2013JD020908, 2013.
 - Gettelman, A., Kinnison, D., Dunkerton, T. J., and Brasseur, G. P.: Impact of monsoon circulations on the upper troposphere and lower stratosphere, J. Geophys. Res., 109, D22101, doi:http://dx.doi.org/10.1029/2004JD00487810.1029/2004JD004878, 2004.
 - Griessbach, S., Hoffmann, L., Spang, R., von Hobe, M., Müller, R., and Riese, M.: MIPAS Volcanic Sulfate Aerosol Observations of the Nabro Eruption, in: Stratospheric Sulfur and its Role in Climate (SSiRC), Atlanta, Georgia, USA, October 2013, SPARC, available at: http://www.sparc-ssirc.org/downloads/Griessbach. pdf, 2013.
 - Highwood, E. J. and Hoskins, B. J.: The tropical tropopause, Q. J. Roy. Meteor. Soc., 124, 1579–1604, 1998.
 - Holton, J. R., Haynes, P., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, Rev. Geophys., 33, 403-439, 1995.
 - Homeyer, C. R., Bowman, K. P., Pan, L. L., Atlas, E. L., Gao, R.-S., and Campos, T. L.: Dynamical and chemical characteristics of tropospheric intrusions observed during START08, J. Geophys. Res., 116, D06111. doi:http://dx.doi.org/10.1029/2010JD01509810.1029/2010JD015098, 2011.

- Hoor, P., Gurk, C., Brunner, D., Hegglin, M. I., Wernli, H., and Fischer, H.: Seasonality and extent of extratropical TST derived¹⁰⁷⁰ from in-situ CO measurements during SPURT, Atmos. Chem. Phys., 4, 1427–1442, doi:http://dx.doi.org/10.5194/acp-4-1427-200410.5194/acp-4-1427-2004, 2004.
- Hsu, C. J. and Plumb, R. A.: Nonaxisymmetric thermally driven circulations and upper-tropospheric monsoon dynamics, J. Atmos.¹⁰⁷⁵ Sci., 57, 1255–1276, 2001.
- Huang, Y., Zhang, W., Zheng, X., Li, J., and Yu, Y.: Modeling methane emission from rice paddies with various agricultural practices, J. Geophys. Res., 109, D8113, doi:http://dx.doi.org/10.1029/2003JD00440110.1029/2003JD0044401, 2004.
 - Hurst, D. F., Oltmans, S. J., Vömel, H., Rosenlof, K. H., Davis, S. M., Ray, E. A., Hall, E. G., and Jordan, A. F.: Strato-
- 1025spheric water vapor trends over Boulder, Colorado: analysis of
the 30 year Boulder record, J. Geophys. Res., 116, D02306,1085
doi:http://dx.doi.org/10.1029/2010JD01506510.1029/2010JD015065,Liu, Y., Wang, Y., Liu, X., Cai, Z., and Chance, K.: Tibetan
middle tropospheric ozone minimum in June discovered1025
- Jackson, D. R., Driscoll, S. J., Highwood, E. J., Harries, J. E., and Russell III, J. M.: Troposphere to stratosphere transport at low latitudes as studies using HALOE observations of water vaponoso 1992–1997, Q. J. Roy. Meteor. Soc., 124, 169–192, 1998.
 - Kaufmann, M., Blank, J., Guggenmoser, T., Ungermann, J., Engel, A., Ern, M., Friedl-Vallon, F., Gerber, D.,, Grooß, J. U.,
- Günther, G., Höpfner, M., Kleinert, A., Latzko, T., Maucher, G., Neubert, T., Nordmeyer, H., Oelhaf, H., Olschewski, F.,1095 Orphal, J., Preusse, P., Schlager, H., Schneider, H., Schuettemeyer, D., Stroh, F., Sumińska-Ebersoldt, O., Vogel, B., Volk, M., Wintel, J., Woiwode, W., and Riese, M.: Re-
- trieval of three-dimensional small scale structures in upper tropospheric/lower stratospheric composition as measured₁₁₀₀ by GLORIA, Atmos. Meas. Tech. Discuss., 7, 4229–4274, doi:http://dx.doi.org/10.5194/amtd-7-4229-201410.5194/amtd-7-4229-2014, 2014.
- Khalil, M. A. K., Rasmussen, R. A., Shearer, M. J., Dalluge, R. W., Ren, L. X., and Duan, C. L.: Measurements of methane emissions105 from rice fields in China, J. Geophys. Res., 103, 25181–25210, 1998.
- Kirk-Davidoff, D. B., Hintsa, E. J., Anderson, J. G., and Keith, D. W.: The effect of climate change on ozone depletion through changes in stratospheric water vapour, Nature, 402, 399–110 401, doi:http://dx.doi.org/10.1038/4652110.1038/46521, 1999.
- Konopka, P., Grooß, J.-U., Günther, G., Ploeger, F., Pommrich, R., Müller, R., and Livesey, N.: Annual cycle of ozone
- at and above the tropical tropopause: observations versus simulations with the Chemical Lagrangian Model of theiris Stratosphere (CLaMS), Atmos. Chem. Phys., 10, 121–132, doi:http://dx.doi.org/10.5194/acp-10-121-201010.5194/acp-10-121-2010, 2010.
- Konopka, P., Ploeger, F., and Müller, R.: Entropy-based and static stability based Lagrangian model grids, in: Lagrangian Modeling₁₁₂₀ of the Atmosphere, Geophysical Monograph Series, 200, AGU Chapman Conference on Advances in Lagrangian Modeling of the Atmosphere, Grindelwald, Switzerland, 9–14 October 2011, 99–109, 2012.
- Kossin, J. P., Emanuel, K. A., and Vecchi, G. A.: The₁₂₅ poleward migration of the location of tropical cyclone maximum intensity, Nature, 509, 349–352,

doi:http://dx.doi.org/10.1038/nature1327810.1038/nature13278, 2014.

- Kunz, A., Müller, R., Homonnai, V., Jánosi, I., Hurst, D., Rap, A., Forster, P., Rohrer, F., Spelten, N., and Riese, M.: Extending water vapor trend observations over Boulder into the tropopause region: trend uncertainties and resulting radiative forcing, J. Geophys. Res., 118, 11269–11284, doi:http://dx.doi.org/10.1002/jgrd.5083110.1002/jgrd.50831, 2013.
- Li, Q., Jiang, J. H., Wu, D. L., Read, W. G., Livesey, N. J., Waters, J. W., Zhang, Y., Wang, B., Filipiak, M. J., Davis, C. P., Turquety, S., Wu, S., Park, R. J., Yantosca, R. M., and Jacob, D. J.: Convective outflow of South Asian pollution: a global CTM simulation compared with EOS MLS observations, Geophys. Res. Lett., 32, L14826, doi:http://dx.doi.org/10.1029/2005GL02276210.1029/2005GL022762, 2005.
- 5, Liu, Y., Wang, Y., Liu, X., Cai, Z., and Chance, K.: Tibetan middle tropospheric ozone minimum in June discovered from GOME observations, Geophys. Res. Lett., 36, L05814, doi:http://dx.doi.org/10.1029/2008GL03705610.1029/2008GL037056, 2009.
- McKenna, D. S., Grooß, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 2. Formulation of chemistry scheme and initialization, J. Geophys. Res., 107, 4256, doi:http://dx.doi.org/10.1029/2000JD00011310.1029/2000JD000113, 2002a.
- McKenna, D. S., Konopka, P., Grooß, J.-U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 1. Formulation of advection and mixing, J. Geophys. Res., 107, 4309, doi:http://dx.doi.org/10.1029/2000JD00011410.1029/2000JD000114, 2002b.
- Munchak, L., Fan, Q., Pan, L., Bian, J., and Bowman, K.: An analysis of transport pathways that contribute to water vapor and ozone profiles measured in the Asian monsoon anticyclone, in: Fall Meeting, Fall Meeting, 13–17 December 2010, AGU, San Francisco, A51B-0080, 2010.
- Orlanski, I. and Sheldon, J. P.: Stages in the energetics of baroclinic systems, Tellus A, 47, 605–628, 1995.
- Park, M., Randel, W. J., Kinnison, D. E., Garcia, R. R., and Choi, W.: Seasonal variation of methane, water vapor, and nitrogen oxides near the tropopause: Satellite observations and model simulations, J. Geophys. Res., 109, D03302, doi:http://dx.doi.org/10.1029/2003JD00370610.1029/2003JD003706, 2004.
- Park, M., Randel, W. J., Gettleman, A., Massie, S. T., and Jiang, J. H.: Transport above the Asian summer monsoon anticyclone inferred from Aura Microwave Limb Sounder tracers, J. Geophys. Res., 112, D16309, doi:http://dx.doi.org/10.1029/2006JD00829410.1029/2006JD008294, 2007.
- Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A., and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, Atmos. Chem. Phys., 8, 757– 764, doi:http://dx.doi.org/10.5194/acp-8-757-200810.5194/acp-8-757-2008, 2008.

- Park, M., Randel, W. J., Emmons, L. K., and Livesey, N. J.: Transport pathways of carbon monoxide in the Asian summer monsoon diagnosed from Model of Ozone and Related Tracers (MOZART), J. Geophys. Res., 114, D08303, doi:http://dx.doi.org/10.1029/2008JD01062110.1029/2008JD010621, 2009.
- F., Konopka, P., Günther, G., Grooß, J.-Ploeger, U., and Müller, R.: Impact of the vertical veloc-1135 ity scheme on modeling transport across the tropical tropopause layer, J. Geophys. Res., 115, D03301,1195 doi:http://dx.doi.org/10.1029/2009JD01202310.1029/2009JD012023, 2010.
- Ploeger, F., Günther, G., Konopka, P., Fueglistaler, S., 1140 Müller, R., Hoppe, C., Kunz, A., Spang, R., Grooß, J.-U., and Riese, M.: Horizontal water vapor transport in the1200 lower stratosphere from subtropics to high latitudes during boreal summer, J. Geophys. Res., 118, 8111-8127, doi:http://dx.doi.org/10.1002/jgrd.5063610.1002/jgrd.50636, 1145 2013.
 - Pommrich, R., Müller, R., Grooß, J.-U., Konopka, P., Ploeger, P., 1205 Vogel, B., Tao, M., Hoppe, C., Günther, G., Hoffmann, L., Pumphrey, H.-C., Viciani, S., D'Amato, F., Volk, M.,

Hoor, P., Schlager, H, and Riese, M.: Tropical tropo-1150 sphere to stratosphere transport of carbon monoxide and long-lived trace species in the Chemical Lagrangian Model₂₁₀ of the Stratosphere (CLaMS), Geosci. Model Dev. Discuss., 7, 5087-5139,doi:http://dx.doi.org/10.5194/gmdd-7-5087-201410.5194/gmdd-7-5087-2014, 2014. 1155

Popovic, J. M. and Plumb, R. A.: Eddy Shedding from the uppertropospheric Asian monsoon anticyclone, J. Atmos. Sci., 58, 93-1215 104, 2001.

- Randel, W. J. and Park, M.: Deep convective influence on the Asian summer monsoon anticyclone and asso-1160 ciated tracer variability observed with Atmospheric Infrared Sounder (AIRS), J. Geophys. Res., 111, D12314,1220 doi:http://dx.doi.org/10.1029/2005JD00649010.1029/2005JD006490, 2006.
- Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., 1165 Walker, K. A., Boone, C., and Pumphrey, H.: Asian monsoon transport of pollution to the stratosphere, Science, 328, 611–613,1225 doi:http://dx.doi.org/10.1126/science.118227410.1126/science.1182274, Campos, T., Pollack, I., Weinheimer, A., Wei, J., Atlas, E. L., 2010.
- Riese, M., Grooß, J.-U., Feck, T., and Rohs, S.: Long-term changes 1170 of hydrogen-containing species in the stratosphere, J. Atmos. Sol-Terr. Phys., 68, 1973-1979, 2006. 1230
- Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS composition and 1175 related radiative effects, J. Geophys. Res., 117, D16305, doi:http://dx.doi.org/10.1029/2012JD01775110.1029/2012JD017251, 2012.
- Riese, M., Oelhaf, H., Preusse, P., Blank, J., Ern, M., Friedl-Vallon, F., Fischer, H., Guggenmoser, T., Höpfner, M., Hoor, P., Kauf-1180 mann, M., Orphal, J., Plöger, F., Spang, R., Suminska-Ebersoldt, O., Ungermann, J., Vogel, B., and Woiwode, W.: Gimballed₁₂₄₀ Limb Observer for Radiance Imaging of the Atmosphere (GLO-RIA) scientific objectives, Atmos. Meas. Tech., 7, 1915-1928, doi:10.5194/amt-7-1915-2014, 2014. 1185

Röckmann, T., Grooß, J.-U., and Müller, R.: The impact of anthropogenic chlorine emissions, stratospheric ozone change and chemical feedbacks on stratospheric water, Atmos. Chem. Phys., 4, 693-699, doi:http://dx.doi.org/10.5194/acp-4-693-200410.5194/acp-4-693-2004, 2004.

- Rohs, S., Schiller, C., Riese, M., Engel, A., Schmidt, U., Wetter, T., Levin, I., Nakazawa, T., and Aoki, S.: Long-term changes of methane and hydrogen in the stratosphere in the period 1978-2003 and their impact on the abundance of stratospheric water vapor, J. Geophys. Res., 111, D14315, doi:http://dx.doi.org/10.1029/2005JD00687710.1029/2005JD006877, 2006.
- Rosenlof, K. H., Tuck, A. F., Kelly, K. K., Russell III, J. M., and McCormick, M. P.: Hemispheric asymmetries in the water vapor and inferences about transport in the lower stratosphere, J. Geophys. Res., 102, 13213-13234, 1997.
- Schiller, C., Krämer, M., Afchine, A., Spelten, N., and Sitnikov, N.: The ice water content of Arctic, mid latitude and tropical cirrus, J. Geophys. Res., 113, D24208, doi:http://dx.doi.org/10.1029/2008JD01034210.1029/2008JD010342, 2008.
- Shindell, D. T.: Climate and ozone response to increased stratospheric water vapor, Geophys. Res. Lett., 28, 1551-1554, doi:http://dx.doi.org/10.1029/1999GL01119710.1029/1999GL011197, 2001
- Smith, C. A., Haigh, J. D., and Toumi, R.: Radiative forcing due to trends in stratospheric water vapour, Geophys. Res. Lett., 28, 179–182, 2001.
- Solomon, S., Rosenlof, K., Portmann, R., Daniel, J., Davis, S., Sanford, T., and Plattner, G.-K.: Contributions of stratospheric water vapor to decadal changes in the rate of global warming, Science, 327, 1219-1223, doi:http://dx.doi.org/10.1126/science.118248810.1126/science.1182488, 2010
- Vogel, B., Feck, T., and Grooß, J.-U.: Impact of stratospheric water vapor enhancements caused by CH₄ and H₂ increase on polar ozone loss, J. Geophys. Res., 116, D05301, doi:http://dx.doi.org/10.1029/2010JD01423410.1029/2010JD014234, 2011a.
- Vogel, B., Pan, L. L., Konopka, P., Günther, G., Müller, R., Hall, W.,
 - and Bowman, K. P.: Transport pathways and signatures of mixing in the extratropical tropopause region derived from Lagrangian model simulations, J. Geophys. Res., 116, D05306, doi:http://dx.doi.org/10.1029/2010JD01487610.1029/2010JD014876, 2011b.
 - Vogel, B., Feck, T., Grooß, J.-U., and Riese, M.: Impact of a possible future global hydrogen economy on Arctic stratospheric ozone loss, Energy Environ. Sci., 5, 6445-6452, doi:http://dx.doi.org/10.1039/c2ee03181g10.1039/c2ee03181g, 2012.
 - Xiong, X., Houweling, S., Wei, J., Maddy, E., Sun, F., and Barnet, C.: Methane plume over south Asia during the monsoon season: satellite observation and model simulation, Atmos. Chem. Phys., 9, 783-794, doi:http://dx.doi.org/10.5194/acp-9-783-200910.5194/acp-9-783-2009, 2009.
 - Zahn, A., Weppner, J., Widmann, H., Schlote-Holubek, K., Burger, B., Kühner, T., and Franke, H.: A fast and precise chemiluminescence ozone detector for eddy flux

1130

- and airborne application, Atmos. Meas. Tech., 5, 363–375, doi:http://dx.doi.org/10.5194/amt-5-363-201210.5194/amt-5-363-2012, 2012.
 - Zöger, M., Schiller, C., and Eicke, N.: Fast in situ hygrometers: a new family of balloonborne and airborne Lyman- α photofrag-
- ¹²⁵⁰ ment fluorescence hygrometers, J. Geophys. Res., 104, 1807– 1816, 1999.

B. Vogel et al.: Fast transport from Asian monsoon anticyclone to Europe

PV [PVU]

Table 1. Number (N_{trajs}) and percentage of all trajectories in the region of interest (in total 432 trajectories), mean and maximum vertical velocity (calculated between 17 and 27 August 2012) along 40 day backward trajectories depending on levels of potential temperature at the air mass origin (Θ_{org}). The 40 day backward trajectories starting at the observation on 26 September 2012 and ending at the origin of the air masses in the past (17 August 2012). The air mass origins in terms of altitude and geographical position are also listed. table

Θ_{org}	N_{trajs}	precentage	mean vertical velocity	max. vertical velocity	altitude origin	geographical position origin
295–320 K	8	2%	very rapid uplift: 31 K day ⁻¹	$41 \mathrm{K}\mathrm{dav}^{-1}$	boundary layer	Southeast Asia
			$(= 371 \text{ hPaday}^{-1})$	$(= 523 \mathrm{hPadav}^{-1})$		
			$(= 6.6 \mathrm{km} \mathrm{day}^{-1})$	$(=9.4 \mathrm{km}\mathrm{day}^{-1})$		
			$13 \mathrm{K}/6 \mathrm{h}$	21 K/6 h		
			(= 144 hPa/6 h)	(= 276 h Pa/6 h)		
			$(= 2.5 \mathrm{km}/6 \mathrm{h})$	$(= 3.7 \mathrm{km}/6 \mathrm{h})$		
320–360 K	15	3%	rapid uplift:		troposphere	mainly West Pacific
			8 K day ⁻¹	13 K day ⁻¹		
			$(= 63 \mathrm{hPa} \mathrm{day}^{-1})$	$(= 139 h Pa day^{-1})$		
			$(= 2.3 \mathrm{km}\mathrm{day}^{-1})$	$(= 4.4 \mathrm{km}\mathrm{day}^{-1})$		
			3K/6R	$\frac{\partial K}{\partial n}$		
			(= 31 HFa/6 h)	$(= 00 \Pi F a / 0 \Pi)$ (= 1.0 km / 6 h)		
360-370 K	50	12%	(= 1.1 Km/0 m) moderately rapid unlift	(= 1.3 km/0 m)	AM anticyclone	mainly South Asia/North Africa
500 570 R	50	12 /0	$2 \mathrm{K} \mathrm{dav}^{-1}$	$5 \mathrm{K}\mathrm{dav}^{-1}$	7 nor unite yerone	manny South Asian torth Annea
			$(= 22 \mathrm{hPa}\mathrm{dav}^{-1})$	$(=41 \mathrm{hPa}\mathrm{dav}^{-1})$		
			$(= 1.0 \mathrm{km} \mathrm{day}^{-1})$	$(= 1.8 \mathrm{km}\mathrm{day}^{-1})$		
			1 K/6 h	$2\mathrm{K}/6\mathrm{h}$		
			(= 11 hPa/6 h)	(= 25 hPa/6 h)		
			(= 0.5 km/6 h)	(= 1.1 km/6 h)		
370–380 K	97	22 %	moderate uplift:		UTLS	mainly edge of the AM anticyclone
			$1 \mathrm{K}\mathrm{day}^{-1}$	$3 \mathrm{K}\mathrm{day}^{-1}$		
			$(= 16 h Pa day^{-1})$	$(=45 \mathrm{hPa}\mathrm{day}^{-1})$		
			$(= 0.8 \mathrm{km}\mathrm{day}^{-1})$	$(= 2.3 \mathrm{km}\mathrm{day}^{-1})$		
			0.5 K/6 h	1 K/6 h		
			(= 0.1 ra/0.1)	(= 20 mPa/6 n)		
380–420 K	262	61 %	(= 0.4 km/6 h) mainly descent	$(= 1.0 \mathrm{km}/6 \mathrm{n})$	lower stratosphere	Northern Hemisphere

26.09.12 12:00:00 θ= 375 K



Fig. 1. The horizontal distribution of PV at 375 K potential temperature over Northern Europe on 26 September 2012. The flight path for the TACTS flight on 26 September 2012 (at flight time) starting and ending in Oberpfaffenhofen (near Munich, 48° N 11° E, Germany) is marked in black. The part of the flightpath discussed within this paper is highlighted in white (cf. Sect. 2 and see Fig. 2). The horizontal winds are indicated by white arrows. PV and horizontal winds are from ERA-Interim reanalysis data. figure



Fig. 2. Time evolution (given in Universal Time Coordinated (UTC)) of potential temperature, CO, O_3 , CH₄, and H₂O measurements for the flight on 26 September 2012. Potential temperature values at the end points of 40 day backward trajectories (17 August 2012) are shown as red dots (upper panel). The part of the flightpath discussed within this paper is highlighted in grey (region of interest) (cf. Sect. 2 and see Fig. 1). Trajectories with origins (Θ_{org}) lower than 360 K potential temperature in the region of interest are shown in Fig. 3.



Fig. 3.

Fig. 3. Different 40 day backward trajectories for Θ_{org} intervals 295–320 K (left) and 320–360 K (right) are shown colour-coded by potential temperature (top) and by days reversed from 26 September 2012 (middle). Further, potential temperature vs. time (in UTC) along 40 day backward trajectories (bottom) are shown. Here, the colour indicate the latitude position of the trajectories. The trajectory positions are plotted every hour (coloured dots). Large distances between single dots indicated very rapid uplift. The geographical position of the origins (Θ_{org}) of the 40 day backward trajectories (red dots) are shown in the embedded longitude-latitude-cut. It should be noted that these geographical positions are at levels of potential temperature in the given Θ_{org} intervals and not at surface level.



Fig. 4. The geographical position of the origins (Θ_{org}) of the 40 day backward trajectories for intervals 360–370 K (top), 370–380 K (middle) and 380–420 K (bottom) are shown colour-coded by potential temperature. The corresponding trajectories are shown in the electronic supplement of this paper.





Fig. 5. Geopotential height in [m] (colour) and horizontal winds in $[ms^{-1}]$ (white arrows) at 500 hPa on 24 August 2012 taken from ERA-Interim reanalysis data (top). To show the position of the typhoons Tembin ($\approx 23^{\circ}$ N 118° E) and Bolaven ($\approx 23^{\circ}$ N 135° E) the mean sea level pressure in [hPa] is shown as blue thick lines. Longitude-height cross-section showing the typhoon Tembin and Bolaven at 23° N (bottom). The vertical velocity (ω) in [hPas⁻¹] (colour), the horizontal winds in [ms⁻¹] (white line (positive values) and white dashed line (negative values)), and potential temperature in Kelvin (black lines) are shown. Very rapid uplift up to $-0.014 hPas^{-1}$ is found at the eastern flank of typhoon Bolaven ($\approx 135^{\circ}$ E) and up to $-0.006 hPas^{-1}$ is found at western flank of typhoon Tembin ($\approx 118^{\circ}$ E).



Fig. 6. Selected forward trajectories starting in the boundary layer around the position the typhoon Bolaven on 23 August 2012 12:00 UTC $(15^{\circ}N-25^{\circ}N, 128^{\circ}E-140^{\circ}E \text{ and below} \approx 310 \text{ K} \text{ potential temperature})$. The trajectories are started at a horizontal grid of $0.5^{\circ} \times 0.5^{\circ}$ and a vertical grid of 1 K. Shown are only trajectories that experience a strong uplift greater than 50 K within the first 48 hours caused by typhoon Bolaven. For simplification only a subset of the trajectories is shown (every 100th trajectory).



Fig. 7. Time sequence of PV fields (colour shading in PVU) over Asia at 370 K potential temperature from 9 to 20 September 2012 based on ERA-Interim reanalysis data. The horizontal winds are indicated by white arrows. To show the position of tropospheric low-pressure systems in particular the super typhoon Sanba (e.g. $\approx 20^{\circ}$ N 130° E on 14 September, $\approx 25^{\circ}$ N 130° E on 15 September, $\approx 30^{\circ}$ N 130° E on 16 September, and $\approx 40^{\circ}$ N 130° E on 17 September) the sea level pressure of 1000 hPa is marked as thick purple line.



Fig. 7. Continued.



Fig. 8. Forward trajectories starting at the position of the typhoon Bolaven $(15^{\circ}N-25^{\circ}N, 128^{\circ}E-140^{\circ}E)$ and below ≈ 310 K potential temperature) marked as grey box on 23 August 2012 12:00 UTC. Shown are only trajectories that reach the region of interest during the flight on 26 September 2012 12:00 UTC (at locations between 365 K and 385 K, between 55^{\circ}N and 70^{\circ}N, and between 20^{\circ}W and 0^{\circ}W). The flightpath shown in black is transformed to synoptic 12:00 UTC locations.



Fig. 9. Different 60 day backward trajectories for the tropospheric signal between 08:05 UTC and 08:23 UTC (see Fig. 2; enhanced values of CH₄, CO, and H₂O and reduced O₃ occurring before the region of interest) originating at low level of potential temperature ($\Theta_{org} = 295-320$ K). The very rapid uplift over Southeast Asia is caused by typhoon Haikui.



Fig. 10. Selected trajectories representing characteristic pathways with different vertical velocities (cf. Table 1) as function of potential temperature and longitude for the time period between 17 August and 26 September 2012. The air mass positions are plotted every hour (coloured dots). Large distances between the single dots indicated very rapid uplift. The classification of the different vertical velocities is derived for the first ten days of the trajectories (17–28 August 2012). For simplification longitudes between 60° W and 20° E are shown twice. The coloured text gives information about the altitude origin and geographical position origin of the different pathways. The figure illustrates different transport pathways of air masses with different origin to the location of the measurement over Northern Europe. The combination of very rapid uplift by a typhoon and eastward eddy shedding from the Asian monsoon anticyclone yield fast transport (≈ 5 weeks) from Southeast Asia boundary layer sources to Northern Europe (red).