Long term halocarbon observations from a coastal and an inland site in Sabah, Malaysian Borneo

- 3
- 4 A. D. Robinson¹, N. R. P. Harris¹, M. J. Ashfold^{1,*}, B. Gostlow¹, N. J. Warwick^{1,2}, L. M.
- 5 O'Brien¹, E. J. Beardmore¹, M. S. M. Nadzir^{3,†}, S. M. Phang³, A. A. Samah³, S. Ong⁴, H.
- 6 E. Ung⁴, L. K. Peng⁵, S. E. Yong⁵, M. Mohamad⁵ and J. A. Pyle^{1,2}
- 7
- 8 [1] {Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW,
 9 UK}
- 10 [2] {National Centre for Atmospheric Science, NCAS, UK}
- 11 [3] {Institute of Ocean & Earth Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia}
- [4] {Global Satria Life Sciences Lab, TB 12188, Taman Megajaya Phase 3, 91000 Tawau, Sabah,
 Malaysia}
- 14 [5] {Malaysian Meteorological Department, Ketua Stesen GAW Lembah Danum, Jabatan
- 15 Meteorologi Malaysia Cawangan Sabah, Lapangan Terbang Wakuba Tawau, Peti Surat 60109,
- 16 91011 Tawau, Sabah, Malaysia}
- 17 [*] {now at: School of Biosciences, University of Nottingham Malaysia Campus, Jalan Broga,
- 18 43500 Semenyih, Selangor Darul Ehsan, Malaysia}
- 19 [[†]] {now at: School of Environmental and Natural Resource Sciences, Faculty of Science and
- 20 Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia}
- 21 Correspondence to: A.D. Robinson (adr22@cam.ac.uk)

22 Abstract

Short lived halocarbons are believed to have important sources in the tropics where rapid vertical transport could provide a significant source to the stratosphere. In this study, quasi-continuous measurements of short-lived halocarbons are reported for two tropical sites in Sabah (Malaysian Borneo), one coastal and one inland (rainforest). We present the observations for C₂Cl₄, CHBr₃, CH₂Br₂^{*} (actually ~80 % CH₂Br₂ and ~20 % CHBrCl₂) and CH₃I from November 2008 to January

1 2010 made using our µDirac gas chromatographs with electron capture detection (GC-ECD). We 2 focus on the first 15 months of observations, showing over one annual cycle for each compound and therefore adding significantly to the few limited-duration observational studies that have been 3 4 conducted thus far in southeast Asia. The main feature in the C₂Cl₄ behaviour at both sites is its 5 annual cycle with the winter months being influenced by northerly flow with higher concentrations, typical of the northern hemisphere, with the summer months influenced by southerly flow and 6 7 lower concentrations representative of the southern hemisphere. No such clear annual cycle is seen for CHBr₃, CH₂Br₂^{*} or CH₃I. The baseline values for CHBr₃ and CH₂Br₂^{*} are similar at the coastal 8 (overall median: CHBr₃ 1.7 ppt; CH₂Br₂^{*} 1.4 ppt) and inland sites (CHBr₃ 1.6 ppt, CH₂Br₂^{*} 1.1 9 ppt), but periods with elevated values are seen at the coast (overall 95th percentile: CHBr₃ 4.4 ppt; 10 CH₂Br₂^{*} 1.9 ppt) presumably resulting from the stronger influence of coastal emissions. Overall 11 median bromine values from $[CHBr_3 \times 3] + [CH_2Br_2^* \times 2]$ are 8.0 ppt at the coast and 6.8 ppt 12 13 inland. The median values reported here are largely consistent with other limited tropical data and 14 imply that southeast Asia generally is not, as has been suggested, a hot-spot for emissions of these 15 compounds. These baseline values are consistent with the most recent emissions found for southeast 16 Asia using the p-TOMCAT model. CH₃I, which is only observed at the coastal site, is the shortest-17 lived compound measured in this study and the observed atmospheric variations reflect this, with 18 high variability throughout the study period.

19 **1** Introduction

20 Reactions involving halogen compounds play an important role in the chemistry of both the 21 stratosphere (e.g. Molina and Rowland, 1974; Yung et al., 1980) and the troposphere (e.g. Read et 22 al., 2008). The sources of halogens are many and varied with important anthropogenic and natural 23 components for many compounds (Montzka and Reimann, 2011). Total atmospheric bromine concentrations are generally dominated by the halons (compounds of only anthropogenic origin 24 25 widely used for fire control) and CH₃Br (used in agriculture, but with many sources). Their atmospheric concentrations are well known and their lifetimes long enough to know with 26 27 reasonable accuracy how much of these gases reaches the stratosphere. It was thus puzzling to find 28 that they could only account for 70-80 % of the bromine observed in the stratosphere (Wamsley et 29 al., 1998; Dorf et al., 2006) though there remains uncertainty as to whether the discrepancy is that 30 large (Kreycy et al., 2013).

Since there are no other known long-lived sources, the finger of suspicion points to the shorter-lived
 CHBr₃ and CH₂Br₂ which are observed in high concentration in the marine boundary layer and are

1 naturally produced in the surface ocean (e.g. Yokouchi et al., 2005; Butler et al., 2007; Zhou et al., 2 2008; O'Brien et al., 2009). For these compounds to be the source of the missing 20-30% of the stratospheric bromine budget, they (or their breakdown products) have to be rapidly transported up 3 4 to the stratosphere. The only credible place for this to occur is at low latitudes where upward 5 transport can be very rapid in the intense tropical convection. This possibility has been investigated 6 in a number of measurement (Schauffler et al., 1999; Laube et al., 2008; Park et al., 2010; 7 Brinckmann et al., 2012) and modelling (Levine et al., 2007, 2008; Aschmann et al., 2009; Hosking 8 et al., 2010; Hossaini et al., 2010, 2012 & 2013) studies and overall it seems likely that the short-9 lived bromocarbons are responsible for the apparent deficit.

One of the limitations on these studies is the scarcity of high quality measurements in the regions of most importance, i.e. in the highly convective areas. The majority of reported tropical boundary layer measurements have been made using whole air samplers or short-term deployments of GC systems (Yokouchi et al., 2005; O'Brien et al., 2009; Pyle et al., 2011), or ship-borne measurements made during research cruises (Butler et al., 2007; Carpenter et al., 2009; Quack et al., 2004, 2007; Yokouchi et al., 1997; Nadzir et al., 2014; Brinckmann et al., 2012). No long-term and continuous measurements in the tropics have been reported to date.

17 Accordingly, we have set up a small network of instruments to make continuous measurements at a 18 number of sites in the West Pacific / southeast Asian region. This region was chosen as it is where 19 convection is strongest, especially during NH winter (e.g. Newell and Gould-Stewart, 1981; 20 Gettelman et al., 2002; Fueglistaler et al., 2009). In addition, an early model study suggested high 21 CHBr₃ emissions and atmospheric concentrations here, due to the many islands and therefore large 22 length of coastlines found in this region (Warwick et al, 2006), although this estimate has since 23 been revised downward (Pyle et al., 2011). The first two instruments were installed in 2008 in the 24 State of Sabah in Malaysian Borneo, one at a coastal site and one inland (Figure 1). Early measurements were reported in Pyle et al. (2011). Here we report measurements from the 25 26 subsequent years focussing on the period from late 2008 to early 2010. In addition to CHBr₃ and $CH_2Br_2^*$, we report our observations of C_2Cl_4 , a manufactured compound which we use to identify 27 air with anthropogenic influence, and CH₃I, a compound produced naturally in the ocean which has 28 29 a short atmospheric lifetime (7 days, Montzka and Reimann, 2011).

In Section 2 we describe the instruments used in this study, their calibration and the measurement quality reported here. Section 3 contains a discussion of the measurement sites including their physical characteristics, while in Section 4 we describe the modelling tools (NAME, p-TOMCAT) 1 used in the interpretation of the data. The measurements for C_2Cl_4 , CHBr₃, CH₂Br₂^{*} and CH₃I are 2 presented in Section 5, with particular emphasis on their variability, and with air mass history maps 3 to aid in interpretation. We also consider the implications of the measurements, including an 4 analysis of the total organic bromine in short-lived bromocarbons at each site. Finally, Section 6 5 summarises our findings and ongoing/future work.

6 2 Methodology

7 2.1 Instrument

8 The instruments used for these observations were the original µDirac gas chromatographs built by 9 us for use on balloons, with electron capture detection (GC-ECD), in a modified configuration 10 appropriate for extended ground-based deployment as described in Pyle et al. (2011). The 11 instrument is described in detail in Gostlow et al. (2010). Briefly, each sample is pre-concentrated using a dual-bed adsorbent trap (1 mg Carboxen-1016 60/80 mesh followed by 1 mg Carboxen-569 12 20/45 mesh). The trap is then heated and the gaseous constituents are separated on a 10 m long, 13 14 0.18 mm i.d. capillary column (Restek MXT 502.2) prior to measurement in the ECD. As the 15 instruments are designed to run unattended, the control software runs chromatograms according to a pre-determined sequence of samples bracketed by calibration and blank chromatograms. The 16 17 sequence includes frequent calibrations using the same volume as the samples (20 ml) for correction 18 of instrument sensitivity drift and precision determination. The sequence also runs calibrations at a 19 range of different volumes so that instrument response curves are generated for each target compound to allow for non-linearity. The response curves are fit using a third order polynomial. 20 21 Chromatograms and system data are stored on a host computer for later analysis using in-house 22 software to determine peak heights for the target compounds which are then converted into mixing 23 ratios by comparison with the calibration standards. Mixing ratios are reported as dry air mole 24 fraction in ppt.

Measurement performance is monitored for each target compound during the analysis of the chromatograms, and the typical performance for the measurement period reported here is summarised in Table 1. For C₂Cl₄ the typical measurement precision (1 s.d.) is 2.5 % and 4.0 % at Bukit Atur and Tawau respectively. The typical measurement precision of CHBr₃ is 6.0 % and 7.5 % at Bukit Atur and Tawau respectively and those for $CH_2Br_2^*$ are 5.0 % and 6.5 %. Sample concentrations for C₂Cl₄, CHBr₃ and CH₂Br₂^{*} are always above the limit of detection at both sites. CH₃I is only measured at Tawau (the µDirac at Bukit Atur did not achieve satisfactory chromatographic separation for reliable quantification) and the typical precision is 11 %. The CH₃I peak is small in the sample chromatograms so particular care is taken in assessing its signal to noise ratio (SNR). The SNR varies over time depending on the instrument condition and environment, and is calculated separately for each analysis period (typically 1-2 weeks) for which concentrations are calculated. Between January and June 2009, the estimated detection limit (defined using a SNR of 10) decreases from 0.9 ppt to 0.3 ppt as a result of reduced noise in the chromatograms and for the rest of 2009 it is below 0.2 ppt.

8 The chromatographic set-up uses a 10 m column for these measurements which provides a good 9 separation for C₂Cl₄ and CHBr₃. However, the CH₂Br₂ peak co-elutes with that of CHBrCl₂ and so we report a combined value called $CH_2Br_2^*$ which we estimate to be 10-30 % higher than the 10 11 CH₂Br₂ value alone (O'Brien et al., 2009). This co-elution problem is further discussed in section 12 5.3 of this paper and has been previously assessed in Gostlow et al. (2010). In a recently improved version of the instrument, where a 20 m long column is used, the CH₂BrCl₂ and CH₂Br₂ peaks are 13 14 partly separated and so individual values can be derived by using two overlapped Gaussian fits. By 15 analysing calibration chromatograms from three of our newer instruments using the longer column, we find that the ECD is significantly more sensitive to CH₂Br₂ than it is to CHBrCl₂, the peak 16 height per ppt for CHBrCl₂ being typically 80 % of that for the CH₂Br₂ peak, confirming the 17 assumption made in O'Brien et al. (2009). Measurements of CH₂Br₂ and CHBrCl₂ using 18 19 instruments with 20 m columns in similar tropical locations show the ratio (10-30 %) to be still 20 valid. For example, in a 2010 cruise from Singapore to New Zealand we found that, for the southeast Asia samples, the average CHBrCl₂ mixing ratio was 0.18 ppt, ~16 % of that for CH₂Br₂ 21 at 1.12 ppt (unpublished data). In this case, had a 10 m column been used, the $CH_2Br_2^*$ value from 22 23 the combined peaks would have been 1.26 ppt (allowing for the 80 % lower sensitivity to CHBrCl₂ 24 compared to CH₂Br₂). This compares favourably to the value of 1.30 ppt for the sum of the 25 separately calculated values for CH₂BrCl₂ and CH₂Br₂. Other studies in the southeast Asia region have also reported similar values for CHBrCl₂ and CH₂Br₂: during the OP3 campaign at Bukit Atur 26 27 in July 2008, GC-MS observations made by the University of East Anglia (Pyle et al., 2011a) found 28 an average CHBrCl₂ mixing ratio of 0.25 ppt, ~27 % of that for CH₂Br₂ (0.91 ppt); during a 2009 29 cruise in the Strait of Malacca, South China and Sulu-Sulawesi Seas where mixing ratios of CHBrCl₂ were on average 33 % of those for CH₂Br₂ (Nadzir et al., 2014). 30

The instrument based at the coastal site developed a problem with desorption efficiency caused by incomplete thermal desorption of the target molecules from the adsorbent. This affected the quantification of the $CH_2Br_2^*$ peak by carry over from one chromatogram to the next. However, we are still able to use calibration chromatograms for CH_2Br_2 as these are preceded by a blank chromatogram. About one in every ten sample chromatograms is also usable for the $CH_2Br_2^*$ peak as these are also preceded by a blank. As a result of this problem $CH_2Br_2^*$ is only reported for about 1 in every 10 samples at Tawau. The C_2Cl_4 and $CHBr_3$ peaks are not affected by this problem at Tawau as these molecules are always completely retained on the weaker Carboxen-1016 at the front of the dual adsorbent bed.

8 2.2 Calibration

9 Samples are calibrated by running frequent chromatograms from cylinders of clean natural air, 10 previously filled at Niwot Ridge (Colorado, USA), at an elevation of 3.5 km above sea level. The cylinder air is subsequently enriched with known concentrations of the target compounds. The 11 cylinders (one for each instrument) are supplied by the Earth System Research Laboratory (ESRL) 12 13 within the National Oceanic and Atmospheric Administration (NOAA) so that the reported mixing ratios are linked directly to the NOAA halocarbon calibration scales. Figure 2 summarises the 14 15 period of data coverage for this study together with a time-line of the calibration cylinders in use at the two sites. The calibration gas in use at Tawau for the entire period of this study is in a high 16 17 quality, humidified stainless steel cylinder (Essex Cryogenics Inc., cylinder SX-3542). The composition of this cylinder for the four compounds of interest is reasonably stable (Table 2a) and 18 19 we plan to send this cylinder back to NOAA for subsequent analysis. The calibration gas in use at Bukit Atur from November 2008 to May 2009 was decanted into a 36 l stainless steel 20 21 electropolished flask from a dried NOAA Aculife cylinder (AAL-70424) in October 2008. 22 Comparisons of this decanted cylinder with the Aculife cylinder at either end of this period reveal a 23 significant change for CH₂Br₂ and CHBr₃ and a smaller one in C₂Cl₄ (Table 2b). The agreement 24 with the Aculife standard is much better in the later comparison which was performed in Sabah 25 under the same conditions as the atmospheric measurements. We do not know why the first comparison should have been poor (it was done in Cambridge just prior to shipping) though we 26 27 suspect it is related to temporary adsorption on the canister walls. Given the closer agreement between the baseline measurements of C₂Cl₄, CHBr₃ and CH₂Br₂^{*} at Tawau and Bukit Atur 28 29 (Section 5.1) using the values derived in the second comparison (last column of Table 2b), we 30 choose to use this for the final analysis, rather than the values from the October 2008 check (or an 31 average of the two checks). From May 2009 the NOAA Aculife cylinder (AAL-70424) used for the 32 decanting is used directly at Bukit Atur. This cylinder appears to be more stable than the decanted

1 cylinder though not as stable as the SX-3542 cylinder in use at Tawau (Table 2c). For the analysis 2 of our Bukit Atur data from May 2009 to January 2010 we use the values from the check against the University of East Anglia (UEA) made in April 2009 using their NOAA SX cylinder, as this is 3 4 closest in time to the period of the measurements. We also have more confidence in the comparison 5 made at UEA (April 2009) under stable laboratory conditions using their GC-MS than in an incomplete comparison made at Bukit Atur (November 2012) over 3 years later. During June and 6 7 July 2008 our measurements at Bukit Atur, analysed using the same NOAA (SX-3542) standard 8 (later used at Tawau), compared well with GC-MS measurements made by UEA (Gostlow et al., 9 2010; Pyle et al., 2011).

In addition, we took part in an inter-comparison of measurements by several laboratories using a round robin NOAA calibration standard (Jones et al., 2011). This showed our laboratory calibration scale to be within the instrumental error (10-20%, 2 sigma) of this new NOAA standard. In July 2008 we ran our two instruments side-by-side at Bukit Atur, measuring C_2Cl_4 , CHBr₃ and CH₂Br₂^{*} in ambient air, using the same calibration gas on both GC-ECDs. For this 4 day period we found that the observations of C_2Cl_4 , CHBr₃ and CH₂Br₂^{*} from the two instruments were within 6 %, 1 % and 15 % of each other respectively.

17 3 Field deployment

18 We have been making halocarbon observations in Sabah (Malaysian Borneo) since April 2008. 19 Initially, the instruments were deployed in the Danum Valley as part of the NERC funded OP3 20 project (Oxidant and Particle Photochemical Processes above a southeast Asian tropical rainforest), Hewitt et al. (2010). Two uDirac GCs were used in OP3: one was based at the Global Atmospheric 21 22 Watch (GAW) station at Bukit Atur (Gostlow et al., 2010) and the other made 'mobile' measurements at several rainforest locations in Danum Valley and during a 4 day visit to the coast 23 at Kunak (Pyle et al., 2011). At the end of the OP3 campaign the uDirac GC at the Bukit Atur 24 25 GAW station was left running and the second 'mobile' instrument was moved to a coastal site near Tawau where it remained for the period of this study. The instrument at Bukit Atur is visited at ~ 2 26 27 weeks intervals (and on request) by staff from the Malaysian Meterological Department (MMD) who manage the site. The Tawau instrument was visited at ~1 month intervals and on request by 28 29 one of our collaborators from Global Satria Sdn. Bhd. (an aquaculture company). Neither instrument was connected to the internet so these visits were essential for data recovery and 30 31 monitoring of instrument performance. Periodic service visits (every 9-12 months) were made to both sites by Cambridge University staff. As a result of this low intensity monitoring, the 32

performance of the instruments varies over time depending on the condition of the instrument and
 its environment. Each site is now described in the following sub-sections.

3 **3.1** Danum Valley rainforest site: the Bukit Atur GAW station

The Danum Valley conservation area covers 438 km² and is one of the largest, most important and 4 best-protected expanses of pristine lowland dipterocarp rainforest remaining in southeast Asia. The 5 6 GAW station located on Bukit Atur (4.980°N, 117.844°E, elevation 426 m) has been in operation since 2004 and is reasonably remote being ~50 linear km away from the nearest town of Lahad 7 8 Datu (Figure 1). The station is equipped with a range of monitoring instruments and an automatic 9 weather station located on a 5 metre roof-top platform. The 100 metre tower, which adjoins the 10 laboratory, has air-intakes and platforms for sampling equipment at various levels and is the tallest 11 instrumented structure on Borneo island. The on-going measurement programme at the Bukit Atur 12 GAW station includes surface ozone, greenhouse gases (flask sampling), carbon dioxide (using a Loflo Mk2 analyser), precipitation chemistry, aerosol (load, chemical and physical properties), 13 14 reactive gases (filterpack sampling), persistent organic pollutants, solar radiation and meteorological parameters. The site is operated by the MMD and staff visit weekly from their 15 nearest office at Tawau airport. The station itself is located on a hill-top ~200 m above the valley 16 17 floor and the inlet for the µDirac instrument is attached 12 m up the tower, several metres above the 18 roof of the site building. The instrument is placed in an air-conditioned room in the main GAW 19 building. The topography in the area consists of a series of hills ~ 400 m above sea level with valley 20 floors down to ~200 m above sea level. One of the largest mountains in the area is Mount Danum at 21 1,093 m above sea level. The rainforest extends to all areas including mountain tops. Primary 22 unlogged rainforest in the conservation area is surrounded by areas of secondary rainforest (which 23 have been logged in recent years). In terms of site meteorology, a clear diurnal cycle in rainfall is 24 observed at the nearby Danum Valley Field Centre (DVFC) with an afternoon peak around 15:00 25 hrs local time resulting from diurnal development of convective cells following the midday peak in temperature (Hewitt at al., 2010). Boundary layer stability also varies diurnally from highly stable 26 27 at night (often leading to dense fog at night) to highly unstable during the day due to the onset of strong convection coupled with low wind speeds (Pearson et al., 2010). 28

29 **3.2** Tawau coastal site: Kampung Batu Payong

The Tawau coastal site is located in the peaceful village of Kampung Batu Payong (4.223°N,
117.997°E, elevation 15 m) and is ~10 km from Tawau centre and ~85 km south of the Bukit Atur

1 rainforest station (Figure 1). Between July 2008 and October 2011 the µDirac instrument here was 2 located in an air-conditioned room at a fish hatchery site maintained by the Global Satria company. There are no seaweed beds in the immediate vicinity of the site. The activity at the site does not 3 4 involve the use of any seaweeds (which are known to emit bromocarbons) and we do not find any 5 evidence of emissions from the hatchery activities. The air inlet is located ~5 m above ground level 6 at the site (about roof-top height) 40 m away from the shoreline. There is an unobstructed view to 7 the sea extending to more than 180° of horizon to the south and the site is situated at the foot of a 8 small group of hills extending to the north, reaching an elevation of 350 m. A low cost weather 9 station at the site gives an indication of wind speed and direction, temperature and humidity. This 10 reveals a typical diurnal variation in wind due to land/sea breezes expected for coastal sites, though 11 this does not obviously influence the observations reported here.

12 **4 Models**

13 Two modeling tools are used in this paper to assist in the interpretation of the observations. They 14 are now described in turn in sub-Sections 4.1 and 4.2

15 **4.1 NAME**

16 The Numerical Atmospheric dispersion Modelling Environment (NAME) is a Lagrangian particle 17 dispersion model developed by the UK Met Office (e.g. Jones et al., 2007), which has been 18 extensively used for analysis of long-term halocarbon data sets (e.g. Manning et al., 2003; 19 Simmonds et al., 2006; Derwent et al., 2007). Abstract particles are moved through the model atmosphere by a combination of 0.5625° longitude by 0.375° latitude mean wind fields calculated 20 by the UK Met Office Unified Model (Davies et al., 2005), and a random walk turbulence scheme. 21 22 NAME can be run backwards in time, to see where the air measured at a particular site may have 23 originated, and forwards to see where air from a particular emission source might go. Each of these 24 capabilities have previously been used to investigate relationships between sources, transport and measurements of pollutants (e.g. Redington and Derwent, 2002; Witham and Manning, 2007). 25

Here, NAME is run backwards in time to reveal the origin of air arriving at Bukit Atur and Tawau. For each 3 hour period thousands of inert particles (carrying an arbitrary mass) are released and travel backwards in time for 12 days. At each 15 min time-step the location of each particle is recorded. Then, at the end of the run this information is used to create an 'air history map'. Here, we only record near surface particles (lowest 100 m of the model) which indicate where air may have been exposed to surface emissions. Monthly mean air histories are then calculated. The air history
 maps made for Tawau are almost identical to those from Bukit Atur so are not shown here.

3 **4.2 p-TOMCAT**

4 The global chemistry transport model (CTM), p-TOMCAT, is used to analyse the causes of seasonal bromoform variability at Bukit Atur. The basic formulation of p-TOMCAT is described in 5 6 Cook et al. (2007) and Hamilton et al. (2008). Tracer transport is based on 6-hourly meteorological fields, including winds and temperatures, derived from European Centre for Medium-Range 7 8 Weather Forecasts operational analysis for the years 2008, 2009 and 2010. Here we use a high 9 resolution (0.5° x 0.5°) version with bromoform tracers 'coloured' according to the region of 10 emission and simple OH oxidation chemistry, described previously in Pyle et al. (2011) and 11 Warwick et al. (2006). The degradation of bromoform is determined using 3-D fields of pre-12 calculated hourly OH values taken from a previous p-TOMCAT integration and photolysis frequencies from an integration of the Cambridge 2-D model using cross section data summarised 13 14 by Sander et al. (2003). For further details see Pyle et al. (2011). Here we compare time series of 15 bromoform from the model output with the measurements.

16 **5** Results and Discussion

In this section we present the measurements of C_2Cl_4 , CHBr₃ and CH₂Br₂^{*} from both sites and CH₃I from Tawau only. We first show that C_2Cl_4 can be used as a good tracer of air mass, differentiating clearly between air masses with anthropogenic and unpolluted origins. This provides a valuable background for the analysis of the three, primarily biogenic, short-lived compounds, CHBr₃, CH₂Br₂^{*} and CH₃I. In this section we also assess the total organic Br coming from CHBr₃ and CH₂Br₂^{*} combined. We discuss the implications of these observations, in particular for the regional emissions of CHBr₃ and CH₂Br₂^{*}.

24 **5.1 Perchloroethene**

Perchloroethene (C_2Cl_4) is an excellent anthropogenic tracer with few (if any) known natural sources. Its main anthropogenic sources are the textile industry, dry-cleaning applications and in vapour degreasing of metals (Montzka and Reimann, 2011). It currently has a northern hemisphere background concentration of 2-5 ppt with a pronounced seasonal variation, likely due to variation in the concentration of its major sink, the OH radical. In the southern hemisphere the background concentration is lower (0.5-1 ppt), though still with a pronounced seasonal variation (see http://agage.eas.gatech.edu/data.htm; Simmonds et al., 2006). C₂Cl₄ has an estimated lifetime of
~90 days (see Table 1-4 in Montzka and Reimann, 2011), making it a good tracer for studying long
range transport.

At Bukit Atur and Tawau the C_2Cl_4 dry air mole fractions ranged typically from 0.3-0.8 ppt in May-August 2009 to 1.0-3.7 ppt from November 2008 to March 2009 (Figure 3a). 'Above baseline' spikes are observed occasionally at Tawau, but for most of the time the data at Bukit Atur and Tawau track each other on monthly and daily timescales. As the two sites are 85 km apart and as both C_2Cl_4 time series track each other closely, it seems there is little local contribution from Sabah to the background mixing ratio.

10 For the 4 month period between December 2008 and March 2009 there is a systematic difference in 11 the C₂Cl₄ observations at the two sites, with the observations at Bukit Atur on average 48 % higher 12 than those at Tawau. It is surprising to find this large systematic difference given that the observations of C₂Cl₄ were within 6 % of each other when the GCs ran side-by-side at the same site 13 14 and with the same calibration gas at Bukit Atur in July 2008 (Section 2 last paragraph). The differences are probably due to a relative drift in the composition of the calibration gases used at 15 16 each site. Any drift in composition is likely to be larger for the decanted calibration gas which was 17 used at Bukit Atur between November 2008 and May 2009 and therefore we have more confidence 18 in the time series from Tawau (see Section 2.2). The C_2Cl_4 observations reported at Bukit Atur 19 between December and March are likely to be an upper estimate of the actual mixing ratios at that 20 time.

21 Although there is a greater frequency of high (local pollution) spikes at Tawau these do not impact 22 significantly on the underlying background which is similar to that at Bukit Atur. The seasonal 23 trend at both sites is clear from box and whisker plots (Figure 4a & d) and the tabulated monthly 24 mean, median and modal values (Table 3a) which show low levels and low variability in the 25 northern hemisphere summer months in contrast with higher levels in winter. To further explore these seasonal differences and site-to-site similarities, for selected months we use probability 26 27 distributions from the C_2Cl_4 observations, using a size interval of 0.1 ppt (Figure 5a & e). In 28 January/February 2009 there is a broad distribution of mixing ratios at both sites ranging from 0.85 29 to 3.75 ppt, tailing towards higher mixing ratios. The Tawau coastal median value in 30 January/February 2009 (seen in Figure 5a) is lower (1.30 ppt) than inland at Bukit Atur (1.91 ppt), 31 though recall that the C₂Cl₄ observations at Bukit Atur are likely an upper estimate. In June/July 32 2009 the distribution at each site has a much narrower range (0.25 to 1.65 ppt) and the median values are similar, though the coastal median (0.45 ppt) is still lower than inland (0.63 ppt). In
January/February 2009 the Bukit Atur instrument ran on the decanted calibration air from the
NOAA Aculife cylinder AAL-70424, whereas in June/July 2009 the calibration came directly from
the Aculife cylinder.

5 The seasonal changes in C₂Cl₄ are linked to changes in the prevailing monsoon flow, and the inter-6 hemispheric gradient typically observed for gases with predominantly northern hemisphere 7 anthropogenic sources. Air history maps generated using NAME are used here to illustrate the 8 seasonal changes in transport. From the months of November through to February the air mass 9 history is influenced strongly by air coming from the north and east (Figure 6a, January 2009). The 10 large urban centres in and around China in this region are expected to contribute to the background 11 level of C₂Cl₄ arriving at Sabah. From March to May 2009 the air mass origin tends to be 12 influenced more by oceanic air from the West Pacific though with still some trajectories passing the 13 vicinity of China (Figure 6b, March 2009). During June, July, August and September 2009 the air 14 mass history indicates that Sabah is influenced strongly from the region to the southeast, 15 characterised by southern hemisphere air originating from the Indonesian islands and Northern Australia (Figure 6c, June 2009). As there are no major sources of C₂Cl₄ across this region and the 16 17 air masses are not influenced by emission from for example Jakarta, we find (as expected) background mixing ratios to be low. October sees a transition between background air influenced 18 19 by the southern hemisphere and northern hemisphere air influenced by China and other large urban 20 centres. It is thus an example of a transition period between the two monsoon regimes (Figure 6d, 21 October 2009).

22 5.2 Bromoform

23 Bromoform (CHBr₃) is an excellent tracer of marine air and is an important source of organic 24 bromine to the atmosphere (Carpenter and Liss, 2000). An open ocean, phytoplankton source has 25 been identified (Tokarczyk and Moore, 1994), and macroalgae found in coastal areas are also well 26 known as major sources of bromoform (Moore and Tokarczyk, 1993). Emission fluxes in coastal regions are thought to be higher than in the open ocean (Quack and Wallace, 2003; Yokouchi et al., 27 28 2005; Butler et al., 2007) though the relative contribution from coasts and from the open ocean to 29 the total global emission remains uncertain. Recent estimates of the global emissions of CHBr₃ are 30 in the range 430-1400 Gg Br/yr (Liang et al., 2010; Pyle et al., 2011; Ordonez et al., 2012 and Table 31 1-8 in Montzka and Reimann, 2011) though much lower estimates of 120 and 200 Gg Br/yr are 32 given by Ziska et al. (2013). Although bromoform emissions are dominated by natural sources there are minor anthropogenic contributions from by-products of drinking water chlorination in the presence of bromide ions, salt water swimming pools and power plant cooling water (Worton et al. 2006). In the atmosphere, CHBr₃ has a lifetime of 24 days (from Table 1-4 in Montzka and Reimann, 2011) which is shorter in the tropics (15-21 days), as discussed in Warwick et al. (2006). The significant loss mechanisms include reaction by photolysis and reaction with OH (Hossaini et al., 2010; Montzka and Reimann, 2011).

7 The background CHBr₃ concentration at both Bukit Atur and Tawau is broadly in the range 0.5 to 3 8 ppt. In contrast to the time series of C_2Cl_4 , there is no unambiguous seasonal variation in the 9 background CHBr₃ concentration (Figure 3b). During the periods when we report CHBr₃ mixing 10 ratios from both sites, the background concentrations are sometimes the same within experimental 11 error (e.g. Figure 5b for January/February 2009), and in some months the coastal background is higher than inland (e.g. Figure 5f for June/July 2009). These probability distributions also indicate 12 13 that we do not observe strong seasonality in the background mixing ratio at Tawau. In contrast, 14 there is a suggestion of a seasonal cycle, with a northern hemisphere winter peak, in the data 15 collected inland at Bukit Atur (e.g. Figure 4b). To emphasise this point, the median mixing ratios for the two sites were similar in January/February 2009 (1.67 ppt at Tawau and 2.04 ppt at Bukit 16 17 Atur), whereas for June/July 2009 the median value for Tawau (1.68 ppt) was higher than at Bukit Atur (0.95 ppt). Clearly, measurements over many years are needed to ascertain whether the 18 19 somewhat different behaviour we have observed at the two sites in Borneo is a persistent feature. 20 See Table 3b for the statistical summaries of all the relevant periods.

A major difference between the two sites is that there are frequent periods (spikes) at Tawau when very high levels (> 10 ppt) of bromoform are observed, whereas at Bukit Atur such episodes are rare, and the highest concentration is just 5.5 ppt. This difference in variability is shown clearly in the box and whisker plots for Bukit Atur and Tawau (Figure 4b & e) where CHBr₃ observations > 10 ppt at Tawau are seen in 12 of the 15 reported months.

Various factors determine the concentrations we observe at these two sites. The sources of CHBr₃ are certainly heterogeneous in space, and so the mere fact that the boundary layer is influenced by different regions in different seasons (Figure 6) leads to a varying influence of recent emissions and hence varying observed concentrations. This heterogeneity occurs due to both natural spatial variations in the populations of algae in coastal and open ocean areas, and due to the presence of commercial aquaculture in the region, whose influence on bromoform emissions needs to be assessed. For example, northeast Sabah and the archipelagos between Sabah and the Philippines are

1 major seaweed farming areas whose output has grown quickly over the last 20 years (Phang, 2006; 2 Perez, 2011). At present it appears that entirely natural emissions of bromoform in the region are significantly larger than those linked to aquaculture (Leedham et al., 2013) but continued rapid 3 growth in the cultivation of many species with commercial potential (Goh and Lee, 2010; Tan et al., 4 5 2013) could change this situation (Phang et al., 2010). Further, the strength of individual sources is likely to vary over time. In the case of seaweeds, it seems likely that their halocarbon emissions will 6 7 depend on environmental factors such as temperature, rainfall, wind, sunlight and if applicable, the 8 phase of the cropping cycle (Seh-Lin Keng et al., 2013, Leedham et al., 2013 and references 9 therein). Given this complexity, without extensive biological field surveys it is difficult to explain much of the variability in detail. 10

However, we are able to examine how well the CTM p-TOMCAT is able to simulate the long-term 11 12 statistical properties of observed bromoform mixing ratios. The model configuration is described in Section 4.2, and the bromoform emissions are those developed by Pyle et al. (2011) as an update to 13 Warwick et al. (2006). These emissions provide the best match with our earlier observations 14 collected at Bukit Atur during OP3 (Pyle et al., 2011). The observed time series for CHBr₃ is 15 consistent with the p-TOMCAT model output using the revised emission estimate as shown for 16 17 Tawau in Figure 7a. The model is unable to capture the very high values seen in the observations, 18 which is to be expected given the model resolution and the uniform distribution of emissions around 19 the coastlines in the model. The model does a reasonable job of reproducing the background 20 observed concentration throughout 2009 at Tawau. The bromoform observations and model output 21 for the four months from December 2008 to March 2009 are compared as a probability distribution in Figure 7b. The observations show only a slightly broader distribution (inter-quartile range 0.84 22 23 ppt) than the model output (inter-quartile range 0.58 ppt) and the median value for the observations 24 of 1.73 ppt is just slightly higher than that for the model output (1.65 ppt).

25 **5.3 Dibromomethane**

It is thought that the marine sources of dibromomethane (CH_2Br_2) are similar to those of CHBr₃, as shown by the strong correlations observed between their atmospheric concentrations (Yokouchi et al., 2005; Carpenter et al., 2009). Emission fluxes are probably higher in coastal regions than over the open ocean (Ziska et al., 2013), but the relative effect on coastal atmospheric measurements is less marked than for CHBr₃ due to a longer lifetime for CH₂Br₂ of ~120 days (Table 1-4 in Montzka and Reimann, 2011). The CH₂Br₂ lifetime is expected to be slightly lower than 120 days in the tropics due to the higher abundance of the OH radical. Open ocean sources are also more important 1 globally than coastal sources. Recent estimates of the global CH_2Br_2 emissions are 50-70 Gg Br/yr

2 (Liang et al., 2010; Ordonez et al., 2012).

The mixing ratios of $CH_2Br_2^*$ measured at Bukit Atur and Tawau are shown in Figure 3c. The 3 observed baseline values of between 1 and 2 ppt for $CH_2Br_2^*$ at the two sites are similar, as 4 reflected in the median values in Table 3. Previous work suggests that 10-30% of the $CH_2Br_2^*$ 5 6 signal may in fact be CH₂BrCl₂. During June and July 2009, Nadzir et al. (2014) collected air 7 samples on a ship cruise around the coast of the Sulu-Sulawesi seas close to Tawau and the average 8 CH₂Br₂ mixing ratio was 1.04 ppt compared to 0.3 ppt for CHBrCl₂. The authors also found that the 9 two compounds tracked each other closely through the entire cruise. In another study, Sala et al. 10 (2014) reported average boundary layer mixing ratios of CH₂Br₂ and CHBrCl₂ of 1.19 and 0.34 ppt 11 respectively, from an aircraft campaign in the Western Pacific region during November/December 12 2011. Both CHBrCl₂ and CH₂Br₂ are known to have a common macroalgal source as found by Seh-13 Lin Keng et al. (2013) and Leedham et al. (2013), in studies of tropical seaweeds from the 14 Malaysian coast. These studies also found that production rates of CH₂Br₂ were generally (though 15 not always) much higher than for CHBrCl₂. In the atmosphere CHBrCl₂ has a lifetime of 78 days compared to 123 days for CH₂Br₂ (Montzka and Reimann, 2011). Thus we would not expect to see 16 any significant difference in the measured CHBrCl₂:CH₂Br₂ ratio between Tawau and Bukit Atur 17 due to differential processing, given the short air mass transit time between the sites of typically 1 18 day. The variability of $CH_2Br_2^*$ at Tawau is much less than for CHBr₃. What may be an important 19 20 factor here in variability differences between the two sites, is the extent to which boundary layer air 21 at Bukit Atur has been diluted with free tropospheric air with a lower ratio of CHBr₂Cl to CH₂Br₂, 22 given that the site is 426 m above sea level. Statistically, the observations at Bukit Atur and Tawau 23 are similar to each other as shown by the box and whisker plots, although the two months at Tawau with distinctly low median values (March/April 2009) are not reflected in the Bukit Atur 24 observations (Figure 4c & f). There is a definite hint of some seasonality in the $CH_2Br_2^*$ 25 26 measurements (Figure 5c & g), though multi-year data are required to determine whether there is a 27 seasonal cycle. Median values in January/February 2009 are almost the same at both sites (1.20 and 28 1.28 ppt at Bukit Atur and Tawau respectively), though in June/July 2009 the median value at 29 Tawau (1.33 ppt) is distinctly higher than at Bukit Atur (1.00 ppt).

We find that in all reported months at Bukit Atur there is always a positive correlation between the observations of $CH_2Br_2^*$ and $CHBr_3$, suggesting a common source (Table 3c). Nadzir et al. (2014) also found a strong positive correlation between $CHBr_3$ and CH_2Br_2 in whole air flask samples 1 taken during a 2009 cruise in the South China and Sulu-Sulawesi Seas (close to the Tawau site) and Strait of Malacca. The strength of the correlation at Bukit Atur and Tawau is highly variable 2 between months. The strongest correlation at Bukit Atur is found in November 2008 with an R^2 3 value of 0.86. At Tawau the correlation between $CH_2Br_2^*$ and $CHBr_3$ is generally much weaker than 4 5 at Bukit Atur, though still positive, with the exception of June 2009. Only March and September 2009 show strong positive correlations at Tawau. As we are only able to report a $CH_2Br_2^*$ 6 7 measurement in about 1 in every 10 samples at Tawau there are less data available for the correlation of CH₂Br₂^{*} and CHBr₃. 8

9 **5.4 Methyl iodide**

10 Methyl iodide (CH₃I) is ubiquitous in the marine boundary layer, with typically reported 11 concentrations of 0.1–5 ppt (e.g., Lovelock et al., 1973; Rasmussen et al., 1982; Yokouchi et al., 2001, 2008, 2011, 2012). There are several known oceanic sources of CH₃I including 12 phytoplankton (Moore and Tokarczyrk, 1993; Manley and dela Cuesta, 1997), macroalgae (Manley 13 14 and Dastoor, 1988; Schall et al., 1994; Giese et al. 1999), methylation of iodine by bacteria (Amachi et al., 2001), picoplankton (Smythe-Wright et al., 2006; Brownell et al., 2010), 15 photochemical reactions (Moore and Zafiriou, 1994) and as a by-product of the ozonolysis of 16 dissolved organic matter (Martino et al., 2009). Methyl iodide has an atmospheric lifetime of 7 days 17 18 (Table 1-4 in Montzka and Reimann, 2011) and has the shortest lifetime of the compounds reported 19 here.

20 The most extensive, continuous measurements of CH₃I reported to date in the tropics or sub-tropics 21 were made hourly at Hateruma Island (24.1°N, 123.8°E) from August 2008 to January 2010 22 (Yokouchi et al., 2011). At Hateruma Island, no significant seasonal variation was observed, in contrast to Cape Ochiishi (43.2°N, 145.5°E) where a clear maximum was observed in late 23 summer/autumn coincident with increased variability. No diurnal variation in CH₃I was observed at 24 either location. These high resolution measurements are complemented by the longer, semi-monthly 25 26 measurements made from the late 1990s to 2011 at 5 remote sites, Alert (82.5°N, 62.5°W), Cape 27 Ochiishi, Happo Ridge (36.7°N, 137.8°E), Hateruma Island and Cape Grim (40.4°S, 144.6°E) as well as regular ship transects of the Pacific (Yokouchi et al., 2012). Summer maxima were seen at 28 29 all the four sites apart from Alert where a wintertime maximum (attributed to transport) was seen. 30 The discrepancy between the continuous and semi-monthly measurements at Hateruma Island 31 presumably results from low frequency sampling of the highly variable mixing ratio in the late summer. No significant seasonal cycle is seen in the baseline of the higher resolution CH_3I time series. Observations of CH_3I in the tropical boundary layer have otherwise been limited to measurements of whole air samples made at specific sites or on research cruises (see summaries in Table 1 in Yokouchi et al. (2008) and Table 2 in Saiz-Lopez et al. (2012)).

5 Our CH_3I record for Borneo (Tawau only) is therefore a unique source of information in the tropics. 6 The measurements for 2009 are shown in Figures 3d, 4g and 5d & h. The CH_3I peak is small and 7 hard to separate (using our current column configuration) from the larger peaks that elute at a 8 similar time. The µDirac at Bukit Atur did not achieve satisfactory chromatographic separation for 9 reliable CH_3I quantification and these data are not reported here.

10 The CH₃I mixing ratios at Tawau (~8400 in total) vary from below 0.1 up to 11 ppt, with the large majority of points towards the lower end of the range. July sees the lowest monthly mean of 0.34 11 12 ppt and the highest monthly mean of 2.50 ppt is in September. The highest values and the greatest variability are observed in September and October. Back-trajectories indicate that air containing 13 14 high amounts of CH₃I had previously passed over large biomass burning events in southern Borneo 15 (as observed by MODIS). Emissions of CH₃I in biomass burning have been reported previously 16 (Andreae et al., 1996; Mead et al., 2008c). However while biomass burning emissions may be 17 significant locally at some times of year, it is not the case globally (Bell et al., 2002; Butler et al., 18 2007). Interestingly some of the lowest observed values (< 0.1 ppt) of CH₃I occur when the air has 19 previously been in the terrestrial boundary layer, indicating that the tropical rainforests were not 20 normally a source of CH₃I. The seasonal cycle in the baseline (i.e. if the high values seen in the late 21 summer/early autumn are excluded) at Tawau is weak, similar to that seen at Hateruma Island. A 22 longer record is needed to identify whether one exists.

23 **5.5 Total bromine**

Here we present and discuss our observations of bromine defined by $[CHBr_3 \times 3] + [CH_2Br_2^* \times 2]$ 24 25 as an approximate measure of total organic bromine from the two sites. For a two week period of 26 the OP3 campaign (5-12 July 2008) we ran our Bukit Atur instrument alongside a GC-MS system 27 from UEA which measured CHBr₃, CH₂Br₂, CHBr₂Cl, CHBrCl₂ and CH₂BrCl (the five main naturally occurring bromine source gases). The GC-MS observations showed that CHBr₃ and 28 29 CH₂Br₂ accounted for 93.5 % of the total Br coming from these five bromine source gases (Pyle et al., 2011a). Total bromine as $[CHBr_3 \times 3] + [CH_2Br_2^* \times 2]$ from our measurements at Tawau over 30 31 the period November 2008 to January 2010 is shown as a time series in Figure 8a. The black line 1 through the data is the running average total bromine (40 point, 50 pass) to highlight average 2 variability. Bromine from these two compounds (accounting for >90 % of the total organic Br) averages 9.7 ppt over the period of observations and ranges from 3.7 ppt (June 2009) to 189.5 ppt 3 4 (December 2008). The probability distribution function for total observed bromine as [CHBr₃] + [CH₂Br₂^{*}] at Bukit Atur is compared with that for Tawau in Figure 8b. The distributions are similar 5 (median values of 6.79 and 7.97 ppt at Bukit Atur and Tawau respectively) and Bukit Atur has a 6 7 narrower inter-quartile range than Tawau (2.06 ppt compared to 2.80 ppt). The extended tail of very 8 high total bromine values (>20 ppt) is evident at Tawau and completely absent from the Bukit Atur 9 observations.

10 As discussed earlier, we are unable to report separate observations of CHBrCl₂ and CH₂Br₂ and are 11 obliged to report combined observations $(CH_2Br_2^*)$. Separate observational studies have indicated 12 that CHBrCl₂ mixing ratios are typically 10 to 30 % of those for CH₂Br₂ (Pyle et al., 2011a; Nadzir 13 et al., 2014; Sala et al., 2014). The total bromine observations at each site are relatively insensitive 14 to this range of possible variation. If we apply this range to the $CH_2Br_2^*$ observations then the 15 impact on the median bromine observations is to reduce total Br at Bukit Atur by between 0.1 and 16 0.3 ppt and at Tawau by between 0.1 and 0.4 ppt.

The fact that our observations of bromine from $[CHBr_3 \times 3] + [CH_2Br_2^* \times 2]$ are so similar at both 17 sites (with the exception of the very high coastal values) shows that an inland site can be used 18 19 effectively to make regional observations suited to model comparison. In fact, as the model we used 20 here was unable to capture the large variability in the observations, presumably due to insufficient 21 information on the spatial and temporal inhomogeneity of the emissions seen at the coast and to 22 model resolution, we suggest that, for evaluating models, an inland site is preferable to a coastal site. We also calculate total modelled bromine coming from $[CHBr_3 \times 3] + [CH_2Br_2 \times 2]$ in p-23 TOMCAT at Tawau and find for the period of our observations a total bromine median value of 24 25 6.56 ppt, somewhat lower than the median value of 7.97 ppt from the measurements.

Our data provide a longer-term context for other reports from short campaign-type observations. There are month-to-month variations in total Br in our data that would be missed by shorter deployments. Overall, our mean value of 9.7 ppt from $[CHBr_3] + [CH_2Br_2^*]$ seems quite consistent with other work in the region. There have been somewhat conflicting reports of the fraction of Br in CHBr₃ and CH₂Br₂ compared to the minor very short-lived substances (VSLS). Montzka and Reimann (2011) suggest a boundary layer median value of 8.4 ppt Br from VSLS, with 83 % from CHBr₃ (1.6 ppt × 3) and CH₂Br₂ (1.1 ppt × 2). Based on data obtained in July 2008 from the OP3 campaign, Pyle et al. (2011a) report 7 ppt (5-10 ppt) Br from VSLS, with over 90 % of this present
as CHBr₃, CH₂Br₂ or CHBrCl₂. Nadzir et al. (2014) report a mean of 8.9 ppt Br from all VSLS
(range 5.2 to 21.4 ppt in individual samples), with ~90% from CHBr₃ and CH₂Br₂ during the Prime
Expedition Scientific Cruise during June and July 2009.

5 6 Summary

6 Until recently, few measurements of short-lived halogenated compounds had been made in 7 southeast Asia. These compounds have significant atmospheric importance, and previous studies 8 have speculated that southeast Asia represents an unusually strong source of the naturally occurring 9 species (e.g. CHBr₃). Here we present more than one year of observations (November 2008 to 10 January 2010) of key short-lived compounds from a coastal (Tawau) and an inland (Bukit Atur) site 11 in this region, which together provide valuable data to the atmospheric science community. The 12 data have been placed on the HalOcAt database for community access (https://halocat.geomar.de/). 13 The data allow seasonal cycles, emissions and the role of local and long range transport to be 14 studied.

By combining the observations with air mass history information generated by NAME we explain the strong seasonal cycle seen in the anthropogenic tracer (C_2Cl_4). From December to February, northerly air masses lead to higher and more variable mixing ratios, ranging from 0.8 to 3.7 ppt. In contrast, from May to August the observed air masses originate from regions to the south (typical mixing ratio 0.5 ppt). The overall median values for C_2Cl_4 are 1.33 ppt and 0.99 ppt at Bukit Atur and Tawau respectively.

The naturally occurring brominated tracers CHBr₃ and CH₂Br₂^{*} do not show unambiguous seasonal 21 variation, though there is a slight hint of a seasonal cycle in CH₂Br₂^{*}. Both compounds nevertheless 22 23 show strong variability over shorter timescales. At the coast, the CHBr₃ observations occasionally 24 peak at over 100 ppt compared with the overall median level of 1.7 ppt. At Bukit Atur, the overall 25 inland median level for CHBr₃ is 1.6 ppt and the occasional peaks are much lower than at the coast with maximum observations of just over 5 ppt. Coastal $CH_2Br_2^*$ mixing ratios occasionally rise to 26 27 over 7 times the median level of 1.4 ppt, whereas inland the maximum level observed is a little over 28 2 ppt. The strong coastal variability of both these compounds is likely related to the strongly 29 heterogeneous nature of their marine sources, which are believed to be a combination of coastal 30 macroalgae and both coastal and oceanic microalgae. Accordingly, the measurements from coastal 31 Tawau are more variable than at Bukit Atur, being made closer to these sources.

The observations of CHBr₃ and CH₂Br₂^{*} also provide a valuable longer term estimate of the 1 2 contribution to total Br in the boundary layer from the main naturally occurring short-lived 3 brominated compounds. At the coast we observe an overall median value of Br from $[CHBr_3 \times 3] +$ $[CH_2Br_2^* \times 2]$ of 8.0 ppt and inland the overall median value is 6.8 ppt. These values are reasonably 4 5 consistent with other recent estimates of this quantity based on much shorter measurement periods. 6 Our data provide evidence that southeast Asia may not in fact be a region of enhanced brominated 7 VSLS emissions into the boundary layer in relation to other tropical locations where similar 8 measurements have been made.

9 Our coastal observations of CH_3I show high variability as expected for a short lived tracer with 10 values from less than 0.1 ppt to over 11 ppt. The overall median value is 0.6 ppt. The highest values 11 are observed in September and October 2009 and are attributed to the influence of biomass burning 12 in southern Borneo that year. In other times of the year, air that passed over land showed some of 13 the lowest values observed, indicating that the tropical rain forests in Borneo are not a source of 14 CH_3I .

15 To date, modelling work has focussed on understanding our CHBr₃ data. The CHBr₃ observations 16 presented here were recently used by Ashfold et al. (2014), along with NAME trajectories and an 17 inversion method, to estimate regional scale CHBr₃ emissions. When extrapolated across the global 18 tropics, the magnitude of their estimated emissions was somewhat lower than most other recent 19 estimates, which have tended to assume strong emissions in southeast Asia. This is consistent with the finding here, that observed CHBr₃ mixing ratios in Borneo (and indeed CH₂Br₂^{*} mixing ratios 20 and total Br estimates) are in fact similar to those measured in other parts of the world. Our 21 22 measurements also provide a means of testing the ability of global models like p-TOMCAT to 23 reproduce boundary layer time series of short-lived compounds in tropical regions. Although p-24 TOMCAT was unable to reproduce the same magnitude of variability seen in the Borneo CHBr₃ 25 observations, it was able to capture background concentrations using the bromoform emission 26 distribution described in Pyle et al., (2011). This emission distribution is an updated version of the 27 distribution described in Warwick et al., (2006), containing substantial reductions in southeast 28 Asian emissions relative to the original scenario.

We continue to make halocarbon measurements in Sabah and have upgraded to the newer version of μ Dirac with a 20 m long column and better resolution of species including CH₂Br₂. In addition, we have recently begun measuring the same group of compounds using upgraded versions of the instruments used here at three other sites in the southeast Asia region: at Bachok on the Malaysian peninsula, Kelantan state; at Gunn Point near Darwin, Australia; a coastal site near Taipei, Taiwan.
 These sites significantly extend the geographic coverage of our measurements.

3

4 Acknowledgements

5 This work was supported by a NERC consortium grant to the OP3 team, by NCAS, by the European Commission through the SCOUT-O3 project (505390-GOCE-CF2004) and by NERC 6 7 West Pacific grant number NE/F020341/1 and NERC CAST grant number NE/J006246/1. Louise 8 O'Brien and Matt Ashfold thank NERC for research studentships. Andrew Robinson acknowledges 9 NERC for their support through small grant project NE/D008085/1. Neil Harris is supported by a 10 NERC Advanced Research Fellowship. We thank The Sabah Foundation, Danum Valley Field 11 Centre and the Royal Society (Glen Reynolds) for field site support. The research leading to these 12 results has received funding from the European Union's Seventh Framework Programme FP7/2007-13 2013 under grant agreement n°226224 - SHIVA. We thank David Oram and Stephen Humphrey at 14 UEA for their assistance in checking the calibration of our Aculife cylinder in May 2009.

15

16 This is paper number 626 of the Royal Society's South East Asian Rainforest Research Programme.

17

22

18 **References**

- Amachi, S., Kamagata, Y., Kanagawa, T. and Muramatsu, Y.: Bacteria mediate methylation of
 iodine in marine and terrestrial environments, Appl. Environ. Microbiol., 67, 2718-2722, 2001.
- 21 Andreae, M.O., Atlas, E., Harris, G.W., Helas, G., de Kock, A., Koppmann, R., Maenhaut, W.,
- halide emissions from savanna fires in southern Africa, J. Geophys. Res., 101, 23,603-23,613, 1996.

Manø, S.M., Pollock, W.H., Rudolph, J., Scharffe, D., Schebeske, G. and Welling, M.: Methyl

- Aschmann, J., Sinnhuber, B.-M., Atlas, E. L., and Schauffler, S. M.: Modeling the transport of very
- 25 short-lived substances into the tropical upper troposphere and lower stratosphere, Atmos. Chem.
- 26 Phys., 9, 9237-9247, doi:10.5194/acp-9-9237-2009, 2009.
- 27 Ashfold, M. J., Harris, N. R. P., Manning, A. J., Robinson, A. D., Warwick, N. J., and Pyle, J. A.:
- Estimates of tropical bromoform emissions using an inversion method, Atmos. Chem. Phys., 14,
 979-994, 10.5194/acp-14-979-2014, 2014.

- 1 Brinckmann, S., Engel, A., Bönisch, H., Quack, B., and Atlas, E.: Short-lived brominated
- 2 hydrocarbons observations in the source regions and the tropical tropopause layer, Atmos. Chem.
- 3 Phys., 12, 1213-1228, doi:10.5194/acp-12-1213-2012, 2012.
- 4 Bell, N., Hsu, L., Jacob, D. J., Schultz, M. G., Blake, D. R., Butler, J. H., King, D. B., Lobert, J. M.,
- 5 and Maier-Reimer, E.: Methyl iodide: Atmospheric budget and use as a tracer of marine convection
- 6 in global models, J. Geophys. Res., 107, 4340, doi:10.1029/2001JD001151, 2002.
- Brownell, D. K., Moore, R. M., and Cullen, J. J.: Production of methyl halides by Prochlorococcus
 and Synechococcus, Global Biogeochem. Cy., 24, 1–7, doi:10.1029/2009GB003671, 2010.
- 9 Butler, J. H., King, D. B., Lobert, J. M., Montzka, S. A., Yvon-Lewis, S. A., Hall, B. D., Warwick,
- 10 N. J., Mondeel, D. J., Aydin, M., and Elkins, J. W.: Oceanic distributions and emissions of short-
- 11 lived halocarbons, Global Biogeochem. Cycles, 21, GB1023, doi: 10.1029/2006GB002732, 2007.
- 12 Carpenter, L. J. and Liss, P. S.: On temperate sources of bromoform and other reactive bromine
- 13 gases, J. Geophys. Res., 105, 20,539 20,547, doi:10.1029/2000JD900242, 2000.
- 14 Carpenter, L. J., Jones, C. E., Dunk, R. M., Hornsby, K. E., and Woeltjen, J.: Air-sea fluxes of 15 biogenic bromine from the tropical and North Atlantic Ocean, Atmos. Chem. Phys., 9, 1805-1816,
- 16 doi:10.5194/acp-9-1805-2009, 2009.
- 17 Cook, P. A., Savage, N. H., Turquety, S., Carver, G. D., O'Connor, F. M., Heckel, A., Stewart, D.,
- 18 Whalley, L. K., Parker, A. E., Schlager, H., Singh, H. B., Avery, M. A., Sachse, G. W., Brune, W.,
- 19 Richter, A., Burrows, J. P., Purvis, R., Lewis, A. C., Reeves, C. E., Monks, P. S., Levine, J. G., and
- Pyle, J. A.: Forest fire plumes over the North Atlantic: p-TOMCAT model simulations with aircraft
 and satellite measurements from the ITOP/ICARTT campaign, J. Geophys. Res., 112, D10S43,
- 22 doi:10.1029/2006JD007563, 2007.
- Davies, T., Cullen, M. J. P., Malcolm, A. J., Mawson, M. H., Staniforth, A., White, A. A., and
 Wood, N.: A new dynamical core for the Met Office's global and regional modelling of the
 atmosphere, Q. J. Roy. Meteorol. Soc., 131, 1759–1782, 2005.
- Derwent, R. G., Simmonds, P. G., Greally, B. R., O'Doherty, S., McCulloch, A., Manning, A. J.,
 Reimann, S., Folini, D., and Vollmer, M. K.: The phase-in and phase-out of European emis- sions
 of HCFC-141b and HCFC-142b under the Montreal Pro- tocol: Evidence from observations at
 Mace Head, Ireland and Jungfraujoch, Switzerland from 1994–2004, Atmos. Environ., 41, 757–
 767, 2007.

- 1 Dorf, M., Butler, J H., Butz, A., Camy-Peyret, C., Chipperfield, M P., Kritten, L., Montzka, S A.,
- 2 Simmes, B., Weidner, F., and Pfeilsticker, K.: Long-term observations of stratospheric bromine
- 3 reveal slow down in growth, Geophys. Res. Lett., 33, L24803, doi:10.1029/2006GL027714, 2006.
- 4 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical
- 5 Tropopause Layer, Rev. Geophys., 47, RG1004, doi:10.1029/2008RG000267, 2009.
- 6 Gettelman, A. and Forster, P. D. F.: A Climatology of the Tropical Tropopause Layer, Journal of
 7 the Meteorological Society of Japan, 80, 911–924, 2002.
- 8 Giese, B., Laturnus, F., Adams, F.C. and Wiencke, C.: Release of volatile iodinated C_1 - C_4 9 hydrocarbons by marine macroalgae from various climate zones, Environ. Sci. Technol., 33(14), 10 2432–2439, doi:10.1021/es980731n, 1999.
- 11 Goh, C. S. and Lee, K.T.: A visionary and conceptual macroalgae-based third-generation bioethanol
- 12 (TGB) biorefinery in Sabah, Malaysia as an underlay for renewable and sustainable development,
- 13 Renewable and Sustainable Energy Reviews, 14, 842–848, doi:10.1016/j.rser.2009.10.001, 2010.
- 14 Gostlow, B., Robinson, A. D., Harris, N. R. P., O'Brien, L. M., Oram, D. E., Mills, G. P., Newton,
- H. M., and Pyle, J. A.: μDirac: An Autonomous Instrument for Halocarbon Measurements, Atmos.
 Meas. Tech., 3, 507-521, 2010.
- 17 Hamilton, J. F., Allen, G., Watson, N. M., Lee, J. D., Saxton, J. E., Lewis, A. C., Vaughan, G.,
- 18 Bower, K. N., Flynn, M. J., Crosier, J., Carver, G. D., Harris, N. R. P., Parker, R. J., Remedios, J. J.,
- 19 and Richards, N. A. D.: Observations of an atmospheric chemical equator and its implications for
- 20 the tropical warm pool region, J. Geophys. Res., 113, D20313, doi:10.1029/2008JD009940, 2008.
- 21 Hewitt, C. N., Lee, J. D., MacKenzie, A. R., Barkley, M. P., Carslaw, N., Carver, G. D., Chappell,
- 22 N. A., Coe, H., Collier, C., Commane, R., Davies, F., Davison, B., DiCarlo, P., Di Marco, C. F.,
- 23 Dorsey, J. R., Edwards, P. M., Evans, M. J., Fowler, D., Furneaux, K. L., Gallagher, M., Guenther,
- 24 A., Heard, D. E., Helfter, C., Hopkins, J., Ingham, T., Irwin, M., Jones, C., Karunaharan, A.,
- 25 Langford, B., Lewis, A. C., Lim, S. F., MacDonald, S. M., Mahajan, A. S., Malpass, S., McFiggans,
- 26 G., Mills, G., Misztal, P., Moller, S., Monks, P. S., Nemitz, E., Nicolas-Perea, V., Oetjen, H., Oram,
- 27 D. E., Palmer, P. I., Phillips, G. J., Pike, R., Plane, J. M. C., Pugh, T., Pyle, J. A., Reeves, C. E.,
- 28 Robinson, N. H., Stewart, D., Stone, D., Whalley, L. K., and Yin, X.: Overview: oxidant and
- 29 particle photochemical processes above a south-east Asian tropical rainforest (the OP3 project):
- 30 introduction, rationale, location characteristics and tools, Atmos. Chem. Phys., 10, 169-199,
- 31 doi:10.5194/acp-10-169-2010, 2010.

- 1 Hosking, J. S., Russo, M. R., Braesicke, P., and Pyle, J. A.: Modelling deep convection and its
- 2 impacts on the tropical tropopause layer, Atmos. Chem. Phys., 10, 11175–11188, doi:10.5194/acp-
- 3 10-11175-2010, 2010.
- 4 Hossaini, R., Chipperfield, M. P., Monge-Sanz, B. M., Richards, N. A. D., Atlas, E., and Blake, D.
- 5 R.: Bromoform and dibromomethane in the tropics: a 3-D model study of chemistry and transport,
- 6 Atmos. Chem. Phys., 10, 719–735, doi:10.5194/acp-10-719-2010, 2010.
- 7 Hossaini, R., Chipperfield, M. P., Feng, W., Breider, T. J., Atlas, E., Montzka, S. A., Miller, B. R.,
- 8 Moore, F., and Elkins, J.: The contribution of natural and anthropogenic very short-lived species to
- 9 stratospheric bromine, Atmos. Chem. Phys., 12, 371-380, doi:10.5194/acp-12-371-2012, 2012.
- Hossaini, R., Mantle, H., Chipperfield, M. P., Montzka, S. A., Hamer, P., Ziska, F., Quack, B.,
 Krüger, K., Tegtmeier, S., Atlas, E., Sala, S., Engel, A., Bönisch, H., Keber, T., Oram, D., Mills,
- 12 G., Ordóñez, C., Saiz-Lopez, A., Warwick, N., Liang, Q., Feng, W., Moore, F., Miller, B. R.,
- Marécal, V., Richards, N. A. D., Dorf, M., and Pfeilsticker, K.: Evaluating global emission
 inventories of biogenic bromocarbons, Atmos. Chem. Phys., 13, 11819-11838, 10.5194/acp-1311819-2013, 2013.
- Jones, A., Thomson, D., Hort, M., and Devenish, B.: The UK Met Office's next-generation
 atmospheric dispersion model, NAME III, in: Air Pollution Modeling and Its Application XVII,
 edited by: Borrego, C. and Norman, A.-L., Springer US, New York, 580–589, doi:10.1007/978-0387-68854-1_62, 2007.
- Jones, C. E., Andrews, S. J., Carpenter, L. J., Hogan, C., Hopkins, F. E., Laube, J. C., Robinson, A.
 D., Spain, T. G., Archer, S. D., Harris, N. R. P., Nightingale, P. D., O'Doherty, S. J., Oram, D. E.,
 Pyle, J. A., Butler, J. H., and Hall, B. D.: Results from the first national UK inter-laboratory
 calibration for very short-lived halocarbons, Atmos. Meas. Techn., 4, 865–874, www.atmos-meastech.net/4/865/2011/, 2011.
- Kreycy, S., Camy-Peyret, C., Chipperfield, M. P., Dorf, M., Feng, W., Hossaini, R., Kritten, L.,
 Werner, B., and Pfeilsticker, K.: Atmospheric test of the J(BrONO₂)/kBrO+NO₂ ratio: implications
 for total stratospheric Bry and bromine-mediated ozone loss, Atmos. Chem. Phys., 13, 6263-6274,
- 28 doi:10.5194/acp-13-6263-2013, 2013.
- Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Worton, D. R., Sturges, W. T., Grunow, K., and
 Schmidt, U.: Contribution of very short-lived organic substances to stratospheric chlorine and

- bromine in the tropics a case study, Atmos. Chem. Phys., 8, 7325–7334, doi:10.5194/acp-8-7325 2008, 2008.
- Leedham, E. C., Hughes, C., Keng, F. S. L., Phang, S.-M., Malin, G., and Sturges, W. T.: Emission
 of atmospherically significant halocarbons by naturally occurring and farmed tropical macroalgae,
- 5 Biogeosciences, 10, 3615-3633, doi:10.5194/bg-10-3615-2013, 2013.
- Levine, J. G., Braesicke, P., Harris, N., Savage, N., and Pyle, J. A: Pathways and timescales for
 troposphere-to-stratosphere transport via the tropical tropopause layer and their relevance for very
 short lived substances, J. Geophys. Res., 112, D04308, doi:10.1029/2005JD006940, 2007.
- 9 Levine, J. G., Braesicke, P., Harris, N. R. P., and Pyle, J. A.: Seasonal and inter-annual variations in
 10 troposphere-to-stratosphere transport from the tropical tropopause layer, Atmos. Chem. Phys., 8,
 11 3689 3703, doi:10.5194/acp-8-3689-2008, 2008.
- 12 Liang, Q., Stolarski, R. S., Kawa, S. R., Nielsen, J. E., Douglass, A. R., Rodriguez, J. M., Blake, D.
- 13 R., Atlas, E. L., and Ott, L. E.: Finding the missing stratospheric Bry: a global modeling study of
- 14 CHBr₃ and CH₂Br₂, Atmos. Chem. Phys., 10, 2269-2286, doi:10.5194/acp-10-2269-2010, 2010.
- Lovelock, J. E., Maggs, R. J. and Wade, R.J.: Halogenated hydro-carbons in and over the Atlantic,
 Nature, 241, 194–196, doi:10.1038/241194a0, 1973.
- Manley, S. L. and de la Cuesta, J. L.: Methyl iodide production from marine phytoplanktoncultures, Limnol. Oceanogr., 42, 142-147, 1997.
- Manley, S. L. and Dastoor M.N.: Methyl iodide production by kelp and associated microbes, Mar.
 Biol, 98, 477-482, 1988.
- Manning, A. J., Ryall, D. B., Derwent, R. G., Simmonds, P. G., and O'Doherty, S.: Estimating
 European emissions of ozone- depleting and greenhouse gases using observations and a modelling
 back-attribution technique, J. Geophys. Res., 108, 4405, doi:10.1029/2002JD002312, 2003.
- Martino, M., Mills, G.P., Woeltjen, J. and Liss, P.S.: A new source of volatile organoiodine
 compounds in surface seawater, Geophys. Res. Lett., 36, L01609, doi:10.1029/2008GL036334,
 2009.
- Mead, M.I., Khan, M.A.H., White, I.R, Nickless G. and Shallcross, D.E.: Methyl halide emission
 estimates from domestic biomass burning in Africa, Atmos. Environ., 42 (21), 5241-5250, doi:
 10.1016/j.at-mosenv.2008.02.066, 2008c.

- Molina, M.J. and Rowland F.S.: Chlorine atom-catalysed destruction of ozone. Nature. 249, 810–
 812. doi:10.1038/249810a0, 1974.
- 3 Montzka, S. A. and Reimann, S.: Ozone-Depleting Substances (ODSs) and Related Chemicals, vol.

4 Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project –

- 5 Report No. 52, chap. 1, World Meteorological Organization (WMO), Geneva, 2011.
- Moore, R. M., and Tokarczyk, R.: Volatile biogenic halocarbons in the northwest Atlantic, Global
 Biogeochem. Cycles, 7, 195-210, 1993.
- Moore, R. M., and Zafiriou, O.C.: Photochemical production of methyl iodide in seawater, J.
 Geophys. Res., 99(D8), 16, 415-16, 420, 1994.
- Nadzir, M. S., Phang, S. M., Abas, M. R., Abdul Rahman, N., Abu Samah, A., Sturges, W. T.,
 Oram, D. E., Mills, G. P., Leedham, E. C., Pyle, J. A., Harris, N. R. P., Robinson, A. D., Ashfold,
 M. J., Mead, M. I., Latif, M. T., Mohd Hanafiah, M., Khan, M. F., and Amiruddin, A. M.:
 Bromocarbons in the tropical coastal and open ocean atmosphere during the Prime Expedition
 Scientific Cruise 2009 (PESC 09), Atmos. Chem. Phys. Discuss., 14, 953-984, doi:10.5194/acpd14-953-2014, 2014.
- 16 Newell, R. and Gould-Stewart, S: A stratospheric fountain?, J. Atmos, Sci., 38, 2789–2796, 1981.
- O'Brien, L. M., Harris, N. R. P., Robinson, A. D., Gostlow, B., Warwick, N., Yang, X., and Pyle, J.
 A.: Bromocarbons in the tropical marine boundary layer at the Cape Verde Observatory –
 measurements and modelling, Atmos. Chem. Phys., 9, 9083-9099, doi:10.5194/acp-9-9083-2009,
 2009.
- Ordóñez, C., Lamarque, J.-F., Tilmes, S., Kinnison, D. E., Atlas, E. L., Blake, D. R., Sousa Santos,
 G., Brasseur, G., and Saiz-Lopez, A.: Bromine and iodine chemistry in a global chemistry-climate
 model: description and evaluation of very short-lived oceanic sources, Atmos. Chem. Phys., 12,
 1423-1447, doi:10.5194/acp-12-1423-2012, 2012.
- Park, S., Atlas, E. L., Jiménez, R., Daube, B. C., Gottlieb, E. W., Nan, J., Jones, D. B. A., Pfister,
 L., Conway, T. J., Bui, T. P., Gao, R.-S., and Wofsy, S. C.: Vertical transport rates and
 concentrations of OH and Cl radicals in the Tropical Tropopause Layer from observations of CO2
 and halocarbons: implications for distributions of long- and short-lived chemical species, Atmos.
 Chem. Phys., 10, 6669–6684, doi:10.5194/acp-10-6669- 2010, 2010.

- 1 Pearson, G., Davies, F., and Collier, C.: Remote sensing of the tropical rain forest boundary layer
- using pulsed Doppler lidar, Atmos. Chem. Phys., 10, 5891-5901, doi:10.5194/acp-10-5891-2010,
 2010.
- 4 Perez, G.: Fisheries Profile, Bureau of Fisheries and Aquatic Resources, The Philippines,
 5 www.bfar.da.gov.ph, 2011.
- Phang, S. M.: Seaweed resources in Malaysia: Current status and future prospects, Aquatic
 Ecosystem Health & Management, 9, 2, 185-202, DOI:10.1080/14634980600710576, 2006.
- Phang, S. M., Yeong, H-Y., Lim, P-E., Rahiman A., Nor Md and Gan, K. T.: Commercial varieties
 of Kappaphycus and Eucheuma in Malaysia, Malaysian Journal of Science 29, 3, 214-224, 2010.
- 10 Pyle, J. A., Ashfold, M. J., Harris, N. R. P., Robinson, A. D., Warwick, N. J., Carver, G. D.,
- 11 Gostlow, B., O'Brien, L. M., Manning, A. J., Phang, S. M., Yong, S. E., Leong, K. P., Ung, E. H.,
- 12 and Ong, S.: Bromoform in the tropical boundary layer of the Maritime Continent during OP3,
- 13 Atmos. Chem. Phys., 11, 529-542, doi:10.5194/acp-11-529-2011, 2011.
- 14 Pyle, J. A., Warwick, N. J., Harris, N. R. P., Abas, M. R., Archibald, A. T., Ashfold, M. J.,
- 15 Ashworth, K., Barkley, M. P., Carver, G. D., Chance, K., Dorsey, J., Fowler, D., Gonzi, S.,
- 16 Gostlow, B., Hewitt, C. N., Kurosu, T. P., Lee, J. D., Langford, B., Mills, G., Moller, S.,
- 17 MacKenzie, A. R., Manning, A.J., Misztal, P., Nadzir, M. S. M., Nemitz, E., Newton, H., O'Brien,
- 18 L. M., Ong, S., Oram, D., Palmer, P. I., Peng, L. K., Phang, S. M., Pike, R., Pugh, T. A. M.,
- 19 Rahman, N. A., Robinson, A. D., Samah, A. A., Sentian, J., Skiba, U., Ung, H. E., Yong, S. E., and
- 20 Young, P.: The impact of local surface changes in Borneo on atmospheric composition at wider
- spatial scales: coastal processes, land-use change and air quality, Phil Trans R Soc B 2011 366:
 3210-3224, 2011a.
- Quack, B. and Wallace, D. W. R.: Air-sea flux of bromoform: Controls, rates and implications,
 Global Biogeochem. Cy., 17(1), 1023, doi:10.1029/2002GB001890, 2003.
- Quack, B., Atlas, E., Petrick, G., Schauffler, S., and Wallace, D.: Oceanic bromoform sources for
 the tropical atmosphere, Geophys. Res. Lett., 31, L23505, doi:10.1029/2004GL020597, 2004.
- Quack, B., Atlas, E., Petrick, G., and Wallace, D. W. R.: Bromoform and dibromomethane above
 the Mauritanian upwelling: Atmospheric distributions and oceanic emissions, J. Geophys. Res.,
- 29 112, D09312, doi:10.1029/2006JD007614, 2007.

- 1 Rasmussen, R. A., Khalil, M. A. K., Gunawardena, R. and Hoyt S. D.: Atmospheric methyl iodide
- 2 (CH₃I), J. Geophys. Res., 87(C4), 3086–3090, doi:10.1029/JC087iC04p03086, 1982.
- 3 Read, K. A., Mahajan, A. S., Carpenter, L. J., Evans, M. J., Faria, B. V. E., Heard, D. E., Hopkins,
- 4 J. R., Lee, J. D., Moller, S.J., Lewis, A. C., Mendes, L., McQuaid, J. B., Oetjen, H., Saiz-Lopez, A.,
- 5 Pilling, M. J., and Plane, J.M. C.: Extensive halogen mediated ozone destruction over the tropical
- 6 Atlantic Ocean, Nature, 453, 1232–1235, doi:10.1038/nature07035, 2008.
- Redington, A. L. and Derwent, R. G.: Calculation of sulphate and nitrate aerosol concentrations
 over Europe using a Lagrangian dispersion model, Atmos. Environ., 36, 4425–4439, 2002.
- 9 Saiz-Lopez, A., Plane, J. M. C., Baker, A. R., Carpenter, L. J., Von Glasow, R., Martín, J. C.
 10 G.,McFiggans, G., and Saunders, R. W.: Atmospheric chemistry of iodine, Chem. Rev., 112,1773–
- 11 804, doi:10.1021/cr200029u, 2012.
- Sala, S., Bönisch, H., Keber, T., Oram, D. E., Mills, G., and Engel, A.: Deriving an atmospheric
 budget of total organic bromine using airborne in-situ measurements from the Western Pacific
 during SHIVA, Atmos. Chem. Phys. Discuss., 14, 4957-5012, doi:10.5194/acpd-14-4957-2014,
 2014.
- Sander, S. P., Finlayson-Pitts, B. J., Friedl, R. R., Golden, D. M., Huie, R. E., Kolb, C. E., Kurylo,
 M. J., Molina, M. J., Moortgat, G. K., Orkin, V. L. and Ravishankara, A. R.: Chemical kinetics and
 photochemical data for use in atmospheric studies, Evaluation 14, JPL Publ. 02-25, Jet Propul. Lab.,
 Pasadena, CA, USA, 2003.
- Schall, C., Laturnus, F. and Heumann, K. G.: Biogenic volatile organoiodine and organobromine
 compounds released from polar macroalgae, Chemosphere, 28(7), 1315–1324, doi:10.1016/00456535(94) 90076-0, 1994.
- Schauffler, S. M., Atlas, E. L., Blake, D. R., Flocke, F., Lueb, R. A., Lee-Taylor, J. M., Stroud, V.,
 and Travnicek, W.: Distributions of brominated organic compounds in the troposphere and lower
 stratosphere, J. Geophys. Res., 104(D17), 21513–21535, doi:10.1029/1999JD900197, 1999.
- Seh-Lin Keng, F., Phang, S. M., Abd Rahman, N., Leedham, E., Hughes, C., Pyle, J. A., Harris, N.
 R. P., Robinson, A., and Sturges, W. T.: Volatile halocarbon emissions by three tropical brown
 seaweeds under different irradiances. J. Appl. Phycol. 25(5), 1377-1386, doi 10.1007/s10811-0139990-x, 2013.

- Simmonds, P. G., Manning, A. J., Cunnold, D. M., McCulloch, A., O'Doherty, S., Derwent, R. G.,
 Krummel, P. B., Fraser, P. J., Dunse, B., Porter, L. W., Wang, R. H. J., Greally, B. R., Miller, B. R.,
 Salameh, P., Weiss, R. F., and Prinn, R. G.: Global trends, seasonal cycles, and European emissions
 of dichloromethane, trichloroethene, and tetrachloroethene from the AGAGE observations at Mace
- 5 Head, Ireland, and Cape Grim, Tasmania, J. Geophys. Res., 111, D18304,
 6 doi:10.1029/2006JD007082, 2006.
- 7 Smythe-Wright, D., Boswell, S. M., Breithaupt, P., Davidson, R. D., Dimmer, C. H. and Diaz L. B.
- 8 E.: Methyl iodide production in the ocean: Implications for climate change, Global Biogeochem.
- 9 Cycles, 20, GB3003, doi:10.1029/2005GB002642, 2006.
- 10 Tan, J., Lim P-E. and Phang S.-M.: Phylogenetic relationship of Kappaphycus Doty and Eucheuma
- 11 J. Agardh (Solieriaceae, Rhodophyta) in Malaysia, J. Appl. Phycol., 25:13-29 DOI
- 12 10.1007/s10811-012-9833-1, 2013.
- Tokarczyk, R. and Moore, R. M.: Production of volatile organohalogens by phytoplankton cultures,
 Geophys. Res. Lett., 21(4), 285–288, 1994.
- 15 Wamsley, P. R., Elkins, J. W., Fahey, D. W., Dutton, G. S., Volk, C. M. Myers, R. C., Montzka, S.
- 16 A., Butler, J. H., Clarke, A. D., Fraser, P. J., Steele, L. P., Lucarelli, M. P., Atlas, E. L., Schauffler,
- 17 S. M., Blake, D. R., Rowland, F. S., Sturges, W. T., Lee, J. M., Penkett, S. A., Engel, A., Stimpfle,
- 18 R. M., Chan, K. R., Weisenstein, D. K., Ko, M. K. W., and Salawitch, R. J.: Distribution of halon-
- 19 1211 in the upper troposphere and lower stratosphere and the 1994 total bromine budget, J.
- 20 Geophys. Res., 103, D1, 1513–1526, 1998.
- Warwick, N. J., Pyle, J. A., Carver, G. D., Yang, X., Savage, N. H., O'Connor, F. M., and Cox, R.
 A.: Global modeling of biogenic bromocarbons, J. Geophys. Res., 111, D24305,
 doi:10.1029/2006JD007264, 2006.
- Witham, C. and Manning, A.: Impacts of Russian biomass burning on UK air quality, Atmos.
 Environ., 41, 8075–8090, 2007.
- 26 Worton, D.R., Sturges, W.T., Schwander, J., Mulvaney, R., Barnola, J.-M. and Chappellaz, J.:
- 27 20th century trends and budget implications of chloroform and related tri- and dihalomethanes
- 28 inferred from firn air, Atmos. Chem. Phys., 6 (10), 2847-2863, 2006.
- 29 Yokouchi, Y., Mukai, H., Yamamoto, H., Otsuki, A., Saitoh, C. and Nojiri, Y.: Distribution of 30 methyl iodide, ethyl iodide, bromoform, and dibromomethane over the ocean (east and southeast

- Asian seas and the western Pacific), J. Geophys. Res., 102(D7), 8805–8809, doi:10.1029/
 96JD03384, 1997.
- Yokouchi, Y., Nojiri, Y., Barrie, L. A., Toom-Sauntry, D. and Fujinuma, Y.: Atmospheric methyl
 iodide: High correlation with surface seawater temperature and its implications on the sea-to-air
 flux, J. Geophys. Res., 106(D12), 12,661–12,668, doi:10.1029/2001JD900083, 2001.
- Yokouchi, Y., Hasebe, F., Fujiwara, M., Takashima, H., Shiotani, M., Nishi, N., Kanaya, Y.,
 Hashimoto, S., Fraser, P., Toom- Sauntry, D., Mukai, H., and Nojiri, Y.: Correlations and emission
 ratios among bromoform, dibromochloromethane, and dibromomethane in the atmosphere, J.
 Geophys. Res., 110, D23309, doi:10.1029/2005JD006303, 2005.
- Yokouchi, Y., Osada, K., Wada, M., Hasebe, F., Agama, M., Murakami, R., Mukai, H., Nojiri, Y.,
 Inuzuka, Y., Toom-Sauntry, D., and Fraser, P.: Global distribution and seasonal concentration
 change of methyl iodide in the atmosphere, J. Geophys. Res., 113, D18311,
 doi:10.1029/2008JD009861, 2008.
- Yokouchi, Y., Saito, T., Ooki, A. and Mukai H.: Diurnal and seasonal variations of iodocarbons
 (CH₂ClI, CH₂I₂, CH₃I, and C₂H₅I) in the marine atmosphere, J. Geophys. Res., 116, D06301,
 doi:10.1029/2010JD015252, 2011.
- Yokouchi, Y., Nojiri, Y., Toom-Sauntry, D., Fraser, P., Inuzuka, Y., Tanimoto, H., Nara, H.
 Murakami, R. and Mukai, H.: Long-term variation of atmospheric methyl iodide and its link to
 global environmental change, Geophys. Res. Lett., 39, L23805, doi:10.1029/2012GL053695, 2012.
- Yung, Y.L., Pinto, J.P., Watson, R.T. and Sander, S.P.: Atmospheric bromine and ozone
 perturbations in the lower stratosphere. J. Atmos. Sci., 37, 339-353, doi:10.1175/1520-0469, 1980.
- Zhou, Y., Mao, H. T., Russo, R. S., Blake, D. R., Wingenter, O. W., Haase, K. B., Ambrose, J.,
 Varner, R. K., Talbot, R., and Sive, B. C.: Bromoform and dibromomethane measurements in the
 seacoast region of New Hampshire, 2002–2004, J. Geophys. Res.-Atmos., 113, D08305,
 doi:10.1029/2007JD009103, 2008.
- Ziska, F., Quack, B., Abrahamsson, K., Archer, S. D., Atlas, E., Bell, T., Butler, J. H., Carpenter, L.
 J., Jones, C. E., Harris, N. R. P., Hepach, H., Heumann, K. G., Hughes, C., Kuss, J., Krüger, K.,
 Liss, P., Moore, R. M., Orlikowska, A., Raimund, S., Reeves, C. E., Reifenhäuser, W., Robinson,
 A. D., Schall, C., Tanhua, T., Tegtmeier, S., Turner, S., Wang, L., Wallace, D., Williams, J.,
 Yamamoto, H., Yvon-Lewis, S., and Yokouchi, Y.: Global sea-to-air flux climatology for

- 1 bromoform, dibromomethane and methyl iodide, Atmos. Chem. Phys., 13, 8915-8934,
- 2 doi:10.5194/acp-13-8915-2013, 2013.

1 Table 1: μ Dirac target compounds and measurement performance at Bukit Atur and Tawau. 2 Atmospheric lifetimes are taken from Table 1-4 in Montzka and Reimann (2011); the lifetime value 3 for CH₂Br₂^{*} is actually that for CH₂Br₂; estimated accuracies are ±1 standard deviation; typical 4 measurement precisions are ±1 standard deviation of 6 calibration chromatograms made in a single 5 day; limit of detection is based on a signal to noise ratio of 10:1; n/a means no data available.

6

Compound name	Formula	Atmospheric lifetime	Estimated accuracy (%)	Typical precision (%)		Limit of detection (ppt)	
				Bukit Atur	Tawau	Bukit Atur	Tawau
tetrachloroethene	C_2Cl_4	90 days	5-10	2.5	4.0	0.04	0.02
bromoform	CHBr ₃	24 days	5-10	6.0	7.5	0.19	0.18
dibromomethane*	$\mathrm{CH}_{2}\mathrm{Br_{2}}^{*}$	123 days	5-10	5.0	6.5	0.14	0.14
methyliodide	CH ₃ I	7 days	10	n/a	11.0	n/a	0.50

7

Table 2(a-c): Calibration mixing ratios of C₂Cl₄, CHBr₃, CH₂Br₂ and CH₃I used for quantifying our data from Bukit Atur and Tawau. Calibration data for (a) the high quality NOAA SX-3542 cylinder based at Tawau; (b) data for the cylinder containing decanted NOAA cal gas (from AAL-70424) based at Bukit Atur from Nov-2008 to May-2009; (c) data for the NOAA AAL-70424 Aculife standard based at Bukit Atur from May-2009 to Jan-2010; n/a means no data available.

6

Table 2a: SX-3542 used at Tawau (November 2008 to January 2010)

Compound name	Formula	NOAA analysis (Nov-2005) (ppt)	UEA check during OP3 (Apr-2008) (ppt)
tetrachloroethene	C_2Cl_4	2.01	n/a
bromoform	CHBr ₃	9.00	8.10
dibromomethane	CH_2Br_2	4.80	4.80
methyliodide	CH ₃ I	3.40	3.30

7

Table 2b: decanted AAL-70424 used at Bukit Atur (Nov 2008 to May 2009)

Compound name	Formula	check using AAL-70424 (Oct-2008) (ppt)	check using AAL-70424 (May-2009) (ppt)
tetrachloroethene	C_2Cl_4	1.98	2.47
bromoform	CHBr ₃	2.12	3.31
dibromomethane	CH_2Br_2	2.41	4.48
methyliodide	CH ₃ I	n/a	n/a

8

 Table 2c: AAL-70424 used at Bukit Atur (May 2009 to January 2010)

Compound name	Formula	NOAA analysis (Nov-2005) (ppt)	UEA check (Apr-2009) (ppt)	check using SX-3542 (Nov-2012) (ppt)
tetrachloroethene	C_2Cl_4	2.22	2.33	1.94
bromoform	CHBr ₃	7.50	4.66	7.30
dibromomethane	CH_2Br_2	4.30	4.06	4.75
methyliodide	CH ₃ I	3.20	n/a	n/a

9

1 Table 3(a-d): Statistical summary of our observations at Bukit Atur and Tawau: monthly and overall mean, mode and median values for the mixing ratios of (a) C₂Cl₄, (b) CHBr₃, (c) CH₂Br₂ and (d) 2 CH₃I. Table 3c also shows the R^2 value for the correlation between CH₂Br₂^{*} and CHBr₃ for each 3 month, the four strongest correlations are shown in **bold**, all correlations give a positive slope 4 5 except June 2009 at Tawau (shown in italics). The total sample number varies between months and between sites due to differences in chromatogram cycle times and periods of instrument downtime. 6 Modal values are based on a bin width of 0.1 ppt. For $CH_2Br_2^*$ at Tawau the sample counts are 7 particularly low as we are only able to use about 1 in 10 sample chromatograms for reliable 8 9 quantification. The overall C_2Cl_4 distribution at both sites is distinctly bi-modal, the secondary 10 mode is shown in parentheses. n/a means no data available

Month	Sample to	tal (n)	Mean (j	Mean (ppt)		Mode (ppt)		Median (ppt)	
	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau	
Nov-2008	416	544	1.00	0.85	0.85	0.75	0.99	0.76	
Dec-2008	1748	1766	1.77	1.30	1.55	1.25	1.68	1.20	
Jan-2009	1746	1910	2.38	1.61	1.85	1.35	2.28	1.49	
Feb-2009	1572	1702	1.72	1.16	1.45	1.05	1.69	1.09	
Mar-2009	1759	1896	1.65	1.01	1.65	0.95	1.63	1.00	
Apr-2009	1216	604	1.01	0.81	0.85	0.75	0.92	0.80	
May-2009	1142	443	0.71	0.55	0.65	0.55	0.69	0.54	
Jun-2009	1314	867	0.63	0.44	0.65	0.45	0.62	0.44	
Jul-2009	1273	735	0.66	0.48	0.65	0.45	0.65	0.47	
Aug-2009	1117	725	0.61	0.74	0.55	0.55	0.58	0.74	
Sep-2009	n/a	678	n/a	0.70	n/a	0.65	n/a	0.64	
Oct-2009	n/a	1382	n/a	0.70	n/a	0.55	n/a	0.62	
Nov-2009	n/a	1296	n/a	1.22	n/a	1.25	n/a	1.22	
Dec-2009	n/a	628	n/a	2.57	n/a	2.55	n/a	2.63	
Jan-2010	n/a	273	n/a	2.31	n/a	2.25	n/a	2.25	
Overall	13303	15449	1.33	1.10	0.65 (1.65)	1.05 (0.55)	1.33	0.99	

Table 3a: Statistical summary for C_2Cl_4 at Bukit Atur and Tawau

Month	Sample to	otal (n)	Mean (ppt)		Mode (ppt)		Median (ppt)	
	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau
Nov-2008	415	544	1.69	5.87	1.25	1.65	1.41	2.63
Dec-2008	1640	1754	1.98	2.89	1.65	1.65	1.85	1.73
Jan-2009	1640	1889	2.39	1.70	1.85	1.65	2.31	1.66
Feb-2009	1526	1676	1.85	2.01	1.35	1.65	1.78	1.68
Mar-2009	1701	1886	1.69	1.72	1.65	1.35	1.66	1.46
Apr-2009	1118	604	1.81	2.38	1.15	1.65	1.57	1.68
May-2009	1020	437	1.67	1.49	1.55	1.25	1.52	1.40
Jun-2009	1247	701	0.93	1.97	0.85	1.15	0.89	1.62
Jul-2009	1305	729	1.11	2.38	0.85	1.65	1.02	1.71
Aug-2009	1119	285	1.35	2.18	1.15	2.45	1.25	2.25
Sep-2009	n/a	679	n/a	3.31	n/a	2.85	n/a	3.21
Oct-2009	n/a	1279	n/a	1.99	n/a	1.15	n/a	1.80
Nov-2009	n/a	1227	n/a	2.29	n/a	1.45	n/a	1.64
Dec-2009	n/a	580	n/a	2.48	n/a	2.05	n/a	2.21
Jan-2010	n/a	201	n/a	2.98	n/a	2.15	n/a	2.33
Overall	12731	14471	1.68	2.32	1.35	1.65	1.57	1.73

Table 3b: Statistical summary for CHBr₃ at Bukit Atur and Tawau

Table 3c: Statistical summary for CH2Br2* at Bukit Atur and Tawau

Month	Sample to	otal (n)	Mean (ppt)	Mode (ppt)	Median	(ppt)	CHBr ₃ v CH ₂	$Br_2^* R^2$
	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau	Bukit Atur	Tawau
Nov-2008	423	36	1.18	1.23	1.05	1.05	1.13	1.14	0.86	0.03
Dec-2008	1793	165	1.06	1.51	1.05	1.35	1.05	1.50	0.53	0.02
Jan-2009	1759	181	1.31	1.35	1.35	1.35	1.31	1.32	0.30	0.41
Feb-2009	1580	153	1.08	1.28	0.95	1.15	1.06	1.21	0.47	0.02
Mar-2009	1760	184	1.01	0.93	1.05	0.85	1.01	0.88	0.34	0.88
Apr-2009	1208	58	0.92	0.95	0.85	0.95	0.89	0.91	0.50	0.14
May-2009	1571	41	0.95	1.31	0.75	1.25	0.92	1.30	0.43	0.53
Jun-2009	1562	77	0.92	1.32	0.85	1.15	0.90	1.27	0.31	<0.01
Jul-2009	1555	70	1.12	1.37	1.05	1.35	1.10	1.37	0.59	0.03
Aug-2009	1134	64	1.20	1.79	1.15	1.75	1.18	1.79	0.76	0.01
Sep-2009	n/a	26	n/a	1.59	n/a	1.55	n/a	1.57	n/a	0.64
Oct-2009	n/a	115	n/a	1.81	n/a	1.55	n/a	1.60	n/a	< 0.01
Nov-2009	n/a	90	n/a	1.56	n/a	1.25	n/a	1.45	n/a	0.01
Dec-2009	n/a	42	n/a	1.49	n/a	1.35	n/a	1.49	n/a	0.07
Jan-2010	n/a	22	n/a	1.72	n/a	1.85	n/a	1.74	n/a	0.15
Overall	14345	1324	1.07	1.37	1.05	1.35	1.05	1.35	0.26	0.01

Table 3d.	Statistical	summarv	for	CH _J I at	Tawan	only
Table Su.	Statistical	summar y	101	CH31 at	Lawau	omy

Month	Sample total (n)	Mean (ppt)	Mode (ppt)	Median (ppt)
Jan-2009	933	0.46	0.25	0.38
Feb-2009	1259	0.58	0.55	0.56
Mar-2009	1459	0.55	0.45	0.51
Apr-2009	396	0.49	0.45	0.46
May-2009	234	0.68	0.55	0.64
Jun-2009	486	0.54	0.55	0.54
Jul-2009	537	0.34	0.25	0.30
Aug-2009	338	0.85	0.65	0.72
Sep-2009	488	2.50	0.95	1.69
Oct-2009	935	1.17	0.75	0.96
Nov-2009	925	0.86	0.55	0.63
Dec-2009	383	0.94	0.85	0.91
Overall	8372	0.78	0.55	0.58



Figure 1: Site location map for Sabah. The inland rainforest site on Bukit Atur is shown by the blue circle, the coastal site near Tawau is a red circle and the coastal site at Kunak (green circle) is also shown as featured in Pyle et al. (2011). The Tawau site is ~85 km to the south of Bukit Atur. The inset (lower left corner of main panel) shows the southeast Asia region with a red square to show coverage of the main map.



Month.Year

2 Figure 2: Timeline plot showing calibration comparisons and data coverage for the two sites. The 3 solid blue line shows the data coverage for Bukit Atur and is divided into two periods where 4 different calibration cylinders were used (decanted AAL-70424 from November 2008 to May 2009 5 and AAL-70424 used directly from May 2009 onwards). The solid red line shows the data coverage 6 for the Tawau site where a high quality calibration cylinder (SX-3542) was used constantly. The 7 arrows show times of the initial calibration by NOAA and the various times of comparisons made 8 for the Bukit Atur site. Other measurement periods not used in this study are shown as dotted lines, 9 including the deployment to Kunak (during the OP3 campaign) shown in green.



1

Figure 3(a-d): Time series of the four short-lived compounds (a) C_2Cl_4 , (b) CHBr₃, (c) CH₂Br₂^{*} and (d) CH₃I at Bukit Atur (blue) and Tawau (red) made between November 2008 and January 2010. High values beyond the scale of the y-axis are not shown, but are included in the box and whisker plots in Figure 4. A guide to measurement precision (±1 s.d.) is shown by the vertical error bars visible in each panel (May 2009) which show the typical precision for each compound, at each site, for the full period of data coverage. Gaps in the data are due to a combination of power outages and instrument breakdown.



Figure 4(a-g): Box and whisker plots for the halocarbon observations at Bukit Atur for (a) C₂Cl₄, 2 (b) CHBr₃, (c) $CH_2Br_2^*$ and at Tawau for (d) C_2Cl_4 , (e) CHBr₃, (f) $CH_2Br_2^*$ and (g) CH_3I . In each 3 panel, the coloured boxes show the data falling between the 25th and 75th percentiles, with the 4 median value shown as a black horizontal line inside each box. The vertical black lines above and 5 below each box indicate the 95th and 5th percentiles respectively. Outliers (values beyond the 95th 6 and 5th percentiles are shown as grey open circles. The mixing ratio (y-axis) scale is logarithmic in 7 8 order to show the extreme high values, such as for CHBr₃ at Tawau (e). The filled circles in the 9 smaller panel above each box and whisker plot show the number of days sampled each month and 10 labels indicate the number of samples used to generate each box and whisker. The total sample 11 count is different at each site due mainly to different periods of instrument down time and 12 differences in the sampling frequency at each site.



2 Figure 5(a-h): Probability distributions and median values at each site for January/February 2009 (merged) for (a) C₂Cl₄, (b) CHBr₃, (c) CH₂Br₂^{*}, (d) CH₃I (Tawau only) and June/July 2009 3 (merged) for (e) C₂Cl₄, (f) CHBr₃, (g) CH₂Br₂^{*} and (h) CH₃I (Tawau only). Bukit Atur distributions 4 5 are shown as blue circles with dotted blue lines and Tawau as red circles with dotted red lines. 6 Median values for inland Bukit Atur are shown by solid vertical blue bars and for coastal Tawau by 7 solid vertical red bars. The bin width for each probability distribution is 0.1 ppt and the area under 8 each curve is equal to 1 (probability of 1). High values beyond the scale of x-axis are not shown, for 9 instance the high tail is not wholly visible for CHBr₃ at Tawau (b & f), though all data are included 10 in the box and whisker plots (Figure 4).



3.2E-11 1.0E-07 3.2E-07 1.0E-06 3.2E-06 1.0E-05 3.2E-05 1.0E-04 3.2E-04 0-100m time integrated particle density / g s m⁻³

1

2 Figure 6(a-d): Air history maps for 2009 for Bukit Atur for selected months: (a) January; (b) March; 3 (c) June; (d) October. Maps for all months were produced but not shown. Each map is compiled 4 from 11000 particles released on an hourly basis by the NAME dispersion model using UK Met 5 Office analysed wind fields at ~50 km horizontal resolution. They show the particle density in the lowest 100 m averaged over the preceding 12 days. From November to February the air mass origin 6 7 is mainly from the northern hemisphere, often passing close to China (a). Between March and May 8 the air mass origin gradually changes towards West Pacific equatorial latitudes, away from large 9 urban regions (b). From June to September the air mass history is influenced from the region to the 10 southeast, characterised by southern hemisphere air, with sparse anthropogenic emissions (c). 11 October sees a transition back to northern hemisphere air masses (d).



1

2 Figure 7(a & b): Time series (a) of observed (red dots) and p-TOMCAT modelled (green line) 3 CHBr₃ at Tawau and probability distributions (b) for observed (red circles with red dotted line) and 4 modelled CHBr₃ (green circles with green dotted line) for December 2008 to March 2009. The 5 black vertical error bar in (panel a during May 2009) shows the typical CHBr₃ measurement 6 precision (±1 s.d.) for the whole observational period. Median CHBr₃ mixing ratios are indicated 7 (b) by the red solid line for the observations and by the green solid line for the model. The model 8 compares well overall with the observations though there are periods when the model 9 underestimates the observations (e.g. October 2009). The model does not capture the full variability 10 of the observations which is expected given the model resolution and the uniform distribution of 11 coastal emissions in the model, this is illustrated by absence of the high tail in the modelled CHBr₃ 12 (b).



Figure 8(a & b): Estimated total Br (a) from observed CHBr₃ and CH₂Br₂^{*} as a time series at Tawau 2 and (b) as probability distribution functions at Bukit Atur and Tawau. Bromine from $CH_2Br_2^*$ is 3 shown by the light red dots (panel a) and from CHBr₃ by the dark red dots, whilst Br summed from 4 5 CHBr₃ and CH₂Br₂^{*} is shown by the black dots and the heavily smoothed black line. The 6 probability distributions for Bukit Atur (blue circles with blue dotted line) and Tawau (red circles 7 with red dotted line) for the 10 month period from November 2008 to August 2009 are similar 8 (panel b). The total Br median values shown as blue (Bukit Atur) and red (Tawau) vertical lines 9 agree with each other to within 1.2 ppt. Occasionally very high total Br observed at Tawau is clearly seen (right of the x-axis break), events higher than 20 ppt were never observed at Bukit 10 11 Atur.