

# Size-dependent wet removal of BC test in Canadian biomass burning plumes

J. W. Taylor<sup>1</sup>, J. D. Allan<sup>1,2</sup>, G. Allen<sup>1</sup>, H. Coe<sup>1</sup>, P. I. Williams<sup>1,2</sup>, M. J. Flynn<sup>1</sup>, M. Le Breton<sup>1</sup>, J. B. A. Muller<sup>1</sup>, C. J. Percival<sup>1</sup>, D. Oram<sup>3</sup>, G. Forster<sup>3</sup>, J. D. Lee<sup>4</sup>, A. R. Rickard<sup>5,a</sup>, M. Parrington<sup>6,b</sup>, and P. I. Palmer<sup>6</sup>

<sup>1</sup>Centre for Atmospheric Science, School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester, UK

<sup>2</sup>National Centre for Atmospheric Science, University of Manchester, Manchester, UK

<sup>3</sup>National Centre for Atmospheric Science (NCAS), School of Environmental Sciences, University of East Anglia, Norwich, UK

<sup>4</sup>National Centre for Atmospheric Science (NCAS), Department of Chemistry, University of York, York, UK

<sup>5</sup>National Centre for Atmospheric Science (NCAS), School of Chemistry, University of Leeds, Leeds, UK

<sup>6</sup>School of GeoSciences, University of Edinburgh, Edinburgh, UK

<sup>a</sup>now at the National Centre for Atmospheric Science (NCAS), Department of Chemistry, University of York, York, UK

<sup>b</sup>now at European Centre for Medium-Range Weather Forecasts, Shinfield Park, Reading, RG2 9AX, UK

*Correspondence to:* Jonathan W. Taylor  
(jonathan.taylor@manchester.ac.uk)

**Abstract.** Wet deposition is the dominant mechanism for removing black carbon (BC) from the atmosphere, and is key in determining its atmospheric lifetime, vertical gradient and global transport. Despite the importance of BC in the climate system, especially in terms of its ability to modulate the radiative energy budget, there are few quantitative case studies of wet removal in ambient environments. We present a case study of BC wet removal by examining aerosol size distributions and BC coating properties sampled in three Canadian boreal biomass burning plumes, one of which passed through a precipitating cloud. ~~In this plume,~~ This depleted the majority of the plume's BC mass, and the largest and most coated BC particles were found to be preferentially removed, suggesting that nucleation scavenging was likely the dominant mechanism. Calculated mass absorption coefficient (MAC) in the plumes showed no significant variation, as the shifts to smaller BC cores and thinner coatings had opposing effects. Similarly, calculated single-scattering albedo (SSA) showed little variation, as a large number of non-BC particles were also present in the precipitation-affected plume. The remaining BC cores were smaller than those observed in previous studies of BC in post-precipitation outflow over Asia, possibly due to the thick coatings associated with the Canadian biomass burning particles. This study provides measurements of BC size, mixing state and removal efficiency to constrain ~~important constraints to~~ model parameterisations of BC wet removal in biomass burning regions, which will help to reduce uncertainty in radiative forcing calculations.

## 1 Introduction

Black carbon (BC) is the dominant absorbing aerosol in the atmosphere and ~~its estimated radiative forcing is second only to CO<sub>2</sub>~~ is an important, ubiquitous climate warming agent (Ramanathan and Carmichael, 2008; Chung et al., 2012; Bond et al., 2013). The optical properties of BC af-

fect the single-scattering albedo (SSA) of an aerosol layer, which determines the sign of its radiative forcing (Haywood and Shine, 1995). Important uncertainties remain regarding global and local emissions of BC, as well as its chemical processing, lifetime in the atmosphere lifetime and optical properties. Observations are required to further constrain and/or validate model parameterisations surrounding BC processes

in the atmosphere. Open biomass burning (BB) is the largest source category of BC, responsible for  $\sim 40\%$  of total emissions in the year 2000 (Bond et al., 2013), and the size distribution and mixing state of BC from this source are known to exhibit systematic differences to fossil fuel emissions (Kondo et al., 2011; Sahu et al., 2012).

Modelling studies suggest wet deposition is the dominant mechanism for BC's removal from the atmosphere (Textor et al., 2006), and consequently determines its lifetime and atmospheric burden to first order. As convection is often accompanied by precipitation, this also affects vertical profiles of number and mass concentration, which are important for determining radiative forcing (Samset et al., 2013). Such profiles are often poorly represented by models (Koch et al., 2009) as calculated removal rates are sensitive to the microphysical schemes used (Textor et al., 2006; Croft et al., 2010; Vignati et al., 2010).

In climate models, complex aerosol-cloud interactions must be simplified to reduce computing requirements, and many microphysical processes are sub-grid in both space and time. In-cloud scavenging occurs in two stages: aerosol activation to form cloud droplets, and removal of droplets by precipitation. Cloud droplet number concentration may be calculated based on empirical relations with updraft velocity and aerosol number concentration (Lohmann et al., 2007). Alternatively, many models use parameterisations designed to emulate size- and composition-based Köhler theory and the competition for water vapour between activated aerosols growing to cloud droplets (Ghan et al., 2011). Precipitation is determined by calculating autoconversion rates (the rates at which cloud water is converted to precipitation) which are tuned to simulate detailed microphysical schemes (Lin et al., 1983; Liu and Daum, 2004). Below-cloud scavenging is calculated by multiplying the precipitation rate by a scavenging coefficient, which may be size resolved, and again may be calculated empirically or based on theoretical considerations (Wang et al., 2010). Some models treat explicit size-resolved, cloud liquid, ice and mixed-phase microphysics (e.g. Jacobson, 2003), and this can generate better agreement with measured vertical profiles. Observations (particularly of size distribution, hygroscopicity and mixing state) are needed to constrain such both types of wet removal schemes (Koch et al., 2011). Explicit microphysical models are also used to investigate smaller scale phenomena (e.g. Kreidenweis et al., 1997), and ambient measurements are similarly required for constraint and comparison.

Fresh BC is generally considered hydrophobic, though it may act as a cloud condensation nucleus (CCN) in liquid cloud if subsequently coated with hydrophilic material (Khalizov et al., 2009; Liu et al., 2013). In biomass burning plumes this coating is thought to occur in the first few hours after emission (Abel, 2003; Akagi et al., 2012), and coatings are generally thicker than in fossil fuel emissions (Kondo et al., 2011; Sahu et al., 2012). BB emissions can be efficient CCN (Latham et al., 2013), and can also act as ice

nuclei (IN) (Petters et al., 2009), though different species in BB emissions may be more effective than others. Several recent studies have suggested that non-nucleation scavenging (i.e. impaction) mechanisms may be more effective for BC removal, both in mixed-phase (Twohy et al., 2010) and ice clouds (Stith et al., 2011; Baumgardner et al., 2008).

The different scavenging mechanisms exhibit different size- and composition-dependent efficiencies. In liquid cloud, BC generally requires a soluble coating to activate, and larger particles are generally more effective CCN (Dusek et al., 2006). Larger particles are also more efficient IN (Hoose and Möhler, 2012), however soluble coatings have been shown to inhibit BC ice nucleation (Koehler et al., 2009). Falling ice and raindrops may scavenge coarse mode particles and liquid droplets, and hence remove particles that have already activated (Miller and Wang, 1991; Seinfeld and Pandis, 1998; Croft et al., 2009). This type of scavenging is also more efficient for Aitken mode aerosols. For typical BC size distributions, which span the Aitken and accumulation modes, impaction scavenging therefore favours smaller BC, whereas nucleation scavenging favours larger (Jacobson, 2003).

Oshima et al. (2012) calculated removal rates of diesel-dominated BC by liquid clouds in convective airmasses over the Yellow and East China Seas. By comparing BC/CO ratios in different air parcels, they calculated the transmission efficiency of BC ( $TE_{BC}$ ), the fraction of BC not removed by precipitation. Moteki et al. (2012) showed that larger BC, which also had more coating material, were more efficiently removed during this process, meaning they must have been scavenged by nucleation. For airmasses in which the most BC was removed ( $TE_{BC} < 0.3$ ) the remaining BC core size distribution was similar to that seen by Schwarz et al. (2010b) in pristine conditions in the remote Pacific, suggesting a fraction of BC is not removed by wet processes and is able to travel long distances. In this manuscript, we present a case study of size-dependent wet removal of biomass burning BC, which may behave differently to diesel-dominated BC due to its contrasting microphysical properties. Firstly, we present an overview of the experiment, and use backtrajectories and weather satellite data to identify three plumes from Canadian boreal fires in the same region, one of which had passed through a region of a precipitating cloud. Franklin et al. (2014) recently highlighted aerosol depletion in similar plumes using remote sensing measurements, but were unable to determine the mechanism or measure the properties of any particles remaining in the plume. We examine the aerosol size distributions and BC coating properties using in-situ measurements sampled in the three plumes to determine the most likely removal mechanism and consider the effect this has on optical properties.

## 2 Experimental

### 2.1 BORTAS-B

BORTAS-B was the second phase of the BORTAS (Quantifying the impact of BOREal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites) project (Palmer et al., 2013), and took place over eastern Canada and the north Atlantic during July–August 2011. The project aimed to study the chemistry of biomass burning plumes using a combination of airborne, ground and remote sensing measurements. This paper uses airborne aerosol measurements made aboard the UK BAe-146-301 Atmospheric Research Aircraft (ARA), and focuses on a single flight (B622) on 20 July 2011. The flight track for B622 is shown on Fig. 1. The ARA took off from Halifax Stanfield International Airport at 14:56 UTC and flew a series of sawtooth and straight and level runs on a generally East-West axis, before landing at Québec City Jean Lesage International Airport at 19:11 UTC.

### 2.2 Instrumentation

#### 2.2.1 The Single Particle Soot Photometer

The Single-Particle Soot Photometer (SP2, Droplet Measurement Technologies, Boulder, CO, USA) uses laser-induced incandescence to measure refractory black carbon (rBC; hereafter referred to as BC) on a single particle basis (Schwarz et al., 2006). Particles are drawn through a high-intensity 1064 nm Nd:YAG laser which heats BC-containing particles to incandescence. The incandescent light is detected by two photomultiplier tubes (PMTs), one fitted with a narrowband filter to determine incandescence temperature, and  $\ddagger$ . During BORTAS these were run at different gain settings to expand the dynamic range of the instrument. The signal these two PMTs detect is proportional to the mass of refractory black carbon present in the particle, regardless of mixing state (Slowik et al., 2007; Moteki and Kondo, 2007), and converted to mass-equivalent core diameter ( $D_C$ ) using a density of  $1.8 \text{ g cm}^{-3}$ . The broadband PMT (measuring particles  $60 \leq D_C \leq 275 \text{ nm}$ ) was calibrated using several sizes of monodisperse Aquadag (Henkel Corporation, Bay Point, CA, USA), with particle mass calculated using the mass-mobility relationship reported by Gysel et al. (2011). This was then corrected for the difference in the response of the instrument compared to atmospheric BC (Baumgardner et al., 2012; Laborde et al., 2012b). The narrowband PMT (measuring  $90 \leq D_C \leq 550 \text{ nm}$ ) was then cross calibrated to the broadband in the overlap region. The instrument was run with sufficient laser power to ensure 100% detection efficiency for BC particles  $\geq 1 \text{ fg}$  (femtogram) (Schwarz et al., 2010a). In plumes, the statistical uncertainty in BC mass concentration was typically  $\sim 10\%$  for an averaging time of 10 s, and accuracy within 20% due to possible differences between

the instrumental response to biomass burning BC and the calibrant (Laborde et al., 2012a).

The instrumental setup of the SP2 on the ARA was described by McMeeking et al. (2010), and is briefly summarised here. The SP2 sampled through a 1/4 inch stainless steel tube connected to a modified Rosemount inlet (Foltescu et al., 1995). Though recent results have shown that Rosemount inlets can enhance aerosol measurements at larger diameters and higher densities, the vast majority of BC mass is present in particles  $< 600 \text{ nm}$ , where the efficiency is close to unity at densities typical of biomass burning emissions (Trembath et al., 2012). Losses between the inlet and the SP2 were minimised using a bypass flow of  $1 \text{ l min}^{-1}$ , and this also reduced the residence time in the inlet.

The SP2 is also capable of measuring light scattering particles (LSP, particles that do not contain BC), and the coating properties of BC-containing particles. Two avalanche photodiodes (APDs) detect scattered light at 1064 nm, and the detected signal is proportional to the scattering cross section of the particle. One APD is position-sensitive, and this provides a measure of the position of the particle in the laser (Gao et al., 2007). This allows calculation of the fractional laser power at the edge of the beam, as the laser has a Gaussian profile when configured correctly. Leading-edge only (LEO) fitting is then used to calculate the scattering cross section of the particle before it begins to evaporate. In this work we considered the leading edge as data from the baseline up to 5% of the maximum laser intensity.

Mie core/shell modelling is then used to infer the coated particle diameter ( $D_P$ ), and hence coating thickness, though this is dependent on the assumed core density, and the refractive index of the BC core and coating (Taylor et al., 2014b). In this analysis we assumed a BC core density of  $1.8 \text{ g cm}^{-3}$  (Bond and Bergstrom, 2006) and core refractive index  $n_C = (2.26 - 1.26i)$  (Moteki et al., 2010). These parameters were shown by Taylor et al. (2014b) to be the most appropriate for use with fossil fuel BC, by comparing Mie calculations to the scattering by thermodenuded particles. As a similar analysis of biomass burning emissions has not been performed, we used these parameters on the assumption that the core properties of fossil fuel and biomass burning BC are broadly similar. For the shell refractive index, we have assumed a value of  $n_S = (1.5 + 0i)$  (Reid et al., 2005a).

#### 2.2.2 Aerosol mass spectrometer

Nonrefractory submicron aerosol mass concentration was measured with a compact time-of-flight Aerosol Mass Spectrometer (AMS, Aerodyne Research Inc, Billerica, MA, USA.) (Drewnick et al., 2005; Canagaratna et al., 2007). The AMS reports Organic Aerosol (OA), sulphate, nitrate, ammonium and chloride mass concentrations, though in this analysis we only consider the OA concentrations. The operation of the AMS on the ARA has been described previously (Crosier et al., 2007; Morgan et al., 2009). The AMS

was connected to a Rosemount inlet and the aerosol flow is considered dry due to ram heating and the increase in temperature as the sample line passes into the cabin.

Data were processed as described by Morgan et al. (2010), using a collection efficiency of 0.5. Unfortunately, after landing on B622 the aircraft lost power, and the AMS had to be shut down before an Ionisation Efficiency (IE) calibration could be performed. For B622 the calibration for the following flight (B623) was applied, which may have introduced a systematic error in the AMS data. As an estimate of the size of this error, for the seven nitrate calibrations performed after B622, the standard deviation of the IE/airbeam was 13% of the mean. The calibrations before this were discarded because of a faulty power supply module in the AMS, which was replaced before B622. Also, in this case study the AMS data is only used to monitor relative changes in the OA/CO ratio, so this increased uncertainty does not affect the analysis method or conclusions.

### 2.2.3 Scanning mobility particle sizer

A custom-made Scanning Mobility Particle Sizer (SMPS) system, sampling from the same inlet as the AMS, measured ~~particle-mobility-size-distributions~~ distributions of particle mobility diameter ( $D_{mob}$ ) divided into 28 logarithmically-spaced bins in the range 20–350 nm. A custom-built low pressure Water-based Condensation Particle Counter (WCPC model 3786-LP, Aerosol Dynamics, Berkeley, CA, USA / Quant Technologies, Blaine, MN, USA) was connected to a TSI 3081 Differential Mobility Analyser (DMA). The WCPC drew  $0.6 \text{ l min}^{-1}$ , half of which was used as the sample flow, and the DMA sheath flow was  $6 \text{ l min}^{-1}$ . The data was inverted using routines developed by Zhou (2001), over a 30 s averaging time.

### 2.2.4 Supporting measurements

Aerosol light scattering coefficient ( $B_{Sca}$ ) was measured using an integrating nephelometer (TSI 3563, St. Paul, MN, USA). Here we only consider the scattering measurement at 550 nm, though data from the instruments other two wavelengths were used to correct for truncation and the non-lambertian light source (Anderson and Ogren, 1998). Relative humidity (RH) measured in the nephelometer remained below 40%, so no scattering enhancements due to swollen aerosol are expected (Magi, 2003). CO was measured by vacuum-UV resonance fluorescence, with accuracy of  $\pm 3\%$  and precision of 1 ppb at 1 Hz (Gerbig et al., 1999).  $\text{CH}_3\text{CN}$  (acetonitrile) was measured with a Proton-Transfer Reaction Mass Spectrometer (PTR-MS) (Murphy et al., 2010) and HCN (hydrogen cyanide) with a Chemical-Ionisation Mass Spectrometer (CIMS) (Le Breton et al., 2013).

## 2.3 Data analysis and plume identification

The SP2 and instruments listed in Sect. 2.2.4 record data at 1 Hz, and were averaged to 10 s. The AMS records data every 8–9 s and so, for comparison, CO was also averaged to the AMS time series. SMPS scans were averaged over the relevant in-plume periods. All aerosol measurements were corrected to standard temperature (273.15 K) and pressure (1013.25 hPa).

We identified three distinct plumes from B622 which were sampled during straight and level runs, flying at fixed heading and altitude. The times and locations of the plumes are listed ~~in the supplementary material~~ in Table S1. These plumes were selected as each plume had a large range of enhanced CO concentration above the regional background and good correlation ( $R^2 \geq 0.55$ ) between CO and  $\text{CH}_3\text{CN}$ , which are commonly-used biomass burning tracers. Out-of-plume data were excluded using the thresholds defined by Palmer et al. (2013); only data with  $\text{CH}_3\text{CN} \geq 150 \text{ ppt}$ ,  $\text{CO} \geq 148 \text{ ppb}$  and  $\text{HCN} \geq 122 \text{ ppt}$  were considered in-plume (see O’Shea et al. (2013) for a further discussion of the BORTAS airborne chemical climatology). These thresholds were determined from the 99th percentile measured in background air on flight B625 on 24 July 2011, on which no biomass burning plumes were detected (Palmer et al., 2013).

## 2.4 Backtrajectories and meteorology (moved from Section 3)

By examining Lagrangian backtrajectories along with fire location, meteorological fields and satellite data, we can place each plume into the context of its contrasting air mass history. Five-day backtrajectories were calculated using the Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPLIT) (Draxler and Hess, 1998), and are shown in Fig. 1. The trajectories were initialised using representative times and co-ordinates from the centre of each plume (defined as the mid-point of each straight-and-level run). The HYSPLIT model was driven with full vertical dynamics using 3-hourly NCEP GDAS reanalysis fields on a 1-degree global grid and 23 model (sigma) levels. A comparison with O’Shea et al. (2013, Fig. 6c), who performed back trajectories along the entire B622 flight track, shows that these trajectories are representative of the air mass history at these altitudes.

Although HYSPLIT (and indeed any Lagrangian model driven by coarse resolution meteorological fields) cannot capture convection (and hence transport) in isolated convective systems, it can effectively model vertical transport in mesoscale and synoptic features such as fronts (Stohl et al., 2001). As we shall discuss later when examining satellite imagery and meteorological fields, passage through cloud in our “precipitating” trajectory in this study was through frontal uplift in the region of active fires, which gives us confidence in the appropriate use of HYSPLIT in this convective regime.

We limit our historical examination of these backtrajectories to five days here due to the rapidly increasing uncertainty in the path of airmasses beyond this time, and the fact that we cannot trust the accuracy of trajectories that are suggested to have spent significant periods in the boundary layer or in close contact to the ground.

Moderate Resolution Imaging Spectrometer (MODIS) fire counts (Giglio et al., 2003) for 18–20 July are also shown in Fig. 1, though this only shows fires that were not masked by cloud. The trajectories for all three plumes are seen to have passed over a large region of forest fires in northwest Ontario, to the southwest of Hudson Bay and east of Lake Winnipeg. Plumes 2 and 3 were sampled approximately  $\sim 1$  day after passing over the fires, while plume 1 was sampled after  $\sim 2$  days. The meteorology data used to calculate the HYSPLIT backtrajectories does not take account of likely pyroconvection from the fires. Therefore the smoke plumes are expected to have formed a vertical emission profile which would be expected to join the trajectories at the altitudes shown in Fig. 1 over the region. It is therefore not possible to identify a particular fire as the source of a plume, but all three plumes are clearly observed to have passed over the same region of fires.

The back trajectory from plume 3 showed calculated precipitation between 11:30–13:00 and 21:00–22:30 UTC on July 19 2011 (approximately 27 and 18 hours before sampling respectively), as indicated by the aqua-coloured regions in Fig. 1. The precipitation rates are calculated by HYSPLIT using the reanalysis meteorology data. Figure 2 shows cloud data measured by the Geostationary Operational Environmental Satellite (GOES) during these two time periods, and highlights the region where the three back trajectories were at these times. The trajectories for plumes 1 and 2 clearly avoided regions of deep convective cloud, though plume 1 was below cloud at 13:00–14:45 UTC. During both periods that showed precipitation, the trajectory for plume 3 was co-located with a cloud system associated with a warm front, with effective cloud height in excess of 10 km and cloud depth up to 7 km, indicative of deep convection, potentially enhanced by pyroconvection at that time. The retrieved Ice Water Path (IWP) gives a qualitative indication of precipitation in these clouds. In Fig. 2, it is clear that the frontal cloud that plume 3 passed through was not only 7 km deep but also has a core of IWP in excess of  $1000 \text{ g m}^{-2}$ , suggesting that this deep convective (frontal) cloud was actively precipitating (Lin and Rossow, 1997).

Uncertainty in the calculated trajectories and the coarse resolution of the meteorology data used by HYSPLIT mean that the calculated precipitation rates cannot necessarily be considered accurate, but do highlight the presence of precipitating clouds in the area near where the plume passed. The GOES data confirm the presence of a deep convective cloud in this region that was likely to have been precipitating. In the next section we examine aerosol emission ratios and demonstrate that most of the aerosol in this plume must have been removed by precipitation.

## 3 Results

### 3.1 Chemical tracer Emission ratios

Figure 3 shows the BC/CO, OA/CO and BC/ $B_{Sca}$  ratios measured in for the three plumes described in Sect. 2.3. The gradients of BC/CO and OA/CO are listed in Table 1, while the BC/ $B_{Sca}$  gradients are listed in Table 3 and discussed later in Sect. 3.3. These ratios are a product of the initial emission ratios and physical processes between emission and measurement.

Both BC and OA may be affected by wet removal, and OA/CO may increase or decrease due to evaporation or condensation of OA (Donahue et al., 2011). While there was some variation between Plumes 1 and 2, which did not show evidence of precipitation, this is not unexpected. Several studies have demonstrated the dependence of aerosol emission ratios on combustion conditions (e.g. McMeeking et al., 2009; Kondo et al., 2011), as BC is more associated with flaming combustion and OA with smouldering. Although CO<sub>2</sub> data was available for the campaign, which can be used in conjunction with CO to characterise combustion efficiency (Ward and Radke, 1993), we did not consider this calculation robust as the variation in CO<sub>2</sub> background was greater than the excess in the plumes, meaning the derived slopes ( $\partial\text{CO}/\partial\text{CO}_2$ ) may be misleading (Yokelson et al., 2013). This issue did not affect the aerosol or CO measurements as the background variations were small compared to the excess levels in the plumes.

Table 1 presents a comparison of tracer emission ratios with several other studies and emission inventory values. This is not intended to be a comprehensive review of all previous measurements emission ratios, but to demonstrate the range of values typically observed. BC/CO in Plume 3 was significantly below the other reported values that were unaffected by precipitation, and was most similar to those measured during ICARTT in Alaskan/Canadian boreal plumes that were also affected by precipitation (Val Martín et al., 2006). BC/CO was an order of magnitude lower in Plume 3 than the average for plumes unaffected by precipitation. While the poor correlation between OA and CO in Plume 3 means this ratio cannot be considered reliably accurate, it is clear that it was also significantly below all other measurements listed. By comparing to literature values and the other plumes in this study, it is clear that both BC/CO and OA/CO in Plume 3 were largely the result of the wet removal itself, rather than the initial combustion conditions.

Figure 3c shows the correlations between BC mass concentration and  $B_{Sca}$  measured by the nephelometer. For Plume 3, this provides a better comparison than BC/OA, as the AMS signal/noise was low at high altitude. There was excellent correlation in all three plumes, meaning there was no significant difference in the removal efficiencies of the bulk of BC and nonrefractory mass. Comparing the slopes in

Fig. 3a and b, it is clear that the majority of BC and organics were removed in Plume 3.

### 3.2 Aerosol size and coating distributions

In this section we examine the effects of the precipitation event on particle size distributions. Biomass burning size distributions at source show a large amount of variation (Reid et al., 2005b). Additionally, aerosol concentrations in biomass burning plumes mean that coagulation can significantly affect the size distribution over timescales of hours (e.g. Capes et al., 2008). It is therefore not always possible to compare size distributions between different fires.

#### 3.2.1 Bulk aerosol size distributions

Figure 4 shows SMPS size distributions for the three plumes in this case study. Plumes 1 and 2 had count median core diameter (CMD) of 196 and 194 nm respectively, whereas Plume 3 had a CMD of 101 nm. The particles in Plumes 1 and 2 were larger than those reported for fresh Canadian boreal BB plumes during ARCTAS-B (Latham et al., 2013), due to a different source profile and/or coagulation. The similarity between Plumes 1 and 2 is likely as they were from the same set of fires and had similar transport times. We can therefore infer that had Plume 3 not been affected by precipitation, it ~~was~~would likely have had a similar size distribution. As the particles in Plume 3 were much smaller than in Plumes 1 and 2, this suggests nucleation scavenging was the wet removal mechanism, as this process favours larger particles (Dusek et al., 2006). The remaining particles are likely interstitial in the cloud (i.e. not activated), and this is qualitatively consistent with the difference between cloud residual and interstitial size distributions seen previously (Allan et al., 2008).

OA comprised over 85% of the aerosol measured in Plumes 1 and 2. The fraction of OA mass measured in the AMS at  $m/z$  44 ( $f_{44}$ ) provides a measure of the oxygenation of the OA fraction (Ng et al., 2010). The mean values of  $f_{44}$  measured in Plumes 1 and 2 were 0.085 and 0.120 respectively, which are indicative of a reasonable degree of oxidation. Previous studies have shown that increased  $f_{44}$  is qualitatively related to increased hygroscopicity (e.g. Duplissy et al., 2011), though this is to an extent system dependent. The  $f_{44}$  in these plumes indicates the OA was likely hygroscopic and may act as a CCN at sufficient supersaturations. The mass of OA measured at  $m/z$  44 in Plume 3 was not sufficient to make a robust calculation of  $f_{44}$ , so it is not possible to make a comparison.

#### 3.2.2 BC core size distributions

Although coagulation can quickly affect bulk size distributions, BC-containing particles represent a small fraction of the total number concentration. Kondo et al. (2011) found

little variation in BC core size distributions from Canadian biomass burning, meaning the effects of BC-BC coagulation are likely to be minor over the timescale of 1–2 days. Figure 5 shows the BC core size distributions in the three plumes, and a comparison to literature values of CMD and mass-median core diameter (MMD) is presented in Table 2. The BC mass distributions of Plumes 1 and 2 were very similar, though there was a smaller mode in the number distribution for Plume 1, which may suggest some limited mixing with fossil fuel emissions in the boundary layer. This is unlikely to affect total BC mass as the mode is not clear in the mass distribution. The BC CMDs for Plumes 1 and 2 therefore show a difference of  $\sim 10$  nm whereas the MMDs are nearly identical.

The BC CMD is a more qualitative measurement than the MMD as it is more sensitive to the detection range of the instrument. The SP2 typically has good detection efficiency only for particles  $\geq 1$  fg, equivalent to  $\sim 100$  nm in diameter (Schwarz et al., 2010a). For smaller BC cores, the detection efficiency drops down to zero, though the rate at which this occurs depends on the laser power. Not only is it possible therefore that the BC CMD is sensitive to the instrumental setup of the SP2, it is also possible that the true CMD is too small to be detectable at all (e.g. Liggio et al., 2012). A significant fraction of the total BC number is  $< 1$  fg, but as the smallest BC particles contribute little to BC mass, the MMD is well above the cutoff of the instrument. We therefore consider MMD a more robust variable to test for variation in BC size distribution.

The literature values of BC MMD in Table 2 range from 194–214 nm, and Plumes 1 and 2 fall within the lower end of this range. These also compare very well to Canadian plumes from ARCTAS, which were sampled close to emission (Kondo et al., 2011). Plume 3 shows a distribution skewed to smaller sizes, with MMD of 152 nm, smaller even than measured in some fossil fuel emissions (Sahu et al., 2012). The similarity of the size-dependence of wet removal between bulk aerosol and black carbon (shown in Figs. 4 and 5 respectively) suggests that BC-containing particles were also removed by nucleation scavenging, rather than another mechanism. As Fig. 3 shows the vast majority of BC mass was removed in the plume, and Fig. 5 shows there was still a significant overlap in the size distributions, this shows that the majority of BC particles across most of the size distribution were removed by the precipitation event. In order to activate, BC requires coating by hydrophilic material (Zhang et al., 2008; Khalizov et al., 2009). We examine coatings measured by the SP2 in the next section.

#### 3.2.3 BC coatings distributions

BC coatings were calculated for BC particles in the range  $130 \leq D_C \leq 230$  nm. This range was determined using the method outlined by Taylor et al. (2014b). Figure 6 shows the measured distributions of coating thickness, expressed

in terms of shell/core ratio and absolute coating thickness. The median coating thicknesses are also listed in Table 2. In all three plumes, nearly all particles had measurable coatings. Plumes 1 and 2 had broadly similar distributions, with similar median shell/core ratios of 2.34 and 2.35 and absolute coating thicknesses of 110 nm and 107 nm respectively. Plume 2 had a slightly higher proportion of less-coated particles, probably due to mixing with a small amount of anthropogenic emissions in the boundary layer. However, this does not appear to have significantly affected the shape of the distribution, or the median values. Plume 3 had thinner coatings overall, with median shell/core ratio of 2.02 and absolute coating thickness of 79 nm. This suggests that the more coated particles were more effective CCN, and hence were removed more efficiently like due to both the increased  $D_p$  and greater soluble content.

Figure 5 compares the BC mass and number distributions to the range of BC core diameters that was used for calculating coating properties,  $130 \leq D_C \leq 230$  nm. For Plumes 1 and 2 this range spans the mode of the mass distribution and is just above the mode of the number distribution, whereas for Plume 3 it borders the mode of the mass distribution and is above the mode of the number distribution. However, as this range is larger than the number mode in all three plumes, the median coating properties are weighted towards the lower end of this range. The range is not significantly far from the mass mode in Plume 3, and so it is likely representative of the bulk of BC mass in all three plumes.

### 3.3 The effect of wet removal on optical properties

Figure 3c shows correlations between BC mass loading and  $B_{Sca}$ , and the slopes are listed in Table 3. While there is over a factor of two spanning the slopes, this does not appear to be governed by the wet removal. The slopes for Plumes 1 and 2 show the greatest difference, meaning this is probably due to the initial combustion conditions and photochemical processing that occurred between emission and measurement, though Plume 1 has a possible minor contribution by mixing with other aerosols in the boundary layer. While the shift to smaller sizes will have reduced the per-particle scattering cross sections, it also reduced the average BC mass, and so this ratio was not as affected as the BC/CO or OA/CO. As noted in Sect. 3.2.2, the majority of BC particles were removed by the precipitation event over most of the size range.

Table 3 also lists the Mass Absorption Coefficient (MAC) for the three plumes, calculated at 550 nm using the SP2 data. This calculation is integrated over the entire measured BC size distribution. Though the coating properties were calculated using a BC core refractive index of  $n_{core} = (2.26 - 1.26i)$  at 1064 nm, this is not appropriate for use at visible wavelengths when using the Mie model (Taylor et al., 2014a). We therefore used  $n_{core} = (1.85 - 0.71i)$ , the central refractive index listed by Bond and Bergstrom (2006). A sensitivity analysis of this assumption is presented in Ta-

ble S2 in the supplementary material. We also assumed that BC coatings and particles not associated with BC did not absorb. While this assumption may not hold at shorter wavelengths, brown carbon absorption at 550 nm is highly uncertain (McNaughton et al., 2011; Kirchstetter and Thatcher, 2012; Lack et al., 2012), and we are unable to constrain this using these observations. We are also unable to constrain any hygroscopic growth the particles may undergo in ambient humidities, and therefore all our calculations are performed under the dry ( $< 40\%$  RH conditions of the sample line).

As the coating distributions were only measured in the range  $130 \leq D_C \leq 230$  nm, coatings at other core sizes must be inferred from this size range. Table 3 presents MAC calculated assuming either a constant distribution of shell/core ratio or a constant distribution of absolute coating thickness, as shown in Fig. 6. A constant shell/core ratio distribution gives thicker absolute coatings to larger cores, whereas a constant absolute coating thickness distribution gives large shell/core ratios to smaller cores. The differences in calculated MAC between the two cases are 15–20%. A more realistic treatment of the variation in BC coatings with core size would require detailed modelling of condensation and coagulation processes, and is not in the scope of this analysis.

For comparison, McNaughton et al. (2011) separated out the different contributions of brown carbon, mineral dust and BC in boreal BB plumes during ARCTAS/ARCPAC, and reported average MAC of BC of  $9.5 \pm 0.6 \text{ m}^2 \text{ g}^{-1}$  at 530 nm. We note however that this was derived using a filter-based absorption measurement, which suffer from systematic errors that must be corrected empirically (Lack et al., 2008; Cappa et al., 2008). MAC calculated in Plumes 1 and 2 using the absolute coating thickness distribution shows good agreement with this average, though absolute agreement using the shell/core ratio distribution is possible if using a smaller (real and imaginary) core refractive index (see Table S2). Additionally, Table 2 shows the BC core size distributions discussed here were smaller than those measured during ARCTAS, which would have increased MAC. The coatings measured here were also thicker than those reported during ARCTAS/ARCPAC by Kondo et al. (2011), which would also increase MAC, though these measurements are not absolutely comparable due to the different methods used (Taylor et al., 2014b).

Regardless of which method was used to incorporate the coating distributions, the modelled MAC in the three plumes varies by  $< 10\%$ . This is in spite of clear differences in the core size and coatings distributions between Plumes 1 and 2 and Plume 3. Figure 7 compares the BC core size distributions to the size-dependent MAC, calculated using the measured distribution of shell/core ratio for each plume. For a given core diameter, Plumes 1 and 2 have generally higher MAC than Plume 3, as they had thicker coatings. However, as the core size distribution in Plume 3 was smaller, this largely

cancels out this change, and therefore the calculated MAC shows little variation between the plumes.

Multiplying the calculated MAC by the BC/CO ratios in Table 1 gives  $B_{Abs}/CO$ , black carbon absorption per ppbv of CO. While there is some variation between Plumes 1 and 2, Plume 3 is an order of magnitude lower, due to its lower BC/CO ratio. As the MAC does not vary significantly between the plumes, BC/CO was the controlling factor determining aerosol absorption in these plumes. Combining these values with the measured  $BC/B_{Sca}$  one can calculate SSA using

$$SSA = \frac{1}{\left(1 + \frac{B_{Abs}}{B_{Sca}}\right)} = \frac{1}{\left(1 + MAC \times \frac{BC}{B_{Sca}}\right)} \quad (1)$$

Calculated bulk SSA are presented in Table 3. SSA was high in all plumes, even for aged smoke (Reid et al., 2005a), though this would be reduced slightly if absorption by brown carbon was included. They also compare well to SSA measured in boreal BB plumes during ARCTAS/ARCPAC (McNaughton et al., 2011). Though there was some variation between the plumes, Plume 3 was the central value. This shows that wet removal had only a minor effect on SSA compared to the burn conditions and secondary condensation.

## 4 Discussion

Moteki et al. (2012) listed CMD for BC core size distributions in Asian outflow after different amounts of wet removal. In free troposphere outflow the CMD was 132 nm (corrected to a BC core density of  $1.8 \text{ g cm}^{-3}$ ) in air masses with transmission efficiency  $TE_{BC} > 80\%$  and 100 nm for  $TE_{BC} < 30\%$ . While this shows excellent agreement with the CMDs measured in this analysis, the CMD is sensitive to the detection efficiency at the lower cut-off of the BC number distribution.

Moteki et al. (2012) also listed  $\langle m \rangle$ , the mean BC mass per BC-containing particle. Based on their fit data, this ranged from 5.87 fg for  $TE_{BC} = 1$  and 2.74 fg for  $TE_{BC} = 0$ . For comparison,  $\langle m \rangle$  was 3.65 fg, 4.17 fg and 1.86 fg for Plumes 1, 2 and 3 respectively. We have not quantified  $TE_{BC}$  in this study, but based on by dividing the BC/CO ratios listed in Table 1 it is  $\sim 0.1$  was  $\sim 0.1$ – $0.2$ . As the initial size distributions and coatings were different it is difficult to say conclusively if the removal efficiency was the same between the two studies. The size distributions of BC remaining after wet removal were qualitatively similar, though the initial and remaining mass distributions were both smaller in this study. What is clear between the two cases is that nucleation scavenging was an effective mechanism for removing coated BC, and that only the smallest particles were left behind. In this study we have also demonstrated that less coated BC are also less effectively removed.

The MMD in Plume 3 was significantly smaller than those measured in remote Pacific air masses by Schwarz et al.

(2010b), which were consistently within a few nm of 186 nm (corrected to BC core density of  $1.8 \text{ g cm}^{-3}$ ). Schwarz et al. (2010b) suggested the consistency (and narrow shape) of this size distribution implied more than one process was responsible for regulating this distribution, one which preferentially removed larger BC cores and ones which removed smaller. Considering our results and those of Moteki et al. (2012), it is clear that nucleation scavenging is an effective mechanism for shifting BC core size distributions to smaller sizes. Additional measurements of BC wet removal in different environments (and involving different mechanisms) are required to further constrain these processes.

The result that wet removal did not significantly affect the SSA and MAC of Plume 3 is important. Many climate models divide BC into two modes with different hydrophilicity, and assume a conversion from the hydrophobic to the hydrophilic mode over time. Bond et al. (2006) recommended a simple optical aging scheme in which the hydrophobic mode had the optical properties of fresh BC, and the hydrophilic had that of aged BC, and some models have implemented this scheme (e.g. Goldenson et al., 2012). This was based on the assumption that coatings were the only factor responsible for BC hydrophilicity. In this case study both hydrophobic and hydrophilic BC had similar MAC, as the BC core size distribution was different between the two modes. This provides a point of reference for models, and highlights the fact that coatings are not the only factor that affects MAC.

The fact that the SSA was not significantly affected by precipitation suggests that some fraction of nonabsorbing particles also did not activate, presumably because they were too small. For comparison, the mean BC number concentration in Plume 3 was  $34 \text{ scm}^{-3}$ , whereas the mean number concentration of LSP measured by the SP2 was  $718 \text{ scm}^{-3}$ . A range of hydrophilicity and size-dependent activation efficiency would be expected of all particles, and these results show that BC activated with similar efficiency to LSP.

## 5 Conclusions

We have presented a case study of black carbon wet removal by comparing three biomass burning plumes from the same region of fires. Through the use of back trajectories and weather satellite data we identified that one of these plumes convected up into a likely precipitating cloud. The BC/CO and OA/CO ratios in this plume were much lower than those in the other two plumes, and almost all those found in literature. The only comparable literature values were BC/CO ratios in plumes that had also been affected by precipitation. Taken together, the back trajectory, satellite and chemical tracer emission ratios suggest that most of the aerosol in this plume had been removed by precipitation. By comparing particles in this plume to the two unaffected by precipitation, we have made deductions about the properties of the particles that were rained out.



Upon examination of the bulk size distributions it was clear that the larger particles were preferentially scavenged, meaning nucleation scavenging must have been the wet removal mechanism. Similarly, the largest and most thickly coated black carbon particles were also removed more efficiently, meaning they must have undergone the same nucleation mechanism. Removing particles with the largest BC cores and also those with the thickest coatings has opposing effects on MAC, and so the calculated values were similar in all three measured plumes. A large number of LSP were also present in the precipitation-affected plume, so similarly the SSA (based on measured scattering and calculated absorption) was not obviously altered by the precipitation event.

The size-dependence of wet removal is an important process for moderating BC lifetime, vertical profile and transport to remote environments. While the observation that larger particles were more efficiently scavenged is similar to those of Moteki et al. (2012), the residual size distribution measured here was smaller. This may be due to thicker coatings on these particles or simply because the original size distribution was smaller than the one described by Moteki et al. (2012). To date there have been very few studies of BC wet removal. Further measurements of BC wet removal in different environments are required to improve model parameterisations and reduce what is currently a major source of uncertainty in radiative forcing calculations.

**Acknowledgements.** The authors wish to thank all those involved in the BORTAS project. Airborne data was obtained using the BAe-146-301 Atmospheric Research Aircraft [ARA] flown by Directflight Ltd and managed by the Facility for Airborne Atmospheric Measurements (FAAM), which is a joint entity of the Natural Environment Research Council (NERC) and the Met Office. Processed data are available from the British Atmospheric Data Centre. Raw data are archived at the University of Manchester and available on request. BORTAS was supported by NERC under grant number NE/F017391/1 and NE/F017200/1. JWT received extra support from the National Centre for Atmospheric Science (NCAS) for travel and subsistence and JWT and MLB were also supported by NERC PhD studentships. PIP acknowledges financial support from the Leverhulme Trust. We thank Patrick Minnis and Chad Thee at NASA Langley Research Centre, USA, for providing the GOES satellite data. The topographical map in Fig. 1 was produced using GTOPO30 data, available at <https://lta.cr.usgs.gov/GTOPO30>

## References

- Abel, S. J.: Evolution of biomass burning aerosol properties from an agricultural fire in southern Africa, *Geophysical Research Letters*, 30, 1783, doi:10.1029/2003GL017342, <http://doi.wiley.com/10.1029/2003GL017342>, 2003.
- Akagi, S. K., Craven, J. S., Taylor, J. W., McMeeking, G. R., Yokelson, R. J., Burling, I. R., Urbanski, S. P., Wold, C. E., Seinfeld, J. H., Coe, H., Alvarado, M. J., and Weise, D. R.: Evolution of trace gases and particles emitted by a chaparral fire in California, *Atmospheric Chemistry and Physics*, 12, 1397–1421, doi:10.5194/acp-12-1397-2012, <http://www.atmos-chem-phys.net/12/1397/2012/acp-12-1397-2012.html>, 2012.
- Allan, J. D., Baumgardner, D., Raga, G. B., Mayol-Bracero, O. L., Morales-García, F., García-García, F., Montero-Martínez, G., Borrmann, S., Schneider, J., Mertes, S., Walter, S., Gysel, M., Dusek, U., Frank, G. P., and Krämer, M.: Clouds and aerosols in Puerto Rico – a new evaluation, *Atmospheric Chemistry and Physics*, 8, 1293–1309, doi:10.5194/acp-8-1293-2008, <http://www.atmos-chem-phys.net/8/1293/2008/acp-8-1293-2008.html>, 2008.
- Anderson, T. L. and Ogren, J. A.: Determining Aerosol Radiative Properties Using the TSI 3563 Integrating Nephelometer, *Aerosol Science and Technology*, 29, 57–69, doi:10.1080/02786829808965551, <http://dx.doi.org/10.1080/02786829808965551>, 1998.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochemical Cycles*, 15, 955–966, doi:10.1029/2000GB001382, <http://doi.wiley.com/10.1029/2000GB001382>, 2001.
- Baumgardner, D., Subramanian, R., Twohy, C., Stith, J., and Kok, G.: Scavenging of black carbon by ice crystals over the northern Pacific, *Geophysical Research Letters*, 35, L22 815, doi:10.1029/2008GL035764, <http://doi.wiley.com/10.1029/2008GL035764>, 2008.
- Baumgardner, D., Popovicheva, O., Allan, J., Bernardoni, V., Cao, J., Cavalli, F., Cozic, J., Diapouli, E., Eleftheriadis, K., Genberg, P. J., Gonzalez, C., Gysel, M., John, A., Kirchstetter, T. W., Kuhlbusch, T. A. J., Laborde, M., Lack, D., Müller, T., Niessner, R., Petzold, A., Piazzalunga, A., Putaud, J. P., Schwarz, J., Sheridan, P., Subramanian, R., Swietlicki, E., Valli, G., Vecchi, R., and Viana, M.: Soot reference materials for instrument calibration and intercomparisons: a workshop summary with recommendations, *Atmospheric Measurement Techniques*, 5, 1869–1887, doi:10.5194/amt-5-1869-2012, <http://www.atmos-meas-tech.net/5/1869/2012/amt-5-1869-2012.html>, 2012.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Science and Technology*, 40, 27–67, doi:10.1080/02786820500421521, 2006.
- Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, *Journal of Geophysical Research*, 111, D20 211, doi:10.1029/2006JD007315, <http://doi.wiley.com/10.1029/2006JD007315>, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 1–173, doi:10.1002/jgrd.50171, <http://doi.wiley.com/10.1002/jgrd.50171>, 2013.
- Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davi-

- dovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer., *Mass spectrometry reviews*, 26, 185–222, doi:10.1002/mas.20115, <http://www.ncbi.nlm.nih.gov/pubmed/17230437>, 2007.
- Capes, G., Johnson, B., McFiggans, G., Williams, P. I., Haywood, J., and Coe, H.: Aging of biomass burning aerosols over West Africa: Aircraft measurements of chemical composition, microphysical properties, and emission ratios, *Journal of Geophysical Research*, 113, D00C15, doi:10.1029/2008JD009845, <http://doi.wiley.com/10.1029/2008JD009845>, 2008.
- Cappa, C. D., Lack, D. A., Burkholder, J. B., and Ravishankara, A. R.: Bias in filter-based aerosol light absorption measurements due to organic aerosol loading: Evidence from laboratory measurements, *Aerosol Science and Technology*, 42, 1022–1032, doi:10.1080/02786820802389285, 2008.
- Chung, C. E., Ramanathan, V., and Decremier, D.: Observationally constrained estimates of carbonaceous aerosol radiative forcing., *Proceedings of the National Academy of Sciences of the United States of America*, 109, 11624–9, doi:10.1073/pnas.1203707109, <http://www.pnas.org/content/109/29/11624>, 2012.
- Croft, B., Lohmann, U., Martin, R. V., Stier, P., Wurzler, S., Feichter, J., Posselt, R., and Ferrachat, S.: Aerosol size-dependent below-cloud scavenging by rain and snow in the ECHAM5-HAM, *Atmospheric Chemistry and Physics*, 9, 4653–4675, doi:10.5194/acp-9-4653-2009, <http://www.atmos-chem-phys.net/9/4653/2009/acp-9-4653-2009.html>, 2009.
- Croft, B., Lohmann, U., Martin, R. V., Stier, P., Wurzler, S., Feichter, J., Hoose, C., Heikkilä, U., van Donkelaar, A., and Ferrachat, S.: Influences of in-cloud aerosol scavenging parameterizations on aerosol concentrations and wet deposition in ECHAM5-HAM, *Atmospheric Chemistry and Physics*, 10, 1511–1543, doi:10.5194/acp-10-1511-2010, <http://www.atmos-chem-phys.net/10/1511/2010/acp-10-1511-2010.html>, 2010.
- Crosier, J., Allan, J. D., Coe, H., Bower, K. N., Formenti, P., and Williams, P. I.: Chemical composition of summertime aerosol in the Po Valley (Italy), northern Adriatic and Black Sea, *Quarterly Journal of the Royal Meteorological Society*, 133, 61–75, doi:10.1002/qj.88, <http://doi.wiley.com/10.1002/qj.88>, 2007.
- Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics, *Atmospheric Chemistry and Physics*, 11, 3303–3318, doi:10.5194/acp-11-3303-2011, <http://www.atmos-chem-phys.net/11/3303/2011/acp-11-3303-2011.html>, 2011.
- Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT\_4 modelling system for trajectories, dispersion and deposition, *Australian Meteorological Magazine*, 47, 295–308, 1998.
- Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S., and Worsnop, D. R.: A New Time-of-Flight Aerosol Mass Spectrometer (TOF-AMS)- Instrument Description and First Field Deployment, *Aerosol Science and Technology*, 39, 637–658, doi:10.1080/02786820500182040, <http://dx.doi.org/10.1080/02786820500182040>, 2005.
- Duplissy, J., DeCarlo, P. F., Dommen, J., Alfarra, M. R., Metzger, A., Barmapadimos, I., Prevot, A. S. H., Weingartner, E., Tritscher, T., Gysel, M., Aiken, A. C., Jimenez, J. L., Canagaratna, M. R., Worsnop, D. R., Collins, D. R., Tomlinson, J., and Baltensperger, U.: Relating hygroscopicity and composition of organic aerosol particulate matter, *Atmospheric Chemistry and Physics*, 11, 1155–1165, doi:10.5194/acp-11-1155-2011, <http://www.atmos-chem-phys.net/11/1155/2011/acp-11-1155-2011.html>, 2011.
- Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S., and Andreae, M. O.: Size matters more than chemistry for cloud-nucleating ability of aerosol particles., *Science*, 312, 1375–8, doi:10.1126/science.1125261, <http://www.sciencemag.org/content/312/5778/1375.abstract>, 2006.
- Foltescu, V., Selin, E., and Below, M.: Corrections for particle losses and sizing errors during aircraft aerosol sampling using a rosemount inlet and the PMS LAS-X, *Atmospheric Environment*, 29, 449–453, doi:10.1016/1352-2310(94)00258-M, [http://dx.doi.org/10.1016/1352-2310\(94\)00258-M](http://dx.doi.org/10.1016/1352-2310(94)00258-M), 1995.
- Franklin, J. E., Drummond, J. R., Griffin, D., Pierce, J. R., Waugh, D. L., Palmer, P. I., Parrington, M., Lee, J. D., Lewis, A. C., Rickard, A. R., Taylor, J. W., Allan, J. D., Coe, H., Walker, K. A., Chisholm, L., Duck, T. J., Hopper, J. T., Blanchard, Y., Gibson, M. D., Curry, K. R., Sakamoto, K. M., Lesins, G., Dan, L., Kliever, J., and Saha, A.: A case study of aerosol depletion in a biomass burning plume over Eastern Canada during the 2011 BORTAS field experiment, *Atmospheric Chemistry and Physics Discussions*, 14, 3395–3426, doi:10.5194/acpd-14-3395-2014, <http://www.atmos-chem-phys-discuss.net/14/3395/2014/acpd-14-3395-2014.html>, 2014.
- Gao, R. S., Schwarz, J. P., Kelly, K. K., Fahey, D. W., Watts, L. A., Thompson, T. L., Spackman, J. R., Slowik, J. G., Cross, E. S., Han, J.-H. H., Davidovits, P., Onasch, T. B., and Worsnop, D. R.: A Novel Method for Estimating Light-Scattering Properties of Soot Aerosols Using a Modified Single-Particle Soot Photometer, *Aerosol Science and Technology*, 41, 125–135, doi:10.1080/02786820601118398, <http://dx.doi.org/10.1080/02786820601118398>, 2007.
- Gerbig, C., Schmitgen, S., Kley, D., Volz-Thomas, A., Dewey, K., and Haaks, D.: An improved fast-response vacuum-UV resonance fluorescence CO instrument, *Journal of Geophysical Research*, 104, 1699, doi:10.1029/1998JD100031, <http://www.agu.org/pubs/crossref/1999/1998JD100031.shtml>, 1999.
- Ghan, S. J., Abdul-Razzak, H., Nenes, A., Ming, Y., Liu, X., Ovchinnikov, M., Shipway, B., Meskhidze, N., Xu, J., and Shi, X.: Droplet nucleation: Physically-based parameterizations and comparative evaluation, *Journal of Advances in Modeling Earth Systems*, 3, M10001, doi:10.1029/2011MS000074, <http://doi.wiley.com/10.1029/2011MS000074>, 2011.
- Giglio, L., Descloitres, J., Justice, C. O., and Kaufman, Y. J.: An Enhanced Contextual Fire Detection Algorithm for {MODIS}, *Remote Sensing of Environment*, 87, 273–282, doi:10.1016/S0034-4257(03)00184-6, <http://www.sciencedirect.com/science/article/pii/S0034425703001846>, 2003.
- Goldenson, N., Doherty, S. J., Bitz, C. M., Holland, M. M., Light, B., and Conley, A. J.: Arctic climate response to forcing from light-absorbing particles in snow and sea ice in CESM, *Atmospheric Chemistry and Physics*, 12, 7903–7920, doi:10.5194/acp-12-7903-2012, <http://www.atmos-chem-phys.net/12/7903/2012/acp-12-7903-2012.html>, 2012.

- Gysel, M., Laborde, M., Olfert, J. S., Subramanian, R., and Gröhn, A. J.: Effective density of Aquadag and fullerene soot black carbon reference materials used for SP2 calibration, *Atmospheric Measurement Techniques*, 4, 2851–2858, doi:10.5194/amt-4-2851-2011, <http://www.atmos-meas-tech.net/4/2851/2011/amt-4-2851-2011.html>, 2011.
- Haywood, J. M. and Shine, K. P.: The effect of anthropogenic sulfate and soot aerosol on the clear sky planetary radiation budget, *Geophysical Research Letters*, 22, 603–606, doi:10.1029/95GL00075, <http://doi.wiley.com/10.1029/95GL00075>, 1995.
- Hecobian, A., Zhang, X., Zheng, M., Frank, N., Edgerton, E. S., and Weber, R. J.: Water-Soluble Organic Aerosol material and the light-absorption characteristics of aqueous extracts measured over the Southeastern United States, *Atmospheric Chemistry and Physics*, 10, 5965–5977, doi:10.5194/acp-10-5965-2010, <http://www.atmos-chem-phys.net/10/5965/2010/acp-10-5965-2010.html>, 2010.
- Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmospheric Chemistry and Physics*, 12, 9817–9854, doi:10.5194/acp-12-9817-2012, <http://www.atmos-chem-phys.net/12/9817/2012/acp-12-9817-2012.html>, 2012.
- Jacobson, M. Z.: Development of mixed-phase clouds from multiple aerosol size distributions and the effect of the clouds on aerosol removal, *Journal of Geophysical Research*, 108, 4245, doi:10.1029/2002JD002691, <http://doi.wiley.com/10.1029/2002JD002691>, 2003.
- Khalizov, A. F., Zhang, R., Zhang, D., Xue, H., Pagels, J., and McMurry, P. H.: Formation of highly hygroscopic soot aerosols upon internal mixing with sulfuric acid vapor, *Journal of Geophysical Research*, 114, D05 208, doi:10.1029/2008JD010595, <http://doi.wiley.com/10.1029/2008JD010595>, 2009.
- Kirchstetter, T. W. and Thatcher, T. L.: Contribution of organic carbon to wood smoke particulate matter absorption of solar radiation, *Atmospheric Chemistry and Physics*, 12, 6067–6072, doi:10.5194/acp-12-6067-2012, <http://www.atmos-chem-phys.net/12/6067/2012/acp-12-6067-2012.html>, 2012.
- Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Bernsten, T., Bond, T. C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O., Easter, R., Fahey, D. W., Feichter, J., Fillmore, D., Freitag, S., Ghan, S., Ginoux, P., Gong, S., Horowitz, L., Iversen, T., Kirkevåg, A., Klimont, Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N., Myhre, G., Penner, J. E., Perlwitz, J., Pitari, G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G., Schwarz, J. P., Seland, O., Stier, P., Takegawa, N., Takemura, T., Textor, C., van Aardenne, J. A., and Zhao, Y.: Evaluation of black carbon estimations in global aerosol models, *Atmospheric Chemistry and Physics*, 9, 9001–9026, 2009.
- Koch, D., Balkanski, Y., Bauer, S. E., Easter, R. C., Ferrachat, S., Ghan, S. J., Hoose, C., Iversen, T., Kirkevåg, A., Kristjansson, J. E., Liu, X., Lohmann, U., Menon, S., Quaas, J., Schulz, M., Seland, O., Takemura, T., and Yan, N.: Soot microphysical effects on liquid clouds, a multi-model investigation, *Atmospheric Chemistry and Physics*, 11, 1051–1064, doi:10.5194/acp-11-1051-2011, <http://www.atmos-chem-phys.net/11/1051/2011/acp-11-1051-2011.html>, 2011.
- Koehler, K. A., DeMott, P. J., Kreidenweis, S. M., Popovicheva, O. B., Petters, M. D., Carrico, C. M., Kireeva, E. D., Khokhlova, T. D., and Shonija, N. K.: Cloud condensation nuclei and ice nucleation activity of hydrophobic and hydrophilic soot particles., *Physical chemistry chemical physics : PCCP*, 11, 7906–20, doi:10.1039/b905334b, <http://www.ncbi.nlm.nih.gov/pubmed/19727498>, 2009.
- Kondo, Y., Matsui, H., Moteki, N., Sahu, L., Takegawa, N., Kajino, M., Zhao, Y., Cubison, M. J., Jimenez, J. L., Vay, S., Diskin, G. S., Anderson, B., Wisthaler, A., Mikoviny, T., Fuelberg, H. E., Blake, D. R., Huey, G., Weinheimer, A. J., Knapp, D. J., and Brune, W. H.: Emissions of black carbon, organic, and inorganic aerosols from biomass burning in North America and Asia in 2008, *Journal of Geophysical Research*, 116, D08 204, doi:10.1029/2010JD015152, <http://doi.wiley.com/10.1029/2010JD015152>, 2011.
- Kreidenweis, S. M., Zhang, Y., and Taylor, G. R.: The effects of clouds on aerosol and chemical species production and distribution: 2. Chemistry model description and sensitivity analysis, *Journal of Geophysical Research*, 102, 23 867, doi:10.1029/97JD00775, <http://doi.wiley.com/10.1029/97JD00775>, 1997.
- Laborde, M., Mertes, P., Zieger, P., Dommen, J., Baltensperger, U., and Gysel, M.: Sensitivity of the Single Particle Soot Photometer to different black carbon types, *Atmospheric Measurement Techniques*, 5, 1031–1043, doi:10.5194/amt-5-1031-2012, <http://www.atmos-meas-tech.net/5/1031/2012/amt-5-1031-2012.html>, 2012a.
- Laborde, M., Schnaiter, M., Linke, C., Saathoff, H., Naumann, K.-H., Möhler, O., Berlenz, S., Wagner, U., Taylor, J. W., Liu, D., Flynn, M., Allan, J. D., Coe, H., Heimerl, K., Dahlkötter, F., Weinzierl, B., Wollny, A. G., Zannata, M., Cozic, J., Laj, P., Hitznerberger, R., Schwarz, J. P., and Gysel, M.: Single Particle Soot Photometer intercomparison at the AIDA chamber, *Atmospheric Measurement Techniques*, 5, 3077–3097, doi:10.5194/amt-5-3077-2012, <http://www.atmos-meas-tech.net/5/3077/2012/amt-5-3077-2012.html>, 2012b.
- Lack, D. A., Cappa, C. D., Covert, D. S., Baynard, T., Massoli, P., Sierau, B., Bates, T. S., Quinn, P. K., Lovejoy, E. R., and Ravishankara, A. R.: Bias in filter-based aerosol light absorption measurements due to organic aerosol loading: Evidence from ambient measurements, *Aerosol Science and Technology*, 42, 1033–1041, doi:10.1080/02786820802389277, 2008.
- Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, *Proceedings of the National Academy of Sciences*, 109, 14 802–14 807, doi:10.1073/pnas.1206575109, <http://www.pnas.org/content/early/2012/08/13/1206575109.abstract>, 2012.
- Latham, T. L., Beyersdorf, A. J., Thornhill, K. L., Winstead, E. L., Cubison, M. J., Hecobian, A., Jimenez, J. L., Weber, R. J., Anderson, B. E., and Nenes, A.: Analysis of CCN activity of Arctic aerosol and Canadian biomass burning during summer 2008, *Atmospheric Chemistry and Physics*, 13, 2735–2756, doi:10.5194/acp-13-2735-2013, <http://www.atmos-chem-phys.net/13/2735/2013/>, 2013.
- Le Breton, M., Bacak, A., Muller, J. B. A., O’Shea, S. J., Xiao, P., Ashfold, M. N. R., Cooke, M. C., Batt, R., Shallcross, D. E., Oram, D. E., Forster, G., Bauguitte, S. J.-

- B., and Percival, C. J.: Airborne hydrogen cyanide measurements using a chemical ionisation mass spectrometer for the plume identification of biomass burning forest fires, *Atmospheric Chemistry and Physics*, 13, 9217–9232, doi:10.5194/acp-13-9217-2013, <http://www.atmos-chem-phys.net/13/9217/2013/acp-13-9217-2013.html>, 2013.
- Liggio, J., Gordon, M., Smallwood, G., Li, S.-M., Stroud, C., Staebler, R., Lu, G., Lee, P., Taylor, B., and Brook, J. R.: Are emissions of black carbon from gasoline vehicles underestimated? Insights from near and on-road measurements., *Environmental science & technology*, 46, 4819–28, doi:10.1021/es2033845, <http://dx.doi.org/10.1021/es2033845>, 2012.
- Lin, B. and Rossow, W. B.: Precipitation water path and rainfall rate estimates over oceans using special sensor microwave imager and International Satellite Cloud Climatology Project data, *Journal of Geophysical Research*, 102, 9359, doi:10.1029/96JD03987, <http://doi.wiley.com/10.1029/96JD03987>, 1997.
- Lin, Y.-L., Farley, R. D., and Orville, H. D.: Bulk Parameterization of the Snow Field in a Cloud Model, *Journal of Climate and Applied Meteorology*, 22, 1065–1092, doi:10.1175/1520-0450(1983)022<1065:BPOTSF>2.0.CO;2, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0450\(1983\)022<1065:BPOTSF>2.0.CO;2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0450(1983)022<1065:BPOTSF>2.0.CO;2), 1983.
- Liu, D., Allan, J., Whitehead, J., Young, D., Flynn, M., Coe, H., McFiggans, G., Fleming, Z. L., and Bandy, B.: Ambient black carbon particle hygroscopic properties controlled by mixing state and composition, *Atmospheric Chemistry and Physics*, 13, 2015–2029, doi:10.5194/acp-13-2015-2013, <http://www.atmos-chem-phys.net/13/2015/2013/acp-13-2015-2013.html>, 2013.
- Liu, Y. and Daum, P. H.: Parameterization of the Auto-conversion Process. Part I: Analytical Formulation of the Kessler-Type Parameterizations, *Journal of the Atmospheric Sciences*, 61, 1539–1548, doi:10.1175/1520-0469(2004)061<1539:POTAPI>2.0.CO;2, [http://journals.ametsoc.org/doi/full/10.1175/1520-0469\(2004\)061<1539:POTAPI>2.0.CO;2](http://journals.ametsoc.org/doi/full/10.1175/1520-0469(2004)061<1539:POTAPI>2.0.CO;2), 2004.
- Lohmann, U., Stier, P., Hoose, C., Ferrachat, S., Kloster, S., Roeckner, E., and Zhang, J.: Cloud microphysics and aerosol indirect effects in the global climate model ECHAM5-HAM, *Atmospheric Chemistry and Physics*, 7, 3425–3446, doi:10.5194/acp-7-3425-2007, <http://www.atmos-chem-phys.net/7/3425/2007/acp-7-3425-2007.html>, 2007.
- Magi, B. I.: Effects of humidity on aerosols in southern Africa during the biomass burning season, *Journal of Geophysical Research*, 108, 8495, doi:10.1029/2002JD002144, <http://www.agu.org/pubs/crossref/2003/2002JD002144.shtml>, 2003.
- McMeeking, G. R., Kreidenweis, S. M., Baker, S., Carrico, C. M., Chow, J. C., Collett, J. L., Hao, W. M., Holden, A. S., Kirchstetter, T. W., Malm, W. C., Moosmüller, H., Sullivan, A. P., and Wold, C. E.: Emissions of trace gases and aerosols during the open combustion of biomass in the laboratory, *Journal of Geophysical Research*, 114, D19 210, doi:10.1029/2009JD011836, <http://doi.wiley.com/10.1029/2009JD011836>, 2009.
- McMeeking, G. R., Hamburger, T., Liu, D., Flynn, M., Morgan, W. T., Northway, M., Highwood, E. J., Krejci, R., Allan, J. D., Minikin, A., and Coe, H.: Black carbon measurements in the boundary layer over western and northern Europe, *Atmospheric Chemistry and Physics*, 10, 9393–9414, doi:10.5194/acp-10-9393-2010, <http://www.atmos-chem-phys.org/10/9393/2010/acp-10-9393-2010.html>, 2010.
- McNaughton, C. S., Clarke, A. D., Freitag, S., Kapustin, V. N., Kondo, Y., Moteki, N., Sahu, L., Takegawa, N., Schwarz, J. P., Spackman, J. R., Watts, L., Diskin, G., Podolske, J., Holloway, J. S., Wisthaler, A., Mikoviny, T., de Gouw, J., Warneke, C., Jimenez, J., Cubison, M., Howell, S. G., Middlebrook, A., Bahreini, R., Anderson, B. E., Winstead, E., Thornhill, K. L., Lack, D., Cozic, J., and Brock, C. A.: Absorbing aerosol in the troposphere of the Western Arctic during the 2008 ARCTAS/ARCPAC airborne field campaigns, *Atmospheric Chemistry and Physics*, 11, 7561–7582, doi:10.5194/acp-11-7561-2011, <http://www.atmos-chem-phys.net/11/7561/2011/acp-11-7561-2011.html>, 2011.
- Miller, N. and Wang, P.: A theoretical determination of the collection rates of aerosol particles by falling ice crystal plates and columns, *Atmospheric Environment. Part A. General Topics*, 25, 2593–2606, doi:10.1016/0960-1686(91)90177-9, [http://dx.doi.org/10.1016/0960-1686\(91\)90177-9](http://dx.doi.org/10.1016/0960-1686(91)90177-9), 1991.
- Morgan, W. T., Allan, J. D., Bower, K. N., Capes, G., Crosier, J., Williams, P. I., and Coe, H.: Vertical distribution of sub-micron aerosol chemical composition from North-Western Europe and the North-East Atlantic, *Atmospheric Chemistry and Physics*, 9, 5389–5401, doi:10.5194/acp-9-5389-2009, <http://www.atmos-chem-phys.net/9/5389/2009/acp-9-5389-2009.html>, 2009.
- Morgan, W. T., Allan, J. D., Bower, K. N., Highwood, E. J., Liu, D., McMeeking, G. R., Northway, M. J., Williams, P. I., Krejci, R., and Coe, H.: Airborne measurements of the spatial distribution of aerosol chemical composition across Europe and evolution of the organic fraction, *Atmospheric Chemistry and Physics*, 10, 4065–4083, doi:10.5194/acp-10-4065-2010, <http://www.atmos-chem-phys.net/10/4065/2010/acp-10-4065-2010.html>, 2010.
- Moteki, N. and Kondo, Y.: Effects of Mixing State on Black Carbon Measurements by Laser-Induced Incandescence, *Aerosol Science and Technology*, 41, 398–417, doi:10.1080/02786820701199728, <http://dx.doi.org/10.1080/02786820701199728>, 2007.
- Moteki, N., Kondo, Y., and Nakamura, S.: Method to measure refractive indices of small nonspherical particles: Application to black carbon particles, *Journal of Aerosol Science*, 41, 513–521, doi:10.1016/j.jaerosci.2010.02.013, 2010.
- Moteki, N., Kondo, Y., Oshima, N., Takegawa, N., Koike, M., Kita, K., Matsui, H., and Kajino, M.: Size dependence of wet removal of black carbon aerosols during transport from the boundary layer to the free troposphere, *Geophysical Research Letters*, 39, L13 802, doi:10.1029/2012GL052034, <http://www.agu.org/pubs/crossref/2012/2012GL052034.shtml>, 2012.
- Murphy, J. G., Oram, D. E., and Reeves, C. E.: Measurements of volatile organic compounds over West Africa, *Atmospheric Chemistry and Physics*, 10, 5281–5294, doi:10.5194/acp-10-5281-2010, <http://www.atmos-chem-phys.net/10/5281/2010/acp-10-5281-2010.html>, 2010.
- Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prevot, A. S. H., Di-

- nar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmospheric Chemistry and Physics*, 10, doi:10.5194/acp-10-4625-2010, 2010.
- 1210 O'Shea, S. J., Allen, G., Gallagher, M. W., Bauguitte, S. J.-B., Illingworth, S. M., Le Breton, M., Muller, J. B. A., Percival, C. J., Archibald, A. T., Oram, D. E., Parrington, M., Palmer, P. I., and Lewis, A. C.: Airborne observations of trace gases over boreal Canada during BORTAS: campaign climatology, air mass analysis and enhancement ratios, *Atmospheric Chemistry and Physics*, 13, 12451–12467, doi:10.5194/acp-13-12451-2013, <http://www.atmos-chem-phys.net/13/12451/2013/acp-13-12451-2013.html>, 2013.
- 1220 Oshima, N., Kondo, Y., Moteki, N., Takegawa, N., Koike, M., Kita, K., Matsui, H., Kajino, M., Nakamura, H., Jung, J. S., and Kim, Y. J.: Wet removal of black carbon in Asian outflow: Aerosol Radiative Forcing in East Asia (A-FORCE) aircraft campaign, *Journal of Geophysical Research*, 117, D03 204, doi:10.1029/2011JD016552, <http://www.agu.org/pubs/crossref/2012/2011JD016552.shtml>, 2012.
- 1225 Palmer, P. I., Parrington, M., Lee, J. D., Lewis, A. C., Rickard, A. R., Bernath, P. F., Duck, T. J., Waugh, D. L., Tarasick, D. W., Andrews, S., Aruffo, E., Bailey, L. J., Barrett, E., Bauguitte, S. J.-B., Curry, K. R., Di Carlo, P., Chisholm, L., Dan, L., Forster, G., Franklin, J. E., Gibson, M. D., Griffin, D., Helmig, D., Hopkins, J. R., Hopper, J. T., Jenkin, M. E., Kin-1230 dred, D., Kliever, J., Le Breton, M., Matthiesen, S., Maurice, M., Moller, S., Moore, D. P., Oram, D. E., O'Shea, S. J., Owen, R. C., Pagniello, C. M. L. S., Pawson, S., Percival, C. J., Pierce, J. R., Punjabi, S., Purvis, R. M., Remedios, J. J., Rotermund, K. M., Sakamoto, K. M., da Silva, A. M., Strawbridge, K. B., Strong, K., Taylor, J., Trigwell, R., Tereszchuk, K. A., Walker, K. A., Weaver, D., Whaley, C., and Young, J. C.: Quantifying the impact of BOREal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites (BORTAS) experiment: design, execution and science overview, *Atmospheric Chemistry and Physics*, 13, 6239–6261, doi:10.5194/acp-13-6239-2013, <http://www.atmos-chem-phys.net/13/6239/2013/acp-13-6239-2013.html>, 2013.
- 1245 Petters, M. D., Parsons, M. T., Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., Carrico, C. M., Sullivan, A. P., McMeeking, G. R., Levin, E., Wold, C. E., Collett, J. L., and Moosmüller, H.: Ice nuclei emissions from biomass burning, *Journal of Geophysical Research*, 114, D07 209, doi:10.1029/2008JD011532, <http://doi.wiley.com/10.1029/2008JD011532>, 2009.
- 1250 Pratt, K. a., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, a. J., Heymsfield, a. J., Seinfeld, J. H., and Prather, K. a.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, *Atmospheric Chemistry and Physics*, 11, 12549–12565, doi:10.5194/acp-11-12549-2011, <http://www.atmos-chem-phys.net/11/12549/2011/>, 2011.
- 1255 Ramanathan, V. and Carmichael, G.: Global and regional climate changes due to black carbon, *Nature Geoscience*, 1, 221–227, doi:10.1038/ngeo156, 2008.
- 1260 Reid, J. S., Eck, T. F., Christopher, S. A., Koppmann, R., Dubovik, O., Eleuterio, D. P., Holben, B. N., Reid, E. A., and Zhang, J.: A review of biomass burning emissions part III: intensive optical properties of biomass burning particles, *Atmospheric Chemistry and Physics*, 5, 827–849, doi:10.5194/acp-5-827-2005, <http://www.atmos-chem-phys.net/5/827/2005/acp-5-827-2005.html>, 2005a.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physics*, 5, 799–825, doi:10.5194/acp-5-799-2005, <http://www.atmos-chem-phys.net/5/799/2005/acp-5-799-2005.html>, 2005b.
- Sahu, L. K., Kondo, Y., Moteki, N., Takegawa, N., Zhao, Y., Cubison, M. J., Jimenez, J. L., Vay, S., Diskin, G. S., Wisthaler, A., Mikoviny, T., Huey, L. G., Weinheimer, A. J., and Knapp, D. J.: Emission characteristics of black carbon in anthropogenic and biomass burning plumes over California during ARCTAS-CARB 2008, *Journal of Geophysical Research*, 117, D16 302, doi:10.1029/2011JD017401, <http://www.agu.org/pubs/crossref/2012/2011JD017401.shtml>, 2012.
- Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Seland, O., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty, *Atmospheric Chemistry and Physics*, 13, 2423–2434, doi:10.5194/acp-13-2423-2013, <http://www.atmos-chem-phys.net/13/2423/2013/acp-13-2423-2013.html>, 2013.
- Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., Wilson, J. C., Reeves, J. M., Darbeheshti, M., Baumgardner, D. G., Kok, G. L., Chung, S. H., Schulz, M., Hendricks, J., Lauer, A., Kärcher, B., Slowik, J. G., Rosenlof, K. H., Thompson, T. L., Langford, A. O., Loewenstein, M., and Aikin, K. C.: Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, *Journal of Geophysical Research*, 111, D16 207, doi:10.1029/2006JD007076, <http://doi.wiley.com/10.1029/2006JD007076>, 2006.
- Schwarz, J. P., Gao, R. S., Spackman, J. R., Watts, L. A., Thomson, D. S., Fahey, D. W., Ryerson, T. B., Peischl, J., Holloway, J. S., Trainer, M., Frost, G. J., Baynard, T., Lack, D. A., de Gouw, J. A., Warneke, C., and Del Negro, L. A.: Measurement of the mixing state, mass, and optical size of individual black carbon particles in urban and biomass burning emissions, *Geophys. Res. Lett.*, 35, doi:10.1029/2008gl033968, <http://dx.doi.org/10.1029/2008GL033968>, 2008.
- Schwarz, J. P., Spackman, J. R., Gao, R. S., Perring, A. E., Cross, E., Onasch, T. B., Ahern, A., Wrobel, W., Davidovits, P., Olfert, J., Dubey, M. K., Mazzoleni, C., and Fahey, D. W.: The Detection Efficiency of the Single Particle Soot Photometer, *Aerosol Science and Technology*, 44, 612–628, doi:10.1080/02786826.2010.481298, <http://dx.doi.org/10.1080/02786826.2010.481298>, 2010a.
- Schwarz, J. P., Spackman, J. R., Gao, R. S., Watts, L. A., Stier, P., Schulz, M., Davis, S. M., Wofsy, S. C., and Fahey, D. W.: Global-scale black carbon profiles observed in the remote atmosphere and compared to models, *Geophysical Research Letters*, 37, L18 812, doi:10.1029/2010GL044372, <http://doi.wiley.com/10.1029/2010GL044372>, 2010b.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics - From Air Pollution to Climate Change* (2nd Edition), 1998.

- 1325 Slowik, J. G., Cross, E. S., Han, J. H., Davidovits, P., Onasch, T. B.,  
Jayne, J. T., Williams, L. R., Canagaratna, M. R., Worsnop, D. R.,  
Chakrabarty, R. K., Moosmuller, H., Arnott, W. P., Schwarz, J. P.,  
1385 Gao, R. S., Fahey, D. W., Kok, G. L., and Petzold, A.: An inter-  
comparison of instruments measuring black carbon content of  
1330 soot particles, *Aerosol Science and Technology*, 41, 295–314,  
doi:10.1080/02786820701197078, 2007.
- Stith, J. L., Twohy, C. H., DeMott, P. J., Baumgardner, D., Cam-  
1390 pos, T., Gao, R., and Anderson, J.: Observations of ice nu-  
clei and heterogeneous freezing in a Western Pacific extratrop-  
ical storm, *Atmospheric Chemistry and Physics*, 11, 6229–6243,  
1335 doi:10.5194/acp-11-6229-2011, <http://www.atmos-chem-phys.net/11/6229/2011/acp-11-6229-2011.html>, 2011.
- Stohl, A., Haimberger, L., Scheele, M. P., and Wernli, H.: An in-  
1395 tercomparison of results from three trajectory models, *Meteorolo-  
gical Applications*, 8, 127–135, <http://adsabs.harvard.edu/abs/2001MeApp...8..127S.2001>.
- 1340 Sullivan, A. P., Peltier, R. E., Brock, C. A., de Gouw, J. A.,  
Holloway, J. S., Warneke, C., Wollny, A. G., and We-  
ber, R. J.: Airborne measurements of carbonaceous aerosol  
soluble in water over northeastern United States: Method  
1345 development and an investigation into water-soluble or-  
ganic carbon sources, *Journal of Geophysical Research*, 111, D23S46,  
doi:10.1029/2006JD007072, <http://doi.wiley.com/10.1029/2006JD007072>, 2006.
- Taylor, J. W., Allan, J. D., Liu, D., Flynn, M., Hayes, P. L., Jimenez,  
1350 J. L., Lefer, B. L., Grossberg, N., Flynn, J., Gilman, J., de Gouw,  
J. A., and Coe, H.: Comparison of measured black carbon mass  
absorption coefficient to modelled values during CalNex 2010.,  
*Journal of Geophysical Research - Atmospheres*, in prep., 2014a.
- Taylor, J. W., Allan, J. D., Liu, D., Flynn, M., Weber, R.,  
1355 Zhang, X., Lefer, B. L., Grossberg, N., Flynn, J., and  
Coe, H.: Assessment of the sensitivity of core/shell pa-  
rameters derived using the single-particle soot photome-  
ter to density and refractive index, *Atmospheric Measure-  
ment Techniques Discussions*, 7, 5491–5532, doi:10.5194/amtd-  
1360 7-5491-2014, <http://www.atmos-meas-tech-discuss.net/7/5491/2014/amtd-7-5491-2014.html>, 2014b.
- 1420 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer,  
S., Bernsten, T., Berglen, T., Boucher, O., Chin, M., Dentener,  
F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Gi-  
1365 noux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang,  
P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A.,  
Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu,  
X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S.,  
Seland, O., Stier, P., Takemura, T., and Tie, X.: Analysis and  
1370 quantification of the diversities of aerosol life cycles within  
AeroCom, *Atmospheric Chemistry and Physics*, 6, 1777–1813,  
doi:10.5194/acp-6-1777-2006, <http://www.atmos-chem-phys.net/6/1777/2006/acp-6-1777-2006.html>, 2006.
- Trembath, J., Bart, M., and Brooke, J.: Efficiencies of Modi-  
1375 fied Rosemount Housings for sampling Aerosol on a Fast At-  
mospheric Research Aircraft., [http://www.faam.ac.uk/index.php/  
component/docman/doc\\_download/1673-inlet-efficiency](http://www.faam.ac.uk/index.php/component/docman/doc_download/1673-inlet-efficiency), 2012.
- Twohy, C. H., DeMott, P. J., Pratt, K. A., Subramanian, R.,  
1380 Kok, G. L., Murphy, S. M., Lersch, T., Heymsfield, A. J.,  
Wang, Z., Prather, K. A., and Seinfeld, J. H.: Relation-  
ships of Biomass-Burning Aerosols to Ice in Orographic  
Wave Clouds, *Journal of the Atmospheric Sciences*, 67, 2437–  
2450, doi:10.1175/2010JAS3310.1, [http://journals.ametsoc.org/  
doi/full/10.1175/2010JAS3310.1](http://journals.ametsoc.org/doi/full/10.1175/2010JAS3310.1), 2010.
- Val Martín, M., Honrath, R. E., Owen, R. C., Pfister, G., Fi-  
1385 alho, P., and Barata, F.: Significant enhancements of nitro-  
gen oxides, black carbon, and ozone in the North Atlantic  
lower free troposphere resulting from North American bo-  
real wildfires, *Journal of Geophysical Research*, 111, D23S60,  
doi:10.1029/2006JD007530, [http://www.agu.org/pubs/crossref/  
2006/2006JD007530.shtml](http://www.agu.org/pubs/crossref/2006/2006JD007530.shtml), 2006.
- Vignati, E., Karl, M., Krol, M., Wilson, J., Stier, P., and Cavalli, F.:  
Sources of uncertainties in modelling black carbon at the global  
1390 scale, *Atmospheric Chemistry and Physics*, 10, 2595–2611,  
doi:10.5194/acp-10-2595-2010, <http://www.atmos-chem-phys.net/10/2595/2010/acp-10-2595-2010.html>, 2010.
- Wang, X., Zhang, L., and Moran, M. D.: Uncertainty assessment of  
1395 current size-resolved parameterizations for below-cloud particle  
scavenging by rain, *Atmospheric Chemistry and Physics*,  
10, 5685–5705, doi:10.5194/acp-10-5685-2010, [http://www.  
atmos-chem-phys.net/10/5685/2010/acp-10-5685-2010.html](http://www.atmos-chem-phys.net/10/5685/2010/acp-10-5685-2010.html),  
2010.
- Ward, D. E. and Radke, L. F.: Emissions measurements from veg-  
1400 etation fires: A comparative evaluation of methods and results,  
John Wiley and Sons, Inc.; John Wiley and Sons Ltd., Ward, D.  
E. Intermountain Res. Station, Forest Serv., U.S. Dep. Agric.,  
P.O. Box 8089, Missoula, MT 59807, USA, 1993.
- Yokelson, R. J., Andreae, M. O., and Akagi, S. K.: Pitfalls with  
1405 the use of enhancement ratios or normalized excess mixing  
ratios measured in plumes to characterize pollution sources and  
aging, *Atmospheric Measurement Techniques*, 6, 2155–2158,  
doi:10.5194/amt-6-2155-2013, <http://www.atmos-meas-tech.net/6/2155/2013/amt-6-2155-2013.html>, 2013.
- Zhang, R., Khalizov, A. F., Pagels, J., Zhang, D., Xue, H.,  
1410 and McMurry, P. H.: Variability in morphology, hygroscopic-  
ity, and optical properties of soot aerosols during atmo-  
spheric processing., *Proceedings of the National Academy  
of Sciences of the United States of America*, 105, 10291–  
6, doi:10.1073/pnas.0804860105, [http://www.pnas.org/content/  
105/30/10291.full](http://www.pnas.org/content/105/30/10291.full), 2008.
- Zhou, J.: Hygroscopic Properties of Atmospheric Aerosol Particles  
1415 in Various Environments, Doctoral dissertation, Lund University,  
Lund, 2001.

**Table 1.** Ratios of BC/CO and OA/CO from this study and several previous measurements. The **emission** ratios from this study were calculated from Fig. 3, and listed errors are the fit errors.

Project/plume	Description	BC/CO ( $\mu\text{g sm}^{-3}/\text{ppmv}$ )	OA/CO ( $\mu\text{g sm}^{-3}/\text{ppmv}$ )	Reference
Plume 1	Canadian boreal forest	$3.28 \pm 0.06$	$174 \pm 5$	This study
Plume 2	Canadian boreal forest	$5.4 \pm 0.5$	$78 \pm 5$	This study
Plume 3	Canadian boreal forest	$0.61 \pm 0.04$	$0.4 \pm 0.3$	This study
ARCTAS	Asian boreal forest	$9 \pm 5$	$110^{\text{a}}$	Hecobian et al. (2010); Kondo et al. (2011)
ARCTAS	Canadian boreal forest	$2 \pm 2$	120	Hecobian et al. (2010); Kondo et al. (2011)
ARCTAS	Californian boreal forest	$3.4 \pm 1.4$	100	Hecobian et al. (2010); Kondo et al. (2011)
TexAQS	Unknown BB	$9 \pm 2$	-	Schwarz et al. (2008)
ICARTT non-precip.	N. American boreal forest	$2.5 - 8.4$	$39.4 - 125.6$	Sullivan et al. (2006); Val Martín et al. (2006)
ICARTT precip.	N. American boreal forest	$0.47 - 2.2$	-	Sullivan et al. (2006); Val Martín et al. (2006)
Prescribed burn	Large shrubs	18	82	Pratt et al. (2011)
Emission inventory	Extratropical forest	$6.5 \pm 3.2$	$150 - 170^{\text{b}}$	Andreae and Merlet (2001)

<sup>a</sup> Estimated from Siberian and mixed Siberian + Asian BB

<sup>b</sup> Assuming OM:OC ratio of 1.5 (Reid et al., 2005b).

**Table 2.** BC core size distribution properties for the three plumes, and a comparison of literature values. The ARCTAS values are averages of multiple fires sampled over North America.


	BC Core CMD (nm) <sup>a</sup>	BC Core MMD (nm) <sup>a</sup>	BC Core $\sigma_{geo}$	Reference
Plume 1	128	196	1.49	This study
Plume 2	137	194	1.38	This study
Plume 3	100	152	1.37	This study
ARCTAS Asian BB	146	214	1.37	Kondo et al. (2011)
ARCTAS Canadian BB	141	194	1.32	Kondo et al. (2011)
ARCTAS-CARB Californian BB	146	200	1.33	Sahu et al. (2012)
TexAQS biomass-burning BB	140	210	-	Schwarz et al. (2008)
CARB fossil fuel	119	175	1.36	Sahu et al. (2012)

<sup>a</sup> Literature values are corrected to BC core density of  $1.8\text{g cm}^{-3}$ .




**Table 3.** BC optical properties (at 532 nm) in the three plumes.  $BC/B_{Sca}$  are the slopes from Fig. 3c, calculated using ODR fits with zero offset. The listed errors are the fit errors. MAC was calculated using the measured BC core size and coating distributions, integrated over the entire BC size distribution, assuming either a constant distribution of shell/core ratio or absolute coating thickness from Fig. 6. The stated values were calculated using BC core and coating refractive indices  $n_{core} = (1.85 - 0.71i)$  and  $n_{coat} = (1.50 - 0i)$ .  $B_{abs}/CO$ , and errors are the standard deviation of the corresponding range in Table S2. SSA were calculated by combining the calculated MAC with measured  $BC/CO$  and  $BC/B_{Sca}$ .


	Calculated MAC ( $m^2 g^{-1}$ )	$B_{abs}/CO$ ( $Mm^{-1}/ppmv$ )	$BC/B_{Sca}$ ( $\mu g sm^{-3}/Mm^{-1}$ )	SSA
Using shell/core ratio distribution				
Plume 1	$10.64 \pm 0.64$	$34.9 \pm 0.6$	$2.38 \pm 0.03$	0.975
Plume 2	$10.95 \pm 0.63$	$59 \pm 5$	$5.55 \pm 0.09$	0.943
Plume 3	$10.24 \pm 0.68$	$6.2 \pm 0.4$	$4.36 \pm 0.15$	0.957
Using coating thickness distribution				
Plume 1	$9.10 \pm 0.53$	$29.8 \pm 1.8$	$2.38 \pm 0.03$	0.979
Plume 2	$9.27 \pm 0.55$	$50 \pm 5$	$5.55 \pm 0.09$	0.951
Plume 3	$8.79 \pm 0.55$	$5.4 \pm 0.5$	$4.36 \pm 0.15$	0.963



**Figure 1.** 5-day HYSPLIT back trajectories, starting at the time and location sampled (the square markers) and with triangular markers every 6 hours. Parts (a) and (b) show the horizontal and vertical tracks respectively. The aqua-coloured parts of the trajectories are regions with HYSPLIT precipitation, and the yellow dots show MODIS fire data between 18–20 July 2011. The grey line shows the flight track, starting from light-grey and ending in dark-grey.



**Figure 2.** Cloud data from the GOES satellite from 19 July 2011. The retrievals are from the the periods when the meteorology data in the back trajectory for plume 3 showed precipitation. Parts (a) and (b) show effective cloud height, (c) and (d) show cloud depth (calculated as the difference between effective cloud height and cloud bottom) and parts (e) and (f) show ice water path. The back trajectories are the same as those shown in Fig. 1, and the circles show the locations of the back trajectories at the time of the GOES retrievals.



**Figure 3.** Correlations of (a) OA/CO, (b) BC/CO and (c) BC/ $B_{\text{Sca}}$  in the three plumes. The slopes (which were calculated by orthogonal distance regression) are listed in Table 1 for (a) and (b) and Table 3 for (c). The peak OA concentration in Plume 3 was  $1.4 \mu\text{g sm}^{-3}$ .







Figure 4. Average normalised bulk size distributions measured by the SMPS.



**Figure 5.** Normalised average BC core size distributions measured by the SP2 during for the three plumes. Part (a) presents the number distribution, and part (b) the mass distribution. The discontinuity at  $\sim 300$  nm is due to saturation of the broadband detector and switching to the narrowband. The grey shaded regions show the range of core diameter used for calculating coating properties.



**Figure 6.** Distributions of measured coating thicknesses, measured by the SP2, expressed in terms of (a) shell/core ratio and (b) absolute coating thickness. The first bin ( $D_P = D_C = 1$ , uncoated particle) also contains all particles with measured scattering less than that predicted for an uncoated core (equivalent to  $D_P < D_C$ ). Such particles are a result of random variations in the detected scattering, however as the result  $D_P < D_C$  is nonphysical they are assigned the nearest realistic value.



**Figure 7.** BC core size distributions measured by the SP2 and calculated size-dependent MAC for the three plumes, calculated using  $n_{core} = (1.85 - 0.71i)$  and assuming a constant distribution of shell/core ratio.