

Modeled black carbon radiative forcing and atmospheric lifetime in AeroCom Phase II constrained by aircraft observations

B. H. Samset, Center for International Climate and Environmental Research – Oslo (CICERO), Oslo,
Norway

G. Myhre, Center for International Climate and Environmental Research – Oslo (CICERO), Oslo,
Norway

A. Herber, Alfred Wegener Institute for Polar and Marine Research in the Helmholtz Association,
Burgermeister-Smidt-Straße 20, 27568 Bremerhaven, Germany

Y. Kondo, Department of Earth and Planetary Science, Graduate School of Science, University of
Tokyo, Tokyo, Japan

S.-M. Li, Air Quality Research Division, Science and Technology Branch, Environment Canada, 4905
Dufferin Street, Toronto, Ontario, M3H 5T4, Canada

N. Moteki, Department of Earth and Planetary Science, Graduate School of Science, University of
Tokyo, Tokyo, Japan

M. Koike, Department of Earth and Planetary Science, Graduate School of Science, University of
Tokyo, Tokyo, Japan

N. Oshima, Meteorological Research Institute, Tsukuba, Ibaraki Japan

J. P. Schwarz, Chemical Sciences Division, Earth System Research Laboratory, National Oceanic
and Atmospheric Administration, Boulder, Colorado, USA and Cooperative Institute for Research in
Environmental Sciences, University of Colorado Boulder, Boulder, Colorado, USA

Y. Balkanski, Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, Gif-
sur-Yvette, France

S. E. Bauer, NASA Goddard Institute for Space Studies and Columbia Earth Institute, New York, NY,
USA

N. Bellouin, Met Office Hadley Centre, Exeter, UK. Now at Department of Meteorology, University
of Reading, Reading, UK

27 **T. K. Berntsen**, Center for International Climate and Environmental Research – Oslo (CICERO),
28 Oslo, Norway

29 **H. Bian**, Joint Center for Earth Systems Technology, University of Maryland Baltimore County, MD,
30 USA

31 **M. Chin**, NASA Goddard Space Flight Center, Greenbelt, MD, USA

32 **T. Diehl**, NASA Goddard Space Flight Center, Greenbelt, MD, USA *and* Universities Space Research
33 Association, Columbia, MD, USA (now at European Commission at the Joint Research Center, Ispra,
34 Italy)

35 **R. C. Easter**, Pacific Northwest National Laboratory, Richland, WA, USA

36 **S. J. Ghan**, Pacific Northwest National Laboratory, Richland, WA, USA

37 **T. Iversen**, Norwegian Meteorological Institute, Oslo, Norway *and* Department of Geosciences,
38 University of Oslo, Oslo, Norway *and* ECMWF, Shinfield Park, RG2 9AX, Reading, UK

39 **A. Kirkevåg**, Norwegian Meteorological Institute, Oslo, Norway

40 **J.-F. Lamarque**, NCAR Earth System Laboratory, National Center for Atmospheric Research,
41 Boulder, CO, USA

42 **G. Lin**, Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann
43 Arbor, Michigan, USA

44 **X. Liu**, Department of Atmospheric Science, University of Wyoming, Laramie, WY, USA

45 **J. E. Penner**, Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann
46 Arbor, Michigan, USA

47 **M. Schulz**, Norwegian Meteorological Institute, Oslo, Norway

48 **Ø. Seland**, Norwegian Meteorological Institute, Oslo, Norway

49 **R. B. Skeie**, Center for International Climate and Environmental Research – Oslo (CICERO), Oslo,
50 Norway

51 **P. Stier**, Department of Physics, University of Oxford, Oxford, UK

52 **T. Takemura**, Research Institute for Applied Mechanics, Kyushu University, Fukuoka, Japan

53 **K. Tsigaridis**, NASA Goddard Institute for Space Studies and Columbia Earth Institute, New York,
54 NY, USA

Abstract

Atmospheric black carbon (BC) absorbs solar radiation, and exacerbates global warming through exerting positive radiative forcing (RF). However, the contribution of BC to ongoing changes in global climate is under debate. Anthropogenic BC emissions, and the resulting distribution of BC concentration, are highly uncertain. In particular, long range transport and processes affecting BC atmospheric lifetime are poorly understood. Here we discuss whether recent assessments may have overestimated present day BC radiative forcing in remote regions. We compare vertical profiles of BC concentration from four recent aircraft measurement campaigns to simulations by 13 aerosol models participating in the AeroCom Phase II intercomparison. An atmospheric lifetime of BC of less than 5 days is shown to be essential for reproducing observations in remote ocean regions, in line with other recent studies. Adjusting model results to measurements in remote regions, and at high altitudes, leads to a 25% reduction in AeroCom Phase II median direct BC forcing, from fossil fuel and biofuel burning, over the industrial era. The sensitivity of modeled forcing to BC vertical profile and lifetime highlights an urgent need for further flight campaigns, close to sources and in remote regions, to provide improved quantification of BC effects for use in climate policy.

Introduction

As an absorber of solar radiation, anthropogenic BC emissions can contribute positively to global radiative forcing through the aerosol direct effect, they can affect clouds through the aerosol indirect and semidirect effects, change albedo of snow and ice, and influence precipitation by changing atmospheric stability and the surface energy balance (Myhre et al.,

2013a;Ramanathan and Carmichael, 2008;Haywood and Shine, 1995). Presently both the magnitude of anthropogenic BC emissions and the resulting global distribution of BC concentrations are highly uncertain. In particular the vertical profile of BC concentration, which strongly affects its total impact on the energy balance of the atmosphere, is poorly constrained (Koffi et al., 2012;Textor et al., 2007;Samset et al., 2013). Comparisons of measurements with model results, with emphasis both on total BC mass, spatio-temporal distribution and vertical structure are therefore essential for constraining estimates of BC effects on climate.

The IPCC AR5 (Boucher et al., 2013) assessed the direct aerosol effect radiative forcing due to anthropogenic BC (defined here as BC from anthropogenic fossil fuel and biofuel sources, BC FF+BF) to be +0.40 [range: +0.05 to +0.80] W m⁻² over the period 1750-2010. That assessment took into account both model based and observational studies. Recently, Phase II of the AeroCom model intercomparison project (Myhre et al., 2013b) also evaluated BC FF+BF radiative forcing (RF), based purely on 15 global aerosol models (Myhre et al., 2013b), and found it to be +0.23 [+0.06 to +0.48] Wm⁻². The uncertainty ranges (5-95%) show that BC is still a major contributor to the total uncertainty on anthropogenic radiative forcing. Further, it has recently been shown that, because the direct radiative forcing per unit mass BC increases strongly with altitude (Ban-Weiss et al., 2011;Zarzycki and Bond, 2010;Samset and Myhre, 2011), the diversity in modeled vertical profiles of BC concentration in the AeroCom Phase II models may account for up to 50% of the model diversity in anthropogenic BC RF (Samset et al., 2013). Climate model simulations, however, indicate that, while direct BC forcing strengthens with altitude, its climate efficacy may decrease, i.e. the surface temperature response to BC in the middle and upper troposphere may be small or even negative (Ban-Weiss et al., 2011;Flanner, 2013). Further, (Samset et al., 2014) concluded that an upward adjustment on the model based uncertainty on total aerosol forcing

may be necessary, in an analysis that looked at multi-model variability in per-species aerosol burdens and optical parameters. BC forcing diversity was found to be a significant component in this analysis.

Recently, several single model studies have investigated other factors that may underlie the intermodel variability in AeroCom Phase II, or assessed its multi-model mean results in light of observations. E.g. (Wang et al., 2014a) compared results from the GEOS-Chem model (<http://geos-chem.org>) with results from the HIPPO flight campaign, and concluded that to reproduce HIPPO, more wet removal was required than is represented in most present models. Based on this, and on revised estimates on direct radiative forcing of BC when taking HIPPO constraints (Schwarz et al., 2013) into account, they argue that previous model estimates may be biased high due to elevated BC concentrations in the free troposphere. Bauer et al. (2013) studied the atmospheric lifetime of BC, which is a combined measure of transport and removal processes, by comparing simulations with the GISS-MATRIX model to HIPPO (Bauer et al., 2010), using CMIP5 emissions. They found that under present day conditions, BC lifetime should be no more than 4 days, which is significantly shorter than what is used in some present models. Hodnebrog et al. (2014) showed that such a reduction in lifetime, when combined with estimates of the impact from BC on atmospheric stability, can lead to major reductions in the global mean climate impact of BC emissions. Based on the above, it is natural to investigate both whether the full AeroCom Phase II RF estimate is biased high, as was found for GEOS-Chem, and what contribution to multimodel variability may be due to differences in modeled BC lifetime.

In the following, we compare vertical concentration profiles from the AeroCom Phase II models to recent aircraft campaigns. The results are used to set multi-model constraints on BC lifetime, and to find limits on the possible high bias of AeroCom Phase II BC RF if constraining to HIPPO results.

Methods

Flight data and BC definition

We have used data from flight campaigns HIAPER Pole-to-Pole Observations (HIPPO) 1-5 (Schwarz et al., 2013), Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) SP2 (Jacob et al., 2010), Polar Airborne Measurements and Arctic Regional Climate Model Simulation Project (PAMARCMiP) (Herber et al., 2012; Stone et al., 2010) and Aerosol Radiative Forcing in East Asia (A-FORCE) (Oshima et al., 2012). (See Table 1)

All flights measured BC concentrations using the single particle soot photometer (SP2) instrument (Schwarz et al., 2010). Hence, in the present work, “BC” in relation to measured data stands for “refractive BC (rBC)” as quantified by SP2, equivalent to properly measured elemental carbon (Kondo et al., 2011).

HIPPO 1-5 (Schwarz et al., 2013) flew mainly pole-to-pole over the Pacific Ocean at various times during 2009-2011. Combining data from all five campaigns yields an approximate annual average. The HIPPO data have been screened against contributions from fires, so as to be representative of the background concentration of BC over the Pacific. Two of the HIPPO campaigns also flew over the North American mainland, allowing also for comparisons closer to anthropogenic BC source regions. HIPPO covers atmospheric pressures from surface values up to 100hPa, meaning that its upper range reaches into the lower stratosphere.

The ARCTAS SP2 campaign (Jacob et al., 2010) was flown in two separate time periods in 2008. During spring, flights were conducted over the northern Pacific Ocean, comparable to parts of the HIPPO region, and also over the North Polar regions. During summer, flights were conducted over the North American continent, again comparable to HIPPO. Part of the ARCTAS motivation was to study fires, so the measured concentrations can be expected to

have a larger contribution from open biomass burning than the HIPPO dataset. ARCTAS data cover the atmospheric segment from the surface up to 250hPa.

The Polar Airborne Measurements and Arctic Regional Climate Model Simulation Project (PAMARCMiP) campaign (Herber et al., 2012; Stone et al., 2010) consisted of a series of flights conducted over the North Polar and Northern Pacific regions, from 2009 through 2012. It covers a vertical range up to 500hPa, and is partially comparable to both ARCTAS and HIPPO. As ARCTAS, PAMARCMiP is partially affected by biomass burning emissions.

Finally, the Aerosol Radiative Forcing in East Asia (A-FORCE) aircraft campaign (Oshima et al., 2012) flew over East Asia (southwestern Japan) in spring 2009, covering a vertical range up to 300hPa. It is not regionally comparable to the other campaigns, but is highly relevant because it represents a region dominated by outflow from mainland China, one of the main sources of anthropogenic BC emissions.

AeroCom Phase II models

The models results in the present study come from 13 state of the art aerosol models, submitted as part of Phase II of the AeroCom model intercomparison project (Myhre et al., 2013b; Samset et al., 2013; Schwarz et al., 2013). Participating models are NCAR-CAM3.5 (Lamarque et al., 2012), CAM4-Oslo (Kirkevåg et al., 2013), CAM5.1 (Liu et al., 2012), GISS-MATRIX (Bauer et al., 2010), GISS modelE (Koch et al., 2011), GMI (Bian et al., 2009), GOCART-v4 (Chin et al., 2009), HadGEM2 (Bellouin et al., 2011), IMPACT (Lin et al., 2012), INCA (Szopa et al., 2012), ECHAM5-HAM (Zhang et al., 2012), OsloCTM2 (Skeie et al., 2011) and SPRINTARS (Takemura et al., 2005). See (Myhre et al., 2013b) and individual model references for further descriptions.

All models submitted monthly mean 3D fields of total BC mass mixing ratios, using year 2000 emissions (Lamarque et al., 2010). Meteorological year was 2006, or model internal

present day (PD) climatology. To calculate BC concentrations from mixing ratios, the models' own monthly mean temperature and pressure fields were used. When discussing modifications to the BC radiative forcing from the direct effect from anthropogenic fossil fuels and biofuels, the models' own monthly mean 2D forcing fields were used, in combination with a preindustrial simulation using year 1850 emissions but still year 2006 or PD meteorology (Myhre et al., 2013b). For consistency with recent literature, forcing results are given for 1750-2010, using scaling factors presented in (Myhre et al., 2013b).

Analysis

To compare flight campaigns and model output, a series of geographical regions were first selected. See Figure 1. For the flight data, only measurements that fell within the regions were kept, and for each region an average profile was constructed. For the models, all output within the selected region was averaged into a single profile for each model. However, to take the seasonality into account, we produced a model profile for each measurement point or profile from the flights. These were then averaged. The result is a set of model profiles that correspond to the flight profiles both geographically and temporally.

From the concentration profiles, we calculated aerosol burdens for both models and flight data. To ensure comparability, model burdens were calculated only in the same vertical range covered by the flight campaign.

Further, we used the methodology presented in Samset et al. (2013) to calculate BC RF from the concentration profiles. Briefly, we use spatially and temporally resolved normalized forcing efficiency profiles (RF exerted per gram of aerosol at a given altitude) calculated from a single model (Samset and Myhre, 2011), and multiplied with profiles of BC burden per model layer. This yields comparable estimates for total BC forcing within the selected regions, for the seasons covered by the respective flight campaigns; thus all RF calculations are

performed with a consistent method. To distinguish the models' own estimates of RF from the RF calculated by this method, we refer to the two as “native RF” and “recalculated RF” respectively. While using the forcing efficiency from a single model (OsloCTM2) will naturally bias the calculated RF towards the forcing strength predicted by that model, it also allows for an estimate of differences in vertical profile shape. Since the forcing efficiency for BC is strongly and monotonically rising with altitude, differences between the overall shape of measured and modeled profiles will cause the ratio of recalculated forcing per burden to differ from unity.

Calculation of RMS values and correlations

To estimate how well a given model reproduces the Pacific HIPPO flight data, as primarily used below, we calculated model bias, root-mean-square (RMS) error values and correlation coefficients. The HIPPO dataset was subdivided into five regions (P1-P5 in Figure 1), and an annual mean profile was constructed for each region as shown in Figure 2. For each model, diagnostics were calculated from the difference between the HIPPO concentration profile at each of its given altitude levels, and the regionally averaged model concentration value interpolated to the corresponding altitude, according to the following equations:

$$Mean\ bias = \frac{1}{N} \sum_{Reg=P1}^{P5} \sum_{Alt=Surf}^{TOA} [C_{HIPPO}(Reg, Alt) - C_{Model}(Reg, Alt)]$$

$$Normalized\ mean\ bias = \frac{1}{N} \sum_{Reg=P1}^{P5} \sum_{Alt=Surf}^{TOA} \frac{[C_{HIPPO}(Reg, Alt) - C_{Model}(Reg, Alt)]}{0.5 \times [C_{HIPPO}(Reg, Alt) + C_{Model}(Reg, Alt)]}$$

$$RMS\ Error = \sqrt{\frac{1}{N} \sum_{Reg=P1}^{P5} \sum_{Alt=Surf}^{TOA} [C_{HIPPO}(Reg, Alt) - C_{Model}(Reg, Alt)]^2}$$

Here C denotes a concentration value and N is the total number of data points. In the present case, this value is 72, determined by the number of altitude bins where HIPPO reported measurements. Further, we calculated the Pearson sample correlation coefficient based on the same dataset.

Derivation of scaled forcing estimates

Two scalings were applied in the present work to assess the potential impact of adjusting models to measured BC concentrations. These scalings were derived by altering the 3D concentration fields of total BC (fossil fuel, biofuel and biomass burning) provided by the AeroCom models, and then applied to the BC FF+BF forcing fields supplied. This method is used to ensure that intermodel variability in RF due to differences in optical parameters of BC, cloud distributions and other factors related to the host model are kept unchanged.

For the “remote ocean” scaling, the concentration fields were altered within the grey boxes shown in Figure 1. Between the surface and 500hPa concentrations were reduced to 1/3, then to 1/8 up to 200hPa, and then to 1/15 up to TOA. These factors were derived from the comparison between AeroCom Phase II and HIPPO 1-5 presented in Schwarz 2013 (Schwarz et al., 2013).

Using the forcing efficiency profile method (Samset and Myhre, 2011), we then calculated global, annual mean BC RF from both scaled and unscaled concentration fields. The ratio of these forcing values is taken as the scaling factor for that particular model.

Finally, we constructed the multi-model median BC FF+BF for all 13 models used for the present study, based on their original 2D forcing fields. These forcing values were scaled with the derived scaling factors, to produce the revised model median forcing.

For the “high altitude” scaling the same procedure was followed, except that the concentration fields were scaled to 1/20 at altitudes between 200hPa and TOA globally.

For the “all scaled” analysis both scalings were applied to the concentration fields, i.e. the fields were scaled to 1/20 at altitudes between 200hPa and TOA globally, and then as described above in the grey marked regions at altitudes below 200hPa.

Results and discussion

Comparisons of flights and models

Here, we constrain the model range of global, annual mean direct radiative forcing by anthropogenic BC, by comparing AeroCom Phase II vertical profiles from 13 models, to recent aircraft campaigns. Figure 1 shows the flight tracks of the four campaigns, the AeroCom multimodel median anthropogenic BC forcing field, and the regions selected for analysis.

Table 2 shows individual model BC RF, recalculated using the forcing efficiency profile method, globally and for the regions in Figure 1. We also show the fraction of exerted above 5km (500hPa),

Figure 2 compares flight campaign data with AeroCom Phase II model output. Panels a-f show the HIPPO1-5 campaigns (Schwarz et al., 2013) for five regions in the remote Pacific Ocean and for western North America, overlain with AeroCom Phase II results. A common pattern is that the models strongly overpredict the HIPPO measurements. Further, the overprediction is more pronounced at high altitudes. Comprised of five campaigns distributed throughout the year, HIPPO represents an approximate annual average. As recently noted (Schwarz et al., 2013), its Pacific measurements indicate that at the highest altitudes studied, BC concentrations converge towards a common background value, here found to be

approximately 0.1 ng m^{-3} , with very low seasonality. Here we also find (Figure 2f) the same background value above western North America.

Panels g-i of Figure 2 show the ARCTAS (Jacob et al., 2010) campaign, which reports significantly higher concentrations than HIPPO. The models mainly underpredict these observations, linked to the fact that ARCTAS encountered biomass burning BC from episodic forest fires (Wang et al., 2011), which HIPPO did not encounter in this region. A notable feature is that above the fire dominated segments, the ARCTAS profiles show a strong decline with altitude. In the P1 (Northern Pacific) region upper tropospheric ARCTAS concentrations are similar to those measured in HIPPO.

Panels j-l show PAMARCMiP (Herber et al., 2012; Stone et al., 2010) data, first over the North Pacific region, and then over two North Polar regions. While the altitude range covered by PAMARCMiP is limited compared to ARCTAS and HIPPO, the concentrations found in the lowest few kilometers of the troposphere are consistent with ARCTAS. Over the NP1 and NP2 region, north of America and Greenland, models underpredict the measurements. As for ARCTAS, this is at least partly due to episodic fires. The Arctic region may however have further sources of BC not adequately represented in the emission inventories used by the models (Stohl et al., 2013).

Panel m shows A-FORCE (Oshima et al., 2012) data in the sea areas around Japan. Here we find good agreement between models and measured concentrations, both in absolute values and in the shape of the vertical profile. The variability between models is also much lower in this region than for the others. The aerosol in the A-FORCE region is mainly sensitive to outflow from mainland China, Korea and Japan. This indicates that in AeroCom Phase II, East Asian BC emissions and outflow are either well represented or, if emissions are still underestimated as discussed for AeroCom Phase I in recent literature (Bond et al.,

2013;Chung et al., 2012), the atmospheric lifetime of BC must be compensatingly long in the models to allow enough BC to be transported into the region sampled by A-FORCE. We note, however, that the A-FORCE data do not extend as far up in the atmosphere as HIPPO did, and that we find significant intermodel variability at $p < 400$ hPa also for the near-source A-FORCE and HIPPO America regions.

While the aircraft data in the present study were taken over the period 2008-2012, the models used emissions from year 2000. BC emissions have increased in the intervening period (Wang et al., 2014b), indicating that any overestimation of concentrations by the models would have been strengthened had they used a more recent emission inventory. One model (CAM4-Oslo) delivered results for both year 2000 and 2006 emissions, reflecting this increase. In remote regions (e.g. the HIPPO regions in Figure 1), the resulting increase in concentration is found to be evenly distributed throughout the vertical profile, except in the range 1000-800hPa where no significant increase was found. It is clear that for future comparisons, model calculations with updated emission inventories are desirable.

Consequences for BC atmospheric lifetime

In the following, we assess the implications of the flight observations on modeled BC lifetime and RF. Episodic biomass burning emissions from fires, even though represented in the model emissions, pose challenges when comparing flight campaigns to monthly mean model data. Arguably fires are also difficult to characterize as anthropogenic. Below, we therefore constrain our discussion to the HIPPO dataset, which was less influenced by episodic fires, reached the highest altitudes, covers the largest geographical area, and represents an approximate annual mean.

Figure 2 shows that some models more closely reproduce the measurements than others, both in magnitude and shape. Several studies have suggested that to reproduce HIPPO data, a low

modeled atmospheric lifetime, or a short ageing timescale, of BC is required (Bauer et al., 2013; Wang et al., 2014a). Here we can test this supposition for a larger set of models. Quantifying the difference between models and data requires care, as absolute concentrations range over several orders of magnitude. Common diagnostic variables include model bias and RMS error. Of these, RMS error and model mean bias (see Methods) will be dominated by high absolute concentrations, i.e. low altitudes in the present case. Model mean normalized bias avoids this, but will be more sensitive to model and measurement uncertainties in high altitude, low concentration ranges. In Figure 3, RMS error and biases are plotted as function of the modeled BC lifetime. Lifetime, also referred to as atmospheric residence time, is here defined as modeled global, annual mean emissions divided by burden (Table 3).

Figure 3 shows that, independent of diagnostic variable, a low BC lifetime is a requirement for good reproduction of absolute modeled concentrations. Regressing bias or RMS error versus lifetime (black, dashed line in Figure 3) gives an intercept at 3 days for RMS error and model mean bias. This value is in line with indications from other recent studies, e.g. (Bauer et al., 2013). In the present dataset, a single model with high lifetime (HadGEM2) represents a significant outlier. That model did not include BC ageing and transition to a hydrophilic state, with the consequence that both BC lifetime and burdens over remote areas become high (Bellouin et al., 2011). To test the impacts of single models on the result, Figure 3 also shows regressions with one model removed (grey lines). For the normalized mean bias, which is less sensitive to high concentrations, the regression with this particular model removed is consistent with the results from RMS error and mean bias.

Bias and RMS error give information on absolute deviations, but less on any covariance in shape. The Pearson correlation coefficient, however, is sensitive to the shape of the BC profiles. In Figure 3, correlation is indicated by symbol size. (See also values in Table 3.) Several models with low lifetimes also yield low correlations. Regressing only the models

with correlation coefficients $\rho > 0.8$ gives similar slopes and intercepts to what we find using all models (red, dashed line). We note that 12 out of 13 models show correlation with the Pacific HIPPO data at significance $p > 0.05$.

Low BC lifetime appears necessary, but not sufficient, to describe the data. Only three models (IMPACT, GMI, GISS-MATRIX) exhibit both a low bias or RMS error and a high correlation, with no single obvious factor linking their aerosol treatments. AeroCom Phase II models use a wide variety of microphysics schemes (Mann et al., 2013). Meteorology and treatment of BC aging and wet scavenging also vary, and will impact the vertical profiles (Kipling et al., 2013; Bauer et al., 2013). Further model experiments, in line with the single model study in (Wang et al., 2014a), are required to address the reasons behind the relationship found in Figure 3.

Consequences for modeled BC FF+FB RF

However, the three models that best reproduce HIPPO in the Pacific all report consistent and relatively low BC FF+BF forcing, exerted close to emission sources as expected from the low lifetime. In these three models very little BC reaches remote ocean regions, or gets lifted above 500hPa, relative to the other models in the ensemble. While the correspondence to HIPPO cannot be used to extract information close to emission sources, it does suggest that scaling down the average modeled forcing aloft and in remote ocean regions has merit.

Compared with HIPPO, the current model ensemble overestimates BC concentrations at all altitudes in remote regions. The overestimation increases with altitude, and is particularly significant at pressures below 200hPa. Further, (Schwarz et al., 2013) suggest that the minimum concentration consistently observed by HIPPO in the upper troposphere, lower stratosphere and tropical transition layer may be a global feature. Interestingly, the models

shown here do, on average, reproduce the general feature of a common background level, but with a concentration that is approximately 20 times higher than indicated by HIPPO.

The HIPPO dataset allows us to test the possible implications of these observations on the multimodel BC RF from AeroCom Phase II, a key basis for BC forcing recently assessed in the IPCCs AR5. We attempt two scalings of the modeled BC concentration fields, to align them with the HIPPO observations. The first assumes that the vertically resolved ratio between models and observations in the Pacific holds for all remote ocean regions, shown as grey shaded areas in Figure 1. The second assumes that the supposition of a globally uniform high altitude BC concentration from (Schwarz et al., 2013) is true. While the present dataset is insufficient to determine if such a supposition is true, it is nevertheless interesting to assess its potential impact to see if efforts to measure high altitude BC concentrations should be prioritized. Figure 4 shows the implications of these scalings on the multimodel direct RF due to BC from fossil fuel and biofuel burning (BC FF+BF). Unscaled, the multimodel median RF found here is $+0.24$ [$+0.17$ to $+0.47$] W m^{-2} . (See Table 3) Applying the remote region scaling reduces the global, annual median BC FF+BF RF to $+0.22$ [$+0.16$ to $+0.39$] W m^{-2} . The high altitude scaling reduces it to $+0.19$ [$+0.15$ to $+0.33$] W m^{-2} . Applying both simultaneously, while ensuring that we do not doubly scale in remote, high altitude regions, yields a BC FF+BF RF of $+0.17$ [$+0.13$ to $+0.28$] W m^{-2} , or a reduction of 25% from the AeroCom Phase II value combined with a strong reduction in the model spread.

A 25% reduction in the direct radiative forcing of BC would have significant implications, placing the entire model based 5-95% range below the central BC RF value recently reported in IPCC AR5 (Boucher et al., 2013). Presently, the remaining uncertainty in BC forcing is heavily driven by scalings such as the ones attempted above. The IPCC AR5 assessment took input both from AeroCom Phase II and other studies. One of these studies (Bond et al., 2013) reported a significantly stronger forcing of 0.51 Wm^{-2} from fossil fuel and biomass burning.

That estimate includes both a gross 15% global downscaling of BC forcing efficiency due to overestimation of BC aloft, and a differentiated regional upscaling of emissions derived by comparing aerosol absorption optical depth from AeroCom Phase I with that from AERONET ground based remote sensing. Their downscaling, based on recent model studies (Bond et al., 2013; Samset and Myhre, 2011; Zarzycki and Bond, 2010) and evaluation of AeroCom Phase I results (Schwarz et al., 2010), is comparable to our 25% reduction in forcing, though our reduction is attributed to remote ocean areas. For near source and remote regions covered in the present study we here find no need for an emission bias related upscaling; however the present data do not cover the regions where the upscaling in that analysis (Bond et al., 2013) was most pronounced. Also, the median anthropogenic BC RF in AeroCom Phase II is already a factor of 2 stronger than in Phase I, in part due to differences in emissions and modeled aerosol optical properties (Myhre et al., 2013b). We note that the above conclusions are broadly consistent with recent findings using the GEOS-Chem model, which is not represented in the present dataset (Wang et al., 2014a). It is clear that further observations of BC concentrations, vertically resolved and both in situ and remote, are imperative for constraining the radiative forcing of black carbon.

Applicability of scaling factors derived from total BC to BC FF+BF fields

A question raised by the scaling analysis is whether we bias the results by deriving scaling factors based on total BC fields, and subsequently applying them to BC FF+BF forcing. At present, no measurement exists that can determine systematic differences between the global distributions of total BC and BC from fossil fuel and biofuel burning. However four of the models participating here (OsloCTM2, CAM3, CAM5 and IMPACT) also supplied full 3D concentration fields from BC FF+BF only, and we have used these to test the applicability of the method.

For this subset of models, which spans the range of predicted BC burdens, we found the ratio of modeled anthropogenic BC FF+BF to total BC concentrations to be approximately constant with altitude in the regions defined as remote. While trends exist for individual models in single regions, for the remote regions as a whole the ratio changes by less than 10% through the atmospheric column. Hence any alteration of the BC vertical profile should equally affect both fields. Further, the fraction of the total global mean forcing found to be exerted at altitudes above 200hPa was, for these models, found to be comparable for total BC and BC FF+BF. (See Table 2.)

These two observations lead us to conclude that we do not strongly bias our results by applying scaling factors derived from total BC fields to BC FF+BF forcing.

Forcing pattern from models with low RMS and good correlation with HIPPO

We have shown that of the models participating in the present comparison, there are three (GMI, GISS-MATRIX, IMPACT) that both show a low RMS error and a good correlation with the Pacific HIPPO data. These three models all have low global mean atmospheric lifetimes of BC, and report among the lowest BC FF+BF RF values in the AeroCom Phase II ensemble.

Figure 5 shows the zonal mean BC FF+BF RF, and total BC forcing density (RF per unit height) vertical profile, from the full model ensemble, and from the three HIPPO-corresponding models only. Total BC is used for the vertical profile as not all models provided full 3D concentration fields, as outlined above. The obvious feature is that for these three models, forcing is exerted primarily closer to the sources, and at lower altitudes, than in the full ensemble. Very little is exerted in the southern ocean, or above 200hPa.

Figure 6 shows the results from the scaling analysis above, compared with results for the three HIPPO-corresponding models only. From the outset they have low forcing, and the scaling

exercise does not significantly affect them as there is already very little forcing in the scaled regions. The final model median, however, is consistent with that from the scaled full model ensemble. This gives a separate indication that a reduction of 25% in anthropogenic BC RF relative to the AeroCom Phase II value is reasonable if we take the HIPPO Pacific measurements as guidance.

Conclusions

We have compared recent aircraft based measurements of BC concentration with state of the art global aerosol-climate models. In remote regions where BC concentration are dominated by long range transport, and at high altitudes, there is a tendency for the models to overestimate the aircraft measurements, where and when the effects of fires are small. For a region sensitive to Asian emission sources, models reproduce the aircraft measurements remarkably well, with no indication of an underestimation in BC emissions. In remote ocean regions, an atmospheric lifetime of anthropogenic BC of less than 5 days seems crucial, but not sufficient, to be able to reproduce measurement data. Scaling the multimodel results to HIPPO measurements, remotely and aloft, and assuming a globally uniform high altitude BC concentration, leads to a reduction of 25% in anthropogenic BC direct RF, relative to the models native values. The revised median of 0.17 Wm^{-2} stands in stark contrast to recent assessments, which report up to 2-3 times stronger present day BC forcing, but is in line with recent single-model studies (Wang et al., 2014a; Bauer et al., 2013). This discrepancy underlines the impact of combining measured BC concentration data with model estimates. To resolve these differences, and better constrain the climate impact of BC, there is an urgent need for further flight campaigns to provide BC vertical concentration profiles over both source regions, and regions where anthropogenic BC concentrations are dominated by transport and wet scavenging.

Acknowledgements

B. H. Samset, G. Myhre and M. Schulz were supported by the Research Council of Norway, through the grants SLAC, AEROCOM-P3 and EarthClim, and the EC Seventh Framework Programme under grant agreement FP7-ENV-2011-282688 (ECLIPSE). R.C. Easter S.J. Ghan, X. Liu and K. Zhang were supported by through Scientific Discovery through Advanced Computing (SciDAC) program funded by U.S. Department of Energy Office of Advanced Scientific Computing Research and Office of Biological and Environmental Research. The Pacific Northwest National Laboratory (PNNL) is operated for the DOE by Battelle Memorial Institute under contract DE-AC06-76RLO 1830. P. Stier's research has been supported by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no. FP7- 280025 and by the UK NERC project GASSP (NE/J022624/1). S. E. Bauer was supported by the NASA MAP program (NN-H-04-Z-YS-008-N and NN-H-08-Z-DA-001-N). Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Center for Climate Simulation (NCCS) . T. Iversen, A. Kirkevåg and Ø. Seland were supported by the Research Council of Norway through the EarthClim (207711/E10) and NOTUR/NorStore projects, CRAICC, and through the EU projects PEGASOS and ACCESS. A. Kirkevåg also received funding from the Norwegian Space Center through the PM-VRAE project. The National Center for Atmospheric Research is operated by the University Corporation for Atmospheric Research under sponsorship of the National Science Foundation. N. Bellouin was supported by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (GA01101).

Figure captions





Figure 1: Flight tracks and regions selected for the analysis. Dots represent either single measurements (ARCTAS, A-FORCE, PAMARCMiP) or pre-averaged profiles (HIPPO) from the flight campaigns. Blue: HIPPO. Green: ARCTAS. Red: PAMARCMiP. Black: A-FORCE. Green boxes show the geographical regions where model data was averaged for comparison to the flight profiles. Grey boxes show the areas defined as “remote ocean”. The background map shows the median, annual mean RF due to fossil fuel and biofuel burning from the 13 models used in the present study.



485

486

Figure 2: Comparison of measurements and model data for all selected regions. For each panel, the left box shows an

487

overlay of the observed total BC concentration profiles (black lines: mean (solid), median (dotted) and mean+1 standard

488

deviation (dashed), and 25th-75th percentile range (grey band)) with the mean BC concentration profiles from individual

AeroCom Phase II models (colored lines, see legend). The three middle boxes show, from left to right, the BC burden (mg/m²), direct radiative forcing (W/m²) and forcing efficiency (W/g) for observations (black) and models (red). The colored diamonds show the individual AeroCom Phase II models. Finally, the rightmost box shows the ratio of models to observations for the burden (green), radiative forcing (blue) and forcing efficiency (red) within the selected region.




Figure 3: Modeled global black carbon lifetime plotted versus root-mean-square error, mean bias and normalized mean bias, between model prediction and observations for the five HIPPO regions (P1-P5). Each RMS or bias calculation is based on 72 data points. The symbol size indicates the corresponding Pearson correlation coefficient. The black dashed line shows a least-squares regression line with all models included. The grey lines show regressions with one model removed. The red dashed line shows a regression using only models with $p > 0.8$.





Figure 4: Model median and 5%-95% range for BC FF+BF forcing, for 1750-2010, with various scalings applied. The yellow bar shows the AeroCom Phase II result (Myhre et al., 2013b). The grey bar shows unscaled values from the present work, then with remote scaling (pink) and high altitude scaling (blue) applied. The khaki bar shows the lower limit on BC FF+BF forcing from the present work, with both scalings applied. Below we compare with the recent estimate in IPCC WG1's AR5.



507

508 **Figure 5: Zonal mean and altitude forcing profiles for AeroCom Phase II models. a) BC FF+BF forcing, zonal mean, for all**
 509 **13 models in the present study (black, solid), and the three models selected based on RMS and correlation vs HIPPO (red,**
 510 **solid). Dotted lines show only forcing over land, dashed show forcing over ocean. b) As a, except total BC (FF+BF+BB)**
 511 **forcing, and global, annual mean vertical profile. Forcing is shown per altitude meter (unit Wm^{-3}), to avoid dependence**
 512 **on model vertical structure.**




Figure 6: As Figure 4, except also showing the BC FF+BF forcing from the three models selected based on RMS and correlation vs HIPPO (hatched boxes).

Supplementary tables:

Table 1: The flight campaigns included in the present work, and the times when they flew.

Campaign	Region	Time	Data/web site
HIPPO 1	Pacific	January 2009	http://hippo.ucar.edu/
HIPPO 2	Pacific	November 2009	
HIPPO 3	Pacific	March/April 2010	
HIPPO 4	Pacific	June 2011	
HIPPO 5	Pacific	August 2011	
A-FORCE	Japan	March/April 2009	
ARCTAS Spring	Northern Pacific, North Polar	April 2008	http://wwwair.larc.nasa.gov/ missions/
ARCTAS Summer	Continental North America	July 2008	arctas/dataaccess.htm
PAMARCMiP 2009	Northern Pacific, North Polar	April 2009	
PAMARCMiP 2011	Northern Pacific, North Polar	April 2011	
PAMARCMiP 2012	North Polar	March - April 2012	

Table 2: Modeled mean forcing and fractions for the regions used in the present analysis (see Figure 1). Mean RF is the forcing within that region. Fraction of global RF is defined as the fraction of energy deposited, on annual mean, within that region. "Remotes" represents the total of all grey marked regions in Figure 1.

Region	Mean RF [Wm-2]	Fraction of global area [1]	Fraction of global RF [1]	Fraction of RF above 200hPa [1]
Global	0.26	1.00	1.00	0.24
America	0.32	0.01	0.01	0.30
Japan	0.98	0.01	0.04	0.20
P1	0.28	0.01	0.01	0.31
P2	0.39	0.02	0.03	0.28
P3	0.17	0.03	0.02	0.32
P4	0.06	0.03	0.01	0.43
P5	0.06	0.02	0.01	0.47
NP1	0.26	0.003	0.003	0.30
NP2	0.25	0.003	0.003	0.31
Remotes	0.17	0.26	0.15	0.35

Table 3: BC FF+BF RF and BC atmospheric lifetime of BC for the models used in the present study. Bias, RMS error and correlation coefficients are for comparisons of each model with HIPPO data for all Pacific regions (P1-P5). Each calculation is based on 72 data points.

Model	BC FF+BF RF [Wm ⁻²]	BC Lifetime [days]	RMS error [ngm ⁻³]	Correlation [1]	Mean bias [ngm ⁻³]	Normalized bias [1]
INCA	0.18	7.1	5.6	0.12	0.002	0.79
GOCART	0.18	7.2	13.3	0.86	0.010	1.34
OsloCTM2	0.28	6.0	4.7	0.64	0.003	0.87
CAM4-Oslo	0.37	8.2	15.9	0.72	0.014	1.44
SPRINTARS	0.21	6.7	4.4	0.62	0.002	0.78
ECHAM-HAM	0.14	5.5	4.5	0.39	-0.002	0.01
HadGEM2	0.19	17.1	34.1	0.84	0.025	1.56
GISS-modeIE	0.21	5.9	6.1	0.67	0.004	0.87
IMPACT	0.14	3.8	2.7	0.82	0.000	0.10
GMI	0.17	4.4	4.1	0.86	0.003	0.68
CAM5	0.2	3.8	4.6	0.37	-0.001	0.15
NCAR-CAM3.5	0.15	5.8	4.1	0.80	0.002	0.71
GISS-MATRIX	0.19	3.5	2.9	0.87	-0.001	-0.49

533 **References**

- 534 Ban-Weiss, G. A., Cao, L., Bala, G., and Caldeira, K.: Dependence of climate forcing and response on
535 the altitude of black carbon aerosols, *Climate Dynamics*, 38, 897-911, 10.1007/s00382-011-1052-y,
536 2011.
- 537 Bauer, S. E., Menon, S., Koch, D., Bond, T. C., and Tsigaridis, K.: A global modeling study on
538 carbonaceous aerosol microphysical characteristics and radiative effects, *Atmos Chem Phys*, 10,
539 7439-7456, DOI 10.5194/acp-10-7439-2010, 2010.
- 540 Bauer, S. E., Bausch, A., Nazarenko, L., Tsigaridis, K., Xu, B. Q., Edwards, R., Bisiaux, M., and
541 McConnell, J.: Historical and future black carbon deposition on the three ice caps: Ice core
542 measurements and model simulations from 1850 to 2100, *J Geophys Res-Atmos*, 118, 7948-7961, Doi
543 10.1002/Jgrd.50612, 2013.
- 544 Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the
545 Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of
546 ammonium nitrate, *J Geophys Res-Atmos*, 116, ArtD20206
Doi 10.1029/2011jd016074, 2011.
- 548 Bian, H., Chin, M., Rodriguez, J. M., Yu, H., Penner, J. E., and Strahan, S.: Sensitivity of aerosol optical
549 thickness and aerosol direct radiative effect to relative humidity, *Atmos Chem Phys*, 9, 2375-2386,
550 2009.
- 551 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G.,
552 Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz,
553 M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M.
554 Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and
555 Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment,
556 *Journal of Geophysical Research: Atmospheres*, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- 557 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo,
558 Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds
559 and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I*
560 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D.
561 Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley
562 (eds.)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- 563 Chin, M., Diehl, T., Dubovik, O., Eck, T. F., Holben, B. N., Sinyuk, A., and Streets, D. G.: Light
564 absorption by pollution, dust, and biomass burning aerosols: a global model study and evaluation
565 with AERONET measurements, *Ann Geophys-Germany*, 27, 3439-3464, 2009.
- 566 Chung, C. E., Ramanathan, V., and Decremier, D.: Observationally constrained estimates of
567 carbonaceous aerosol radiative forcing, *P Natl Acad Sci USA*, 109, 11624-11629, DOI
568 10.1073/pnas.1203707109, 2012.
- 569 Flanner, M. G.: Arctic climate sensitivity to local black carbon, *J Geophys Res-Atmos*, 118, 1840-1851,
570 Doi 10.1002/Jgrd.50176, 2013.
- 571 Haywood, J. M., and Shine, K. P.: The Effect of Anthropogenic Sulfate and Soot Aerosol on the Clear-
572 Sky Planetary Radiation Budget, *Geophys Res Lett*, 22, 603-606, 1995.
- 573 Herber, A. B., Haas, C., Stone, R. S., Bottenheim, J. W., Liu, P., Li, S.-M., Staebler, R. M., Strapp, J. W.,
574 and Dethloff, K.: Regular airborne surveys of Arctic sea ice and atmosphere, *Eos, Transactions*
575 *American Geophysical Union*, 93, 41, 10.1029/2012eo040001, 2012.
- 576 Hodnebrog, O., Myhre, G., and Samset, B. H.: How shorter black carbon lifetime alters its climate
577 effect, *Nature Communications* (Accepted, in press.), 2014.
- 578 Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A.,
579 Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R.,
580 and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and

581 Satellites (ARCTAS) mission: design, execution, and first results, *Atmos Chem Phys*, 10, 5191-5212,
582 10.5194/acp-10-5191-2010, 2010.

583 Kipling, Z., Stier, P., Schwarz, J. P., Perring, A. E., Spackman, J. R., Mann, G. W., Johnson, C. E., and
584 Telford, P. J.: Constraints on aerosol processes in climate models from vertically-resolved aircraft
585 observations of black carbon, *Atmos Chem Phys*, 13, 5969-5986, DOI 10.5194/acp-13-5969-2013,
586 2013.

587 Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjánsson, J. E., Struthers, H., Ekman, A. M. L., Ghan,
588 S., Griesfeller, J., Nilsson, E. D., and Schulz, M.: Aerosol-climate interactions in the Norwegian Earth
589 System Model – NorESM1-M, *Geosci. Model Dev.*, 6, 207-244, 10.5194/gmd-6-207-2013, 2013.

590 Koch, D., Bauer, S. E., Del Genio, A., Faluvegi, G., McConnell, J. R., Menon, S., Miller, R. L., Rind, D.,
591 Ruedy, R., Schmidt, G. A., and Shindell, D.: Coupled Aerosol-Chemistry-Climate Twentieth-Century
592 Transient Model Investigation: Trends in Short-Lived Species and Climate Responses, *J Climate*, 24,
593 2693-2714, Doi 10.1175/2011jcli3582.1, 2011.

594 Koffi, B., Schulz, M., Breon, F. M., Griesfeller, J., Winker, D., Balkanski, Y., Bauer, S., Berntsen, T., Chin,
595 M. A., Collins, W. D., Dentener, F., Diehl, T., Easter, R., Ghan, S., Ginoux, P., Gong, S. L., Horowitz, L.
596 W., Iversen, T., Kirkevåg, A., Koch, D., Krol, M., Myhre, G., Stier, P., and Takemura, T.: Application of
597 the CALIOP layer product to evaluate the vertical distribution of aerosols estimated by global models:
598 AeroCom phase I results, *J Geophys Res-Atmos*, 117, ArtD10201
599 Doi 10.1029/2011jd016858, 2012.

600 Kondo, Y., Sahu, L., Moteki, N., Khan, F., Takegawa, N., Liu, X., Kolke, M., and Miyakawa, T.:
601 Consistency and Traceability of Black Carbon Measurements Made by Laser-Induced Incandescence,
602 Thermal-Optical Transmittance, and Filter-Based Photo-Absorption Techniques, *Aerosol Sci Tech*, 45,
603 295-312, Doi 10.1080/02786826.2010.533215, 2011.

604 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A.,
605 Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R.,
606 Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical
607 (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols:
608 methodology and application, *Atmos Chem Phys*, 10, 7017-7039, DOI 10.5194/acp-10-7017-2010,
609 2010.

610 Lamarque, J. F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E.
611 A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and
612 evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci*
613 *Model Dev*, 5, 369-411, DOI 10.5194/gmd-5-369-2012, 2012.

614 Lin, G., Penner, J., Sillmann, S., Taraborrelli, D., and Lelieveld, J.: Global modeling of SOA formation
615 from dicarbonyls, epoxides, organic nitrates and peroxides, *Atmos Chem Phys*, 12, 4743-4774,
616 10.5194/acp-12-4743-2012, 2012.

617 Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison,
618 H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins,
619 W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation
620 of aerosols in climate models: description and evaluation in the Community Atmosphere Model
621 CAM5, *Geosci Model Dev*, 5, 709-739, 10.5194/gmd-5-709-2012, 2012.

622 Mann, G. W., Carslaw, K. S., Reddington, C. L., Pringle, K. J., Schulz, M., Asmi, A., Spracklen, D. V.,
623 Ridley, D. A., Woodhouse, M. T., Lee, L. A., Zhang, K., Ghan, S. J., Easter, R. C., Liu, X., Stier, P., Lee, Y.
624 H., Adams, P. J., Tost, H., Lelieveld, J., Bauer, S. E., Tsigaridis, K., van Noije, T. P. C., Strunk, A., Vignati,
625 E., Bellouin, N., Dalvi, M., Johnson, C. E., Bergman, T., Kokkola, H., von Salzen, K., Yu, F., Luo, G.,
626 Petzold, A., Heintzenberg, J., Clarke, A., Ogren, J. A., Gras, J., Baltensperger, U., Kaminski, U., Jennings,
627 S. G., O'Dowd, C. D., Harrison, R. M., Beddows, D. C. S., Kulmala, M., Viisanen, Y., Ulevicius, V.,
628 Mihalopoulos, N., Zdimal, V., Fiebig, M., Hansson, H. C., Swietlicki, E., and Henzig, J. S.:
629 Intercomparison and evaluation of aerosol microphysical properties among AeroCom global models
630 of a range of complexity, *Atmospheric Chemistry and Physics Discussions*, 13, 30841-30928,
631 10.5194/acpd-13-30841-2013, 2013.

632 Myhre, G., Myhre, C., Samset, B. H., and Storelvmo, T.: Aerosols and their Relation to Global Climate
 633 and Climate Sensitivity, Nature Knowledge Project, 4, 2013a.
 634 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N.,
 635 Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S.,
 636 Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Luo, G., Ma, X., Penner, J. E., Rasch, P. J., Seland, Ø., Skeie,
 637 R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J. H., Zhang, K., Zhang,
 638 H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations,
 639 Atmos Chem Phys, 13, 1-25, doi:10.5194/acp-13-1-2013, 2013b.
 640 Oshima, N., Kondo, Y., Moteki, N., Takegawa, N., Koike, M., Kita, K., Matsui, H., Kajino, M., Nakamura,
 641 H., Jung, J. S., and Kim, Y. J.: Wet removal of black carbon in Asian outflow: Aerosol Radiative Forcing
 642 in East Asia (A-FORCE) aircraft campaign, Journal of Geophysical Research, 117,
 643 10.1029/2011jd016552, 2012.
 644 Ramanathan, V., and Carmichael, G.: Global and regional climate changes due to black carbon, Nat
 645 Geosci, 1, 221-227, Doi 10.1038/Ngeo156, 2008.
 646 Samset, B. H., and Myhre, G.: Vertical dependence of black carbon, sulphate and biomass burning
 647 aerosol radiative forcing, Geophys Res Lett, 38, Artn L24802
 648 Doi 10.1029/2011gl049697, 2011.
 649 Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N.,
 650 Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X.,
 651 Penner, J. E., Seland, O., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon
 652 vertical profiles strongly affect its radiative forcing uncertainty, Atmos Chem Phys, 13, 2423-2434,
 653 DOI 10.5194/acp-13-2423-2013, 2013.
 654 Samset, B. H., Myhre, G., and Schulz, M.: Upward adjustment needed for aerosol radiative forcing
 655 uncertainty, Nat Clim Change, 4, 230-232, 10.1038/nclimate2170, 2014.
 656 Schwarz, J. P., Spackman, J. R., Gao, R. S., Watts, L. A., Stier, P., Schulz, M., Davis, S. M., Wofsy, S. C.,
 657 and Fahey, D. W.: Global scale black carbon profiles observed in the remote atmosphere and
 658 compared to models, Geophys Res Lett, 37, 2010.
 659 Schwarz, J. P., Samset, B. H., Perring, A. E., Spackman, J. R., Gao, R. S., Stier, P., Schulz, M., Moore, F.
 660 L., Ray, E. A., and Fahey, D. W.: Global-scale seasonally resolved black carbon vertical profiles over
 661 the Pacific, Geophys Res Lett, 40, 5542-5547, Doi 10.1002/2013gl057775, 2013.
 662 Skeie, R. B., Bernsten, T., Myhre, G., Pedersen, C. A., Strom, J., Gerland, S., and Ogren, J. A.: Black
 663 carbon in the atmosphere and snow, from pre-industrial times until present, Atmos Chem Phys, 11,
 664 6809-6836, 10.5194/acp-11-6809-2011, 2011.
 665 Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V. P., Kopeikin, V. M., and Novigatsky, A.
 666 N.: Black carbon in the Arctic: the underestimated role of gas flaring and residential combustion
 667 emissions, Atmos Chem Phys, 13, 8833-8855, DOI 10.5194/acp-13-8833-2013, 2013.
 668 Stone, R. S., Herber, A., Vitale, V., Mazzola, M., Lupi, A., Schnell, R. C., Dutton, E. G., Liu, P. S. K., Li, S.
 669 M., Dethloff, K., Lampert, A., Ritter, C., Stock, M., Neuber, R., and Maturilli, M.: A three-dimensional
 670 characterization of Arctic aerosols from airborne Sun photometer observations: PAM-ARCMIP, April
 671 2009, Journal of Geophysical Research, 115, 10.1029/2009jd013605, 2010.
 672 Szopa, S., Balkanski, Y., Schulz, M., Bekki, S., Cugnet, D., Fortems-Cheiney, A., Turquety, S., Cozic, A.,
 673 Déandres, C., Hauglustaine, D., Idelkadi, A., Lathière, J., Lefevre, F., Marchand, M., Vuolo, R., Yan, N.,
 674 and Dufresne, J. L.: Aerosol and ozone changes as forcing for climate evolution between 1850 and
 675 2100, Climate Dynamics, 40, 2223-2250, 10.1007/s00382-012-1408-y, 2012.
 676 Takemura, T., Nozawa, T., Emori, S., Nakajima, T. Y., and Nakajima, T.: Simulation of climate response
 677 to aerosol direct and indirect effects with aerosol transport-radiation model, J Geophys Res-Atmos,
 678 110, Artn D02202
 679 Doi 10.1029/2004jd005029, 2005.
 680 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher,
 681 O., Chin, M., Dentener, F., Diehl, T., Feichter, J., Fillmore, D., Ginoux, P., Gong, S., Grini, A., Hendricks,

682 J., Horowitz, L., Huang, P., Isaksen, I. S. A., Iversen, T., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J.
 683 E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy,
 684 M. S., Seland, O., Stier, P., Takemura, T., and Tie, X.: The effect of harmonized emissions on aerosol
 685 properties in global models - an AeroCom experiment, *Atmos Chem Phys*, 7, 4489-4501, 2007.
 686 Wang, Q., Jacob, D. J., Fisher, J. A., Mao, J., Leibensperger, E. M., Carouge, C. C., Le Sager, P., Kondo,
 687 Y., Jimenez, J. L., Cubison, M. J., and Doherty, S. J.: Sources of carbonaceous aerosols and deposited
 688 black carbon in the Arctic in winter-spring: implications for radiative forcing, *Atmos Chem Phys*, 11,
 689 12453-12473, 2011.
 690 Wang, Q. Q., Jacob, D. J., Spackman, J. R., Perring, A. E., Schwarz, J. P., Moteki, N., Marais, E. A., Ge, C.,
 691 Wang, J., and Barrett, S. R. H.: Global budget and radiative forcing of black carbon aerosol:
 692 Constraints from pole-to-pole (HIPPO) observations across the Pacific, *J Geophys Res-Atmos*, 119,
 693 195-206, Doi 10.1002/2013jd020824, 2014a.
 694 Wang, R., Tao, S., Shen, H. Z., Huang, Y., Chen, H., Balkanski, Y., Boucher, O., Ciais, P., Shen, G. F., Li,
 695 W., Zhang, Y. Y., Chen, Y. C., Lin, N., Su, S., Li, B. G., Liu, J. F., and Liu, W. X.: Trend in Global Black
 696 Carbon Emissions from 1960 to 2007, *Environ Sci Technol*, 48, 6780-6787, 2014b.
 697 Zarzycki, C. M., and Bond, T. C.: How much can the vertical distribution of black carbon affect its
 698 global direct radiative forcing?, *Geophys Res Lett*, 37, L20807, Artn L20807
 699 Doi 10.1029/2010gl044555, 2010.
 700 Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J.,
 701 Wan, H., Rast, S., and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2:
 702 sensitivity to improvements in process representations, *Atmos. Chem. Phys.*, 12, 8911-8949,
 703 10.5194/acp-12-8911-2012, 2012.

704