

Responses to reviews of Loh et al., Simulations of atmospheric methane for Cape Grim, Tasmania, to constrain South East Australian methane emissions

November 26, 2014

Reply to review 1

The review is in italics and our responses are in normal font. Line and page numbers for manuscript changes are in bold and correspond to the pdf version without marked up changes.

... *The authors discuss the main differences between the model ratios and the observations, and attribute the discrepancies between the models and observation to the representation of wetland methane. But they do not mention differences between models and other specific mismatches, this leaves questions.*

We have assumed that these 'questions' are those detailed below and that in addressing those comments we have answered this more general point.

The manuscript is descriptive with the simulation results with the prescribed emission scenarios. It would have been nice to see additional model experiments to support their implications. From the comparison between the simulation results and observations, the authors conclude that the austral winter wetland methane emissions are overestimated, but suggesting the springtime maximum of wetland emissions. To support the idea, the authors show the satellite measurements (Fraser et al., 2011), but it would be more interesting and supportive to try to assess (at least partially) the sensitivity of seasonality in wetland emission to the simulated residual CH₄ concentrations, as well as the ratio of CH₄ and radon.

While we anticipate that a future study will further investigate our hypothesis of greater springtime methane emissions, we have performed an additional sensitivity experiment with CCAM in which we shifted the CTL fluxes (across the whole globe) later in time by 3 months. Since the only significant seasonality in the CTL fluxes for SE Australia is the wetland component, this gives an initial indication of the likely impact of moving wetland emissions from winter to spring. We find that this gives largest methane to radon ratios in September-November instead of May-August with slightly smaller magnitude (around 12 ppb/Bqm⁻³ instead of around 13 ppb/Bqm⁻³). This is a better fit to the observations although a further

one month shift in wetland fluxes would likely give a further improvement and the early winter CH₄/radon ratios remain too high. This may be due to large methane fluxes in February which are just north of our region of interest and in this test get shifted to May. We have rewritten the first paragraph of Section 5 (now split into 4 paragraphs) to include this information (in the third paragraph, **p16, line 26 to p17, line 12**) and added Figure 8 (**p 36**) to show the methane to radon ratio seasonality for this new simulation.

2 Detailed comments

- *Section 3.1: The authors state that the maximums of WLBB and EXTRA are in December and January due to biomass burning, as seen Figure 3. I wonder how much the annual wetland emission in WLBB and EXTRA and how the seasonality looks like, compared to the wetland fluxes in CTL and CTL-E₄ and BB which has annual total of 1.24Tg/yr with high emissions for May to October. How about the CH₄ emission from rice cultivation? How significant the rice emission among the total CH₄ emissions and in the seasonality? The seasonal variation of wetland (and rice) CH₄ emission is one of the key points in this study. These would be helpful to understand the results discussed in the later sections.*

For the TransCom-CH₄ model intercomparison, the emission scenarios were provided to modellers as the sum of component emissions. For the CTL case only, the individual components were also made available for analysis purposes and it was from this information that we were able to determine that wetland emissions dominated the seasonal cycle of CTL fluxes for SE Australia (while rice emissions are much smaller, approximately 0.13 Tg/y in summer and zero in winter). In theory the wetland component of WLBB should be deducible from WLBB - BB + CTL(wetland component). We have done this but the resulting flux shows a residual biomass burning signal (identifiable by its interannual variability), possibly due to a slightly different scaling being applied when each scenario was created, in order to match a global total flux. This makes it impossible to exactly determine the wetland and rice component fluxes in the WLBB and EXTRA scenarios although they appear to be close to zero for SE Australia or they would not be obscured by the residual biomass burning signal. This examination of the emission scenarios highlighted a couple of mistakes in the description of the emissions in Sec 3.1. These have now been corrected in the text (**p8, line 3 improve clarity on BB emissions and p8, line 7 correct source of wetland emissions for EXTRA**) and in Table 1 (**p26, EXTRA row, Wetland column changed from Ringeval et al, to Ito and Inatomi**).

- *Table 2: The difference between WLBB and EXTRA is only rice CH₄ emission. The inter-annual variation periods of wetland emissions are different between WLBB and EXTRA in Table 2? Why?*

The period of interannual variation is correct in Table 2. The error was in Table 1 (and the text). The wetland emissions for EXTRA are from the VISIT model (as for rice emissions) and not from Ringeval et al as used in WLBB. Manuscript changes are as previous point.

- *Section 3.2: The description on ACCESS is not clear. How different is the run for this study from the run for TransCom-CH₄? Why this ACCESS run can be analysed on synoptic variations for statistical purpose, but not for individual events? It would be helpful to give more description of the ACCESS model, and to explain why the two models used in this study.*

We have rewritten the paragraph describing the ACCESS model (now paragraph 3 of Sec 3.2, **p10, line 10-21**) to give more detail about the model simulation and how it compares to the one submitted for TransCom-CH₄ (only the horizontal resolution was changed). A sentence has been added to the first paragraph of Sec 3.2 (**p9, line 29 to p10, line 2**) to note why two models have been used (to determine whether the analysis is sensitive to transport model error).

- *Section 4: "...exceptionally high methane concentration in January and February... (BB, WLBB and EXTRA)" But it seems inconsistent with Figure 4, where high concentrations in January and February can be seen in BB and WLBB and INV in (a)ACCESS, and BB and INV in (b)CCAM. EXTRA shows the lowest concentrations among the simulations by both model runs.*

Sorry for the confusion. The text was correct but the EXTRA and INV lines in Figure 4 were incorrectly labelled. This has been fixed. The figure has also been extended to include 2002 and 2004 to provide some context for the large summer concentrations in 2003. (**p32**)

- *Figure 4: BB and EXTRA are in similar colors. It is hard to distinguish.*

Agreed. We have redrawn the figure using the same colour scheme as Figure 3. (**p32**)

- *Section 4.1: the model experiments with injection levels of fire methane have been conducted. The results are described very briefly, but it is not clear. Does the simulation with fire methane emitted in higher model level give comparable amplitude to the observation?*

The sensitivity tests are not directly comparable to the observations since they model only one component of the CH₄ flux (biomass burning). Nevertheless, they do illustrate how the amplitude can be reduced substantially when emissions are inserted higher into the atmosphere. An extra paragraph has been added to Sec 4.1 (**p12, line 25 to p13, line 9**) to describe the sensitivity tests in more detail including giving one measure of amplitude for comparison between cases.

- *Section 4.2: the discussion here is mainly regarding the difference of wetland emissions. But in Figure 7, the model runs with CTL-E₄ show the difference each other, unlike the other simulations. So I would like to see some explanation on CTL-E₄. How different are anthropogenic emissions between EDGAR 3.2 and EDGAR 4.0 for southeastern Australia? I understand both are annually constant with no seasonal variation. If the difference is only increasing trend (seen in Figure 3b), CTL-E₄ simulation results can be expected to be similar to CTL in this study which is focused on the non-baseline events? But the results of CTL-E₄ are different from CTL, especially in ACCESS run. Furthermore, the seasonal cycles of both model ratios*

are not close to the observations, but the springtime ratios are more comparable to the observations than other emission scenarios.

We have added a paragraph at the end of Section 4.2 (**p15, line 26 to p16, line 8**) to discuss the CTL_E4 case. The difference in winter to summer ratios is smaller with CTL_E4 (despite larger ratios overall). There are differences in the spatial distribution of fluxes between CTL and CTL_E4 and we think this is the most likely explanation for the slightly weakened seasonality. We have looked at the trend in ratio over time between the different cases to see if the increasing emissions from CTL_E4 are supported by the observations, but the large seasonal and interannual variability in CH₄/radon ratio means that we have little confidence in any calculated trend.

- *P.21205: The discussion on Wang and Bentley (2002) is not easy to follow. They suggested the large reduction of CH₄ emission in the region of interest in this study? In Section 3, however, it can read that the CH₄ emission estimated by Wang and Bentley (2002) is comparable or slightly larger than those by Fraser et al. (2011) and EDGAR.*

This text was confusing because it was not clear that Wang and Bentley provide both an inventory based flux estimate and one based on an inversion (which is much lower than the inventory). We quoted the inventory based flux early in the paper but discussed the inversion based estimate here. We have hopefully removed this confusion by adding a table of flux estimates from different sources (as requested by reviewer 2) (Table 3, **p28**) referenced in Sec 3.1 (**p8, line 28**) and modifying the text in the second last paragraph in Sec 5 (**p18, line 18-28**).

- *p.21205, L24: Please spell out NSW.*

Done (**p18, line 17**).

- *Figure 7, "tracers" should be "emission scenarios"? It is hard to see the bars of standard deviation for January and December. Their presentations need to be improved.*

The figure caption has been changed. The x axis has been extended to make January and December easier to read and the error bars for each curve have been offset slightly to avoid overlap. (**p35**)

Reply to review 2

The review is in italics and our responses are in normal font. Changes to the manuscript are noted in bold using the page and line number from the pdf without marked up changes.

Major comments:

The authors should show how well such low-resolution global models reproduce local tracer transport phenomena (here, local does not mean the southeastern Australia region, but means a much smaller region). If the models cannot simulate local transport phenomena, it cannot be ruled out that the model-observation discrepancies are

caused by the model error. If the models fail to reproduce local transport and some local emission contributed seasonal CH₄ variations, a different conclusion could be made. Even if CH₄ data are excluded at times when CCAM fails to reproduce observed high-radon concentration, are seasonal patterns of CH₄ concentration and CH₄-radon ratio still similar to those shown in the manuscript?

Low resolution global models (with low resolution input fluxes) are unlikely to reproduce local tracer transport phenomena. Additionally, using offshore grid-points to represent Cape Grim, aims to minimise the influence of local fluxes to preferentially focus on regional signals. Given this modelling strategy, the reviewer asks a valid question as to whether the analysis is sensitive to local flux signals in the observations and whether local signals contribute to any of the discrepancy seen between model and observed seasonality. To test this, we have repeated the calculation of the monthly observed methane-radon ratios but excluding all methane-radon pairs for times when the observed wind speed is lower than 3 m/s at 10m. Since local signals are more likely to dominate when wind speed is low, this should minimise the influence of local fluxes on the analysis. We find that there is no significant change to the observed seasonality of methane to radon ratio. Monthly ratios change by at most 0.4 ppb/(Bq m⁻³) or less than 6%. Given the small impact of this wind speed filter, we have chosen not to modify the text.

Minor comments:

It would be helpful that the CH₄ budget estimates for southeastern Australia (by this study and other previous papers) are summarized by a table.

This has been added (Table 3, **p28**) and is referenced in Sec 3.1 (**p8, line 28**) and Sec 5 (**p18, line 20**).

P21193, L4: ‘Cape Grim has been operating...’ =_j may be like ‘The Cape Grim station has been operating...’

Done (**p4, line 28 to p5, line 1**)

Figure 2: Why is the year 2003 chosen for showing CH₄ and radon observation time series, though radon data are missing in one month?

There was no specific reason for showing 2003. We have changed the figure to 2006 which has more complete records. The text in paragraph 2, Section 2 (**p5, line 4**) has been modified accordingly.

P21193, L25: Which inlet data did the authors use for their analyses, 75m or 10m?

Data from both inlets are used. This information has been added to the text (Sec 2, paragraph 3, **p5, line 21**: "The data from both inlets are used in this study").

P21196, L16: Please clarify that EXTRA has the same IAV biomass burning emission as BB and WLBB.

This information has been added (Sec 3.1, paragraph 2, **p8, line 6**: "EXTRA uses the same biomass burning as BB and interannually-varying model generated wetlands and rice emissions from the VISIT model").

P21198, L5: Maybe, the wetland emission included in the EXTRA scenario is from VISIT (Ito and Inatomi, 2012), but not from Ringeval et al. (2010).

The wetland emission in EXTRA had been incorrectly attributed to Ringeval et al. The text (Sec 3.1, paragraph 2, as point above) and table 1 (**p26**) have been corrected.

P21199, L5: Please elaborate why did the authors choose the model output points that are located slightly to the north of Cape Grim. If the model grids were chosen by seeing how well radon concentrations are simulated (as stated in P21202, L10-12), please describe it here.

For CCAM this was the nearest ocean grid point. For ACCESS the choice was made based on the radon simulation. Two sentences have been added to the appropriate paragraph in Sec 3.2 (was paragraph 3, now paragraph 4, **p10, line 27 to p11, line 1**).

P21199, L15: When are ‘two periods’?

This information has been added (Sec 4, paragraph 1, **p11, line10**: "The first feature was two periods, December 2002 to February 2003 and November 2006 to January 2007, with very high peak...").

P21199, L16-18: Is the feature of the large winter CH₄ differences coherently found in all the years?

Yes. ‘each’ has been added before ‘winter’ in the appropriate sentence (**p11, line 13**) to make this clear and Figure 4 (**p32**) has been extended to show three years to further illustrate this point.

P2199, L25-27: ‘the remaining emission : : . Bousquet et al. (2006)’ should be changed to like ‘the remaining emission scenarios using modified wetland emissions (WLBB and EXTRA) or that based on the inversion of Bousquet et al. (2006) (INV)’.

Done (**p11, line 21-22**).

P21200, L3: ‘H₂’ =_g ‘hydrogen (H₂)’, ‘CO’ =_g ‘carbon monoxide (CO)’

Done (**p11, line 25**).

P21205, L14: ‘WLBB’ might be ‘WLBB/EXTRA’

Done (**p18, line 8**).

Simulations of atmospheric methane for Cape Grim, Tasmania, to constrain South East Australian methane emissions

**Z. M. Loh¹, R. M. Law¹, K. D. Haynes^{1,*}, P. B. Krummel¹, L. P. Steele¹, P. J. Fraser¹,
S. D. Chambers², and A. G. Williams²**

¹Centre for Australian Weather and Climate Research, CSIRO Oceans and Atmosphere Flagship,
Private Bag 1, Aspendale, Vic 3195, Australia

²Australian Nuclear Science and Technology Organisation, Locked Bag 2001, Kirrawee DC, NSW
2232, Australia

*now at: Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA

Correspondence to: R. M. Law (rachel.law@csiro.au)

Abstract

This study uses two climate models and six scenarios of prescribed methane emissions to compare modelled and observed atmospheric methane between 1994 and 2007, for Cape Grim, Australia (40.7° S, 144.7° E). The model simulations follow the TransCom-CH₄ protocol and use the Australian Community Climate and Earth System Simulator (ACCESS) and the CSIRO Conformal-Cubic Atmospheric Model (CCAM). Radon is also simulated and used to reduce the impact of transport differences between the models and observations. Comparisons are made for air samples that have traversed the Australian continent. All six emission scenarios give modelled concentrations that are broadly consistent with those observed. There are three notable mismatches, however. Firstly, scenarios that incorporate interannually varying biomass burning emissions produce anomalously high methane concentrations at Cape Grim at times of large fire events in southeastern Australia, most likely due to the fire methane emissions being unrealistically input into the lowest model level. Secondly, scenarios with wetland methane emissions in the austral winter overestimate methane concentrations at Cape Grim during wintertime while scenarios without winter wetland emissions perform better. Finally, all scenarios fail to represent a methane source in austral spring implied by the observations. It is possible that the timing of wetland emissions in the scenarios is incorrect with recent satellite measurements suggesting an austral spring (September-October-November), rather than winter, maximum for wetland emissions.

1 Introduction

Methane (CH₄) is an important greenhouse gas whose atmospheric concentration has more than doubled since the 18th century (MacFarling Meure et al., 2006), with considerable variations in its growth rate over recent decades (Rigby et al., 2008; Dlugokencky et al., 2009; Sussmann et al., 2012). Methane has both anthropogenic and natural emissions, while the main sink for methane is through reaction with hydroxyl radical (OH) in the tropo-

sphere and by photolysis in the stratosphere. On a global scale, considerable uncertainty remains about the causes of recent changes in the methane growth rate (Kirschke et al., 2013), and on a regional level, significant discrepancies have been found between “bottom up” inventory estimates of emissions and “top down” atmospheric inverse modelling studies. For example, in Miller et al. (2013), the authors find that the inventories underestimate emissions by up to 2.7 times in the south-central USA.

Atmospheric concentrations of methane are measured approximately weekly at more than 150 flask-sampling sites worldwide and continuously at a smaller number of in situ instrumented sites. Some of the longer in situ records are those of the Advanced Global Atmospheric Gases Experiment (AGAGE) network (Cunnold et al., 2002), including the record from Cape Grim, Australia. A number of global modelling studies have used such methane atmospheric records to constrain the methane budget for example, Fung et al. (1991), Houweling et al. (1999) and Bousquet et al. (2011). Rigby et al. (2008) used the AGAGE in situ methane and CSIRO flask methane data, together with the AGAGE methyl chloroform records to deduce the Northern and Southern Hemisphere changes in methane emissions required to account for the measured increase in methane mole fraction growth rate from 2007.

Recently a model intercomparison, “TransCom-CH₄”, has been run for methane (Patra et al., 2011), with a focus on understanding how transport model differences contributed to variations in the methane simulations. All model simulations used the same prescribed methane emissions and modelled methane loss using prescribed, climatological OH fields. We have used the simulations defined for this intercomparison to compare simulated and measured methane at Cape Grim, with the goal of evaluating the accuracy of the prescribed methane emissions for southeastern Australia. The model intercomparison also simulated radon, sulfur hexafluoride (SF₆) and methyl chloroform (CH₃CCl₃) as additional tests for different components of the transport model. Radon (²²²Rn) is emitted reasonably uniformly in both space and time from land surfaces at much higher rates than from oceans and decays radioactively with a half-life of 3.8 days. Due to its short lifetime, radon is often used as a tracer of recent contact with land and to explore vertical mixing through the lower

atmosphere (Zahorowski et al., 2004; Williams et al., 2011, 2013; Chambers et al., 2011). Since radon fluxes are usually assumed to be better characterized than fluxes of other trace gases, radon has been used, via tracer ratio methods, to estimate regional carbon dioxide, methane and nitrous oxide or other greenhouse gas fluxes (Schmidt et al., 1996; 5 Wilson et al., 1997; Biraud et al., 2000; Zahorowski et al., 2004; Wada et al., 2013).

An Australian methane budget was described by Wang and Bentley (2002), who used 1997 Cape Grim atmospheric methane measurements in an inversion to constrain the methane fluxes from southern Australia. Their study suggested that the methane inventory overestimated southeastern Australian fluxes. A more recent modelling study by 10 Fraser et al. (2011) ran forward model simulations for 2005–2008 and separated methane into different regional and process-based components, focusing on better parameterising methane fluxes from seasonal wetlands in tropical Australia. At Cape Grim, they found their simulated methane was dominated by animal, landfill and ocean fluxes. They also noted that their model reproduced background, baseline methane concentrations at Cape Grim well, but the model was less successful in reproducing concentrations influenced by local 15 emissions. The concentration excursions driven by local emissions are typically large in magnitude and are referred to as “non-baseline” events. These events contain information regarding regional fluxes and are the focus of this study.

To investigate non-baseline events, this study uses both observations at Cape Grim and 20 forward model simulations with prescribed emissions. Sections 2 and 3 describe the observations at Cape Grim and the model simulations, respectively. Section 4 focuses on seasonal scale results, exploring an apparent anomaly between modelled and observed methane in austral spring. The implications for regional methane fluxes are discussed in Sect. 5.

25 **2 Observations**

Cape Grim is located at the top of a 90 m cliff on the north-west coast of Tasmania (40.7° S, 144.7° E), which is separated from mainland Australia by Bass Strait (Fig. 1).

~~Cape Grim~~ The Cape Grim station has been operating since the 1970s and now has the most comprehensive monitoring program in the Southern Hemisphere for greenhouse gases (Langenfelds et al., 2014), ozone-depleting gases (Krummel et al., 2014), and radon (Zahorowski et al., 2014).

5 Figure 2 shows methane and radon observations made at Cape Grim during ~~2003-2006~~. Observations made at Cape Grim are characterised by baseline periods when the wind is from the south-west, (typically 30 % of the time), and air parcels have had long trajectories over the Southern Ocean. During these periods, concentrations are reasonably steady displaying underlying seasonal variations and long-term trends. Baseline periods are punctuated by non-baseline events characterised by higher concentrations that vary rapidly in time, 10 indicative of air parcel trajectories that have travelled over mainland Australia or Tasmania, and consequently have been influenced by regional and local emissions. The seasonal cycles in both methane and radon are apparent in Fig. 2a and b which shows a full year of data. ~~(Unfortunately, there are no radon data available for the month of April in 2003.)~~ In 15 Fig. 2c and d, showing just one month of data, the ~~clear~~ difference between baseline periods and non-baseline events can be more clearly seen and one gets a sense of the degree to which methane and radon are correlated.

Flask measurements of CH₄ began at Cape Grim in 1984, while the record from the current AGAGE GC-MD (multi-gas chromatograph, multi-detector) in situ instrument system, 20 which incorporates a Carle GC fitted with a flame ionisation detector (see Prinn et al., 2000) began late in 1993. Ambient methane measurements are made on discrete air samples every 40 min, taken alternately from a 75 and 10 m inlet for the majority of the study period. The data from both inlets are used in this study. Ambient samples are bracketed by analysis of a calibration standard, and the resulting CH₄ record is reported on the Tohoku University 25 scale (Aoki et al., 1992; Prinn et al., 2000).

During the period of observations used for this study, Cape Grim radon measurements were made using a number of detectors. From 1994 to 1997, a 9000 L two filter radon detector featuring a particle generator was used, operating at a nominal flow rate of 200 L min⁻¹ and with a response time of approximately 90 min (Whittlestone and Zahorowski, 1995).

From 1997, a newly-designed 5000 L dual flow loop, two filter radon detector was commissioned, operating at a nominal flow rate of 285 L min^{-1} and with a response time of 45 min (Whittlestone and Zahorowski, 1998). Later developments saw this new detector enhanced from a single-head design to two and eventually four heads in 2004, with corresponding improvements to sensitivity and lower limit of detection. Air was sampled from a 75 m inlet, the same height as the upper CH_4 observations. Raw radon counts were collected half-hourly and aggregated to hourly values during post processing. Detector sensitivity ranged from 0.6 to 1.2 counts per second per Bq m^{-3} during the period of measurements. Calibrations were performed monthly using a Pylon flow-through radon source ($20.9 \pm 0.8 \text{ kBq Radium-226}$), traceable to US National Institute of Standards and Technology (NIST) standards, and instrumental background checks were performed approximately every 3 months. The lower limit of detection for the Cape Grim radon detectors, defined as the radon concentration below which the statistical counting error exceeds 30 %, ranged from 6 to 10 mBq m^{-3} .

For this study, CH_4 observations from 1994–2007 have been processed using the following steps:

1. We have linearly interpolated between the discrete measurements of atmospheric CH_4 every 40 min, to generate hourly CH_4 data, to facilitate comparisons with the hourly radon data, and the hourly values from the model simulations.
2. The CH_4 observations were then selected for baseline conditions by excluding all hours when the coincident radon measurement was greater than 100 mBq m^{-3} .
3. A smooth curve was then found through these baseline-selected CH_4 observations using the methodology described in Thoning et al. (1989). Specifically, the baseline data are fitted with a function consisting of a second order polynomial and four harmonics. This function fit is then subtracted from the baseline data and the resulting residuals are then filtered with a band-pass filter with a short term cut-off of 80 days. The original function fit is then added back to the filtered residuals to give a smooth curve fit through the data. This procedure is performed iteratively, and in each itera-

tion, the individual hours that lie outside twice the standard deviation around the fit are excluded until the fit converges.

4. Lastly, the smooth curve fitted to the baseline data is subtracted from all CH₄ observations to give a timeseries of residuals. In most cases it is only the residuals from hours that have been flagged as non-baseline that are used for comparison with the model simulations.

3 Model simulations

This study uses simulation experiments that were run for the TransCom-CH₄ model intercomparison to investigate various methane flux estimates. The TransCom-CH₄ model intercomparison involved running nine tracers in a global atmospheric model for the years 1988–2007. The first six tracers used different methane emission scenarios. The remaining three tracers were radon, sulfur hexafluoride and methyl chloroform. In each methane case, chemical loss of methane was simulated using prescribed OH fields (with seasonal variations but no interannual variations) and prescribed loss rates to represent photolysis in the stratosphere. The emissions are described in detail and a global analysis of the results is presented in Patra et al. (2011). Other details can also be found online in the TransCom-CH₄ protocol (Patra et al., 2010).

3.1 Methane emission scenarios

The six methane scenarios were created by combining various estimates of anthropogenic, rice, biomass burning and wetland components in different ways (Table 1). Details are given in Patra et al. (2011) and we use the same emission scenario labels as used in that paper and across the TransCom-CH₄ project. The control (CTL) scenario uses anthropogenic fluxes as specified in the Emissions Database for Global Atmospheric Research (EDGAR) inventory, version 3.2 (Olivier and Berdowski, 2001) and includes fossil fuel, industrial, animal, fire, waste and biofuel emissions. Added to these fluxes are ~~the flux estimates of~~

seasonally-varying (no interannual variability) natural fluxes comprising biomass burning from ~~the Global Fire Emissions Database (GFED), version 2~~, Fung et al. (1991), wetland emissions (Matthews and Fung, 1987; Fung et al., 1991) and rice (Yan et al., 2009).

Four alternative emissions scenarios change one or more of the CTL component fluxes. CTL_E4 uses EDGAR 4.0 (van Aardenne et al., 2001) for the anthropogenic component: BB (biomass burning) uses ~~interannually-varying~~ biomass burning emissions ~~when available from the Global Fire Emissions Database, version 2~~ (van der Werf et al., 2006) (including interannual variations when available); WLBB (wetland and biomass burning) additionally includes interannually varying wetland emissions (Ringeval et al., 2010); EXTRA uses the same biomass burning as BB and interannually-varying model generated wetlands ~~(as per WLBB)~~ and rice emissions from the VISIT model (Ito and Inatomi, 2012). A final emissions scenario, INV, does not use fluxes from inventories or process models but those estimated by the atmospheric inversion of Bousquet et al. (2006). All six emissions scenarios use the same soil sink. Table 2 details which components of each scenario include interannual variability and in which years of the simulations.

Figure 3a shows the aggregated emissions for the region of southeastern Australia bounded by 135–155° E and 45–30° S which is the region shown in Fig. 1. The emissions show a seasonal cycle that is dominated by the wetland component for CTL, CTL_E4 and BB with higher emissions from May to October. The seasonality is much smaller for the WLBB and EXTRA scenarios with maximum fluxes in December and January due to biomass burning. The INV emissions show similar seasonality to the CTL emissions, though with smaller amplitude. It is worth noting that the inversion used to generate these fluxes included only baseline CH₄ data at Cape Grim, so there is no reason to expect the INV fluxes to fit the non-baseline record at Cape Grim better than the other flux scenarios.

Figure 3b shows the interannual variability of the prescribed emissions, again aggregated over the region shown in Fig. 1. The CTL fluxes show almost no change in time, while the CTL_E4 fluxes increase over time. The three fluxes that include interannually varying biomass burning; BB, WLBB and EXTRA, show peaks in 2003 and 2006 associated with

significant summer fires in southeastern Australia. This will be discussed in more detail in Sect. 4.1. The INV fluxes show the greatest interannual variability in this region.

The prescribed TransCom-CH₄ methane emissions can be compared with those reported in Wang and Bentley (2002) and Fraser et al. (2011) (Table 3). To approximate our region of interest, from Wang and Bentley (2002) we sum their regions A, C and D, which extend slightly further west and north than our region. This gives total anthropogenic emissions (agriculture including cattle, the energy and transport sectors and waste management) of 3.39 Tg y⁻¹, using the methodology of the Australian National Greenhouse Gas Inventory (NGGI) coupled with statistical data for 1997 to give a spatially explicit representation of methane emissions. By adding in an estimate of methane uptake by Australian soils, the net anthropogenic flux used in Wang and Bentley (2002) across southeastern Australia is 3.09 Tg y⁻¹. Wang and Bentley (2002) also adjust their inventory-based estimate by fitting atmospheric CH₄ at Cape Grim using an inversion technique. This gives a substantially lower flux estimate (1.93 Tg y⁻¹).

From Fraser et al. (2011) we sum their anthropogenic emissions (agriculture including cattle (and rice, which is very small), the energy sector, waste management and a small amount of prescribed burning) from five regions; New South Wales, Australian Capital Territory, Victoria, Tasmania and South Australia, to give total anthropogenic emissions of 2.88 Tg y⁻¹ for 2008. Fraser et al. (2011) take emissions from the EDGAR 3.2 inventory (Olivier et al., 2005), and scale them to the Australian NGGI. Both inventories have total annual CH₄ emissions which are closer to the annual emissions of the lower set of methane scenarios used here, but the inventories do not include natural fluxes while the TransCom methane scenarios do. In southeastern Australia, the major natural flux is from wetlands, but the magnitude of this flux is uncertain. The CTL based emission scenarios (CTL, CTL_E4 and BB) include a large 1.24 Tg y⁻¹ component from wetlands for southeastern Australia, taken from Matthews and Fung (1987) and Fung et al. (1991). The WLBB and EXTRA emission scenarios take their estimates of wetland emissions from Ringeval et al. (2010) , which and Ito and Inatomi (2012), which are close to zero for southeastern Aus-

tralia, ~~are much lower estimates than those of and~~. Wetland emissions will be discussed further in Sects. 4.2 and 5.

3.2 Atmospheric models

We have run the TransCom-CH₄ simulations with two models: the CSIRO Conformal-Cubic Atmospheric Model (CCAM) (McGregor, 2005; McGregor and Dix, 2008) and the Australian Community Climate and Earth System Simulator (ACCESS) (Corbin and Law, 2011). ~~CCAM~~ Using two models allows us to better understand any sensitivity in the analysis to model transport error.

The first model, CCAM, has an approximately uniformly spaced conformal cubic grid and was run with a horizontal resolution of approximately 220 km, and 18 levels in the vertical. The horizontal components of the wind were nudged (Thatcher and McGregor, 2009) to NCEP analyses (Kalnay et al., 1996; Collier, 2004). This helps to ensure that simulated atmospheric concentrations of trace gases can be more realistically compared to observations on synoptic timescales. The CCAM simulations analysed here are the same as those submitted to the TransCom-CH₄ experiment (Patra et al., 2011).

The second model, ACCESS is derived from the UK Met Office Unified Model but has the land surface scheme replaced by the Community Atmosphere Biosphere Land Exchange (CABLE) model. For this study, ACCESS was run at 1.875° longitude by 1.25° latitude, with 38 levels in the vertical. ~~Note that the~~ This is higher horizontal resolution (to better represent the region around Cape Grim) than the ACCESS case submitted to the TransCom-CH₄ ~~archive experiment, which~~ was run at ~~lower resolution to meet the TransCom deadlines.~~ 3.75° longitude by 2.5° latitude. In both cases, ACCESS was run without any nudging to analysed winds or temperature, so that the ~~only forcing is through~~ tracer transport is dependent on the ACCESS simulation of meteorological fields forced with observed monthly sea surface temperatures. Consequently the output from this model ~~can only be compared to observations on~~ is not expected to reproduce observed day to day variations in methane concentration and comparison with the observations is limited to seasonal or

longer timescales, ~~or by analysing the statistics of synoptic behaviour~~. The ACCESS run used a 360 day calendar with 12 months of equal length.

It is important to consider where the model output is sampled to be most comparable with Cape Grim observations, as well as to minimise differences between the two model simulations. The model sampling locations are shown in Fig. 1. The inset in Fig. 1 shows the spatial extent of each of the grid cells chosen to represent Cape Grim, giving a sense of their relative size. For both models we have sampled slightly to the north of Cape Grim. For CCAM, this grid cell was chosen as it is the nearest ocean grid point to the location of Cape Grim. For ACCESS, the grid cell was chosen based on a radon simulation; the grid cell to the north of Cape Grim gave a better simulated seasonal cycle amplitude for radon than grid cells to the south or west. It is worth noting that in CCAM grid-cells are either all land or all ocean, whereas in ACCESS fractional land area is allowed.

Model timeseries are output hourly. The simulated concentrations are processed in the same manner as for the observations. Firstly coincident radon concentrations are used to select for baseline CH₄, a smooth curve is fitted to the baseline data, and the baseline fit is removed from the timeseries. The residual concentrations are used for comparison with the observations.

4 Results

Initial analysis of the simulated CH₄ at Cape Grim highlighted two features. The first feature was two periods ~~of 1–2 months duration~~, December 2002 to February 2003 and November 2006 to January 2007, with very high peak CH₄ concentrations for the flux scenarios that included interannally varying biomass burning. The second feature was a large difference in CH₄ concentrations in each winter between different flux scenarios.

As an example of these features, Fig. Figure 4 shows the monthly mean methane residuals in 2003 for for three years (2002-2004) for the observations and all emission scenarios for both models. Both features mentioned above are most obvious in the CCAM results (Fig. 4b), where the model predicts exceptionally high methane concentrations in January

and February 2003 for those tracers that include interannually varying biomass burning (BB, WLBB and EXTRA). Over the winter months, the three emission scenarios with default wetland emissions, (CTL, CTL_E4 and BB) all produce ~~markedly~~ higher methane concentrations than the observations or the remaining emission scenarios using modified wetland emissions ~~or those (WLBB and EXTRA) or~~ based on the inversion of Bousquet et al. (2006) (INV).

4.1 Biomass burning

Figure 5 shows observed and CCAM simulated CH₄ residuals for January 2003 as well as observed hydrogen (H₂) and carbon monoxide (CO). High H₂ and CO are signatures of air influenced by biomass burning. Two simulated CH₄ tracers are shown, one which includes interannually varying biomass burning emissions (WLBB) and one that does not (CTL). The inclusion of “hot spots” of biomass burning emissions produces very large methane concentrations, much larger than those observed. Significant fires did occur during this period; in the eastern Victorian alpine region starting on 8 January 2003 and burning around 1.3 million hectares over close to two months (<http://www.depi.vic.gov.au/fire-and-emergencies/managing-risk-and-learning-about-managing-fire/bushfire-history/maps-of-past-bushfires>) and around Canberra between 18 and 21 January 2003. The observations indicate that the Victorian fire was likely seen briefly at Cape Grim on 11 January, when CH₄, H₂ and CO all had elevated concentrations. It is less clear what contribution biomass burning makes to other elevated methane events later in the month, when only small CO elevations are seen and H₂ signals do not rise above the instrumental noise (except perhaps around 25 January).

There are a number of reasons why the models may overestimate the impact of this fire at Cape Grim. Firstly, the biomass burning emissions are specified at the middle of each month and interpolated to the middle of the previous and following months. This means that a January fire is spread temporally into December and February. Indeed, WLBB also shows very large CH₄ concentrations in December 2002 and February 2003 (not shown). Secondly, the fire emissions were provided on a 1° × 1° grid and have been regridded to the

lower resolutions of the atmospheric models. In reality, the active fire at any given time would have covered a much smaller area. Finally, the fire itself would modify the local circulation, with emissions likely distributed not just near the surface but throughout the entire lower troposphere. This is not captured in our simulations where the emissions are input only to the lowest model level. For instance, Sofiev et al. (2013), find that under Australian fire conditions, 90 % of mass is emitted from the surface up to 3 km altitude.

~~We~~

To test the sensitivity to the height that emissions are input to the atmosphere, we have performed some short tests, running CCAM from December 2002 to February 2003, with only biomass burning emissions ~~and releasing them into different model levels or spreading them through a vertical column~~. Six tests were performed, releasing the emissions into model levels 1, 3, 5, 7, and 9 in turn (centred at approximately 40, 470, 1420, 2880, 4870 m respectively), or distributed through all levels 1-9. In general we find ~~the same that the simulated timing of~~ elevated CH₄ events ~~simulated~~ at Cape Grim ~~but with smaller magnitude as the emissions are inserted is similar across the six tests but the amplitude of the events varies with emission insertion height.~~ For example, the mean January CH₄ concentration at Cape Grim is reduced from 16 ppb for emissions inserted into level 1 to 12, 7, 2 and 0.3 ppb as emissions are input higher into the atmosphere. For the case where emissions are spread between levels 1 and 9, the mean January CH₄ concentration is 5 ppb. While these simulations (modelling only the biomass burning component) are not directly comparable to the observations, they clearly illustrate one reason why the amplitude of events at Cape Grim could be overestimated. The frequency of events appears to be more strongly controlled by the temporal and spatial (horizontal) resolution of the emissions.

4.2 Relationship between methane and radon: seasonal cycle

Given the variation in simulated winter CH₄ concentrations from the different flux scenarios, we would like to determine which gives a better comparison with the observations. To avoid biasing our analysis of the seasonality of methane by the very large concentrations

caused by the interannually varying biomass burning fluxes, we remove the biomass burning contribution from BB, WLBB and EXTRA for the months December 2002 to February 2003 and also November 2006 to January 2007, when another large fire gives unrealistically high simulated concentrations of CH₄ at Cape Grim. This was achieved by subtracting (BB-CTL) from each of the three affected tracers, during the months in question.

To minimise the impact of any errors in modelled atmospheric transport, we consider the ratio of methane residual concentrations to radon concentrations. The timeseries of methane residuals at Cape Grim is reasonably well correlated with radon, with both showing significantly elevated concentrations when air parcels have travelled over continental Australia. For the period 1994–2007, the correlation coefficient between observed methane residuals and observed radon is 0.65. In the model simulations, correlations between residual methane and radon vary across methane scenarios between 0.81–0.87 for ACCESS and 0.88–0.89 for CCAM. The higher correlation coefficients for the modelled data set compared to the observed data set presumably reflect the reduced spatial variability of both CH₄ and radon in the gridded models compared to the real world.

The strategy of using a residual methane to radon ratio to minimise transport errors relies on the assumption that both sources are similarly ~~distrubuted~~distributed. We note that this assumption is only loosely true. The methane emission scenarios we assess here have considerable spatial variability (though doubtless less than the real world), while the TransCom-CH₄ specified radon emissions are uniform over land. Land surface emissions of radon do vary (Griffiths et al., 2010), though much less than for methane. As a check on our sensitivity to radon spatial variability, we ran ACCESS for three years using a series of different radon flux fields taken from Griffiths et al. (2010). In general, neither the spatial variability nor the interannual variability appeared to have a significant impact on our sampled concentrations at Cape Grim. We found that the choice of grid cell was much more influential in the modelled radon results matching the observed radon results, hence our choice of ocean grid cells to the north of Cape Grim. One of the reasons for this is likely to be the coarseness of the grid cells. See Sect. 3.2 and refer to the inset in Fig. 1.

We examine the seasonal relationship between methane residuals and radon by fitting a linear relationship to all hourly non-baseline methane-radon pairs in a given month across the 14 years of the simulation for which we have observational data for comparison (1994–2007). For clarity, Fig. 6 shows this fit for a single month only, January 2006, for observations and modelled CTL cases. We take this approach in order to compare the observations to both the CCAM and ACCESS simulations. At the time of this work, ACCESS meteorology was forced only with sea surface temperatures and ran on a 360 day calendar, as described in Sect. 3. This means that we do not expect the timing of individual “events” (non-baseline periods) in ACCESS to match the observations well enough for a direct comparison. We do however expect that the seasonal scale meteorology will be realistic enough to provide seasonal fetch changes that are comparable to the real meteorology and fetch patterns at Cape Grim. Averaging results up to a seasonal timescale allows direct comparison of all three datasets (the observations and both models). The observations show more scatter than the model output, and this is reflected in lower R^2 values, in this case 0.61 for the observations and 0.92 and 0.74 for CCAM and ACCESS respectively. The slope of the line gives the methane residual to radon ratio. We determine the ratio this way to preserve the temporal pairing of the methane and radon concentrations. We also calculate linear fits for each individual month as per Fig. 6, and use the standard deviation of the slopes for each of the 14 months as a measure of the uncertainty on the methane-radon ratios.

Figure 7 shows the mean seasonal cycle of the methane residual-radon ratio for each methane scenario, compared to observed ratios. The observed ratios are smallest in winter (June to August) and are largest in spring and summer (October to February). The modelled ratios show three patterns depending on the methane scenario. The CTL, CTL_E4 and BB scenarios show maximum ratios in winter, while the WLBB and EXTRA scenarios show minimum ratios in winter, which is more consistent with the observations. The INV scenario is intermediate between the other cases, also showing somewhat elevated winter ratios. For any given scenario, the ratios for the two different models compare well, much better than if either of the individual trace gases (methane or radon) are directly compared, as was the case in Fig. 4 where the monthly mean methane is shown for both models. This illustrates

the benefit of using the radon simulation to account for some of the transport differences between the models.

Overall the WLBB and EXTRA scenarios give ratios that are a better fit to those observed. For the southeastern Australian region, the major difference from the group of CTL-based cases is in the representation of wetlands. These results suggest that the large winter wetland fluxes in the CTL-based scenarios taken from Matthews and Fung (1987) and Fung et al. (1991) are not realistic and annual mean fluxes should be close to the anthropogenic only inventory estimates noted in Sect. 3.1. For all scenarios (except perhaps CTL E4) there is a discrepancy between the observed and modelled ratios in the austral spring (September to November) with the observed ratios being larger than the modelled ones. This suggests that the methane flux scenarios tested here underestimate methane fluxes in spring. We will discuss this further in Sect. 5.

While the ratios from the CTL E4 scenario agree reasonably with those observed in September-October, this occurs mainly because the CTL E4 scenario gives generally higher ratios all year round compared to the CTL case. This is expected since the CTL E4 fluxes for southeastern Australia gradually increase in time relative to the CTL fluxes (Fig. 3b). There is weak evidence of an increase in ratio over time for CTL E4 compared to CTL but the large seasonal and interannual variability means that we can have little confidence in the calculated trends and which might compare better to the observations. The seasonality of ratios is similar between CTL and CTL E4 (and a poor fit to observed seasonality) although the difference between summer and winter ratios is smaller in CTL E4 especially in the ACCESS case. This may be due to changes in the spatial distribution of fluxes between CTL and CTL E4. Overall CTL_4 does not agree well with the observations.

5 Discussion and concluding remarks

Forward modelling of a range of methane flux scenarios gives us the opportunity to compare those results with the measurement record at Cape Grim for the same period, offering

insights into which of the flux scenarios appear to be most representative of southeastern Australian methane fluxes. The WLBB and EXTRA flux scenarios appear to be the best fit to the observed data. Like Fraser et al. (2011), we find that anthropogenic methane emissions taken from inventories for southeastern Australia look quite reasonable in magnitude at around 2.5 Tg y^{-1} . However, there remain questions about the scale and timing of a wetland component to CH_4 emissions.

Our analysis reveals that mismatches in the CTL, CTL_E4 and BB scenarios were due to high wetland emissions during the winter, suggesting that the wintertime wetland flux is overestimated in these scenarios. The Cape Grim observations point to somewhat larger springtime fluxes than are represented in the WLBB or EXTRA emissions scenarios (or indeed any of the other tracers). Although the wintertime maxima of the CTL-style emissions scenarios (driven by a wetland emission component) is clearly not warranted by the observational data, ~~they look to be about the right magnitude to explain the “shortfall” in spring for all the emission scenarios. A shift in the wetland emissions from austral winter to spring might result in a better fit to the observations.~~

To test this we have run CCAM from December 1, 1993 to December 31, 2007 for radon and two methane scenarios, the standard CTL case and with the same CTL fluxes shifted forward in time by three months (across the whole globe). For southeastern Australia, this means that methane fluxes are elevated between August and January instead of between May and October. The output from the simulation has been processed similarly to the previous cases and Fig. 8 shows the seasonal cycle of methane to radon ratio. The case with temporally shifted fluxes clearly fits the observations better than the original CTL fluxes, although a four month shift in the fluxes might improve the comparison with observations further. The shifted fluxes case does not simulate a low enough methane minimum in winter. This may be a consequence of shifting the CTL fluxes across the whole globe rather than just for southeastern Australia. For example a large flux just to the north of the region we define occurs in February in the CTL fluxes and consequently gets shifted to May in the sensitivity test performed here. Nevertheless, this simulation supports the need for a shift in wetland emissions from austral winter to spring ~~may be~~ and summer.

A shift of this nature is plausible given that wetland methane emissions have both a soil moisture and soil temperature dependence, making it possible that southeastern Australian methane emissions from wetlands are highest in springtime when there is available moisture and warmer temperatures. Indeed, Bloom et al. (2012) use satellite column observations of CH₄ from the SCanning Imaging Absorption spectroMeter for Atmospheric CHar-
5 tographyY (SCIAMACHY) coupled with a measure of equivalent water height from the Gravity Recovery and Climate Experiment (GRACE) to model seasonal variability in wetland methane emissions. For our region of interest across the years 2003–2008, they find the minimum in methane emissions from southeastern Australian wetlands occurs in late au-
10 tumn and winter with a rapid rise through spring giving a maximum in October and November, (A. Fraser, personal communication, 2012) in accordance with the Cape Grim observations. However, the magnitude they predict for this *wetland* flux is around 2.5 Tg y⁻¹. This seems larger than indicated by the Cape Grim data; of the six emission scenarios considered in this work, annual means for *total* CH₄ flux (including all anthropogenic emissions)
15 in our defined region range between 2.4 (WLBB) and 4 (CTL) Tg y⁻¹ (Fig. 3b). Moreover, the additional 1.6 Tg y⁻¹ in the CTL emission scenario comes from a wintertime wetland flux that we find no evidence for in the observations. It should be noted however that in Fraser et al. (2011) the GRACE data used were scaled to match a prior emissions estimate. We therefore find that the GRACE data have plausible seasonality for southeastern
20 Australia but that the magnitude is too large to offer a realistic assessment of the scale of southeastern Australian wetland emissions given the Cape Grim observations. Nevertheless, we believe the seasonality of the GRACE data lends credibility to the idea that a springtime wetland emission of around the magnitude represented in the CTL-style emissions scenarios for winter may be responsible for the discrepancy between our modelled
25 WLBB/EXTRA results and the observations.

Although our hypothesis that the timing of wetland methane emissions in the inventories may be off by ~~2–3~~3–4 months is plausible and supported by other data, we cannot entirely rule out another source for the additional austral spring methane emissions. For instance, ruminant emissions from cattle are the single biggest contributor to Australia's

anthropogenic methane emissions. Seasonality in ruminant emissions that is not captured by the inventories, due to changes in feed or cattle number, might also be responsible for the austral springtime maximum observed. This explanation is offered in Wang and Bentley (2002) to account for ~~the spike in emissions from their region D (a springtime spike in their estimated emissions from a region~~ roughly equivalent to Victoria and ~~NSW) during the springtime. However, although New South Wales (their region D) when fitting the Cape Grim CH₄ data. While this would concur with our results, overall~~ Wang and Bentley find ~~an austral springtime peak in that coincides with the timing of the maximum noted in this work, they find overall~~ a significant reduction ~~in the (around 40%) in the inversion-estimated CH₄ inventory fluxes~~ for southeastern Australia ~~is required to reconcile it to the Cape Grim observations for 1997. The reduction proposed by Wang and Bentley of around 40% in the largest region, D is at odds with~~ compared to the 1997 inventory (Table 3). ~~This does not agree with this work or with~~ Fraser et al. (2011) ~~and this work~~, which both suggest that the total methane emissions in the inventories for southeastern Australia are consistent with the Cape Grim data. The methodology in Wang and Bentley (2002) is to invert a series of 22 individual non-baseline “events” each lasting between 2 and 11 days. The inversion results for each event show considerable variability ~~giving fluxes ranging from 0-7 Tg y⁻¹ for region D. Such high variability may be indicative of transport error would appear to be unrealistic and may be caused by errors~~ in the modeling ~~and thus flux estimates that are unrepresentative~~ of CH₄ ~~concentration at Cape Grim for a given Australian flux. Thus the inversion estimated fluxes are unlikely to be representative of~~ the true flux.

This study used 14 years of observational data, and output from two models (CCAM and ACCESS) to investigate methane emissions data for southeastern Australia. Inclusion of biomass burning emissions produces unrealistically high CH₄ concentrations at Cape Grim, but this is most likely due to the coarse spatio-temporal resolution of the models and the unrealistic injection of these emissions into the lowest model level. In future, continuous CH₄ measurements made from unmanned aerial vehicles (UAVs) around large fire plumes may be a better way to verify the scale of emissions from large biomass burning events. By comparing a range of methane emission scenarios run in the models, we find that the

large wintertime wetland flux in the CTL-style scenarios is unrealistic, but also that there is a deficit in spring in all six emission scenarios. This deficit is present in even the WLBB and EXTRA scenarios which otherwise provide a good fit to the observational data. It is notable that these two emission scenarios have a very small wetland emission component compared to the CTL-style scenarios. We suggest then, that it may be a springtime wetland emission that is missing from these scenarios. Finally, we note that given the size and uncertainty associated with the biogenic CH₄ fluxes, it is difficult to make assessments about changes in anthropogenic CH₄ emissions in southeastern Australia from the Cape Grim data set alone using this approach. Additional in situ instrumented sites for the continuous measurement of methane on the Australian mainland would help to answer questions about the scale and timing of wetland emissions as well as providing more stringent constraints on changes to the anthropogenic flux from southeastern Australia.

Acknowledgements. We thank AGAGE and Cape Grim staff, along with all the TransCom-CH₄ participants, but in particular Prabir Patra for his efforts in setting up the experiment. This work has been undertaken as part of the Australian Climate Change Science Program, funded jointly by the Department of the Environment, the Bureau of Meteorology and CSIRO. Model simulations were undertaken on the NCI National Facility in Canberra, Australia, which is supported by the Australian Commonwealth Government. [We thank the reviewers for their helpful comments.](#)

References

- Aoki, S., Nakazawa, T., Murayama, S., and Kawaguchi, S.: Measurements of atmospheric methane at the Japanese Antarctic station, Syowa, *Tellus B*, 44, 273–281, 1992.
- Biraud, S., Ciais, P., Ramonet, M., Simmonds, P., Kazan, V., Monfray, P., O'Doherty, S., Spain, T. G., and Jennings, S. G.: European greenhouse gas emissions estimated from continuous atmospheric measurements and radon 222 at Mace Head, Ireland, *J. Geophys. Res.*, 105, 1351–1366, 2000.
- Bloom, A. A., Palmer, P. I., Fraser, A., and Reay, D. S.: Seasonal variability of tropical wetland CH₄ emissions: the role of the methanogen-available carbon pool, *Biogeosciences*, 9, 2821–2830, doi:10.5194/bg-9-2821-2012, 2012.

- 5 Bousquet, P., Ciais, P., Miller, J. B., Dlugokencky, E. J., Hauglustaine, D. A., Prigent, C., Van der Werf, G. R., Peylin, P., Brunke, E. G., Carouge, C., Langenfelds, R. L., Lathière, J., Papa, F., Ramonet, M., Schmidt, M., Steele, L. P., Tyler, S. C., and White, J.: Contribution of anthropogenic and natural sources to atmospheric methane variability, *Nature*, 443, 439–443, doi:10.1038/nature05132, 2006.
- 10 Bousquet, P., Ringeval, B., Pison, I., Dlugokencky, E. J., Brunke, E.-G., Carouge, C., Chevallier, F., Fortems-Cheiney, A., Frankenberg, C., Hauglustaine, D. A., Krummel, P. B., Langenfelds, R. L., Ramonet, M., Schmidt, M., Steele, L. P., Szopa, S., Yver, C., Viovy, N., and Ciais, P.: Source attribution of the changes in atmospheric methane for 2006–2008, *Atmos. Chem. Phys.*, 11, 3689–3700, doi:10.5194/acp-11-3689-2011, 2011.
- 15 Chambers, S., Williams, A. G., Zadorowski, W., Griffiths, A., and Crawford, J.: Separating remote fetch and local mixing influences on vertical radon measurements in the lower atmosphere, *Tellus B*, 63, 843–859, doi:10.1111/j.1600-0889.2011.00565.x, 2011.
- Collier, M. A.: The CSIRO NCEP/NCAR/DOE *R-1/R-2* archive, CSIRO Atmospheric Research technical paper 68, Aspendale, Victoria, Australia, available at: http://www.cmar.csiro.au/e-print/open/collier_2004a.pdf, 2004.
- 20 Corbin, K. D. and Law, R. M.: Extending atmospheric CO₂ and tracer capabilities in ACCESS, CAWCR Technical Report 035, Aspendale, Victoria, Australia, available at: http://www.cawcr.gov.au/publications/technicalreports/CTR_035.pdf, 2011.
- Cunnold, D., Steele, L., Fraser, P., Simmonds, P., Prinn, R., Weiss, R., Porter, L., O'Doherty, S., Langenfelds, R., Krummel, P., Wang, H., Emmons, L., Tie, X., and Dlugokencky, E.: In situ measurements of atmospheric methane at GAGE/AGAGE sites during 1985–2000 and resulting source inferences, *J. Geophys. Res.-Atmos.*, 107, 4225, doi:10.1029/2001JD001226, 2002.
- 25 Dlugokencky, E. J., Bruhwiler, L., White, J. W. C., Emmons, L. K., Novelli, P. C., Montzka, S. A., Masarie, K. A., Lang, P. M., Crotwell, A. M., Miller, J. B., and Gatti, L. V.: Observational constraints on recent increases in the atmospheric CH₄ burden, *Geophys. Res. Lett.*, 36, L18803, doi:10.1029/2009GL039780, 2009.
- Fraser, A., Chan Miller, C., Palmer, P. I., Deutscher, N. M., Jones, N. B., and Griffith, D. W. T.: The Australian methane budget: interpreting surface and train-borne measurements using a chemistry transport model, *J. Geophys. Res.*, 116, D20306, doi:10.1029/2011JD015964, 2011.
- 30 Fung, I., John, J., Lerner, J., Matthews, E., Prather, M., Steele, L. P., and Fraser, P. J.: Three-dimensional model synthesis of the global methane cycle, *J. Geophys. Res.*, 96, 13033–13065, 1991.

- Griffiths, A. D., Zahorowski, W., Element, A., and Werczynski, S.: A map of radon flux at the Australian land surface, *Atmos. Chem. Phys.*, 10, 8969–8982, doi:10.5194/acp-10-8969-2010, 2010.
- Houweling, S., Kaminski, T., Dentener, F., Lelieveld, J., and Heimann, M.: Inverse modeling of methane sources and sinks using the adjoint of a global transport model, *J. Geophys. Res.*, 104, 26137–26160, 1999.
- Ito, A. and Inatomi, M.: Use of a process-based model for assessing the methane budgets of global terrestrial ecosystems and evaluation of uncertainty, *Biogeosciences*, 9, 759–773, doi:10.5194/bg-9-759-2012, 2012.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, 1996.
- Kirschke, S., Bousquet, P., Ciais, P., Saunoy, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J.-F., Langenfelds, R. L., Le Quérec, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/NGEO1955, 2013.
- Krummel, P., Fraser, P., Steele, L., Derek, N., Rickard, C., Ward, J., Somerville, N. T., Cleland, S. J., Dunse, B., Langenfelds, R., Baly, S. B., and Leist, M.: The AGAGE in situ program for non-CO₂ greenhouse gases at Cape Grim, 2009–2010, in: *Baseline Atmospheric Program (Australia), 2009–2010*, edited by: Derek, N., Krummel, P. B., and Cleland, S. J., Australian Bureau of Meteorology and CSIRO Marine and Atmospheric Research, Aspendale, Victoria, Australia, 56–70, 2014.
- Langenfelds, R., Steele, L., Gregory, R. L., Krummel, P., Spencer, D., and Howden, R.: Atmospheric methane, carbon dioxide, hydrogen, carbon monoxide and nitrous oxide from Cape Grim flask air samples analysed by gas chromatography, in: *Baseline Atmospheric Program (Australia), 2009–2010*, edited by: Derek, N., Krummel, P. B., and Cleland, S. J., Australian Bureau of Meteorology and CSIRO Marine and Atmospheric Research, Aspendale, Victoria, Australia, 45–49, 2014.

- MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., van Ommen, T., Smith, A., and Elkins, J.: Law dome CO₂, CH₄ and N₂O ice core records extended to 2000 years BP, *Geophys. Res. Lett.*, 33, L14810, doi:10.1029/2006GL026152, 2006.
- 5 Matthews, E. and Fung, I.: Methane emission from natural wetlands: global distribution, area, and environmental characteristics of sources, *Global Biogeochem. Cy.*, 1, 61–86, doi:10.1029/GB001i001p00061, 1987.
- McGregor, J. L.: C-CAM: Geometric aspects and dynamical formulation, CSIRO Atmospheric Research Technical Paper 70, Aspendale, Victoria, Australia, available at: http://www.cmar.csiro.au/e-print/open/mcgregor_2005a.pdf, 2005.
- 10 McGregor, J. L. and Dix, M. R.: An updated description of the Conformal Cubic Atmospheric Model, in: *High Resolution Numerical Modelling of the Atmosphere and Ocean*, edited by: Hamilton, K. and Ohfuchi, W., Springer, Berlin, 51–76, 2008.
- Miller, S. M., Wofsy, S. C., Michalak, A. M., Kort, E. A., Andrews, A. E., Biraud, S. C., Dlugokencky, E. J., Eluszkiewicz, J., Fischer, M. L., Janssens-Maenhout, G., Miller, B. R., Miller, J. B., 15 Montzka, S. A., Nehrkorn, T., and Sweeney, C.: Anthropogenic emissions of methane in the United States, *P. Natl. Acad. Sci. USA*, 110, 20018–20022, doi:10.1073/pnas.1314392110, 2013.
- Olivier, J. G. J. and Berdowski, J. J. M.: Global emissions sources and sinks, in: *The Climate System*, edited by: Berdowski, J. J. M., Guicherit, R., and Heij, B. J., A. A. Balkema Publishers/Swets & Zeitlinger Publishers, Lisse, The Netherlands, 33–78, 2001.
- 20 Olivier, J. G. J., van Aardenne, J. A., Dentener, F., Ganzeveld, L., and Peters, J. A. H. W.: Recent trends in global greenhouse gas emissions: regional trends and spatial distribution of key sources, in: *Non-CO₂ Greenhouse Gases (NCGG-4)*, edited by: van Amstel, A., Millpress, Rotterdam, 325–330, 2005.
- Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Bruhwiler, L., and Jacob, D.: Protocol for TransCom CH₄ intercomparison, Version 7, available at: http://transcom.project.asu.edu/pdf/transcom/T4.methane.protocol_v7.pdf (last access: August 2014), 2010.
- 25 Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Corbin, K., Fortems-Cheiney, A., Fraser, A., Gloor, E., Hess, P., Ito, A., Kawa, S. R., Law, R. M., Loh, Z., Maksyutov, S., Meng, L., Palmer, P. I., Prinn, R. G., Rigby, M., Saito, R., and Wilson, C.: TransCom model simulations of CH₄ and related species: linking transport, surface flux and chemical loss with CH₄ variability in the troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 11, 12813–12837, doi:10.5194/acp-11-12813-2011, 2011.
- 30

- Prinn, R., Weiss, R., Fraser, P., Simmonds, P., Cunnold, D., Alyea, F., O'Doherty, S., Salameh, P., Miller, B., Huang, J., Wang, R., Hartley, D., Harth, C., Steele, L., Sturrock, G., Midgley, P., and McCulloch, A.: A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *J. Geophys. Res.-Atmos.*, 105, 17751–17792, doi:10.1029/2000JD900141, 2000.
- 5 Rigby, M., Prinn, R. G., Fraser, P. J., Simmonds, P. G., Langenfelds, R. L., Huang, J., Cunnold, D. M., Steele, L. P., Krummel, P. B., Weiss, R. F., O'Doherty, S., Salameh, P. K., Wang, H. J., Harth, C. M., Muehle, J., and Porter, L. W.: Renewed growth of atmospheric methane, *Geophys. Res. Lett.*, 35, L22805, doi:10.1029/2008GL036037, 2008.
- 10 Ringeval, B., de Noblet-Ducoudré, N., Ciais, P., Bousquet, P., Prigent, C., Papa, F., and Rossow, W. B.: An attempt to quantify the impact of changes in wetland extent on methane emissions on the seasonal and interannual time scales, *Global Biogeochem. Cy.*, 24, GB2003, doi:10.1029/2008GB003354, 2010.
- Schmidt, M., Graul, R., Sartorius, H., and Levin, I.: Carbon dioxide and methane in continental Europe: a climatology, and ²²²Radon-based emission estimates, *Tellus B*, 48, 457–473, 1996.
- 15 Sofiev, M., Vankevich, R., Ermakova, T., and Hakkarainen, J.: Global mapping of maximum emission heights and resulting vertical profiles of wildfire emissions, *Atmos. Chem. Phys.*, 13, 7039–7052, doi:10.5194/acp-13-7039-2013, 2013.
- Sussmann, R., Forster, F., Rettinger, M., and Bousquet, P.: Renewed methane increase for five years (2007–2011) observed by solar FTIR spectrometry, *Atmos. Chem. Phys.*, 12, 4885–4891, doi:10.5194/acp-12-4885-2012, 2012.
- Thatcher, M. J. and McGregor, J. L.: Using a scale-selective filter for dynamical downscaling with the Conformal Cubic Atmospheric Model, *Mon. Weather Rev.*, 137, 1742–1752, 2009.
- Thoning, K. W., Tans, P. P., and Komhyr, W. D.: Atmospheric carbon dioxide at Mauna Loa Observatory, 2, Analysis of the NOAA/GMCC data, 1974–1985, *J. Geophys. Res.*, 94, 8549–8565, 1989.
- van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Klein Goldewijk, C. G. M., and Lelieveld, J.: A $1^\circ \times 1^\circ$ resolution data set of historical anthropogenic trace gas emissions for the period 1890–1990, *J. Geophys. Res.*, 15, 909–928, 2001.
- 30 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

- Wada, A., Matsueda, H., Murayama, S., Taguchi, S., Hirao, S., Yamazawa, H., Moriizumi, J., Tsuboi, K., Niwa, Y., and Sawa, Y.: Quantification of emission estimates of CO₂, CH₄ and CO for East Asia derived from atmospheric radon-222 measurements over the western North Pacific, *Tellus B*, 65, 18037, doi:10.3402/tellusb.v65i0.18037, 2013.
- 5 Wang, Y. P. and Bentley, S. T.: Development of a spatially explicit inventory of methane emissions from Australia and its verification using atmospheric concentration data, *Atmos. Environ.*, 36, 4965–4975, 2002.
- Whittlestone, S. and Zahorowski, W.: The Cape Grim Huge Radon detector, in: *Baseline Atmospheric Program (Australia) 92*, edited by: Dick, A. L. and Fraser, P. J., Australian Bureau of Meteorology and CSIRO Marine and Atmospheric Research, Aspendale, Victoria, Australia, 26–30, 10
1995.
- Whittlestone, S. and Zahorowski, W.: Baseline radon detectors for shipboard use: development and deployment in the First Aerosol Characterisation experiment (ACE 1), *J. Geophys. Res.*, 103, 16743–16751, 1998.
- 15 Williams, A. G., Zahorowski, W., Chambers, S., Griffiths, A., Hacker, J. M., Element, A., and Werczynski, S.: The vertical distribution of radon in clear and cloudy daytime terrestrial boundary layers, *J. Atmos. Sci.*, 68, 155–174, doi:10.1175/2010JAS3576.1, 2011.
- Williams, A. G., Chambers, S., and Griffiths, A.: Bulk mixing and decoupling of the nocturnal stable boundary layer characterized using a ubiquitous natural tracer, *Bound.-Lay. Meteorol.*, 149, 381–
20 402, doi:10.1007/s10546-013-9849-3, 2013.
- Wilson, S., Dick, A., Fraser, P., and Whittlestone, S.: Nitrous oxide flux estimates for South-Eastern Australia, *J. Atmos. Chem.*, 26, 169–188, 1997.
- Yan, X., Akiyama, H., Yagi, K., and Akimoto, H.: Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Inter-
25 governmental Panel on Climate Change Guidelines, *Global Biogeochem. Cy.*, 23, GB2002, doi:10.1029/2008GB003299, 2009.
- Zahorowski, W., Chambers, S. D., and Henderson-Sellers, A.: Ground based radon-222 observations and their application to atmospheric studies, *J. Environ. Radioact.*, 76, 3–33, 2004.
- Zahorowski, W., Williams, A. G., Chambers, S. D., Crawford, J., and Griffiths, A.: Radon and radon daughters, in: *Baseline Atmospheric Program (Australia), 2009–2010*, edited by: Derek, N. and Krummel, P. B. and Cleland, S. J., Australian Bureau of Meteorology and CSIRO Marine and
30 Atmospheric Research, Aspendale, Victoria, Australia, 33–38, 2014.

Table 1. The broad components of the methane emission scenarios. The anthropogenic category includes fossil fuel, industrial, animal, fire, waste and biofuel emissions. See Patra et al. (2011) for further details of how the emission scenarios were constructed.

Code	Anthropogenic	Rice	Biomass burning	Wetlands
CTL	EDGAR3.2	Yan et al. (2009)	Fung et al. (1991)	Matthews and Fung (1987); Fung et al. (1991)
CTL_E4	EDGAR4.0	Yan et al. (2009)	Fung et al. (1991)	Matthews and Fung (1987); Fung et al. (1991)
BB	EDGAR3.2	Yan et al. (2009)	van der Werf et al. (2006)	Matthews and Fung (1987); Fung et al. (1991)
WLBB	EDGAR3.2	Yan et al. (2009)	van der Werf et al. (2006)	Ringeval et al. (2010)
EXTRA	EDGAR3.2	Ito and Inatomi (2012)	van der Werf et al. (2006)	Ito and Inatomi (2012)
INV			LSCE Inversion	Bousquet et al. (2006)

Table 2. Methane emission scenarios, indicating which components have interannual variations over different periods. A mean seasonal cycle is used outside the listed periods.

Code	Anthropogenic	Rice	Biomass burning	Wetlands
CTL	1990 : 1995 : 2000	No	No	No
CTL_E4	1990–2005	No	No	No
BB	As CTL	No	1996–2008	No
WLBB	As CTL	No	As BB	1994–2000
EXTRA	As CTL	1988–2008	As BB	1988–2008
INV		1988–2005		

Table 3. Annual flux emissions (in Tg y^{-1}) from southeastern Australia for different years and from different sources.

<u>Source</u>	<u>Year</u>	<u>Flux</u>
Wang and Bentley (2002) <u>inventory</u>	<u>1997</u>	<u>3.09</u>
Wang and Bentley (2002) <u>inversion</u>	<u>1997</u>	<u>1.93</u>
Fraser et al. (2011)	<u>2008</u>	<u>2.88</u>
<u>This study</u>	<u>1997</u>	<u>2.38 – 3.82</u>
<u>This study</u>	<u>2008</u>	<u>2.34 – 4.16</u>

Figure 1. Map of South East Australia, showing the location of the Cape Grim Baseline Air Pollution Station, along with the ACCESS (blue) and CCAM (red) grid points selected to best represent Cape Grim. The choice of these grid points is discussed in Sect. 4.2. The inset shows the extent of each of the grid cells.

Figure 2. Cape Grim observations **(a)** ~~2003–2006~~ methane data **(b)** ~~2003–2006~~ radon data **(c)** January ~~2003–2006~~ methane data and **(d)** January ~~2003–2006~~ radon data.

Figure 3. Seasonal cycle **(a)** and annual mean **(b)** methane fluxes for 1994–2007 integrated over the SE Australian region (135–155° E, 30–45° S). Six flux scenarios are shown as listed in the key.

Figure 4. Monthly mean methane residuals [for 2002-2004](#) at Cape Grim for the observations (black) and six emission scenarios (colours shown in key) **(a)** ACCESS **(b)** CCAM. Note that the observational mean residual for April [2003](#) is missing, due to lack of radon data with which to define the baseline threshold (rather than a lack of methane data). [Refer Fig. 2.](#)

Figure 5. Cape Grim data for January 2003. Upper panel: observed (black) CH₄ residuals; CCAM simulated residuals for tracers CTL (red) and WLBB (blue). Middle panel: observed CO. Lower panel: observed H₂.

Figure 6. Scatter plots of methane to radon, with linear fits for January 2006. The upper panel shows the observational data, the middle panel the CCAM data and the lower panel the ACCESS data.

Figure 7. Mean seasonal cycle of residual methane to radon ratios for each of the six methane tracersemission scenarios. Observations are shown in black, ACCESS results in blue and CCAM results in red.

Figure 8. Mean seasonal cycle of residual methane to radon ratios for the CTL methane scenario (red, solid) and the CTL methane fluxes shifted forward in time by three months (red, dashed) compared to the observations (black). Both simulations used the CCAM model.