

1 **Overview of the Focused Isoprene eXperiments at California Institute of Technology**  
2 **(FIXCIT): mechanistic chamber studies on the oxidation of biogenic compounds**

3

4 Tran B. Nguyen,<sup>1</sup> John D. Crouse,<sup>1</sup> Rebecca H. Schwantes,<sup>1</sup> Alexander P. Teng,<sup>1</sup> Kelvin H.  
5 Bates,<sup>2</sup> Xuan Zhang,<sup>1</sup> Jason M. St. Clair,<sup>1</sup> William H. Brune,<sup>3</sup> Geoffrey S. Tyndall,<sup>4</sup> Frank N.  
6 Keutsch,<sup>5</sup> John H. Seinfeld,<sup>2,6</sup> and Paul O. Wennberg<sup>1,6</sup>

7

8 1. Division of Geological and Planetary Sciences, California Institute of Technology,  
9 Pasadena, California, USA

10 2. Division of Chemistry and Chemical Engineering, California Institute of Technology,  
11 Pasadena, California, USA

12 3. Department of Meteorology, Pennsylvania State University, University Park,  
13 Pennsylvania, USA

14 4. Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder,  
15 Colorado, USA

16 5. Department of Chemistry, University of Wisconsin – Madison, Madison, Wisconsin,  
17 USA

18 6. Division of Engineering and Applied Science, California Institute of Technology,  
19 Pasadena, California, USA

20

1 **Abstract**

2 The Focused Isoprene eXperiment at the California Institute of Technology (FIXCIT) was a  
3 collaborative atmospheric chamber campaign that occurred during January 2014. FIXCIT is the  
4 laboratory component of a synergistic field and laboratory effort aimed toward (1) better  
5 understanding the chemical details behind ambient observations relevant to the Southeastern  
6 United States, (2) advancing the knowledge of atmospheric oxidation mechanisms of important  
7 biogenic hydrocarbons, and (3) characterizing the behavior of field instrumentation using  
8 authentic standards. Approximately 20 principal scientists from 14 academic and government  
9 institutions performed parallel measurements at a forested site in Alabama and at the  
10 atmospheric chambers at Caltech. During the four-week campaign period, a series of chamber  
11 experiments was conducted to investigate the dark- and photo-induced oxidation of isoprene,  $\alpha$ -  
12 pinene, methacrolein, pinonaldehyde, acylperoxy nitrates, isoprene hydroxy nitrates (ISOPN),  
13 isoprene hydroxy hydroperoxides (ISOPOOH), and isoprene epoxydiols (IEPOX) in a highly-  
14 controlled and atmospherically-relevant manner. Pinonaldehyde and isomer-specific standards of  
15 ISOPN, ISOPOOH, and IEPOX were synthesized and contributed by campaign participants,  
16 which enabled explicit exploration into the oxidation mechanisms and instrument responses for  
17 these important atmospheric compounds. The present overview describes the goals, experimental  
18 design, instrumental techniques, and preliminary observations from the campaign. This work  
19 provides context for forthcoming publications affiliated with the FIXCIT campaign. Insights  
20 from FIXCIT are anticipated to significantly aid in interpretation of field data and the revision of  
21 mechanisms currently implemented in regional and global atmospheric models.

22

# 1        **Introduction**

## 2        **1.1. Background**

3            Biogenically-produced isoprenoids (hydrocarbons comprised of  $C_5H_8$  units) have global  
4 emission rates to the atmosphere surpassing those of anthropogenic hydrocarbons and methane  
5 (Guenther et al., 1995; Guenther et al., 2012). The biogenic carbon emission flux is dominated  
6 by isoprene ( $C_5H_8$ ) and monoterpenes ( $C_{10}H_{16}$ ), which account for approximately 50% and 30%  
7 of the OH reactivity over land, respectively (Fuentes et al., 2000). Further, it has been suggested  
8 that the atmospheric oxidation of isoprene, in particular, can buffer the oxidative capacity of  
9 forested regions by maintaining levels of the hydroxyl radical (OH) under lower nitric oxide  
10 (NO) conditions (Lelieveld et al., 2008). Due to their large abundances, isoprene and  
11 monoterpenes also dominate the global budget of secondary organic aerosol (SOA) (Henze et al.,  
12 2008). Thus, the accurate representation of detailed chemistry for isoprene and monoterpene is  
13 necessary for meaningful simulations of atmospheric  $HO_x$  (OH +  $HO_2$ ),  $NO_x$  (NO +  $NO_2$ ),  
14 surface ozone ( $O_3$ ), trace gas lifetimes, and SOA.

15            Unsaturated hydrocarbons like isoprene and monoterpenes are primarily oxidized by OH,  
16  $O_3$ , and the nitrate ( $NO_3$ ) radical in the atmosphere. OH-oxidation is the dominant fate for  
17 isoprene but  $O_3$  and  $NO_3$  oxidation can dominate reactivity for monoterpenes and sesquiterpenes.  
18 Our understanding of the OH-initiated isoprene oxidation mechanism has significantly improved  
19 during the last decade, following the first suggestion of the capacity of isoprene to produce SOA  
20 (Claeys et al., 2004). The mechanistic developments have been propelled by technological  
21 advancements in instrumentation (Hansel et al., 1995; Crounse et al., 2006; Jordan et al., 2009;  
22 Junninen et al., 2010), enabling the detection of more-complex oxidation products derived from  
23 isoprene and other biogenic hydrocarbons. However, the scientific understanding of these

1 biogenic oxidation mechanisms is far from complete. It is outside the scope of this overview to  
2 describe comprehensively the isoprene and monoterpene oxidation mechanisms. Rather, we  
3 provide a brief background of the oxidation of biogenic hydrocarbons, which includes a “state-  
4 of-the-science” knowledge, to motivate the study. The mechanisms described here are illustrated  
5 in Scheme 1.

6 **OH oxidation:** OH predominantly adds to either of the double bonds of isoprene,  
7 followed by the reversible addition of O<sub>2</sub> (Peeters et al., 2009) to produce several isomers of  
8 alkylperoxyl radicals (RO<sub>2</sub>). In the atmosphere, these RO<sub>2</sub> react mainly with HO<sub>2</sub> and NO to form  
9 stable products, although self-reaction can be non-negligible under certain conditions. The stable  
10 products are often termed oxidized volatile organic compounds (OVOCs). In urban-influenced  
11 areas, the “high-NO” pathway is more important and in more pristine environments, the “low-  
12 NO” or HO<sub>2</sub>-dominated pathway is more important. The high-NO pathway generates isoprene  
13 hydroxy nitrates (ISOPN) that act as reservoirs for NO<sub>x</sub>, as well as other products such as methyl  
14 vinyl ketone (MVK), methacrolein (MAC), and hydroxyacetone (HAC) (Paulot et al., 2009a).  
15 For conditions with sufficiently high NO<sub>2</sub>-to-NO ratios, as is mainly the case in the atmospheric  
16 boundary layer outside of cities, methacryloyl peroxyxynitrate (MPAN) is formed from the  
17 photooxidation of MAC. Further oxidation of MPAN can generate SOA (Chan et al., 2010,  
18 Surratt et al., 2010). The low-NO pathway generates isoprene hydroxy hydroperoxides  
19 (ISOPOOH) in almost quantitative yields, and further OH-oxidation of ISOPOOH produces the  
20 epoxydiols in an OH-conserving mechanism (Paulot et al., 2009b). In unpolluted atmospheres,  
21 when the RO<sub>2</sub> lifetimes are sufficiently long (~ 100 s in a forest), isomerization of the RO<sub>2</sub>  
22 followed by reaction with O<sub>2</sub> becomes an important fate, producing the isoprene hydroperoxy  
23 aldehydes (HPALDs) and other products (Peeters et al., 2009; Crouse et al., 2011). These RO<sub>2</sub>

1 isomerization reactions are a type of rapid oxygen incorporation chemistry (Vereecken et al.,  
2 2007; Crouse et al., 2013; Ehn et al., 2014) that is thought to be responsible for the prompt  
3 generation of low-volatility SOA components. Further generations of OH-oxidation in isoprene  
4 are currently being explored owing to recent success with chemical syntheses of important  
5 OVOCs (Wolfe et al., 2012; Jacobs et al., 2013; Bates et al., 2014; Lee et al., 2014b). It has been  
6 found that the OH-oxidation of IEPOX and ISOPN, surprisingly under both low-NO and high-  
7 NO conditions, results primarily in fragmentation of the C<sub>5</sub> skeleton.

8         Despite extensive work on the isoprene + OH mechanism, large uncertainties persist,  
9 some of which directly translate into uncertainties in atmospheric model predictions. These  
10 uncertainties stem from, for example, the large range in reported yields for isoprene nitrates (4 –  
11 15%) (Paulot et al., 2009a), disagreements up to 90% in reported MAC and MVK yields from  
12 the low-NO pathway (Liu et al., 2013, and references therein), various proposed sources of SOA  
13 from the high-NO pathway (Chan et al., 2010; Kjaergaard et al., 2012; Lin et al., 2013), missing  
14 contributions to SOA mass from the low-NO pathway (Surratt et al., 2010), uncharacterized fates  
15 of oxidized species like HPALDs (which may have isomer-dependence), incomplete  
16 understanding of oxygen incorporation (Peeters et al., 2009; Crouse et al., 2013), and under-  
17 characterized impact of RO<sub>2</sub> lifetimes on chamber results (Wolfe et al., 2012). The OH oxidation  
18 of  $\alpha$ -pinene (Eddingsaas et al., 2012) and other monoterpenes is less well-characterized than that  
19 of isoprene, but in general, proceeds through analogous steps.

20         **Ozone oxidation:** Ozonolysis is a significant sink for unsaturated hydrocarbons and a  
21 large nighttime source of OH, particularly in urban-influenced areas. Reaction with ozone is  
22 more important for monoterpenes than isoprene, due to the faster rate coefficients (Atkinson and  
23 Carter, 1984) and the nighttime emission profile for the monoterpenes. Further, monoterpene

1 ozonolysis is highly efficient at converting VOC mass to SOA (Hoffmann et al., 1997; Griffin et  
2 al., 1999). There is a general consensus that ozonolysis occurs via the Criegee mechanism  
3 (Criegee, 1975), wherein ozone adds to a hydrocarbon double bond to form a 5-member primary  
4 ozonide that quickly decomposes to a stable carbonyl product and an energy-rich Criegee  
5 intermediate (CI). In  $\alpha$ -pinene oxidation, ozonolysis,  $\text{NO}_3$ -initiated, and OH-initiated reactions  
6 all produce pinonaldehyde ( $\text{C}_{10}\text{H}_{16}\text{O}_2$ ) as a major product (Wängberg et al., 1997; Atkinson and  
7 Arey, 2003), whereas major first-generation products from isoprene ozonolysis include MAC,  
8 MVK, and formaldehyde. The “hot” Criegee can promptly lose OH (Kroll et al., 2001) while  
9 ejecting an alkyl radical, or become stabilized by collision with atmospheric gases to form a  
10 stabilized Criegee intermediate (sCI) with long enough lifetimes to react bimolecularly. The  
11 subsequent reactions of sCIs produce both carbonyl products and non-carbonyl products such as  
12 hydroperoxides. The *syn* and *anti* conformers of CIs and sCI can have substantially different  
13 reactivities (Kuwata et al., 2010; Anglada et al., 2011), with *syn* conformers more likely to  
14 decompose unimolecularly, possibly through a vinyl hydroperoxide intermediate (Donahue et al.,  
15 2011).

16         It has been suggested that reaction with water molecules is a major (if not dominant)  
17 bimolecular fate of sCI in the atmosphere due to the overwhelming abundance of atmospheric  
18 water (Fenske et al., 2000). This suggestion is supported by observations of high mixing ratios  
19 (up to 5 ppbv) of hydroxymethyl hydroperoxide (HMHP), a characteristic product of reactions of  
20 the smallest sCI ( $\text{CH}_2\text{OO}$ ) with water (Neeb et al., 1997), over forested regions and in biomass  
21 burning plumes (Gäb et al., 1985; Lee et al., 1993; Lee et al., 2000; Valverde-Canossa et al.,  
22 2006). Although HMHP and other hydroperoxides produced from ozonolysis are important  
23 atmospheric compounds, their yield estimates are highly uncertain (Becker et al., 1990; Neeb et

1 al., 1997; Sauer et al., 1999; Hasson et al., 2001; Huang et al., 2013). This may be attributable to  
2 the fact that hydroperoxide yields have mainly been determined by offline methods or under  
3 conditions with highly-elevated hydrocarbon loadings. Further, little empirical data exist on the  
4 humidity dependence of product branching in this reaction. Lastly, the rate coefficients for the  
5 sCI+H<sub>2</sub>O reaction, and other sCI reactions, are still uncertain by several orders of magnitude  
6 (Johnson and Marston, 2008; Welz et al., 2012), precluding the assessment of their atmospheric  
7 importance.

8         **Nitrate oxidation:** NO<sub>3</sub>-oxidation also produces RO<sub>2</sub> radicals by addition to alkenes in  
9 the presence of O<sub>2</sub>. Owing to its high reaction rate coefficient coupled to atmospheric abundance,  
10 α-pinene is expected to be an important sink for NO<sub>3</sub> in many areas. The NO<sub>3</sub>-derived RO<sub>2</sub>  
11 radicals react with (a) NO<sub>3</sub> to form alkoxy radicals (RO) that lead primarily to the production of  
12 nitrooxy carbonyls (b); with other RO<sub>2</sub> radicals to form RO radicals, nitrooxy carbonyls, hydroxy  
13 nitrates, and nitrooxy peroxy dimers; and (c) with HO<sub>2</sub> to form nitrooxy hydroperoxides. Further  
14 generation NO<sub>3</sub>-oxidation produces dinitrates, amongst other products. As the NO<sub>3</sub> addition  
15 initiates the reaction, the thermodynamically-preferred organic hydroxy nitrates produced  
16 through nighttime oxidation may be structurally different than those produced in the daytime  
17 through OH oxidation. During nighttime oxidation, tropospheric HO<sub>2</sub> mixing ratios often surpass  
18 those of NO<sub>3</sub> (Mao et al., 2012), implicating HO<sub>2</sub> reaction to be a common fate for NO<sub>3</sub>-derived  
19 RO<sub>2</sub>. However, previous studies of this reaction have maintained conditions where minimal HO<sub>2</sub>  
20 + RO<sub>2</sub> chemistry occurs and the dominant fate of RO<sub>2</sub> is reaction with NO<sub>3</sub> and RO<sub>2</sub> (Ng et al.,  
21 2008; Perring et al., 2009; Rollins et al., 2009; Kwan et al., 2012). This may be one of the  
22 reasons why nitrooxy hydroperoxides (the RO<sub>2</sub> + HO<sub>2</sub> product) are observed with much higher  
23 relative abundances in ambient air (Beaver et al., 2012) than in chamber studies.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**1.2. Scientific goals**

The 2014 Focused Isoprene eXperiment at the California Institute of Technology (FIXCIT) is a collaborative atmospheric chamber campaign focused on advancing the understanding of biogenic hydrocarbon oxidation in the atmosphere. The campaign was motivated by the communal need for a tight coupling of field and laboratory efforts toward understanding the mechanistic details responsible for ambient observations, exploring explicit chemistry as driven by the fate of RO<sub>2</sub> radicals through well-controlled experiments, and fully characterizing instrumental response to important trace gases using authentic standards to guide data interpretation. To accomplish these goals, a suite of instruments typically deployed for field missions was used to perform parallel measurements at a forested site in Alabama and then in the atmospheric chambers at Caltech. This overview provides an account of the goals and conditions for the experiments performed during the campaign. A key component of FIXCIT is the re-design of “typical chamber experiments” to recreate the ambient atmosphere with higher fidelity so that results from laboratory studies can be implemented in models and used to interpret ambient observations with higher confidence.

**Understanding ambient observations:** FIXCIT was designed as a sister investigation to the 2013 Southern Oxidant and Aerosol Study (SOAS). During SOAS (June – July 2013), a select sub-suite of instruments recorded ambient observations above the forest canopy on top of a metal walk-up tower 20 m in height. The sampling site, located in Brent, Alabama at the Centreville (CTR) SEARCH location managed by the Electric Power Research Institute (CTR, Latitude 32.90289 Longitude -87.24968), was surrounded by a temperate mixed forest (part of the Talladega National Forest) that was occasionally impacted by anthropogenic emission. CTR



1 was characterized by high atmospheric water content (2.4 – 3 vol. % typically), elevated  
2 temperatures (28 – 30°C during the day), high SOA loadings (particulate organics ~ 4 – 10  $\mu\text{g m}^{-3}$ ;  
3 sulfate ~ 2  $\mu\text{g m}^{-3}$ ), high isoprene mixing ratios (4 – 10 ppbv), high ozone (40 – 60 ppbv), low-  
4 to-moderate nitrogen oxides ( $[\text{NO}] \sim 0.3 - 1.5$  ppbv,  $[\text{NO}_2] \sim 1 - 5$  ppbv), occasional plumes of  
5  $\text{SO}_2$  from nearby power plants, and occasional biomass burning events during the SOAS  
6 campaign.

7 The first goal of the chamber campaign was to further investigate the more interesting  
8 observations at SOAS. Due to the ability of laboratory experiments to study the chemistry of a  
9 single reactive hydrocarbon in a controlled setting, it was possible to test hypotheses during  
10 FIXCIT in a systematic manner. Below we list some relevant questions from the SOAS  
11 campaign that were explored during FIXCIT.

- 12 1. *Which reactions or environmental conditions control the formation and destruction of*  
13 *OVOCs in the Southeast U.S.?*
- 14 2. *Are  $\text{RO}_2$  isomerization and other rapid oxygen incorporation mechanisms of key*  
15 *hydrocarbons important during SOAS?*
- 16 3. *How do anthropogenic influences, e.g.,  $\text{NO}_x$ ,  $\text{O}_3$ , and  $(\text{NH}_4)_2\text{SO}_4$ , impact atmospheric*  
17 *chemistry over the forest?*
- 18 4. *How much does the  $\text{NO}_3$ -initiated reaction control nighttime chemistry during SOAS?*
- 19 5. *How do environmental conditions in the Southeast U.S. affect ozonolysis end*  
20 *products, which are known to be water-sensitive?*
- 21 6. *Which reactions or environmental conditions most significantly impact SOA mass and*  
22 *composition?*

23  
24 **Updating the isoprene and monoterpene mechanisms:** Several experiments were  
25 designed to “fill in the gaps” of the isoprene oxidation mechanisms by leveraging the

1 comprehensive collection of sophisticated instrumentation at FIXCIT. We targeted the following  
2 acknowledged open questions.

- 3 7. *What are the products of the photochemical reactions stemming from OVOCs like*  
4 *ISOPOOH, IEPOX, ISOPN, and pinonaldehyde?*
- 5 8. *What is the impact of photolysis vs. photooxidation for photolabile compounds?*
- 6 9. *What is the true yield of isoprene nitrates from the high-NO photooxidation*  
7 *pathway?*
- 8 10. *What is the product distribution and true yield of nitrooxy hydroperoxides from the*  
9 *NO<sub>3</sub>-oxidation reaction of isoprene and monoterpenes under typical atmospheric*  
10 *conditions?*
- 11 11. *How do products and yields change as RO<sub>2</sub> lifetimes in chamber studies approach*  
12 *values estimated to be prevalent in the troposphere?*

13

14 **Instrument characterization:** A final goal of FIXCIT was to evaluate, compare, and  
15 identify biases in field instrumentation by isolating one variable at a time. We focused on the  
16 following objectives.

- 17 12. *Identify the causal factor(s) producing the “OH interference” (Mao et al., 2012)*  
18 *that has been observed in various biogenically-impacted regions by some gas-*  
19 *expansion laser-induced-fluorescence (LIF) techniques.*
- 20 13. *Characterize the performance of newly commercially-available CIMS*  
21 *instrumentation with respect to the detection of OVOCs by using authentic*  
22 *standards.*
- 23 14. *Compare similar measurements (e.g., OH reactivity, hydrocarbons, OVOCs) made*  
24 *with different techniques.*

25

## 26 **2. Scope of the campaign**

27 **2.1. Facilities:** Experiments were performed in the Caltech Atmospheric Chamber Facility  
28 within a one-month period in January 2014. The facility contains several in-house gas- and

1 aerosol-phase instruments and an 8 x 5 m insulated enclosure, housing two side-by-side Teflon  
2 atmospheric chambers that are suspended from the ceiling. The chambers were manufactured  
3 from Fluorinated Ethylene Propylene (FEP) Teflon. The chamber volume was measured  
4 regularly by quantitative transfer of highly-volatile organics such as isoprene by an externally  
5 calibrated GC-FID. Quantitative transfer was checked via injections of a measured quantity of  
6 isoprene (checked by gravimetric, volumetric, and FT-IR methods) into a pillow bag with known  
7 volume by timing a calibrated mass flow of air into the pillow bag. For most experiments, the  
8 chamber volume was between 23 – 24 m<sup>3</sup>. The spatial configuration of instruments in the  
9 chamber facility during FIXCIT is shown in Fig. 1. The instruments, contributors, and  
10 identifying abbreviations used in this work are described in Table 1. A total of 320 UV black  
11 lamps (broadband  $\lambda_{\text{max}} \sim 350$  nm) are mounted on the walls of the enclosure. The lamps are  
12 located behind Teflon films so that the heat produced from the operation of the lamps can be  
13 removed by recirculating cool air. The interior of the enclosure is covered with reflective  
14 aluminum sheets. Light intensities can be tuned to 100 %, 50 %, 10 %, and 1 %.  $J_{\text{NO}_2}$  was  
15 measured to be  $7 \times 10^{-3} \text{ s}^{-1}$  at 100 % light intensity. Light fluxes at several locations within the  
16 chamber (e.g., center, corner, right, left, high, low) did not vary more than 15%. Temperature  
17 controls in the chamber enclosure are tunable from 10 – 50 °C (typically set at 25°C) and did not  
18 fluctuate more than 1°C except during periods when the temperature was explicitly changed or  
19 during a 30 minute period immediately following a change in the light intensities (up to 2 °C  
20 increase was observed from switching on 100 % lights.)

21         The chamber experiments were operated in batch mode throughout the campaign.  
22 Temperature and RH were monitored continuously inside the chamber by a Vaisala HMM211  
23 probe calibrated with saturated salt solutions in the RH range of 11 – 95%. In the range  $\text{RH} <$

1 11%, water vapor measurements were provided by the TripCIMS. The chambers were flushed at  
2 least 24 h before each use with ultra-purified air (purified in-house via a series of molecular  
3 sieves, activated carbon, Purafil™ media, and particulate filters), at elevated temperature when  
4 needed (~ 40 °C), so that the backgrounds on gas- and particle-phase instrumentation are at  
5 baseline levels. As a reference, NO levels before each run were typically < 100 pptv (from NO-  
6 CL measurements) and particle concentrations were < 0.01 µg m<sup>-3</sup>. Flushing rates, as balanced  
7 by exhaust rates, were typically 250 std. L min<sup>-1</sup> (SLM) or ~ 0.6 chamber volumes per hour.  
8 Chambers were mixed on the timescale of minutes by injecting high-pressure pulses of air during  
9 the beginning of experiments.

10 Chamber 1 was reserved for low-NO experiments, so that the walls did not contact  
11 elevated levels of nitric acid and organic nitrates during the lifetime of the chamber, while  
12 Chamber 2 was reserved for moderate-to-high-NO experiments. Experiments were carried out  
13 daily in alternating chambers to allow for the full flushing period of the previously-used  
14 chamber. Each chamber was characterized separately prior to the campaign for vapor and  
15 particle wall loss rates. Typically, wall loss rates for gas-phase species are slightly higher in the  
16 high-NO chamber than the low-NO chamber due to the greater acidity of the walls. Particle wall  
17 loss rates were not significantly different between chambers. Measurements of the particle wall  
18 loss rates were performed by injecting ammonium sulfate (AS) seed aerosols into the chamber  
19 and monitoring the decay over the course of 10 – 24 h. Particles were injected via atomization of  
20 dilute salt solutions (e.g., AS 0.06 M) through a <sup>210</sup>Po neutralizer and water trap. Measurements  
21 of vapor wall loss rates were performed by injecting OVOC standards (e.g., IEPOX, HMHP,  
22 etc.) into the chamber. Both particle and vapor wall loss characterizations were performed at

1 several RH conditions (4 – 85 % RH). These characterizations have been described in more  
2 detail previously (Loza et al., 2010; Nguyen et al., 2014).

3       Organic compounds were injected into the chamber by two methods: (1) For volatile  
4 compounds, a measured volume was injected with a micro-syringe through a septum into a clean  
5 glass bulb and the evaporated standard was quantitatively transferred into the chamber by dry  
6 purified air. Gas introduction of VOCs (done for isoprene and methacrolein) by filling an  
7 evacuated bulb with the chemical vapor, backfilling with nitrogen gas, and characterizing with  
8 Fourier-transform infrared spectrometry before injecting did not produce significantly different  
9 results than volume injection. (2) For semi-volatile compounds, the solid or liquid standard was  
10 placed inside a two-neck flask, which was heated by a water bath (35 – 65°C), and the headspace  
11 was carried into the chamber by dry purified air. The ToFCIMS or TripCIMS instruments  
12 measured the gas-phase mixing ratio of the semi-volatiles in real time as the compounds entered  
13 the chamber, and injection was halted when a satisfactory quantity was introduced. OVOCs were  
14 calibrated by the ToFCIMS and TripCIMS by methods described earlier (Paulot et al., 2009a).  
15 The desired RH inside the chamber was achieved by flowing dry purified air through a water-  
16 permeable (Nafion) membrane humidifier (FC200, Permapure LLC), kept moist by recirculating  
17 27 °C ultra-purified (18 MΩ, 3 ppb TOC) water (Milli-Q, Millipore Corp). Particles were  
18 atomized into the chamber as described for particle wall loss experiments. When hydrated  
19 particles were needed for experiments, particles were injected via an in-line, heated, wet-wall  
20 denuder into a chamber that has RH above the efflorescence point of the particular salt (Martin,  
21 2000).

22

1        **2.2. Instrumentation and sampling modifications:** Instruments were connected via  
2 sampling lines to both chambers through port holes in the enclosure as shown in Fig. 1. Sampling  
3 lines were capped when not in use. Inlet and tubing material were instrument-specific and  
4 included: stainless steel (GTHOS and ToF-AMS), heated stainless steel and quartz (TDLIF),  
5 electro-polished steel and FEP Teflon (NO<sub>3</sub>CIMS), polyetheretherketone (PEEK) and Teflon  
6 (SRI-ToFMS), and perfluoroalkoxy polymer (PFA) Teflon (other instruments).

7        The duration of each experiment (i.e., the level of oxidation that can be probed) was  
8 critically dependent on the net sampling flow rates at which air was withdrawn from the  
9 chamber. Sampling strategies were developed to minimize the effective sampling flow rate from  
10 each instrument, in such a way that instrument responses were not significantly different than  
11 during field campaigns. In many cases, a common high-flow Teflon sampling line was used to  
12 minimize the residence time of gases through tubing, and smaller flows were sampled  
13 orthogonally by each instrument. In some cases, a duty cycle was used as needed.

14        Several modifications from field designs were utilized for chamber sampling. The  
15 modifications included: (1) the GTHOS detection system was located between the chambers  
16 inside of the enclosure to minimize the residence time of HO<sub>x</sub> inside the instrument (Fig. 1). The  
17 detection system was connected to the laser on the outside of the enclosure via a 3 m fiber optic  
18 cable fed through the side port hole. The sampling flow rate was similar to field flows (6 SLM);  
19 however, the fast-flow inlet was situated horizontally (~ 2 m height) instead of vertically. The  
20 inlet was adapted to each bag directly, by attaching it to a Teflon plate that was in turn secured to  
21 the chamber walls via a large o-ring. The GTHOS inlet switched from Chamber 1 to Chamber 2  
22 as needed. Chemical zeroing was performed by releasing hexafluoropropene (C<sub>3</sub>F<sub>6</sub>) into the inlet  
23 as an OH scrubber, and dark zeroing by measuring the difference between online and offline

1 signals. Chemical and dark zeroing methods were used to distinguish between OH present in the  
2 chamber or atmosphere (chemical OH) and OH that may have been produced after the gas  
3 stream enters the instrument, which is additional to the chemical OH signal; (2) LIF-OHR was  
4 diluted a factor of 10 with nitrogen gas (effective flow 6 SLM); (3) NO<sub>3</sub>CIMS was diluted a  
5 factor of 5 with scrubbed zero air (effective flow 2 SLM); (4) GlyLIP and FormLIF both  
6 operated at 5 SLM instead of the usual 17 and 10 SLM, respectively; and (5) SRI-ToFMS (1.5  
7 SLM) and GCFID (0.1 SLM) occasionally sampled through a 0.125 – 0.25” OD PFA Teflon  
8 tube that was submerged in a cold bath kept at – 40°C in order to remove interferences from  
9 certain OVOC (see Section 2.3).

10 GC-ToFCIMS, first described in Bates et al (2014), is an extension of the ToFCIMS.  
11 Analyte gas samples were focused with a cold trap onto the head of a RTX 1701 column  
12 (Restek) and eluted with a temperature ramping program (30 – 130°C) in the oven before  
13 reaching the ToFCIMS for mass spectrometry detection. GC-ToFCIMS recorded data only when  
14 isomer separation was needed, because its operation took the standard scanning mode of the  
15 ToFCIMS offline. All other instruments operated normally with the following sampling flows:  
16 TDLIF (4 SLM), ToFCIMS and TripCIMS (2 SLM), CRM-OHR (0.5 SLM), NO-CL (1 SLM),  
17 and IACIMS (2 SLM). Frequencies of zeroing (with dry N<sub>2</sub> or ultrazero air) and calibration  
18 (various methods) were instrument-specific, with some instruments zeroing once per hour and  
19 calibrating once per few hours and others performing zeroing/calibration between experiments.

20

### 21 **2.3. Experimental design**



22 The experiments performed at FIXCIT can be divided into several categories, each  
23 probing one or more specific science questions outlined in Section 1.2. Every experiment

1 included successful elements from past studies, but with a special focus on extending to  
2 atmospheric conditions. One example is reducing the occurrence of RO<sub>2</sub> + RO<sub>2</sub> side reactions in  
3 chamber experiments, which can lead to yields of atmospherically-relevant products that are  
4 biased low. Enabled by the high sensitivity of field instruments, photooxidation was performed  
5 with precursor mixing ratios as low as 12 ppbv. Certain instruments that required extensive  
6 dilution in a chamber setting, e.g., LIF-OHR, had poorer-quality data for low-loading  
7 experiments. Experimental durations were typically 4 – 6 h, with the exception of overnight runs  
8 where the majority of instruments sampled briefly to establish starting conditions, then were  
9 taken offline during the nighttime and resumed sampling in the morning. The typical reaction  
10 time for an overnight experiment was ~ 15 h. Experimental details are reported in Table 2. OH  
11 concentrations were derived from hydrocarbon decay data from GCFID, SRI-ToFMS, or  
12 ToFCIMS, when available, using published rate coefficients (Atkinson et al., 2006; Lee et al.,  
13 2014b; Bates et al., 2014). Otherwise, preliminary GTHOS chemical-zeroing data were used.  
14 The following types of experiments were included in the study:

15 **a. Blank** (Exp. 4b and 5b): Blank experiments were designed to investigate background signals  
16 present in experiments that may have sources other than gas-phase chemistry of the injected  
17 hydrocarbon, e.g., from heterogeneous oxidation of residual organics on the chamber walls.  
18 OH precursors, such as hydrogen peroxide, were added to each chamber, the UV lamps were  
19 turned on, and sampling occurred as usual. Further, the temperatures inside the chambers  
20 were ramped from 25 to 35° C to explore the extent to which elevated temperatures change  
21 the chamber background signals due to increased volatilization of organics. Blank  
22 experiments were performed under dry conditions. Common background compounds  
23 produced from heterogeneous wall reactions are formic acid and acetic acid.



1 **b. Low-NO photooxidation** (Exp. 2, 10, 17, 19, and 25): The “low-NO” experiments that have  
2 been extensively investigated in atmospheric chamber studies were designed to be relevant to  
3 the pristine troposphere, and certain conditions at SOAS, where HO<sub>2</sub> reactions dominate the  
4 RO<sub>2</sub> fate. Experiments were initiated by H<sub>2</sub>O<sub>2</sub> photolysis as a NO<sub>x</sub>-free source of OH and  
5 HO<sub>2</sub>:



6  
7  
8 The execution of these experiments requires precise engineering to closely simulate the  
9 troposphere. One outstanding challenge of “low-NO” experiments is the variation in initial  
10 NO levels across different chamber settings and on different days. Because typical HO<sub>2</sub>  
11 levels in a chamber environment do not typically exceed ~ 200 pptv from the self-limiting  
12 HO<sub>2</sub> recombination, NO should be ~ 40 pptv during the reaction (a factor of 5 less abundant)  
13 in order for the C<sub>5</sub> RO<sub>2</sub> reactions to be dominated by HO<sub>2</sub> by a factor of 10 ( $k_{\text{RO}_2+\text{HO}_2} \sim 1.6 \times$   
14  $10^{-11}$  and  $k_{\text{RO}_2+\text{NO}} \sim 8.5 \times 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  at 298K (Atkinson et al., 2006)). Thus,  
15 experimental variations in NO that can lead to discrepancies in “low-NO” kinetics typically  
16 elude quantification by commercially-available NO chemiluminescence instruments, owing to  
17 their high limits of detection (~500 pptv).



18 NO levels in the Caltech chambers were suppressed by continually flushing with filtered  
19 air on the inside and outside the chamber walls. Initial NO levels of < 40 pptv were typically  
20 achieved during experiments. The NO-CL instrument available during FIXCIT (Table 1) has  
21 a limit of detection better than 25 pptv, and the GTHOS instrument provided online HO<sub>2</sub>  
22 quantification at the pptv-level. Another common challenge for “low-NO” experiments (even  
23 when [NO] < [HO<sub>2</sub>]) is that homogeneous or cross RO<sub>2</sub> + RO<sub>2</sub> reactions may dominate the

1 RO<sub>2</sub> reactivity ( $k_{\text{RO}_2+\text{RO}_2} \sim 10^{-15} - 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  at 298K (Atkinson et al., 2006)).  
2 These experiments may be more correctly characterized as “low-NO, high-RO<sub>2</sub>”. For  
3 experiments using [H<sub>2</sub>O<sub>2</sub>] as an OH precursor, RO<sub>2</sub>+RO<sub>2</sub> reactions were largely minimized  
4 by using reaction conditions that ensures [HO<sub>2</sub>] > [RO<sub>2</sub>] (e.g., [H<sub>2</sub>O<sub>2</sub>]<sub>0</sub>/[ISOP]<sub>0</sub> ~ 10<sup>5</sup> and  
5  $J[\text{H}_2\text{O}_2] \sim 4 - 5 \times 10^{-6} \text{ s}^{-1}$ ). Thus, the peroxy radical self-reaction channels are minor  
6 compared to RO<sub>2</sub>+RO<sub>2</sub> chemistry. We estimate that the “low-NO” experiments were HO<sub>2</sub>-  
7 dominated by at least a factor of 10 in RO<sub>2</sub> reactivity by monitoring tracers of chemistry  
8 stemming from high-NO (isoprene nitrates), high-RO<sub>2</sub> (C<sub>5</sub> diols and other products), and  
9 low-NO (ISOPOOH and IEPOX) pathways. The molar yield of the low-NO products  
10 ISOPOOH + IEPOX (measured within the first 15 minutes of reaction) was estimated to be  
11 95%, supporting the dominance of RO<sub>2</sub> + HO<sub>2</sub> chemistry over other channels. The  
12 structurally isomeric ISOPOOH and IEPOX that were formed from the HO<sub>2</sub>-dominated  
13 isoprene photooxidation were distinguished by TripCIMS, and the sum was measured by  
14 ToFCIMS, IACIMS, and NO<sub>3</sub>CIMS. These experiments were performed with isoprene,  $\alpha$ -  
15 pinene, 4,3-ISOPOOH and MAC precursors.

16 **c. High-NO photooxidation** (Exp. 3, 11, 22, and 24): “High-NO” experiments are also  
17 commonly performed in chamber studies. These experiments were designed to be relevant to  
18 the urban-influenced troposphere, such as some cases at SOAS, where NO can dominate RO<sub>2</sub>  
19 reactions. Experiments were typically initiated by H<sub>2</sub>O<sub>2</sub> with added NO during FIXCIT, but  
20 have been performed using HONO or other precursors elsewhere. It is easier to ensure that  
21 reaction with NO is the main fate of RO<sub>2</sub>, even with higher hydrocarbon loadings, because  
22 NO mixing ratios are typically in excess of both HO<sub>2</sub> and RO<sub>2</sub> by hundreds of ppbv. Hydroxy  
23 nitrate products were measured by TDLIF, IACIMS, ToFCIMS, and GC-ToFCIMS.

1 Functionalized carbonyl products were measured by SRI-ToFMS and ToFCIMS. Glyoxal  
2 and formaldehyde, also important high-NO products, were measured by the GlyLIP and  
3 FormLIF, respectively. This well-studied experiment was important for multiple reasons,  
4 including calibration, diagnostics, and for determining the hydroxy nitrate yields from  
5 alkenes within the first few minutes of photooxidation. However, it should be noted that the  
6 experimental result represents a boundary condition that may not fully represent NO-  
7 influenced reactions in the atmosphere due to the extremely short RO<sub>2</sub> lifetimes (< 0.01 s at  
8 500 ppbv NO). These experiments were performed with isoprene, α-pinene, and the 4,3-  
9 ISOPN standard synthesized by the Caltech group.

10 **d. “Slow chemistry” photooxidation** (Exp. 7, 16, 18, and 27): The slow chemistry experiment  
11 is designed to extend RO<sub>2</sub> lifetimes closer to atmospheric values when both NO and HO<sub>2</sub>  
12 impact RO<sub>2</sub> reactivity (~ 3 - 30 s, assuming 1500 – 100 pptv NO and 40 pptv HO<sub>2</sub>). This was  
13 achieved by employing low radical mixing ratios. With relevant RO<sub>2</sub> lifetimes, the RO<sub>2</sub>  
14 isomers may be closer to their equilibrium distribution because of the reversible addition of  
15 oxygen (Peeters et al., 2009). Figure 2 shows the progress of a representative slow chemistry  
16 experiment. The “slow” portion of experiments was performed under a low light flux ( $J_{\text{NO}_2} \sim$   
17  $4 \times 10^{-5} \text{ s}^{-1}$ ) with methyl nitrite as the OH precursor (Atkinson et al., 1981):



20 These reactions produce a steady-state OH concentration of  $[\text{OH}]_{\text{ss}} \sim 0.4 - 1 \times 10^5 \text{ molec cm}^{-3}$   
21 and an atmospherically-relevant ratio of NO/HO<sub>2</sub> (2 – 3) that is stable throughout the  
22 majority of the experiment. Further, we aimed to simulate the summer conditions at SOAS,  
23 where RO<sub>2</sub> isomerization is competitive with RO<sub>2</sub> + HO<sub>2</sub> and RO<sub>2</sub> + NO chemistry. Thus,

1 most experiments of this type were performed at elevated temperatures ( $T \sim 40 - 45^\circ\text{C}$ ) to  
2 facilitate the isoprene  $\text{RO}_2$  isomerization to HPALDs (Crouse et al., 2011), as measured by  
3 ToFCIMS. The atmospheric  $\text{RO}_2$  fates were qualitatively deduced by observations of their  
4 respective products during SOAS (forthcoming papers) and during other campaigns (Paulot  
5 et al., 2009b; Wolfe et al., 2011; Beaver et al., 2012).





6 The fate of HPALDs is not known, but has been suggested to be strongly influenced by  
7 photolysis based on reactions of chemical analogues (Wolfe et al., 2012). After the slow  
8 chemistry period, 20 – 100% lights were turned on in order to diagnose the effects of direct  
9 photolysis and OH-oxidation on the product compounds, which is especially instructive when  
10 coupled with photochemical modeling. Table 2 reports conditions only for the  $\leq 1\%$  light  
11 period and the 20% light period due to availability of hydrocarbon decay data. When  
12  $\text{CH}_3\text{ONO}$  experiments were performed with higher light flux from the start, the NO-to- $\text{HO}_2$   
13 reactivities were still competitive but the OH mixing ratios were higher. These experiments  
14 were performed with isoprene,  $\alpha$ -pinene, and *trans*  $\beta$ -IEPOX precursors.


15 **e. Ozonolysis** (Exp. 6, 14, 23, and 29): Ozonolysis reactions were performed in the dark, with  
16 and without the use of excess cyclohexane (50 ppmv) as a scavenger for OH (Atkinson,  
17 1995). Ozone reacts with isoprene and  $\alpha$ -pinene with rate coefficients of  $k_{\text{ISO}+\text{O}_3} = 1.3 \times 10^{-17}$   
18  $\text{molec cm}^{-3}$  and  $k_{\alpha\text{-PIN}+\text{O}_3} = 9.0 \times 10^{-17} \text{ molec cm}^{-3}$  at 298 K, respectively (Atkinson et al.,  
19 2006). After the first few steps of the reaction, however, little agreement exists in the  
20 literature for product yields, product distribution, or rate coefficients stemming from  
21 reactions of stabilized Criegee intermediates (sCI). This may be due to the large differences  
22 among studies in the hydrocarbon loadings ( $[\text{ISO}]_i = 40 - 10000 \text{ ppbv}$ ), ozone-to-isoprene  
23 ratios ( $< 0.5$  to  $> 100$ ), water vapor content ( $< 10 - 20000 \text{ ppmv}$ ), reaction pressures (4 –

1 760 torr), analytical methods used for product analysis (GC, HPLC, FTIR, direct OH vs.  
2 scavenging, etc.), and methods used to generate sCI ( $\text{CH}_2\text{I}_2 + \text{h}\nu$  vs. gas-phase ozonolysis)  
3 (Simonaitis et al., 1991; Neeb et al., 1997; Sauer et al., 1999; Hasson et al., 2001; Kroll et al.,  
4 2002; Johnson and Marston, 2008; Drozd and Donahue, 2011; Welz et al., 2012; Huang et  
5 al., 2013).

6 We designed the ozonolysis experiments to have similar ozone-to-isoprene ratios to those  
7 observed during SOAS (~ 5 – 7), and performed the experiments under dry (RH ~ 4%) and  
8 moderately-humid (RH ~ 50%) conditions. The ozonolysis experiments at FIXCIT primarily  
9 focused on studying unimolecular and bimolecular chemistry of sCI that affects the yields of  
10 OH, hydroperoxides, organic acids, and aldehydes/ ketones under humid vs. dry conditions.  
11 These experiments represent the first coupling between direct OH observations from  
12 GTHOS, aldehyde/ketone measurements from GCFID and SRI-ToFMS, online  
13 formaldehyde measurements from FormLIF, and online hydroperoxide measurements from  
14 the various CIMS instruments present to provide the most comprehensive picture thus far on  
15 the humidity-dependent ozonolysis of isoprene.

16 **f. Competitive- $\text{HO}_2$  nitrate ( $\text{NO}_3$ ) oxidation (Exp. 9 and 13):** The  $\text{NO}_3$ -initiated experiments  
17 during the campaign were performed in the dark, under dry conditions. Excess formaldehyde  
18 ( $[\text{HCHO}]_i \sim 4 - 8$  ppmv) was used as a dark  $\text{HO}_2$  precursor in order to elevate the  
19 contributions of  $\text{RO}_2 + \text{HO}_2$  reactions in the  $\text{NO}_3$  chemistry:





2 This process produces an HO<sub>2</sub>/NO<sub>3</sub> ratio of approximately 2 (determined by photochemical  
3 modeling from the mechanism described in Paulot et al (2009)), a ratio more relevant to the  
4 troposphere during nighttime oxidation. As α-pinene has a higher NO<sub>3</sub> loss rate compared to  
5 isoprene, a factor of 2 greater mixing ratio of initial formaldehyde was used. The  
6 consequence of the experimental design is that the isoprene nitrooxy hydroperoxide (INP)  
7 and monoterpene nitrooxy hydroperoxide (MTNP) are major products, in contrast to  
8 experiments performed under RO<sub>2</sub> + RO<sub>2</sub> or RO<sub>2</sub> + NO<sub>3</sub> dominated conditions (Ng et al.,  
9 2008; Perring et al., 2009; Kwan et al., 2012). The focus of these experiments was the  
10 quantification of INP and MTNP with the various CIMS and with TDLIF, and further  
11 exploration of their loss channels to OH oxidation (simulating sunrise) or to dry AS seed  
12 particles by measuring organic aerosol growth on the ToF-AMS. These experiments were  
13 performed with isoprene and α-pinene precursors.

14 **g. High NO<sub>2</sub>/NO photooxidation** (Exp. 26 and 30): The high NO<sub>2</sub>-to-NO ratios in the lower  
15 troposphere in most regions of the globe favor the production of acylperoxy nitrates (APNs)  
16 from the OH-initiated reaction of aldehydes like methacrolein and pinonaldehyde (Bertman  
17 and Roberts, 1991; Nozière and Barnes, 1998). Unlike the APN from methacrolein (MPAN),  
18 the APN from pinonaldehyde has never been measured in the atmosphere (Nouaime et al.,  
19 1998; Roberts et al., 1998; Wolfe et al., 2009). The OH oxidations of aldehydes were  
20 performed with an NO<sub>2</sub>/NO ratio > 10, and NO<sub>2</sub> was replenished as it was reacted away.  
21 These reactions were initiated by CH<sub>3</sub>ONO photolysis under higher light flux, producing  
22 [OH] > 3 x 10<sup>6</sup> molec cm<sup>-3</sup>. Certain APNs were monitored with ToFCIMS, and total peroxy  
23 nitrates (ΣPNs) were monitored with TDLIF. A major focus of the high-NO<sub>2</sub> experiments

1 was to investigate the SOA-formation potential and mechanisms from atmospherically-  
2 relevant APNs, which is expanded in **h**.

3 **h. SOA-formation chemistry** (Exp. 19, 24, 26, and 30): Experiments aimed specifically at  
4 studying chemistry leading to SOA formation have overlapping goals with those described  
5 above. One focus was the evaluation of the SOA-formation route from APNs by the  
6 proposed dioxo ketone, lactone, and epoxide mechanisms (Chan et al., 2010; Kjaergaard et  
7 al., 2012 ; Lin et al., 2013), none of which have yet been validated by independent studies.  
8 However, the proposed epoxide chemistry has been integrated into some studies published  
9 soon after the proposal by Lin et al. (2013) (Worton et al., 2013; Pye et al., 2013). After  
10 MPAN was formed from the high-NO<sub>2</sub> reaction of MAC + OH, a synthesized standard of  
11 methacrylic acid epoxide (MAE, provided by the UNC group), the proposed epoxide  
12 intermediate, was added to discern the SOA-forming potential of MAE vs. other reactive  
13 intermediates in the MPAN reaction. Following the injection and stabilization of MAE, water  
14 vapor was added until the reaction mixture reached ~ 40% RH. Then wet AS seeds were  
15 injected to investigate any SOA mass growth, as quantified by ToF-AMS.

16 SOA formation from ISOPN high-NO photooxidation and isoprene low-NO  
17 photooxidation products were investigated in the presence of wet AS seeds (40 – 50%  
18 particle liquid water by volume), meant to simulate the high particle liquid water and sulfate  
19 quantities during SOAS. For these experiments, the chambers were humidified to 40 – 50%  
20 RH, and hydrated AS particles were injected through a wet-wall denuder so that the seed  
21 particles retain liquid water above the efflorescence point of AS (Biskos et al., 2006). In the  
22 ISOPN high-NO photooxidation, the potential for forming organics that will likely condense  
23 onto seed particles, e.g., dinitrates and IEPOX, was recently suggested (Lee et al., 2014b;

1 Jacobs et al., 2014). The dinitrate pathway was investigated as a potential source of particle-  
2 phase organic nitrogen. In the low-NO isoprene photooxidation, IEPOX reactive uptake onto  
3 acidic Mg<sub>2</sub>SO<sub>4</sub> particles (Lin et al., 2012) and non-acidified AS particles (Nguyen et al.,  
4 2014), both with non-zero liquid water content, were recently demonstrated. We focused on  
5 AS particles with no added acid. The impact of the partitioning of IEPOX on the gas-phase  
6 mixing ratios was examined as a potential reason for the differences in observed IEPOX in  
7 dry and humid regions.

8 **i. Cross-calibrations** (Exp. 4a, 5a, 24, 27, and 30): Newly commercially-available negative-  
9 ion CIMS (Junninen et al., 2010; Lee et al., 2014a) may become common tools for  
10 monitoring complex OVOCs in the atmosphere, similarly to the widespread adoption of  
11 positive ion CIMS (PTR-MS based instruments). Some of the new negative ion CIMS  
12 instruments were deployed for the first time in field campaigns occurring in recent years.  
13 During FIXCIT, synthesized standards of 8 isomer-specific compounds were available for  
14 cross calibrations with different CIMS in order to better understand the chemical sources of  
15 ambient signals during SOAS and in other field campaigns. Table 3 shows the structures,  
16 abbreviations, and contributors of the synthesized chemicals. The TripCIMS and the GC-  
17 ToFCIMS separated structural isomers through collision-induced dissociation (CID) and  
18 through chromatography, respectively. Figure 3 shows a GC-ToFCIMS separation of isomers  
19 of the ISOPN synthesized standards, as well as ISOPNs present in a complex photooxidation  
20 mixture. SRI-ToFMS and IACIMS tested the switchable reagent ion sources for preferential  
21 detection of one or more isomers of compounds with the same molecular formula.

22 For certain cross-calibrations experiments, standards were injected into an inflatable  
23 pillow bag (~ 0.2 – 0.3 m<sup>3</sup>) that was filled with dry N<sub>2</sub> to a known volume. The purities of the



1 standards were quantified by nuclear magnetic resonance (NMR) or Fourier-transform  
2 infrared spectrometry (FT-IR). The injected material was measured by vapor pressure,  
3 quantitative volume transfer, or by ToFCIMS and TripCIMS that were calibrated using  
4 techniques described elsewhere (Garden et al., 2009; Paulot et al., 2009a; Lee et al., 2014b;  
5 Bates et al., 2014 ). Some experiments, such as the IEPOX photooxidation experiment, also  
6 served a dual purpose for cross-calibration. For example, *trans*  $\beta$ -IEPOX was injected into a  
7 clean chamber and instruments were allowed to sample for  $\sim 1$  h to cross-calibrate before an  
8 oxidant precursor was injected. Both LIF-OHR and CRM-OHR were able to measure the OH  
9 reactivity of these OVOC compounds individually, which aids in determining the known and  
10 unknown OH reactivity in ambient environments.

11 **j. GTHOS test experiments:** The OH interference in GTHOS, and possibly other gas-  
12 expansion LIF techniques, has been shown to systematically bias OH measurements high in  
13 some biogenically-influenced areas unless chemical zeroing was performed (Mao et al.,  
14 2012). The excess OH was demonstrated not to be produced by the GTHOS laser itself (308  
15 nm) but rather, more likely, in the low-pressure flow zone within the nozzle of the  
16 instrument. During FIXCIT, several hypotheses proposed by Mao et al (2012), and some  
17 original proposals based on field observations, were tested. The interference precursor  
18 candidates were: (i) ozonolysis intermediates – tested with ozonolysis experiments and with  
19 ozone injection into the GTHOS inlet; (ii) biogenic peroxides like ISOPOOH or HMHP –  
20 tested with synthesized standards; (iii) background chemistry such as  $\text{NO}_2 + \text{O}_3$  – tested by  
21 the nitrate-oxidation experiment and by sequential injection of  $\text{NO}_2$  and  $\text{O}_3$  separately; (iv)  
22 dry and humid  $\text{HO}_2 + \text{O}_3$  reaction – tested by formaldehyde photolysis with ozone injection  
23 during a separate experiment (2/1/14, not shown in Table 2); (v) beta-hydroxy  $\text{RO}_2$  radicals

1 formed from OH + alkene – tested with the photooxidation of 2-methyl-2-butanol and  
2 compared with 2,2-dimethylbutane (2/2/14 and 1/31/14, not shown in Table 2); and (vi) heat-  
3 mediated decomposition of thermally-unstable species – tested by temperature ramping to 35  
4 – 40 °C inside the chamber. Often, single variables (like ozone or heat) were isolated by  
5 incremental additions toward the end of an experiment.

6 The experiments not described in Table 2 (to test iv and v) were performed after the  
7 formal experiments; thus, not all investigators were present. Only GTHOS, ToFCIMS,  
8 TripCIMS, ToF-AMS, GCFID, O<sub>3</sub> monitor and NO<sub>x</sub> monitor were collecting data. The HO<sub>2</sub>  
9 + O<sub>3</sub> test experiment (2/1/14) was performed by injecting ~ 600 ppbv of ozone, then ~ 50  
10 ppbv of cyclohexane as an OH tracer for CIMS (monitored by the formation of cyclohexyl  
11 hydroperoxide). UV lights were turned on and then 4 ppmv of formaldehyde was injected,  
12 which photolyzed to produce 550 pptv of HO<sub>2</sub>. The HO<sub>2</sub> reaction with formaldehyde  
13 produced a small yield of HMHP (Niki et al., 1980). Water vapor was injected to diagnose  
14 the effect of humidity. Experiments to test the effects of RO<sub>2</sub> structure utilized CH<sub>3</sub>ONO to  
15 oxidize ~ 50 ppbv of either 2-methyl-2-butanol and 2,2-dimethylbutane with OH. Ozone (~  
16 600 ppbv), water vapor (until RH ~ 30 – 40%), and NO<sub>2</sub> (400 ppbv) were added sequentially  
17 at toward the end of the photooxidation. Finally severally hundred ppb of NO was added to  
18 titrate away the ozone.

#### 19 20 **2.4. Analytical challenges**

21 Throughout the campaign, several sources of analytical interferences or systematic  
22 biases were discovered. Some challenges resulted from the integration of field instruments to  
23 a chamber setting, where high concentrations of certain chemicals were used to engineer

1 extremely specific conditions. Thus, these issues do not affect ambient sampling. For  
2 example: (1) High  $\text{NO}_2$  levels in some experiments affected the normal operation of TDLIF  
3 because the  $\Sigma\text{ANs}$  and  $\Sigma\text{PNs}$  measurements were determined by subtraction of  $\text{NO}_2$ . When  
4  $\text{NO}_2$  is much higher than  $\Sigma\text{ANs}$  and  $\Sigma\text{PNs}$ , the measurement by difference contains large  
5 uncertainties; (2) High  $\text{H}_2\text{O}_2$  for low- $\text{NO}$  conditions affected the operation of some CIMS  
6 instruments because the ppmv mixing ratios of  $\text{H}_2\text{O}_2$  depleted a non-negligible quantity of  
7 reagent ions. In order to correct for this, the CIMS instruments needed to calibrate as a  
8 function of  $\text{H}_2\text{O}_2$  in addition to traditional methods, or account for the true reagent ion signal  
9 (which was anti-correlated with  $\text{H}_2\text{O}_2$  concentration). High  $\text{H}_2\text{O}_2$  also affected GTHOS due to  
10 photolysis-derived OH production by the laser. GTHOS corrected for this effect by removing  
11 the OH background that was determined by sampling when only  $\text{H}_2\text{O}_2$  was present; (3) High  
12 formaldehyde, cyclohexane, or  $\text{H}_2\text{O}_2$  dominated the OH reactivity for certain experiments. In  
13 experiments where ppmv levels of volatile compounds were used, LIF-OHR and CRM-OHR  
14 did not operate. In contrast, high ozone and NO levels did not appear to affect the operation  
15 of any instruments. Temperature and humidity effects on ion sensitivities have been  
16 corrected for by ToFCIMS and TripCIMS as standard procedure. Other CIMS are actively  
17 characterizing these effects for analytes of interest.

18 Yet other analytical challenges were not unique to laboratory studies. It was found  
19 that chemical artifacts were produced from the decomposition of multifunctional OVOC  
20 (e.g., ISOPN, ISOPOOH, IEPOX, and pinonaldehyde) under normal operating conditions in  
21 some instruments; thus, possibly affecting ambient sampling and field data interpretation.  
22 Scheme 2 shows the proposed decomposition pathways of certain isomers of isoprene-  
23 derived OVOC to form MAC and MVK. We are aware of MAC and MVK interference only

1 from the 1,2- and 4,3- isomers of ISOPOOH, the 1,2- and 4,3- isomers and ISOPN, and the  
2 beta isomers of IEPOX (i.e., the peroxide, nitrate, and epoxide groups are secondary or  
3 tertiary). Unfortunately, these isomers are expected to be the most abundant in the  
4 atmosphere, e.g., the beta IEPOXs are estimated to represent > 97% of atmospheric IEPOX  
5 (Bates et al., 2014). The extent of decomposition and product distribution may also vary  
6 based on the operating conditions of the particular analytical method. In general, the  
7 decomposition was exacerbated by instruments with harsher sampling conditions, i.e., high  
8 ionization energy (e.g., the standard  $\text{H}_3\text{O}^+$  mode of SRI-ToFMS), high temperatures, and/or  
9 materials incompatible with organics (e.g., the hot stainless steel sample loop and inlet of  
10 GCFID). OVOCs from the low-NO isoprene photooxidation have been shown to decompose  
11 to MAC and MVK in commercial PTRMS instruments (Liu et al., 2013), but the exact  
12 identities of the compounds were unclear. During FIXCIT, it was observed that ISOPOOH,  
13 IEPOX, and pinonaldehyde were detected at  $m/z$  71.050 in the SRI-ToFMS in PTR mode  
14 (the sum of MAC+MVK). Switchable reagent ions show promise for removing certain  
15 biases, but more work is needed to characterize the chemistry that forms interfering ions.  
16 Further, we observed that the decomposition interference also affected GCFID, the other  
17 commonly-used detection method for MAC and MVK in ambient samples. ISOPOOH,  
18 IEPOX, and ISOPN were detected as either MAC or MVK in the GCFID, depending on the  
19 specific isomer. The interferences may not be localized to this particular GCFID, and a more  
20 detailed account is forthcoming (Rivera et al., 2014). Conversion efficiencies of OVOCs to  
21 the  $\text{C}_4$  carbonyls in the Caltech GCFID range in order of ISOPOOH > IEPOX > ISOPN, and  
22 can be almost quantitative for ISOPOOH because of the facile cleavage of the weak O-O

1 bond. Lastly, ISOPN were found to be converted to NO with a small yield in the NO-CL and  
2 a larger yield in commercial NO<sub>x</sub> analyzers.

3 All decomposition-derived artifacts can be avoided by collecting the air sample through a  
4 length of tubing submerged in a cold bath (– 40 °C), which trapped OVOCs that are less  
5 volatile than authentic MAC and MVK. Liu et al (2013) implemented this technique  
6 successfully in their laboratory study using SRI-ToFMS, resulting in a lower yield than  
7 previously reported for MAC and MVK in the low-NO oxidation of isoprene. Field  
8 application may prove more challenging, however, as the trapping is labor-intensive and  
9 requires careful humidity control to avoid ice buildup and blockage. During FIXCIT, both  
10 GCFID and SRI-ToFMS employed trapping techniques at various times to avoid biases in  
11 the detection and interpretation of MAC and MVK data.

12 **3. Preliminary results and atmospheric implications:** Forthcoming papers will discuss  
13 campaign results in detail. Here, we summarize a few interesting observations that appeared  
14 to be robust based on preliminary data analysis of the laboratory and field work.

- 15 • *Nighttime chemistry of alkenes, as controlled by the NO<sub>3</sub> radical, leads to several*  
16 *organic nitrates that are unique compared to daytime high-NO photooxidation. A*  
17 *significant product is the nitrooxy hydroperoxide, the atmospheric importance of*  
18 *which has likely been significantly underestimated in past chamber studies. The*  
19 *nitrooxy hydroperoxide reacts further in the daytime through a currently-unknown*  
20 *mechanism.*
- 21 • *The high-NO hydroxy nitrate yield from isoprene is closer to the high end of the*  
22 *spectrum (range 4 – 15%), important for the accurate simulations of volatile nitrogen*  
23 *in the atmosphere.*

- 1       • *Observed mixing ratios of isoprene low-NO photooxidation products are impacted by*  
2       *heterogeneous chemistry that appears to be mediated by aqueous processes, which*  
3       *has implications for the interpretation of IEPOX observations in dry vs. humid areas*  
4       *of the world.*
- 5       • *Environmental conditions in many locations, including within a biomass burning*  
6       *plume, are favorable for the H-shift RO<sub>2</sub> isomerization chemistry that produces*  
7       *compounds like HPALDs and very low-volatility oxygenates. The atmospheric fate of*  
8       *HPALDs is highly impacted by direct photolysis that recycles OH, as well as other*  
9       *complex chemistry and physical processes.*
- 10      • *The ozonolysis reaction of isoprene produces a high yield of C<sub>1</sub> compounds that are*  
11      *also observed with considerable abundance during ambient sampling. The*  
12      *hydroperoxide and acid yields appear to be underestimated by previous studies that*  
13      *detected these compounds via offline techniques. The OH yield may not follow the*  
14      *same trend with RH as the hydroperoxide and acid yields.*
- 15      • *APNs are efficient SOA precursors. SOA formation was prompt, and organic mass*  
16      *growth occurred quickly without the addition of inorganic seeds, i.e., the SOA*  
17      *intermediate(s) from APN+OH condensed onto predominantly-organic SOA*  
18      *material. Injections of the MAE standard did not increase the SOA mass growth.*
- 19      • *Several experiments produced significant amounts of excess OH, as measured by the*  
20      *GTHOS instrument, providing further avenues for investigation. These experiments*  
21      *also ruled out several candidates for the OH interference. More work is underway to*  
22      *comprehensively characterize the phenomenon.*

1 • *Calibrations with several synthesized standards of OVOC (Table 3) significantly aid*  
2 *in data interpretation from OHR and new CIMS instruments. Sampling these OVOC*  
3 *through standard instrumentation may interfere with some routine field and chamber*  
4 *measurements (depends on the run conditions and instrument setup), but may be*  
5 *mediated by cold-trapping methods. This is likely a contributing factor in the high*  
6 *discrepancies in MAC and MVK yields from low-NO isoprene photooxidation*  
7 *previously reported. For example, we find the preliminary low-NO yields of MVK ( $6$*   
8  *$\pm 3\%$ ) and MAC ( $4 \pm 2\%$ ), determined by GC-FID, from photooxidation of isoprene*  
9 *are consistent with Liu et al (2013) when cold-trapping methods were employed (Exp*  
10 *#21). However, the low-NO “yields” of MVK and MAC are each  $> 40\%$  when*  
11 *sampled directly by the GC-FID from the chamber (Exp #2) due to interferences by*  
12 *isomers of ISOPROOH (Rivera et al., 2014) and possibly other OVOCs.*

13 Final data from the FIXCIT campaign will be made publicly available on archives  
14 hosted by the U.S. National Oceanic and Atmospheric Administration (NOAA,  
15 <http://esrl.noaa.gov>) in January 2016. Data will be submitted in the ICARTT format,  
16 standardized by the U.S. National Aeronautics and Space Administration (NASA,  
17 <http://www-air.larc.nasa.gov/missions/etc/IcarttDataFormat.htm>).

#### 18 **4. Summary**

19 Although data analysis is ongoing, the goals of the FIXCIT campaign appear to have  
20 been met during the campaign period. The insights gained from experimental observations  
21 under well-controlled laboratory conditions have already proved valuable for understanding  
22 ambient observations from SOAS. The community effort to pursue atmospherically-  
23 important chemistry with sensitive ambient techniques and custom-synthesized chemicals

1 has elevated our understanding of atmospheric oxidation for a number of biogenic  
2 compounds. Novel mechanistic information obtained during FIXCIT will be helpful to  
3 update chemical mechanisms currently implemented in large-scale chemistry-coupled  
4 transport models. Instrumental inter-comparisons, an important aspect of the campaign, have  
5 demonstrated that a thorough characterization of new and standard ambient sampling  
6 techniques using authentic standards is necessary for accurate data interpretation.

7 Chamber experiments are clearly invaluable to the field of atmospheric chemistry, as  
8 the results feed directly into models that are used to ascertain regional and global climate and  
9 chemistry feedbacks. Further, chamber data aid in the interpretation of complex results  
10 obtained from field studies. Yet, it can be difficult to decipher the conditions under which  
11 chamber experiments are most relevant, and a standard protocol for data reporting may be  
12 needed. For example, best estimates of oxidation conditions in chambers (i.e., if reactions are  
13 HO<sub>2</sub>-dominated, low-NO but RO<sub>2</sub>-dominated, high-NO, high-NO<sub>x</sub> but low-NO, and so on)  
14 would greatly aid in comparisons of these experiments and others. The experiments in this  
15 campaign were fundamentally focused on the fate of the RO<sub>2</sub> radical as a delineation between  
16 chemical regimes. FIXCIT experiments (Table 2) can be further improved or tailored to the  
17 specific needs of the scientist. It has been demonstrated, here and elsewhere, that chamber  
18 studies that include chemistry representative of the atmosphere and well-characterized  
19 instrumental methods can accurately reproduce observations in the ambient environment.  
20 The results from FIXCIT make a case for future synergistic integration of laboratory studies  
21 with field campaigns, which maximizes the level of mechanistic understanding and data  
22 confidence obtained from the combination of both types of studies.


23



1 **Acknowledgements:** We acknowledge the collaborative efforts of FIXCIT participants (Tables  
2 2 – 3), as well as the organizers and logistics personnel for SOAS 2013. FIXCIT was made  
3 possible by the support from multiple agencies: U.S. National Science Foundation (NSF) under  
4 grants AGS-1240604 (Caltech), AGS-1246918 (PSU), AGS-1247421 (UWM), AGS-1243354  
5 (CU/ARI), AGS-1240611 (CSU), AGS-1120076 (UCB); U.S. Department of Energy under grant  
6 DE-SC0006626 (Caltech); U.S. Environmental Protection Agency (EPA) under STAR grant  
7 835407 (PNNL/UCB/SUNY). TBN (Caltech) was supported by NSF Postdoctoral Research  
8 Fellowship program, award AGS-1331360. NCAR is operated under the sponsorship of the NSF.

9


# 1 Schemes



2  
3 **Scheme 1:** Representative mechanism from the OH-, O<sub>3</sub>- and NO<sub>3</sub>-initiated oxidation of isoprene.  
4 The most abundant isomers of a particular pathway are shown. Red and blue arrows in the OH-  
5 oxidation scheme denote the NO-dominated and HO<sub>2</sub>-dominated RO<sub>2</sub> reactions, respectively. For  
6 the ozonolysis reaction, only the C<sub>1</sub> sCI and its reaction with water are shown as further-  
7 generation chemistry. For the NO<sub>3</sub>-oxidation pathway, only one isomer each of R and RO<sub>2</sub>  
8 radicals is shown for brevity. Abbreviations are defined in the text.

9

10



1

2 **Scheme 2:** Select proposed mechanism for the decomposition of OVOCs to carbonyls on contact

3 with metal surfaces or high ionization energies within instrumentation. Other decomposition

4 pathways likely exist and the branching ratios are dependent on instrument operation conditions.

5 Cleavage sites are indicated by dashed lines.

6

7

1 **Table 1:** List of participating instruments, principle investigators (PIs), and institutions. Key  
 2 acronyms: laser-induced fluorescence (LIF), laser-induced phosphorescence (LIP), high-  
 3 resolution time-of-flight (HRTof), compact time-of-flight (CTof), MS (mass spectrometer), and  
 4 CIMS (chemical ionization mass spectrometer).

Instrument	Instr. ID	PI(s)	Institutions	Measurements	Ref.
Ground-based Hydrogen Oxides Sensor	GTHOS	W.H. Brune	Pennsylvania State University (PSU)	OH, HO <sub>2</sub> , RO <sub>2</sub>	Brune et al., 1995
LIF OH reactivity monitor	LIF-OHR	W.H. Brune	PSU	OH reactivity by decay of OH	Mao et al., 2009
Thermal dissociation LIF NO <sub>2</sub> monitor	TDLIF	R.C. Cohen	University of California, Berkeley (UCB)	NO <sub>2</sub> , sum of organic nitrates (ΣANs), sum of peroxy nitrates (ΣPNs), particulate org. nitrates (pANs)	Day et al., 2002
Switchable iodide and acetate ion HRTof-CIMS	IACIMS	D.K. Farmer	Colorado State University (CSU)	Oxidized VOCs (organic nitrates, organic acids, etc.)	Lee et al., 2014a
NO <sub>3</sub> <sup>-</sup> HRTof-CIMS	NO <sub>3</sub> CIMS	M.R. Canagaratna, D.R. Worsnop, J.L. Jimenez	Aerodyne Research, Inc. (ARI) and Univ. of Colorado, Boulder (CUB)	Low-volatility organic compounds	Junninen et al., 2010
LIP glyoxal monitor	GlyLIP	F.N. Keutsch	University of Wisconsin, Madison (UWM)	Glyoxal	Huisman et al., 2008
LIF formaldehyde monitor	FormLIF	F.N. Keutsch	UWM	Formaldehyde	Hottle et al., 2008; DiGangi et al., 2011
Comparative rate method OH reactivity monitor	CRM-OHR	S. Kim, A.B. Guenther	Univ. of California, Irvine (UCI) and Pacific NW National Lab (PNNL)	OH reactivity by decay of hydrocarbons	Sinha et al., 2008
Switchable reagent ion (H <sub>3</sub> O <sup>+</sup> /NO <sup>+</sup> /O <sub>2</sub> <sup>+</sup> ) HRTof-MS	SRI-ToFMS	A.B. Guenther, J.E. Mak, A.H. Goldstein	PNNL, SUNY Stonybrook (SUNY), and UCB	Hydrocarbons, carbonyls, alcohols, etc.	Jordan et al., 2009
Chemical luminescence NO monitor	NO-CL	G.S. Tyndall, D.D. Montzka, A.J. Weinheimer	National Center for Atmospheric Research (NCAR)	NO (> 25 pptv)	Ridley and Grahek, 1990
CF <sub>3</sub> O <sup>-</sup> triple quadrupole CIMS	TripCIMS	P.O. Wennberg	California Institute of Technology (Caltech)	ISOPOOH, IEPOX, glycolaldehyde, acetic acid, methyl hydroperoxide	St. Clair et al., 2010

CF <sub>3</sub> O <sup>-</sup> CTof-CIMS	ToFCIMS	P.O. Wennberg	Caltech	Oxygenated VOCs (hydroperoxides, organic nitrates, multifunctional compounds)	Crouse et al., 2006
Gas chromatograph with ToFCIMS	GC-ToFCIMS	P.O. Wennberg	Caltech	Isomers for oxygenated VOCs	Bates et al., 2014
HRTof- aerosol mass spectrometer	ToF-AMS	J.H. Seinfeld	Caltech	Aerosol composition and size distribution	DeCarlo et al., 2006; Canagaratna et al., 2007
Gas chromatograph with flame-ionization detector	GCFID	J.H. Seinfeld	Caltech	Isoprene, methacrolein, methyl vinyl ketone, cyclohexane	N/A
Thermocouple and membrane probe	T/RH probe	J.H. Seinfeld	Caltech	Temperature and relative humidity	N/A
UV-absorption ozone monitor	O <sub>3</sub> monitor	J.H. Seinfeld	Caltech	O <sub>3</sub> (> 1000 pptv)	N/A
Chemical luminescence NOx detector	NOx monitor	J.H. Seinfeld	Caltech	NO (> 500 pptv), and NO <sub>2</sub> (catalytic conversion to NO)	N/A

1









2

1 **Table 2:** Formal experiments and reaction conditions during the campaign. Chemical  
2 abbreviations are defined in Table 3. Other abbreviations are: C1 = Chamber 1, C2 = Chamber 2,  
3 ISOP=isoprene,  $\alpha$ -PIN =  $\alpha$ -pinene, HP = hydrogen peroxide, MN = methyl nitrite, CHX =  
4 cyclohexane, HCHO = formaldehyde, AS = ammonium sulfate seeds. Exp. types are defined in  
5 the text. Exp. # corresponds to the date in Jan. 2014 when the experiment was performed.

#	Exp. type	HC precursor	[HC] (ppb)	Ox	Ox source	[OH] <sub>ss</sub> (# cm <sup>-3</sup> )	[O <sub>3</sub> ] <sub>i</sub> (ppb)	[NO] <sub>i</sub> (ppb)	[NO <sub>2</sub> ] <sub>i</sub> (ppb)	[NO] [HO <sub>2</sub> ]	Add'l Inj.	Rxn T (°C)	RH (%)
2	b	ISOP	45	OH	HP + hv	1.5 x 10 <sup>6</sup>	< 5	< 0.04	< 2	1/7	–	27	< 5
3	c	ISOP	100	OH	HP + hv	2.4 x 10 <sup>6</sup>	< 5	500	15	> 100	–	26	< 5
4a	i	ISOPOOHs	250	–	–	–	–	–	–	–	–	24	< 3
4b	a	Blank C1	0	OH	HP + hv	2.0 x 10 <sup>6</sup>	< 5	< 0.04	< 3	1/6	–	27 – 33	< 5
5a	i	ISOPNs	< 13	–	–	–	–	–	–	–	–	24	< 3
5b	a	Blank C2	0	OH	HP + hv	2.0 x 10 <sup>6</sup>	< 5	< 0.04	< 2	1/5	–	27	< 5
6	e	ISOP	91	O <sub>3</sub>	O <sub>3</sub> rxn	[OH] <sub>i</sub> ~ 1 x 10 <sup>6</sup>	615	< 0.04	< 3	–	–	25	< 5
7*	d	ISOP	30	OH	MN + hv	4.1 x 10 <sup>4</sup> , 4.8 x 10 <sup>6</sup>	< 5	0.08	45	2, 6	–	40, 40	< 5
9	f	ISOP	18	NO <sub>3</sub>	NO <sub>2</sub> /O <sub>3</sub>	3.8 x 10 <sup>8</sup>	55	0.10	100	2 – 3	HCHO	26	< 5
10	b	$\alpha$ -PIN	30	OH	HP + hv	2.0 x 10 <sup>6</sup>	< 5	< 0.04	< 2	1/10	–	27	< 5
11	c	$\alpha$ -PIN	30	OH	HP + hv	2.5 x 10 <sup>6</sup>	< 5	85	10	> 100	–	26	< 5
13	f	$\alpha$ -PIN	30	NO <sub>3</sub>	NO <sub>2</sub> /O <sub>3</sub>	4 x 10 <sup>8</sup>	75	0.17	150	1.5 – 8	HCHO	25	< 5
14	e	ISOP	100	O <sub>3</sub>	O <sub>3</sub> rxn	[OH] ~ 0	605	< 0.04	< 3	–	CHX	25	< 5
16*	d	$\alpha$ -PIN	30	OH	MN + hv	6 x 10 <sup>4</sup> , 4 x 10 <sup>6</sup>	< 5	0.08	< 3	2 – 3, 10	–	40, 40	< 5
17	b, i	4,3- ISOPOOH	60	OH	HP + hv	1.2 x 10 <sup>6</sup>	< 5	< 0.04	< 3	1/5	–	26	< 5
18*	d	ISOP	28	OH	MN + hv	1.0 x 10 <sup>5</sup> , 4.3 x 10 <sup>6</sup>	< 5	0.08	< 3	2 – 3, > 100	–	25, 26	< 5
19	b, h	ISOP	60	OH	HP + hv	1.0 x 10 <sup>6</sup>	< 5	< 0.04	< 4	1/5	wet AS	28	51
21	b	ISOP	22	OH	HP + hv	2.0 x 10 <sup>6</sup>	< 5	< 0.04	< 2	1/10	–	27	< 5
22	c	ISOP	100	OH	HP + hv	2.3 x 10 <sup>6</sup>	< 5	430	15	> 100	–	27	< 5
23	e	ISOP	90	O <sub>3</sub>	O <sub>3</sub> rxn	[OH] <sub>i</sub> ~ 1 x 10 <sup>6</sup>	600	< 0.04	< 3	–	–	25	50
24	c, h, i	4,3 - ISOPN	12	OH	HP + hv	3 x 10 <sup>6</sup>	7	115	55	> 100	wet AS	26	52
25	b	MAC	43	OH	HP + hv	3 x 10 <sup>6</sup>	< 5	< 0.03	< 3	1/10	–	28	< 5
26	g, h	MAC	45	OH	MN + hv	2 x 10 <sup>7</sup>	< 5	3.5	50	10 – 20	MAE, wet AS	26	< 5, 40
27	d, i	<i>trans</i> $\beta$ - IEPOX	60	OH	MN + hv	7.3 x 10 <sup>6</sup>	< 5	0.25	< 3	2 – 5	–	25	< 5
29	e	ISOP	91	O <sub>3</sub>	O <sub>3</sub> rxn	[OH] ~ 0	610	< 0.04	< 4	–	CHX	25	58
30	g, h, i	Pinonald.	15	OH	MN + hv	3.5 x 10 <sup>6</sup>	< 5	0.50	< 3	4 – 8	–	26	< 5

\* 1% lights, 20% lights, then 100% lights

1 **Table 3:** List of contributed synthesized chemical standards for experiments and calibration.


Synthesized standard	PIs	Institutions	Molecular structure	Atmos. Relevance	Synthesis Ref.
Isoprene 1-hydroxy 2-hydroperoxide (1,2 – ISOPOOH)	F.N. Keutsch	UWM		Major 1 <sup>st</sup> gen low-NO Isoprene + OH product	Rivera et al., 2014
Isoprene 3-hydroxy 4-hydroperoxide (3,4 – ISOPOOH)	F.N. Keutsch	UWM		Major 1 <sup>st</sup> gen low-NO Isoprene + OH product	Rivera et al., 2014
<i>trans</i> Isoprene 2-epoxydiol ( <i>trans</i> β-IEPOX)	P.O. Wennberg and J.H. Seinfeld	Caltech		Major 2 <sup>nd</sup> gen low-NO Isoprene + OH product	Bates et al., 2014
<i>cis</i> Isoprene 2-epoxydiol ( <i>cis</i> β-IEPOX)	P.O. Wennberg and J.H. Seinfeld	Caltech		Major 2 <sup>nd</sup> gen low-NO Isoprene + OH product	Bates et al., 2014
Isoprene 4-hydroxy 3-nitrate (4,3 – ISOPN)	R.C. Cohen, P.B. Shepson, A.S. Hasson, and P.O. Wennberg	UCB, Purdue Univ, CSU Fresno (CSUF), and Caltech		Major 1 <sup>st</sup> gen high-NO Isoprene + OH product	Lee et al., 2014b
Isoprene 2-hydroxy 1-nitrate (2,1 – ISOPN)	A.S. Hasson	CSUF		Minor 1 <sup>st</sup> gen high-NO Isoprene + OH product	N/A
Pinonaldehyde	P.O. Wennberg and J.H. Seinfeld	Caltech		Major 1 <sup>st</sup> gen α-Pinene + OH and O <sub>3</sub> product	Griesbaum et al., 1997
Methacrylic acid epoxide (MAE)	J.D. Surratt and A. Gold	Univ. of NC Chapel Hill (UNC)		Minor product and possible SOA precursor from MAC + OH + NO <sub>2</sub> reaction	Lin et al., 2013

2

1 **Figures**

2

3




4

5 **Figure 1:** Arrangement of instruments at the Caltech Atmospheric Chamber Facility during the  
6 campaign. Instrument IDs are in Table 1.

7


8





1  
 2 **Figure 2:** Progress of the slow chemistry experiment performed on 1/7/14. Isoprene data were  
 3 provided by GCFID. The red dashed line in the OH plot is the steady-state OH concentration  
 4 derived from the decay of isoprene as monitored by GCFID. OH and HO<sub>2</sub> preliminary data were  
 5 provided by GTHOS, using chemical zeroing, although the steady-state value of  $(0.4 - 1) \times 10^5$   
 6 molec cm<sup>-3</sup> was below the detection limit of GTHOS. OH preliminary data were averaged to  
 7 reduce noise. NO data were provided by NO-CL and OVOC data were provided by ToFCIMS.

8  
 9



1  
 2 **Figure 3:** GC-ToFCIMS chromatogram of ISOPNs from an isoprene high-NO photooxidation  
 3 experiment (black), and from the introduction of 2,1-ISOPN standard synthesized by CSUF  
 4 (cyan) and 4,3-ISOPN standards synthesized by Caltech (magenta), CSUF (green), UCB (blue),  
 5 and Purdue (red). The rightmost four peaks apparent in the photooxidation chromatogram are  
 6 preliminarily identified as the *cis* and *trans* 1,4-ISOPN and *cis* and *trans* 4,1-ISOPN, although  
 7 the elution order is not clear. Asterisks (\*) denote impurities in synthesized samples of  
 8 corresponding color.

9

## 1 References

- 2 Anglada, J. M., Gonzalez, J., and Torrent-Sucarrat, M.: Effects of the substituents on the reactivity of  
3 carbonyl oxides. A theoretical study on the reaction of substituted carbonyl oxides with water,  
4 Phys. Chem. Chem. Phys., 13, 13034-13045, 10.1039/c1cp20872a, 2011.
- 5 Atkinson, R., Carter, W. P., Winer, A. M., and Pitts Jr, J. N.: An experimental protocol for the  
6 determination of OH radical rate constants with organics using methyl nitrite photolysis as an OH  
7 radical source, J. Air Pollut. Contr. Assoc., 31, 1090-1092, 1981.
- 8 Atkinson, R., and Carter, W. P. L.: Kinetics and mechanisms of the gas-phase reactions of ozone with  
9 organic compounds under atmospheric conditions, Chem. Rev., 84, 437-470, 1984.
- 10 Atkinson, R., and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: A  
11 review, Atmospheric Environment, 37, 197-219, 2003.
- 12 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E.,  
13 Rossi, M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for  
14 atmospheric chemistry: Volume II - gas phase reactions of organic species, Atmos. Chem. Phys.,  
15 6, 3625-4055, 10.5194/acp-6-3625-2006, 2006.
- 16 Atkinson, R., E.C. Tuazon, and S.M. Aschmann: Products of the gas-phase reactions of O<sub>3</sub> with alkenes  
17 Environ. Sci. Technol., 29, 1860-1866, 1995.
- 18 Bates, K. H., Crouse, J. D., St. Clair, J. M., Bennett, N. B., Nguyen, T. B., Seinfeld, J. H., Stoltz, B. M.,  
19 and Wennberg, P. O.: Gas phase production and loss of isoprene epoxydiols, J. Phys. Chem. A,  
20 118, 1237-1246, 10.1021/jp4107958, 2014.
- 21 Beaver, M. R., Clair, J. M. S., Paulot, F., Spencer, K. M., Crouse, J. D., LaFranchi, B. W., Min, K. E.,  
22 Pusede, S. E., Wooldridge, P. J., Schade, G. W., Park, C., Cohen, R. C., and Wennberg, P. O.:  
23 Importance of biogenic precursors to the budget of organic nitrates: Observations of  
24 multifunctional organic nitrates by cims and TD-LIF during BEARPEX 2009, Atmos. Chem.  
25 Phys., 12, 5773-5785, 10.5194/acp-12-5773-2012, 2012.
- 26 Becker, K. H., Brockmann, K. J., and Bechara, J.: Production of hydrogen peroxide in forest air by  
27 reaction of ozone with terpenes, Nature, 346, 256-258, 1990.
- 28 Bertman, S. B., and Roberts, J. M.: A PAN analog from isoprene photooxidation, Geophys. Res. Lett., 18,  
29 1461-1464, 1991.
- 30 Biskos, G., Paulsen, D., Russell, L. M., Buseck, P. R., and Martin, S. T.: Prompt deliquescence and  
31 efflorescence of aerosol nanoparticles, Atmos. Chem. Phys., 6, 4633-4642, 10.5194/acp-6-4633-  
32 2006, 2006.
- 33 Brune, W. H., Stevens, P. S., and Mather, J. H.: Measuring OH and HO<sub>2</sub> in the troposphere by laser-  
34 induced fluorescence at low pressure, J. Atmos. Sci., 52, 3328-3336, 10.1175/1520-  
35 0469(1995)052<3328:moahit>2.0.co;2, 1995.
- 36 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B.,  
37 Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway,  
38 M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and  
39 microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer,  
40 Mass Spectrom. Rev., 26, 185-222, 10.1002/mas.20115, 2007.
- 41 Chan, A. W. H., Chan, M. N., Surratt, J. D., Chhabra, P. S., Loza, C. L., Crouse, J. D., Yee, L. D.,  
42 Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Role of aldehyde chemistry and NO<sub>x</sub>  
43 concentrations in secondary organic aerosol formation, Atmos. Chem. Phys., 10, 7169-7188,  
44 doi:10.5194/acp-10-7169-2010, 2010.
- 45 Claeys, M., Wang, W., Ion, A. C., Kourtev, I., Gelencser, A., and Maenhaut, W.: Formation of  
46 secondary organic aerosols from isoprene and its gas-phase oxidation products through reaction  
47 with hydrogen peroxide, Atmos. Environ., 38, 4093-4098, 10.1016/j.atmosenv.2004.06.001,  
48 2004.
- 49 Criegee, R.: Mechanism of ozonolysis, Angew. Chem., 87, 765-771, 1975.

- 1 Crouse, J. D., McKinney, K. A., Kwan, A. J., and Wennberg, P. O.: Measurement of gas-phase  
2 hydroperoxides by chemical ionization mass spectrometry, *Anal. Chem.*, 78, 6726-6732,  
3 10.1021/ac0604235, 2006.
- 4 Crouse, J. D., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: Peroxy radical isomerization in the  
5 oxidation of isoprene, *Phys. Chem. Chem. Phys.*, 13, 13607-13613, 2011.
- 6 Crouse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., and Wennberg, P. O.: Autoxidation of  
7 organic compounds in the atmosphere, *J. Phys. Chem. Lett.*, 4, 3513-3520, 10.1021/jz4019207,  
8 2013.
- 9 Day, D. A., Wooldridge, P. J., Dillon, M. B., Thornton, J. A., and Cohen, R. C.: A thermal dissociation  
10 laser-induced fluorescence instrument for in situ detection of NO<sub>2</sub>, peroxy nitrates, alkyl nitrates,  
11 and HNO<sub>3</sub>, *J. Geophys. Res.*, 107, 4046, 10.1029/2001jd000779, 2002.
- 12 DeCarlo, P.F., Kimmel, J.R., Trimborn, A., Northway, M.J., Jayne, J.T., Aiken, A.C., Gonin, M., Fuhrer,  
13 K., Horvath, T., Docherty, K., Worsnop, D.R. and Jimenez, J.L. Field-Deployable, High-  
14 Resolution, Time-of-Flight Aerosol Mass Spectrometer. *Anal. Chem.*, 78: 8281-8289, 2006.
- 15 DiGangi, J., Boyle, E., Karl, T., Harley, P., Turnipseed, A., Kim, S., Cantrell, C., Maudlin Iii, R., Zheng,  
16 W., and Flocke, F.: First direct measurements of formaldehyde flux via eddy covariance:  
17 Implications for missing in-canopy formaldehyde sources, *Atmos. Chem. Phys.*, 11, 10565-  
18 10578, 2011.
- 19 Donahue, N. M., Drozd, G. T., Epstein, S. A., Presto, A. A., and Kroll, J. H.: Adventures in ozoneland:  
20 Down the rabbit-hole, *Phys. Chem. Chem. Phys.*, 13, 10848-10857, 2011.
- 21 Drozd, G. T., and Donahue, N. M.: Pressure dependence of stabilized Criegee intermediate formation  
22 from a sequence of alkenes, *J. Phys. Chem. A*, 115, 4381-4387, 10.1021/jp2001089, 2011.
- 23 Eddingsaas, N., Loza, C., Yee, L., Seinfeld, J., and Wennberg, P.: A-pinene photooxidation under  
24 controlled chemical conditions—part 1: Gas-phase composition in low-and high-NO<sub>x</sub>  
25 environments, *Atmos. Chem. Phys.*, 12, 6489-6504, 2012.
- 26 Ehn, M., Thornton, J. A., Kleist, E., Sipila, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F.,  
27 Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T.,  
28 Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurten, T., Nielsen, L. B.,  
29 Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petaja, T., Wahner,  
30 A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source  
31 of low-volatility secondary organic aerosol, *Nature*, 506, 476-479, 10.1038/nature13032, 2014.
- 32 Fenske, J. D., Hasson, A. S., Ho, A. W., and Paulson, S. E.: Measurement of absolute unimolecular and  
33 bimolecular rate constants for CH<sub>3</sub>CHO generated by the trans-2-butene reaction with ozone in  
34 the gas phase, *J. Phys. Chem. A*, 104, 9921-9932, 2000.
- 35 Fuentes, J. D., Lerda, M., Atkinson, R., Baldocchi, D., Bottenheim, J. W., Ciccioli, P., Lamb, B., Geron,  
36 C., Gu, L., Guenther, A., Sharkey, T. D., and Stockwell, W.: Biogenic hydrocarbons in the  
37 atmospheric boundary layer: A review, *Bull. Amer. Meteor. Soc.*, 81, 1537-1575, 2000.
- 38 Gäb, S., Hellpointner, E., Turner, W. V., and Korte, F.: Hydroxymethyl hydroperoxide and  
39 bis(hydroxymethyl) peroxide from gas-phase ozonolysis of naturally occurring alkenes, *Nature*,  
40 316, 535-536, 1985.
- 41 Garden, A. L., Paulot, F., Crouse, J. D., Maxwell-Cameron, I. J., Wennberg, P. O., and Kjaergaard, H.  
42 G.: Calculation of conformationally weighted dipole moments useful in ion–molecule collision  
43 rate estimates, *Chem. Phys. Lett.*, 474, 45-50, 2009.
- 44 Griesbaum, K., Dong, Y., and McCullough, K. J.: Ozonolyses of acetylenes: Trapping of a-oxo carbonyl  
45 oxides by carbonyl compounds and stabilization of a-oxo ozonides by derivatizations, *J. Org.*  
46 *Chem.*, 62, 6129-6136, 1997.
- 47 Griffin, R. J., Cocker, D. R., III, Flagan, R. C., and Seinfeld, J. H.: Organic aerosol formation from the  
48 oxidation of biogenic hydrocarbons, *J. Geophys. Res.*, 104, 3555-3567, 1999.
- 49 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerda,  
50 M., and et al.: A global model of natural volatile organic compound emissions, *J. Geophys. Res.*,  
51 100, 8873-8892, 1995.

1 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.:  
2 The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): An  
3 extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-  
4 1492, 10.5194/gmd-5-1471-2012, 2012.

5 Hansel, A., Jordan, A., Holzinger, R., Prazeller, P., Vogel, W., and Lindinger, W.: Proton transfer  
6 reaction mass spectrometry: Online trace gas analysis at the ppb level, *Int. J. Mass. Spectrom. Ion*  
7 *Processes*, 149/150, 609-619, 1995.

8 Hasson, A. S., Ho, A. W., Kuwata, K. T., and Paulson, S. E.: Production of stabilized Criegee  
9 intermediates and peroxides in the gas phase ozonolysis of alkenes 2. Asymmetric and biogenic  
10 alkenes, *J. Geophys. Res.*, 106, 34143-34153, 2001.

11 Henze, D. K., Seinfeld, J. H., Ng, N. L., Kroll, J. H., Fu, T. M., Jacob, D. J., and Heald, C. L.: Global  
12 modeling of secondary organic aerosol formation from aromatic hydrocarbons: High- vs. Low-  
13 yield pathways, *Atmos. Chem. Phys.*, 8, 2405-2420, 10.5194/acp-8-2405-2008, 2008.

14 Hoffmann, T., Odum, J. R., Bowman, F., Collins, D., Klockow, D., Flagan, R. C., and Seinfeld, J. H.:  
15 Formation of organic aerosols from the oxidation of biogenic hydrocarbons, *J. Atmos. Chem.*, 26,  
16 189-222, 1997.

17 Hottle, J. R., Huisman, A. J., DiGangi, J. P., Kamrath, A., Galloway, M. M., Coens, K. L., and Keutsch,  
18 F. N.: A laser induced fluorescence-based instrument for in-situ measurements of atmospheric  
19 formaldehyde, *Environ. Sci. Technol.*, 43, 790-795, 10.1021/es801621f, 2008.

20 Huang, D., Chen, Z., Zhao, Y., and Liang, H.: Newly observed peroxides and the water effect on the  
21 formation and removal of hydroxyalkyl hydroperoxides in the ozonolysis of isoprene, *Atmos.*  
22 *Chem. Phys.*, 13, 5671-5683, doi:10.5194/acp-13-5671-2013, 2013.

23 Huisman, A. J., Hottle, J. R., Coens, K. L., DiGangi, J. P., Galloway, M. M., Kamrath, A., and Keutsch,  
24 F. N.: Laser-induced phosphorescence for the in situ detection of glyoxal at part per trillion  
25 mixing ratios, *Anal. Chem.*, 80, 5884-5891, 2008.

26 Jacobs, M. I., Darer, A. I., and Elrod, M. J.: Rate constants and products of the OH reaction with  
27 isoprene-derived epoxides, *Environ. Sci. Technol.*, 47, 12868-12876, 2013.

28 Jacobs, M. I., Burke, W. J., and Elrod, M. J.: Kinetics of the reactions of isoprene-derived  
29 hydroxynitrates: Gas phase epoxide formation and solution phase hydrolysis, *Atmos. Chem.*  
30 *Phys. Discuss.*, 14, 12121-12165, 10.5194/acpd-14-12121-2014, 2014.

31 Johnson, D., and Marston, G.: The gas-phase ozonolysis of unsaturated volatile organic compounds in the  
32 troposphere, *Chem. Soc. Rev.*, 37, 699-716, 2008.

33 Jordan, A., Haidacher, S., Hanel, G., Hartungen, E., Herbig, J., Märk, L., Schotchkowsky, R., Seehauser,  
34 H., Sulzer, P., and Märk, T.: An online ultra-high sensitivity proton-transfer-reaction mass-  
35 spectrometer combined with switchable reagent ion capability (PTR+ SRI-MS), *Int. J. Mass*  
36 *Spectrom.*, 286, 32-38, 2009.

37 Junninen, H., Ehn, M., Petäjä, T., Luosujärvi, L., Kotiaho, T., Kostianinen, R., Rohner, U., Gonin, M.,  
38 Fuhrer, K., Kulmala, M., and Worsnop, D. R.: A high-resolution mass spectrometer to measure  
39 atmospheric ion composition, *Atmos. Meas. Tech.*, 3, 1039-1053, 10.5194/amt-3-1039-2010,  
40 2010.

41 Kjaergaard, H. G., Knap, H. C., Ørnsø, K. B., Jørgensen, S., Crounse, J. D., Paulot, F., and Wennberg, P.  
42 O.: Atmospheric fate of methacrolein. 2. Formation of lactone and implications for organic  
43 aerosol production, *J. Phys. Chem. A*, 116, 5763-5768, 10.1021/jp210853h, 2012.

44 Kroll, J. H., Clarke, J. S., Donahue, N. M., Anderson, J. G., and Demerjian, K. L.: Mechanism of HO<sub>x</sub>  
45 formation in the gas-phase ozone-alkene reaction. 1. Direct, pressure-dependent measurements of  
46 prompt OH yields, *J. Phys. Chem. A*, 105, 1554-1560, 2001.

47 Kroll, J. H., Donahue, N. M., Cee, V. J., Demerjian, K. L., and Anderson, J. G.: Gas-phase ozonolysis of  
48 alkenes: Formation of OH from *anti* carbonyl oxides, *J. Amer. Chem. Soc.*, 124, 8518-8519,  
49 2002.

- 1 Kuwata, K. T., Hermes, M. R., Carlson, M. J., and Zogg, C. K.: Computational studies of the  
2 isomerization and hydration reactions of acetaldehyde oxide and methyl vinyl carbonyl oxide, *J.*  
3 *Phys. Chem. A*, 114, 9192-9204, 10.1021/jp105358v, 2010.
- 4 Kwan, A., Chan, A., Ng, N. L., Kjærgaard, H. G., Seinfeld, J. H., and Wennberg, P. O.: Peroxy radical  
5 chemistry and OH radical production during the NO<sub>3</sub>-initiated oxidation of isoprene, *Atmos.*  
6 *Chem. Phys.*, 12, 7499-7515, 2012.
- 7 Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Kurtén, T., Worsnop, D. R., and Thornton, J. A.: An iodide-  
8 adduct high-resolution time-of-flight chemical-ionization mass spectrometer: Application to  
9 atmospheric inorganic and organic compounds, *Environ. Sci. Technol.*, 10.1021/es500362a,  
10 2014a.
- 11 Lee, J. H., Leahy, D. F., Tang, I. N., and Newman, L.: Measurement and speciation of gas phase  
12 peroxides in the atmosphere, *J. Geophys. Res.*, 98, 2911-2915, 10.1029/92jd02514, 1993.
- 13 Lee, L., Teng, A. P., Wennberg, P. O., Crounse, J. D., and Cohen, R. C.: On rates and mechanisms of OH  
14 and O<sub>3</sub> reactions with isoprene-derived hydroxy nitrates, *J. Phys. Chem. A*, 10.1021/jp4107603,  
15 2014b.
- 16 Lee, M., Heikes, B. G., and O'Sullivan, D. W.: Hydrogen peroxide and organic hydroperoxide in the  
17 troposphere: A review, *Atmos. Environ.*, 34, 3475-3494, [http://dx.doi.org/10.1016/S1352-](http://dx.doi.org/10.1016/S1352-2310(99)00432-X)  
18 [2310\(99\)00432-X](http://dx.doi.org/10.1016/S1352-2310(99)00432-X), 2000.
- 19 Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Harder, H.,  
20 Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation  
21 capacity sustained by a tropical forest, *Nature*, 452, 737-740, 10.1038/nature06870, 2008.
- 22 Lin, Y.-H., Zhang, Z., Docherty, K. S., Zhang, H., Budisulistiorini, S. H., Rubitschun, C. L., Shaw, S. L.,  
23 Knipping, E. M., Edgerton, E. S., Kleindienst, T. E., Gold, A., and Surratt, J. D.: Isoprene  
24 epoxydiols as precursors to secondary organic aerosol formation: Acid-catalyzed reactive uptake  
25 studies with authentic compounds, *Environ. Sci. Technol.*, 46, 250-258, 10.1021/es202554c,  
26 2012.
- 27 Lin, Y.-H., Zhang, H., Pye, H. O. T., Zhang, Z., Marth, W. J., Park, S., Arashiro, M., Cui, T.,  
28 Budisulistiorini, S. H., Sexton, K. G., Vizuete, W., Xie, Y., Luecken, D. J., Piletic, I. R., Edney,  
29 E. O., Bartolotti, L. J., Gold, A., and Surratt, J. D.: Epoxide as a precursor to secondary organic  
30 aerosol formation from isoprene photooxidation in the presence of nitrogen oxides, *Proc. Nat.*  
31 *Acad. Sci.*, 10.1073/pnas.1221150110, 2013.
- 32 Liu, Y. J., Herdinger-Blatt, I., McKinney, K. A., and Martin, S. T.: Production of methyl vinyl ketone  
33 and methacrolein via the hydroperoxyl pathway of isoprene oxidation, *Atmos. Chem. Phys.*, 13,  
34 5715-5730, 10.5194/acp-13-5715-2013, 2013.
- 35 Loza, C. L., Chan, A. W. H., Galloway, M. M., Keutsch, F. N., Flagan, R. C., and Seinfeld, J. H.:  
36 Characterization of vapor wall loss in laboratory chambers, *Environ. Sci. Technol.*, 44, 5074-  
37 5078, 10.1021/es100727v, 2010.
- 38 Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, A., Huey, L. G., Cohen, R. C.,  
39 Heikes, B., Singh, H. B., Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R., and Shetter, R.  
40 E.: Airborne measurement of OH reactivity during INTEX-B, *Atmos. Chem. Phys.*, 9, 163-173,  
41 10.5194/acp-9-163-2009, 2009.
- 42 Mao, J., Ren, X., Zhang, L., Van Duin, D. M., Cohen, R. C., Park, J. H., Goldstein, A. H., Paulot, F.,  
43 Beaver, M. R., Crounse, J. D., Wennberg, P. O., DiGangi, J. P., Henry, S. B., Keutsch, F. N.,  
44 Park, C., Schade, G. W., Wolfe, G. M., Thornton, J. A., and Brune, W. H.: Insights into hydroxyl  
45 measurements and atmospheric oxidation in a California forest, *Atmos. Chem. Phys.*, 12, 8009-  
46 8020, 10.5194/acp-12-8009-2012, 2012.
- 47 Martin, S. T.: Phase transitions of aqueous atmospheric particles, *Chem. Rev.*, 100, 3403-3453, 2000.
- 48 Neeb, P., Sauer, F., Horie, O., and Moortgat, G. K.: Formation of hydroxymethyl hydroperoxide and  
49 formic acid in alkene ozonolysis in the presence of water vapor, *Atmos. Environ.*, 31, 1417-1423,  
50 1997.

- 1 Ng, N. L., Kwan, A. J., Surratt, J. D., Chan, A. W. H., Chhabra, P. S., Sorooshian, A., Pye, H. O. T.,  
2 Crounse, J. D., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol  
3 (SOA) formation from reaction of isoprene with nitrate radicals ( $\text{NO}_3$ ), *Atmos. Chem. Phys.*, 8,  
4 4117-4140, 2008.
- 5 Nguyen, T. B., Coggon, M. M., Bates, K. H., Zhang, X., Schwantes, R. H., Schilling, K. A., Loza, C. L.,  
6 Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Organic aerosol formation from the reactive  
7 uptake of isoprene epoxydiols (IEPOX) onto non-acidified inorganic seeds, *Atmos. Chem. Phys.*,  
8 14, 3497-3510, 10.5194/acp-14-3497-2014, 2014.
- 9 Niki, H., Maker, P. D., Savage, C. M., and Breitenbach, L. P.: Further ir spectroscopic evidence for the  
10 formation of hydroperoxyhydroxymethane in the gas phase reaction of hydroperoxy radical with  
11 formaldehyde, *Chem. Phys. Lett.*, 75, 533-535, 1980.
- 12 Nouaime, G., Bertman, S., Seaver, C., Elyea, D., Huang, H., Shepson, P., Starn, T., Riemer, D., Zika, R.,  
13 and Olszyna, K.: Sequential oxidation products from tropospheric isoprene chemistry: MACR  
14 and MPAN at a  $\text{NO}_x$ -rich forest environment in the southeastern United States, *J. Geophys. Res.*,  
15 103, 22463-22471, 1998.
- 16 Nozière, B., and Barnes, I.: Evidence for formation of a PAN analogue of pinonic structure and  
17 investigation of its thermal stability, *J. Geophys. Res.*, 103, 25587-25597, 1998.
- 18 Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kroll, J. H., Seinfeld, J. H., and Wennberg, P. O.: Isoprene  
19 photooxidation: New insights into the production of acids and organic nitrates, *Atmos. Chem.*  
20 *Phys.*, 9, 1479-1501, 10.5194/acp-9-1479-2009, 2009a.
- 21 Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kurten, A., St. Clair, J. M., Seinfeld, J. H., and Wennberg,  
22 P. O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325,  
23 730-733, 10.1126/science.1172910, 2009b.
- 24 Peeters, J., Nguyen, T. L., and Vereecken, L.: HO<sub>x</sub> radical regeneration in the oxidation of isoprene,  
25 *Phys. Chem. Chem. Phys.*, 11, 5935-5939, 2009.
- 26 Perring, A. E., Wisthaler, A., Graus, M., Wooldridge, P. J., Lockwood, A. L., Mielke, L. H., Shepson, P.  
27 B., Hansel, A., and Cohen, R. C.: A product study of the isoprene+ $\text{NO}_3$  reaction, *Atmos. Chem.*  
28 *Phys.*, 9, 4945-4956, 2009.
- 29 Pye, H. O. T., Pinder, R. W., Piletic, I. R., Xie, Y., Capps, S. L., Lin, Y.-H., Surratt, J. D., Zhang, Z.,  
30 Gold, A., Luecken, D. J., Hutzell, W. T., Jaoui, M., Offenberg, J. H., Kleindienst, T. E.,  
31 Lewandowski, M., and Edney, E. O.: Epoxide pathways improve model predictions of isoprene  
32 markers and reveal key role of acidity in aerosol formation, *Environ. Sci. Technol.*, 47, 11056–  
33 11064, 10.1021/es402106h, 2013.
- 34 Ridley, B., and Grahek, F.: A small, low flow, high sensitivity reaction vessel for NO chemiluminescence  
35 detectors, *J. Atmos. Ocean. Tech.*, 7, 307-311, 1990.
- 36 Rivera, J., Apel, E., Crounse, J. D., de Gouw, J. A., Hansel, A., Jud, W., Kaser, L., Kaiser, J. B., Nguyen,  
37 T. B., St. Clair, J. M., Wennberg, P. O., Wisthaler, A., and Keutsch, F. N.: Investigation of  
38 pristine isoprene oxidation products reveals observational bias toward modern conditions,  
39 Submitted, *Geophys. Res. Lett.*, 2014.
- 40 Roberts, J. M., Williams, J., Baumann, K., Buhr, M. P., Goldan, P. D., Holloway, J., Hübler, G., Kuster,  
41 W. C., McKeen, S. A., and Ryerson, T. B.: Measurements of PAN, PPN, and MPAN made  
42 during the 1994 and 1995 Nashville intensives of the southern oxidant study: Implications for  
43 regional ozone production from biogenic hydrocarbons, *J. Geophys. Res.*, 103, 22473-22490,  
44 1998.
- 45 Rollins, A. W., Kiendler-Scharr, A., Fry, J. L., Brauers, T., Brown, S. S., Dorn, H. P., Dubé, W. P.,  
46 Fuchs, H., Mensah, A., Mentel, T. F., Rohrer, F., Tillmann, R., Wegener, R., Wooldridge, P. J.,  
47 and Cohen, R. C.: Isoprene oxidation by nitrate radical: Alkyl nitrate and secondary organic  
48 aerosol yields, *Atmos. Chem. Phys.*, 9, 6685-6703, 10.5194/acp-9-6685-2009, 2009.
- 49 Sauer, F., Schäfer, C., Neeb, P., Horie, O., and Moortgat, G. K.: Formation of hydrogen peroxide in the  
50 ozonolysis of isoprene and simple alkenes under humid conditions, *Atmos. Environ.*, 33, 229-  
51 241, 1999.

- 1 Simonaitis, R., Olszyna, K., and Meagher, J.: Production of hydrogen peroxide and organic peroxides in  
2 the gas phase reactions of ozone with natural alkenes, *Geophys. Res. Lett.*, 18, 9-12, 1991.
- 3 Sinha, V., Williams, J., Crowley, J., and Lelieveld, J.: The comparative reactivity method—a new tool to  
4 measure total OH reactivity in ambient air, *Atmos. Chem. Phys.*, 8, 2213-2227, 2008.
- 5 St. Clair, J. M., McCabe, D. C., Crouse, J. D., Steiner, U., and Wennberg, P. O.: Chemical ionization  
6 tandem mass spectrometer for the in situ measurement of methyl hydrogen peroxide, *Rev. Sci.  
7 Instrum.*, 81, 094102-094106, 2010.
- 8 Surratt, J., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan,  
9 R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic  
10 aerosol formation from isoprene, *Proc. Nat. Acad. Sci.*, 107, 6640-6645, 2010.
- 11 Valverde-Canossa, J., Ganzeveld, L., Rappenglück, B., Steinbrecher, R., Klemm, O., Schuster, G., and  
12 Moortgat, G.: First measurements of H<sub>2</sub>O<sub>2</sub> and organic peroxides surface fluxes by the relaxed  
13 eddy-accumulation technique, *Atmos. Environ.*, 40, 55-67, 2006.
- 14 Vereecken, L., Müller, J.-F., and Peeters, J.: Low-volatility poly-oxygenates in the OH-initiated  
15 atmospheric oxidation of  $\alpha$ -pinene: Impact of non-traditional peroxy radical chemistry, *Phys.  
16 Chem. Chem. Phys.*, 9, 5241-5248, 2007.
- 17 Welz, O., Savee, J. D., Osborn, D. L., Vasu, S. S., Percival, C. J., Shallcross, D. E., and Taatjes, C. A.:  
18 Direct kinetic measurements of Criegee intermediate (CH<sub>2</sub>OO) formed by reaction of CH<sub>2</sub>I with  
19 O<sub>2</sub>, *Science*, 335, 204-207, 10.1126/science.1213229, 2012.
- 20 Wängberg, I., Barnes, I., and Becker, K. H.: Product and mechanistic study of the reaction of NO<sub>3</sub>  
21 radicals with  $\alpha$ -pinene, *Environ. Sci. Technol.*, 31, 2130-2135, 10.1021/es960958n, 1997.
- 22 Wolfe, G. M., Thornton, J. A., Yatavelli, R. L. N., McKay, M., Goldstein, A. H., LaFranchi, B., Min, K.  
23 E., and Cohen, R. C.: Eddy covariance fluxes of acyl peroxy nitrates (PAN, PPN and MPAN)  
24 above a ponderosa pine forest, *Atmos. Chem. Phys.*, 9, 615-634, 10.5194/acp-9-615-2009, 2009.
- 25 Wolfe, G. M., Thornton, J. A., Bouvier-Brown, N. C., Goldstein, A. H., Park, J. H., McKay, M., Matross,  
26 D. M., Mao, J., Brune, W. H., LaFranchi, B. W., Browne, E. C., Min, K. E., Wooldridge, P. J.,  
27 Cohen, R. C., Crouse, J. D., Faloon, I. C., Gilman, J. B., Kuster, W. C., de Gouw, J. A.,  
28 Huisman, A., and Keutsch, F. N.: The chemistry of atmosphere-forest exchange (CAFE) model –  
29 part 2: Application to BEARPEX-2007 observations, *Atmos. Chem. Phys.*, 11, 1269-1294,  
30 10.5194/acp-11-1269-2011, 2011.
- 31 Wolfe, G. M., Crouse, J. D., Parrish, J. D., Clair, J. M. S., Beaver, M. R., Paulot, F., Yoon, T. P.,  
32 Wennberg, P. O., and Keutsch, F. N.: Photolysis, OH reactivity and ozone reactivity of a proxy  
33 for isoprene-derived hydroperoxyenals (HPALDs), *Phys. Chem. Chem. Phys.*, 14, 7276-7286,  
34 2012.
- 35 Worton, D. R., Surratt, J. D., LaFranchi, B. W., Chan, A. W. H., Zhao, Y., Weber, R. J., Park, J.-H.,  
36 Gilman, J. B., de Gouw, J., Park, C., Schade, G., Beaver, M., Clair, J. M. S., Crouse, J.,  
37 Wennberg, P., Wolfe, G. M., Harrold, S., Thornton, J. A., Farmer, D. K., Docherty, K. S.,  
38 Cubison, M. J., Jimenez, J.-L., Frossard, A. A., Russell, L. M., Kristensen, K., Glasius, M., Mao,  
39 J., Ren, X., Brune, W., Browne, E. C., Pusede, S. E., Cohen, R. C., Seinfeld, J. H., and Goldstein,  
40 A. H.: Observational insights into aerosol formation from isoprene, *Environ. Sci. Technol.*, 47,  
41 11403-11413, 10.1021/es4011064, 2013.

42  
43

44