Observational estimates of detrainment and entrainment in non-precipitating
 shallow cumulus

- ³ Matthew S. Norgren¹, Jennifer D. Small², Haflidi H. Jonsson³, Patrick Y. Chuang^{4*}
- 4 1. Dept. of Physics, University of California Santa Cruz, Santa Cruz, CA, USA
- 5 2. Department of Meteorology, University of Hawaii at Manoa, Honolulu, HI, USA
- 6 3. Center for Interdisciplinary Remotely-Piloted Aircraft Studies, Naval Postgraduate School,

7 CA, USA

- 8 4. Earth and Planetary Sciences, University of California Santa Cruz, Santa Cruz, CA, USA
- ${}_{9}$ *Corresponding author email: pchuang@ucsc.edu
- 10

11 Abstract

Vertical transport associated with cumulus clouds is important to the redistribution of gases. 12 particles and energy, with subsequent consequences for many aspects of the climate system. 13 Previous studies have suggested that detrainment from clouds can be comparable to the 14 updraft mass flux, and thus represents an important contribution to vertical transport. In 15 this study, we describe a new method to deduce the amounts of gross detrainment and 16 entrainment experienced by non-precipitating cumulus clouds using aircraft observations. 17 The method utilizes equations for three conserved variables: cloud mass, total water and 18 moist static energy. Optimizing these three equations leads to estimate of the mass fractions 19 of adiabatic mixed-layer air, entrained air and detrained air that the sampled cloud has 20 experienced. The method is applied to six flights of the CIRPAS Twin Otter during the Gulf 21 of Mexico Atmospheric Composition and Climate Study (GoMACCS) which took place in the 22 Houston, Texas region during the summer of 2006 during which 176 small, non-precipitating 23 cumulus were sampled. Using our novel method, we find that, on average, these clouds 24 were comprised of 30 to 70% mixed-layer air, with entrained air comprising most of the 25 remainder. The mass fraction of detrained air was usually very small, less than 2%, although 26 values larger than 10% were found in 15% of clouds. Entrained and detrained air mass 27 fractions both increased with altitude, consistent with some previous observational studies. 28 The largest detrainment events were almost all associated with air that was at their level 29 of neutral buoyancy, which has been hypothesized in previous modeling studies. This new 30 method could be readily used with data from other previous aircraft campaigns to expand 31 our understanding of detrainment for a variety of cloud systems. 32

33 1 Introduction

One of the important ways cumulus clouds affect the atmosphere is through vertical trans-34 port. The redistribution of gases, particles and energy that originate at or near the Earth's 35 surface to altitudes above the mixed layer is important for a range of phenomena relevant 36 to Earth's atmosphere and climate. For example, the vertical profile of water vapor is crit-37 ical to longwave heating and cooling profiles, as well as to the subsequent development and 38 evolution of clouds [Malkus, 1954]. The long-range transport and atmospheric lifetime of 39 particulates and trace gases are enhanced when they are at higher altitudes due to decreased 40 probability of wet deposition. Aerosol scattering and absorption are also altitude-dependent, 41 in particular their altitude relative to that of any cloud layers e.g. Liao and Seinfeld, 1998; 42 Chand et al., 2009; Samset and Myhre, 2011]. The amount of air that passes through a 43 cloud strongly impacts the degree to which aerosols and gases can be processed via in-cloud 44 liquid-phase reactions. Lack of understanding of the effects of vertical transport is a primary 45 source of uncertainty in climate models [Rougier et al., 2009]. 46

In cumulus clouds, vertical transport can be approximately separated into two modes: (1) the detrainment of cloudy air to the surrounding environment during the cloud's active period, i.e. when there is dynamical support for the cloud; and (2) the mixed-layer air that remains after the cloud loses dynamical support and dissipates. While there is some ambiguity in separating these two modes, it's helpful to make this distinction because the first has historically been the subject of greater study, even though the latter can potentially dominate [*Wang and Geerts*, 2011].

Detrainment is typically used to describe the process by which cloudy air is transferred outside of the cloud volume, i.e. to the surrounding environment [*Dawe and Austin*, 2011]. Detrainment has been divided into two types [*de Rooy and Siebesma*, 2010]. The first is *turbulent detrainment* and is due to turbulent mixing along the cloud boundary. When cloudy air turbulently mixes with unsaturated environmental air such that the resulting parcel is

unsaturated and not completely surrounded by cloud (i.e. is connected to the sub-saturated 59 cloud environment), then the cloudy air has been detrained. A second kind of detrainment 60 has been termed dynamical detrainment (or cloud outflow) because it is driven by organized 61 circulations comparable to the length scale of the cloud rather than smaller turbulent eddies. 62 Such detrainment has been related to buoyancy gradient profiles that cause deceleration and 63 flow divergence [Bretherton and Smolarkiewicz, 1989; de Rooy and Siebesma, 2010], and also 64 to the flow structure of a shedding thermal [Taylor and Baker, 1991; Blyth, 1993; Zhao and 65 Austin, 2005; Blyth et al., 2005]. 66

There is not an extensive history of observational studies of detrainment in clouds (Wang and 67 Geerts, 2011; see also a recent review by de Rooy et al., 2013), and the various methods and 68 clouds types from these studies have yielded a range of views on the process. Some obser-69 vational estimates come from mass budget studies where, using aircraft flying closed circuits 70 around individual cumulus (Cu), mass and moisture budgets are inferred, from which en-71 trainment and detrainment rates at different levels of the cloud are deduced [Raymond and 72 Wilkening, 1982, 1985; Raga et al., 1990; Raymond et al., 1991; Barnes et al., 1996]. These 73 studies typically find that the net detrainment mass flux (defined as the difference between 74 the gross detrainment and entrainment mass fluxes) can be comparable in magnitude to the 75 updraft mass flux, albeit with strong variability with height and in time. One important 76 mechanism of detrainment deduced from these studies is a detraining outflow in collapsing 77 turrets, where air sinks until reaching its level of neutral buoyancy and then diverges out-78 wards from the cloud, causing detrainment to occur only at specific altitudes. Using aircraft 79 observations of summertime cumuli off of Hawaii (with typical cloud depths of $\sim 2 \text{ km}$), Raga 80 et al. [1990] found that net detrainment occurred only in the top one-third of the cloud, 81 with the lower parts exhibiting net entrainment. Raymond et al. [1991] combined aircraft 82 and radar observations of summertime thunderstorm clouds over New Mexico (cloud depths 83 ranging between 6 and 12 km) and found a similar vertical pattern of detrainment predomi-84 nantly in the upper portion of clouds. Barnes et al. [1996] studied summertime cumulus and 85

cumulus congestus (cloud depths up to 4 km) near coastal Florida. USA, using two coordi-86 nated aircraft flying at different altitudes. They found that detrainment varied greatly with 87 time, with the same layer changing from net entrainment to net detrainment, or vice versa, 88 on the order of a few minutes. Perry and Hobbs [1996] found evidence for regions of enhanced 89 humidity "halos" in shallow maritime cumulus (typical cloud depths between 0.5 and 2.5 km) 90 off the coast of northeast continental USA, particularly on the downshear side. These regions 91 exhibiting enhanced humidities were typically 1 to 2 cloud radii in length, and increased in 92 size with cloud age. This result is suggestive of active detrainment in cumulus clouds, al-93 though the results do not rule out the possibility that these halos are remnants of previous 94 clouds. In contrast, Wang and Geerts [2011] studied orographic cumulus mediocris in Ari-95 zona, USA (typical cloud depth of 2 km) and found no evidence for continuous detrainment; 96 their measurements downwind of a cloud field are instead consistent with vertical transport 97 dominated by evaporation of the clouds themselves rather than active detrainment by the 98 clouds. We note that these studies are performed in different environments (e.g. clear air 99 relative humidity) with varying cumulus cloud sizes, and thus the results are not necessarily 100 expected to be consistent with each other. 101

One assumption that mass budget-based studies make is that the accumulation term is neg-102 ligible, i.e. the cloud is at steady state with respect to mass. However, Carpenter et al. 103 [1998b, a] find that the accumulation term can be dominant which implies a potentially large 104 source of uncertainty for the inferred detrainment rates in some observational studies. An-105 other limitation is that these mass budget studies only yield net entrainment or detrainment; 106 these values are not necessarily reflective of gross entrainment and detrainment rates which 107 could be much higher than the net value. For example, there could be no net detrainment 108 (mass loss) from a cloud if it is exactly balanced elsewhere by an equal amount of entrain-109 ment. Gross detrainment values are, however, of greater relevance for understanding vertical 110 transport. 111

¹¹² Entrainment, in comparison to detrainment, is a much more familiar topic in the cloud physics

literature and thus we only highlight a few studies out of many. Entrainment can be defined 113 as the incorporation of air originating outside the cloud volume into the cloud, thus increasing 114 total cloud mass and volume. It is one of the key processes governing the microphysical struc-115 ture and macrophysical properties of a cloud, and along with precipitation, is responsible for 116 the depletion of cloud water mixing ratio and thus is relevant to cloud lifetime. Entrainment, 117 as with detrainment, can be similarly divided into turbulent and dynamical forms *Houghton* 118 and Cramer, 1951, and evidence exists supporting the importance of both processes. En-119 trainment associated with organized flow has been described using observations [e.g. Stith, 120 1992; Damiani and Vali, 2007] and models [e.g. Zhao and Austin, 2005; Blyth et al., 2005]. 121 Through analysis of aircraft observations, Wang et al. [2009] show that the outermost 10% of 122 cumulus clouds, i.e. cloud edges, are on average strongly depleted in liquid water relative to 123 the interior of the cloud, supporting the idea that turbulent entrainment occurs along outer 124 surface of the cloud, but not ruling out the possibility of localized entrainment that is then 125 transported to other regions by, e.g., the descending outer shell. 126

In this study, we will use a novel approach to estimate total *gross* detrainment and entrainment that has occurred in shallow, non-precipitating cumulus clouds. This method is not able to inform the mechanism for detrainment and entrainment (e.g. cloud-scale dynamical features versus small-scale turbulence), and instead focuses on quantifying the amount of each as a function of height.

132 2 Method

133 2.1 Aircraft Data

Data gathered during August and September 2006 as part of the Gulf of Mexico Atmospheric Composition and Climate Study (GoMACCS) is used in this study. The GoMACCS field campaign included 22 research flights carried out by the Twin Otter aircraft [Lu et al., 2008]

operated by the Center for Interdisciplinary Remotely-Piloted Aircraft Studies (CIRPAS). 137 The flights were conducted over land in a region outside of Houston, Texas. Of 22 total flight 138 days, data from six days (Sept. 1, 2, 8, 11, 14, 15) are analyzed in this study. These six 139 days are selected due to a sufficient number of randomly sampled clouds, and all relevant 140 instrumentation functioned properly during the flights. The sampled clouds are small, warm, 141 non-precipitating continental Cu that typically first form in the late-morning due to surface 142 heating. Sampled cloud sizes are typically 1 to 2 km in width and depth (see Table 1). Later 143 in the afternoon, deeper convection can be triggered but these events were avoided during 144 these flights. Because of the proximity of the flights to a very large city (Houston, TX) and 145 the many industrial activities in the region, aerosol concentrations are high (accumulation 146 mode aerosol concentrations ranging from 400 to 1600 $\rm cm^{-3}$) and contribute to the lack of 147 precipitation from these clouds. More information about the conditions encountered during 148 these flights can be found in $Lu \ et \ al.$ [2008]. 149

The clouds are sampled in random fashion during a series of constant altitude legs, each 150 about 10 min in duration. This is done by flying the Twin Otter through approximately the 151 center of the nearest appropriate cloud as judged visually by the pilots, with factors such 152 as aircraft turn capabilities, and cloud size and appearance being considered. Of course, 153 clouds are irregularly shaped so exactly where the pilot chooses to penetrate each cloud is 154 not easily defined. This introduces uncertainty in our analysis (as discussed in more detail 155 in Section 2.4 below). Figure 1 shows the altitude profile for the Sept. 8 flight, which is 156 representative of all flight days. A number of level legs can be seen in the altitude profile. 157 For each flight, between 3 to 5 of these correspond to the cloud layer and therefore include 158 a number of cloud penetrations. Note that because of this statistical sampling strategy, no 159 effort is specifically made to sample a cloud more than once. Also of note is the continuous 160 ascent from below cloud base, ~ 300 m, to above cloud top, ~ 4800 m, which is utilized in 161 the analysis as our clear-air sounding and which we assume is representative of clear air in 162 the vicinity of all our sampled clouds over the course of the sampling period. Variation of 163

this sounding, either in space or over time, can cause uncertainties in our analysis. Typical aircraft speed is 55 m s⁻¹, and we primarily employ 10 Hz (or 5.5 m) data sets.

In situ measurements of temperature, specific humidity (q_v) and liquid water content (LWC) 166 are needed for our analysis. Temperature was measured using a Rosemount 102E4AL sen-167 sor with 0.4°C accuracy. In clear air, specific humidity is derived from 1-Hz dew point 168 temperature measurements made by a chilled mirror dew point hydrometer with (Edgetech 169 Dewpointer 137-C3 with 0.2°C accuracy). In-cloud specific humidity values are assumed to 170 be saturated at the measured temperature. Clouds are identified using a minimum LWC 171 threshold of 0.05 g kg⁻¹, as measured by a Gerber Particle Volume Monitor 100A instrument 172 [Gerber et al., 1994]. Total specific water (q_t) is the sum of q_v and LWC (none of the sam-173 pled clouds was cold enough for ice to form). Cloud penetrations with LWC satisfying the 174 threshold requirements for a minimum of six seconds, which corresponds to an approximate 175 cloud sample length of 330 m, are identified as clouds and used for this study. The minimum 176 cloud size requirement is imposed so that the clouds used in the study contain enough data 177 points to conduct analyses with reasonable statistics. Figure 1 shows the LWC profile for the 178 flight on Sept. 8. On this day, 27 clouds were sampled across the various levels, with a mean 179 cloud penetration length of 660 m. Table 1 gives cloud number and size information for each 180 flight day. 181

In-cloud wetting of the aircraft temperature probe does not appear to be a significant issue 182 during GoMACCS [Small et al., 2009]. Further analysis was done for this study to confirm 183 this result. We see no statistically significant bias in clear-air temperature prior to entering 184 a cloud and after leaving the same clouds as would be expected for a wetted temperature 185 probe; $\Delta T = T_{exit} - T_{entry}$ has values of 0.18 K, 0.20 K and 0.20 K when comparing the 1 s, 186 3 s and 5 s intervals prior to cloud entry and after departure from the same cloud, which 187 are smaller than the 1σ values of ΔT of 0.35 K, 0.35 K and 0.34 K, respectively. The biases 188 are also smaller than the probe accuracy of 0.4 K. Wetting of the Rosemount probe has 189 been found to result from impaction of drops on the probe housing, which leads to liquid 190

water build up and eventually drops spraying in random directions, some of which reach 191 the probe sensor [Lawson and Cooper, 1990]. Wetting biases have been found to depend on 192 cloud LWC and aircraft speed. In the case of GoMACCS, cloud LWC is generally below 193 0.8 g m^{-3} , from which a typical bias of 0.4 K would be deduced based on clouds sampled by 194 Lawson and Cooper [1990]. However, in this study the aircraft speed is about half of that 195 from Lawson and Cooper [1990], which will reduce drop impaction to the probe housing. 196 Impaction also depends on drop size, and during GoMACCS drops are small due to the large 197 aerosol concentrations, with typical sizes much less than 20 μ m [Small et al., 2009]. Lastly, 198 wetting also depends on the cloud width. GoMACCS clouds exhibit widths between 1 and 199 2 km, which are on the small side for cumuli. We attribute the lack of any statistically 200 significant wetting of the Rosemount during GoMACCS to the combination of all of these 201 factors. 202

203 2.2 Adiabatic Clouds

In order to develop a model of gross entrainment and detrainment, we first explore their effects on an idealized adiabatic cloud. If a parcel of air rises adiabatically, by definition it will exchange neither mass nor energy with the environment. Thus, the mass and energy of the air parcel will be conserved. This also implies that the moist static energy (or MSE) of the parcel also is conserved.

Entrainment/detrainment, precipitation, and radiation are the primary processes which can cause cloudy air parcels to deviate from adiabaticity. Entrainment increases the total mass of the cloud while decreasing mean q_t and MSE. This occurs because, relative to clear air at the same altitude, cloudy air is generally warmer (because it is positively buoyant) and moister (because it is cloudy) [e.g. *Wang et al.*, 2009], although the former may not always be true during the cumulus dissipation stage. For a cloud experiencing detrainment, the total mass of the cloud decreases. In our analysis, we assume that the properties of the detrained air are a function of the cloudy air and adiabatic air properties, which tends to cause the cloud MSE and q_t to either stay constant or decrease (depending on the exact set of assumptions; see Section 2.4 below for more details). However, the potential decrease in MSE and q_t differs for the same amount of entrained or detrained air, which allows the analysis to distinguish between the two processes.

Precipitation could affect cloud properties, but the focus of this study is on non-precipitating clouds, so this is not an important consideration. The clouds sampled did not precipitate due to the combination of polluted aerosol conditions from the Houston region and the limited depth of the clouds which limits cloud liquid water path [*Small et al.*, 2009].

Net emitted radiation from a cloud causes cooling and therefore decreases MSE, while net 225 absorption warms. During the daytime (when the research flights took place), the net ra-226 diative balance for each cloud is determined by the difference between longwave cooling and 227 shortwave heating, which tend to be similar in magnitude. We will assume no net change 228 due to radiation. The bias in cloud temperature, and hence MSE, caused by this assumption 229 is likely to be very small. If we assume a $20 \text{ W} \text{m}^{-2}$ imbalance, and a mean cloud lifetime of 230 30 min, the mean temperature change for a 1-km deep cloud will be a few hundredths of a 231 Kelvin and thus unlikely to be a large source of uncertainty in this analysis. 232

In the absence of substantial effects by precipitation and radiation, we are left with only entrainment and detrainment as the processes capable of altering clouds mass, MSE and q_t from the initial adiabatic values.

236 2.3 Conserved Variables

Our analysis of detrainment and entrainment in cumulus clouds is based on the conservation of three variables: mass, q_t , and moist static energy. The total mass of a cloud, M_c , is the sum of all gases, liquids, and solids contained within the volume of the cloud. The total specific water of a cloud parcel (q_t) is the sum of the liquid water and the water vapor, given 241 by:

$$q_t = q_v + q_l \tag{1}$$

where q_v is the specific humidity and q_l is the specific liquid water, both in units of g kg⁻¹. Again, these clouds are warm, so Eq. 1 excludes ice. Total water is conserved for an adiabatic process because there is no mass exchange with the environment, and therefore q_t is constant. Moist static energy s is a measure of an air parcel's energy in units J kg⁻¹ and to good approximation is conserved during adiabatic ascent/descent:

$$s = c_p T + gh + q_v L_v \tag{2}$$

where T is absolute temperature, the heat capacity of moist air $c_p = c_p(q_v) = c_{pd}(1 + 0.9q_v)$ where c_{pd} is the heat capacity of dry air (assumed to be a constant value 1005 J kg⁻¹ K⁻¹), g is the gravitational acceleration, h is the height of the air parcel above sea level, q_v is the specific humidity, and $L_v = 2260$ kJ kg⁻¹ is the latent heat of vaporization of water (we ignore the effects of temperature on L_v because they are small). Typical uncertainties in calculated s are a few tenths of a percent based on instrumental uncertainties.

As a cloud parcel is lifted along the dry adiabat, the increase in potential energy is accompanied by a decrease in the sensible heat term; the parcel cools as it increases in height. If the parcel is saturated and liquid water is present, the decrease in q_v due to condensation is offset by the release of latent heat, increasing the parcel temperature. We have argued above (Section 2.2) that processes such as precipitation and net radiation flux divergence that can cause MSE to not be conserved are likely negligible in this study.

259 2.4 Conservation Equations

For the clouds chosen in this analysis, we assume that each cloud has a mass that is determined by the balance of three terms (see Figure 2 for a schematic): (a) air that has been adiabatically lifted from near the surface; (b) air that has entrained into the cloud; and (c) air that has detrained from the cloud. Starting with this simple model, we make two important assumptions in order to proceed with the analysis:

1. Entrainment occurs perfectly laterally, so that all the entrained air in the cloud at
the aircraft sampling altitude originates from clear air at the same altitude. A recent
review paper [de Rooy et al., 2013] argues that "lateral entrainment is the dominant
mixing mechanism in comparison with the cloud-top entrainment in shallow cumulus
convection," an idea with a long history (see references and discussion in de Rooy et al.,
2013) supported by recent LES-based studies [Heus et al., 2008; Yeo and Romps, 2013].
We will test the sensitivity of our results to this assumption.

2. Two end-member scenarios for detrainment are (a) that detrainment occurred exactly 272 at the same time as the aircraft penetration of the cloud, i.e. detrainment happened at 273 the last possible moment; and (b) that detrainment occurred when the cloud properties 274 were nearly adiabatic (before substantial entrainment has occurred), i.e. detrainment 275 happened very early during cloud formation. The corresponding properties of the de-276 trained air for these end-members would be (a) detrained air has the identical properties 277 as the cloud at the sampled level and (b) detrained air has the identical properties as 278 the adiabatic mixed layer air. In this analysis, we assume that the detrained air has 279 properties represented by the *mean* of these two end-members, which is intended to 280 represent a middle scenario. We will again test the sensitivity of our results to this 281 assumption. 282

With these assumptions, we can now write conservation equations describing our system. We apply our analysis to each cloud penetration because, as previously stated, each cloud is only sampled once. Thus, the analysis results apply only to each cloud at the level of aircraft sampling, as illustrated in Figure 2, and not to the entire cloud. By mass conservation, the mass of the thin cloud slice M_c can be given by:

$$M_a + M_o - M_d = M_c \tag{3}$$

where the subscript a is mixed-layer air risen <u>a</u>diabatically, o is laterally entrained air (air originating <u>outside</u> the cloud), d is laterally <u>d</u>etrained air, and c is aircraft-sampled <u>cloudy</u> air. Dividing Eq. 3 by M_c , we obtain:

$$m_a + m_o - m_d = 1 \tag{4}$$

where we have now written the equation in terms of mass fractions $m_a = M_a/M_c$, $m_o = M_o/M_c$, and $m_d = M_d/M_c$. Working with mass fractions is more convenient and useful for the purpose of comparing results among different clouds because the results do not explicitly depend on the cloud mass. Furthermore, given our cloud sampling method, we would need to make assumptions about cloud shape in order to determine M_c , introducing more sources of error.

We note that in Section 2.2, detrainment was defined as an active process of turbulence or 297 organized circulations removing air from a cloud. By defining the conservation of mass as 298 we do in Eq. 4, any air that is within the cloud but then later becomes external to the cloud 299 is considered detrained air. Thus, detrainment as defined by this analysis can occur either 300 actively, where cloudy air is transferred outside the cloud via organized flow or turbulence, 301 or passively, where enough air is entrained into the cloud to lower the LWC below our cloud 302 threshold LWC. The latter would not normally be considered detrainment but rather cloud 303 dissipation, but it is relevant to vertical mass transport as described in the Introduction. 304

We can also construct a conservation equation for the moist static energy of our sampled cloud:

$$s_a M_a + s_o M_o - s_d M_d = s_c M_c \tag{5}$$

where s is MSE and the same subscripts from Eq. 3 apply. The adiabatic air MSE, s_a , is 307 computed from the lowest (by altitude) 200 data points on each given flight day. These points 308 are all in the surface mixed layer, which is generally well-mixed because all flights occurred 309 around the middle of the day when the continental convective boundary layer exhibits strong 310 turbulence. The MSE of entrained air s_o is taken from the clear air sounding acquired 311 during each flight. Due to our assumption of lateral entrainment, s_o is taken to be the MSE 312 value of the clear air at the altitude of the cloud penetration. The MSE of the cloud slice 313 \boldsymbol{s}_c is determined as the mean MSE derived from the aircraft observations for each cloud 314 penetration. By assumption #2 above, the MSE of the air that detrains is $s_d = (s_a + s_c)/2$. 315 Again dividing by M_c to write in terms of mass fractions: 316

$$s_a m_a + s_o m_o - s_d m_d = s_c \tag{6}$$

Eq. 6 thus contains the same unknowns, m_a , m_o and m_d , as Eq. 4, but with MSE coefficients that are determined from aircraft measurements. A third equation based on conservation of total specific water can also be derived in the same way as for MSE:

$$q_a m_a + q_o m_o - q_d m_d = q_c \tag{7}$$

The conservation equations are re-written as a set of non-linear equations in order to restrict the mass fractions to positive, physically-plausible solutions:

$$x^2 + y^2 - z^2 - 1 = 0 \tag{8}$$

$$c_1 x^2 + c_2 y^2 - c_3 z^2 - 1 = 0 (9)$$

323

$$d_1x^2 + d_2y^2 - d_3z^2 - 1 = 0 (10)$$

where $x^2 = m_a$, $y^2 = m_o$, and $z^2 = m_d$ are the three unknowns, while the coefficients are computed from aircraft observations as: $c_1 = s_a/s_c$, $c_2 = s_o/s_c$, $c_3 = s_d/s_c$, and $d_1 = q_a/q_c$, $d_2 = q_o/q_c$, $d_3 = q_d/q_c$. To solve for the three unknowns m_a , m_o and m_d , we perform the following:

(a) For <u>each cloud penetration</u>, we use in-cloud observations to compute the mean moist static energy s_c and mean total specific water q_c .

(b) We use aircraft observations to compute the properties of the end-member air masses, i.e. s_a and q_a (adiabatic mixed-layer air), s_o and q_o (entrained air) and s_d and q_d (detrained air). See the discussion following Eq. 5 for details on how this is done.

(c) Using the results from (a) and (b), we can calculate all the coefficients c_i and d_i in Eqs. 9 and 10, respectively.

(d) We use non-linear optimization which minimizes the residuals for the system of Equations 8 to 10 to determine a best estimate for $x^2 = m_a$, $y^2 = m_o$ and $z^2 = m_d$ for each penetration. The magnitude of the total residual is an estimate of the uncertainty in the solution.

This method weights each data point of the cloud penetration equally in calculating mean penetration values of c_i and d_i in Eqs. 9 and 10. However, this can potentially bias the results because in reality a cloud slice is two-dimensional, whereas the penetration is onedimensional. If we assume the cloud slice is circular in cross-section, air sampled during the penetration near the cloud edge is representative of a much larger area than air sampled at the cloud center. Our analysis, then, potentially biases the data towards values near the center of the cloud and under-represents data from cloud edges [Hoffmann et al., 2014]. However, the aircraft may not always sample the exact center of a cloud, and still assuming clouds are circular in shape, a cloud penetration not through the center of the cloud may possibly over-represent the cloud edge data. To evaluate these potential effects on our analysis, we also solve for m_a , m_o , and m_d using only the cloud properties from the first and last second (~55 m) of the cloud penetration (i.e. in computing c_i and d_i in Eqs. 9 and 10), which focuses the analysis strictly on air near the cloud edge.

352 3 Results and Discussion

353 3.1 Individual flight day results

Figs. 3 and 4 show example results from the optimizations for two of the six flight days. On each plot, the left panel plots the mass fraction of detrained air m_d (in units of percent), while the right panel plots the mass fraction of entrained air into the cloud, m_o , both as a function of altitude, with one point for each cloud penetration. There are a total of 176 penetrations over the six days analyzed. The clear-air soundings of MSE and q_t for the flight day are also given on the left and right side, respectively.

The success of the optimization is measured by deviation of the three conservation equations (Eqs. 8 to 10) from zero. The combined total error is calculated as:

$$\epsilon_T = \sqrt{\epsilon_M^2 + \epsilon_E^2 + \epsilon_Q^2} \tag{11}$$

where ϵ_T represents the total root-mean square error associated with the individual residuals from the mass, MSE and moisture equations (ϵ_M , ϵ_E and ϵ_Q respectively). The cloud marker sizes in Figs. 3 and 4 for m_d and m_o are inversely proportional to the value of ϵ_T . Therefore, the largest markers correspond to clouds with optimizations that yielded the smallest residuals in Eqs. 8 to 10. Note that these equations are all order unity due to the normalization. For all clouds sampled, ϵ_T had a median value of 0.07, a mean value of 0.15, and a standard deviation of 0.11.

369 3.1.1 Detrained air

Our analysis indicates that the sampled non-precipitating cumulus clouds exhibit m_d values 370 that are below 2%, although there are a number of cases when some substantially higher m_d 371 values are inferred. Figure 5 shows the distribution of m_d for all flight days (176 clouds). 372 The majority (78%) of cloud penetrations exhibit m_d values below 2%, while 15% of clouds 373 have a m_d value above 10%. Only two events exhibit m_d values larger than 18%, and the 374 largest m_d value was 68%. On almost all days (results not all shown), the biggest m_d values 375 are found at the highest sampling altitudes. The one exception is on Sept. 11 when some 376 larger m_d values are found in the middle part of the clouds. Small (<2%) m_d values were 377 found at all levels, but made up a larger fraction of the observations at lower portions of the 378 clouds. 379

To compare the vertical distribution of detrained air among different days, all cloud penetra-380 tion altitudes are normalized with respect to cloud base and cloud top altitude for each flight 381 day. The clouds are then sorted into 5 evenly spaced normalized altitude (\hat{z}) bins, and for 382 each bin a mean \hat{z} and m_d is computed. All clouds were weighted equally, and the penetration 383 length through each cloud was not factored into the mean m_d calculation. The uncertainty in 384 \hat{z} on a day-by-day basis is likely small compared to the \hat{z} bin spacing. Cloud base altitude is 385 easily determined within ~ 100 m from in situ measurements. Cloud top altitude is less easily 386 determined by the pilot, but the uncertainty is likely modest compared to the total cloud 387 layer depth as cloud top is usually constrained by a temperature and/or humidity inversion. 388 Figure 6 shows that, in the mean, m_d does tend to increase with altitude, although the upper 389 portions of the cloud tend to exhibit a lot of variability. The mean values are not large at any 390 altitude, with the smallest value of 1% closest to cloud base and a maximum in the highest 391

 \hat{z} bin of less than 5%, and an overall mean of 3%.

It is noteworthy that few large m_d values are observed, with only one value over 25%. All 393 clouds analyzed here primarily dissipate by evaporation because they are not precipitating. 394 At the end of a cloud's life, we expect m_d to be equal m_c , since at this point the cloud has 395 dissipated. While completely dissipated clouds are not the target for this analysis, we might 396 expect to see some high m_d values associated with clouds near the end of their life cycle. 397 However, high values of m_d were inferred only once in this study. One potential reason is 398 that the pilots may have considered strongly dissipating clouds to be visually unappealing 399 targets. In a cloud field with many choices of cloud targets, such a bias in pilot judgment 400 could bias our statistical sampling. The constraint that clouds must have sample lengths over 401 330 m to be considered for analysis may also contribute to limiting m_d values. A dissipating 402 cloud whose diameter shrinks to less than 330 m will not yet have reached the point where 403 $m_d = m_c$. Alternately, as noted earlier, previous studies [e.g. Carpenter et al., 1998a] have 404 inferred that detrainment occurs at specific levels within clouds. Because we only sampled 405 one level of each cloud, we may not have been sampling at the level that detrainment was 406 occurring. 407

408 3.1.2 Entrained air

The mass fraction of entrained air within a cloud, m_o , typically ranges from 30 to 70% (illustrated in Figs. 3 and 4). Figure 7 shows the m_o distribution for all flight days. The median m_o is 45%, the mean is 49%, and a standard deviation of 14%. The full range is between 20 and 90%. The amount of entrained air is considerably more than than the mass of detrained air composing a cloud, and there is only one cloud that exhibits m_d greater than m_o .

⁴¹⁵ A vertical profile of m_o for each day is created in the same manner as the one for m_d and is ⁴¹⁶ shown in Fig. 8. This plot shows that m_o tends to be larger in the upper portion of clouds, with mean values between 50 and 55% in the upper half of the clouds (normalized altitudes $\hat{z} > 0.5$), compared to mean values around 40 to 45% in the lower half of the clouds. As with the detrainment fraction, there is substantial variability at each level.

These results in general seem physically reasonable. The large values of m_o are consistent 420 with Barnes et al. [1996] which showed that the entrainment fluxes can be similar to or larger 421 than the vertical mass fluxes. Relatively large values of m_o can occur within these clouds 422 because the high humidity of the surrounding environmental air in south Texas ($q_t \sim 10$ to 423 16 $g kg^{-1}$) in the cloud layer means that the drying effect from entrainment is not as strong as 424 it would be in much drier environments such as New Mexico or Colorado (which have been the 425 setting for numerous previous cumulus studies). In a drier environment, a large entrainment 426 fraction would lead to the complete dissipation of the cloud. The wide range of m_o values is 427 consistent with having sampled clouds at different stages of their life cycle, which one would 428 expect from random aircraft sampling of clouds (even considering the possible bias against 429 strongly dissipating clouds discussed above). The increase in m_o with altitude is consistent 430 with the common observation that the adiabaticity (ratio of the measured cloud LWC to 431 adiabatic LWC) in these clouds decreases with height (e.g. Lu et al., 2008), although drying 432 of the environmental air with altitude may also play a role. Greater entrainment in the 433 upper-portion of the cloud is also consistent with the shedding thermal picture of cumulus 434 growth [e.g. Kitchen and Caughey, 1981; Blyth et al., 2005], where entrained air creates the 435 subsiding shell of cold air at the periphery of the cloud. This air is entrained into the cloud 436 somewhere below cloud top, and is subsequently transported to higher levels in the buoyant 437 updraft. 438

The overall picture that emerges from our analysis, then, is that the sampled clouds are composed of roughly equal parts entrained air and adiabatic mixed-layer air, and have detrained relatively little of their mass, although a minority (15%) exhibit appreciable amounts of detrainment (above 10% mass fraction). Both entrainment and detrainment mass fractions tend to increase with altitude. We next examine how robust these results are to the assumptions made in the analysis.

445 3.2 Sensitivity tests

446 3.2.1 Cloud-edge only

A straight-line penetration of a cloud can potentially misrepresent the area-averaged cloud properties by biasing the measurements in a number of ways. As described in section 2.4, one such bias is to emphasize the interior of the cloud at the expense of cloud edge. To see how much an effect this has on the optimized parameters, we re-ran the optimizations using data only sampled from the outermost 50 m at the edge of the cloud. The resulting ranges of m_d and m_o (not shown) are not changed significantly, suggesting that such a bias did not affect our analysis.

454 3.2.2 Entrainment source level

We previously made the assumption that entrainment occurs only laterally at each sampling level. Although this is an oversimplification of the entrainment process, and thus is a limitation of this model, there exists justification for this assumption. As discussed above (Section 2.4), support for lateral entrainment as the primary mechanism has gained substantial support [de Rooy et al., 2013].

We performed sensitivity tests of our model to the assumed source level of entrained air. In simulations of cumulus congestus with cloud height of 8 km, *Yeo and Romps* [2013] find that entrained air within the cloud at each height can be traced to air in the environment at an altitude of 1 to 2 km lower, at least during the mature and dissipating stages. If we assume self-similarity in the vertical direction, then for the clouds in this study (with depths of 1 to 2 km), the equivalent entrainment altitude is a few hundred meters below the sampling level. Thus, we test the sensitivity of our results by performing the optimization was using MSE and q_t soundings that are shifted upwards or downwards in altitude by 400 m. Since the MSE and q_t soundings, in general, exhibit a decrease with height, this has the effect of changing the MSE and q_t of the source of entrained air.

An example of this sensitivity test is shown in Fig. 9, where the entrainment altitude is shifted upwards by 400 m for the flight of Sept. 8 (compare these results with Fig. 4). There is some increase in m_d for some of the penetrations, although for others, lower m_d is deduced. The mean m_d is nearly the same, with mean and σ of 2.6% and 5.1% for the original analysis, and 2.4% and 5.0% for the shifted sounding analysis. Using geometric mean instead of arithmetic mean also yields strong similarity between the two analyses.

Shifting the source level of entrained air upwards decreases the entrainment mass fraction 476 m_o . The decrease in m_o is expected, because by effectively entraining air from a higher 477 altitude into the cloud, the energy and water content of the entrained air source decreases, 478 and therefore the clouds need to entrain less air (compared to the normal sounding case) in 479 order to generate the same MSE and q_t decrease from adiabatic cloud values. For Sept. 8 480 (Fig. 9), mean m_o decreases from 52% to 37% with the upward-shift in entrainment level. The 48 standard deviation of m_o remains similar, with values of 14% and 12% respectively. These 482 tests suggest that our analysis is robust with respect to our assumption of lateral entrainment. 483 Detrainment mass fractions change rather little, while entrainment mass fractions change 484 moderately in the expected manner. 485

Another assumption made in this method is that the mixed-layer air comprising the cloud is an unbiased sample. Updrafts, however, are biased towards warmer temperatures so T_a is likely biased slightly low. If we were to use only the top 10% of temperature values in calculating MSE, the temperature bias estimated from the data is about 0.5 K, which leads to a s_a bias of 0.5 kJ/kg. This is much smaller than the change in MSE caused by shifting the entrainment levels by 400 m, which is around 3 to 4 kJ/kg depending on the day. Thus, we do not expect the mixed-layer air bias in MSE to substantively change the results.

493 3.2.3 Detrained air properties

The issue of detrainment is made more complex because we only sample each cloud at one 494 level, and therefore we have no information about any single cloud's properties at different 495 altitudes or time (as opposed to entrainment where we have a clear-air sounding that pro-496 vides information at all altitudes). We have previously assumed that the detrained air has 497 properties that are the average of the sampled cloud and the adiabatic air (Section 2.4); see 498 Eqs. 6 and 7. This is rationalized because detrainment from the cloud could have occurred 499 at any time in the past, at which time the cloud would have been closer to adiabatic than 500 at the moment of the aircraft cloud penetration. Here, we change the assumption to one 501 where detrainment occurred when the cloud properties are exactly that at the moment of 502 the penetration, i.e. $q_d = q_c$ and $s_d = s_c$. Figure 10 shows the detrained and entrained air 503 mass fractions when this is assumed. The mean values of m_d are still small, and in fact are 504 smaller than the results shown in Fig. 5. The other difference from the base case detrainment 505 scenario is that the large detrainment events no longer exist; the maximum value of m_d is 506 3%. Physically, this seems to be less plausible than the results from our base case, but does 507 illustrate that the detrainment values deduced by this method exhibit some sensitivity to 508 the assumption of the properties of the detrained air. The corresponding entrainment mass 509 fractions m_o under this assumption are 25 to 60% as compared to 30 to 70% in the base case, 510 a small shift that does not change the qualitative picture of the mass fluxes in these clouds. 511 These sensitivity tests show that our results do depend on the assumed detrained air prop-512 erties, mainly in the fraction of large m_d events, although we consider our base case analysis 513 to be more realistic regarding detrainment than the model used in this sensitivity analysis. 514 The overall picture is consistent between these two analyses: detrainment is generally a weak 515 process in these summertime shallow cumulus clouds. 516

517 4 Relationship with buoyancy profiles

Previous studies have suggested that detrainment is related to cloud buoyancy profiles. For example, a modeling study by *Carpenter et al.* [1998b] found that cold descending air will sink until it reaches its level of neutral buoyancy, at which point it will diverge and detrain. *Bretherton and Smolarkiewicz* [1989] suggest that changes in the gradient of the buoyancy of the cloudy air causes entrainment or detrainment. While our observations can not inform the latter, the former hypothesis can be tested in our observations.

To test these ideas, we compare the environmental density profile along with the measured 524 penetration cloudy air density, both expressed as virtual potential temperature θ_v . Figures 11 525 and 12 illustrate results for two of the six days. The detrainment mass fraction m_d for 526 each penetration is indicated by both color and size of the data marker. In general, the 527 results show that the cloudy air either exhibits θ_v values that are equal to or larger than the 528 environment. This is consistent with the formation of cumulus clouds by air that is positively 529 buoyant relative to the environment. While one expects a shell of cold, negatively buoyant, 530 descending air to be present around the periphery of the cloud, this is offset in the mean by the 531 warm, positively buoyant air inside this shell, at least for actively growing clouds. For those 532 cloud slices that are substantially positively buoyant relative to the environmental sounding, 533 the maximum difference in θ_v is less than 2 K, with most within 1 K. There are a handful 534 of penetrations where the cloudy air is negatively buoyant relative to the environment; the 535 difference in θ_v in these cases appears to be smaller than for the positively buoyant cases, 536 though the small sample size makes it difficult to reach any statistically significant conclusion. 537 The small fraction of negatively buoyant penetrations also suggests that sampling is biased 538 against dissipating clouds as speculated above. 539

If we focus on only those cases with largest m_d values $(m_d > 10\%)$, we find that almost all of these cloud penetrations (20 out of 22 cases) exhibit mean θ_v values that are (within uncertainty) the same as the environmental θ_v , i.e. the cloudy air is, on average, at its level

of neutral buoyancy. This finding is consistent with the hypothesis of detrainment occuring 543 at the level of neutral buoyancy [Carpenter et al., 1998b]. There are two counter-examples 544 over all six days; one of these is illustrated in Fig. 11 (near 2100 m altitude and $\theta_v = 308$ K) 545 where the cloudy air is warmer by ~ 0.5 K. In contrast, the fraction of events at low m_d which 546 exhibit θ_v values that are substantially warmer than the sounding is much greater, perhaps 547 indicating younger, growing clouds which have detrained very little air over their history. At 548 these low m_d values, though, the most likely case is still one where the cloudy air θ_v very 549 closely matches the environment. 550

Lastly, we also see no obvious trend of large m_d events correlated to any change in shape of the environmental sounding. If we had, it may have been an indication that the mechanism proposed by *Bretherton and Smolarkiewicz* [1989] is relevant to these observations; the lack of such a correlation, though, neither proves nor disproves this mechanism as we have no vertical profiles of in-cloud buoyancy to properly test it.

556 5 Summary and Conclusions

We have proposed a novel method to estimate the amounts of gross detrainment and en-557 trainment using aircraft observations. The method optimizes conservation equations for 558 cloud mass, moist static energy and total moisture to solve for the mass fractions of adi-559 abatic, entrained and detrained air (termed m_a , m_o and m_d respectively) for each aircraft 560 cloud penetration. In warm, shallow, non-precipitating cumuli, we find that these clouds are 561 comprised of approximately equal parts of surface-layer air that has been lifted adiabatically 562 and entrained air, the latter comprising between 30 and 70% of the cloud mass, with a me-563 dian of 45%. Detrainment mass fractions are found to be typically quite low, with 78% of 564 our cases exhibiting $m_d < 2\%$. In about 15% of our aircraft cloud penetrations, however, 565 we estimate $m_d > 10\%$. These low values may be inconsistent with budget studies in tower-566 ing/congestus cumuli, which infer detrainment mass fluxes comparable to the upward mass 567

flux of surface-layer air Raymond and Wilkening, 1982, 1985; Raga et al., 1990; Barnes et al., 568 1996; Carpenter et al., 1998a]. These results are more consistent with those from Wang and 569 Geerts [2011], who find no evidence of active detrainment; their study, along with this one, 570 suggest that vertical transport is dominated by the air that remains after dissipation of the 571 cloud, with little active detrainment to the environment during the cloud's active phase. The 572 incompatibility of these results with other previous studies could be explained if detrainment 573 fluxes in cumulus clouds are controlled by parameters that differ among these studies. Such 574 controlling parameters might include cloud type (e.g. cumulus mediocris vs. congestus) and 575 surrounding dynamic and thermodynamic environmental properties (e.g. subsidence rate; 576 T and humidity profiles). Differences in study methodology may also play a role, of which 577 we highlight a few: uncertainties in mass budgets; possible biases in our aircraft sampling 578 towards younger, more vigorous clouds; detrainment occuring at specific layers that are not 579 easily sampled by aircraft; and strong variability of detrainment with cloud height or cloud 580 age. 581

Vertical profiles of detrainment show a trend of increasing m_d with height in the cloud, consistent with *Raga et al.* [1990]. Vertical profiles of entrainment also show an increase in the upper-half of the cloud as compared to the lower-half, which fits with the common observation that adiabaticity in cumulus tends to decrease with height (e.g. *Lu et al.*, 2008). Our confidence in our new method is increased because the inferred vertical trends are physically sensible.

We also find that more than 90% of the larger detrainment events $(m_d > 10\%)$ are associated with cloudy air that has θ_v equal to that of the environmental sounding. This is consistent with *Carpenter et al.* [1998b] that found that descending air will detrain when it reaches its level of neutral buoyancy. In contrast, clouds with low m_d were much more frequently associated with air that was positively buoyant relative to the environment.

A number of assumptions were made as part of this analysis. Most notably, we assume that entrainment occurs laterally at the level of observation, and that detrained air has properties that are the average of adiabatic air and the air sampled by the aircraft. Sensitivity tests show that the former does not dramatically change the qualitative results of this study. Changing the latter assumption to one where detrained air has exactly the same properties as the cloudy air at the same sampling level causes all the detrainment events to shift to small (< 2%) values. Testing this methodology in high resolution models is an important future step to gain further confidence in these results.

Compared to entrainment, detrainment is far less-studied despite its importance to understanding clouds, its role in atmospheric transport and, consequently, weather and climate. The dearth of previous studies of gross detrainment hampers our ability to evaluate these results within a broader context, especially when we expect detrainment to depend on cloud type and environmental conditions. Developing a deeper understanding of detrainment from clouds, and its controlling parameters, will likely require combining a variety of approaches, of which this study is one example, in a variety of settings.

$_{608}$ Acknowledgements

The authors thank NOAA for funding participation in the GoMACCS aerosol-cloud field experiment. We also thank the CIRPAS Twin Otter team for all their dedicated work to make this field program a great success. We are indebted to John Seinfeld and Rick Flagan for their scientific leadership during this experiment, and the rest of the Twin Otter science team for various discussions along the way.

Date	Number of	Avg Penetration	Takeoff	Cloud	Cloud
	Clouds	${ m Length}$	time [UTC]	base [m]	top [m]
Sept. 1	15	890 m	16:52	1330	2400
Sept. 2	42	730 m	16:02	1460	2600
Sept. 8	27	660 m	16:54	1322	2400
Sept. 11	44	$590 \mathrm{~m}$	14:29	655	3100
Sept. 14	27	$630 \mathrm{~m}$	16:55	969	2600
Sept. 15	21	630 m	15:59	1068	2800

Table 1: Summary of clouds sampled on each flight day. Local time is UTC minus 5 hrs (Central daylight time).



Figure 1: Aircraft altitude and cloud liquid water content as a function of time for the Sept. 8 flight. There were 27 clouds sampled on this day. The clear air sounding occurs from approximately minute 20 to 50.



Figure 2: A sketch showing the sources of air that are assumed in this analysis to comprise a cloud. M_a rises adiabatically from cloud base, M_o is entrained laterally at the altitude the cloud is sampled, and M_d is detrained air.



Figure 3: Mass fractions of detrained and entrained air as a function of altitude, along with clear air soundings of MSE and q_t , for Sept. 2, 2006. Larger circles indicate smaller optimization residuals, i.e. less uncertainty in estimated m_d and m_o .



Figure 4: Mass fractions of detrained and entrained air as a function of altitude, along with clear air soundings of MSE and q_t , for Sept. 8, 2006. Larger circles indicate smaller optimization residuals, i.e. less uncertainty in estimated m_d and m_o .



Figure 5: Histogram of detrained air mass fractions for all flight days.



Figure 6: Vertical detrainment mass fraction profile for all flight days. Altitude for each flight day is normalized to an altitude set ranging from cloud base to cloud top.



Figure 7: Histogram of entrained air mass fractions for all flight days.



Figure 8: Vertical entrainment mass fraction profile for all flight days. Altitude for each flight day is normalized to an altitude set ranging from cloud base to cloud top.



Figure 9: Mass fractions of detrained and entrained air as a function of altitude using shifted clear air soundings of MSE and q_t , for Sept. 8, 2006. Large diamonds indicate smaller optimization residuals, i.e. less uncertainty in estimated m_d and m_o . The soundings used in this case were shifted upwards by 400 m.



Figure 10: Histograms of detrained (left) and entrained (right) air mass fractions under the assumption that the detrained air has exactly the same properties as the air sampled during the aircraft penetration.



Figure 11: Virtual potential temperature θ_v of the environmental air (grey dots) from an aircraft sounding and mean θ_v (colored diamonds) for the air during each cloud penetration on Sept. 2, 2006. The detrainment mass fraction m_d for each penetration is indicated by both color and size of the diamond symbol.



Figure 12: Virtual potential temperature θ_v of the environmental air (grey dots) from an aircraft sounding and mean θ_v (colored diamonds) for the air during each cloud penetration on Sept. 8, 2006. The detrainment mass fraction m_d for each penetration is indicated by both color and size of the diamond symbol.

621 References

- Barnes, G. M., J. C. Fankhauser, and W. D. Browning, Evolution of the vertical mass flux
 and diagnosed net lateral mixing in isolated convective clouds, *Mon. Wea. Rev.*, 124(12),
 2764-2784, 1996.
- Blyth, A. M., Entrainment in cumulus clouds, J. Appl. Meteor., 32(4), 626–641, 1993.
- Blyth, A. M., S. G. Lasher-Trapp, and W. A. Cooper, A study of thermals in cumulus clouds, *Quart. J. Roy. Meteor. Soc.*, 131 (607), 1171–1190, 2005.
- Bretherton, C. S., and P. K. Smolarkiewicz, Gravity-waves, compensating subsidence and
 detrainment around cumulus clouds, J. Atmos. Sci., 46(6), 740–759, 1989.
- Carpenter, R., K. Droegemeier, and A. Blyth, Entrainment and detrainment in numerically
 simulated cumulus congestus clouds. Part II: Cloud budgets, J. Atmos. Sci., 55(23), 3433–
 3439, 1998a.
- Carpenter, R. L., K. K. Droegemeier, and A. M. Blyth, Entrainment and detrainment in
 numerically simulated cumulus congestus clouds. Part III: Parcel analysis, J. Atmos. Sci.,
 55(23), 3440–3455, 1998b.
- Chand, D., R. Wood, T. L. Anderson, S. K. Satheesh, and R. J. Charlson, Satellite-derived
 direct radiative effect of aerosols dependent on cloud cover, *Nature Geoscience*, 2(3), 181–
 184, doi:10.1038/NGEO437, 2009.
- Damiani, R., and G. Vali, Evidence for tilted toroidal circulations in cumulus, J. Atmos. Sci.,
 64 (6), 2045–2060, doi:10.1175/JAS3941.1, 2007.
- Dawe, J. T., and P. H. Austin, The influence of the cloud shell on tracer budget measurements
 of LES cloud entrainment, J. Atmos. Sci., 68(12), 2909–2920, doi:10.1175/2011JAS3658.1,
 2011.

- de Rooy, W., P. Bechtold, K. Froehlich, C. Hohenegger, D. Jonker, H. J. J.and Mironov,
 A. Siebesma, J. Teixeira, and J. Yano, Entrainment and detrainment in cumulus convection: an overview, Q. J. R. Meteorol. Soc., 139(670, A), 1–19, doi:10.1002/qj.1959, 2013.
- de Rooy, W. C., and A. P. Siebesma, Analytical expressions for entrainment and detrainment in cumulus convection, *Quart. J. Roy. Meteor. Soc.*, 136(650), 1216–1227, doi:
 10.1002/qj.640, 2010.
- Gerber, H., B. G. Arends, and A. S. Ackerman, New microphysics sensor for aircraft use, *Atmos. Res.*, 31(4), 235–52, 1994.
- Heus, T., G. Van Dijk, H. J. J. Jonker, and H. Van den Akker, Mixing in shallow cumulus
 clouds studied by lagrangian particle tracking, J. Atmos. Sci., 65(8), 2581–2597, doi:
 10.1175/2008JAS2572.1, 2008.
- Hoffmann, F., H. Siebert, J. Schumacher, T. Riechelmann, J. Katzwinkel, B. Kumar, P. Goetzfried, and S. Raasch, Entrainment and mixing at the interface of shallow cumulus clouds:
 Results from a combination of observations and simulations, *Meteorol. Z.*, 23(4), 349–368,
 doi:10.1127/0941-2948/2014/0597, 2014.
- Houghton, H. G., and H. E. Cramer, A theory of entrainment in convective currents, J. *Meteor.*, 8(2), 95–102, 1951.
- Kitchen, M., and S. J. Caughey, Tethered-balloon observations of the structure of small
 cumulus clouds, *Quart. J. Roy. Meteor. Soc.*, 107(454), 853–874, 1981.
- Lawson, R. P., and W. A. Cooper, Performance of some airborne thermometers in clouds, J.
 Atmos. Oceanic Technol., 7(3), 480–494, 1990.
- Liao, H., and J. H. Seinfeld, Effect of clouds on direct aerosol radiative forcing of climate, J.
 Geophys. Res., 103(D4), 3781–3788, doi:10.1029/97JD03455, 1998.

- Lu, M. L., G. Feingold, H. H. Jonsson, P. Y. Chuang, H. Gates, R. C. Flagan, and J. H. Seinfeld, Aerosol-cloud relationships in continental shallow cumulus, *J. Geophys. Res. Atmos.*, 113(D15), doi:10.1029/2007JD009354, 2008.
- Malkus, J. S., Some results of a trade-cumulus cloud investigation, J. Meteor., 11(3), 220–
 237, 1954.
- Perry, K. D., and P. V. Hobbs, Influences of isolated cumulus clouds on the humidity of their
 surroundings, J. Atmos. Sci., 53(1), 159–174, 1996.
- Raga, G. B., J. B. Jensen, and M. B. Baker, Characteristics of cumulus band clouds off the
 coast of Hawaii, J. Atmos. Sci., 47(3), 338–355, 1990.
- Raymond, D. J., and M. Wilkening, Characteristics of mountain-induced thunderstorms and
 cumulus congestus clouds from budget measurements, J. Atmos. Sci., 42(8), 773–783,
 1985.
- Raymond, D. J., and M. H. Wilkening, Flow and mixing in New Mexico mountain cumuli,
 J. Atmos. Sci., 39(10), 2211–2228, 1982.
- Raymond, D. J., R. Solomon, and A. M. Blyth, Mass fluxes in new-mexico mountain thunderstorms from radar and aircraft measurements, *Quart. J. Roy. Meteor. Soc.*, 117(499),
 587-621, doi:10.1002/qj.49711749909, 1991.
- Rougier, J., D. M. H. Sexton, J. M. Murphy, and D. Stainforth, Analyzing the climate sensitivity of the HadSM3 climate model using ensembles from different but related experiments, *J. Climate*, 22(13), 3540–3557, doi:10.1175/2008JCLI2533.1, 2009.
- Samset, B. H., and G. Myhre, Vertical dependence of black carbon, sulphate and
 biomass burning aerosol radiative forcing, *Geophys. Res. Lett.*, 38, L24,802, doi:
 10.1029/2011GL049697, 2011.

- Small, J. D., P. Y. Chuang, G. Feingold, and H. L. Jiang, Can aerosol decrease cloud lifetime?,
 Geophys. Res. Lett., 36, doi:10.1029/2009GL038888, 2009.
- Stith, J. L., Observations of cloud-top entrainment in cumuli, J. Atmos. Sci., 49 (15), 1334–
 1347, 1992.
- Taylor, G. R., and M. B. Baker, Entrainment and detrainment in cumulus clouds, J. Atmos.
 Sci., 48(1), 112–121, 1991.
- Wang, Y., and B. Geerts, Observations of detrainment signatures from non-precipitating orographic cumulus clouds, Atmos. Res., 99(2), 302–324, doi:10.1016/j.atmosres.2010.10.023,
 2011.
- Wang, Y., B. Geerts, and J. French, Dynamics of the cumulus cloud margin: An observational
 study, J. Atmos. Sci., 66(12), doi:10.1175/2009JAS3129.1, 2009.
- Yeo, K., and D. Romps, Measurement of Convective Entrainment Using Lagrangian Particles,
 J. Atmos. Sci., 70(1), 266-277, doi:10.1175/JAS-D-12-0144.1, 2013.
- ⁷⁰³ Zhao, M., and P. H. Austin, Life cycle of numerically simulated shallow cumulus clouds. Part
- ⁷⁰⁴ II. Mixing dynamics, J. Atmos. Sci., 62(5), doi:10.1175/JAS3415.1, 2005.