

Iodine oxide in the global marine boundary layer

C. Prados-Roman¹, C. A. Cuevas¹, T. Hay¹, R. P. Fernandez^{1,*}, A. S. Mahajan^{1,**},
S-J. Royer², M. Galí^{2,†}, R. Simó², J. Dachs³, K. Großmann⁴, D. E. Kinnison⁵, J-F.
Lamarque⁵ and A. Saiz-Lopez¹

[1]{Atmospheric Chemistry and Climate Group, Institute of Physical Chemistry Rocasolano (CSIC), Madrid, Spain}.

[2]{Institute of Marine Sciences, (CSIC), Barcelona, Spain}.

[3]{Institute of Environmental Assessment and Water Research (CSIC), Barcelona, Spain}.

[4]{Institute of Environmental Physics, University of Heidelberg, Germany}.

[5]{Atmospheric Division, NCAR, Boulder, CO, USA}.

[*]{now at: National Scientific and Technical Research Council (CONICET), UTN-FR Mendoza/ICB-UNCuyo, Mendoza, Argentina}.

[**]{now at: Indian Institute of Tropical Meteorology, Pune, India}.

[†]{now at: Takuvik (UL/CNRS), Quebec, Canada}.

Correspondence to: a.saiz@csic.es.

Abstract

Emitted mainly by the oceans, iodine is a halogen compound important for atmospheric chemistry due to its high ozone depletion potential and effect on the oxidizing capacity of the atmosphere. Here we present a comprehensive dataset of iodine oxide (IO) measurements in the open marine boundary layer (MBL) made during the Malaspina 2010 circumnavigation. Results show IO mixing ratios ranging from 0.4 to 1 pmol mol⁻¹ (30% uncertainty) and, complemented with additional field campaigns, this dataset confirms through observations the ubiquitous presence of reactive iodine chemistry in the global marine environment. We use a global model with organic (CH₃I, CH₂ICl, CH₂I₂ and CH₂IBr) and inorganic (HOI and I₂)

1 iodine ocean emissions to investigate the contribution of the different iodine source gases to
2 the budget of IO in the global MBL. In agreement with previous estimates, our results
3 indicate that, globally averaged, the abiotic precursors contribute about 75% to the iodine
4 oxide budget. However, this work reveals a strong geographical pattern in the contribution of
5 organic vs. inorganic precursors to reactive iodine in the global MBL.

6

7 **1 Introduction**

8 The atmospheric relevance of reactive halogens became clear decades ago when their
9 potential to catalytically destroy ozone (O_3) was first recognised in the polar stratosphere
10 (Molina and Rowland, 1974) and later on in the troposphere (e.g. Barrie et al., 1988).
11 Halogens are also known to affect the NO_x (NO , NO_2) and HO_x (HO , HO_2) partitioning and
12 the lifetime of organic compounds, to alter the sulphur and mercury cycles and, in the case of
13 iodine oxides, to form ultra-fine particles in coastal areas (Saiz-Lopez and von Glasow, 2012
14 and references therein).

15 Since the first study to deal with the tropospheric relevance of inorganic iodine (Chameides
16 and Davis, 1980), major efforts have been made to detect reactive iodine species in their main
17 source region: the oceans (Saiz-Lopez et al., 2012 and references therein). Several field
18 campaigns in scattered marine environments have aimed at detecting iodine oxide - beacon
19 for the presence of active iodine chemistry- and determining the nature and strength of
20 organic and inorganic source gases of iodine (referenced hereafter as OSG, ISG, respectively).
21 Air-sea fluxes of iodocarbons (CH_3I , CH_2I_2 , CH_2ICl and CH_2IBr , C_2H_5I , 1- C_3H_7I , 2- C_2H_7I)
22 have been reported (Carpenter et al., 2012), but in general observations were insufficient to
23 explain measured IO concentrations in the MBL, implying the existence of an abiotic ocean
24 source of iodine (Mahajan et al., 2010; Jones et al., 2010; Mahajan et al., 2012; Gómez Martín
25 et al., 2013a; Großmann et al., 2013; Lawler et al., 2014). In several one-dimensional model
26 studies simulated emissions of molecular iodine (I_2) were used to fit IO observations (e.g.
27 Mahajan et al., 2010; Großmann et al., 2013), however the recent work of Lawler et al. (2014)
28 with the first observation of I_2 in the remote MBL, confirmed that the emission of I_2 is still
29 insufficient to explain the observed levels of IO. Recently, the study of Carpenter et al. (2013)
30 has experimentally confirmed that not only I_2 is emitted naturally from the oceans but also,
31 and mainly, hypoiodous acid (HOI). In that study and in the subsequent work of MacDonald
32 et al. (2014), the authors have confirmed through laboratory work that the oceanic emission of

1 ISG (HOI and I₂) follows the deposition of tropospheric O₃ to the oceans and its reaction with
2 aqueous iodide (I_{aq}⁻, Garland et al., 1980), and they proposed a parameterisation for ocean
3 ISG emissions dependent on O₃, wind speed (ws) and sea surface temperature (SST).

4 In this work, we present a comprehensive map of IO observations in the global MBL showing
5 the ubiquity of this radical in the marine environment. Moreover, by means of a global model
6 including OSG and ISG oceanic emissions; we investigate the geographical emission patterns
7 of both iodine precursors and their contribution to the IO budget in the marine environment.
8 Section 2 details the measurement campaign of Malaspina 2010 and provides information on
9 the chemical model used throughout this work. Section 3 presents the results of the IO
10 observations and the modelling studies, and Section 4 concludes this work.

11

12 **2 Measurements and model**

13 In the following we present the setup of the O₃ and IO measurements during the Malaspina
14 2010 expedition as well as the model schemes used in this study.

15 **2.1 Measurements during the Malaspina 2010 circumnavigation**

16 From December 2010 until July 2011 the Spanish research vessel Hesperides
17 circumnavigated the World's oceans within the framework of the Malaspina 2010 project.
18 The main objectives of this interdisciplinary campaign were to investigate the
19 biogeochemistry, physical properties and microbiological biodiversity of the oceans; the
20 genetic diversity of the deep-ocean and the exchange of trace gases and pollutants with the
21 atmosphere; and assessing the impact of global change in the ocean. The different legs of the
22 cruise and the docking dates are indicated in Table 1.

23 A Multi-Axis Differential Optical Absorption Spectroscopy instrument (MAX-DOAS; Platt
24 and Stutz, 2008) and a commercial 2B-205 ozone monitor, along with a GPS, were deployed
25 aboard the vessel in order to investigate the presence of atmospheric trace gases such as IO,
26 O₃, BrO, HCHO and CHOCHO in the MBL. Herein we focus on the observations of IO and
27 O₃ during the campaign.

1 **2.1.1 Surface ozone**

2 The ozone monitor was installed in the ship's bridge with a 5 m long Teflon-lined inlet tube
3 from the upper deck, well forward of the exhaust stacks (~15 m above sea level (m.a.s.l.)).
4 The inlet was placed just above the railing in the air coming from the front of the ship,
5 avoiding sampling air from the ship's boundary layer. Due to GPS communication errors, our
6 data compilation started on 21/02/2011 (2nd leg) and finished on 12/07/2011.

7 The ozone volume mixing ratios (vmr) observed during Malaspina 2010 are presented in Fig.
8 1a along with the ancillary measurements of relevance for the present work (i.e., SST and ws;
9 Fig. 1 and Table 2). Simulations of the 5-day backward trajectories of the air masses arriving
10 at the ship's track are provided in Fig. 2, showing the typical non-continental origin of the air
11 masses sensed during the cruise.

12 **2.1.2 Iodine oxide**

13 Aiming at the detection of IO along the Malaspina's track, a MAX-DOAS instrument was
14 installed on the second deck near the rear of the ship (~10 m.a.s.l.). Briefly, these instruments
15 measure the intensity of scattered light in the UV-VIS range entering a scanning telescope at
16 several precise viewing angles and have been widely used for atmospheric composition
17 research (Platt and Stutz, 2008). Thus only a summary of the particular MAX-DOAS
18 instrument mounted on the Hesperides research vessel is given hereafter. For details regarding
19 the MAX-DOAS technique please refer to the work of, e.g., Platt and Stutz (2008) and
20 Hönninger et al. (2004) and for further details of our ship-based MAX-DOAS instrument
21 please see Mahajan et al. (2012).

22 Briefly, in the case of the Malaspina's MAX-DOAS instrument, the scanning telescope was
23 housed in a weatherproof metal chamber with a flat UV-transmitting acrylic window, with a
24 sunshade to reduce spectral effects on the window. The telescope unit (built by the New
25 Zealander National Institute of Water and Atmospheric Research-NIWA) was mounted on a
26 gimbal table to compensate for the pitch and roll of the ship. The gimbal dampened the
27 effective oscillations in telescope elevation angle to $\pm 1^\circ$ for most of the cruise and $\pm 2^\circ$ in
28 rough conditions. In addition, a high accuracy ($\pm 0.1^\circ$), fast response (0.3 s) inclinometer was
29 used to log the residual oscillations in order to correct the elevation angles. Only true angles
30 within 0.2° of each prescribed elevation angle were used for analysis. The azimuth viewing
31 direction was towards the ship's bow (20° anticlockwise) to minimize exhaust emissions in the

1 line of sight. The scanning telescope consisted of a rotating diagonal mirror driven by a
2 stepper motor and a 50.8 mm diameter fused silica lens with a focal length of 200 mm, giving
3 a field of view of 0.5°. The light was focused onto a 5 m long 19 optic fibre bundle leading to
4 a Princeton Instruments SP500i spectrometer with a Princeton Instruments Pixis 400B CCD
5 camera. A 600 grooves mm⁻¹ grating was used, giving approximately an 80 nm spectral
6 window and a spectral resolution of 0.5 nm FWHM. Spectra were recorded for a short
7 exposure time of 1 s at each discrete elevation angle (2, 4, 6, 8, 10, 15, 30 and 90°) in order to
8 minimize potential deviations in angle due to the ship's movement. The scan sequence was
9 repeated every 2 min and after every 10 cycles the grating was shifted between the two
10 wavelength regions, centred on 358 nm (UV spectral range) and 440 nm (VIS spectral range).
11 Results presented in this work correspond to the VIS channel, where IO could be measured
12 (see Sect. 3.1).

13 **2.2 Modelling the oceanic emissions of reactive iodine precursors**

14 We implemented the experimentally derived ocean fluxes of ISG (Carpenter et al., 2013;
15 MacDonald et al., 2014) into the global chemistry-climate model CAM-Chem (Community
16 Atmospheric Model with Chemistry, version 4.0; Lamarque et al., 2012), which already
17 included a validated OSG emissions inventory and a state-of-the-art halogen chemistry
18 scheme (Ordóñez et al., 2012). The on-line ISG flux formulation, based on the studies of
19 Carpenter et al. (2013) and MacDonald et al. (2014), was performed considering the
20 instantaneous modelled levels of surface O₃, SST and ws in each of the model grid-boxes over
21 the oceans (i.e., imposing an ocean mask). In the following we summarise the model schemes
22 used in this work. Further details on the particular implementation of the ISG
23 parameterisation into the CAM-Chem model are given in Prados-Roman et al. (2014),
24 whereas the general model setup is described in the study of Lamarque et al. (2012).

25 **2.2.1 Model schemes**

26 Throughout this work, two different pairs of simulations were performed in order to evaluate
27 the model, to identify the contribution of OSG/ISG fluxes and to estimate the iodine burden of
28 the MBL. A brief description of the simulations used in this study is given below.

29 (1) *Base-Organic runs*. In the *Base run*, simulations were performed considering the
30 oceanic emission of organic and inorganic iodine precursors. Based on previous
31 publications, the OSG inventory of very-short lived iodocarbons (OSG= CH₃I,

1 CH₂I₂, CH₂I₂Br and CH₂I₂Cl) was considered (Ordóñez et al., 2012), while the ISG
2 computation of HOI and I₂ was used as described in the study of Prados-Roman et
3 al., 2014. In order to distinguish the contribution of the inorganic and the organic
4 iodine source gases to the IO budget in the MBL, the *Organic* scheme included only
5 the abovementioned OSG (by forcing the inorganic emissions to be null). Hence, the
6 contribution of ISG to the IO budget in the MBL (i.e., IO_{ISG}) was defined as the
7 difference between the IO vmr obtained in the MBL after the *Base run*
8 (IO=IO_{ISG}+IO_{OSG}), and the IO vmr obtained after the *Organic run* (i.e., IO_{OSG}). That is
9 IO_{ISG}= (IO) - (IO_{OSG}); and the relative contribution of ISG to IO was defined as
10 (IO_{ISG})/(IO) in percentage. Similarly, the contribution of each individual iodocarbon
11 to the budget of IO was investigated.

12 (2) *NoPhot-Phot runs*. It is known that the self-reaction of IO in pristine conditions
13 yields the formation of higher oxides (I₂O_x, x=2, 3 or 4). However, once formed, the
14 reaction pathways of these compounds are still not well understood. One possibility
15 is their nucleation into ultra-fine particles as observed in coastal areas (Gómez
16 Martín et al., 2013b). Those conditions were however not representative of the
17 Malaspina expedition since most of the marine atmosphere crossed was
18 representative of open ocean environment. A possible pathway for Malaspina's
19 conditions was the photodissociation of those I₂O_x into OIO+I, OIO+IO or OIO+OIO
20 as previously modelled for the Antarctic (Saiz-Lopez et al., 2008) and global marine
21 troposphere (Saiz-Lopez et al., 2014), which would therefore result in additional
22 reactive iodine in the MBL. The so-called *Phot run* included I₂O_x photolysis while
23 the simulation excluding such photolysis was referred to as *NoPhot run*. Note that,
24 unless stated otherwise, in the aforementioned *Base-Organic* schemes the I₂O_x were
25 not allowed to photolyse but, once formed, they were lost by thermal decomposition
26 or to pre-existing aerosols instead.

27 All simulations were performed with a horizontal grid resolution of 1.9° (latitude) × 2.5°
28 (longitude) and 26 hybrid vertical levels (0-40 km), and considered the SST and sea-ice
29 boundary conditions representative of year 2000 (Rayner et al., 2003). Note that, since the
30 model was not run with specified dynamics, simulations are not representative of the
31 meteorology of any specific year. Thus, unless stated the opposite, the model results presented
32 in this work correspond to 24 h annual averages.

33

1 **3 Results and discussions**

2 In this section we present the observations of IO in the MBL and compare them to different
3 model runs. Furthermore, we investigate the contribution of the OSG and ISG fluxes to the IO
4 budget in the MBL.

5 **3.1 Observations of IO in the global marine boundary layer**

6 During Malaspina 2010 IO was detected above instrumental detection limit ($1.2\text{-}3.5 \times 10^{13}$
7 molec cm⁻²) in all marine environments sampled. Figure 3 shows a typical IO spectral fit
8 during that expedition and the IO differential slant column densities (dSCD) measured along
9 the cruise track. Note that diverse filters were used in this dataset for quality assurance (e.g.,
10 cloud and wind direction filters). Following previous studies and using only IO dSCD above
11 the quality filters, the IO mixing ratios were inferred by the well-established "O₄ method"
12 (Wagner et al., 2004), after validating results of several days with a radiative transfer model
13 (RTM) (e.g., Mahajan et al., 2012; Gómez Martín et al., 2013a). Particular details on these
14 procedures (IO spectral and vmr retrieval) as well as the quality filters applied are provided in
15 the Supplementary Information (SI).

16 Overall, during the Malaspina expedition the IO radical was constantly observed in the
17 daytime MBL over three oceans and both hemispheres. The IO vmr integrated in the MBL
18 ranged between 0.4 and 1 pmol mol⁻¹ (detection limit of ~ 0.2 pmol mol⁻¹), with lower values
19 measured over the South Atlantic waters and with the highest levels in the marine region west
20 of Mexico. Figure 4 shows the averaged daytime IO vmr of the Malaspina dataset, along with
21 IO vmr obtained from former field campaigns: Cape Verde (Read et al., 2008; Mahajan et al.,
22 2010), HaloCAST-P (Mahajan et al., 2012), CHARLEX (Gómez Martín et al., 2013a) and
23 TransBrom (Großmann et al., 2013). Note that the IO vmr reported for each of these
24 campaigns are by definition intrinsically linked to the specific viewing geometry of each
25 DOAS instrument (Platt and Stutz, 2008). During the Cape Verde campaign a Long Path-
26 DOAS instrument was used with a fixed light path at 10 m.a.s.l. (Read et al., 2008; Mahajan
27 et al., 2010). In all the other campaigns shown in Fig. 4, MAX-DOAS instruments were
28 employed. Given the different viewing elevation angles and instrumental setup, each of those
29 MAX-DOAS instruments sensed a different part of the MBL (Platt and Stutz, 2008;
30 Hönninger et al., 2004; Wagner et al., 2004). Although sensitivity RTM studies performed
31 during each of those MAX-DOAS campaigns agreed on a decreasing vertical profile of IO in

1 the MBL, the generally poor information content of the measurements hindered the vertical
2 resolution of the inferred IO vmr vertical profiles and the reported vmr were therefore linked
3 to a given sensed layer; particularly 0-200 m during HaloCAST-P and TransBrom (Mahajan
4 et al., 2012; Großmann et al., 2013), 0-1200 m during CHARLEX (Gómez Martín et al.,
5 2013a) and 0-600 m during Malaspina (this work, SI). Therefore the values reported in Fig. 4
6 should be considered as the mean IO vmr in each of the aforementioned altitude ranges linked
7 to a given elevation angle (e.g., 2° in the case of Malaspina). Note that, despite these
8 unavoidable retrieval limitations, Fig. 4 proves the ubiquity of IO in the global MBL and
9 hence the presence of reactive iodine chemistry in all sub-polar marine environments.

10 **3.2 Observations vs. model**

11 Figure 4 shows the most comprehensive map of IO observations in the remote marine
12 environment. We now use these observations together with the CAM-Chem model to evaluate
13 the geographical distribution of IO in the MBL. The performance of the model was evaluated
14 by comparing modelled and observed IO mixing ratios in the MBL for the aforementioned
15 particular altitude range sensed during each campaign. Note that, as mentioned above, a key
16 parameter in the model setup is the flux of ISG, which depends mainly on O₃ and ws
17 (Carpenter et al., 213; MacDonald et al., 2014). Hence, even though Fig. 4 shows IO
18 measurements from 5 different field campaigns, surface O₃, ws and IO were not measured
19 simultaneously during HaloCAST-P and TransBrom campaigns. Thus, only the campaigns of
20 Malaspina, CHARLEX and Cape Verde were chosen for comparison with the model. Figure 5
21 presents this comparison exercise, where the IO vmr observations in three oceans and both
22 hemispheres are juxtaposed to the model output after the *Organic* scheme and after the *Base*
23 run considering the *NoPhot* and the *Phot* schemes. For this exercise the model was sampled at
24 the same time (month) of the year and geolocation as the measurements (considering the
25 model grid resolution of 1.9° latitude x 2.5° longitude). Note that the low IO vmr resulting
26 after the *Organic* run remains basically unaltered despite the photolysis scheme considered.
27 Thus for simplicity only the *Organic-NoPhot* output (i.e., *Organic* run) is shown in Fig. 5.
28 Considering the ISG emissions, along with OSG, the model reproduces satisfactorily the IO
29 observations (Fig. 5). Note that, as found in the *Organic* run, the emission of OSG alone
30 explains on average only ~25% of the IO levels observed over the different oceans,
31 percentage that agrees well with previous one-dimensional model studies performed at
32 specific marine environments (Mahajan et al., 2010; Jones et al., 2010; Mahajan et al., 2012;

1 Gómez Martín et al., 2013a; Großmann et al., 2013; Lawler et al, 2014). This result points out
2 the importance of including ISG emissions in global models. Regarding the *Base* run results,
3 in general the *NoPhot* run reproduces the observations although in some regions the *Phot*
4 scheme is closer to the measurements (Fig. 5). Note that the modelled IO vmr in the *Phot*
5 scheme- likely to be a more realistic scheme for I_2O_x (Saiz-Lopez et al., 2014)- can even
6 double the IO vmr given by the *NoPhot* scheme, stressing the need of further efforts from the
7 community to investigate the fate of these higher iodine oxides. However, since the photolysis
8 rates of I_2O_x are currently subject to uncertainty (Saiz-Lopez et al., 2014), hereafter only the
9 *NoPhot* scheme is considered bearing, thus, the results presented as lower limits.

10

11 **3.3 Sources of IO in the global marine boundary layer**

12 After analysing the consistency of modelled vs. measured IO, in this section we investigate
13 the sensitivity of the IO levels towards the different modelled iodine precursors. Considering
14 the OSG emission inventory (Ordóñez et al., 2012) and the ISG (Prados-Roman et al., 2014),
15 the modelled OSG/ISG ratio allows quantifying the individual sources and total oceanic
16 emissions of iodine to the atmosphere. Results indicate that, globally averaged, the total
17 oceanic iodine emissions yield 2.3 Tg y^{-1} . From these, only 17% (0.4 Tg y^{-1}) originate from
18 organic sources, which are related to bacteria, microalgae, phytoplankton, etc. (Carpenter et
19 al., 2012). On a global average, nearly half (43%) of the organic flux derives from CH_3I , 29%
20 from CH_2ICl , 19% from CH_2I_2 and 9% from CH_2IBr , although their temporal and spatial
21 distribution varies with, e.g., the solar radiation at sea surface and the properties of the ocean
22 mixed layer (Bell et al., 2002; Carpenter et al., 2012; Ordóñez et al., 2012). The sea-air
23 exchange of iodine is thus driven mainly by abiotic sources. Our results indicate that, globally
24 averaged, $1.9 \text{ Tg (I) y}^{-1}$ (i.e., 83% of the total oceanic iodine fluxes) are emitted to the MBL
25 as a result of the reaction of tropospheric O_3 with I_{aq}^- in the ocean surface; and that the
26 partitioning of these ISG emissions is directed by HOI (95% HOI, 5% I_2 ; Prados-Roman et
27 al., 2014).

28 Figure 6a provides the annually averaged burden of IO in the global MBL, with values
29 ranging from less than $0.05 \text{ pmol mol}^{-1}$ in the sub-polar waters; to $\sim 0.9 \text{ pmol mol}^{-1}$ above
30 waters offshore the Baja California peninsula. Figure 6b shows the geographical pattern of the
31 contribution of ISG to the IO budget (i.e., IO_{ISG}). The model results indicate that, globally
32 averaged, about 75% of the IO in the MBL derives from inorganic precursors. As mentioned

1 in the previous section, as an averaged value, this result is indeed consistent with previous
2 estimates at given transects along the Pacific Ocean or offshore waters of Cape Verde and
3 Galapagos Islands (Mahajan et al., 2010; Jones et al., 2010; Mahajan et al., 2012; Gómez
4 Martín et al., 2013a; Großmann et al., 2013; Lawler et al., 2014). However, our model results
5 show the uneven geographical distribution of IO_{ISG} , e.g., marine tropical regions in the
6 southern hemisphere where IO_{ISG} is of 40%; or regions of ozone-related pollution outflow
7 such the Bay of Bengal or the Gulf of Mexico (Myhre et al., 2013, see also SI) where, as a
8 consequence of the $\text{O}_3\text{-I}_{\text{aq}}^-$ interaction, IO_{ISG} can be more than 90%. Figure 7 shows the
9 contribution of each of the four modelled iodocarbons to IO in the MBL, indicating that in the
10 biological active regions of the tropics IO derives mainly from the dihalomethanes ($\text{CH}_2\text{ICl} >$
11 $\text{CH}_2\text{I}_2 > \text{CH}_2\text{IBr}$) and to a lesser extend to CH_3I . Out of those regions CH_3I dominates the
12 organic contribution to IO in the MBL, increasing with latitude as a result of its longer
13 lifetime (Bell et al., 2002). Note however that the model simulations presented here do not
14 include iodine emissions, organic or inorganic, from ice surfaces. Also, the strong dependence
15 of the ISG flux with SST considerably reduces the inorganic iodine emissions over the cold
16 waters in the high latitudes. Furthermore, as detailed in the study of MacDonald et al. (2014),
17 the uncertainty on the parameterisation of ISG increases with decreasing SST. Thus, in the
18 polar marine regions our simulated inorganic contribution to the IO budget should be
19 regarded with caution. Despite these uncertainties, overall the main source of IO in the MBL
20 at a global scale is HOI. However, as shown in Fig. 6b and Fig. 7, this is subject to strong
21 spatial patterns in emission with regions in the southern hemisphere where the OSG can
22 account for up to 50% of the modelled IO levels.

23

24 **4 Summary**

25 Here we present a comprehensive set of observations of iodine oxide mixing ratios in the
26 marine boundary layer obtained after the Malaspina 2010 circumnavigation covering three
27 non-polar oceans and both hemispheres. Complementing this dataset with measurements
28 gained after campaigns in the tropical Atlantic Ocean and in the Eastern and Western Pacific
29 Ocean, we provide field evidence for the ubiquitous presence of IO, and thus reactive iodine
30 chemistry, in the global marine environment. By comparing these measurements with model
31 results, we confirm the need of including the inorganic oceanic emissions of iodine into
32 global models, and also stress the need for further laboratory and theoretical studies about the

1 atmospheric fate of I_2O_x . In particular, the model results indicate that 83% of the total oceanic
2 natural emissions of iodine are inorganic (mainly HOI) following the reaction of iodide with
3 ozone at the sea surface; and these inorganic emissions are indeed necessary to reproduce the
4 observations of IO in all marine environments. Finally, our results show that the contribution
5 of the organic/inorganic source gases to IO levels in the global MBL is geographically highly
6 variable, existing regions of ozone-rich outflow where the inorganic contribution to IO can be
7 more than 90%. This combined observational and modelling exercise strengthens the need of
8 including both the organic and the inorganic oceanic emissions of iodine into global models
9 for a more accurate assessment of the oxidizing capacity of the marine troposphere.

10

11 **Acknowledgements**

12 The authors would like to thank everyone involved in the Malaspina 2010 expedition, funded
13 by the Spanish Ministry of Economy and Competitiveness. In particular the Marine
14 Technology Unit (UTM-CSIC) for facilitating the ancillary data as well as the Hesperides
15 R/V crew. We also thank Alan Thomas, John Robinson and Dave Humphries (NIWA) for
16 building the MAX-DOAS tracker and the atmospheric group of the Institute of Environmental
17 Physics (Heidelberg) for facilitating the gimbal table. We also thank Markus Rex for his
18 comments on the ozone measurements during the TransBrom campaign, and Klaus
19 Pfeilsticker and John Plane for helpful discussions. The Indian Institute of Tropical
20 Meteorology is supported by the Ministry of Earth Sciences, Government of India. R.P.F
21 would like to thank ANPCyT (PICT-PRH 2009-0063) for financial support. The data
22 supporting this article can be requested from the corresponding author A.S-L (a.saiz@csic.es).

23

24

25 **References**

26 Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone
27 depletion and photochemical reactions at polar sunrise in the lower Arctic atmosphere,
28 Nature, 334, 138-141, 1988.

29 Bell, N., Hsu, L., Jacob, D. J., Schultz, M. G., Blake, D. R., Butler, J. H., King, D. B., Lobert,
30 J. M., and Maier-Reimer, E.: Methyl iodide: Atmospheric budget and use as a tracer of marine

1 convection in global models, *Journal of Geophysical Research: Atmospheres*, 107, 4340,
2 10.1029/2001JD001151, 2002.

3 Carpenter, L. J., Archer, S. D., and Beale, R.: Ocean-atmosphere trace gas exchange, *Chem.*
4 *Soc. Rev.*, 41, 6473-6506, 10.1039/C2CS35121H, 2012.

5 Carpenter, L. J., MacDonald, S. M., Shaw, M. D., Kumar, R., Saunders, R. W., Parthipan, R.,
6 Wilson, J., and Plane, J. M. C.: Atmospheric iodine levels influenced by sea surface emissions
7 of inorganic iodine, *Nature Geosci.*, 6, 108-111, 2013.

8 Chameides, W. L., and Davis, D. D.: Iodine: Its possible role in tropospheric photochemistry,
9 *J. Geophys. Res.*, 85, 7383-7398, 1980.

10 Draxler, R.R. and Rolph, G.D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated
11 Trajectory) Model access via NOAA ARL READY Website
12 (<http://ready.arl.noaa.gov/HYSPLIT.php>). NOAA Air Resources Laboratory, Silver Spring,
13 MD, 2014.

14 Garland, J. A., Elzerman, A. W., and Penkett, S. A.: The Mechanism for Dry Deposition of
15 Ozone to Seawater Surfaces, *J. Geophys. Res.*, 85, 7488-7492, 1980.

16 Gómez Martín, J. C., Mahajan, A. S., Hay, T. D., Prados-Román, C., Ordóñez, C.,
17 MacDonald, S. M., Plane, J. M. C., Sorribas, M., Gil, M., Paredes Mora, J. F., Agama Reyes,
18 M. V., Oram, D. E., Leedham, E., and Saiz-Lopez, A.: Iodine chemistry in the eastern Pacific
19 marine boundary layer, *Journal of Geophysical Research: Atmospheres*, 118, 887-904,
20 10.1002/jgrd.50132, 2013a.

21 Gómez Martín, J. C., Gálvez, O., Baeza-Romero, M. T., Ingham, T., Plane, J. M. C., and
22 Blitz, M. A.: On the mechanism of iodine oxide particle formation, *Phys. Chem. Chem. Phys.*,
23 15, 15612-15622, 10.1039/C3CP51217G, 2013b.

24 Großmann, K., Frieß, U., Peters, E., Wittrock, F., Lampel, J., Yilmaz, S., Tschritter, J.,
25 Sommariva, R., von Glasow, R., Quack, B., Krüger, K., Pfeilsticker, K., and Platt, U.: Iodine
26 monoxide in the Western Pacific marine boundary layer, *Atmos. Chem. Phys.*, 13, 3363-3378,
27 10.5194/acp-13-3363-2013, 2013.

28 Hönninger, G., Friedeburg, C. v., and Platt, U.: Multi axis differential optical absorption
29 spectroscopy (MAX-DOAS), *Atmos. Chem. Phys.*, 4, 231-254, 2004.

1 Jones, A. E., Anderson, P. S., Wolff, E. W., Roscoe, H. K., Marshall, G. J., Richter, A.,
2 Brough, N., and Colwell, S. R.: Vertical structure of Antarctic tropospheric ozone depletion
3 events: characteristics and broader implications, *Atmos. Chem. Phys.*, 10, 7775-7794,
4 10.5194/acp-10-7775-2010, 2010.

5 Lamarque, J. F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C.
6 L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.:
7 CAM-chem: description and evaluation of interactive atmospheric chemistry in the
8 Community Earth System Model, *Geosci. Model Dev.*, 5, 369-411, 10.5194/gmd-5-369-2012,
9 2012.

10 Lawler, M. J., Mahajan, A. S., Saiz-Lopez, A., and Saltzman, E. S.: Observations of I₂ at a
11 remote marine site, *Atmos. Chem. Phys.*, 14, 2669-2678, 10.5194/acp-14-2669-2014, 2014.

12 MacDonald, S. M., Gómez Martín, J. C., Chance, R., Warriner, S., Saiz-Lopez, A., Carpenter,
13 L. J., and Plane, J. M. C.: A laboratory characterisation of inorganic iodine emissions from the
14 sea surface: dependence on oceanic variables and parameterisation for global modelling,
15 *Atmos. Chem. Phys.*, 14, 5841-5852, 10.5194/acp-14-5841-2014, 2014.

16 Mahajan, A. S., Plane, J. M. C., Oetjen, H., Mendes, L., Saunders, R. W., Saiz-Lopez, A.,
17 Jones, C. E., Carpenter, L. J., and McFiggans, G. B.: Measurement and modelling of
18 tropospheric reactive halogen species over the tropical Atlantic Ocean, *Atmos. Chem. Phys.*,
19 10, 4611-4624, 10.5194/acp-10-4611-2010, 2010.

20 Mahajan, A. S., Gómez Martín, J. C., Hay, T. D., Royer, S. J., Yvon-Lewis, S., Liu, Y., Hu,
21 L., Prados-Roman, C., Ordóñez, C., Plane, J. M. C., and Saiz-Lopez, A.: Latitudinal
22 distribution of reactive iodine in the Eastern Pacific and its link to open ocean sources,
23 *Atmos. Chem. Phys.*, 12, 11609-11617, 10.5194/acp-12-11609-2012, 2012.

24 Mahajan, A. S., Prados-Roman, C., Hay, T. D., Lampel, J., Pöhler, D., Großmann, K.,
25 Tschritter, J., Frieß, U., Platt, U., Johnston, P., Kreher, K., Wittrock, F., Burrows, J. P., Plane,
26 J. M. C., and Saiz-Lopez, A.: Glyoxal observations in the global marine boundary layer,
27 *Journal of Geophysical Research: Atmospheres*, 119, 2013JD021388,
28 10.1002/2013JD021388, 2014.

29 Molina, M. J., and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes: Chlorine-
30 atom catalysed destruction of ozone, *Nature*, 249, 810-812, 1974.

1 Myhre, G., D. Shindell, D., Bréon, F.-M., Collins, W., Fuglestvedt, J., Huang, J., Koch, D.,
2 Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T.
3 and Zhang, H.: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The*
4 *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of*
5 *the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M.
6 Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)].
7 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, Chap.
8 8, 2013.

9 Ordóñez, C., Lamarque, J. F., Tilmes, S., Kinnison, D. E., Atlas, E. L., Blake, D. R., Sousa
10 Santos, G., Brasseur, G., and Saiz-Lopez, A.: Bromine and iodine chemistry in a global
11 chemistry-climate model: description and evaluation of very short-lived oceanic sources,
12 *Atmos. Chem. Phys.*, 12, 1423-1447, 10.5194/acp-12-1423-2012, 2012.

13 Platt, U., and Stutz, J.: Differential Absorption Spectroscopy, in: *Differential Optical*
14 *Absorption Spectroscopy, Physics of Earth and Space Environments*, Springer, Berlin
15 Heidelberg, 135-174, 2008.

16 Prados-Roman, C., Cuevas, C., Fernandez, R. P., Kinnison, D. E., Lamarque, J.-F., and Saiz-
17 Lopez, A.: A negative feedback between anthropogenic ozone pollution and enhanced ocean
18 emissions of iodine, accepted for ACPD, 2014.

19 Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P.,
20 Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night
21 marine air temperature since the late nineteenth century, *Journal of Geophysical Research:*
22 *Atmospheres*, 108, 4407, 10.1029/2002JD002670, 2003.

23 Read, K. A., Mahajan, A. S., Carpenter, L. J., Evans, M. J., Faria, B. V. E., Heard, D. E.,
24 Hopkins, J. R., Lee, J. D., Moller, S. J., Lewis, A. C., Mendes, L., McQuaid, J. B., Oetjen, H.,
25 Saiz-Lopez, A., Pilling, M. J., and Plane, J. M. C.: Extensive halogen-mediated ozone
26 destruction over the tropical Atlantic Ocean, *Nature*, 453, 1232-1235, 2008.

27 Saiz-Lopez, A., Plane, J. M. C., Mahajan, A. S., Anderson, P. S., Bauguitte, S. J.-B., Jones, A.
28 E., Roscoe, H. K., Salmon, R. A., Bloss, W. J., Lee, J. D., and Heard, D. E.: On the vertical
29 distribution of boundary layer halogens over coastal Antarctica: implications for O₃, HO_x,
30 NO_x and the Hg lifetime, *Atmos. Chem. Phys.*, 8, 887-900, 2008.

31 Saiz-Lopez, A., and von Glasow, R.: Reactive halogen chemistry in the troposphere, *Chem.*
32 *Soc. Rev.*, 41, 6448-6472, 10.1039/C2CS35208G, 2012.

1 Saiz-Lopez, A., Plane, J. M. C., Baker, A. R., Carpenter, L. J., Von Glasow, R., Gómez
2 Martín, J. C., McFiggans, G., and Saunders, R. W.: Atmospheric Chemistry of Iodine, Chem.
3 Rev. (Washington, DC, U. S.), 112, 1773-1804, 10.1021/cr200029u, 2012.

4 Saiz-Lopez, A., Fernandez, R. P., Ordóñez, C., Kinnison, D. E., Gómez Martín, J. C.,
5 Lamarque, J. F., and Tilmes, S.: Iodine chemistry in the troposphere and its effect on ozone,
6 Atmos. Chem. Phys. Discuss., 14, 19985-20044, 10.5194/acpd-14-19985-2014, 2014.

7 Wagner, T., Dix, B., Friedeburg, C. v., Frieß, U., Sanghavi, S., Sinreich, R., and Platt, U.:
8 MAX-DOAS O₄ measurements: A new technique to derive information on atmospheric
9 aerosols—Principles and information content, Journal of Geophysical Research:
10 Atmospheres, 109, D22205, 10.1029/2004JD004904, 2004.

11

1 Table 1. Description of the different legs of the Malaspina 2010 expedition. Due to technical
 2 problems, the O₃ and IO measurements presented in this work correspond to the period from
 3 21/02/2011 to 12/07/2011.

Legs	Docking places	Docking dates (dd/mm/yyyy)
1	Cadiz (Spain) - Rio de Janeiro (Brazil)	14/12/2010-13/01/2011
2	Rio de Janeiro (Brazil) – Cape Town (South Africa)	17/01/2011-06/02/2011
3	Cape Town (South Africa) – Perth (Australia) – Sydney (Australia)	11/02/2011-13/03/2011-30/03/2011
4	Sydney (Australia) – Auckland (New Zealand) – Honolulu (Hawaii)	04/04/2011-13/04/2011-08/05/2011
5	Honolulu (Hawaii) – Panama (Panama) - Cartagena de Indias (Colombia)	13/05/2011-10/06/2011-13/06/2011
6	Cartagena de Indias (Colombia) – Cartagena (Spain)	19/06/2011-14/07/2011


4

5


1 Table 2. Summary of the O₃ and ancillary parameters measured during Malaspina 2010. The
2 data correspond to daytime average values concurrent with the IO measurements gathered
3 during the expedition (Fig. 1).

Parameter	Mean	Std	Minimum	Maximum
O ₃ (nmol mol ⁻¹)	16.0	9.4	3.4	42.4
ws (m s ⁻¹)	7.0	2.0	3.3	11.6
SST (K)	298.9	2.9	291.6	303.0

4



19 Figure 1. Observations of surface ozone and ancillary parameters during Malaspina 2010
 20 (daily average). a. O₃ mixing ratios. b. Sea surface temperature. c. Wind speed. See also Table
 21 2.
 22




1

2 Figure 2. Backward trajectories of the air masses arriving at noon on every day of the
 3 Malaspina's cruise. They were calculated using HYbrid Single-Particle Lagrangian Integrated
 4 Trajectory (HYSPLIT, Draxler and Rolph, 2014).


5

a



1


b



2


3 Figure 3. Retrieval of IO during the Malaspina 2010 circumnavigation. a. Example of a
 4 typical IO spectral fit during the expedition. The particular spectrum was taken on the 31st
 5 May 2011 (3pm LT, 53° SZA) in the Eastern Pacific for a 2° elevation angle. The black line
 6 represents the measured IO optical density and the red line the retrieved one after the DOAS
 7 retrieval. This fit resulted in an IO dSCD of $(3.8 \pm 0.3) \times 10^{13}$ molecules cm^{-2} (i.e., IO vmr of
 8 0.8 ± 0.1 pmol mol^{-1}), with a residual optical density of 3.9×10^{-4} (root mean square). b.
 9 Timeline of the IO dSCD observed during the expedition. Statistically relevant data (i.e., data
 10 above the quality filters, SI) are shown with filled circles, and the non relevant data with

- 1 empty squares. The inset shows the daily evolution of IO dSCD for the 31st of May 2011. The
- 2 colour code indicates the elevation angle of the measurements.
- 3




1
2 Figure 4. Iodine oxide observations in the global marine boundary layer. IO mixing ratios (in
3 pmol mol⁻¹) are shown for five different field campaigns (Malaspina (this work), CHARLEX
4 (Gómez Martín et al., 2013a), TransBrom (Großmann et al., 2013), HaloCAST-P (Mahajan et
5 al., 2012), and Cape Verde (Read et al., 2008; Mahajan et al., 2010). For the three ship
6 campaigns (Malaspina, HaloCAST-P and TransBrom) daytime averaged values are shown.
7 For the long-term measurements on the Galapagos and the Cape Verde Islands (referred to as
8 CHARLEX and Cape Verde, respectively), the mean daytime IO values observed throughout
9 the campaigns are given.


10



1
2 Figure 5. Measured and modelled IO mixing ratios in different field campaigns and oceans.
3 For the two long-term campaigns on islands (Cape Verde- in green- and CHARLEX- in blue)
4 the mean daytime IO mixing ratio observed during the whole campaign period is given (filled
5 squares), together with the minimum and maximum observed values (dashed rectangles)
6 (Read et al., 2008; Mahajan et al., 2010; Jones et al., 2010). In the case of the Malaspina
7 circumnavigation (in red), daytime averaged IO mixing ratios are provided (filled squares)
8 along with their error (see also the SI). The shaded areas represent the standard deviation of
9 the modelled fields for the *NoPhot* (gray) and *Phot* (cyan) *Base* scheme. For comparison
10 purposes the IO vmr modelled considering only the organic iodine precursors (*Organic* run)
11 are also included (solid black line).
12





1
2



3
4
5
6
7
8



Figure 6. Simulated IO in the global marine environment (annually averaged). a. Geographical distribution of the total IO budget in the MBL (i.e., $\text{IO}_{\text{ISG}+\text{OSG}}$), in units of vmr (pmol mol^{-1}). b. Percentage contribution of the ISG emissions to the budget of IO in the global MBL.

a




1


b



2



1



2

3 Figure 7. Simulated percentage contribution of the different short-lived iodocarbons (a. CH_3I .
 4 b. CH_2ICl . c. CH_2I_2 . d. CH_2IBr) to the IO budget in the marine environment. Note that, for
 5 comparison purposes, the colour code is the same in the four panels. Also note that these
 6 model simulations do not include iodine emissions from ice surfaces. For the absolute values
 7 of the OSG emissions, please refer to the study of Ordóñez et al., 2012.

8