



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Determination of interfacial parameters of a soluble particle in a nonideal solution from measured deliquescence and efflorescence humidities

O. Hellmuth¹ and A. K. Shchekin²

¹Leibniz Institute for Tropospheric Research (TROPOS), 04318 Leipzig, Germany

²St Petersburg State University, Department of Statistical Physics, 198504 St Petersburg, Russia

Received: 28 May 2014 – Accepted: 3 August 2014 – Published: 8 September 2014

Correspondence to: O. Hellmuth (olaf@tropos.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In order to study the growth/shrinking of a hygroscopic nanoparticle during hydration/dehydration in an atmosphere of water vapour we have employed a thermodynamic approach proposed by Shchekin et al. (2008). This approach uses the mechanic and thermodynamic concept of disjoining pressure of thin films and allows, among others, the prediction of the humidity growth factor of both (i) a homogeneous solution droplet with completely dissolved residual core, and (ii) a heterogeneous solution droplet with partially dissolved residual core as a function of the ambient relative humidity. For application to a nanometric sodium chloride particle we have extended the original approach by (i) consideration of the nonideality of the solution through the dependence of molecular volumes of the solvent and solute molecules and the solute and solvent activities on the solution concentration, by (ii) derivation of an equation for the estimation of the efflorescence properties of a homogeneous solution droplet, and by (iii) combination with the empirical power law fittings for the size dependence of the deliquescence and efflorescence relative humidity values by Biskos et al. (2006a). It was demonstrated how the solution/solute interface energy and the correlation length of a thin solution film can be determined from a combination of experimentally determinable efflorescence and deliquescence humidities with the present calculus. The solution/solute interface energy was found to be in close agreement with some previous values reported in the literature, while it strongly differs from data of some other sources. The calculated deliquescence humidity shows a low sensitivity against the choice of the numerical value for the film correlation length. The estimated film correlation length of 1 nm for a nanometric sodium chloride particle with dry particle radius of 5 nm was found to be reconcilable with available a priori estimates of the correlation length from the literature when the measurement uncertainty of the deliquescence humidity is considered. Considering the combination of an extensive calculus, a comprehensive set of thermophysical constraints, and independent measurements of the deliquescence and efflorescence humidities as functions of dry particle radius, the ob-

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tained values of the solution/solute interface energy and the correlation length are in close agreement with previous estimations. The humidification of sodium chloride particles in the initial hydration stages was found to be very sensitive against the specification of the disjoining pressure. The enhancement of the wettability of the particle surface leads to an earlier onset of hygroscopic growth.

1 Introduction

Soluble substances (mostly salts) have a strong affinity for moisture, i.e., they are able to absorb water molecules from the ambient air. When the ambient relative humidity (RH) is sufficiently high, the water uptake continues until the hygroscopic particle completely dissolves in the absorbed water, in this way forming a pure solution droplet. The transition from partial to complete dissolution is called deliquescence, and the RH threshold at which a soluble particle completely dissolves is called deliquescence humidity (DRH). For highly soluble salts such as sodium chloride (NaCl) and small condensation nuclei, deliquescence occurs at subsaturation of the ambient partial water vapour with respect to a flat surface of pure bulk water ($RH < 100\%$). In the case of low soluble and larger nuclei, deliquescence can also occur in supersaturated vapour.

The reverse process, called efflorescence (or crystallisation), leads to hydrosol formation by precipitation of solid crystals from the solution within existing droplets. Crystallisation proceeds via two steps: (i) nucleation and (ii) crystal growth. In the nucleation step, solute molecules, dispersed in the solvent, start to form clusters on the nanometer scale. These clusters need to reach a critical size (embryonic stage) in order to freely grow up to solid precipitates within the maternal solution droplet. The critical size of the embryo depends on temperature, supersaturation, and the droplet size. The crystallisation is driven by the supersaturation of the solution, which depends on both ambient relative humidity and temperature. The relative humidity at which efflorescence sets in is called efflorescence humidity (ERH). The difference between DRH and ERH results in a hysteresis effect. Owing to the high variability of the relative humidity and temper-

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ature in the ambient atmosphere, e.g., due to multiscale turbulence, hysteresis effects effectively impact the partitioning between solid and aqueous phases of tropospheric NaCl and $(\text{NH}_4)_2\text{SO}_4$ particles. In ambient air, hygroscopic particles may exist in homogeneous and heterogeneous states. The homogeneous state can be either a pure solid core or a pure solution droplet. The heterogeneous state is a thermodynamically stable configuration in which a solid core substrate may coexist in thermodynamic equilibrium with an enveloping aqueous solution. The solid core can either consist of completely soluble matter or be a mixture of some insoluble fraction, coated with a soluble mantle of certain thickness. Here, we want to restrict our consideration to soluble one-component solid cores.

The atmospheric impact of hygroscopic particles is threefold:

- Hygroscopic particles may impact the global energy balance by a cooling effect (Wise et al., 2012, see references therein for the role of sea salt aerosol). The radiative properties of such particles depend, among others, on the liquid/solid phase partitioning within the particles (Wang et al., 2008a, 2008b). Aqueous particles have a larger mass extinction efficiency but a smaller backscattered fraction than their solid counterparts. Owing to the high variability of the relative humidity in the planetary boundary layer, hysteresis effects were found to effectively impact the partitioning between solid and aqueous phases of tropospheric salts on a global scale. For sulphate-ammonium particles Wang et al. (2008a) reported a significant impact of the solid/liquid phase partitioning on the sulphate direct climate forcing (SDFC). Depending on how these effects were parameterised in a global 3-D chemical transport model, an uncertainty in the SDFC of 23 % on the global scale (but of much higher values on regional scales) has been found from sensitivity studies (Wang et al., 2008b).
- Sea salt aerosol (SSA) is a major contributor to natural global aerosol emission and a main source of hygroscopic particles. According to Seinfeld and Pandis

**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(2006, see p. 61, Table 2.22 therein)¹ the best estimate of the SSA emission flux amounts $10\,100\text{ Tg yr}^{-1}$. This is nearly the forty-fold of the best estimate of the total (primary plus secondary) anthropogenic aerosol flux ($\approx 263\text{ Tg yr}^{-1}$), and 75% of the best estimate of the total (natural plus anthropogenic) aerosol flux ($\approx 13\,466\text{ Tg yr}^{-1}$). Natural SSA particles have a complex chemical composition. As many physical properties of SSA are controlled by NaCl (e.g., light scattering efficiency), several authors, e.g., Wise et al. (2012, see references therein) use NaCl as a proxy for SSA².

- Hygroscopic particles from SSA may serve as ice nuclei and influence cirrus formation. The mechanism of deliquescent-homogeneous and heterogeneous freezing and its atmospheric relevance has been intensively studied by Khvorostyanov and Sassen (1998a, b, 2002); Khvorostyanov and Curry (2000, 2004a, b, 2005, 2007, 2009, 2012); Khvorostyanov et al. (2001, 2003, 2006), and Curry and Khvorostyanov (2012). Recently, also efflorescent-heterogeneous nucleation onto SSA particles has attracted increasingly interest (Wise et al., 2012; Wagner and Möhler, 2013). Investigating the freezing capability of SSA particles Wise et al. (2012) pointed out the temperature dependence of phase composition of SSA particles: a brine solution and solid sodium chloride dihydrate ($\text{NaCl} \cdot 2\text{H}_2\text{O(s)}$) are stable

¹In the cited table only the global emission estimates for the major aerosol classes have been considered.

²The mining of salt in Europe can be traced back to about 4000 years ago (Schmiedel, 1984). Justus von Liebig characterised salt as the most valuable among all precious stones, and Cassiodor, roman statesman (6th century), stated that gold is dispensable but salt is not (Bontjer, 2006). In the middle ages salt was called “white gold”. Despite of its importance in daily life, environmental science, biophysics, catalysis etc., many fundamental physical and chemical properties of NaCl surfaces are still unknown and deserve a better understanding (Li, 2009, see chap. 3 therein).

**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from the triple point of water to the eutectic temperature³. Below the eutectic temperature solid ice and sodium chloride dihydrate are stable. The authors found the following efflorescence behaviour of aqueous NaCl particles: (i) at temperatures above the eutectic temperature solution particles effloresced into anhydrous solid NaCl particles, (ii) at temperatures below the eutectic temperature but above 236 K solution drops effloresced to a mixture of anhydrous and hydrated solid particles, (iii) at temperatures below 236 K solutions drops effloresced into hydrated solid particles. Wise and coworkers demonstrated the high efficiency of depositional ice nucleation on both crystalline anhydrous and hydrated NaCl particles, with an even better ice nucleation capability of hydrated particles. The authors referred to field observations of the chemical composition of anvil cirrus residue near the Florida peninsula, which revealed a SSA fraction of anvil ice residue of 26%. In the present case, however, we will restrict our consideration to a temperature well above the eutectic temperature, i.e., “eutectic solidification” will be excluded.

- Hygroscopic particles may act as chemical microreactors for hydrolysis, secondary organic aerosol formation, micellisation, and heterogeneous surface reactions.

From evaluation of nine years of radiosonde observations at five stations representing oceanic conditions (years 1980–1988, latitude from 14.3° S–57.2° N), Liu et al. (1991,

³“A eutectic system is a mixture of chemical compounds or elements that have a single chemical composition that solidifies at a lower temperature than any other composition made up of the same ingredients. This composition is known as the eutectic composition and the temperature at which it solidifies is known as the eutectic temperature” (taken from http://en.wikipedia.org/wiki/Eutectic_system, visited on 25 May 2014). A liquid with the eutectic composition solidifies at the eutectic temperature without prior solidification of one of the components of the mixture. Water and sodium chloride form a eutectic mixture. The eutectic temperature of aqueous sodium chloride solution amounts $T_E = 252\text{ K}$, the eutectic mass fraction of solute 23.3%.

Determination of interfacial parametersO. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fig. 2 therein) derived vertical profiles of the 9 years means of the relative humidity (inclusive standard deviations of individual measurements and standard deviations of the monthly means). These data reveal some common features of the humidity profile at all stations. The RH surface values were reported to vary between the stations in the range $\overline{RH}_0 \approx 75\text{--}82\%$ rh. The humidity profiles revealed their maximum values (spatially varying in the range $\overline{RH}_{\max} \approx 84\text{--}90\%$ rh) just below the height of the 900 hPa-surface. From the level of RH maxima, the mean relative humidity decreased to minimum values (spatially varying in the range $\overline{RH}_{\min} \approx 30\text{--}40\%$ rh) in the upper troposphere. These observations revealed that the vertical variation of the tropospheric relative humidity over the ocean safely exceeds the humidity interval fixed by the deliquescence and efflorescence humidities of sodium chloride particles (Seinfeld and Pandis, 2006, see Sect. 10.2 therein; Biskos et al., 2006b). Thus, in the marine troposphere sodium chloride particles are expected to exist both in homogeneous (either as an aqueous solution or as pure solid) and in heterogeneous (a solid core enveloped by a liquid solution film) states depending on the particle size.

The study of the deliquescence behaviour of hygroscopic particles in the limit of bulk properties is well understood (Seinfeld and Pandis, 2006, see, e.g., Sect. 10.2 therein). In contrast to this, the investigation of size-dependent effects in both deliquescence and efflorescence is comparatively rare and not completely understood so far.

McGraw and Lewis (2009) investigated the size-dependent deliquescence/efflorescence phase transformation for particles down to several nanometers in size. The authors analysed the thermodynamic properties of inorganic salt particles, coated with aqueous solution layers of varying thickness and surrounded by vapour. In order to define a limiting deliquescence relative humidity (RH_D) for small particles a thin layer criterion (TLC) has been introduced, which is based on the fulfillment of two equilibrium conditions: (1) equality of chemical potentials between salt in an undissolved core, and thin adsorbed solution layer, and (2) equality of chemical potentials between water in the thin layer and vapour phase. For the limit of large dry particle size the bulk deliquescence properties were obtained. Nanosize particles were

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reported to deliquesce at relative humidity just below the RH_D on crossing a nucleation barrier, located at a critical solution layer thickness. This nucleation barrier will vanish at the RH_D defined by the TLC. The authors employed concepts and methods from nucleation theory including the kinetic potential, self-consistent nucleation theory, nucleation theorems, and the Gibbs dividing surface, altogether providing the theoretical foundation of their approach. McGraw and Lewis (2009) emphasised the existence of unifying features of small particle deliquescence/efflorescence processes such as common thermodynamic area constructions and a common free-energy surface. However, the approach of McGraw and Lewis (2009) is based on the employment of the classical capillary drop model, but does not explicitly consider measurable thin film quantities such as the disjoining pressure, which is indispensable for the description of the initial stages of hygroscopic growth from the formation of first monolayers up to thick films. Here, we can already antedate that the approach of McGraw and Lewis (2009) and the one presented in this paper essentially differ in the way of “metrisation” of the small particle deliquescence/efflorescence processes leading to complementary information about these processes.

Bruzewicz et al. (2011) characterised the behaviour of NaCl nanoparticles as a function of RH. The authors measured the heights of particles deposited on a prepared hydrophobic surface by means of noncontact environmental atomic force microscopy and found a cubic shape of NaCl nanoparticles with sides of 35 and 80 nm. These nanoparticles were observed to reversibly take up water with increasing RH well below the bulk DRH of 75 %rh at 23 °C, forming of a liquid-like surface layer of thickness 2 to 5 nm with measurable uptake (> 2 nm increase in particle height) beginning at 70 %rh. The maximum thickness of the layer was observed to increase with increasing RH and increasing particle size over the investigated size range. The liquid-like layer appeared as a reversible rounding at the upper surface of the particles with the shape of the layer fitting to a parabolic cross-section. The ratio of particle height to maximum radius of curvature was reported to increase from zero (flat top) at 68 %rh to 0.7 ± 0.3 at 74 %rh. From their observations Bruzewicz et al. (2011) concluded the occurrence of

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mass reorganisation on the solid NaCl nanocrystal at RH below the DRH, suggesting that the deliquescence of NaCl nanoparticles “is more complex than an abrupt first-order phase transition”. The height measurements were found to be consistent with a phenomenological model that assumes favourable contributions to the free energy of formation of a liquid layer on solid NaCl due to both van der Waals interactions (depending partly upon the Hamaker constant of the interaction between the thin liquid film and the solid NaCl), and to a longer-range electrostatic interaction over a characteristic length of persistence. The authors derived best fit values for the Hamaker constant and the persistence length scale.

Using the thermodynamic theory of thin solution films developed by Shchekin et al. (1993); Kuni et al. (2001); Djikaev et al. (2001a, b) and Djikaev (2002), which is based on the mechanical and thermodynamic concept of the disjoining pressure, Shchekin and Shabaev (2005); Shchekin and Rusanov (2008) and Shchekin et al. (2008) derived for the limit of ideality of the aqueous electrolyte solution generalisations of the Gibbs–Kelvin–Köhler (GKK) equation of the theory of nucleation on soluble particles and of the Ostwald–Freundlich (OF) equation of the theory of solutions (hereafter called SSR theory according to the initials of the authors Shchekin, Shabaev, and Rusanov). These equations allow (i) the determination of the humidity growth factor of a hygroscopic particle⁴ as a function of ambient temperature, relative humidity, dry particle radius and of different thermophysical and interfacial parameters of the system, and (ii) the determination of the DRH.

Based on the SSR theory, the present study pursues the following aims:

1. First, the generalised GKK and OF equations as predicted by the SSR theory will be extended to determine the humidity growth factor and DRH also for nonideal aqueous solutions.

⁴The humidity growth factor is defined as the ratio of the radius of the hygroscopic particle to the dry particle radius.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2. In contrast to the DRH determination, the ERH cannot be obtained from a pure thermodynamic approach alone, but requires a kinetic consideration (e.g., Onasch et al., 2000, Pant et al., 2006, Parsons et al., 2006, Shchekin and Shabaev, 2010⁵, and Shchekin et al., 2013). Therefore second, a governing equation for determination of the ERH on the base of the classical nucleation theory, applied to homogeneous salt crystallisation from a supersaturated solution, will be derived.

3. Third, it will be demonstrated how the extended SSR theory in combination with CNT can be used to determine the specific interfacial parameters, namely the solution/salt interfacial energy, and the characteristic correlation length of a thin film, which are difficult to obtain from direct measurements. In the present approach the mentioned interfacial parameters will be estimated by considering size-dependent DRH and ERH values obtainable from measurements.

4. Fourth, by application of the calculus to a nanometric NaCl particle in a water vapour environment we will show that the approach presented here is able to theoretically describe the phenomenon observed by Bruzewicz et al. (2011) at least in a qualitative way.

The paper is structured as follows. In Sect. 2 we will briefly describe the model, in Sect. 3 we will present the method for the estimation of interfacial properties, in Sect. 4 we will present and discuss the model results, and the last section contains the summary and conclusions. *A comprehensive detailed outline of the calculus can be found in the Supplement (SM).*

⁵Shchekin and Shabaev extended their approach to the determination of the activation barriers for the complete dissolution of a condensation nucleus and its reverse crystallisation in droplets in a subsaturated solvent vapour between DRH and ERH. The authors showed that the coexistence of drops with partly and completely dissolved nuclei is impossible.

2 Model description

2.1 Determination of humidity growth factor and the deliquescence humidity according to SSR theory

2.1.1 Governing equations

5 The basic scenario and assumptions are described in SM/Sect. S1.1. In the initial state, the system consists of a soluble solid core (phase γ) with dry particle radius R_N , embedded in a solvent vapour (phase β) within fixed volume V at absolute temperature T . In the final state, a liquid droplet (phase α) of radius R is formed, depending on the ambient RH either including a partially dissolved residual core of radius R'_N enveloped
10 by a liquid film of thickness $h = R - R'_N$ (phase α), or the residual core being completely dissolved, $R'_N = 0$. The formation work ΔF , required to form a heterogeneous droplet from a vapour, reads (see SM/Sect. 1.3, Eq. S43):

$$\begin{aligned}\Delta F &= -\Delta N^{(\beta \rightarrow \alpha)} \left[\mu^{(\beta)} - \mu_*^{(\beta \rightarrow \alpha)} \right] + \Delta N^{(\gamma \rightarrow \alpha)} \left[\mu_*^{(\gamma \rightarrow \alpha)} - \mu_\infty^{(\gamma)} \right] + 4\pi R^2 \sigma_\infty^{(\alpha\beta)} + \Delta F_W, \\ \Delta F_W &= 4\pi R_N'^2 \left[\sigma_\infty^{(\alpha\gamma)} + l^{(*)} \Pi_D(h) \right] - 4\pi R_N^2 \sigma_\infty^{(\beta\gamma)}, \\ \mu_*^{(\beta \rightarrow \alpha)}(x) &= \mu_\infty^{(\beta)} + k_B T \ln \left[\frac{a_{\text{solvent}}(x)}{a_{\text{solvent}}(x=0)} \right], \\ \mu_*^{(\gamma \rightarrow \alpha)}(x) &= \mu_\infty^{(\gamma)} + k_B T \ln \left[\frac{a_{\text{solute}}(x)}{a_{\text{solute}}(x_\infty)} \right].\end{aligned}\quad (1)$$

15 Here, $\Delta N^{(\beta \rightarrow \alpha)}$ denotes the number of solvent molecules condensed from the vapour into the solution film, and $\Delta N^{(\gamma \rightarrow \alpha)}$ is the number of solute molecules dissolved from the solute into the solution film, respectively. The quantity $\mu^{(\beta)}(T, S^{(\beta)})$ is the chemical potential of the solvent vapour, which is a function of temperature and vapour saturation ratio $S^{(\beta)}$; $\mu_\infty^{(\beta)}$ is the value of $\mu^{(\beta)}$ at equilibrium of the vapour with a pure bulk

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

liquid. The chemical potential differences $\mu_*^{(\beta \rightarrow \alpha)}(x) - \mu_\infty^{(\beta)}$ and $\mu_*^{(\gamma \rightarrow \alpha)}(x) - \mu_\infty^{(\gamma)}$ represent the activity contributions to the chemical potentials of solvent and solute molecules in the solutions, $\mu^{(\beta \rightarrow \alpha)}$ and $\mu^{(\gamma \rightarrow \alpha)}$, respectively, and $\mu_\infty^{(\gamma)}$ is the equilibrium value of the chemical potential of the solid matter at the flat interface between the solid substrate and a bulk solution at saturation vapour pressure. The quantity $a_{\text{solvent}}(x)$ is the solvent (water) activity as a function of the relative solute concentration $x = mM_W / (1 + mM_W)$ in the film with m denoting the molality of the solution and M_W the molar mass of water. Analogously, $a_{\text{solute}}(x)$ is the solute (salt) activity as a function of the relative solute concentration in the film, and x_∞ is the relative solute concentration corresponding to the solubility of the solute.

The term ΔF_W in Eq. (1) represents the work of wetting of a soluble nucleus. The derivation of this term is based on the negligence of adsorption of salt at the droplet surface. Furthermore, all surface tensions and parameters of the disjoining pressure have been taken as independent of salt concentration. The function $\Pi_D(h) = K^{(*)} \exp(-h/l^{(*)})$ appearing in the term ΔF_W is the disjoining pressure of a thin solution film with $K^{(*)} = s/l^{(*)}$ denoting a characteristic pressure scale of the thin film. The quantity $s = \sigma_\infty^{(\beta\gamma)} - (\sigma_\infty^{(\alpha\beta)} + \sigma_\infty^{(\alpha\gamma)})$ therein is the nonequilibrium spreading coefficient with $\sigma_\infty^{(\alpha\beta)}$, $\sigma_\infty^{(\alpha\gamma)}$, and $\sigma_\infty^{(\beta\gamma)}$ denoting the surface tension of the bulk solution (in the absence of vapour adsorption), the solution/solute and vapour/solute interfacial energies, respectively (Shchekin et al., 2008, Eqs. 28 and 29 therein). The quantity $l^{(*)}$ is the film correlation length, which is of the order of the intermolecular separation in the film (Rusanov et al., 1994). Furthermore, k_B denotes the Boltzmann constant. The equilibrium state of the droplet is controlled by two thermodynamic conditions corresponding to the locations of the minima and saddle points of the formation work (see SM/Sect. S1.4):

- the “external solvent equilibrium” between the solvent in the vapour phase β and the solvent in the liquid phase α ;

- the “internal solute equilibrium” between the solute in the residual solid phase γ' and the solute in the liquid phase α .

At external equilibrium, the thermodynamic driving force for the transfer of solvent molecules from the vapour phase into the liquid phase must vanish. From this condition one can arrive at the following *generalisation of the Gibbs–Kelvin–Köhler equation of the theory of nucleation on soluble particles* (see SM/ Sect. S1.4, Eq. S54):

$$\mathcal{F}_{\text{GKK}} \left(S^{(\beta)}, R'_N, R \right) \Big|_{\substack{T=\text{const} \\ R_N=\text{const}}} = -k_B T \ln S^{(\beta)} + k_B T \ln \left[\frac{a_{\text{solvent}}(x(R, R'_N))}{a_{\text{solvent}}(0)} \right] + v^{(\beta \rightarrow \alpha)}(x) \left[\frac{2\sigma_{\infty}^{(\alpha\beta)}}{R} - \left(\frac{R'_N}{R} \right)^2 \Pi_D(h) \right] = 0. \quad (2)$$

In Eq. (2), $S^{(\beta)}$ denotes the ratio of the actual water vapour pressure to the saturated water vapour pressure over bulk water (hereafter called water vapour saturation ratio in accordance with Seinfeld and Pandis, 2006, p. 813 therein), and $v^{(\beta \rightarrow \alpha)}(x)$ is the partial molecular volume of solvent molecules in the solution.

At internal equilibrium, the thermodynamic driving force for the transfer of solute molecules from the solid phase into the liquid phase must vanish. This leads to the following *generalisation of the Ostwald–Freundlich equation of the theory of solutions* (see SM/ Sect. S1.4, Eq. S56):

$$\mathcal{F}_{\text{OF}} \left(R'_N, R \right) \Big|_{\substack{T=\text{const} \\ R_N=\text{const}}} = k_B T \ln \left[\frac{a_{\text{solute}}(x(R, R'_N))}{a_{\text{solute}}(x_{\infty})} \right] + v^{(\gamma \rightarrow \alpha)}(x) \left[\frac{2\sigma_{\infty}^{(\alpha\beta)}}{R} - \left(\frac{R'_N}{R} \right)^2 \cdot \Pi_D(h) \right] - v^{(\gamma)} \left[\frac{2\sigma_{\infty}^{(\alpha\beta)}}{R} - \left(\frac{R'_N}{R} \right)^2 \Pi_D(h) + \frac{2\sigma_{\infty}^{(\alpha\gamma)}}{R'_N} + \left(1 + \frac{2l^{(*)}}{R'_N} \right) \Pi_D(h) \right] = 0. \quad (3)$$

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In Eq. (3), $v^{(y)}$ and $v^{(y \rightarrow \alpha)}(x)$ are the partial molecular volumes of solute molecules in the solid phase and of dissolved solute molecules in the solution film, respectively.

Equations (2) and (3) are extensions of the generalised GKK &OF equations as derived by Shchekin et al. (2008, Eqs. 25 and 26 therein) for the limit of an ideal solution⁶ and serve as governing equations for determination of the humidity growth factor and DRH.

2.1.2 Determination of the humidity growth factor

As demonstrated by Shchekin et al. (2008, see Sect. V, Fig. 6 therein), the dependence of the droplet radius R on the vapour saturation ratio $S^{(\beta)}$ at equilibrium between the droplet and the ambient vapour can be fully determined from simultaneous solution of the generalised GKK and OF Eqs. (2) and (3) (see SM/Sect. S1.5). Knowing the relation $R = f(T, R_N, S^{(\beta)})$, the measurable vapour condensation growth factor GF can be directly determined:

$$GF = \frac{R}{R_N} = R_N^{-1} f(T, R_N, S^{(\beta)}). \quad (4)$$

⁶In derivation of their generalised GKK Eq. (25), Shchekin et al. (2008) employed the approximations $a_{\text{solvent}}(x) \approx 1 - x$, (with the Taylor expansion $\ln a_{\text{solvent}}(x) \approx -x$) and $a_{\text{solute}}(x) \approx x$ for the limit of an ideal solution. Furthermore, the molality-dependent partial molecular volume of solvent molecules in the solution, $v^{(\beta \rightarrow \alpha)}(x)$, has been approximated by an average value for the molecular volume of solution molecules in the solution, $\bar{v}^{(\alpha)}$. Analogously, in derivation of their generalised OF Eq. (26) the authors employed the ideal-solution limit $a_{\text{solute}}(x) \approx x$ and approximated the molality-dependent partial molecular volume of solute molecules in the solution, $v^{(y \rightarrow \alpha)}(x)$, by an average value $\bar{v}^{(y \rightarrow \alpha)}$. Note that Eqs. (25) and (26) of Shchekin et al. (2008) can be obtained as special cases of Eqs. (2) and (3), but, vice versa, the latter cannot be obtained by formal replacement of quantities referring to the ideal-solution limit in Eqs. (25) and (26) of Shchekin et al. (2008).

2.1.3 Determination of the deliquescence humidity

Prompt transition from the equilibrium state with the partially dissolved residual core to the equilibrium state with the completely dissolved residual core (“deliquescence transition”) is a barrierless process. As demonstrated by Shchekin et al. (2008, see Sect. V, Figs. 4–6 therein), the determination of the vapour saturation ratio at deliquescence, $S_{\text{th}}^{(\beta)}$ = DRH/100%, proceeds as follows (see SM/Sect. S1.6):

- Firstly, the drop radius is determined as a function of the residual radius $R = R(R'_N)$ from the numerical solution of the transcendental OF Eq. (3).
- Secondly, we employ the expression for the chemical potential of solvent molecules in the liquid film given in thermal units, $b^{(\beta \rightarrow \alpha)}$ (see SM/Sect. S1.6, Eq. S62):

$$b^{(\beta \rightarrow \alpha)}(R, R'_N) = \frac{\mu^{(\beta \rightarrow \alpha)} - \mu_{\infty}^{(\beta)}}{k_B T} = \ln \left[\frac{a_{\text{solvent}}(x(R, R'_N))}{a_{\text{solvent}}(0)} \right] + \frac{v^{(\beta \rightarrow \alpha)}(x(R, R'_N))}{k_B T} \times \left[\frac{2\sigma_{\infty}^{(\alpha\beta)}}{R} - \left(\frac{R'_N}{R} \right)^2 \Pi_D(h(R, R'_N)) \right]. \quad (5)$$

Here, $\mu^{(\beta \rightarrow \alpha)}$ and $\mu_{\infty}^{(\beta)}$ are the chemical potentials of the solvent molecules in the liquid film and of the vapour at equilibrium with a pure bulk liquid having a flat interface. Inserting function $R'_N = R'_N(R)$ into Eq. (5) delivers the chemical potential of solvent molecules in the solution as a function of drop radius, $b^{(\beta \rightarrow \alpha)} = b^{(\beta \rightarrow \alpha)}(R)$.

- Thirdly, from $b^{(\beta \rightarrow \alpha)} = b^{(\beta \rightarrow \alpha)}(R)$ the maximum value $b_{\text{th}}^{(\beta \rightarrow \alpha)}$ is determined, which allows us to find the deliquescence vapour saturation ratio according to $S_{\text{th}}^{(\beta)} = \exp(b_{\text{th}}^{(\beta \rightarrow \alpha)})$ and the deliquescence relative humidity defined by DRH = 100% $\times S_{\text{th}}^{(\beta)}$.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Determination of the efflorescence humidity according to CNT

The prompt transition from a homogeneous solution droplet to a solution droplet with a stable crystal core is called efflorescence transition. It performs via homogeneous crystallisation, i.e., precipitation of solute crystals from a supersaturated solution droplet. Supersaturation of the solution can be achieved by decreasing the chemical potential of the solvent vapour (or by decreasing the temperature). Here, for a such confined system as a submicron droplet, the efflorescence is assumed to be the result of only one successful crystallisation event inside the solution droplet.

Based on the classical nucleation theory (CNT) (here, homogeneous nucleation) and the efflorescence condition employed by e.g., Gao et al. (2007); Onasch et al. (2000); Pant et al. (2006) and Parsons et al. (2006), one can derive the following governing equation for the determination of the relative solute concentration of a solution droplet with completely dissolved residual core ($R'_N = 0$) at efflorescence transition, $x_{\text{effl}} = x(\text{ERH})$ (see SM/Sect. S2.2, Eqs. S81 and S82):

$$\mathcal{F}_{\text{CNT}}(R(x_{\text{effl}})) \Big|_{\substack{T=\text{const} \\ R'_N=\text{const} \\ R'_N=0}} = \left\{ \frac{16\pi}{3} \left[\frac{\sigma_\infty^{(\alpha\gamma)}}{k_B T} \right]^3 \frac{[v^{(\gamma)}]^2}{\ln \left[\frac{4}{3} \pi [R(x_{\text{effl}})]^3 J_{\text{hom},0} \tau_{\text{ind}} \right]} \right\}^{1/2} - \ln \left[\frac{a_{\text{solute}}(x_{\text{effl}})}{a_{\text{solute}}(x_\infty)} \right] = 0, \quad (6)$$

$$J_{\text{hom},0} = [v^{(\gamma)}]^{-1} \left(\frac{k_B T}{h} \right) \exp \left(- \frac{\Delta F_{\text{act}}}{k_B T} \right).$$

In Eq. (6), $J_{\text{hom},0}$ denotes the kinetic prefactor in the nucleation rate expression. The terms $k_B T/h$ and $\Delta F_{\text{act}}/(k_B T)$ are the frequency and free energy barrier (in thermal units) of a solute molecule to jump from the solution to the crystalline nucleus, respectively, where h denotes the Planck constant. The quantity τ_{ind} is the induction time,

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



defined as the time the system spent at constant droplet composition prior to homogeneous nucleation of the first supercritical nucleus.

To find the root of Eq. (6), i.e., the droplet radius at the onset of efflorescence, one has to consider a closure condition to determine the relative solute concentration as a function of the initial and residual core radii and the droplet radius. The closure condition is given by the condition of preserving the salt mass in the droplet, written in form of a composition-geometry implicit relation $\mathcal{F}_{\text{comp-geom}}(x, R_N, R'_N, R) = 0$. With consideration of this relation, one can determine x_{eff} and $R_{\text{eff}} = R(\text{ERH})$ from the transcendental Eq. (6). For known R_{eff} , the efflorescence humidity, $\text{ERH} = 100\% \times S_{\text{eff}}^{(\beta)}$, is the root of the relation $R = f(T, R_N, S^{(\beta)})$ referring to the solution droplet with a completely dissolved residual core ($R'_N = 0$) (see SM/Sect. S2.2).

2.3 Thermophysical parameters for the aqueous sodium chloride system

For application of the calculus to an aqueous sodium chloride system various thermodynamic and interfacial quantities must be known as input parameters, which were taken from the literature and/or derived employing additional assumptions:

- The solvent and solute activities $a_{\text{solvent}}(x)$ and $a_{\text{solute}}(x)$ for nonideal NaCl solution are determined according to SM/Sect. S3.1.
- The partial molecular volumes $v^{(\beta \rightarrow \alpha)}(x)$, $v^{(\gamma \rightarrow \alpha)}(x)$, and $v^{(\gamma)}$ are determined according to SM/Sect. S3.2.
- The composition–geometry relation $\mathcal{F}_{\text{comp-geom}}(x, R_N, R'_N, R) = 0$ has been derived in SM/Sect. S3.3 and is given by Eq. (S120).
- The relative solute concentration of sodium chloride solubility in water, x_{∞} , is given in SM/Sect. S3.4.
- The activation energy for solution/solute interface-crossing diffusion, $\Delta F_{\text{act}}(T, x)$, has been determined according to SM/Sect. S3.5.

22731

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- For the induction time a value of $\tau_{\text{ind}} = 1$ s has been used (Gao et al., 2007).
- For the surface tension of an aqueous NaCl solution a fixed value of $\sigma_{\infty}^{(\alpha\beta)} = 0.083 \text{ J m}^{-2}$ according to Russell and Ming (2002) was used (see SM/Sect. S3.6).
- Based on computer simulations of Li (2009), the reference value of the vapour/solid interfacial energy was specified as $\sigma_{\infty}^{(\beta\gamma)} = 0.24 \text{ J m}^{-2}$, but varied within the framework of sensitivity studies (see SM/Sect. S3.6).
- Owing to their large uncertainties both the solution/solute interfacial energy, $\sigma_{\infty}^{(\alpha\gamma)}$, and the film correlation length, $l^{(*)}$, have been estimated from measured ERH and DRH values (see SM/Sects. S3.6 and S3.7).

3 Estimation of the solution/solute interfacial energy and the film correlation length

3.1 Determination of the solution/solute interfacial energy

The “best guess” estimation of $\sigma_{\infty}^{(\alpha\gamma)}$ proceeds in three steps:

- At first, from combination of the generalised GKK equation, \mathcal{F}_{GKK} , and the composition–geometry relation, $\mathcal{F}_{\text{comp-geom}}$, specified to the case of a completely dissolved residual core ($R'_N = 0$), one gets the efflorescence radius, R_{effl} , as a function of the measured vapour saturation ratio at efflorescence, $S_{\text{effl, meas}}^{(\beta)}(R_N)$.
- Secondly, from $\mathcal{F}_{\text{comp-geom}}$ one gets the relative solute concentration at efflorescence, χ_{effl} , as a function of R_{effl} .
- Thirdly, the solution/solute interface energy is adjusted in such a way that at the measured ERH exactly one crystallite forms inside the volume of the homogeneous droplet, $V_{R, \text{effl}}$, during the induction time.

The expression for the homogeneous nucleation rate, J_{hom} , is given in SM/ Sect. S2.2, Eq. (S81). On this base, one can define a suitable cost function $\mathcal{F}_{\text{cost}}^{(\sigma)}$ for the determination of $\sigma_{\infty}^{(\alpha\gamma)}$. The whole calculus reads:

$$\begin{aligned} \mathcal{F}_{\text{GKK}} + \mathcal{F}_{\text{comp-geom}} &\rightsquigarrow R_{\text{effl}} = f(T, R_N, S_{\text{effl, meas}}^{(\beta)}) \\ \mathcal{F}_{\text{comp-geom}} &\rightsquigarrow X_{\text{effl}} = X_{\text{effl}}(R_{\text{effl}}) \\ \mathcal{F}_{\text{cost}}^{(\sigma)} \left\{ \sigma_{\infty}^{(\alpha\gamma)} \right\} &= \ln \left[\underbrace{J_{\text{hom}} \left\{ \sigma_{\infty}^{(\alpha\gamma)}, X_{\text{effl}}, \dots \right\}}_{\stackrel{!}{=} 1\#} V_{R, \text{effl}} \tau_{\text{ind}} \right]. \end{aligned} \quad (7)$$

The sought-after value of $\sigma_{\infty}^{(\alpha\gamma)}$ is found if for given $S_{\text{effl, meas}}(R_N)$ the constraint $\mathcal{F}_{\text{cost}}^{(\sigma)} \left\{ \sigma_{\infty}^{(\alpha\gamma)} \right\} \stackrel{!}{=} 0$ is fulfilled.

3.2 Determination of the correlation length

Similarly, the “best guess” estimation of $l^{(*)}$ proceeds in the following three steps:

- At first, $\sigma_{\infty}^{(\alpha\gamma)}$ obtained from the previous step is used to determine the spreading coefficient s .
- Secondly, the theoretical deliquescence humidity, DRH_{SSR} , is determined using the SSR theory according to Sect. 2.1.3.
- Thirdly, $l^{(*)}$ is adjusted in such a way that the theoretical value DRH_{SSR} equals the measured value $\text{DRH}_{\text{meas}}(R_N)$ for given radius of the dry particle.

On this basis, one can define another cost function $\mathcal{F}_{\text{cost}}^{(l)}$ for determination of $l^{(*)}$:

$$\mathcal{F}_{\text{cost}}^{(l)} \left\{ l^{(*)} \right\} = \text{DRH}_{\text{SSR}} \left\{ l^{(*)}, \dots \right\} - \text{DRH}_{\text{meas}}. \quad (8)$$

22733

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The sought-after value of $f^{(*)}$ is found if for given $\text{DRH}_{\text{meas}}(R_N)$ the constraint $\mathcal{F}_{\text{cost}}^{(l)} \{f^{(*)}\} \stackrel{!}{=} 0$ is fulfilled.

3.3 Size dependence of deliquescence and efflorescence humidities

Based on laboratory studies Biskos et al. (2006a) derived the following power law fits for the size dependence of the deliquescence and efflorescence relative humidity values, DRH and ERH, of NaCl particles with dry mobility equivalent diameters in the range $6 < D_{\text{m,dry}}/\text{nm} < 60$ (see Fig. 1):

$$\frac{\text{DRH}_{\text{Biskos}}}{\% \text{rh}} = 213 \left(\frac{D_{\text{m,dry}}}{\text{nm}} \right)^{-1.6} + 76, \quad (9)$$

$$\frac{\text{ERH}_{\text{Biskos}}}{\% \text{rh}} = 213 \left(\frac{D_{\text{m,dry}}}{\text{nm}} \right)^{-1.65} + 44.$$

For the present purposes, we have replaced the dry mobility equivalent diameter, $D_{\text{m,dry}}$, by the dry volume equivalent diameter, $D_{\text{V,dry}}$ ⁷. For the relative humidity mea-

⁷The relation between the dry mobility equivalent diameter and the dry volume equivalent diameter, $D_{\text{V,dry}}$, reads (e.g., Romakkaniemi et al., 2001, Hämeri et al., 2001):

$$D_{\text{V,dry}} = \left[\frac{1}{\chi} \frac{C(D_{\text{V,dry}})}{C(D_{\text{m,dry}})} \right] D_{\text{m,dry}}, \quad \chi = 1.08,$$

$$C(D) = 1 + \left(\frac{133}{D/\text{nm}} \right) \times \left[1.142 + 0.588 \exp \left(-\frac{0.999 D/\text{nm}}{133} \right) \right].$$

Here, χ denotes the shape factor and $C(D)$ the Cunningham slip correction factor. For $D_{\text{m,dry}} \approx D_{\text{V,dry}} = 30 \text{ nm}$ we obtain $\text{DRH}_{\text{Biskos}} = 76.92\%$ and $\text{ERH}_{\text{Biskos}} = 44.78\%$. With consideration of the Cunningham slip correction we obtain $D_{\text{m,dry}} = 31.2 \text{ nm}$, $\text{DRH}_{\text{Biskos}} = 76.87\%$ and

22734

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



surements at several points throughout the apparatus using capacitance sensors, Biskos et al. (2006b, see p. 98 therein) communicated an “accuracy” of $\pm 2.5\%$ rh. In their abstract, however, the authors stated that the power law fits were calibrated with “less than 1%rh uncertainty.” Within the framework of sensitivity studies we have varied the efflorescence and deliquescence humidities as follows:

$$\begin{aligned} \text{ERH} &= \text{ERH}_0 + \Delta\text{ERH}, \\ \text{DRH} &= \text{DRH}_0 + \Delta\text{DRH}. \end{aligned} \quad (10)$$

Here, ERH_0 and DRH_0 are the nominal values of the efflorescence and deliquescence humidity according to Eq. (9), and ΔERH and ΔDRH are the corresponding uncertainties, which are assumed to be independent from each other. Within the framework of sensitivity studies we have assumed three different values for the uncertainties: -1% rh, 0% rh, $+1\%$ rh.

4 Results and discussion

4.1 Reference case calculation

The subsequent calculations were performed at $T = 298\text{ K}$ for a dry NaCl particle radius of $R_N = 5\text{ nm}$, corresponding to $\text{DRH} \approx 81\%$ rh $\text{ERH} \approx 47\%$ rh according to Eq. (9).

At first, we start with the determination of $\sigma_\infty^{(\alpha\gamma)}$ with consideration of the uncertainty in the ERH measurement (see Table 1 and Fig. 2). Figure 2 shows the cost function $\mathcal{F}_{\text{cost}}^{(\sigma)}$ as a function of the solution/solute interfacial energy $\sigma_\infty^{(\alpha\gamma)}$ according to Eq. (7) for three different ΔERH values corresponding to graphs 1, 2, and 3, respectively, in Fig. 2. For the best guess of the solution/solute interfacial energy at the nominal ERH a value of $\sigma_\infty^{(\alpha\gamma)} \approx 0.088\text{ J m}^{-2}$ was obtained, which is considerably larger than the value

$\text{ERH}_{\text{Biskos}} = 44.73\%$. The difference to the approximative results is rather small and can be neglected for the present purposes.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of $\sigma_{\infty}^{(\alpha\gamma)} = 0.029 \pm 0.020 \text{ J m}^{-2}$ given by Russell and Ming (2002, see references therein), but which is close to the value of $\sigma_{\infty}^{(\alpha\gamma)} \approx 0.0895 \text{ J m}^{-2}$ obtained by Gao et al. (2007). Secondly, we determined $l^{(*)}$ with consideration of the uncertainty in both the ERH and DRH measurements (see Table 2 and Fig. 3). Figure 3 shows the cost function $\mathcal{F}_{\text{cost}}^{(l)}$ as a function of the correlation length $l^{(*)}$ according to Eq. (8) for different ERH and DRH values. The sensitivity of the correlation length against uncertainties in the ERH measurements is very small. However, for the assumed DRH uncertainty the correlation length varies in the range $l^{(*)} \approx (0.7\text{--}1.5) \text{ nm}$ with a best guess value of $l^{(*)} \approx 1 \text{ nm}$ at the nominal DRH value. This value is somewhat larger than the value proposed by Marčelja and Radić (1976) ($l^{(*)} \approx 0.3 \text{ nm}$) and the values obtained from density functional theory and discussed in detail in SM/Sect. S3.7 ($l^{(*)} \approx (0.4\text{--}0.54) \text{ nm}$). However, considering (i) that the present approach is based on completely different assumptions than DFT, and (ii) that the present approach employs a comprehensive set of experimental thermophysical data of the sodium chloride/water system, it is justified to assess the achieved agreement as very good⁸. The application of the employed here theory to experimental data allows the determination of interface parameters which are reconcilable with predictions from molecular theories. The subsequent calculations have been performed using the best-guess values of $\sigma_{\infty}^{(\alpha\gamma)}$ and $l^{(*)}$.

The efflorescence radius, i.e., the radius of a homogeneous solution droplet at the onset of crystallisation, was found to be $R(\text{ERH}) \approx 7.3 \text{ nm}$, corresponding to a molality of $m(\text{ERH}) \approx 15.6 \text{ mol kg}^{-1}$ and a solution saturation ratio of $S^{(\alpha)} \approx 2.53$ required for homogeneous crystallisation to occur. Figure 4 displays the dependence of the solute mass fraction as a function of the residual core radius at the efflorescence radius. A vanishing residual core, $R'_N \rightarrow 0$, corresponds to a homogenous solution droplet in

⁸It should be mentioned, of course, that the realisation of an adopted RH uncertainty of only 1 %rh is a very great experimental challenge.

which the solute mass fraction attains its maximum. According to Fig. 4, the solute mass fraction at efflorescence amounts $x_m \approx 0.48$.

Figure 5 displays the chemical potential of solvent molecules in the liquid film given in thermal units, $b^{(\beta \rightarrow \alpha)}$, as a function of the film thickness, h , according to Eq. (5).

The difference between the actual dimensionless chemical potential of the vapour, $b^{(\beta)} = \ln S^{(\beta)}$, and $b^{(\beta \rightarrow \alpha)}$ can be interpreted as a driving force of vapour condensation. It appears that in the initial stages of water uptake, when the film is still very thin, the values of $b^{(\beta \rightarrow \alpha)}$ are very low, i.e., the condensation driving force is very large. This effect originates from the contribution of the disjoining pressure. Without the consideration of the disjoining pressure, the water uptake by hygroscopic particles at RH well below DRH cannot be explained (see Kuni et al., 1996). Condensation occurs as long as the actual value of $b^{(\beta)}$ is larger than $b^{(\beta \rightarrow \alpha)}$. The maximum value of $b^{(\beta \rightarrow \alpha)}$ defines the DRH value. In the present case, deliquescence sets in at a film thickness of $h(\text{DRH}) \approx 2.5 \text{ nm}$ at $R_N = 5 \text{ nm}$.

The numerical solution of the OF Eq. (3) delivers the drop radius R as a function of the residual radius R'_N , which is depicted in Fig. 6 for $R_N = 5 \text{ nm}$. The maximum of the function $R(R'_N)$ represents the stability limit of the heterogeneous droplet, i.e., above this value a heterogeneous solution droplet cannot exist in thermodynamic equilibrium with a residual core. Below the maximum droplet radius and above the efflorescence radius, $R(\text{ERH})$, there exist two numerical solutions of the OF equation, i.e., the equilibrium droplet radius R is reconcilable with two different sizes of the residual core radius R'_N . The lower of the two horizontal solid lines depicted in Fig. 6 indicates the efflorescence radius, $R(\text{ERH})$, as determined from previous steps⁹.

Figure 7 shows the humidity growth factor GF as function of RH according to Eq. (4) for $R_N = 5 \text{ nm}$. The solid line refers to the homogeneous solution droplet with completely

⁹The solution of the OF equation (thick solid line in Fig. 6) is intrinsically not restricted to droplet sizes above the efflorescence radius. In the present approach, however, the efflorescence radius has been considered as an additional physical constraint imposing a lower limit on the physically accessible drop radius.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dissolved residual core, $R'_N = 0$, and the dotted curves refer to the heterogeneous solution droplet with partially dissolved residual core, $R'_N \neq 0$. Let us consider a dry particle which is placed in an atmosphere free of vapour at the beginning. Upon humidification by increasing RH the dry particle initially remains seemingly “inert” against hydration until about 40 %rh. However, the adsorbed film exist even in this stage as a very thin film in equilibrium with vapour. Close to this humidity value the dry particle starts to take up remarkable amounts of water molecules, as seen from the rapid increase of the growth factor; at this stage the film is still thermodynamically stable but already thicker. At the DRH, a liquid film forms without an activation barrier, the residual core fast vanishes, and upon further increase of the RH the growth factor evolves according to the solid line referring to the homogeneous droplet ¹⁰. Upon dehumidification of the ambient atmosphere the droplet shrinks. As the mass of the completely dissolved residual core remains conserved inside the droplet, the solute mass fraction in the droplet will increase until a solution supersaturation is achieved at which homogeneous crystallisation sets in. As described previously, the determination of the conditions of efflorescence relies on the employment of CNT and the use of several thermophysical parameters, among which the kinetic prefactor in the nucleation rate expression is subject of some uncertainties owing to limited availability of data on the energy barrier for interface-crossing diffusion. In the range $ERH \leq RH \leq DRH$ there exist two thermodynamically stable numerical solutions of the calculus; one refers to a heterogeneous, the other to a homogeneous droplet. For the heterogeneous droplet in turn there exist two branches corresponding to two different GF values (symbols “x” and “•”), but as shown by Shchekin and Shabaev (2010) they cannot coexist. The efflorescence transition from the homogeneous to a heterogeneous droplet corresponds to a jump from

¹⁰We emphasise that at all relative humidities below the DRH a thermodynamically stable film forms without an activation barrier. In these cases the stable equilibrium of the film corresponds to a deep minimum of the work of formation. In the vicinity of the DRH, the barrier for transition to a droplet with completely dissolved nucleus vanishes. Hence, one can say that at the DRH a homogeneous droplet forms without an activation barrier.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the solid line to the dotted line (symbol “•”). Figure 7 shows that the specification of the temperature and humidity is in fact a necessary but not sufficient condition for the full characterisation of the thermodynamic state of the system; the full characterisation requires also the specification of the “history” of the thermohygric conditions.

Figure 8 shows the dimensionless 2-D work of droplet formation, $\Delta F/(k_B T)$, as a function of drop radius R and the residual core radius R'_N according to Eq. (1) for at RH = 78 %rh, i.e., just below the deliquescence stage. One can clearly see the existence of two minima in the formation work, which are separated by a saddle point.

4.2 Sensitivity studies

4.2.1 Influence of the solution surface tension

The derivation of the calculus relies on the assumption that the vapour/solution surface tension and the solution/solute interfacial energy are independent of the solution concentration. In order to study the sensitivity of the model results against the special choice of the surface tension of the droplet, we have performed a comparative run with $\sigma_\infty^{(\alpha\beta)} = \sigma_W = 0.072 \text{ J m}^{-2}$, where σ_W is the surface tension of pure water (calculations have been performed for the nominal values of experimental ERH and DRH). For a solution droplet in which a NaCl core with radius $R_N = 5 \text{ nm}$ has been completely dissolved the efflorescence radius was found to be $R(\text{ERH}) \approx 7.4 \text{ nm}$ (slightly higher than its reference case (r.c.) value), corresponding to a molality of $m(\text{ERH}) \approx 15.2 \text{ mol kg}^{-1}$ (slightly lower than its r.c. value) and a solution saturation ratio of $S^{(\alpha)} \approx 2.47$ (slightly lower than the r.c. value) required for homogeneous crystallisation to occur. At efflorescence, the solute mass fraction amounts $x_m \approx 0.47$ (slightly lower than its r.c. value). For the best guess of the solution/solute interface energy a value of $\sigma_\infty^{(\alpha\gamma)} = 0.0866 \text{ J m}^{-2}$ has been found (slightly lower than its r.c. value), which results in a spreading coefficient of $s = 0.081 \text{ J m}^{-2}$ (higher than its r.c. value). For the correlation length the best guess value is $l^{(*)} \approx 5.4 \text{ \AA}$ (lower than its r.c. value). The

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



higher spreading coefficient and lower correlation length scale results in a higher value of the disjoining scaling pressure, and should promote the disjoining pressure effect in the initial stages of wetting, i.e., when the first condensed monolayers will form on the hygroscopic particle. The influence of $\sigma_{\infty}^{(\alpha\beta)}$ on the humidity growth factor can be seen from comparison of Figs. 7 and 9. The change of the droplet surface tension has seemingly a dramatic impact on the affinity of the hygroscopic particle to take up water. The dry particle starts to take up water already at about 20 %rh, which is much “earlier” than in the reference case. However, the influence of the solution surface tension on the growth factor of the homogeneous droplet is insignificant, hence the recognised sensitivity of the growth factor of the heterogeneous droplet is intrinsically an effect of the disjoining pressure.

4.2.2 Influence of the vapour/solid interface energy

Previous studies suggest a large uncertainty in estimates of the solid/vapour interface energy of NaCl, $\sigma_{\infty}^{(\beta\gamma)} \approx (0.1\text{--}0.39)\text{ J m}^{-2}$ (see SM/ Sect. S3.6). Figure 10 shows the calculated correlation length, spreading coefficient, and characteristic disjoining pressure, $l^{(*)}$, s , and $K^{(*)} = s/l^{(*)}$, respectively, as a function of $\sigma_{\infty}^{(\beta\gamma)}$, constraint by the condition $s > 0$. An increase of the surface energy of a solid enhances the affinity of that solid to take up water by both (i) promoting the spreading of the condensate on the surface, and by (ii) lowering the correlation length and enhancing the disjoining pressure. This result underpins the need for careful estimations of interface energies of hygroscopic matter for quantification of the water uptake. The present approach has been applied to the estimation of $\sigma_{\infty}^{(\alpha\gamma)}$ and $l^{(*)}$ from measured ERH and DRH values. The additional estimation of $\sigma_{\infty}^{(\beta\gamma)}$ would require an further governing condition for a well-measurable observable.

4.2.3 “Non-ideal” vs. “ideal” solution model

The original SSR approach was derived adopting the ideal-solution approximation of the activities (see footnote ⁶), i.e., the authors employed a linearisation of the solvent and solute activities that is valid for diluted solutions. In doing so, Shchekin, Shabaev and Rusanov predicted the first time theoretical curves for the humidity growth factor as function of the RH. However, for the special NaCl system considered here and with consideration of the constraints set by the employed thermophysical properties (see SM/Sect. S3) we could by no means get a numerical solution using the ideal-solution approximation of the activities, that is reconcilable with the observed DRH data of Biskos et al. (2006a). For example, replacing in the reference case run the “nonideal-ity” activity model with the ideal-solution approximation we found a solid/liquid interface energy of $\sigma_{\infty}^{(\alpha\gamma)} \approx 0.0593 \text{ J m}^{-2}$ for the observed ERH value (see Fig. 11). For the corresponding efflorescence radius we obtained $R(\text{ERH}) = 5.7 \text{ nm}$, which results in a very high value of the molality at the onset of efflorescence, $m(\text{ERH}) \approx 82 \text{ mol kg}^{-1}$. This value corresponds to an extremely high solution supersaturation and is far beyond an experimentally accessible state of the system. In contrast to the possibility to obtain a value for $\sigma_{\infty}^{(\alpha\gamma)}$ which formally fulfills the adopted constraints, it was impossible to decrease the difference between predicted and observed DRH values to a value smaller than 10 %rh by adjusting the correlation length (see Fig. 12).

However, compared to the ideal-solution limit employed by Shchekin, Shabaev, and Rusanov, the present result is essentially based on consideration of additional physical and physicochemical constraints: (i) employment of the CNT, (ii) using specific hygric, thermodynamic, and interfacial properties obtained from different experimental and modelling sources. Hence, the nonapplicability of the ideal-solution limit does not restrict the physical generality of the SSR approach. A possible candidate for enhancement of the “degrees of freedom” of the present approach is the solubility of the solute in the liquid film. However, as we do not have a profound physical justification

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

for changing the solubility, the “ideal solution” approximation was not further examined here.

5 Conclusions

1. In order to study the growth/shrinking of a hygroscopic nanoparticle during hydration/dehydration in a solvent vapour atmosphere, we have employed a thermodynamic theory proposed by Shchekin, Shabaev, and Rusanov which allows the prediction of the humidity growth factor of both (i) a homogeneous solution droplet with completely dissolved residual core, and (ii) a heterogeneous solution droplet with partially dissolved residual core as a function of the ambient relative humidity.
2. For application to a nanometric sodium chloride particle placed in a water vapour atmosphere, we have extended the theory as follows: (i) consideration of the non-ideality of the solution through the dependence of the molecular volumes of the solvent and solute molecules in the solution on the solution concentration, (ii) consideration of the nonideality of the solution in the determination of the solute and solvent activities, (iii) derivation of a CNT-based governing equation for the estimation of the efflorescence properties of a homogeneous solution droplet.
3. It was demonstrated how the solution/solute interface energy and the correlation length of a thin solution film can be determined from a combination of experimentally determinable efflorescence and deliquescence humidities with the present calculus. The solution/solute interface energy was found to be in close agreement with the value proposed by Gao et al. (2007); the correlation length was found to be about 1 nm, which is somewhat higher than previous estimated from the literature. Considering the combination of an extensive calculus, a comprehensive set of thermophysical constraints, and independent ERH/DRH measurements as function of dry particle size, the obtained values of both the solution/solute interface energy and the correlation length scale are surprisingly close to previous

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimations. This reflects a satisfying degree of “self-consistency” of the present approach allowing the retrieval of microscopic interfacial parameters from macroscopic humidity measurements. It also shows the crucial role that humidity metrology plays for the theoretical description of the water uptake by nanometric particles.

4. The humidification of sodium chloride particles in the initial hydration stages is very sensitive against the specification of the disjoining pressure. While the overall shape of the humidity growth factor dependence on humidity does not change, the onset of water uptake strongly depends on the disjoining pressure scale, which has been parameterised in terms of the spreading coefficient and the correlation length scale. The enhancement of the wettability of the particle surface leads to an earlier onset of hygroscopic growth. However, it remains to be investigated, how an explicit consideration of the dependence of the solution surface tension on the solution concentration in the theory will quantitatively affect the prediction of the humidity growth factor¹¹. Also a systematic sensitivity study using “alternative” formulations of the disjoining pressure remains to be done.
5. The calculations performed here give a theoretical explanation of the hysteresis effect during hydration/dehydration of a hygroscopic particle originating from the onset of the deliquescence and efflorescence transitions at different ambient humidities. Such hysteresis brings some “memory effect” into atmospheric phase transitions and was previously shown to affect aerosol-induced radiative forcing.
6. The retrieval of physically meaningful values of the solution/solute interface energy and the correlation length for a tiny NaCl particle with a dry particle radius of

¹¹It should be noted that the solution surface tension in the final equations cannot be a posteriori replaced by its concentration-dependent counterpart. Instead, the concentration-dependence of the surface tension must be already considered in the physical formulation of the chemical potentials and the energy balance itself. Moreover, the dependence of surface tension on concentration requires a consistent concentration dependent disjoining pressure.

5 nm from experimentally determined ERH/DRH values required the consideration of the “nonideality” of the film solution. The employment of the “ideal solution” limit of the solvent and solute activities did not result in physically sound numerical solutions of the calculus, regardless of the use of alternative, physically possible interfacial parameters (according to parameter uncertainties).

7. The application of the theory to supercooled states down to the eutectic temperature requires the availability of the corresponding thermophysical data (e.g., solvent/solute activities, mass density of the solution, interfacial parameters, viscosity). Sources of corresponding experimental data and formulations are the IAPWS pool of thermophysical properties (<http://www.iapws.org/>), the new Thermodynamic Equation Of Seawater-2010 (TEOS-10, <http://www.teos-10.org/>), and the FREZCHEM database (<http://www.dri.edu/frezchem>).

The Supplement related to this article is available online at doi:10.5194/acpd-14-22715-2014-supplement.

15 *Acknowledgements.* The contribution of O. Hellmuth was provided within the framework of the main research topic II “Impact of tropospheric aerosol on clouds and radiation budget” of the Leibniz Institute for Tropospheric Research (TROPOS), Leipzig. A. K. Shchekin thanks for St Petersburg State University Grants 11.0.59.2010 and 11.37.183.2014.

References

- 20 Biskos, G., Malinowski, A., Russell, L. M., Buseck, P. R., and Martin, S. T.: Nanosize effect on the deliquescence and the efflorescence of sodium chloride particles, *Aerosol Sci. Tech.*, 40, 97–106, doi:10.1080/02786820500484396, 2006a. 22716, 22734, 22741
- Biskos, G., Paulsen, D., Russell, L. M., Buseck, P. R., and Martin, S. T.: Prompt deliquescence and efflorescence of aerosol nanoparticles, *Atmos. Chem. Phys.*, 6, 4633–4642, doi:10.5194/acp-6-4633-2006, 2006b. 22721, 22735

22744

ACPD

14, 22715–22762, 2014

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bontjer, A.: Kochsalz. Chemie, Eigenschaften und Bedeutung, Vortrag, Phillips-Universität Marburg, Fachbereich Chemie, 12 January 2006, available at: http://www.chids.de/veranstaltungen/uebungen_experimentalvortrag.html (last access: 5 September 2014), presentation number 700, 2014. 22719
- 5 Bruzewicz, D. A., Checco, A., Ocko, B. M., Lewis, E. R., McGraw, R. L., and Schwartz, S. E.: Reversible uptake of water on NaCl nanoparticles at relative humidity below deliquescence point observed by noncontact environmental atomic force microscopy, *J. Chem. Phys.*, 134, 044702, doi:10.1063/1.3524195, 2011. 22722, 22724
- 10 Curry, J. A. and Khvorostyanov, V. I.: Assessment of some parameterizations of heterogeneous ice nucleation in cloud and climate models, *Atmos. Chem. Phys.*, 12, 1151–1172, doi:10.5194/acp-12-1151-2012, 2012. 22719
- Djikaev, Y. S.: Kinetics of fluctuational deliquescence, *J. Chem. Phys.*, 116, 9865–9874, doi:10.1063/1.1475743, 2002. 22723
- 15 Djikaev, Y. S., Bowles, R., and Reiss, H.: Role of constraints in the thermodynamics of heterogeneous condensation on solid soluble particles: failure of the capillarity approximation, *Physica A*, 298, 155–176, 2001a. 22723
- Djikaev, Y. S., Bowles, R., Reiss, H., Hämeri, K., Laaksonen, A., and Väkevä, M.: Theory of size dependent deliquescence of nanoparticles: relation to heterogeneous nucleation and comparison with experiments, *J. Phys. Chem. B*, 105, 7708–7722, doi:10.1021/jp010537e, 2001b. 22723
- 20 Gao, Y., Chen, S. B., and Yu, L. E.: Efflorescence relative humidity of airborne sodium chloride particles: a theoretical investigation, *Atmos. Environ.*, 41, 2019–2023, doi:10.1016/j.atmosenv.2006.12.014, 2007. 22730, 22732, 22736, 22742
- Hämeri, K., Laaksonen, A., Väkevä, M., and Suni, T.: Hygroscopic growth of ultrafine sodium chloride particles, *J. Geophys. Res.*, 106, 20749–20757, 2001. 22734
- 25 Khvorostyanov, V. I. and Curry, J. A.: A new theory of heterogeneous ice nucleation for application in cloud and climate models, *Geophys. Res. Lett.*, 27, 4081–4084, 2000. 22719
- Khvorostyanov, V. I. and Curry, J. A.: Thermodynamic theory of freezing and melting of water and aqueous solutions, *J. Phys. Chem. A*, 108, 11073–11085, doi:10.1021/jp048099, 2004a. 22719
- 30 Khvorostyanov, V. I. and Curry, J. A.: The theory of ice nucleation by heterogeneous freezing of deliquescent mixed CCN. Part I: Critical radius, energy, and nucleation rate, *J. Atmos. Sci.*, 61, 2676–2691, 2004b. 22719

**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Khvorostyanov, V. I. and Curry, J. A.: The theory of ice nucleation by heterogeneous freezing of deliquescent mixed CCN. Part II: Parcel model simulation, *J. Atmos. Sci.*, 62, 261–285, 2005. 22719

5 Khvorostyanov, V. I. and Curry, J. A.: Refinements to the Köhler's theory of aerosol equilibrium radii, size spectra, and droplet activation: effects of humidity and insoluble fraction, *J. Geophys. Res.*, 112, D05206, doi:10.1029/2006JD007672, 2007. 22719

Khvorostyanov, V. I. and Curry, J. A.: Critical humidities of homogeneous and heterogeneous ice nucleation: inferences from extended classical nucleation theory, *J. Geophys. Res.*, 114, D04307, doi:10.1029/2008JD011197, 2009. 22719

10 Khvorostyanov, V. I. and Curry, J. A.: Parameterization of homogeneous ice nucleation for cloud and climate models based on classical nucleation theory, *Atmos. Chem. Phys.*, 12, 9275–9302, doi:10.5194/acp-12-9275-2012, 2012. 22719

Khvorostyanov, V. I. and Sassen, K.: Cirrus cloud simulation using explicit microphysics and radiation. Part I: model description, *J. Atmos. Sci.*, 55, 1808–1821, 1998a. 22719

15 Khvorostyanov, V. I. and Sassen, K.: Toward the theory of homogeneous ice nucleation and its parameterization for cloud models, *Geophys. Res. Lett.*, 25, 3155–3158, 1998b. 22719

Khvorostyanov, V. I. and Sassen, K.: Microphysical processes in cirrus and their impact on radiation: a mesoscale modeling perspective, in: *Cirrus*, edited by Lynch, D. K., Sassen, K., Starr, D. O., and Stephens, G., Oxford University Press, Oxford, chap. 19, 397–432, 2002. 22719

20 Khvorostyanov, V. I., Curry, J. A., Pinto, J. O., Shupe, M., Baker, B. A., and Sassen, K.: Modeling with explicit spectral water and ice microphysics of a two-layer cloud system of altostratus and cirrus observed during FIRE Arctic Clouds Experiment, *J. Geophys. Res.*, 106, 15099–15112, 2001. 22719

25 Khvorostyanov, V. I., Curry, J. A., Gultepe, I., and Strawbridge, K.: A springtime cloud over the Beaufort Sea polynya: three-dimensional simulation with explicit spectral microphysics and comparison with observations, *J. Geophys. Res.*, 108, 4296, doi:10.1029/2001JD001489, 2003. 22719

30 Khvorostyanov, V. I., Morrison, H., Curry, J. A., Baumgardner, D., and Lawson, P.: High supersaturation and modes of ice nucleation in thin tropopause cirrus: simulation of the 13 July 2002 Cirrus Regional Study of Tropical Anvils and Cirrus Layers case, *J. Geophys. Res.*, 111, D02201, doi:10.1029/2004JD005235, 2006. 22719

**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kuni, F. M., Shchekin, A. K., Rusanov, A. I., and Widom, B.: Role of surface forces in heterogeneous nucleation on wettable nuclei, *Adv. Colloid Interfac.*, 65, 71–124, 1996. 22737
- Kuni, F. M., Shchekin, A. K., and Grinin, A. P.: Theory of heterogeneous nucleation for vapour undergoing a gradual metastable state formation process, *Physics Uspekhi*, 44, 331–370, 2001. 22723
- Li, B.: Density-Functional Theory and Quantum Chemistry Studies on “dry” and “wet” NaCl(001), Ph.D. thesis, Technical University of Berlin, Faculty of Mathematics and Natural Sciences, Berlin, 2009. 22719, 22732
- Liu, W. T., Tang, W., and Niler, P. P.: Humidity profiles over the ocean, *J. Climate*, 4, 1023–1034, 1991. 22720
- Marčelja, S. and Radić, N.: Repulsion of interfaces due to boundary water, *Chem. Phys. Lett.*, 42, 129–130, 1976. 22736
- McGraw, R. and Lewis, E.: Deliquescence and efflorescence of small particles, *J. Chem. Phys.*, 131, 194705, doi:10.1063/1.3251056, 2009. 22721, 22722
- Onasch, T. B., McGraw, R., and Imre, D.: Temperature-dependent heterogeneous efflorescence of mixed ammonium sulfate/calcium carbonate particles, *J. Phys. Chem. A*, 104, 10797–10806, doi:10.1021/jp0024064, 2000. 22724, 22730
- Pant, A., Parsons, M. T., and Bertram, A. K.: Crystallization of aqueous ammonium sulfate particles internally mixed with soot and kaolinite: crystallization relative humidities and nucleation rates, *J. Phys. Chem. A*, 110, 8701–8709, doi:10.1021/jp060985s, 2006. 22724, 22730
- Parsons, M. T., Riffell, J. L., and Bertram, A. K.: Crystallization of aqueous inorganic–malonic acid particles: nucleation rates, dependence on size, and dependence on the ammonium-to-sulfate ratio, *J. Phys. Chem. A*, 110, 8108–8115, doi:10.1021/jp057074n, 2006. 22724, 22730
- Romakkaniemi, S., Hämeri, K., Väkevä, M., and Laaksonen, A.: Adsorption of water on 8–15 nm NaCl and $(\text{NH}_4)_2\text{SO}_4$ aerosols measured using an ultrafine tandem differential mobility analyzer, *J. Phys. Chem. A*, 105 (35), 8183–8188, doi:10.1021/jp0106471, 2001. 22734
- Rusanov, A. I., Kuni, F. M., and Shchekin, A. K.: On the thermodynamics of thin wetting films, *Colloid J. Russ. Acad.*, 56, 172–173, 1994. 22726
- Russell, L. M. and Ming, Y.: Deliquescence of small particles, *J. Chem. Phys.*, 116, 311–321, doi:10.1063/1.1420727, 2002. 22732, 22736
- Schmiedel, H.: *Rund ums Salz – mit Tips und Rezepten*, VEB Fachbuchverlag Leipzig, Leipzig, 1984 22719

**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics. From Air Pollution to Climate Change, John Wiley & Sons Inc., New York, 2006. 22718, 22721, 22727
- Shchekin, A. K. and Rusanov, A. I.: Generalization of the Gibbs–Kelvin–Köhler and Ostwald–Freundlich equations for a liquid film on a soluble nanoparticle, *J. Chem. Phys.*, 129, 154116, doi:10.1063/1.2996590, 2008. 22723
- Shchekin, A. K. and Shabaev, I. V.: Thermodynamics and kinetics of deliquescence of small soluble particles, in: *Nucleation Theory and Applications*, edited by: Schmelzer, J. W. P., Röpke, G., and Priezhev, V. B., JINR Joint Institute for Nuclear Research, Bogoliubov Laboratory of Theoretical Physics, Dubna, 267–291, 2005. 22723
- Shchekin, A. K. and Shabaev, I. V.: Activation barriers for the complete dissolution of condensation nucleus and its reverse crystallization in droplets in the undersaturated solvent vapor, *Colloid J.*, 72, 432–439, doi:10.1134/S1061933X1003018X, 2010. 22724, 22738
- Shchekin, A. K., Rusanov, A. I., and Kuni, F. M.: Thermodynamics of condensation on soluble nuclei of colloidal surfactants, *Colloid J. Russ. Acad.*, 55, 227–234, 1993. 22723
- Shchekin, A. K., Shabaev, I. V., and Rusanov, A. I.: Thermodynamics of droplet formation around a soluble condensation nucleus in the atmosphere of a solvent vapor, *J. Chem. Phys.*, 129, 214111, doi:10.1063/1.3021078, 2008. 22723, 22726, 22728, 22729
- Shchekin, A. K., Shabaev, I. V., and Hellmuth, O.: Thermodynamic and kinetic theory of nucleation, deliquescence and efflorescence transitions in the ensemble of droplets on soluble particles, *J. Chem. Phys.*, 138, 054704, doi:10.1063/1.4789309, 2013. 22724
- Wagner, R. and Möhler, O.: Heterogeneous ice nucleation ability of crystalline sodium chloride dihydrate particles, *J. Geophys. Res.*, 118, 4610–4622, doi:10.1002/jgrd.50325, 2013. 22719
- Wang, J., Hoffmann, A. A., Park, R. J., Jacob, D. J., and Martin, S. T.: Global distribution of solid and aqueous sulfate aerosols: effect of the hysteresis of particle phase transitions, *J. Geophys. Res.*, 113, D11206, doi:10.1029/2007JD009367, 2008a. 22718
- Wang, J., Jacob, D. J., and Martin, S. T.: Sensitivity of sulfate direct climate forcing to the hysteresis of particle phase transitions, *J. Geophys. Res.*, 113, D11207, doi:10.1029/2007JD009368, 2008b. 22718
- Wise, M. E., Baustian, K. J., Koop, T., Freedman, M. A., Jensen, E. J., and Tolbert, M. A.: Depositional ice nucleation onto crystalline hydrated NaCl particles: a new mechanism for ice formation in the troposphere, *Atmos. Chem. Phys.*, 12, 1121–1134, doi:10.5194/acp-12-1121-2012, 2012. 22718, 22719

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Determination of the efflorescence radius $R(\text{ERH})$, the efflorescence molality $m(\text{ERH})$, the efflorescence mass fraction $x_m(\text{ERH})$, and the liquid/solid interface energy $\sigma_\infty^{(\alpha\gamma)}$ according to Eq. (7) for three different ΔERH values (see Fig. 2).

Graph	ΔERH %rh	$R(\text{ERH})$ nm	$m(\text{ERH})$ mol kg ⁻¹	$x_m(\text{ERH})$ 1	$\sigma_\infty^{(\alpha\gamma)}$ J m ⁻²
1	-1.0	7.28	15.95	0.4825	0.0894
2	0.0	7.32	15.58	0.4765	0.088
3	+1.0	7.36	15.20	0.4705	0.0866

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Determination of the correlation length $l^{(*)}$ according to Eq. (7) for six possible combinations of ΔERH and ΔDRH values (see Fig. 3). The spreading coefficient s is added.

Graph	ΔERH %rh	ΔDRH %rh	$l^{(*)}$ nm	s $J m^{-2}$
11	−1.0	−1.0	1.52	0.0676
21	0.0	−1.0	1.51	0.069
31	+1.0	−1.0	1.5	0.07
12	−1.0	0.0	0.98	0.0676
22	0.0	0.0	0.98	0.069
32	+1.0	0.0	1.05	0.07
13	−1.0	+1.0	0.7	0.0676
23	0.0	+1.0	0.7	0.069
33	+1.0	+1.0	0.7	0.07

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 1. Empirically derived power law fits of the size dependence of the deliquescence and efflorescence relative humidity values, DRH and ERH, of NaCl particles with dry mobility equivalent diameters in the range $6 < D_{m,dry}/\text{nm} < 60$ according to Eq. (9).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 2. Cost function $\mathcal{F}_{\text{cost}}^{(\sigma)}$ as a function of the solution/solute interfacial energy $\sigma_{\infty}^{(\alpha\gamma)}$ according to Eq. (7) for three different ERH values $\text{ERH} = \text{ERH}_0 + \Delta\text{ERH}$ with ERH_0 denoting the nominal value of ERH according to the power law fit given by Eq. (9) with ΔERH optionally assuming -1% rh, 0% rh, and $+1\%$ rh, corresponding to graphs 1, 2, and 3, respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin





Figure 3. Cost function $\mathcal{F}_{\text{cost}}^{(l)}$ as a function of the correlation length $l^{(*)}$ according to Eq. (8).

**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin**Figure 4.** Solute mass fraction x_m as a function of the residual radius R'_N .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 5. Chemical potential of solvent molecules in the liquid film given in thermal units, $b^{(\beta \rightarrow \alpha)}$, as a function of the film thickness, h , according to Eq. (5) at $R_N = 5$ nm.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Determination of
interfacial
parameters**O. Hellmuth and
A. K. Shchekin

Figure 6. Drop radius R as a function of residual radius R'_N as the numerical solution of the OF equation (3) at $R_N = 5$ nm.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 7. Humidity growth factor GF as function of RH according to Eq. (4) for $R_N = 5$ nm. The solid line represents the numerical solution of the GKK equation for the homogeneous solution droplet with completely dissolved residual core, $R'_N = 0$. The dotted curves (symbols “x” and “•”) represent the numerical solution of both the OF and GKK equations for the heterogeneous droplet with partially dissolved residual core, $R'_N \neq 0$.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 8. Dimensionless 2-D work of droplet formation, $\Delta F/(k_B T)$, as a function of drop radius R and the residual core radius R'_N according to Eq. (1) for at RH = 78 %rh.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 9. As in Fig. 7 but using the surface tension of pure water, $\sigma_{\infty}^{(\alpha\beta)} = 0.072 \text{ J m}^{-2}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 10. Calculated interfacial parameters as functions of $\sigma_{\infty}^{(\beta\gamma)}$ (with constraint $s > 0$). Top: spreading coefficient s and correlation length $l^{(*)}$. Bottom: characteristic disjoining pressure $K^{(*)} = s/l^{(*)}$.

Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 11. Cost function $\mathcal{F}_{\text{cost}}^{(\sigma)}$ as a function of the solution/solute interfacial energy $\sigma_{\infty}^{(\alpha\gamma)}$ according to Eq. (7) employing the “ideal solution” limit of the activity model for two different $\sigma_{\infty}^{(\alpha\beta)}$ values. Graph 1 using $\sigma_{\infty}^{(\alpha\beta)} = 0.083 \text{ J m}^{-2}$, graph 2 using $\sigma_{\infty}^{(\alpha\beta)} = 0.072 \text{ J m}^{-2}$.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Determination of interfacial parameters

O. Hellmuth and
A. K. Shchekin




Figure 12. Cost function $\mathcal{F}_{\text{cost}}^{(l)}$ as a function of the correlation length $l^{(*)}$ according to Eq. (8), employing the “ideal solution” limit of the activity model for two different $\sigma_{\infty}^{(\alpha\beta)}$ values. Graph 1 using $\sigma_{\infty}^{(\alpha\beta)} = 0.083 \text{ J m}^{-2}$, graph 2 using $\sigma_{\infty}^{(\alpha\beta)} = 0.072 \text{ J m}^{-2}$.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
