

1 **Effects of molecular weight and temperature on liquid-liquid phase separation in particles**  
2 **containing organic species and inorganic salts**

3 Yuan You and Allan K. Bertram<sup>1</sup>

4 Department of Chemistry, University of British Columbia, 2036 Main Mall, Vancouver, BC,  
5 V6T 1Z1, Canada

6 <sup>1</sup>To whom correspondence should be addressed:

7 Email: bertram@chem.ubc.ca

8 **Abstract**

9 Atmospheric particles containing organic species and inorganic salts may undergo liquid-liquid  
10 phase separation when the relative humidity varies between high and low values. To better  
11 understand the parameters that affect liquid-liquid phase separation in atmospheric particles, we  
12 studied the effects of molecular weight and temperature on liquid-liquid phase separation in  
13 particles containing one organic species mixed with either ammonium sulfate or ammonium  
14 bisulfate. In the molecular weight dependent studies, we measured liquid-liquid phase separation  
15 relative humidity (SRH) in particles containing ammonium sulfate and organic species with large  
16 molecular weights (up to 1153 Da). These results were combined with recent studies of liquid-  
17 liquid phase separation in the literature to assess if molecular weight is a useful parameter for  
18 predicting SRH. The combined results, which include results from 33 different particle types,  
19 illustrate that SRH does not depend strongly on molecular weight (i.e. a clear relationship  
20 between molecular weight and SRH was not observed). In the temperature dependent studies, we  
21 measured liquid-liquid phase separation in particles containing ammonium sulfate mixed with 20  
22 different organic species at  $244 \pm 1$  K,  $263 \pm 1$  K, and  $278 \pm 1$  K, as well as  $290 \pm 1$  K for a few  
23 of these particle types. These new results were combined with previous measurements of the  
24 same particle types at  $290 \pm 1$  K. The combined SRH data illustrate that for the organic-  
25 ammonium sulfate particles studied the SRH does not depend strongly on temperature. At most  
26 the SRH varied by 9.7 % as the temperature varied from 290 to 244 K. The high SRH values (>  
27 65%) in these experiments may explain the lack of temperature dependence. Since water is a  
28 plasticizer, high relative humidities can lead to high water contents, low viscosities and high  
29 diffusion rates in the particles. For these cases, unless the temperature is very low, liquid-liquid  
30 phase separation is not expected to be kinetically inhibited. The occurrence of liquid-liquid phase  
31 separation and SRH did depend strongly on temperature over the range of 290-244 K for  
32 particles containing  $\alpha$ ,4-dihydroxy-3-methoxybenzeneacetic acid mixed with ammonium  
33 bisulfate. For this particle type, a combination of low temperatures and low water content likely  
34 favored kinetic inhibition of the liquid-liquid phase separation by slow diffusion rates in highly  
35 viscous particles. The combined results suggest that liquid-liquid phase separation is likely a  
36 common occurrence in atmospheric particles at temperatures from 244-290 K, although particles  
37 that do not undergo liquid-liquid phase separation are also likely common.

## 38 **1 Introduction**

39 A large fraction of particles in the atmosphere contains both organic species and inorganic salts  
40 (Murphy and Thomson, 1997; Murphy et al., 1998; Middlebrook et al., 1998; Buzorius et al.,  
41 2002; Murphy, 2005; Murphy et al., 2006; Tolocka et al., 2005; Chen et al., 2009; Pratt and  
42 Prather, 2010). Ammonium sulfate and ammonium bisulfate are both common inorganic salts in  
43 these particles (Dibb et al., 1996; Huebert et al., 1998; Talbot et al., 1998; Dibb et al., 2000; Lee  
44 et al., 2003). The types of organic species in these particles are numerous with only around 10%  
45 by mass of these organic species identified at the molecular level (Hamilton et al., 2004;  
46 Goldstein and Galbally, 2007; Decesari et al., 2006; Hallquist et al., 2009). Organic functional  
47 groups found in mixed organic-inorganic salt particles include carboxylic acids, alcohols,  
48 oxidized aromatic compounds, ethers, and esters (Finlayson-Pitts and Pitts, 2000; Seinfeld and  
49 Pandis, 2006; Decesari et al., 2006; Hallquist et al., 2009; Takahama et al., 2011; Rogge et al.,  
50 1993; Saxena and Hildemann, 1996; Finlayson-Pitts and Pitts, 1997; Day et al., 2009; Gilardoni  
51 et al., 2009; Liu et al., 2009; Russell et al., 2009; Russell et al., 2011; Fu et al., 2011; Fuzzi et al.,  
52 2001). The molecular weight of organic molecules in these mixed particles is thought to range  
53 from less than 100 Da to as high as 1000 Da (Gao et al., 2004; Tolocka et al., 2004; Kalberer et  
54 al., 2004; Nguyen et al., 2010). In addition, the average oxygen-to-carbon elemental ratios (O:C)  
55 of the organic material in ambient particles generally range from 0.1 to 1.0 (Chen et al., 2009;  
56 Aiken et al., 2008; DeCarlo et al., 2008; Jimenez et al., 2009; Hawkins et al., 2010; Heald et al.,  
57 2010; Ng et al., 2010; Takahama et al., 2011), while the average organic-to-inorganic mass ratios  
58 (OIR) of atmospheric particles generally range from 0.2 to 3.5 (Chen et al., 2009; Zhang et al.,  
59 2007; Jimenez et al., 2009).

60 When the relative humidity varies in the atmosphere, mixed organic-inorganic salt particles can  
61 undergo different phase transitions, including efflorescence and deliquescence (Martin, 2000;  
62 Brooks et al., 2002; Choi and Chan, 2002; Brooks et al., 2003; Chan and Chan, 2003; Wise et al.,  
63 2003; Braban and Abbatt, 2004; Pant et al., 2004; Parsons et al., 2004a; Badger et al., 2006;  
64 Chang and Pankow, 2006; Erdakos et al., 2006; Parsons et al., 2006; Salcedo, 2006; Ling and  
65 Chan, 2008; Treuel et al., 2009; Bodsworth et al., 2010). More recent laboratory measurements  
66 and calculations have focused on liquid-liquid phase separation (Clegg et al., 2001; Pankow,  
67 2003; Marcolli and Krieger, 2006; Anttila et al., 2007; Buajjarern et al., 2007; Zuend et al., 2008;  
68 Ciobanu et al., 2009; Kwamena et al., 2010; Prisle et al., 2010; Zuend et al., 2010; Bertram et al.,  
69 2011; Reid et al., 2011; Smith et al., 2011; Zuend et al., 2011; Krieger et al., 2012; Pöhlker et al.,  
70 2012; Smith et al., 2012; Song et al., 2012a, b; You et al., 2012; Zuend and Seinfeld, 2012;  
71 Drozd et al., 2013; Shiraiwa et al., 2013; Song et al., 2013; Veghte et al., 2013; You et al., 2013;  
72 Zuend and Seinfeld, 2013; Veghte et al., 2014). Liquid-liquid phase separation in the  
73 atmospheric particles can influence the partitioning of organic molecules between the gas and the  
74 particle phase (Seinfeld et al., 2001; Chang and Pankow, 2006; Zuend et al., 2010; Shiraiwa et  
75 al., 2013) and influence the reactive uptake of important gas-phase molecules, such as N<sub>2</sub>O<sub>5</sub>, into  
76 atmospheric particles (Anttila et al., 2006; Folkers et al., 2003; Escoreia et al., 2010; Cosman and

77 Bertram, 2008; Park et al., 2007; Thornton and Abbatt, 2005; McNeill et al., 2006; Gaston et al.,  
78 2014). Liquid-liquid phase separation can also influence the deliquescence and efflorescence  
79 relative humidity in mixed organic-inorganic salt particles (Bertram et al., 2011; Smith et al.,  
80 2012; Song et al., 2013). In addition, liquid-liquid phase separation can alter the ice nucleation  
81 properties of particles (Schill and Tolbert, 2013).

82 Recent work has investigated different parameters that influence liquid-liquid phase separation in  
83 particles containing mixtures of organic species and inorganic salts (You et al., 2014).  
84 Understanding the parameters that affect these transitions is necessary for predicting these phase  
85 transitions in atmospheric particles. Studies have shown that O:C is an important parameter for  
86 predicting liquid-liquid phase separation in these particles, with this phase transition always  
87 observed for O:C less than 0.5, frequently observed for O:C values between 0.5 and 0.8, and  
88 never observed for O:C values greater than 0.8 (Bertram et al., 2011; Song et al., 2012a, b; You  
89 et al., 2013). Measurements have also illustrated that the type of functional groups can also  
90 influence the relative humidity required for liquid-liquid phase separation when the O:C of the  
91 organic material is in the range of approximately 0.5 to 0.8 (Song et al., 2012b). On the other  
92 hand, the occurrence of liquid-liquid phase separation is not a strong function of the OIR or the  
93 number of organic species (Marcolli and Krieger, 2006; Ciobanu et al., 2009; Bertram et al.,  
94 2011; Song et al., 2012a, b; Schill and Tolbert, 2013). In addition, the occurrence of liquid-liquid  
95 phase separation may not be a strong function of the type of inorganic salt for  $O:C \geq 0.8$  and  $\leq$   
96 0.5, but, in the range of 0.5 to 0.8, the occurrence of liquid-liquid phase separation can depend on  
97 the salt type (You et al. 2013).

98 In the following paper, we investigate if molecular weight of the organic material and  
99 temperature of the particles influence liquid-liquid phase separation in particles containing  
100 organic species and inorganic salts. Molecular weight of the organic molecules is of interest  
101 because viscosity of organic-inorganic salt mixtures can depend roughly on the molecular weight  
102 of the organic species (Greener et al., 1993; Zobrist et al., 2008; Koop et al., 2011; Saukko et al.,  
103 2012), and at high molecular weights, liquid-liquid phase separation may become kinetically  
104 inhibited in highly viscous solutions. The effect of molecular weight of the organic material on  
105 the occurrence of liquid-liquid phase separation in mixed organic-inorganic salt particles has not  
106 been explored.

107 Temperature is of interest since temperature ranges from approximately 220 to 300 K in the  
108 troposphere, and temperature can influence the thermodynamics and kinetics of liquid-liquid  
109 phase separation. As an example of the effect of temperature on the thermodynamics of liquid-  
110 liquid phase transitions consider the binary mixture of water and butanol. At room temperature  
111 the binary mixture is partially immiscible. But as the temperature increases the region of  
112 immiscibility decreases until the upper critical solution temperature is reached. At higher  
113 temperatures the binary mixture is completely miscible. Certain mixtures can also have lower  
114 critical solution temperatures due to an increase in hydrogen bonding as the solution temperature  
115 decreases (Levine. 2009). Temperature may also influence the kinetics of liquid-liquid phase

116 transitions. Some mixtures of organic species and inorganic salts can become highly viscous at  
117 lower temperatures and low relative humidities (Tong et al., 2011; Zobrist et al., 2008; Koop et  
118 al., 2011; Zobrist et al., 2011; Murray, 2008; Mikhailov et al., 2009; Saukko et al., 2012). At  
119 these low temperatures and relative humidities, liquid-liquid phase separation may be kinetically  
120 inhibited due to diffusion limitations. Only two studies have investigated liquid-liquid phase  
121 separation in mixed organic-inorganic salt particles at temperatures below 290 K. Bertram et al.  
122 reported that SRH results were similar at 273 K and 290 K for particles containing ammonium  
123 sulfate and 1,2,6-hexanetriol (Bertram et al., 2011). Schill and Tolbert reported that SRH results  
124 were similar for temperatures from 240 to 265 K for particles containing ammonium sulfate  
125 mixed with 1,2,6-hexanetriol and particles containing ammonium sulfate, 1,2,6-hexanetriol, and  
126 2,2,6,6-tetrakis(hydroxymethyl)cyclohexanol (Schill and Tolbert, 2013).

127 Here we carried out a systematic study of the effect of molecular weight of the organic material  
128 on liquid-liquid phase separation at  $290 \pm 1$  K. We first studied liquid-liquid phase separation in  
129 particles containing ammonium sulfate mixed with one of ten organic species, with molecular  
130 weights up to 1153 Da. The data from these studies were then combined with recent studies of  
131 liquid-liquid phase separation in particles reported in the literature (You et al., 2013) to assess if  
132 molecular weight is a useful parameter to predict the occurrence of liquid-liquid phase separation  
133 and the liquid-liquid phase separation relative humidity (SRH).

134 To gain a better understanding of the effect of temperature on liquid-liquid phase separation in  
135 mixed organic-inorganic salt particles, we investigated liquid-liquid phase separation in particles  
136 containing ammonium sulfate mixed with one of twenty organic species at  $244 \pm 1$  K,  $263 \pm 1$  K,  
137 and  $278 \pm 1$  K, respectively. Some of these particle types were also studied at  $290 \pm 1$  K. These  
138 new data were combined with previous measurements of liquid-liquid phase separation at  $290 \pm$   
139  $1$  K by You et al. (You et al., 2013) to assess the effect of temperature on liquid-liquid phase  
140 separation in mixed organic-ammonium sulfate particles.

141 We also investigated liquid-liquid phase separation in particles containing ammonium bisulfate  
142 mixed with  $\alpha$ ,4-dihydroxy-3-methoxybenzeneacetic acid at temperatures of  $244 \pm 1$  K,  $263 \pm 1$   
143 K, and  $278 \pm 1$  K. This system was studied since it has a relatively low SRH-value (38%) at  $290$   
144  $\pm 1$  K and, hence, has relatively low water content at the onset of liquid-liquid phase separation.  
145 In addition to temperature and molecular weight, water content is likely important for the  
146 kinetics of liquid-liquid phase separations, as the viscosity and diffusion rates within the particles  
147 are expected to be sensitive to water content (Koop et al. 2011; Renbaum-Wolff et al. 2013;  
148 Power et al. 2013; Kidd et al. 2014).

## 149 **2 Experimental**

### 150 **2.1 Sample preparation and apparatus.**

151 Solutions of ammonium sulfate or ammonium bisulfate and one organic species were prepared in  
152 high purity water (Millipore,  $18.2 \text{ M}\Omega \text{ cm}$ ) with OIR of  $2.0 \pm 0.1$ . The solutions were then

153 nebulized to produce submicron particles, which impacted onto a hydrophobic glass slide  
154 (Hampton Research) and coagulated into super-micron droplets. Water was then evaporated to  
155 generate mixed organic-inorganic salt particles with lateral dimensions ranging from 10 to 35  
156  $\mu\text{m}$ .

157 The glass slide supporting the mixed organic-inorganic salt particles was mounted to a  
158 temperature and relative humidity controlled flow cell, which was coupled to an optical  
159 reflectance microscope (Zeiss AxioTech; 50 $\times$  objective) (Koop et al., 2000; Parsons et al., 2004b;  
160 Pant et al., 2006; Bodsworth et al., 2010). To control the relative humidity in the flow cell, dry  
161 and humidified nitrogen gas flows were combined and continuously passed through the cell. The  
162 total flow rate was approximately 1.5 L  $\text{min}^{-1}$ . While the RH was decreased, liquid-liquid phase  
163 separation was identified by monitoring the change in morphology of the particles. The relative  
164 humidity of the gas was determined with a chilled mirror hygrometer (General Eastern, Model  
165 1311DR). The hygrometer was calibrated prior to experiments at each temperature ( $244 \pm 1$  K,  
166  $263 \pm 1$  K,  $278 \pm 1$  K, and  $290 \pm 1$  K) by measuring the deliquescence relative humidity (DRH)  
167 of ammonium sulfate particles, and comparing the measured DRH value to the DRH valued  
168 predicted with the Extended Aerosol Inorganic Model (E-AIM model) (Clegg et al. 1998). The  
169 uncertainty ( $2\sigma$ ) of the hygrometer was  $\pm 2.5$  % RH after calibration based on reproducibility of  
170 the DRH measurements.

## 171 **2.2 Molecular weight dependent studies.**

172 Particles containing ammonium sulfate mixed with one of ten organic species were studied at  
173  $290 \pm 1$  K (see Table 1). Most of previous laboratory studies of liquid-liquid phase separation in  
174 particles containing organic species mixed with ammonium sulfate used organic species with  
175 molecular weight less than 200 Da (Bertram et al., 2011; Song et al., 2012a, b; You et al., 2013).  
176 To complement these previous studies, in the current study we investigated particles containing  
177 ammonium sulfate and organic species with molecular weight ranging from 180 to 1153 Da (see  
178 Table 1). The specific organic species selected for these studies (Table 1) also had a relatively  
179 wide range of O:C values. The glass transition temperatures of the organics studied covered at  
180 least the range of 223 K to 412 K (see Table 1). In all the studies, the organic-to-inorganic mass  
181 ratio (OIR) was  $2.0 \pm 0.1$ . This value was chosen so that the current studies could be compared  
182 with the previous studies by You et al. (2013), who also used an OIR =  $2.0 \pm 0.1$ . In addition,  
183 this OIR value is in the range of OIR values observed in many field studies (Zhang et al., 2007;  
184 Jimenez et al., 2009).

185  
186 In a typical experiment, the RH in the cell was ramped down at a rate of 0.4-0.6% RH  $\text{min}^{-1}$ ,  
187 while the temperature of the cell was held at  $290 \pm 1$  K. At the same time, images of the particles  
188 were captured continuously until one of the following conditions occurred: liquid-liquid phase  
189 separation was observed, the particles effloresced, or the RH reached  $\leq 0.5\%$ . For each type of  
190 particle, experiments were repeated at least three times. All of the organic species were

191 purchased from Sigma-Aldrich and had purities  $\geq 95\%$ , except for maltoheptaose, which had a  
192 purity  $\geq 90\%$ .

193

### 194 **2.3 Temperature dependent studies.**

195 Particles consisting of ammonium sulfate mixed with one of twenty organic species were studied  
196 (see Table 2) at  $244 \pm 1$  K,  $263 \pm 1$  K, and  $278 \pm 1$  K, as well as  $290 \pm 1$  K for a few of these  
197 particle types. For these studies, we chose the same particle types previously studied by You et  
198 al. (2013) at  $290 \pm 1$  K. These particle types cover a wide range of O:C values and included  
199 functional groups found in atmospheric particles. Also, ammonium sulfate is one of the most  
200 common inorganic salts found in the atmosphere. In addition to picking the particles types  
201 previously studied by You et al. (2013), we also studied two additional organic species mixed  
202 with ammonium sulfate. These two organic species were raffinose and poly(ethylene glycol)  
203 diacrylate. These two species were chosen since they had relatively high molecular weights. An  
204 OIR value of  $2.0 \pm 0.1$  was used in all the studies, consistent with You et al. (2013). The glass  
205 transition temperatures of the organic species studied cover the range of at least 192 K to 396 K.

206 In addition to studying particles consisting of ammonium sulfate mixed with one of twenty  
207 organic species, we also studied particles containing ammonium bisulfate mixed with  $\alpha,4$ -  
208 dihydroxy-3-methoxybenzeneacetic acid at temperatures of  $244 \pm 1$  K,  $263 \pm 1$  K, and  $278 \pm 1$   
209 K. As mentioned in the introduction, this system was studied since it has a low SRH-value (38%)  
210 at  $290 \pm 1$  K and hence has relatively low water content at the onset of liquid-liquid phase  
211 separation.

212 The organic species studied were purchased from Sigma-Aldrich with purities  $\geq 98\%$ . All  
213 organics were used without further purification. In a typical temperature dependent experiment,  
214 the RH was ramped down at a rate of around  $0.1$ -  $0.5$  % RH  $\text{min}^{-1}$ . Images were recorded in the  
215 same way as the molecular weight dependent studies.

## 216 **3 Results and Discussion**

### 217 **3.1 Effect of molecular weight on liquid-liquid phase separation**

218 Particles containing ammonium sulfate mixed with one of ten organic species were studied at  
219  $290 \pm 1$  K. In these studies organic species with large molecular weights (180 to 1153 Da) were  
220 used. The organic-to-inorganic mass ratio (OIR) was  $2.0 \pm 0.1$  in all the studies. The results from  
221 these studies are listed in Table 3 as well as the data from You et al., who also studied particles  
222 with an OIR of  $2.0 \pm 0.1$  at  $290 \pm 1$  K (You et al., 2013). The combined data set in Table 3,  
223 which includes results of liquid-liquid phase separation for 33 different particle types, was used  
224 to determine the importance of molecular weight of the organic species on liquid-liquid phase  
225 separation.

226 In Figure 1, the data from Table 3 were plotted as a function of O:C and molecular weight. Open  
227 circles indicate liquid-liquid phase separation was observed and stars indicate liquid-liquid phase  
228 separation was not observed. No clear relationship between molecular weight and the occurrence  
229 of liquid-liquid phase separation was observed, however a relationship between occurrence of  
230 liquid-liquid phase separation and O:C was clear: liquid-liquid phase separation was always  
231 observed when  $O:C < 0.57$  (orange hatched region), was never observed when  $O:C > 0.83$  (green  
232 hatched region), and was frequently observed when O:C ranged from 0.57 to 0.83. This range is  
233 slightly different than the range previously measured (0.5 to 0.8) (Bertram et al. 2011; Song et al.  
234 2012a, b; You et al. 2013), which is not surprising since this range is expected to change slightly  
235 with the type of organic molecules studied. As such, there is a small uncertainty in this range. In  
236 Figure 2, the SRH data from Table 3 were plotted as a function of molecular weight of the  
237 organic species. The colors of the symbols indicate the O:C of organic species in the mixed  
238 particles. Data at SRH = 0 % indicate liquid-liquid phase separation was not observed even at the  
239 lowest relative humidity studied. Similar to Figure 1, no correlation with molecular weight was  
240 apparent. For contrast, in Figure 3, we show the same SRH data plotted as a function of O:C of  
241 organic species with the color of the symbols representing the molecular weight of organic  
242 species. Consistent with previous results (Bertram et al., 2011; Song et al., 2012b), a correlation  
243 between SRH and O:C is apparent. These results suggest that O:C is more important for  
244 predicting the occurrence of liquid-liquid phase separation in atmospherically relevant mixed  
245 organic-ammonium sulfate particles compared with molecular weight.

### 246 **3.2 Effect of temperature on liquid-liquid phase separation**

247 In the temperature dependent experiments, particles containing ammonium sulfate mixed with  
248 one of twenty organic species were studied at  $244 \pm 1$  K,  $263 \pm 1$  K, and  $278 \pm 1$  K. Some of the  
249 particle types were also studied at  $290 \pm 1$  K. The temperature dependent results are included in  
250 Table 4 as well as results from You et al., who studied most of the same types of particles but  
251 only at  $290 \pm 1$  K (You et al., 2013). The combined SRH data from Table 4, which cover the  
252 temperature range of 290-244 K, were plotted in Figure 4A as a function of O:C of the organic  
253 species. The temperature dependent results show that for all the particle types studied and at all  
254 the temperatures studied liquid-liquid phase separation was always observed when  $O:C < 0.57$ ,  
255 frequently observed when  $0.57 \leq O:C < 0.8$ , and never observed when  $O:C \geq 0.8$ . We conclude  
256 that the O:C range at which liquid-liquid phase separation was observed at 290-293 K in  
257 previous studies (Ciobanu et al., 2009; Bertram et al., 2011; Song et al., 2012b; You et al., 2013),  
258 is reasonably consistent with the range observed at temperatures down to  $244 \pm 1$  K.

259 Figure 5 shows the same data as in Figure 4A, but displayed in a slightly different way. The SRH  
260 results for the twelve types of mixed organic-ammonium sulfate particles that underwent liquid-  
261 liquid phase separation in Figure 4A, are shown as a function of temperature, with the colors of  
262 the symbols representing the O:C values of the organic species in the particles. For all the  
263 particle types included in Figure 5, the SRH varied by less than 9.7 % RH as the temperature  
264 varied from 244 to 290 K. Figure 4A and 5 illustrate that SRH is not a strong function of

265 temperature for the particle types investigated. These results are consistent with earlier studies by  
266 Bertram et al. and Schill and Tolbert discussed in the Introduction. In the Supplementary  
267 Material we carried out a linear regression analysis to determine the level of significance of the  
268 temperature dependent trends observed in Figure 5. In short, five out of the twelve systems  
269 studied had a high correlation coefficient ( $r \geq 0.94$ ) between temperature and SRH and an  
270 associated low p-value ( $\leq 0.06$ ). A decrease in SRH with a decrease in temperature may be  
271 expected if there is a closed loop miscibility gap and the measurements are probing the lower  
272 temperature region of the closed loop.

273 Based on our results, particles containing poly(propylene glycol) and ammonium sulfate had  
274 SRH values ranging from 87.7 to 94.1 % as the temperature ranged from 244 to 290 K. Solutions  
275 of water and poly(propylene glycol) show a lower critical solution temperature of approximately  
276 50 °C (Malcom and Rowlinson, 1957). A comparison between this result and our current results  
277 indicate that the lower critical solution temperature is decreased significantly when ammonium  
278 sulfate is added to mixtures of poly(propylene glycol) and water.

279 As mentioned in the introduction, some mixtures of organic species and inorganic salts can  
280 become highly viscous at lower temperatures and low relative humidities, and at these low  
281 temperatures and relative humidities, liquid-liquid phase separation may be kinetically inhibited  
282 due to diffusion limitations. A possible reason that a stronger dependence on temperature was  
283 not observed in the studies discussed above may be because for the systems studied the particles  
284 either didn't undergo liquid-liquid phase separation or the SRH was relatively high ( $> 65\%$  RH).  
285 For the cases where liquid-liquid phase separation did not occur, an increase in viscosity from a  
286 decrease in temperature is not expected to change the results. For cases where the SRH was  
287 relatively high ( $> 65\%$  RH), the water content in the particles was also likely relatively high.  
288 Since water is a plasticizer, high water contents can lead to low viscosities and high diffusion  
289 rates. In this case, unless the temperature is very low, liquid-liquid phase separation is not  
290 expected to be kinetically inhibited.

291 Although SRH does not appear to be a strong function of the temperature for the organic-  
292 ammonium sulfate particle types studied over the temperature range of 244 to 290 K, SRH may  
293 be a strong function of temperature for these particles at temperatures lower than 244 K due to  
294 kinetic limitations or thermodynamic reasons. Additional studies of SRH at temperatures lower  
295 than 244 K are still needed for these particle types.

296 In Figure 6, SRH as a function of temperature is plotted for particles containing  $\alpha$ ,4-dihydroxy-3-  
297 methoxybenzeneacetic acid mixed with ammonium bisulfate. At  $263 \pm 1$  K and warmer no  
298 strong dependence on temperature is observed, with an average SRH value of approximately  
299 37.6 %. At 244 K, however, liquid-liquid phase separation was not observed. A likely  
300 explanation for this drastic decrease in the SRH is likely kinetic inhibition of the liquid-liquid  
301 phase separation. Related, previous research has shown that efflorescence is inhibited in particles  
302 containing organic species and inorganic salts at low temperatures and low relative humidities



303 likely due to diffusion limitations (Bodsworth et al., 2010). The studies with particles containing  
304 ammonium bisulfate and  $\alpha$ ,4-dihydroxy-3-methoxybenzeneacetic acid suggests that a  
305 combination of low temperatures (e.g. 244 K) and low relative humidities (e.g. approximately  
306 38% RH) can lead to kinetic inhibition of liquid-liquid phase separation.

307 The average O:C of organic material in atmospheric particles has been measured at many  
308 locations in the Northern Hemisphere and in the Amazon and has been shown to range from 0.1  
309 to 1.0 (Chen et al., 2009; Aiken et al., 2008; DeCarlo et al., 2008; Jimenez et al., 2009; Hawkins  
310 et al., 2010; Heald et al., 2010; Ng et al., 2010; Takahama et al., 2011). This range of O:C values  
311 is indicated in Figure 4B. The range of average O:C values measured in the atmosphere overlaps  
312 with the range of O:C values where liquid-liquid phase separation was observed at temperatures  
313 ranging from 244 to 290 K (Figure 4 B). This overlap suggests that liquid-liquid phase separation  
314 is likely a common occurrence in the atmosphere over this temperature range. Particles that do  
315 not undergo liquid-liquid phase separation are also expected to be common based on a  
316 comparison between our data and the range of O:C values found in the atmosphere.

#### 317 **4 Conclusions**

318 The occurrence of liquid-liquid phase separation and SRH did not depend strongly on the  
319 molecular weight of the organic species at  $290 \pm 1$  K, at least for the particle types studied. The  
320 occurrence of liquid-liquid phase separation and SRH also did not depend strongly on  
321 temperature over the range of 290- 244 K for particles containing ammonium sulfate mixed with  
322 one organic species. The SRH varied by at most 9.7 % RH as the temperature varied from  $290 \pm$   
323  $1$  K to  $244 \pm 1$  K for the particle types studied. The high SRH values ( $> 65\%$ ) in these  
324 experiments may explain the lack of temperature dependence. Since water is a plasticizer, high  
325 relative humidities can lead to high water contents, low viscosities and high diffusion rates. For  
326 these cases, unless the temperature is very low, liquid-liquid phase separation is not expected to  
327 be kinetically inhibited. The occurrence of liquid-liquid phase separation and SRH did depend  
328 strongly on temperature over the range of 290-244 K for particles containing  $\alpha$ ,4-dihydroxy-3-  
329 methoxybenzeneacetic acid mixed with ammonium bisulfate. For this particle type, a  
330 combination of low temperatures and low water content likely favored kinetic inhibition of the  
331 liquid-liquid phase separation by slow diffusion rates in highly viscous particles.

332 The combined results suggest that liquid-liquid phase separation is likely a common occurrence  
333 in the atmospheric particles at 290-244 K, although particles that do not undergo liquid-liquid  
334 phase separation are also likely common.

335

336 **Acknowledgements**

337 This research was supported by the Natural Science and Engineering Research Council of  
338 Canada. We thank two anonymous referees for their very helpful suggestions.

339

340 **References**

- 341 Aiken, A. C., Decarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M.,  
342 Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J.,  
343 Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prévôt, A. S. H., Dommen, J., Duplissy, J., Metzger,  
344 A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient  
345 organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*,  
346 42, 4478-4485, doi:10.1021/es703009q, 2008.
- 347 Anttila, T., Kiendler-Scharr, A., Tillmann, R., and Mentel, T. F.: On the reactive uptake of gaseous  
348 compounds by organic-coated aqueous aerosols: Theoretical analysis and application to the  
349 heterogeneous hydrolysis of N<sub>2</sub>O<sub>5</sub>, *J. Phys. Chem. A*, 110, 10435-10443, doi:10.1021/jp062403c, 2006.
- 350 Anttila, T., Kiendler-Scharr, A., Mentel, T. F., and Tillmann, R.: Size dependent partitioning of organic  
351 material: evidence for the formation of organic coatings on aqueous aerosols, *J. Atmos. Chem.*, 57, 215-  
352 237, doi:10.1007/s10874-007-9067-9, 2007.
- 353 Badger, C. L., George, I., Griffiths, P. T., Braban, C. F., Cox, R. A., and Abbatt, J. P. D.: Phase  
354 transitions and hygroscopic growth of aerosol particles containing humic acid and mixtures of humic acid  
355 and ammonium sulfate, *Atmos. Chem. Phys.*, 6, 755-768, 2006.
- 356 Bertram, A. K., Martin, S. T., Hanna, S. J., Smith, M. L., Bodsworth, A., Chen, Q., Kuwata, M., Liu, A.,  
357 You, Y., and Zorn, S. R.: Predicting the relative humidities of liquid-liquid phase separation,  
358 efflorescence, and deliquescence of mixed particles of ammonium sulfate, organic material, and water  
359 using the organic-to-sulfate mass ratio of the particle and the oxygen-to-carbon elemental ratio of the  
360 organic component, *Atmos. Chem. Phys.*, 11, 10995-11006, doi:10.5194/acp-11-10995-2011, 2011.
- 361 Bodsworth, A., Zobrist, B., and Bertram, A. K.: Inhibition of efflorescence in mixed organic-inorganic  
362 particles at temperatures less than 250 K, *Phys. Chem. Chem. Phys.*, 12, 12259-12266,  
363 doi:10.1039/c0cp00572j, 2010.
- 364 Braban, C. F., and Abbatt, J. P. D.: A study of the phase transition behavior of mixed ammonium sulfate -  
365 malonic acid aerosols, *Atmos. Chem. Phys.*, 4, 1451, 2004.
- 366 Brooks, S. D., Wise, M. E., Cushing, M., and Tolbert, M. A.: Deliquescence behavior of  
367 organic/ammonium sulfate aerosol, *Geophys. Res. Lett.*, 29, 1917, doi:10.1029/2002GL014733, 2002.
- 368 Brooks, S. D., Garland, R. M., Wise, M. E., Prenni, A. J., Cushing, M., Hewitt, E., and Tolbert, M. A.:  
369 Phase changes in internally mixed maleic acid/ammonium sulfate aerosols, *J. Geophys. Res.*, 108, 4478,  
370 doi:10.1029/2002jd003204, 2003.
- 371 Buajarnern, J., Mitchem, L., and Reid, J. P.: Characterizing multiphase Organic/Inorganic/Aqueous aerosol  
372 droplets, *J. Phys. Chem. A*, 111, 9054-9061, doi: 10.1021/jp074366a, 2007.
- 373 Buzorius, G., Zelenyuk, A., Brechtel, F., and Imre, D.: Simultaneous determination of individual ambient  
374 particle size, hygroscopicity and composition, *Geophys. Res. Lett.*, 29, 1974, doi:  
375 10.1029/2001GL014221, 2002.
- 376 Chan, M. N., and Chan, C. K.: Hygroscopic properties of two model humic-like substances and their  
377 mixtures with inorganics of atmospheric importance, *Environ. Sci. Technol.*, 37, 5109-5115,  
378 doi:10.1021/es034272o, 2003.
- 379 Chang, E. I., and Pankow, J. F.: Prediction of activity coefficients in liquid aerosol particles containing  
380 organic compounds, dissolved inorganic salts, and water - Part 2: Consideration of phase separation  
381 effects by an X-UNIFAC model, *Atmos. Environ.*, 40, 6422-6436, doi:10.1016/j.atmosenv.2006.04.031,  
382 2006.

383 Chen, Q., Farmer, D. K., Schneider, J., Zorn, S. R., Heald, C. L., Karl, T. G., Guenther, A., Allan, J. D.,  
384 Robinson, N., Coe, H., Kimmel, J. R., Pauliquevis, T., Borrmann, S., Pöschl, U., Andreae, M. O., Artaxo,  
385 P., Jimenez, J. L., and Martin, S. T.: Mass spectral characterization of submicron biogenic organic  
386 particles in the Amazon Basin, *Geophys. Res. Lett.*, 36, L20806, doi:10.1029/2009GL039880, 2009.

387 Choi, M. Y., and Chan, C. K.: The effects of organic species on the hygroscopic behaviors of inorganic  
388 aerosols, *Environ. Sci. Technol.*, 36, 2422-2428, doi:10.1021/es0113293, 2002.

389 Ciobanu, V. G., Marcolli, C., Krieger, U. K., Weers, U., and Peter, T.: Liquid-liquid phase separation in  
390 mixed organic/inorganic aerosol particles, *J. Phys. Chem. A*, 113, 10966-10978, doi: 10.1021/jp905054d,  
391 2009.

392 Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: A thermodynamic model of the  $\text{H}^+\text{-NH}_4^+\text{-SO}_4^{2-}\text{-NO}_3^-$   
393  $\text{-H}_2\text{O}$  at tropospheric temperatures. *J. Phys. Chem. A*, 102, 2137-2154, doi: 10.1021/Jp973042r, 1998.

394 Clegg, S. L., Seinfeld, J. H., and Brimblecombe, P.: Thermodynamic modelling of aqueous aerosols  
395 containing electrolytes and dissolved organic compounds, *J. Aerosol. Sci.*, 32, 713-738, doi:  
396 10.1016/S0021-8502(00)00105-1, 2001.

397 Cosman, L. M., and Bertram, A. K.: Reactive uptake of  $\text{N}_2\text{O}_5$  on aqueous  $\text{H}_2\text{SO}_4$  solutions coated with 1-  
398 component and 2-component monolayers, *J. Phys. Chem. A*, 112, 4625-4635, doi: 10.1021/Jp8005469,  
399 2008.

400 Day, D. A., Takahama, S., Gilardoni, S., and Russell, L. M.: Organic composition of single and  
401 submicron particles in different regions of western North America and the eastern Pacific during INTEX-  
402 B 2006, *Atmos. Chem. Phys.*, 9, 5433-5446, 2009.

403 DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crouse, J., Wennberg, P. O.,  
404 Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D., Weinheimer,  
405 A. J., Montzka, D. D., Campos, T., and Jimenez, J. L.: Fast airborne aerosol size and chemistry  
406 measurements above Mexico City and Central Mexico during the MILAGRO campaign, *Atmos. Chem.*  
407 *Phys.*, 8, 4027-4048, 2008.

408 Decesari, S., Fuzzi, S., Facchini, M. C., Mircea, M., Emblico, L., Cavalla, F., Maenhaut, W., Chi, X.,  
409 Schkolnik, G., Falkovich, A., Rudich, Y., Claeys, M., Pashynska, V., Vas, G., Kourtchev, I., Vermeylen,  
410 R., Hoffer, A., Andreae, M. O., Tagliavini, E., Moretti, F., and Artaxo, P.: Characterization of the organic  
411 composition of aerosols from Rondonia, Brazil, during the LBA-SMOCC 2002 experiment and its  
412 representation through model compounds, *Atmos. Chem. Phys.*, 6, 375-402, 2006.

413 Dibb, J. E., Talbot, R. W., Klemm, K. I., Gregory, G. L., Singh, H. B., Bradshaw, J. D., and Sandholm, S.  
414 T.: Asian influence over the western North Pacific during the fall season: Inferences from lead 210,  
415 soluble ionic species and ozone, *J. Geophys. Res.*, 101, 1779-1792, doi: 10.1029/94JD03117, 1996.

416 Dibb, J. E., Talbot, R. W., and Scheuer, E. M.: Composition and distribution of aerosols over the North  
417 Atlantic during the Subsonic Assessment Ozone and Nitrogen Oxide Experiment (SONEX), *J. Geophys.*  
418 *Res.*, 105, 3709-3717, doi: 10.1029/1999JD900424, 2000.

419 Drozd, G. T., Woo, J. L., and McNeill, V. F.: Self-limited uptake of  $\alpha$ -pinene oxide to acidic aerosol: the  
420 effects of liquid-liquid phase separation and implications for the formation of secondary organic aerosol  
421 and organosulfates from epoxides, *Atmos. Chem. Phys.*, 13, 8255-8263, doi: 10.5194/acp-13-8255-2013,  
422 2013.

423 Erdakos, G. B., Chang, E. I., Pankow, J. F., and Seinfeld, J. H.: Prediction of activity coefficients in liquid  
424 aerosol particles containing organic compounds, dissolved inorganic salts, and water - Part 3: Organic  
425 compounds, water, and ionic constituents by consideration of short-, mid-, and long-range effects using  
426 X-UNIFAC.3, *Atmos. Environ.*, 40, 6437-6452, doi:10.1016/j.atmosenv.2006.04.001, 2006.

427 Escoreia, E. N., Sjostedt, S. J., and Abbatt, J. P. D.: Kinetics of N<sub>2</sub>O<sub>5</sub> hydrolysis on secondary organic  
428 aerosol and mixed ammonium bisulfate-secondary organic aerosol particles, *J. Phys. Chem. A*, 114,  
429 13113-13121, doi: 10.1021/jp107721v, 2010.

430 Faucher, J. A.; Koleske, J. V.; Santee, E. R.; Stratta, J. J.; Wilson, C. W.: Glass transitions of ethylene  
431 oxide polymers, *J. Appl. Phys.* 37, 3962-3964, 1966.

432 Finlayson-Pitts, B. J., and Pitts, J. N.: Tropospheric air pollution: Ozone, airborne toxics, polycyclic  
433 aromatic hydrocarbons, and particles, *Science*, 276, 1045-1052, doi: 10.1126/science.276.5315.1045,  
434 1997.

435 Finlayson-Pitts, B. J., and Pitts, J. N.: Chemistry of the upper and lower atmosphere: theory, experiments  
436 and applications, Academic Press, San Diego, California. ; London, pp.969, 2000.

437 Folkers, M., Mentel, T. F., and Wahner, A.: Influence of an organic coating on the reactivity of aqueous  
438 aerosols probed by the heterogeneous hydrolysis of N<sub>2</sub>O<sub>5</sub>, *Geophys. Res. Lett.*, 30, 1644,  
439 doi:10.1029/2003GL017168, 2003.

440 Fu, P. Q., Kawamura, K., and Miura, K.: Molecular characterization of marine organic aerosols collected  
441 during a round-the-world cruise, *J. Geophys. Res.*, 116, D13302, doi: 10.1029/2011jd015604, 2011.

442 Fuzzi, S., Decesari, S., Facchini, M. C., Matta, E., Mircea, M., and Tagliavini, E.: A simplified model of  
443 the water soluble organic component of atmospheric aerosols, *Geophys. Res. Lett.*, 28, 4079-4082, doi:  
444 10.1029/2001gl013418, 2001.

445 Gao, S., Keywood, M., Ng, N. L., Surratt, J., Varutbamkul, V., Bahreini, R., Flagan, R. C., and Seinfeld,  
446 J. H.: Low-molecular-weight and oligomeric components in secondary organic aerosol from the  
447 ozonolysis of cycloalkenes and  $\alpha$ -pinene, *J. Phys. Chem. A*, 108, 10147-10164, doi: 10.1021/jp047466e,  
448 2004.

449 Gaston, C. J., Thornton, J. A., and Ng, N. L.: Reactive uptake of N<sub>2</sub>O<sub>5</sub> to internally mixed inorganic and  
450 organic particles: the role of organic carbon oxidation state and inferred organic phase separations,  
451 *Atmos. Chem. Phys.*, 14, 5693-5707, 2014.

452 Gilardoni, S., Liu, S., Takahama, S., Russell, L. M., Allan, J. D., Steinbrecher, R., Jimenez, J. L., De  
453 Carlo, P. F., Dunlea, E. J., and Baumgardner, D.: Characterization of organic ambient aerosol during  
454 MIRAGE 2006 on three platforms, *Atmos. Chem. Phys.*, 9, 5417-5432, 2009.

455 Goldstein, A. H., and Galbally, I. E.: Known and unexplored organic constituents in the earth's  
456 atmosphere, *Environ. Sci. Technol.*, 41, 1514-1521, doi: 10.1021/es072476p, 2007.

457 Greener, J., Gillmor, J. R., and Daly, R. C. : Melt rheology of a class of polyester ionomers,  
458 *Macromolecules*, 26, 6416-6424, doi: 10.1021/ma00076a018, 1993.

459 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J.,  
460 Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y.,  
461 Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F.,  
462 Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation,  
463 properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9,  
464 5155-5236, 2009.

465 Hamilton, J. F., Webb, P. J., Lewis, A. C., Hopkins, J. R., Smith, S., and Davy, P.: Partially oxidised  
466 organic components in urban aerosol using GCXGC-TOF/MS, *Atmos. Chem. Phys.*, 4, 1279-1290, 2004.

467 Hawkins, L. N., Russell, L. M., Covert, D. S., Quinn, P. K., and Bates, T. S.: Carboxylic acids, sulfates,  
468 and organosulfates in processed continental organic aerosol over the southeast Pacific Ocean during  
469 VOCALS-REx 2008, *J. Geophys. Res.*, 115, D13201, doi:10.1029/2009jd013276, 2010.

470 Heald, C. L., Kroll, J. H., Jimenez, J. L., Docherty, K. S., DeCarlo, P. F., Aiken, A. C., Chen, Q., Martin,  
471 S. T., Farmer, D. K., and Artaxo, P.: A simplified description of the evolution of organic aerosol  
472 composition in the atmosphere, *Geophys. Res. Lett.*, 37, L08803, doi:10.1029/2010gl042737, 2010.

473 Huebert, B. J., Howell, S. G., Zhuang, L., Heath, J. A., Litchy, M. R., Wylie, D. J., Kreidler-Moss, J. L.,  
474 Coppicus, S., and Pfeiffer, J. E.: Filter and impactor measurements of anions and cations during the First  
475 Aerosol Characterization Experiment (ACE 1), *J. Geophys. Res.*, 103, 16493-16509,  
476 doi:10.1029/98jd00770, 1998.

477 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P.  
478 F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P.,  
479 Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J.,  
480 Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M.,  
481 Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P.  
482 I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo,  
483 D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y.  
484 M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L.  
485 R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of  
486 organic aerosols in the atmosphere, *Science*, 326, 1525-1529, doi: 10.1126/science.1180353, 2009.

487 Kalberer, M., Paulsen, D., Sax, M., Steinbrecher, M., Dommen, J., Prevot, A. S. H., Fisseha, R.,  
488 Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major  
489 components of atmospheric organic aerosols, *Science*, 303, 1659-1662, doi: 10.1126/science.1092185,  
490 2004.

491 Kidd, C., Perraud, V., Wingen, L. M., and Finlayson-Pitts, B. J.: Integrating phase and composition of  
492 secondary organic aerosol from the ozonolysis of  $\alpha$ -pinene, *P. Natl. Acad. Sci. USA*, 111, 7552-7557,  
493 doi:10.1073/pnas.1322558111, 2014.

494 Koop, T., Kapilashrami, A., Molina, L. T., and Molina, M. J.: Phase transitions of sea-salt/water mixtures  
495 at low temperatures: Implications for ozone chemistry in the polar marine boundary layer, *J. Geophys.*  
496 *Res.*, 105, 26393-26402, doi: 10.1029/2000jd900413, 2000.

497 Koop, T., Bookhold, J., Shiraiwa, M., and Pöschl, U.: Glass transition and phase state of organic  
498 compounds: dependency on molecular properties and implications for secondary organic aerosols in the  
499 atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19238-19255, doi: 10.1039/C1cp22617g, 2011.

500 Krieger, U. K., Marcolli, C., and Reid, J. P.: Exploring the complexity of aerosol particle properties and  
501 processes using single particle techniques, *Chem. Soc. Rev.*, 41, 6631-6662, doi: 10.1039/C2cs35082c,  
502 2012.

503 Kwamena, N. O. A., Buajarern, J., and Reid, J. P.: Equilibrium Morphology of Mixed  
504 Organic/Inorganic/Aqueous Aerosol Droplets: Investigating the Effect of Relative Humidity and  
505 Surfactants, *J. Phys. Chem. A*, 114, 5787-5795, doi: 10.1021/jp1003648, 2010.

506 Lee, Y.-N., Weber, R., Ma, Y., Orsini, D., Maxwell-Meier, K., Blake, D., Meinardi, S., Sachse, G.,  
507 Harward, C., Chen, T. Y., Thornton, D., Tu, F.-H., and Bandy, A.: Airborne measurement of inorganic  
508 ionic components of fine aerosol particles using the particle-into-liquid sampler coupled to ion  
509 chromatography technique during ACE-Asia and TRACE-P, *J. Geophys. Res.*, 108, 8646,  
510 doi:10.1029/2002JD003265, 2003.

511 Levine, I. N.: *Physical Chemistry*, 6th edition. McGraw-Hill, New York, 2009

512 Ling, T. Y., and Chan, C. K.: Partial crystallization and deliquescence of particles containing ammonium  
513 sulfate and dicarboxylic acids, *J. Geophys. Res.*, 113 (D14), D14205, doi:10.1029/2008JD009779, 2008.

514 Liu, S., Takahama, S., Russell, L. M., Gilardoni, S., and Baumgardner, D.: Oxygenated organic functional  
515 groups and their sources in single and submicron organic particles in MILAGRO 2006 campaign, *Atmos.*  
516 *Chem. Phys.*, 9, 6849-6863, 2009.

517 Malcolm, G. N. and Rowlinson, J. S.: The Thermodynamic Properties of Aqueous Solutions of  
518 Polyethylene Glycol, Polypropylene Glycol and Dioxane, *T. Faraday Soc.*, 53, 921-931, doi:  
519 10.1039/Tf9575300921, 1957.

520 Marcolli, C., and Krieger, U. K.: Phase changes during hygroscopic cycles of mixed organic/inorganic  
521 model systems of tropospheric aerosols, *J. Phys. Chem. A*, 110, 1881-1893, doi: 10.1021/jp0556759,  
522 2006.

523 Martin, S. T.: Phase transitions of aqueous atmospheric particles, *Chem. Rev.*, 100, 3403-3453, doi:  
524 10.1021/cr990034t, 2000.

525 McNeill, V. F., Patterson, J., Wolfe, G. M., and Thornton, J. A.: The effect of varying levels of surfactant  
526 on the reactive uptake of N<sub>2</sub>O<sub>5</sub> to aqueous aerosol, *Atmos. Chem. Phys.*, 6, 1635-1644, 2006.

527 Middlebrook, A. M., Murphy, D. M., and Thomson, D. S.: Observations of organic material in individual  
528 marine particles at Cape Grim during the First Aerosol Characterization Experiment (ACE 1), *J. Geophys.*  
529 *Res.*, 103, 16475-16483, doi: 10.1029/97jd03719, 1998.

530 Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol  
531 particles interacting with water vapor: conceptual framework and experimental evidence for restructuring,  
532 phase transitions and kinetic limitations, *Atmos. Chem. Phys.*, 9, 9491-9522, 2009.

533 Murphy, D. M., and Thomson, D. S.: Chemical composition of single aerosol particles at Idaho Hill:  
534 Negative ion measurements, *J. Geophys. Res.*, 102, 6353-6368, doi: 10.1029/96jd00859, 1997.

535 Murphy, D. M., Thomson, D. S., and Mahoney, T. M. J.: In situ measurements of organics, meteoritic  
536 material, mercury, and other elements in aerosols at 5 to 19 kilometers, *Science*, 282, 1664-1669, doi:  
537 10.1126/science.282.5394.1664, 1998.

538 Murphy, D. M.: Something in the air, *Science*, 307, 1888-1890, doi: 10.1126/science.1108160, 2005.

539 Murphy, D. M., Cziczo, D. J., Froyd, K. D., Hudson, P. K., Matthew, B. M., Middlebrook, A. M., Peltier,  
540 R. E., Sullivan, A., Thomson, D. S., and Weber, R. J.: Single-particle mass spectrometry of tropospheric  
541 aerosol particles, *J. Geophys. Res.*, 111, D23S32, doi:10.1029/2006JD007340, 2006.

542 Murray, B. J.: Inhibition of ice crystallisation in highly viscous aqueous organic acid droplets, *Atmos.*  
543 *Chem. Phys.*, 8, 5423-5433, 2008.

544 Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty,  
545 K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M.,  
546 DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol  
547 components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem.*  
548 *Phys.*, 10, 4625-4641, doi:10.5194/acp-10-4625-2010, 2010.

549 Nguyen, T. B., Bateman, A. P., Bones, D. L., Nizkorodov, S. A., Laskin, J., and Laskin, A.: High-  
550 resolution mass spectrometry analysis of secondary organic aerosol generated by ozonolysis of isoprene,  
551 *Atmos. Environ.*, 44, 1032-1042, doi: 10.1016/j.atmosenv.2009.12.019, 2010.

552 Pankow, J. F.: Gas/particle partitioning of neutral and ionizing compounds to single and multi-phase  
553 aerosol particles. 1. Unified modeling framework, *Atmos. Environ.*, 37, 3323-3333, doi:10.1016/S1352-  
554 2310(03)00346-7, 2003.

555 Pant, A., Fok, A., Parsons, M. T., Mak, J., and Bertram, A. K.: Deliquescence and crystallization of  
556 ammonium sulfate-glutaric acid and sodium chloride-glutaric acid particles, *Geophys. Res. Lett.*, 31,  
557 L12111, doi:10.1029/2004GL020025, 2004.

558 Pant, A., Parsons, M. T., and Bertram, A. K.: Crystallization of aqueous ammonium sulfate particles  
559 internally mixed with soot and kaolinite: Crystallization relative humidities and nucleation rates, *J. Phys.*  
560 *Chem. A*, 110, 8701-8709, doi: 10.1021/jp060985s, 2006.

561 Park, S. C., Burden, D. K., and Nathanson, G. M.: The inhibition of N<sub>2</sub>O<sub>5</sub> hydrolysis in sulfuric acid by 1-  
562 butanol and 1-hexanol surfactant coatings, *J. Phys. Chem. A*, 111, 2921-2929, doi: 10.1021/Jp068228h,  
563 2007.

564 Parsons, M. T., Knopf, D. A., and Bertram, A. K.: Deliquescence and crystallization of ammonium sulfate  
565 particles internally mixed with water-soluble organic compounds, *J. Phys. Chem. A*, 108, 11600-11608,  
566 doi: 10.1021/jp0462862, 2004a.

567 Parsons, M. T., Mak, J., Lipetz, S. R., and Bertram, A. K.: Deliquescence of malonic, succinic, glutaric,  
568 and adipic acid particles, *J. Geophys. Res.*, 109, D06212, doi:10.1029/2003JD004075, 2004b.

569 Parsons, M. T., Riffell, J. L., and Bertram, A. K.: Crystallization of aqueous inorganic-malonic acid  
570 particles: Nucleation rates, dependence on size, and dependence on the ammonium-to-sulfate, *J. Phys.*  
571 *Chem. A*, 110, 8108-8115, doi: 10.1021/jp057074n, 2006.

572 Pöhlker, C., Wiedemann, K. T., Sinha, B., Shiraiwa, M., Gunthe, S. S., Smith, M., Su, H., Artaxo, P.,  
573 Chen, Q., Cheng, Y. F., Elbert, W., Gilles, M. K., Kilcoyne, A. L. D., Moffet, R. C., Weigand, M.,  
574 Martin, S. T., Pöschl, U., and Andreae, M. O.: Biogenic Potassium Salt Particles as Seeds for Secondary  
575 Organic Aerosol in the Amazon, *Science*, 337, 1075-1078, doi: 10.1126/science.1223264, 2012.

576 Power, R. M., Simpson, S. H., Reid, J. P., and Hudson, A. J.: The transition from liquid to solid-like  
577 behaviour in ultrahigh viscosity aerosol particles, *Chem Sci*, 4, 2597-2604, doi: 10.1039/C3sc50682g,  
578 2013.

579 Pratt, K. A., and Prather, K. A.: Aircraft measurements of vertical profiles of aerosol mixing states, *J.*  
580 *Geophys. Res.*, 115, D11305, doi:10.1029/2009JD013150, 2010.

581 Prisle, N. L., Engelhart, G. J., Bilde, M., and Donahue, N. M.: Humidity influence on gas-particle phase  
582 partitioning of  $\alpha$ -pinene + O<sub>3</sub> secondary organic aerosol, *Geophys. Res. Lett.*, 37, L01802,  
583 doi:10.1029/2009GL041402, 2010.

584 Reid, J. P., Dennis-Smith, B. J., Kwamena, N. O. A., Miles, R. E. H., Hanford, K. L., and Homer, C. J.:  
585 The morphology of aerosol particles consisting of hydrophobic and hydrophilic phases: hydrocarbons,  
586 alcohols and fatty acids as the hydrophobic component, *Phys. Chem. Chem. Phys.*, 13, 15559-15572, doi:  
587 10.1039/c1cp21510h, 2011.

588 Renbaum-Wolff, L., Grayson, J. W., Bateman, A. P., Kuwata, M., Sellier, M., Murray, B. J., Shilling, J.  
589 E., Martin, S. T., and Bertram, A. K.: Viscosity of  $\alpha$ -pinene secondary organic material and implications  
590 for particle growth and reactivity, *P. Natl. Acad. Sci. USA*, 110, 8014-8019,  
591 doi:10.1073/pnas.1219548110, 2013.

592 Rogge, W. F., Mazurek, M. A., Hildemann, L. M., Cass, G. R., and Simoneit, B. R. T.: Quantification of  
593 Urban Organic Aerosols At a Molecular-Level: Identification, Abundance and Seasonal-Variation,  
594 *Atmos. Environ., Part A* 27, 1309-1330, doi: 10.1016/0960-1686(93)90257-Y, 1993.

595 Russell, L. M., Takahama, S., Liu, S., Hawkins, L. N., Covert, D. S., Quinn, P. K., and Bates, T. S.:  
596 Oxygenated fraction and mass of organic aerosol from direct emission and atmospheric processing  
597 measured on the R/V Ronald Brown during TEXAQS/GoMACCS 2006, *J. Geophys. Res.*, 114, D00F05,  
598 doi:10.1029/2008JD011275, 2009.

599 Russell, L. M., Bahadur, R., and Ziemann, P. J.: Identifying organic aerosol sources by comparing  
600 functional group composition in chamber and atmospheric particles, *P. Natl. Acad. Sci. USA*, 108, 3516-  
601 3521, doi: 10.1073/pnas.1006461108, 2011.



602 Salcedo, D.: Equilibrium phase diagrams of aqueous mixtures of malonic acid and sulfate/ammonium  
603 salts, *J. Phys. Chem. A*, 110, 12158-12165, doi: 10.1021/jp063850v, 2006.

604 Saukko, E., Lambe, A. T., Massoli, P., Koop, T., Wright, J. P., Croasdale, D. R., Pedernera, D. A.,  
605 Onasch, T. B., Laaksonen, A., Davidovits, P., Worsnop, D. R., and Virtanen, A.: Humidity-dependent  
606 phase state of SOA particles from biogenic and anthropogenic precursors, *Atmos. Chem. Phys.*, 12, 7517-  
607 7529, doi: 10.5194/acp-12-7517-2012, 2012.

608 Saxena, P., and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical review of  
609 the literature and application of thermodynamics to identify candidate compounds, *J. Atmos. Chem.*, 24,  
610 57-109, doi: 10.1007/BF00053823, 1996.

611 Schill, G. P., and Tolbert, M. A.: Heterogeneous ice nucleation on phase-separated organic-sulfate  
612 particles: effect of liquid vs. glassy coatings, *Atmos. Chem. Phys.*, 13, 4681-4695, doi: 10.5194/acp-13-  
613 4681-2013, 2013.

614 Seinfeld, J. H., Erdakos, G. B., Asher, W. E., and Pankow, J. F.: Modeling the formation of secondary  
615 organic aerosol (SOA). 2. The predicted effects of relative humidity on aerosol formation in the  $\alpha$ -pinene-  
616  $\beta$ -pinene-, sabinene-,  $\Delta^3$ -Carene-, and cyclohexene-ozone systems, *Environ. Sci. Technol.*, 35, 1806-  
617 1817, doi: 10.1021/es001765+, 2001.

618 Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air pollution to Climate  
619 Change*, 2nd ed., Wiley-Interscience, Hoboken, New Jersey, 2006.

620 Shiraiwa, M., Zuend, A., Bertram, A. K., and Seinfeld, J. H.: Gas-particle partitioning of atmospheric  
621 aerosols: interplay of physical state, non-ideal mixing and morphology, *Phys. Chem. Chem. Phys.*, 15,  
622 11441-11453, doi: 10.1039/C3cp51595h, 2013.

623 Slade, L. and Levine, H.: Water and the glass-transition - dependence of the glass-transition on  
624 composition and chemical-structure -special implications for flour functionality in cookie baking, *J. Food  
625 Eng.*, 22, 143-188, doi 10.1016/0260-8774(94)90029-9, 1994.

626 Smith, M. L., Kuwata, M., and Martin, S. T.: Secondary organic material produced by the dark ozonolysis  
627 of  $\alpha$ -pinene minimally affects the deliquescence and efflorescence of ammonium sulfate, *Aerosol Sci.  
628 Technol.*, 45, 244-261, doi: 10.1080/02786826.2010.532178, 2011.

629 Smith, M. L., Bertram, A. K., and Martin, S. T.: Deliquescence, efflorescence, and phase miscibility of  
630 mixed particles of ammonium sulfate and isoprene-derived secondary organic material, *Atmos. Chem.  
631 Phys.*, 12, 9613-9628, doi:10.5194/acp-12-9613-2012, 2012.

632 Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid-liquid phase separation and  
633 morphology of internally mixed dicarboxylic acids/ammonium sulfate/water particles, *Atmos. Chem.  
634 Phys.*, 12, 2691-2712, doi:10.5194/acp-12-2691-2012, 2012a.

635 Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid-liquid phase separation in aerosol  
636 particles: Dependence on O:C, organic functionalities, and compositional complexity, *Geophys. Res.  
637 Lett.*, 39, L19801, doi:10.1029/2012GL052807, 2012b.

638 Song, M., Marcolli, C., Krieger, U. K., Lienhard, D. M., and Peter, T.: Morphologies of mixed  
639 organic/inorganic/aqueous aerosol droplets, *Faraday Discuss.*, 165, 289-316, doi:10.1039/C3FD00049D,  
640 2013.

641 Takahama, S., Schwartz, R. E., Russell, L. M., Macdonald, A. M., Sharma, A., and Leitch, W. R.:  
642 Organic functional groups in aerosol particles from burning and non-burning forest emissions at high-  
643 elevation mountain site, *Atmos. Chem. Phys.*, 11, 6367-6386, doi:10.5194/acp-11-6367-2011, 2011.

644 Talbot, R. W., Dibb, J. E., and Loomis, M. B.: Influence of vertical transport on free tropospheric aerosols  
645 over the central USA in springtime, *Geophys. Res. Lett.*, 25, 1367-1370, doi: 10.1029/98GL00184, 1998.

646 Thornton, J. A., and Abbatt, J. P. D.:  $\text{N}_2\text{O}_5$  reaction on submicron sea salt aerosol: Kinetics, products, and  
647 the effect of surface active organics, *J. Phys. Chem. A*, 109, 10004-10012, doi: 10.1021/Jp054183t, 2005.

648 Tolocka, M. P., Jang, M., Ginter, J. M., Cox, R. J., Kamens, R. M., and Johnston, M. V.: Formation of  
649 oligomers in secondary organic aerosol, *Environ. Sci. Technol.*, 38, 1428-1434, doi: 10.1021/es035030r,  
650 2004.

651 Tolocka, M. P., Lake, D. A., Johnston, M. V., and Wexler, A. S.: Size-resolved fine and ultrafine particle  
652 composition in Baltimore, Maryland, *J. Geophys. Res.*, 110, D07S04, doi:10.1029/2004JD004573, 2005.

653 Tong, H. J., Reid, J. P., Bones, D. L., Luo, B. P., and Krieger, U. K.: Measurements of the timescales for  
654 the mass transfer of water in glassy aerosol at low relative humidity and ambient temperature, *Atmos.*  
655 *Chem. Phys.*, 11, 4739-4754, doi: 10.5194/acp-11-4739-2011, 2011.

656 Treuel, L., Pederzani, S., and Zellner, R.: Deliquescence behaviour and crystallisation of ternary  
657 ammonium sulfate/dicarboxylic acid/water aerosols, *Phys. Chem. Chem. Phys.*, 11, 7976-7984, doi:  
658 10.1039/b905007h, 2009.

659 Veghte, D. P., Altaf, M. B., and Freedman, M. A.: Size Dependence of the Structure of Organic Aerosol,  
660 *J. Am. Chem. Soc.*, 135, 16046-16049, doi: 10.1021/Ja408903g, 2013.

661 Veghte, D. P., Bittner, D. R., and Freedman, M. A.: Cryo-Transmission Electron Microscopy Imaging of  
662 the Morphology of Submicrometer Aerosol Containing Organic Acids and Ammonium Sulfate, *Anal.*  
663 *Chem.*, 86, 2436-2442, doi:10.1021/ac403279f, 2014.

664 Wise, M. E., Surratt, J. D., Curtis, D. B., Shilling, J. E., and Tolbert, M. A.: Hygroscopic growth of  
665 ammonium sulfate/dicarboxylic acids, *J. Geophys. Res.*, 108, 4638, doi:10.1029/2003JD003775, 2003.

666 You, Y., Renbaum-Wolff, L., Carreras-Sospedra, M., Hanna, S. J., Hiranuma, N., Kamal, S., Smith, M.  
667 L., Zhang, X. L., Weber, R. J., Shilling, J. E., Dabdub, D., Martin, S. T., and Bertram, A. K.: Images  
668 reveal that atmospheric particles can undergo liquid-liquid phase separations, *P. Natl. Acad. Sci. USA*,  
669 109, 13188-13193, doi: 10.1073/pnas.1206414109, 2012.

670 You, Y., Renbaum-Wolff, L., and Bertram, A. K.: Liquid-liquid phase separation in particles containing  
671 organics mixed with ammonium sulfate, ammonium bisulfate, ammonium nitrate or sodium chloride,  
672 *Atmos. Chem. Phys.*, 13, 11723-11734, doi: 10.5194/acp-13-11723-2013, 2013.

673 You, Y., Smith, M. L., Song, M., Martin, S. T., and Bertram, A. K.: Liquid-liquid phase separation in  
674 atmospherically relevant particles consisting of organic species and inorganic salts, *International Reviews*  
675 *in Physical Chemistry*, 33, 43-77, doi: 10.1080/0144235X.2014.890786, 2014.

676 Zobrist, B., Marcolli, C., Pedernera, D. A., and Koop, T.: Do atmospheric aerosols form glasses?, *Atmos.*  
677 *Chem. Phys.*, 8, 5221-5244, 2008.

678 Zobrist, B., Soonsin, V., Luo, B. P., Krieger, U. K., Marcolli, C., Peter, T., and Koop, T.: Ultra-slow  
679 water diffusion in aqueous sucrose glasses, *Phys. Chem. Chem. Phys.*, 13, 3514-3526,  
680 doi:10.1039/c0cp01273d, 2011.

681 Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic  
682 aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559-4593, 2008.

683 Zuend, A., Marcolli, C., Peter, T., and Seinfeld, J. H.: Computation of liquid-liquid equilibria and phase  
684 stabilities: implications for RH-dependent gas/particle partitioning of organic-inorganic aerosols, *Atmos.*  
685 *Chem. Phys.*, 10, 7795-7820, doi:10.5194/acp-10-7795-2010, 2010.

686 Zuend, A., Marcolli, C., Booth, A. M., Lienhard, D. M., Soonsin, V., Krieger, U. K., Topping, D. O.,  
687 McFiggans, G., Peter, T., and Seinfeld, J. H.: New and extended parameterization of the thermodynamic  
688 model AIOMFAC: calculation of activity coefficients for organic-inorganic mixtures containing carboxyl,

- 689 hydroxyl, carbonyl, ether, ester, alkenyl, alkyl, and aromatic functional groups, *Atmos. Chem. Phys.*, 11,  
690 9155-9206, doi: 10.5194/acp-11-9155-2011, 2011.
- 691 Zuend, A., and Seinfeld, J. H.: Modeling the gas-particle partitioning of secondary organic aerosol: the  
692 importance of liquid-liquid phase separation, *Atmos. Chem. Phys.*, 12, 3857-3882, doi: 10.5194/acp-12-  
693 3857-2012, 2012.
- 694 Zuend, A., and Seinfeld, J. H.: A practical method for the calculation of liquid-liquid equilibria in  
695 multicomponent organic-water-electrolyte systems using physicochemical constraints, *Fluid Phase*  
696 *Equilib.*, 337, 201-213, doi: 10.1016/j.fluid.2012.09.034, 2013.

697 **Tables**

698 **Table 1.** List of the ten organic species used in molecular weight dependent measurements. Each  
 699 organic species was separately mixed with ammonium sulfate to make particles, and liquid-  
 700 liquid phase separation was studied in these particles at  $290 \pm 1$  K.

Compounds	Formula	Molecular weight (Da)	O:C	Functional group(s)	Glass transition temperature, $T_g$ (K)	References for $T_g$
Glucose	$C_6H_{12}O_6$	180.2	1	alcohol, ether	$296.1 \pm 3.1$	Zobrist et al. 2008
Poly(ethylene glycol) bis(carboxymethyl) ether	$C_{2n+4}H_{4n+6}O_{n+5}$	250	0.83	alcohol, ether	-	-
Sucrose	$C_{12}H_{22}O_{11}$	342.3	0.92	alcohol, ether	$335.7 \pm 3.6$	Zobrist et al. 2008
Poly(ethylene glycol)	$C_{2n}H_{4n+2}O_{n+1}$	400	0.56	alcohol, ether	223	Faucher et al. 1966
Ouabain	$C_{29}H_{44}O_{12}$	584	0.41	alcohol, ether, ester, C-C double bonds	$373^a$	Koop et al. 2011
Raffinose	$C_{18}H_{32}O_{16}$	594.5	0.89	alcohol, ether	$395.7 \pm 21.6$	Zobrist et al. 2008
Poly(ethylene glycol)	$C_{2n}H_{4n+2}O_{n+1}$	600	0.54	alcohol, ether	231	Faucher et al. 1966
Maltopentaose	$C_{30}H_{52}O_{26}$	829	0.87	alcohol, ether	398	Slade and Levine 1994
Poly(ethylene glycol)	$C_{2n}H_{4n+2}O_{n+1}$	900	0.53	alcohol, ether	231-245	Faucher et al. 1966
Maltoheptaose	$C_{42}H_{72}O_{36}$	1153	0.86	alcohol, ether	412	Slade and Levine 1994

701

702 <sup>a</sup>  $T_g$  values are estimates based on the melting points of those organic compounds using the results from  
 703 Koop et al. 2011.

704 - Literature data not available

705

706 **Table 2.** Summary of the twenty organic species used in temperature dependent experiments.  
 707 Each organic species was separately mixed with ammonium sulfate to make particles, and liquid-  
 708 liquid phase separation was studied in these particles at  $244 \pm 1$  K,  $263 \pm 1$  K and  $278 \pm 1$  K.  
 709 Two of the organic species (poly(ethylene glycol) diacrylate and raffinose) mixed with  
 710 ammonium sulfate were also studied at  $290 \pm 1$  K.

Compounds	Formula	O:C	Molecular weight (Da)	Functional group(s)	Glass transition temperature $T_g$ (K)	References for $T_g$
2,5-hexanediol	$C_6H_{14}O_2$	0.33	118.2	alcohol	$192.4 \pm 1.3$	Zobrist et al. 2008
Poly(propylene glycol)	$C_{3n}H_{6n+2}O_{n+1}$	0.38	425	alcohol, ether	-	-
Poly(ethylene glycol) diacrylate	$C_{2n+6}H_{4n+6}O_{n+3}$	0.5	575	ester, ether, C-C double bond	-	-
Poly(ethylene glycol) 900	$C_{2n}H_{4n+2}O_{n+1}$	0.53	900	alcohol, ether	$231-245^a$	Faucher et al. 1966
$\alpha$ ,4-dihydroxy-3-methoxybenzoic acid	$C_9H_{10}O_5$	0.56	198.2	alcohol, aromatic, carboxylic acid, ether	$293.6 \pm 6.6$	Zobrist et al. 2008
Diethylmalonic acid	$C_7H_{12}O_4$	0.57	160.2	carboxylic acid	$282^b$	Koop et al. 2011
3,3-dimethylglutaric acid	$C_7H_{12}O_4$	0.57	160.2	carboxylic acid	$261^b$	Koop et al. 2011
2,5-hydroxybenzoic acid	$C_7H_6O_4$	0.57	154.2	carboxylic acid, aromatic	$225^b$	Koop et al. 2011
Poly(ethylene glycol) 300	$C_{2n}H_{4n+2}O_{n+1}$	0.58	300	alcohol, ether	203-223	Faucher et al. 1966
Poly(ethylene glycol) 200	$C_{2n}H_{4n+2}O_{n+1}$	0.63	200	alcohol, ether	203	Faucher et al. 1966
Poly(ethylene glycol) bis(carboxymethyl) ether	$C_{2n+4}H_{4n+6}O_{n+5}$	0.63	600	alcohol, ether	-	-
2-methylglutaric acid	$C_6H_{10}O_4$	0.67	146.1	carboxylic acid	$247^b$	Koop et al. 2011
2,2-dimethylsuccinic acid	$C_6H_{10}O_4$	0.67	146.2	carboxylic acid	$289^b$	Koop et al. 2011
Diethyl-L-tartrate	$C_8H_{14}O_6$	0.75	206.2	alcohol, ether	-	-
Glycerol	$C_3H_8O_3$	1.00	92.1	alcohol	$193.3 \pm 1.1$	Zobrist et al. 2008
Glutaric acid	$C_5H_8O_4$	0.8	132.1	carboxylic acid	$258^b$	Koop et al. 2011

Levoglucozan	$C_6H_{10}O_5$	0.83	162.1	alcohol, ester	319 <sup>b</sup>	Koop et al. 2011
Raffinose	$C_{18}H_{32}O_{16}$	0.89	594.5	alcohol, ester	395.7 ± 21.6	Zobrist et al. 2008
Citric acid	$C_6H_8O_7$	1.17	192.1	alcohol, carboxylic acid	280.1	Bodsworth et al. 2010
Malonic acid	$C_3H_4O_4$	1.33	104.1	carboxylic acid	286 <sup>b</sup>	Koop et al. 2011

711

712 <sup>a</sup> T<sub>g</sub> value is an estimate based on the molecular weight of the ploy (ethylene glycol 900) from Faucher et  
713 al. 1966.

714 <sup>b</sup> T<sub>g</sub> values are estimates based on the melting points of those organic compounds using the results from  
715 Koop et al. 2011.

716 - Literature data not available

717

718 **Table 3.** Combined data set used to assess the effect of molecular weight on SRH in mixed  
 719 organic-ammonium sulfate particles. This data set includes the ten types of particles studied here  
 720 (see Table 1) and the SRH results of twenty-three types of particles containing single organic  
 721 species and ammonium sulfate studied by You et al. (2013). OIR =  $2.0 \pm 0.1$  in all the  
 722 experiments. Uncertainties are 95% confidence intervals considering  $\sigma$  of multiple SRH  
 723 measurements and the uncertainty from the calibration.

Compounds	Formula	Molecular weight (Da)	O:C	SRH (%)	Reference
Glycerol	C <sub>3</sub> H <sub>8</sub> O <sub>3</sub>	92.1	1	Not observed	You et al. (2013)
Malonic acid	C <sub>3</sub> H <sub>4</sub> O <sub>4</sub>	104.1	1.33	Not observed	You et al. (2013)
Maleic acid	C <sub>4</sub> H <sub>4</sub> O <sub>4</sub>	116.1	1	Not observed	You et al. (2013)
2,5-hexanediol	C <sub>6</sub> H <sub>14</sub> O <sub>2</sub>	118.2	0.33	88.8 ± 7.1	You et al. (2013)
Glutaric acid	C <sub>5</sub> H <sub>8</sub> O <sub>4</sub>	132.1	0.8	Not observed	You et al. (2013)
Malic acid	C <sub>4</sub> H <sub>6</sub> O <sub>5</sub>	134.1	1.25	Not observed	You et al. (2013)
1,2,6-hexanetriol	C <sub>6</sub> H <sub>14</sub> O <sub>3</sub>	134.2	0.5	76.7 ± 6.2	You et al. (2013)
2-methylglutaric acid	C <sub>6</sub> H <sub>10</sub> O <sub>4</sub>	146.1	0.67	75.3 ± 6.4	You et al. (2013)
2,2-dimethylsuccinic acid	C <sub>6</sub> H <sub>10</sub> O <sub>4</sub>	146.2	0.67	Not observed	You et al. (2013)
2,5-hydroxybenzoic acid	C <sub>7</sub> H <sub>6</sub> O <sub>4</sub>	154.2	0.57	Not observed	You et al. (2013)
Diethylmalonic acid	C <sub>7</sub> H <sub>12</sub> O <sub>4</sub>	160.2	0.57	89.2 ± 4.2	You et al. (2013)
3,3-dimethylglutaric acid	C <sub>7</sub> H <sub>12</sub> O <sub>4</sub>	160.2	0.57	89.1 ± 6.9	You et al. (2013)
Levogluconan	C <sub>6</sub> H <sub>10</sub> O <sub>5</sub>	162.1	0.83	Not observed	You et al. (2013)
Glucose	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	180.2	1	Not observed	Current study
Suberic acid monomethyl ester	C <sub>9</sub> H <sub>16</sub> O <sub>4</sub>	188.2	0.44	100 ± 6.2	You et al. (2013)
Citric acid	C <sub>6</sub> H <sub>8</sub> O <sub>7</sub>	192.1	1.17	Not observed	You et al. (2013)
α,4-dihydroxy-3-methoxybenzeneacetic acid	C <sub>9</sub> H <sub>10</sub> O <sub>5</sub>	198.2	0.56	72.6 ± 6.3	You et al. (2013)
Poly(ethylene glycol)	C <sub>2n</sub> H <sub>4n+2</sub> O <sub>n+1</sub>	200	0.63	79.8 ± 7.4	You et al. (2013)
Diethyl-L-tartrate	C <sub>8</sub> H <sub>14</sub> O <sub>6</sub>	206.2	0.75	90.2 ± 6.6	You et al. (2013)
Poly(ethylene glycol) bis(carboxymethyl) ether	C <sub>2n+4</sub> H <sub>4n+6</sub> O <sub>n+5</sub>	250	0.83	67.6 ± 6.2	Current study
Diethyl sabacate	C <sub>14</sub> H <sub>26</sub> O <sub>4</sub>	258.4	0.29	100 ± 6.2	You et al. (2013)
Poly(ethylene glycol)	C <sub>2n</sub> H <sub>4n+2</sub> O <sub>n+1</sub>	300	0.58	86.7 ± 6.4	You et al. (2013)

Sucrose	$C_{12}H_{22}O_{11}$	342.3	0.92	Not observed	Current study
Poly(ethylene glycol)	$C_{2n}H_{4n+2}O_{n+1}$	400	0.56	$88.3 \pm 6.6$	Current study
Poly(propylene glycol)	$C_{3n}H_{6n+2}O_{n+1}$	425	0.38	$94.1 \pm 6.7$	You et al. (2013)
Poly(ethylene glycol) diacrylate	$C_{2n+6}H_{4n+6}O_{n+3}$	575	0.5	$94.7 \pm 6.2$	You et al. (2013)
Ouabain	$C_{29}H_{44}O_{12}$	584	0.41	$90.1 \pm 6.8$	Current study
Raffinose	$C_{18}H_{32}O_{16}$	594.5	0.89	Not observed	Current study
Poly(ethylene glycol)	$C_{2n}H_{4n+2}O_{n+1}$	600	0.54	$89.5 \pm 6.3$	Current study
Poly(ethylene glycol) bis(carboxymethyl) ether	$C_{2n+4}H_{4n+6}O_{n+5}$	600	0.63	$92.0 \pm 6.3$	You et al. (2013)
Maltopentaose	$C_{30}H_{52}O_{26}$	829	0.87	Not observed	Current study
Poly(ethylene glycol)	$C_{2n}H_{4n+2}O_{n+1}$	900	0.53	$92.9 \pm 6.4$	Current study
Maltoheptaose	$C_{42}H_{72}O_{36}$	1153	0.86	Not observed	Current study

724

725




726 **Table 4.** Combined data set used to determine the effect of temperature on SRH. This includes  
 727 the measurements at  $244 \pm 1$  K,  $263 \pm 1$  K and  $278 \pm 1$  K in the current studies (Table 2) and  
 728 results from You et al. (2013) at  $290 \pm 1$  K. OIR =  $2.0 \pm 0.1$  in all the experiments. Uncertainties  
 729 are 95% confidence intervals considering  $\sigma$  of multiple SRH measurements and the uncertainty  
 730 from the calibration.

Compound	O:C	SRH (%) at different temperatures			
		$244 \pm 1$ K	$263 \pm 1$ K	$278 \pm 1$ K	$290 \pm 1$ K
2,5-hexanediol	0.33	$84.0 \pm 8.3$	$84.6 \pm 6.5$	$88.0 \pm 6.2$	$88.8 \pm 7.1^a$
Poly(propylene glycol)	0.38	$87.7 \pm 6.5$	$89.4 \pm 6.7$	$92.5 \pm 6.2$	$94.1 \pm 6.7^a$
Poly(ethylene glycol) diacrylate	0.5	$88.5 \pm 7.0$	$90.4 \pm 7.2$	$95.0 \pm 6.2$	$94.7 \pm 6.2$
Poly(ethylene glycol) 900	0.53	$89.2 \pm 9.2$	$88.7 \pm 6.2$	$91.8 \pm 6.3$	$92.9 \pm 6.4^a$
$\alpha$ ,4-dihydroxy-3-methoxybenzeneacetic acid	0.56	$76.0 \pm 6.3$	$67.3 \pm 8.7$	$72.3 \pm 6.4$	$72.6 \pm 6.3^a$
Diethylmalonic acid	0.57	$89.8 \pm 5.6$	$87.0 \pm 6.7$	$88.6 \pm 6.3$	$89.2 \pm 4.2^a$
3,3-dimethylglutaric acid	0.57	$98.2 \pm 4.3$	$88.5 \pm 6.3$	$88.5 \pm 6.3$	$89.1 \pm 6.9^a$
2,5-hydroxybenzoic acid	0.57	Not observed	Not observed	Not observed	Not observed <sup>a</sup>
Poly(ethylene glycol) 300	0.58	$85.6 \pm 9.7$	$83.6 \pm 6.5$	$85.6 \pm 6.4$	$87.6 \pm 6.4^a$
Poly(ethylene glycol) 200	0.63	$71.3 \pm 6.2$	$77.2 \pm 6.3$	$79.7 \pm 6.2$	$79.8 \pm 7.4^a$
Poly(ethylene glycol) bis(carboxymethyl) ether 600	0.63	$87.8 \pm 7.1$	$90.4 \pm 6.2$	$90.4 \pm 6.6$	$92.0 \pm 6.3^a$
2-methylglutaric acid	0.67	$76.7 \pm 6.6$	$76.2 \pm 6.3$	$76.6 \pm 6.2$	$75.3 \pm 6.4^a$
2,2-dimethylsuccinic acid	0.67	Not observed	Not observed	Not observed	Not observed <sup>a</sup>
Diethyl-L-tartrate	0.75	$86.9 \pm 7.3$	$85.0 \pm 6.5$	$87.4 \pm 4.1$	$90.2 \pm 6.6^a$
Glutaric acid	0.8	Not observed	Not observed	Not observed	Not observed <sup>a</sup>
Levogluconan	0.83	Not observed	Not observed	Not observed	Not observed <sup>a</sup>
Raffinose	0.89	Not observed	Not observed	Not observed	Not observed
Glycerol	1	Not observed	Not observed	Not observed	Not observed <sup>a</sup>
Citric acid	1.17	Not observed	Not observed	Not observed	Not observed <sup>a</sup>
Malonic acid	1.33	Not observed	Not observed	Not observed	Not observed <sup>a</sup>

731

732 <sup>a</sup> Data taken from You et al. (2013). All other data are from this study


733



735


736

737 **Figure 1** The effect of molecular weight and O:C of the organic species on the occurrence of  
738 liquid-liquid phase separation in mixed organic-ammonium sulfate particles ( $\text{OIR} = 2.0 \pm 0.1$ ).  
739 Data plotted are from the current study and You et al. (2013), and are summarized in Table 3.  
740 Open circles indicate liquid-liquid phase separation was observed, while stars indicate liquid-  
741 liquid phase separation was not observed. The orange hatched region corresponds to the  
742 molecular weight and O:C of the organic species when liquid-liquid phase separation was always  
743 observed, and the green hatched region corresponds to the molecular weight and O:C of the  
744 organic species when liquid-liquid phase separation was never observed.  
745



747  
 748 **Figure 2.** SRH as a function of molecular weight of organic species in the particles at  $290 \pm 1$  K.  
 749 The SRH results are from the current study and You et al. (2013) (see Table 3). The colors  
 750 represent the O:C of different organic species. Squares represent SRH of particles in which  
 751 liquid-liquid phase separation was observed. Bars for the squares are 95% confidence intervals  
 752 considering  $\sigma$  of multiple SRH measurements and the uncertainty from the calibration. Stars  
 753 indicate that liquid-liquid phase separation was not observed. OIR =  $2.0 \pm 0.1$  in all the  
 754 experiments.


755



756  
 757  
 758  
 759  
 760  
 761  
 762  
 763  
 764  
 765


**Figure 3.** SRH as a function of O:C of the organic species at  $290 \pm 1$  K. The SRH results are from the current study and You et al. (2013) (see Table 3). The colors represent the molecular weight of the different organic species. Squares represent the SRH of particles in which liquid-liquid phase separation was observed. Bars for the squares are 95% confidence intervals considering  $\sigma$  of multiple SRH measurements and the uncertainty from the calibration. Stars indicate liquid-liquid phase separation was not observed. OIR =  $2.0 \pm 0.1$  in all the experiments.






766  
 767 **Figure 4.** A SRH of mixed organic-ammonium sulfate particles as a function of O:C measured at  
 768 four different temperatures. Different symbols represent the different temperatures. Bars for the  
 769 data are 95% confidence intervals considering  $\sigma$  of multiple SRH measurements and the  
 770 uncertainty from the calibration. Data at SRH= 0% indicate liquid-liquid phase separation was  
 771 not observed. Data plotted are summarized in Table 4. The OIR =  $2.0 \pm 0.1$  in all the  
 772 experiments. B Range of the average O:C of organic material in particles from measurements at  
 773 many locations in the Northern Hemisphere and the Amazon.

774



775  
 776 **Figure 5.** Summary of SRH as a function of temperature for twelve types of mixed organic-  
 777 ammonium sulfate particles which underwent liquid-liquid phase separation. Data plotted were  
 778 taken from Table 4. Bars for the data are 95% confidence intervals considering  $\sigma$  of multiple  
 779 SRH measurements and the uncertainty from the calibration. Colors represent the O:C values of  
 780 different organic species in the particles. OIR =  $2.0 \pm 0.1$  in all the experiments.

781



782

783 **Figure 6** SRH as a function of temperature for particles containing  $\alpha,4$ -dihydroxy-3-  
784 methoxybenzeneacetic acid mixed with ammonium bisulfate. Bars for the data are 95%  
785 confidence intervals considering  $\sigma$  of multiple SRH measurements and the uncertainty from the  
786 calibration. Data at  $290 \pm 1$  K was taken from You et al. 2013.

787