



Vertical structure of
BrO

P. K. Peterson et al.

Meteorological controls on the vertical distribution of bromine monoxide in the lower troposphere

P. K. Peterson¹, W. R. Simpson¹, K. A. Pratt^{2,*}, P. B. Shepson^{2,3}, U. Frieß⁴, J. Zielcke⁴, U. Platt⁴, S. J. Walsh¹, and S. V. Nghiem⁵

¹Department of Chemistry and Biochemistry, Geophysical Institute, University of Alaska Fairbanks, Fairbanks, Alaska, USA

²Department of Chemistry, Purdue University, West Lafayette, Indiana, USA

³Department of Earth, Atmospheric and Planetary Sciences, Purdue University, West Lafayette, Indiana, USA

⁴Institute for Environmental Physics, University of Heidelberg, Germany

⁵Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

* now at: Department of Chemistry and Department of Earth & Environmental Sciences, University of Michigan, Ann Arbor, Michigan, USA

Received: 26 August 2014 – Accepted: 2 September 2014 – Published: 16 September 2014

Correspondence to: W. R. Simpson (wrsimpson@alaska.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Multiple axis differential absorption spectroscopy (MAX-DOAS) measurements of bromine monoxide (BrO) probed the vertical structure of halogen activation events during March–May 2012 at Barrow, Alaska. An analysis of the BrO averaging kernels and degrees of freedom obtained by optimal-estimation-based inversions from raw MAX-DOAS measurements reveals the information is best represented by reducing the retrieved BrO profile to two quantities, the integrated column from the surface through 200 m (VCD_{200m}), and the lower tropospheric vertical column density (LT-VCD) which represents the integrated column of BrO from the surface through 2 km. The percentage of lower-tropospheric BrO in the lowest 200 m was found to be highly variable ranging from shallow layer events, where BrO is present primarily in the lowest 200 m to distributed column events where BrO is observed at higher altitudes. The highest observed LT-VCD events occurred when BrO was distributed throughout the lower troposphere, rather than concentrated near the surface. Atmospheric stability in the lowest 200 m influenced the percentage of LT-VCD that is in the lowest 200 m, with inverted temperature structures having a first-to-third quartile range (Q1–Q3) of $VCD_{200m}/LT-VCD$ from 15–39% while near neutral temperature structures had a Q1–Q3 range of 7–13%. Data from this campaign show no clear influence of wind speed on either lower-tropospheric bromine activation (LT-VCD) or the vertical distribution of BrO, while examination of seasonal trends and the temperature dependence of the vertical distribution supported the conclusion that the atmospheric stability affects the vertical distribution of BrO.

1 Introduction

The seasonal return of sunlight during late winter in the polar regions is associated with production of reactive halogens (e.g. Br, BrO, Cl) from saline ice surfaces (Abbatt et al., 2012; Saiz-Lopez and von Glasow, 2012). These halogen species influence boundary

ACPD

14, 23949–23994, 2014

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



layer chemistry through phenomena such as boundary layer ozone depletion events (ODEs) (Barrie et al., 1988; Simpson et al., 2007b) and mercury deposition events (MDEs) (Schroeder et al., 1998; Steffen et al., 2008). While the production mechanism of these halogens is not fully understood, modeling (e.g. Fan and Jacob, 1992; Mozurkewich, 1995; Lehrer et al., 2004; Piot and von Glasow, 2007; Yang et al., 2010; Parrella et al., 2012; Toyota et al., 2014), laboratory (e.g. Fickert et al., 1999; Huff and Abbatt, 2000; Wren et al., 2013) and field studies (e.g. Foster et al., 2001; Spicer et al., 2002; Toom-Sauntry and Barrie, 2002; Krnavek et al., 2012; Pratt et al., 2013; Liao et al., 2014) point to heterogeneous chemistry involving salts present on ice surfaces (e.g. NaBr, NaCl) as important sources of reactive halogens.

Multiple studies have examined the vertical extent of ODEs that result from halogen chemistry and have shown ODE vertical structure to be highly variable (Bottenheim et al., 2002b; Jones et al., 2010; Helmig et al., 2012; Oltmans et al., 2012). Jones et al. (2010) suggest a link between ozone depletion at higher altitudes and low pressure systems (storms) from Antarctic observations, but also point out that this relationship may not be the case in the Arctic due to differing meteorology. Ozonesonde data from ARCTAS/ARCPAC and OASIS campaigns presented in Oltmans et al. (2012) show the vertical extent of ozone depletion varies between 200 and 1000 m, and that this difference may be tied to the origin of the observed air masses in addition to the local meteorology. Although the profile of ozone depletion gives some insights into the vertical structures of the reactive halogens that deplete ozone, the differing chemical lifetimes of ozone (a day to longer) and reactive halogens (minutes to hour unless recycled by heterogeneous chemistry on particles) may mean that halogen vertical profiles differ from ozone depletion profiles. Tackett et al. (2007) present a view from tethered balloon measurements of a stratified lower boundary layer, in which the rates of ozone and Hg depletion are determined by downward mixing into the halogen-chemistry-active surface layer. Moore et al. (2014) found that both ozone and mercury levels recovered to higher levels when vertical mixing induced by sea ice leads had recently influenced the observed airmasses, but the effect of this mixing on reactive halogens

is not known. The work of Frieß et al. (2011) used MAX-DOAS observations along with optimal-estimation-based inversions to retrieve full BrO profiles and showed significant variability in vertical structure with reactive halogens typically being in the lowest ~ 1 km of the atmosphere, similar to the ozone depleted layer.

5 The physical mechanisms for the transport of reactive halogens from the snow pack to the boundary layer and recycling of those halogens aloft are still poorly understood. Given the short lifetime of BrO in the absence of heterogeneous recycling (McConnell et al., 1992; Platt and Hönninger, 2003), it is likely that some sort of aerosol particles (Fan and Jacob, 1992), either from blowing snow or other sources are required to sus-
10 tain halogen activation aloft. Recent literature in particular has focused on the role of blowing snow events in halogen activation (e.g. Jones et al., 2009, 2010; Yang et al., 2010; Frieß et al., 2011). In particular, Jones et al. (2009) used O₃ observations at Hal-
ley station in Antarctica, to derive a model for the dependence of ODEs on wind speed. Jones et al. (2009) suggest that there are two environmental regimes that favor local
15 halogen activation and subsequent ozone depletion. One regime consists of low wind speeds and a stable boundary layer, the other requires high winds and the presence of blowing snow. Blowing snow events occur when high winds loft snow, potentially enhancing available snow surface area available for reaction. Blowing snow may shift
20 reactive halogens aloft through reactions on the blowing snow or aerosol particles produced by sublimed blowing snow (Jones et al., 2009, 2010; Yang et al., 2010; Frieß et al., 2011). Comparison of satellite-observed column BrO data with modeling results by Yang et al. (2010) suggests that blowing snow events may be sufficient to explain the majority of halogen activation events observed via satellite. In contrast, Halfacre et al. (2014) conducted an analysis at a variety of locations across the Arctic and found
25 no relationship between observed ozone and wind speed, suggesting that high wind speeds, and consequently blowing snow, may not play a major role in ozone depletion. Again, because of the differing chemical lifetimes of ozone and reactive halogens, using ozone depletion as a proxy for halogen activation requires caution.

**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To examine these ideas further, it is necessary to observe both the total amount of reactive halogens, and their distribution throughout the lower troposphere. While satellite measurements (e.g. Wagner and Platt, 1998; Richter et al., 1998; Chance, 1998) provide extensive spatial coverage of the Arctic, they only measure the total column of BrO, and do not provide vertical profile or near surface information. Multiple methods exist to separate total column measurements into stratospheric and tropospheric components (e.g. Theys et al., 2011; Salawitch et al., 2010; Koo et al., 2012; Sihler et al., 2012). However, discrepancies still exist between ground based and satellite based measurements, potentially due to clouds masking portions of the tropospheric BrO column from satellite views. Additionally, Theys et al. (2009) and Salawitch et al. (2010) showed that some of the variability of total column BrO arises from stratospheric variability, thus not all satellite-detected BrO “hotspots” are actually lower tropospheric halogen activation events. In-situ chemical ionization mass spectrometry (e.g. Liao et al., 2011; Pratt et al., 2013; Liao et al., 2014) and long path differential optical absorption spectroscopy (LP-DOAS) (e.g. Tuckermann et al., 1997; Pöhler et al., 2010) fail to capture the vertical extent of halogen activation, except in the case of limited aircraft-based campaigns (e.g. Neuman et al., 2010; Liao et al., 2012), and thus can not address questions about the total amount of halogen activation taking place at a given time. However, the Tackett et al. (2007) work implied that much of the halogen chemistry is occurring in the lowest ~ 200 m of the boundary layer.

Multiple axis differential optical absorption spectroscopy (MAX-DOAS) has the ability to retrieve vertical profiles of BrO in the lowest few kilometers, and thus can examine the variability of the vertical distribution of BrO over time (Frieß et al., 2011). The present work builds on the method of Frieß et al. (2011) by considering the BrO averaging kernels, which provide information on the sensitivity of these BrO measurements (Payne et al., 2009), to identify what observations are independent of the a priori assumed profile of BrO and creating a consistent timeseries of these data over a field campaign. Through this analysis, we identify two relevant properties of the BrO vertical profile, the partial vertical column density (VCD) of BrO in the lowest 200 m (termed VCD_{200m})

**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and the lower-tropospheric partial VCD (termed LT-VCD). The present work explores the dependence of these BrO profile properties on environmental factors to investigate activation of BrO and its propagation outside of the near surface layer over time. This present work focuses on measurements occurring near Barrow, Alaska, during the Bromine, Ozone, and Mercury Experiment (BROMEX) campaign (Nghiem et al., 2013) from 6 March to 15 May 2012. The primary MAX-DOAS observations were made from the roof of the Barrow Arctic Research Center (BARC) building at 71.325° N, 156.668° W, which is about 6 km northeast of the city of Barrow, Alaska.

2 Methods

2.1 MAX-DOAS measurements

Multiple-Axis Differential Optical Absorption Spectroscopy (MAX-DOAS) probes vertical distributions of trace gases above a measurement site through a combination of measurements of scattered sunlight and modeling of vertical profiles consistent with those observations (Hönninger et al., 2004). The instrument measures spectra of scattered sunlight in a region where the molecule of interest absorbs as a function of elevation angle of a telescope receiving the light. Measurements at all elevation angles contain absorption features due to stratospheric and higher tropospheric gases, and because most sunlight is scattered in the lower atmosphere, these stratospheric absorption features are common to observations at all elevation angles. Measurements at low elevation angles contain enhanced absorption for gases present near the ground, due to the tangent geometry of the light's final path to enter the telescope at a low elevation. Therefore, analysis of the relative absorption spectrum comparing light at low elevation angles to high elevation angles provides high sensitivity to near-ground absorbers with greatly reduced sensitivity to stratospheric and upper tropospheric gases.

The instrument used in this study is similar to the one described in Carlson et al. (2010) with a few improvements. In this study, we measured scattered sunlight us-

**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing a QE-65000 (Ocean Optics) spectrometer with spectral range from 309–397 nm and optical resolution of 0.39 nm full width at half maximum. This new spectrometer is improved by using a thermoelectrically cooled detector, stabilized at -15°C . The spectrometer optical bench is heated to 38°C to improve optical alignment stability.

Another improvement of the instrument is the use of a MEMS tiltmeter (SignalQuest SQ-SI-360DA-3.3R-HMP-HP-IND-S) mounted on the moving telescope that directly measures the elevation angle of the observation. The scan pattern measures scattered sunlight at elevation angles of 90° (zenith), 20° , 10° , 5° , 2° , and 1° over a period of approximately 30 min.

We used QDOAS software (Fayt et al., 2011) to fit the relative absorption spectrum between near temporally coincident low elevation and zenith observations to a linear combination of possible absorbing spectra to quantify the differential slant column density (dSCD) of each absorbing gas at each elevation angle. We performed this analysis in the wavelength window between 346 and 364 nm. The absorber cross sections are detailed in Table 1 and are convoluted with an instrumental function measured from the 334 nm line of a low pressure mercury lamp. In addition to gaseous absorbers, we include a third-order polynomial to account for Rayleigh and Mie scattering by gases and particles in the atmosphere, and a spectral offset to account for stray light within the spectrometer. The mean residual root mean square (RMS) of our dSCD retrieval was 3.9×10^{-4} . Data collected during low sight conditions (solar zenith angle $> 85^{\circ}$), $\text{RMS} > 10^{-3}$, or when frost is detected on the instrument's optical window are discarded. The dSCD fitting errors (1σ) over the campaign average 1.9×10^{13} molecules cm^{-2} for BrO and 5.1×10^{41} molecules 2 cm^{-5} for O_4 .

2.2 Retrieval of vertical profiles

Retrieval of trace gas vertical profiles from dSCD data is a two-step procedure. First, we determine the aerosol particle extinction vertical profile from O_4 dSCD measurements using an optimal estimation procedure detailed in Frieß et al. (2006) and briefly described here. The observed O_4 dSCD values are strongly dependent on atmospheric

**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



visibility and light scattering, which we are able to deduce because we know the vertical profile of O_4 . Measured O_4 dSCDs are compared to values modeled using the SCIA-TRAN radiative transfer model (Rozanov et al., 2005), which depend on a variable aerosol particle extinction profile and static parameters such as the viewing geometry and aerosol particle light scattering properties. Because the measurement data alone do not provide enough information to retrieve the profile, an assumed a priori profile is required to further constrain the solution, as described by Rodgers (2000). We used a variable a priori profile that peaks at the surface and exponentially decays with altitude. Initially, the profile peaks at the surface with an aerosol particle extinction value of 0.05 km^{-1} and exponentially decays with a scale height of 1 km. In following iterations, the a priori is set to the retrieved profile and smoothed using a boxcar average with a width of 300 m (Frieß et al., 2006). The final output of this first step is a vertical profile of aerosol particle extinction.

The procedure to retrieve the BrO profile from BrO dSCD measurements is described in detail in Frieß et al. (2011). Briefly, we used the above-retrieved aerosol particle extinction profiles as input to a Monte Carlo radiative transfer model (McArtim, Deutschmann et al., 2011) that simulates BrO dSCDs as a function of a variable BrO vertical concentration profile. The vertical profile of BrO is varied to give best fit to the BrO dSCD observations, again by optimal estimation. In the case of BrO, the a priori profile peaks at the surface with a value of 10 pmol mol^{-1} and exponentially decays with a scale height of 400 m. The result of the second step is the average BrO mixing ratio every 100 m from the ground to an altitude of two kilometers. An example of the retrieved profile is shown in green in Fig. 1. Note that while our general retrieval uses a 400 m scale height for the a priori profile of BrO, Fig. 1 also includes the effect of modifying the a priori profile to have a different (1000 m) scale height (dashed blue line), as described below.

2.3 Information content of the profile

The retrieved BrO profiles contain average mixing ratios in twenty 100 m thick layers from the surface to 2 km; however, they are based on only five low-elevation spectral measurements of BrO dSCD. Therefore, it is clear that our observations contain less information about the true vertical profile than is represented in Fig. 1. The optimal estimation method used the assumed a priori information to constrain the fit in the absence of further information, but the actual vertical profile of BrO is not known. In fact, the results of this study indicate that both the BrO abundance and vertical profile shape vary over time. Therefore, the full profile, as retrieved by optimal estimation, lends itself to overinterpretation of the observations and we seek to find a method to determine the actual information content, as well as reduce the number of derived properties to more accurately reflect the true observations.

As described in Rodgers (2000), the retrieved profile contains true profile information as well as a bias toward the assumed a priori profile. The extent to which the retrieved profile depends on the dSCD measurement data is determined by examining the averaging kernel matrix. An example of the BrO averaging kernels for a MAX-DOAS retrieval is given in the left panel of Fig. 2. If the value of the diagonal element of the averaging kernel at some altitude were unity (1), it would indicate that all of the information content at that layer is constrained by observational data and none of the information at that altitude comes from the a priori assumption. Non-zero off-diagonal elements indicate that the retrieved parameter in that layer is not independently resolved and is influenced by results in other layers. There are two peaks in the averaging kernels, one at the surface and one aloft at an altitude of about 300 m. We can understand the sharp peak at the surface easily because MAX-DOAS spectroscopy always has highest sensitivity and greatest vertical resolution in the layer in which the observer resides. The broader peak aloft, centered at about 300 m, in this example, comes from the observations at higher elevation angles, which view through those aloft layers, but do not provide enough information to independently resolve mixing ratios in 100 m thick

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



layers at higher altitudes. The trace (sum of diagonal elements) of the averaging kernel matrix gives the degrees of freedom (DOF), which represents the total number of independent pieces of information contained in the retrieval. For BrO retrieval using this technique and our instrumentation, the distribution of DOF is well fit by a Gaussian function, with a mean of 2.0 DOF and $\sigma = 0.27$. Our retrievals also exhibit a distinct lack of sensitivity to changes in individual 100 m layers at altitudes over 1000 m. The DOF for this example is 1.99, implying our retrieved profile contains two independent pieces of information, rather than the 20 we retrieved.

To illustrate the influence of our a priori assumptions, Fig. 1 also shows a retrieved profile with a different assumed BrO scale height, 1000 m instead of the normal 400 m assumption. The influence of a priori selection in the retrieved BrO mixing ratio in the lowest 200 m is minimal, showing that our instrument is most independent of a priori assumptions about the BrO profile near the surface. However, aloft the effects of a priori selection are clearly visible. What appeared to be a relatively sharp peak at ≈ 650 m when the 400 m a priori was assumed becomes much broader with the 1000 m assumed scale height. Note that both optimally estimated results fit the observational data with similar figure of merit (χ^2), so the difference in profile is not due to poor fitting in one case or the other, but instead a consequence of the a priori influencing the resultant BrO profile.

2.4 Reduction of the full profile

A method for mapping the BrO profile retrieved on a fine grid to a course grid, such that the DOF are represented as physical quantities without the influence of a priori information is outlined in von Clarmann and Grabowski (2007). They suggest moving to a variable coarse altitude grid where one would select grid points such that each grid point would represent one DOF. The lowest layer's upper boundary would be determined by summing the diagonal elements of the BrO averaging kernel matrix until the sum reached one. This process would be repeated to obtain additional layer boundary altitudes. However, this approach is not practical for long-term time series analysis

**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



because the vertical resolution would vary over time, making comparisons difficult. As a compromise, we calculated coarse grid points by examining the ensemble of profile retrievals over this campaign and splitting them into two layers by determining the average altitudes such that each layer contains one degree of freedom. This method led us to represent one degree of freedom as the abundance of BrO from surface through 200 m, and another as the abundance of BrO from 200 m to 2.0 km. These two layers differ in thickness due to the enhanced vertical resolution at the surface and decreased vertical resolution aloft, so we choose to represent the abundance in each layer as a partial vertical column density (VCD) within each layer. The vertical column density is also useful because satellite total column retrievals of BrO use the same unit of abundance. Thus, our two quantities ($VCD_{200\text{m}}$ and VCD_{Aloft}) represent a surface layer VCD and a lofted layer VCD respectively.

Figure 1 shows the partial VCDs in each layer derived from the grid coarsening procedure, for an example retrieval. While the profiles look fairly different, particularly aloft, the partial VCDs in each of the layers are only around ten percent different from each other as an effect of the assumed a priori profile. This example shows how the grid coarsening procedure gives results that are less dependent on the unknown but assumed BrO profile.

Payne et al. (2009) describe how to calculate the BrO averaging kernel for the coarsened grid, and an example of the coarse grid BrO averaging kernels is shown in Fig. 2. The DOF has been reduced, meaning we have lost some information in the transfer to a coarse grid, however, both parameters have averaging kernels that peak within the desired layer and at a value close to unity, implying most of the information comes from the observations. Payne et al. (2009) suggested a cutoff of 0.7 for the diagonal element of the averaging kernel within a layer as sufficiently close to unity to consider the observations as derived from the data and independent of the a priori assumption. We chose to adopt the same cutoff of 0.7 degrees of freedom in the present work.

The right panel of Fig. 3 shows the dependence of the aloft information content on the aerosol optical depth. As Fig. 3 shows, our ability to retrieve information beyond

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the near surface layer is heavily dependent on the aerosol optical depth, while our ability to retrieve information near the surface is less influenced by aerosol particles. In addition to the degrees of freedom cutoff, over the course of the campaign we also observed limited times when the retrieved aerosol optical depth was large, in excess of 2, which indicates that the visibility is insufficient to obtain an accurate profile of aerosol particle extinction or BrO. During these periods, we do not consider the VCD_{Aloft} due to insufficient degrees of freedom, and we also disregard the VCD_{200m} . This approach is consistent with methods used in comparison of near-surface MAX-DOAS and LP-DOAS BrO measurements by Frieß et al. (2011).

When both quantities have averaging kernels peaking about 0.7 or higher, we refer to the sum as the lower tropospheric vertical column density (LT-VCD). It is important to note that this quantity does not represent a new degree of freedom, as it is derived from and not independent of the other two measurements. To evaluate the vertical distribution of BrO, we examine the ratio of the $VCD_{200m}/LT-VCD$ as well as the total amount of observed activation in the lower tropospheric (LT-VCD) and near surface amounts (VCD_{200m}). A time series of these three quantities, along with the aerosol optical depth over a portion of the campaign is presented in Fig. 4. The rest of the timeseries can be found in the supplemental material. This timeseries will be discussed below, but here we note that the fraction of BrO in the lower layer varies between a few percent and up to $\approx 80\%$. The a priori profile, exponentially decaying with a scale height of 400 m, would have a constant fraction of BrO in the lowest 200 m, approximately 34%. It is clear that the data show substantially more vertical profile variability than the assumed profile. Therefore, we use the grid-coarsened data to represent the information content from our MAX-DOAS observations over this long time period.

2.5 Other field sites

Figure 5 shows locations for various measurements used in this study. While this paper focuses on MAX-DOAS measurements from the BARC building, Sect. 3.3 makes use of data from multiple field sites to evaluate the effectiveness of our near-surface BrO

3 Results

3.1 Relationship between visibility and retrieval information content

The ability of MAX-DOAS to observe BrO beyond the near surface layer is heavily influenced by the optical properties of the atmosphere (e.g. visibility) at the time of the measurement. The methods outlined above allow us to identify times when MAX-DOAS measurements are sensitive to BrO beyond the near surface layer and restrict our further analysis of the vertical distribution of BrO to these periods. Figure 3 details the relationship between the aerosol optical depth and the information content of the measurement. As the aerosol optical depth increases, the ability to observe the full LT-VCD sharply decreases. As seen in Fig. 3, we can retrieve the full LT-VCD 50% of the time. We excluded periods where we are unable to fully observe the LT-VCD from further analysis of the vertical distribution of BrO.

3.2 Influence of ozone

Ambient ozone levels influence the partitioning of reactive halogen species between atomic (e.g. Br atoms) and oxide (e.g. BrO) forms. MAX-DOAS measurements observe only BrO and thus only constrain one component of BrO_x ($\text{BrO} + \text{Br}$). Therefore, it is important to consider effects of gas-phase partitioning of BrO_x on subsequent conclusions about halogen activation. To examine this influence, we examined ozone data and meteorological data from Barrow, provided by the National Oceanic and Atmospheric Administration (NOAA) Earth Systems Research Laboratory/Global Monitoring Division (ESRL/GMD). Under ordinary ozone conditions (ozone mixing ratios of $30\text{--}40\text{ nmol mol}^{-1}$), BrO will make up the majority of BrO_x ; however, as ozone levels approach zero (a common occurrence during the ozone depletion event season), our BrO-only measurements could be insufficient to observe reactive halogens, as shown in Helmig et al. (2012). We term this effect “ozone titration” and seek to avoid interpretation of BrO data below some threshold where the effect dominates. The threshold

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



depends on the photolysis rate of BrO, which varies depending on cloud cover and solar zenith angle, among other factors. For the purposes of this study, we used an ozone threshold of $< 1 \text{ nmol mol}^{-1}$ to identify potentially titrated air masses to exclude from further analysis (Simpson et al., 2007a). Figure 4 shows times when ozone is below 1 nmol mol^{-1} as shaded in grey. Note that, for the most part, LT-VCD and $\text{VCD}_{200\text{m}}$ are small during ozone titration, as would be expected, and also observed by Neuman et al. (2010).

3.3 Performance of surface BrO retrieval

To evaluate the effectiveness of the BrO retrieval near the surface, we used in situ BrO measurements made using chemical ionization mass spectrometry (CIMS) to evaluate our retrieval in the lowest 100 m. While the mixing ratio in the lowest 100 m is not one of the grid coarsened quantities, because MAX-DOAS is most sensitive near the surface, the near surface mixing ratio has similar information content to the $\text{VCD}_{200\text{m}}$ while minimizing the effects of vertical averaging on comparison with in situ measurements. Figure 6, shows a timeseries of the average BrO mixing ratio retrieved in the lowest 100 m by MAX-DOAS (Red) and in situ CIMS measurements taken 1 m above the snow pack (Green). The error associated with 100 m mixing ratios retrieved using MAX-DOAS varies depending on the state of the atmosphere at the time of the measurement. The error associated with each retrieval is determined by examining the sensitivity of the retrieval to dSCD errors associated with the measurement data (Rodgers, 2000). Average errors (2σ) were $4.2 \text{ pmol mol}^{-1}$ for our MAX-DOAS measurements. Uncertainties are estimated at $30\% + 0.3 \text{ pmol mol}^{-1}$ for CIMS measurements.

Because CIMS measures in one discrete location just above the snowpack, while MAX-DOAS measurements are spatially averaged, both horizontally over 10–20 km and vertically over approximately 100 m, it is expected that large changes in the environment in the view direction of the MAX-DOAS will cause differences between the CIMS and MAX-DOAS observations. We also expect that given the snowpack source of Br_2 (Pratt et al., 2013), there may be a significant near-surface BrO gradient. Fig-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 6 shows a time series of near surface BrO measurements from the CIMS, as well as MAX-DOAS measurements at the BARC building. Since the BARC data originate at ~ 4 m, and average over the lowest 100 m, we might expect these data to be lower than the CIMS, which measures closer to the snowpack. Figure 6 shows agreement for the majority of concurrent observations, excluding the period shaded in gray, during which we observed substantial gradients in the near surface amounts of BrO at IL1 (blue line) and BARC (red line), which could explain discrepancies between the MAX-DOAS measurements at the BARC building and the CIMS measurements. Figure 7 shows the overall correlation between MAX-DOAS measurements at the BARC building and the CIMS, each averaged to hourly bins. Points occurring during the gray period marked on Fig. 6 are shown in red open circles. Outside times of disagreement between the two MAX-DOAS instruments, shown in gray on Fig. 6, the correlation observed between the DOAS data and CIMS data is $R = 0.70$, which is similar to correlation of MAX-DOAS and long path DOAS measurements observed by Frieß et al. (2011). The CIMS and LP-DOAS were previously shown to agree well, particularly during times of moderate wind speeds ($3\text{--}8\text{ m s}^{-1}$) and low NO ($< 100\text{ pmol mol}^{-1}$) (Liao et al., 2011).

3.4 Relationship between amount observed and vertical distribution of BrO

Figure 4 shows the timeseries of the $\text{VCD}_{200\text{m}}$ and LT-VCD. LT-VCD values over the course of this study ranged from 0 to 8×10^{13} molecules cm^{-2} with an average of 1.46×10^{13} molecules cm^{-2} and the top quartile being above 2.0×10^{13} molecules cm^{-2} , which we will later consider to be “highly activated” events. LT-VCD retrieval errors had a mean of 3.9×10^{12} molecules cm^{-2} over the course of the campaign. Figure 8 shows the percentage of the LT-VCD present in the lowest 200 m ranges from shallow layer events observed primarily below 200 m to distributed column events that are also observed at higher altitudes. The third panel of Fig. 4 shows a timeseries of this ratio. Surface layer percentages over this campaign ranged between 0 and 80 %, with the highest LT-VCD observed during distributed column events with lowest 200 m percentages of around 10 %. As the coloration of Fig. 8 indicates, high near surface mixing ratios do not always

imply high LT-VCD. In most cases, high LT-VCDs are associated with distributed column events.

3.5 Dependence on local meteorology

To examine atmospheric stability, we obtained meteorological balloon sounding data from the Barrow Airport (PABR) to calculate the temperature gradient (dT/dz) near the surface. We calculated the slope of a linear fit of the first three data points of the sounding. Typically, the first three sounding points corresponded to measurements between the surface and 200 m altitude. Soundings take place twice daily, at 00:00 UTC, which is in the daylight hours, 15 AKST, at Barrow, and 12:00 UTC. Because MAX-DOAS data are daylight only, to compare our half-hourly BrO measurements with the sounding data, we used daily averages of the fraction of near-surface BrO to LT-VCD from each day and examined the relationship between daily averaged ratios and the corresponding temperature gradients obtained from the local daytime sounding. Figure 9 shows the relationship between these two quantities. This box and whisker plot is generated by splitting daily observations into three bins of about 20 days each. The median is given by the red lines and shows the general trend that more stable atmospheres with inverted temperature structures (right bin) have more BrO present in the surface layer, while unstable atmospheres (left bin) tend to have BrO more distributed throughout the lower troposphere. The short diurnal time scales of the temperature gradient do not allow the use of daily sounding data to determine the functional relationship between the temperature gradient and where an event would fall on the spectrum of BrO vertical distributions.

The left panel of Fig. 10 illustrates the relationship between surface temperature and the amount of BrO in the near surface layer. On average, there is a higher fraction of BrO in the near surface layer when temperatures are below -25°C , with more distributed column BrO events being observed aloft at higher temperatures. Figure 10 (right panel) shows that activation events in excess of 2×10^{13} molecules cm^{-2} occur at various temperatures.

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vertical structure of
BrO**

P. K. Peterson et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Additionally, we examine the response of both the total activation as measured by the LT-VCD and the vertical distribution of BrO to changes in wind speed. Figure 11 (left panel) shows LT-VCD values in excess of 2×10^{13} molecules cm^{-2} at various wind speeds and, as shown by the red line, there is no obvious relationship between the LT-VCD and wind speed. The observed lack of dependence of the LT-VCD on wind speed and temperature is consistent with the observations of Halfacre et al. (2014). However, the vertical distribution of BrO is influenced by windspeed. The right panel of Fig. 11 shows the majority of shallow layer events occur during times when wind speeds are between $3\text{--}6 \text{ m s}^{-1}$. Additionally, events at wind speeds in excess of 8 m s^{-1} tend to be more distributed throughout the column.

4 Discussion

4.1 Role of atmospheric stability

The third panel of Fig. 4 shows the vertical structure of BrO was highly variable over the course of this study. Events range from shallow layer events where most of the observed BrO is found in the lowest 200 m to distributed column events with BrO present throughout the lower troposphere. Figure 8 shows the highest observed LT-VCDs were observed during distributed column events while shallow layer events, despite their enhanced near surface mixing ratios, did not result in large amounts of total activation, as measured by the LT-VCD. This points out the difficulty in comparing satellite BrO column retrievals with surface observations (e.g. Sihler et al., 2012). Figure 9 shows the vertical distribution of BrO is clearly influenced by the observed atmospheric stability. During times of inversion, bromine activation events tend toward the shallow layer end of the spectrum. When the atmosphere is less stable, events tend to be distributed throughout the column. Given that this study only uses daily sounding data, and knowing that convective mixing has a strong diurnal variability, it is likely this study represents a lower bound on the influence of atmospheric stability.

**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ozone depletion events have been shown to be highly variable in vertical extent (e.g. Bottenheim et al., 2002b; Tarasick and Bottenheim, 2002; Jones et al., 2010; Oltmans et al., 2012; Helmig et al., 2012). More recently, Moore et al. (2014) observed that opening of leads enhanced vertical mixing leading to a recovery of ozone and gaseous elemental mercury in the near surface layer. This mixing potentially impacts the vertical distribution of halogens as well. McElroy et al. (1999) observed high total tropospheric BrO events and interpreted them as arising from convective vertical mixing associated with open leads that brought near-surface BrO to the free troposphere. Therefore, it is reasonable that vertical mixing or hindered mixing due to atmospheric stability affects the vertical structure of BrO.

Seasonal and temperature trends are consistent with atmospheric stability's influence on the BrO vertical structure. Figure 10 shows that at lower temperatures, which are more prevalent early in the season, events tend to be observed in shallow layers and transition to more vertically distributed as the surface temperature increases. These observations are consistent with the idea that the distribution of BrO is primarily influenced by atmospheric stability.

The right panel of Fig. 10 indicates that significant activation events happen at a variety of temperatures. Pöhler et al. (2010) observed there was a linear trend of decreasing near surface maximum BrO, observed with LP-DOAS, as temperature increased up to -15°C . When examining the whole LT-VCD, rather than a near surface concentration, this trend does not appear, however, Fig. 10 (left panel) shows a distinct preference toward near surface distributions of BrO at lower temperatures, which as Fig. 8 shows, often contain higher near surface amounts of BrO, but have a lower overall LT-VCD. Figure 12 illustrates the diurnal behavior of both the LT-VCD, shown in blue, and the $\text{VCD}_{200\text{m}}$ shown in red. The $\text{VCD}_{200\text{m}}$ observed on Fig. 12 is similar to surface concentration data observed by Pöhler et al. (2010). The MAX-DOAS observations of the present study allow observation of the BrO LT-VCD, which Fig. 12 shows increases to a peak in the late afternoon, mirroring typical diurnal boundary-layer growth. Therefore, the diurnal pattern of the vertical profile of BrO appears to reflect boundary layer

dynamics. The observation of Pöhler et al. (2010) of decreasing surface BrO mixing ratio at higher temperatures could therefore reflect increased vertical mixing of surface-sourced BrO that dilutes the surface concentration upon warming rather than being indicative of decreased halogen activation chemistry at higher temperatures.

4.2 Role of wind speed

Figure 11 (left panel) shows that activation events with BrO LT-VCDs in the top quartile occur across a spectrum of wind speeds, rather than at just the low wind speeds associated with stable boundary layers, or high wind speeds associated with blowing snow events. The right panel of Fig. 11 shows a peak in shallow layer events between $3\text{--}6\text{ m s}^{-1}$ suggesting that some ventilation of the snowpack, which has been shown to be an effective source of Br₂ (Pratt et al., 2013), is required for the release of BrO during shallow layer events. For the most part, observed wind speeds exceeding 8 m s^{-1} , led to vertically distributed events, but beyond that there is little evidence to support the need for high winds or blowing snow to sustain halogen activation aloft, consistent with the conclusions of Halfacre et al. (2014). This suggests that, during the course of this study, high wind events and blowing snow were not required to observe significant halogen activation. Activation events in the top quartile were observed across a range of wind speeds, particularly at more moderate wind speeds, with 66 % of highly activated events occurring at wind speeds between 3 and 8 m s^{-1} . This distribution suggests that enhanced ventilation of the snowpack, and subsequent vertical mixing of snowpack-influenced airmasses, is likely responsible for the majority of elevated LT-VCDs observed over the course of this study.

Jones et al. (2009) and Yang et al. (2010) rely on satellite observations of BrO rather than ground-based measurements to draw conclusions about the role of wind in halogen activation events. The differing viewing geometry potentially becomes an issue when clouds are present, as shallow layer events occurring below clouds would potentially be masked from the satellite view, but observable with ground-based MAX-DOAS. In contrast, BrO above a cloud that would be potentially seen by a satellite would not

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be detected by ground-based MAX-DOAS, although the profile reduction procedure should minimize these issues. Sihler et al. (2012) observed that satellite measurements could underestimate the amount of BrO during shallow layer events, implying events with distributed column vertical structure are more likely to be satellite detectable. This idea that shallow layer events may not be satellite detectable is also supported by the modeling work of Toyota et al. (2014). Enhanced visibility of the larger in magnitude and more vertically distributed BrO events could potentially explain why satellite remote sensing of BrO may show an apparent relationship to wind speed. These differences should be considered when trying to resolve differences between ground-based and satellite-based measurements.

4.3 Relationship between activation and aerosol particles

The near-surface aerosol particle extinction retrieved from our MAX-DOAS measurements is a proxy for suspended particulate surface area that enhances recycling of halogens, allowing us to examine the relationship between BrO and aerosol particles. The left panel of Fig. 13 shows an increase in the VCD_{200m} with increased aerosol particle extinction, while the right panel shows that the relationship between aerosol particles and the LT-VCD is less clear. Laboratory studies (e.g. Huff and Abbatt, 2000; Wren et al., 2013) and field studies (e.g. Pratt et al., 2013) indicate that pH is an important control on heterogeneous recycling of halogens. Although we are unable to determine the pH of these particles by MAX-DOAS, we simply point out that, not only is the presence of suspended surface area important, but also that the chemical composition of that surface is likely a controlling factor.

Prior to using these data to consider the role of blowing snow in observed halogen activation during this study, it is important to point out that the lack of LT-VCD observations at high aerosol optical depth, shown in Fig. 3, could potentially lead to preferential exclusion of blowing snow events from further analysis of the total amount of activation as measured by the LT-VCD. To examine the potential impacts of data exclusions, we examined the amount of data at high wind speeds discarded by the

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



application of methods described in Sect. 2.4. Of the 147 measurements occurring when wind speeds exceeded 8 m s^{-1} threshold used for blowing snow in Jones et al. (2009), 86 of those were rejected due to insufficient information content. If one were to presume these were all blowing snow events and belonged in the top quartile of observed activation events, they would represent 9.6 % of LT-VCD observations and 29 % of events where the LT-VCD exceeded $2 \times 10^{13} \text{ molecules cm}^{-2}$. While we end up discarding a significant amount of data, these filtered data still provide a fair idea of the role of high wind events, blowing snow or otherwise, in halogen activation.

While the increase in $\text{VCD}_{200\text{m}}$ during times of high aerosol particle extinction supports blowing snow as one source of halogens, it is interesting to note that LT-VCD above $2 \times 10^{13} \text{ molecules cm}^{-2}$ often occur at wind speeds well below typical thresholds for blowing snow (8 m s^{-1} , Jones et al., 2009), as shown in the left panel of Fig. 11. Additionally, examining the dependence of the aerosol particle extinction on wind speed shown by the coloration in the left panel of Fig. 13, there is no apparent increase in extinction at high wind speeds, which we would expect if blowing snow was the only source of particulate surface area for halogen recycling. The observation of aerosol particle extinction values in excess of 10^{-1} km^{-1} across a variety of wind speeds suggests that blowing snow is not the only source of the aerosol particles needed to sustain halogen activation aloft. While blowing snow is not simply a univariate function of wind speed (Sturm and Stuefer, 2013), given 9 % of observations occurred when wind speeds exceeded 8 m s^{-1} and halogen activation events in excess of $2 \times 10^{13} \text{ molecules cm}^{-2}$ occurred 25 % of the time, immediate activation during blowing snow events is not the sole driver of halogen activation over the course of this study, and likely some other aerosol particle source is required. Frieß et al. (2011) postulated that aerosol particles produced by sublimation of blowing snow could transport and increase observed aerosol particle extinction while also providing a suspended surface for the recycling of halogens aloft. Further, in the Arctic springtime, Hara et al. (2002) observed bromide loss from coarse sea salt particles and addition to fine sulfate particles, supporting the suggestion of bromine recycling on aerosol particles.

5 Conclusions

The methods described in this paper outline the reduction of vertical profiles retrieved from MAX-DOAS observations using optimal estimation to produce two quantities, the lower tropospheric vertical column density (LT-VCD), and the near surface vertical column density in the lowest 200 m (VCD_{200m}), that appropriately reflect the information content of ground-based MAX-DOAS measurements for timeseries analysis. Consideration of BrO averaging kernels and degrees of freedom resulting from optimal estimation validates that these quantities are most appropriate to express the two degrees of freedom observed. This method allows for identification of time periods when MAX-DOAS measurements contain information about BrO beyond the near surface layer, and evaluate how the vertical structure of BrO responds to various environmental factors. Application of these methods shows retrieval of LT-VCDs by ground-based MAX-DOAS is highly dependent on visibility. During this study, we retrieved the LT-VCD 50 % of the time.

The vertical structure of BrO is highly variable, with the fraction of lower-tropospheric BrO in the lowest 200 m varying from near zero to nearly 80 %. The vertical distribution of activation events is clearly influenced by the atmospheric stability. This influence also manifests itself when examining the influence of temperature and seasonality. Later in the season, higher temperatures, and likely enhanced vertical mixing, tend to result in more BrO aloft than early in the season, where activation takes place in shallower layers. LT-VCD values in excess of 2×10^{13} molecules cm^{-2} occur across a variety of temperatures and wind speeds, but they are typically associated with distributed column events, rather than shallow layer events. Events that have larger observed near surface mixing ratios are typically of the shallow layer variety and do not typically have large amounts of BrO spread through the lower troposphere, potentially implying less overall ozone depletion and deposition of mercury from these events. This implies that both surface and upper profile measurements are necessary to appropriately identify

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the impact of environmental variables on O₃ depletion and Hg oxidation rates and extent.

High wind events were not common over the course of this study. While high winds did lead to some of the highest measured columns of BrO, given the low frequency of these events, high wind events, blowing snow or otherwise, are not the sole driver of halogen activation over the course of this study, suggesting that mechanisms requiring only moderate wind speeds (e.g. wind pumping) are important for halogen activation in the Arctic.

**The Supplement related to this article is available online at
doi:10.5194/acpd-14-23949-2014-supplement.**

Acknowledgements. The research at the University of Alaska was supported by the National Aeronautics and Space Administration (NASA) Cryospheric Sciences Program (CSP), and partial financial support for MAX-DOAS analysis methods was provided by the National Science Foundation under grant ARC-1023118. The Purdue group recognizes NSF support through grant ARC-1107695. K. A. Pratt was supported by a NSF Postdoctoral Fellowship in Polar Regions Research. The Purdue group acknowledges field assistance from Kyle Custard (Purdue Univ.), David Tanner (Georgia Tech), and L. Gregory Huey (Georgia Tech). The research at the Jet Propulsion Laboratory, California Institute of Technology, was supported by the NASA CSP. The authors gratefully acknowledge Chris Moore for helpful discussions, as well as Alexei Rozanov from IUP Bremen for providing the SCIATRAN radiative transfer code. The authors also wish to thank UMIAQ for logistical support, and Bristow Air for providing a helicopter for the deployment of IL1.

References

Abbatt, J. P. D., Thomas, J. L., Abrahamsson, K., Boxe, C., Granfors, A., Jones, A. E., King, M. D., Saiz-Lopez, A., Shepson, P. B., Sodeau, J., Toohey, D. W., Toubin, C.,

ACPD

14, 23949–23994, 2014

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



von Glasow, R., Wren, S. N., and Yang, X.: Halogen activation via interactions with environmental ice and snow in the polar lower troposphere and other regions, *Atmos. Chem. Phys.*, 12, 6237–6271, doi:10.5194/acp-12-6237-2012, 2012. 23950

Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138–141, doi:10.1038/334138a0, 1988. 23951

Bottenheim, J. W., Fuentes, J. D., Tarasick, D. W., and Anlauf, K. G.: Ozone in the Arctic lower troposphere during winter and spring 2000 (ALERT2000), *Atmos. Environ.*, 36, 2535–2544, doi:10.1016/S1352-2310(02)00121-8, 2002. 23951, 23967

Carlson, D., Donohoue, D., Platt, U., and Simpson, W. R.: A low power automated MAX-DOAS instrument for the Arctic and other remote unmanned locations, *Atmos. Meas. Tech.*, 3, 429–439, doi:10.5194/amt-3-429-2010, 2010. 23954

Chance, K.: Analysis of BrO measurements from the global ozone monitoring experiment, *Geophys. Res. Lett.*, 25, 3335–3338, 1998. 23953

Chance, K. V. and Spurr, R. J.: Ring effect studies: Rayleigh scattering, including molecular parameters for rotational Raman scattering, and the Fraunhofer spectrum, *Appl. Optics*, 36, 5224–5230, 1997. 23981

Deutschmann, T., Beirle, S., Frieß, U., Grzegorski, M., Kern, C., Kritten, L., Platt, U., Prados-Román, C., Puškite, J., Wagner, T., Werner, B., and Pfeilsticker, K.: The Monte Carlo atmospheric radiative transfer model McArtim: Introduction and validation of Jacobians and 3D features, *J. Quant. Spectrosc. Ra.*, 112, 1119–1137, doi:10.1016/j.jqsrt.2010.12.009, 2011. 23956

Fan, S.-M., and Jacob, D. J.: Surface ozone depletion in Arctic spring sustained by bromine reactions on aerosols, *Nature*, 359, 522–524, doi:10.1038/359522a0, 1992. 23951, 23952

Fayt, C., De Smedt, I., Letocart, V., Merlaud, A., Pinardi, G., Van Roozendael, M., and Roozendael, M. V. A. N.: QDOAS Software User Manual, available at: <http://uv-vis.aeronomie.be/software/QDOAS/index.php> (last access: 10 March 2014), 2011. 23955

Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br₂ and BrCl via uptake of HOBr onto aqueous salt solutions, *J. Geophys. Res.*, 104, 23719–23727, doi:10.1029/1999JD900359, 1999. 23951

Foster, K. L., Plastridge, R. A., Bottenheim, J. W., Shepson, P. B., Finlayson-Pitts, B. J., and Spicer, C. W.: The role of Br₂ and BrCl in surface ozone destruction at polar sunrise, *Science*, 291, 471–474, 2001. 23951

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Frieß U., Monks, P. S., Remedios, J. J., Rozanov, A., Sinreich, R., Wagner, T., and Platt, U.: MAX-DOAS O₄ measurements: a new technique to derive information on atmospheric aerosols: 2. Modeling studies, *J. Geophys. Res.*, 111, D14203, doi:10.1029/2005JD006618, 2006. 23955, 23956
- 5 Frieß U., Sihler, H., Sander, R., Pöhler, D., Yilmaz, S., and Platt, U.: The vertical distribution of BrO and aerosols in the Arctic: measurements by active and passive differential optical absorption spectroscopy, *J. Geophys. Res.*, 116, D00R04, doi:10.1029/2011JD015938, 2011. 23952, 23953, 23956, 23960, 23964, 23970
- Halfacre, J. W., Knepp, T. N., Shepson, P. B., Thompson, C. R., Pratt, K. A., Li, B., Peterson, P. K., Walsh, S. J., Simpson, W. R., Matrai, P. A., Bottenheim, J. W., Netcheva, S., Perovich, D. K., and Richter, A.: Temporal and spatial characteristics of ozone depletion events from measurements in the Arctic, *Atmos. Chem. Phys.*, 14, 4875–4894, doi:10.5194/acp-14-4875-2014, 2014. 23952, 23966, 23968
- 10 Hara, K., Osada, K., Matsunaga, K., Iwasaka, Y., Shibata, T., and Furuya, K.: Atmospheric inorganic chlorine and bromine species in Arctic boundary layer of the winter/spring, *J. Geophys. Res.*, 107, 4361, doi:10.1029/2001JD001008, 2002. 23970
- Helmig, D., Boylan, P., Johnson, B., Oltmans, S., Fairall, C., Staebler, R., Weinheimer, A., Orlando, J., Knapp, D. J., Montzka, D. D., Flocke, F., Frieß, U., Sihler, H., and Shepson, P. B.: Ozone dynamics and snow-atmosphere exchanges during ozone depletion events at Barrow, Alaska, *J. Geophys. Res.-Atmos.*, 117, D20303, doi:10.1029/2012JD017531, 2012. 23951, 23962, 23967
- 20 Hermans, C., Vandaele, A., Coquart, B., Jenouvrier, A., Merienne, M. F., Fally, S., Carleer, M., and Colin, R.: Absorption bands of O₂ and its collision-induced bands in the 30 000–7500 cm⁻¹ wavenumber region, in: *IRS 2000: Current Problems in Atmospheric Radiation*, edited by: Smith, W. L. and Timofeyev, Y. M., Hampton, VA, Deepak, 639–642, 2001. 23981
- Hönninger, G., von Friedeburg, C., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS), *Atmos. Chem. Phys.*, 4, 231–254, doi:10.5194/acp-4-231-2004, 2004. 23954
- 25 Huff, A. K. and Abbatt, J. P. D.: Gas-phase Br₂ production in heterogeneous reactions of Cl₂, HOCl, and BrCl with halide-ice surfaces, *J. Phys. Chem. A*, 104, 7284–7293, doi:10.1021/jp001155w, 2000. 23951, 23969
- Jones, A. E., Anderson, P. S., Begoin, M., Brough, N., Hutterli, M. A., Marshall, G. J., Richter, A., Roscoe, H. K., and Wolff, E. W.: BrO, blizzards, and drivers of polar tropospheric ozone

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



depletion events, *Atmos. Chem. Phys.*, 9, 4639–4652, doi:10.5194/acp-9-4639-2009, 2009. 23952, 23968, 23970

Jones, A. E., Anderson, P. S., Wolff, E. W., Roscoe, H. K., Marshall, G. J., Richter, A., Brough, N., and Colwell, S. R.: Vertical structure of Antarctic tropospheric ozone depletion events: characteristics and broader implications, *Atmos. Chem. Phys.*, 10, 7775–7794, doi:10.5194/acp-10-7775-2010, 2010. 23951, 23952, 23967

Koo, J.-H., Wang, Y., Kurosu, T. P., Chance, K., Rozanov, A., Richter, A., Oltmans, S. J., Thompson, A. M., Hair, J. W., Fenn, M. A., Weinheimer, A. J., Ryerson, T. B., Solberg, S., Huey, L. G., Liao, J., Dibb, J. E., Neuman, J. A., Nowak, J. B., Pierce, R. B., Natarajan, M., and Al-Saadi, J.: Characteristics of tropospheric ozone depletion events in the Arctic spring: analysis of the ARCTAS, ARCPAC, and ARCIONS measurements and satellite BrO observations, *Atmos. Chem. Phys.*, 12, 9909–9922, doi:10.5194/acp-12-9909-2012, 2012. 23953

Krnavek, L., Simpson, W. R., Carlson, D., Domine, F., Douglas, T. A., and Sturm, M.: The chemical composition of surface snow in the Arctic: examining marine, terrestrial, and atmospheric influences, *Atmos. Environ.*, 50, 349–359, doi:10.1016/j.atmosenv.2011.11.033, 2012. 23951

Lehrer, E., Hönninger, G., and Platt, U.: A one dimensional model study of the mechanism of halogen liberation and vertical transport in the polar troposphere, *Atmos. Chem. Phys.*, 4, 2427–2440, doi:10.5194/acp-4-2427-2004, 2004. 23951

Liao, J., Sihler, H., Huey, L. G., Neuman, J. A., Tanner, D. J., Friess, U., Platt, U., Flocke, F. M., Orlando, J. J., Shepson, P. B., Beine, H. J., Weinheimer, A. J., Sjostedt, S. J., Nowak, J. B., Knapp, D. J., Staebler, R. M., Zheng, W., Sander, R., Hall, S. R., and Ullmann, K.: A comparison of Arctic BrO measurements by chemical ionization mass spectrometry and long path-differential optical absorption spectroscopy, *J. Geophys. Res.*, 116, 1–14, doi:10.1029/2010JD014788, 2011. 23953, 23961, 23964

Liao, J., Huey, L. G., Tanner, D. J., Flocke, F. M., Orlando, J. J., Neuman, J. A., Nowak, J. B., Weinheimer, A. J., Hall, S. R., Smith, J. N., Fried, A., Staebler, R. M., Wang, Y., Koo, J.-H., Cantrell, C. A., Weibring, P., Walega, J., Knapp, D. J., Shepson, P. B., and Stephens, C. R.: Observations of inorganic bromine (HOBr, BrO, and Br₂) speciation at Barrow, Alaska, in spring 2009, *J. Geophys. Res.*, 117, D00R16, doi:10.1029/2011JD016641, 2012. 23953

Liao, J., Huey, L. G., Liu, Z., Tanner, D. J., Cantrell, C. A., Orlando, J. J., Flocke, F. M., Shepson, P. B., Weinheimer, A. J., Hall, S. R., Ullmann, K., Beine, H. J., Wang, Y., Ingall, E. D., Stephens, C. R., Hornbrook, R. S., Apel, E. C., Riemer, D., Fried, A., Mauldin III, R. L.,

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smith, J. N., Staebler, R. M., Neuman, J. A., and Nowak, J. B.: High levels of molecular chlorine in the Arctic atmosphere, *Nat. Geosci.*, 7, 91–94, doi:10.1038/ngeo2046, 2014. 23951, 23953

Malicet, J., Daumont, D., Charbonnier, J., Parisse, C., Chakir, A., and Brion, J.: Ozone UV spectroscopy. II. Absorption cross-sections and temperature dependence, *J. Atmos. Chem.*, 21, 263–273, doi:10.1007/BF00696758, 1995. 23981

McConnell, J. C., Henderson, G. S., Barrie, L., Bottenheim, J., Niki, H., Langford, C. H., and Templeton, E. M. J.: Photochemical bromine production implicated in Arctic boundary-layer ozone depletion, *Nature*, 355, 150–152, doi:10.1038/355150a0, 1992. 23952

McElroy, C. T., McLinden, C. A., and McConnell, J. C.: Evidence for bromine monoxide in the free troposphere during the arctic polar sunrise, *Nature*, 397, 338–341, 1999. 23967

Moore, C. W., Obrist, D., Steffen, A., Staebler, R. M., Douglas, T. A., Richter, A., and Nghiem, S. V.: Convective forcing of mercury and ozone in the Arctic boundary layer induced by leads in sea ice., *Nature*, 506, 81–84, doi:10.1038/nature12924, 2014. 23951, 23967

Mozurkewich, M.: Mechanisms for the release of halogens from sea-salt particles by free radical reactions, *J. Geophys. Res.*, 100, 14199–14207, 1995. 23951

Neuman, J. A., Nowak, J. B., Huey, L. G., Burkholder, J. B., Dibb, J. E., Holloway, J. S., Liao, J., Peischl, J., Roberts, J. M., Ryerson, T. B., Scheuer, E., Stark, H., Stickel, R. E., Tanner, D. J., and Weinheimer, A.: Bromine measurements in ozone depleted air over the Arctic Ocean, *Atmos. Chem. Phys.*, 10, 6503–6514, doi:10.5194/acp-10-6503-2010, 2010. 23953, 23963

Nghiem, S. V., Clemente-Colón, P., Douglas, T., Moore, C., Obrist, D., Perovich, D. K., Pratt, K. A., Rigor, I. G., Simpson, W., Shepson, P. B., Steffen, A., and Woods, J.: Studying bromine, ozone, and mercury chemistry in the Arctic, *EOS T. Am. Geophys. Un.*, 94, 289–291, doi:10.1002/2013EO330002, 2013. 23954

Oltmans, S. J., Johnson, B. J., and Harris, J. M.: Springtime boundary layer ozone depletion at Barrow, Alaska: meteorological influence, year-to-year variation, and long-term change, *J. Geophys. Res.*, 117, D00R18, doi:10.1029/2011JD016889, 2012. 23951, 23967

Parrella, J. P., Jacob, D. J., Liang, Q., Zhang, Y., Mickley, L. J., Miller, B., Evans, M. J., Yang, X., Pyle, J. A., Theys, N., and Van Roozendaal, M.: Tropospheric bromine chemistry: implications for present and pre-industrial ozone and mercury, *Atmos. Chem. Phys.*, 12, 6723–6740, doi:10.5194/acp-12-6723-2012, 2012. 23951

Payne, V. H., Clough, S. A., Shephard, M. W., Nassar, R., and Logan, J. A.: Information-centered representation of retrievals with limited degrees of freedom for signal: application

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to methane from the tropospheric emission spectrometer, *J. Geophys. Res.*, 114, D10307, doi:10.1029/2008JD010155, 2009. 23953, 23959

Piot, M. and von Glasow, R.: The potential importance of frost flowers, recycling on snow, and open leads for ozone depletion events, *Atmos. Chem. Phys.*, 8, 2437–2467, doi:10.5194/acp-8-2437-2008, 2008. 23951

Platt, U., and Hönninger, G.: The role of halogen species in the troposphere, *Chemosphere*, 52, 325–338, doi:10.1016/S0045-6535(03)00216-9, 2003. 23952

Pöhler, D., Vogel, L., Friess, U., and Platt, U.: Observation of halogen species in the Amundsen Gulf, Arctic, by active long-path differential optical absorption spectroscopy, *P. Natl. Acad. Sci. USA*, 107, 6582–6587, doi:10.1073/pnas.0912231107, 2010. 23953, 23967, 23968

Pratt, K. A., Custard, K. D., Shepson, P. B., Douglas, T. A., Pöhler, D., General, S., Zielcke, J., Simpson, W. R., Platt, U., Tanner, D. J., Gregory Huey, L., Carlsen, M., and Stirm, B. H.: Photochemical production of molecular bromine in Arctic surface snowpacks, *Nat. Geosci.*, 6, 351–356, doi:10.1038/ngeo1779, 2013. 23951, 23953, 23963, 23968, 23969

Richter, A., Wittrock, F., Eisinger, M., and Burrows, J. P.: GOME observations of tropospheric BrO in Northern Hemispheric spring and summer 1997, *Geophys. Res. Lett.*, 25, 2683–2686, 1998. 23953

Rodgers, C. D.: *Inverse Methods For Atmospheric Sounding: Theory and Practice*, World Scientific, Singapore, 2000. 23956, 23957, 23963

Rozanov, A., Bovensmann, H., Bracher, A., Hrechanyy, S., Rozanov, V., Sinnhuber, M., Stroh, F., and Burrows, J.: NO₂ and BrO vertical profile retrieval from SCIAMACHY limb measurements: sensitivity studies, *Adv. Space Res.*, 36, 846–854, doi:10.1016/j.asr.2005.03.013, 2005. 23956

Saiz-Lopez, A. and von Glasow, R.: Reactive halogen chemistry in the troposphere, *Chem. Soc. Rev.*, 41, 6448–6472, 2012. 23950

Salawitch, R. J., Canty, T., Kurosu, T., Chance, K., Liang, Q., da Silva, A., Pawson, S., Nielsen, J. E., Rodriguez, J. M., Bhartia, P. K., Liu, X., Huey, L. G., Liao, J., Stickel, R. E., Tanner, D. J., Dibb, J. E., Simpson, W. R., Donohoue, D., Weinheimer, A., Flocke, F., Knapp, D., Montzka, D., Neuman, J. A., Nowak, J. B., Ryerson, T. B., Oltmans, S., Blake, D. R., Atlas, E. L., Kinnison, D. E., Tilmes, S., Pan, L. L., Hendrick, F., Van Roozendaal, M., Kreher, K., Johnston, P. V., Gao, R. S., Johnson, B., Bui, T. P., Chen, G., Pierce, R. B., Crawford, J. H., and Jacob, D. J.: A new interpretation of total column BrO during Arctic spring, *Geophys. Res. Lett.*, 37, L21805, doi:10.1029/2010GL043798, 2010. 23953

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Schroeder, W., Anlauf, K., Barrie, L., and Lu, J.: Arctic springtime depletion of mercury, *Nature*, 16–17, 394, doi:10.1038/28530, 1998. 23951
- Sihler, H., Platt, U., Beirle, S., Marbach, T., Kühl, S., Dörner, S., Verschaeve, J., Frieß, U., Pöhler, D., Vogel, L., Sander, R., and Wagner, T.: Tropospheric BrO column densities in the Arctic derived from satellite: retrieval and comparison to ground-based measurements, *Atmos. Meas. Tech.*, 5, 2779–2807, doi:10.5194/amt-5-2779-2012, 2012. 23953, 23966, 23969
- Simpson, W. R., Carlson, D., Hönninger, G., Douglas, T. A., Sturm, M., Perovich, D., and Platt, U.: First-year sea-ice contact predicts bromine monoxide (BrO) levels at Barrow, Alaska better than potential frost flower contact, *Atmos. Chem. Phys.*, 7, 621–627, doi:10.5194/acp-7-621-2007, 2007a. 23963
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, doi:10.5194/acp-7-4375-2007, 2007b. 23951
- Spicer, C. W., Plastridge, R. A., Foster, K. L., Finlayson-Pitts, B. J., Bottenheim, J. W., Grannas, A. M., and Shepson, P. B.: Molecular halogens before and during ozone depletion events in the Arctic at polar sunrise: concentrations and sources, *Atmos. Environ.*, 36, 2721–2731, doi:10.1016/S1352-2310(02)00125-5, 2002. 23951
- Steffen, A., Douglas, T., Amyot, M., Ariya, P., Aspmo, K., Berg, T., Bottenheim, J., Brooks, S., Cobbett, F., Dastoor, A., Dommergue, A., Ebinghaus, R., Ferrari, C., Gardfeldt, K., Goodsite, M. E., Lean, D., Poulain, A. J., Scherz, C., Skov, H., Sommar, J., and Temme, C.: A synthesis of atmospheric mercury depletion event chemistry in the atmosphere and snow, *Atmos. Chem. Phys.*, 8, 1445–1482, doi:10.5194/acp-8-1445-2008, 2008. 23951
- Sturm, M. and Stuefer, S.: Wind-blown flux rates derived from drifts at arctic snow fences, *J. Glaciol.*, 59, 21–34, doi:10.3189/2013JoG12J110, 2013. 23970
- Tackett, P. J., Cavender, A. E., Keil, A. D., Shepson, P. B., Bottenheim, J. W., Morin, S., Deary, J., Steffen, A., and Doerge, C.: A study of the vertical scale of halogen chemistry in the Arctic troposphere during Polar Sunrise at Barrow, Alaska, *J. Geophys. Res.*, 112, D07306, doi:10.1029/2006JD007785, 2007. 23951, 23953

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tarasick, D. W. and Bottenheim, J. W.: Surface ozone depletion episodes in the Arctic and Antarctic from historical ozonesonde records, *Atmos. Chem. Phys.*, 2, 197–205, doi:10.5194/acp-2-197-2002, 2002. 23967
- 5 Theys, N., Van Roozendael, M., Errera, Q., Hendrick, F., Daerden, F., Chabrillat, S., Dorf, M., Pfeilsticker, K., Rozanov, A., Lotz, W., Burrows, J. P., Lambert, J.-C., Goutail, F., Roscoe, H. K., and De Mazière, M.: A global stratospheric bromine monoxide climatology based on the BASCOE chemical transport model, *Atmos. Chem. Phys.*, 9, 831–848, doi:10.5194/acp-9-831-2009, 2009. 23953
- 10 Theys, N., Van Roozendael, M., Hendrick, F., Yang, X., De Smedt, I., Richter, A., Begoin, M., Errera, Q., Johnston, P. V., Kreher, K., and De Mazière, M.: Global observations of tropospheric BrO columns using GOME-2 satellite data, *Atmos. Chem. Phys.*, 11, 1791–1811, doi:10.5194/acp-11-1791-2011, 2011. 23953
- Toom-Saunty, D. and Barrie, L. A.: Chemical composition of snowfall in the high Arctic: 1990–1994, *Atmos. Environ.*, 36, 2683–2693, 2002. 23951
- 15 Toyota, K., McConnell, J. C., Staebler, R. M., and Dastoor, A. P.: Air–snowpack exchange of bromine, ozone and mercury in the springtime Arctic simulated by the 1-D model PHANTAS – Part 1: In-snow bromine activation and its impact on ozone, *Atmos. Chem. Phys.*, 14, 4101–4133, doi:10.5194/acp-14-4101-2014, 2014. 23951, 23969
- 20 Tuckermann, M., R., A., Golz, C., Lorenzen-Schmidt, H., Senne, T., Stutz, J., Trost, B., Unold, W., and Platt, U.: DOAS-observation of halogen radical-catalysed arctic boundary layer ozone destruction during the ARCTOC-campaigns 1995 and 1996 in Ny-Alesund, Spitsbergen, *Tellus B*, 49, 533–555, doi:10.1034/j.1600-0889.49.issue5.9.x, 1997. 23953
- Vandaele, A., Hermans, C., Simon, P., Carleer, M., Colin, R., Fally, S., Mérianne, M., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 25 42 000 cm⁻¹ to 10 000 cm⁻¹ (238–1000 nm) at 220 K and 294 K, *J. Quant. Spectrosc. Ra.*, 59, 171–184, doi:10.1016/S0022-4073(97)00168-4, 1998. 23981
- von Clarmann, T. and Grabowski, U.: Elimination of hidden a priori information from remotely sensed profile data, *Atmos. Chem. Phys.*, 7, 397–408, doi:10.5194/acp-7-397-2007, 2007. 23958
- 30 Wagner, T. and Platt, U.: Satellite mapping of enhanced BrO concentrations in the troposphere, *Nature*, 395, 486–490, 1998. 23953

Wilmouth, D. M., Hanisco, T. F., Donahue, N. M., and Anderson, J. G.: Fourier transform ultraviolet spectroscopy of the $A^2\Pi_{3/2} \leftarrow X^2\Pi_{3/2}$ transition of BrO, *J. Phys. Chem. A*, 103, 8935–8945, doi:10.1021/jp991651o, 1999. 23981

Wren, S. N., Donaldson, D. J., and Abbatt, J. P. D.: Photochemical chlorine and bromine activation from artificial saline snow, *Atmos. Chem. Phys.*, 13, 9789–9800, doi:10.5194/acp-13-9789-2013, 2013. 23951, 23969

Yang, X., Pyle, J. A., Cox, R. A., Theys, N., and Van Roozendael, M.: Snow-sourced bromine and its implications for polar tropospheric ozone, *Atmos. Chem. Phys.*, 10, 7763–7773, doi:10.5194/acp-10-7763-2010, 2010. 23951, 23952, 23968

ACPD

14, 23949–23994, 2014

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vertical structure of
BrO**

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Absorber cross sections used in the MAX-DOAS fitting.

Species	Cross Section
BrO (223 K)	Wilmouth et al. (1999)
O ₃ (243 K)	Malicet et al. (1995)
NO ₂ (220 K)	Vandaele et al. (1998)
O ₄	Hermans et al. (2001)
Ring	Determined from zenith spectra using Chance and Spurr (1997)




Figure 1. A sample profile retrieval of BrO at the BARC building. The blue line represents the profile retrieved using an a priori profile that exponentially decays with a scale height of 1000 m, while the green represents quantities retrieved using an a priori profile that exponentially decays with a scale height of 400 m (a priori used for this study). The dashed lines indicate layer boundaries determined from the ensemble of retrievals over the course of this study. The partial VCD values noted on the figure correspond to layer selections used in this study. The $\text{VCD}_{200\text{m}}$ is the integral of the retrieved profile from 0–200 m and the $\text{VCD}_{\text{Aloft}}$ is integrated from 200 to 2000 m.

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. An illustration of the grid coarsening of the BrO averaging kernels. These averaging kernels correspond to the retrieval shown in green on Fig. 1. Averaging kernels for the original retrieval (left) and for the coarse grid (right).

Vertical structure of BrO

P. K. Peterson et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vertical structure of
BrO

P. K. Peterson et al.




Figure 3. The left panel shows the degrees of freedom associated with the VCD_{200m} as a function of near-surface aerosol particle extinction. The right panel shows the degrees of freedom associated with the $VCD_{Residual}$ as a function of the aerosol optical depth. The dashed line illustrates the cutoff we have chosen for sufficient information content.



Vertical structure of
BrO

P. K. Peterson et al.




Figure 4. A portion of the timeseries of BrO observed during this study. The top panel represents the VCD_{200} , the second panel represents the LT-VCD, both of which have units of molecules cm^{-2} . The third panel shows the percentage of the LT-VCD observed in the lowest 200 m, while the bottom panel shows the aerosol optical depth over the course of this study. In the third panel, ratios are not calculated for events that have a LT-VCD below 5×10^{12} molecules cm^{-2} . Shaded areas represent potentially titrated air masses near the surface (Ozone $< 1 \text{ nmol mol}^{-1}$). The full timeseries can be found in the Supplement.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Vertical structure of
BrO**

P. K. Peterson et al.



Figure 5. Field locations during the BROMEX field campaign overlain on the Environmental Systems Research Institute (ESRI) world image. These locations are Barrow Arctic Research Center (BARC), Chemical Ionization Mass Spectrometry (CIMS), NOAA Global Monitoring Division (NOAA GMD) station, IceLander 1 (IL1) buoy, and Barrow Airport (PABR), nearby where most of Barrow's population resides. The distance between BARC and IL1 is about 36 km.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 6. A timeseries of 30 min averaged BrO measured using CIMS, compared with average BrO in the lowest 100 m measured by MAX-DOAS at two different sites. The grey area indicates times when there were significant differences in BrO observed between MAX-DOAS sites at the BARC Building and on land fast ice 36 km NE of the BARC building (IL1).

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 7. The correlation of 60 min averaged CIMS BrO with BrO retrieved in the lowest 100 m using MAX-DOAS. Red circles indicate times when we observed a chemical gradient between field sites, which are excluded from this correlation.




Figure 8. The relationship between the total BrO activation, as measured by the LT-VCD, and the vertical structure of the activation event. The red line represents the mean percentage of BrO in the lowest 200 m as a function of the LT-VCD, calculated from deciles on the x axis, while the shaded region represents one standard deviation. The coloration indicates the VCD_{200m} . In this plot LT-VCD values below 5×10^{12} molecules cm^{-2} are eliminated to avoid calculating percentages from near zero amounts of BrO.

Vertical structure of BrO

P. K. Peterson et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Vertical structure of
BrO

P. K. Peterson et al.




Figure 9. The relationship between daily estimated temperature gradients vs. the average percentage of BrO observed in the near surface layer during that day is shown here using a box and whisker plot. The data are split such that each bin comprises ≈ 20 days worth of data. The red lines show the median value for each bin, while the blue box encloses the 25–75th percentile and the whiskers show the full range of data excluding outliers (\times). Outliers are points that are outside 3σ for the corresponding group.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 10. The relationship between BrO and temperature. The left panel shows the percentage of BrO in the lowest 200 m vs. the near-surface temperature, while the right panel shows the LT-VCD vs. the near-surface temperature. The red line represents the mean, calculated from deciles on the x axis, while the shaded region represents one standard deviation. In the left panel, LT-VCD values below 5×10^{12} molecules cm^{-2} are eliminated to avoid calculating percentages from near zero amounts of BrO.

Vertical structure of
BrO

P. K. Peterson et al.




Figure 11. The left panel shows the relationship between the BrO LT-VCD and wind speed, while the right panel shows percentage of BrO in the lowest 200 m vs. the wind speed. In both cases, the red line represents the mean as a function of wind speed, calculated from deciles on the x axis, while the shaded region represents one standard deviation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Vertical structure of
BrO

P. K. Peterson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 12. The diurnal cycling of the BrO LT-VCD (blue) and VCD_{200m} (red). The solid lines represent the mean VCD as a function of time of day, while the shaded region represents one standard deviation. The shaded regions are symmetric about the mean in both cases, however the cycling of the VCD_{200m} is overlain on the cycling of the LT-VCD, partially obscuring the standard deviation of the LT-VCD.

Vertical structure of
BrO

P. K. Peterson et al.




Figure 13. The left panel shows the relationship between the BrO VCD_{200m} and the near-surface aerosol particle extinction. The right panel shows the relationship between the BrO LT-VCD and the aerosol optical depth. In both plots the x axis is a log scale. The red line represents the mean VCD, calculated from deciles on the x axis, while the shaded region represents one standard deviation. In both cases, the coloration indicates the wind speed.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)