

800 year ice-core record of nitrogen deposition in Svalbard linked to ocean productivity and biogenic emissions

I. A. Wendl^{1,2,3}, A. Eichler^{1,2}, E. Isaksson⁴, T. Martma⁵, M. Schwikowski^{1,2,3}

[1]{Laboratory of Radiochemistry and Environmental Chemistry, Paul Scherrer Institut, 5232 Villigen PSI, Switzerland}

[2]{Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland}

[3]{Department of Chemistry and Biochemistry, University of Bern, Bern, Switzerland}

[4]{Norwegian Polar Institute, Framsenteret, 9296 Tromsø, Norway}

[5]{Institute of Geology, Tallinn University of Technology, Estonia}

Correspondence to: M. Schwikowski (margit.schwikowski@psi.ch)

Abstract

We present the records of the two nitrogen species nitrate (NO_3^-) and ammonium (NH_4^+) analysed in a new ice core from Lomonosovfonna, Svalbard, in the Eurasian Arctic covering the period 1222-2009. We investigate the emission sources and find that during the 20th century both records are influenced by anthropogenic pollution from Eurasia. In pre-industrial times NO_3^- is highly correlated with methane-sulfonate (MSA) on decadal time-scales, which we explain by a fertilising effect. Enhanced atmospheric NO_3^- concentrations and the corresponding nitrogen input to the ocean trigger the growth of dimethyl-sulfide-(DMS)-producing phytoplankton. Increased DMS production results in elevated fluxes to the atmosphere where it is oxidised to MSA. Eurasia was presumably the main source area also for pre-industrial NO_3^- , but a more exact source apportionment could not be performed based on our data. This is different for NH_4^+ , where biogenic ammonia (NH_3) emissions from Siberian boreal forests were identified as the dominant source of pre-industrial NH_4^+ . Changes in melt at the Lomonosovfonna glacier are assumed to have a negligible effect on the decadal variations of the investigated compounds.

1 **1 Introduction**

2 The Arctic is generally a nutrient limited region (Dickerson, 1985). Nutrients originate from
3 lower latitudes and reach the remote polar areas via long-range transport, local sources are
4 sparse. The major source for bio-available nitrogen in the Arctic is the deposition of reactive
5 atmospheric nitrogen that is present primarily as nitrate (NO_3^-) and ammonium (NH_4^+)
6 (Björkman et al., 2013; Kühnel et al., 2011). Those species are predominantly removed from
7 the atmosphere by wet deposition (Bergin et al., 1995). NO_3^- is the oxidation product of
8 emitted NO_x (NO and NO_2). At a global scale, major NO_3^- sources include biomass burning,
9 emissions from microbial processes in soils, ammonia oxidation, stratospheric injection,
10 lightning, as well as fossil fuel and biofuel combustion, and aircraft emissions (Fibiger et al.,
11 2013; Galloway et al., 2004; Hastings et al., 2004; Wolff, 2013). NH_4^+ derives from biogenic
12 emissions of ammonia (NH_3) from terrestrial and marine sources, biomass burning,
13 agriculture, and livestock breeding (Fuhrer et al., 1996; Galloway et al., 2004; Wolff, 2013).
14 Both NO_3^- and NH_4^+ concentrations in the atmosphere have varied greatly with time and
15 space due to changing emissions and the short atmospheric lifetimes of a few days (Adams et
16 al., 1999; Feng and Penner, 2007). Generally, concentrations were low in pre-industrial times
17 and increased due to stronger emissions with beginning of the industrialisation and
18 intensification of agricultural activities (Galloway et al., 2004). The deposition of NO_3^- and
19 NH_4^+ in the Arctic is an important nutrient source. Varying concentrations thus greatly affect
20 the nitrogen budget in the Arctic where nutrient supply is limited.

21 Ice cores represent an invaluable archive of past atmospheric composition. Ice core studies
22 from the Arctic clearly reveal an anthropogenic influence on the concentrations of NO_3^- and
23 NH_4^+ approximately during the last 150 years (Fischer et al., 1998; Fuhrer et al., 1996; Goto-
24 Azuma and Koerner, 2001; Kekonen et al., 2002, 2005; Legrand and Mayewski, 1997;
25 Matoba et al., 2002; Simões and Zagorodnov, 2001). North America was identified as major
26 pollutant source for south Greenland, both North America and Eurasia for central and north
27 Greenland, and Eurasia for Svalbard (Goto-Azuma and Koerner, 2001; Hicks and Isaksson,
28 2006). However, the pre-industrial sources of NO_3^- and NH_4^+ are still fairly unknown
29 (Legrand and Mayewski, 1997; Wolff, 2013). Eichler et al. (2011) identified forest fires as
30 major source of NO_3^- in a Siberian Altai ice core from the mid-latitudes. In studies on
31 Greenland ice NO_3^- was also associated with forest fires (Whitlow et al., 1994; Wolff et al.,
32 2008). Pre-industrial NH_4^+ in ice cores from the mid-latitudes was attributed to biogenic

1 emissions (Eichler et al., 2009; Kellerhals et al., 2010). Similarly, long-term trends in
2 Greenland ice cores have been attributed to changing biogenic emission from North America,
3 whereas short-term NH_4^+ changes were found to correlate with forest fires (Fuhrer et al.,
4 1996; Whitlow et al., 1994; Zennaro et al., 2014).

5 Whereas a few records exist from Greenland, there is less information available from the
6 Eurasian Arctic. The NO_3^- and NH_4^+ records of a previous ice core from Lomonosovfonna,
7 Svalbard, retrieved in 1997 (Lomo97, for location see Fig. 1), cover the last 1000 years
8 (Divine et al., 2011; Kekonen et al., 2002, 2005). For both species a clear anthropogenic
9 impact is observed in the second half of the 20th century, but the pre-industrial sources remain
10 largely unidentified due to potential runoff that biased the ion records before the mid-16th
11 century (Kekonen et al., 2002, 2005). Nevertheless, the fairly stable concentrations in the
12 NO_3^- record from the mid-16th to the mid-19th century are interpreted as input from natural
13 NO_3^- sources (Kekonen et al., 2002). An anthropogenic influence in the 20th century is also
14 visible in the NO_3^- and NH_4^+ records of other Eurasian Arctic ice cores (see Fig. 1 for
15 locations) from Holtedahlfonna (Holte05), Svalbard (Beaudon et al., 2013), Snøfjellafonna,
16 Svalbard (Goto-Azuma and Koerner, 2001), and Severnaya Zemlya (Weiler et al., 2005). The
17 industrial records from these cores are discussed in detail, but pre-industrial sources and
18 concentration changes of the inorganic nitrogen species remain unexplained.

19 The interpretation of NO_3^- and NH_4^+ as paleo-environmental proxies may be hampered by the
20 fact that both undergo post-depositional processes leading to loss from or relocation within
21 the snow pack even at temperatures well below the melting point (Pohjola et al., 2002). NO_3^-
22 can be relocated or lost by photolysis and/or evaporation of nitric acid (HNO_3) (Honrath et al.,
23 1999; Röthlisberger et al., 2002). This loss can be severe at low accumulation sites such as
24 Dome C, Antarctica (Röthlisberger et al., 2000, 2002). At sites with higher accumulation rates
25 such as Summit in Greenland or Weissfluhjoch in the European Alps the majority of NO_3^- is
26 preserved (Baltensperger et al., 1993; Fibiger et al., 2013). Many studies reveal that NH_4^+ and
27 NO_3^- are preserved in snow and firn cores with respect to percolating melt water (Eichler et
28 al., 2001; Ginot et al., 2010; Moore and Grinsted, 2009; Pohjola et al., 2002), but others report
29 a preferential elution of these species compared to other major ions (Brimblecombe et al.,
30 1985; Moore and Grinsted, 2009; Pohjola et al., 2002). The underlying mechanism is not well
31 understood, except from the fact that it depends on the overall ion composition.

1 In this paper we discuss the records of the two nitrogen species NO_3^- and NH_4^+ analysed in a
2 new ice core drilled on Lomonosovfonna, Svalbard, in 2009. The study focuses on the
3 investigation of the major sources of NO_3^- and NH_4^+ deposited in the Eurasian Arctic which
4 highly affects the nutrient budget in the region, along with the effect of melt on the
5 geochemical records of these nitrogen species which will gain importance due to the ongoing
6 global warming.

7

8 **2 Methods**

9 **2.1 Drilling site and meteorological setting**

10 In 2009, a 149.5 m long ice core was drilled on Lomonosovfonna, Svalbard (1202 m asl;
11 $78^\circ49'24''\text{N}$, $17^\circ25'59''\text{E}$, Lomo09), using the Fast Electromechanical Lightweight Ice
12 Coring System (FELICS) (Ginot et al., 2002). The 2009 drilling site is 4.6 km south of that in
13 1997 (Lomo97) (Isaksson et al., 2001). Bedrock was not reached but a radar survey suggested
14 it to be at around 200 m (Pettersson, unpublished data). Measured borehole temperatures in
15 the upper 42 m (between -1.7°C and -4.3°C) at the Lomo09 drill site are in good agreement
16 with the average borehole temperature at the Lomo97 site of -2.8°C with a nearly isothermal
17 profile (Van de Wal et al. 2002). Previous studies indicate that summer melt water in the
18 study area is refrozen mostly within the previous winter's snow, and the remainder within the
19 next two to three lower annual layers (Samuelson, 2001). Percolation length was found to be
20 up to 8 years only in the warmest years during the 20th century (Kekonen et al., 2005; Moore
21 et al., 2005).

22 Svalbard is located at a climatically sensitive area being surrounded by the Arctic Ocean, the
23 Barents Sea and the Atlantic Ocean, and situated at the southerly edge of the permanent
24 Arctic sea ice and close to the over-turning point of the North Atlantic thermohaline
25 circulation. Further, it is relatively close to the industrialised areas of Eurasia which were
26 found to highly affect the chemical composition of air reaching the archipelago, especially in
27 spring during the Arctic Haze (Eleftheriadis et al., 2009; Eneroth et al., 2003; Forsström et al.,
28 2009; Goto-Azuma and Koerner, 2001; Law and Stohl, 2007; Stohl et al., 2007). The Arctic
29 Haze describes a phenomenon of increased aerosol concentration in the end of winter to early
30 spring (Greenaway, 1950; Quinn et al., 2007; Shaw, 1995). At that time of the year
31 temperatures in the Arctic become very low which leads to a thermally very stable

1 stratification with strong surface inversions (Shaw, 1995; Stohl, 2006). This cold stratified air
2 forms a dome over the Arctic that hinders warm air masses from lower latitudes to enter. The
3 boundary of this dome that acts as a transport barrier is called Arctic or Polar Front whose
4 position shifts between summer and winter due to temperature. In summer only the more
5 northern parts of the Northern Hemisphere are cold enough to cause a stable stratification of
6 the atmosphere, whereas in winter temperatures in more southern parts are cold enough so
7 that the Arctic Front is located as far south as 40°N. Then large areas of Eurasia and partly
8 North America are included in the Arctic dome, facilitating transport of pollution from those
9 regions. In addition, since both dry and wet deposition is reduced within the Arctic dome in
10 winter, aerosol related species have very long lifetimes once within the Arctic dome (Stohl,
11 2006).

12 **2.2 Sampling and analyses**

13 The Lomonosovfonna 2009 ice core was processed in the cold room (-20°C) at Paul Scherrer
14 Institut, Switzerland, resulting in 3997 samples with a depth resolution of 3-4 cm (details on
15 the method in Eichler et al., 2000). The resolution was adapted to layer thinning with depth,
16 so that even in the deepest and oldest part of the core each year is at least represented by one
17 sample. The inner part of the core was sampled for the analysis of water soluble major ions
18 and the water stable isotopes $\delta^{18}\text{O}$ and δD . Outer core sections were analysed for ^3H and ^{210}Pb
19 used for dating purposes (Eichler et al., 2000).

20 Concentrations of water soluble major ions, including NO_3^- and NH_4^+ , were determined using
21 ion chromatography (Metrohm 850 Professional IC combined with a 872 Extension Module
22 and a 858 Professional Sample Processor autosampler). Prior to analyses ice samples were
23 melted in a N_2 atmosphere to reduce contamination from laboratory air. A list of the measured
24 ionic species, their detection limits and median concentrations are given in Table 1.

25 **2.3 Ice core dating**

26 The Lomo09 ice core covers the time period of 1222 to 2009 (Fig. 2). It was dated with a
27 combination of reference horizons, annual layer counting (ALC), ^{210}Pb decay, and a simple
28 glacier flow model. The reference horizons include the tritium peak indicating the year 1963,
29 and the major volcanic eruptions of Bezymianny (1956), Katmai (1912), Tambora (1815),
30 Laki (1783), Hekla (1766), Kuwae (1458/59; Sigl et al., 2013), and Samalas (1257/58;

1 Lavigne et al., 2013) marked by high non-sea-salt sulphate concentrations and high values for
2 the sulphate-residual of the multiple linear regression of all measured ions, a method
3 previously described in Moore et al. (2012). Annual layer counting was performed down to a
4 depth of ~ 79.7 m weq (= 1750) using the pronounced seasonality of $\delta^{18}\text{O}$ and Na^+
5 (Supplement Fig. S1). A simple glacier flow model (Thompson et al., 1998) was fitted
6 through the volcanic reference horizons. This was used to date the core below ~ 79.7 m weq
7 where ALC was limited due to strong layer thinning. The dating uncertainty for the core down
8 to a depth of ~ 68 m weq is estimated to be ± 1 year within ± 10 years of the reference horizons
9 and increases to ± 3 years in between. Down to a depth of ~ 80 m weq the dating uncertainty
10 enlarges to ± 3 years also in proximity of the reference horizons, and below ~ 80 m weq it
11 increases to ± 10 years. This was calculated using the difference of the year of the volcanic
12 eruptions and the modelled date. The average annual accumulation rate is 0.58 ± 0.13 m weq.

13 **2.4 Calculation of annual melt percent**

14 Melt features are formed when surface snow melts and the melt water percolates into deeper
15 layers where it fills the pores and refreezes under the formation of a layer of ice poor or free
16 of air bubbles. The percentage of annual melt in the Lomo09 core was calculated from the
17 thickness of melt features observed during processing of the core (similar to Henderson et al.,
18 2006). Clear and bubbly ice appears as transparent area when the core is backlit. If the melt
19 did not affect the whole core diameter, this was accounted for by multiplying the length of the
20 melt feature with the percentage of the core diameter it covered. If for example a melt feature
21 was 20 cm long but only affected one fifth of the core diameter, this melt feature would count
22 the same as a four centimetre long melt feature affecting the whole core diameter. The
23 observed melt features were then summed up per year to calculate the annual melt percent
24 (Fig. 3).

25

26 **3 Results and discussion**

27 The records of NO_3^- and NH_4^+ of the Lomo09 core both show the highest concentrations
28 during the period of approximately 1940 to 1980 (Figs. 3 and 4), similar to findings from
29 other Arctic sites (Goto-Azuma and Koerner, 2001). This clearly indicates a strong influence
30 of anthropogenic emissions in recent decades on the chemical composition of aerosols
31 reaching Lomonosovfonna. Both records show a significant decrease after 1980, a trend

1 similarly observed in the NO_3^- and NH_4^+ records of ice cores from the Siberian Altai (Eichler
2 et al., 2009, 2011) (Fig. 5) and Severnaya Zemlya (Opel et al., 2013; Weiler et al., 2005)
3 influenced mainly by Eurasian pollution. The Siberian Altai core ends in 2001, explaining
4 why the continuation of the decrease seen at Lomonosovfonna is not recorded there. The
5 nitrate trend in the Siberian Altai fits well with NO_x emissions as discussed in detail in Eichler
6 et al. (2009) where the Eastern European NO_x emissions are shown. In contrast, NO_3^-
7 concentrations in records from Summit, Greenland, and Colle Gnifetti, Swiss Alps (see Fig. 1
8 for locations), affected by Northern American and Western European air masses, respectively,
9 kept rising into the 21st century (Fig. 5). This suggests that the major sources for the increased
10 concentrations of NO_3^- and NH_4^+ in the Lomo09 core are similar to those for the Siberian
11 Altai and Severnaya Zemlya, whereas the influence of emissions in North America and
12 Europe is of minor importance. Our finding is in agreement with results from transport
13 modelling and trajectory analysis, identifying Eurasia as source region for Svalbard (Hirdman
14 et al., 2010a and b). Eurasian pollution dominates especially during the autumn and winter
15 period characterised by Arctic Haze episodes, in contrast to summer months when North
16 Atlantic air masses prevail (Tunved et al., 2013). We thus attribute the observed trend in NO_3^-
17 to higher NO_x emissions from traffic, energy production, and industrial activities, and in NH_4^+
18 to enhanced NH_3 emissions from agriculture and livestock mainly in Eurasia (Eichler et al.,
19 2009; Weiler et al., 2005). The anthropogenic impact is also seen in the NO_3^- and- less
20 pronounced- in the NH_4^+ record of the Lomo97 core (Divine et al., 2011; Kekonen et al.,
21 2005) (Fig. 4), which underlines the spatial representativeness of the Lomo09 ice core data.
22 The NO_3^- records of the Lomo09 and Lomo97 cores agree well. This is not the case for the
23 NH_4^+ records, where the Lomo97 shows higher concentrations, especially before 1900 (Fig.
24 4). We cannot explain this difference, but NH_4^+ is known to be prone to contamination during
25 analysis (Jauhiainen et al., 1999; Kaufmann et al., 2010; Legrand et al., 1984, 1993, 1999;
26 Udisti et al., 1994). Three observations indicate that the Lomo09 NH_4^+ concentrations are
27 robust: 1) The preindustrial Lomo09 values are generally lower than those of Lomo97 and
28 therefore contamination seems unlikely, 2) the Lomo09 preindustrial ion balance is close to
29 zero (Fig. S2), and 3) the 300 year records of NO_3^- and NH_4^+ from Holtedahlfonna (Beaudon
30 et al., 2013) are in reasonable agreement with the Lomo09 data (Fig. 4).

31 In order to investigate sources of NH_4^+ and NO_3^- and other ionic species in the Lomo09 ice
32 core we performed a principal component analysis (PCA). We used 10-year-averages to
33 account for dating uncertainties and smoothing effects by melt-water relocation. Previous

1 studies on the Lomo97 core suggested that the percolation lengths at the site can reach two to
2 eight annual layers in the warmest years (Moore et al., 2005; Pohjola et al., 2002). The
3 average annual melt percent of the Lomo97 core was 41% (Pohjola et al., 2002) compared to
4 31% of the Lomo09 core. We thus assume that the maximum percolation lengths in the
5 Lomo09 core do not exceed the eight annual layers determined for the Lomo97 core. As a
6 conservative estimate we used the 10-year-average record of melt percent in the PCA to
7 examine the influence of melt on the NH_4^+ and NO_3^- records. The PCA was performed only
8 for pre-industrial times (1222-1859) to exclude anthropogenic influences on the ion
9 concentrations. Sulphate (SO_4^{2-}) from anthropogenic sources has been shown to increase
10 already during the second half of the 19th century (Moore et al., 2006).

11 We obtained six principal components (PCs) from the PCA (Table 2). PC1 has high loadings
12 of sodium (Na^+), potassium (K^+), magnesium (Mg^{2+}), and chloride (Cl^-). This component
13 explains 38% of the total variance and contains species that are directly emitted by sea spray.
14 PC2 has high loadings of methane-sulfonate ($\text{MSA} = \text{CH}_3\text{SO}_3^-$) and NO_3^- . MSA has a strictly
15 marine biogenic source. It results from the oxidation of gaseous dimethyl-sulphide (DMS)
16 which is produced by phytoplankton and emitted from the ocean to the atmosphere. This gas
17 release across the sea-air interface differs distinctly from the way sea salt species are emitted
18 to the atmosphere via sea spray because no droplets are involved (Stefels et al., 2007; Vogt
19 and Liss, 2009). PC3 has a high loading of NH_4^+ , representing biogenic emissions. Calcium
20 (Ca^{2+}) is the only species that has a high loading in PC4. This suggests that PC4 represents a
21 mineral dust component. The melt percent is the only parameter that has a high loading in
22 PC5. This suggests that on the considered decadal time scale the influence of melt on the ion
23 concentration averages is negligible, which is in agreement with Pohjola et al. (2002) and
24 Moore et al. (2005). PC6 has a high loading of SO_4^{2-} , indicating a volcanic source because the
25 marine part of SO_4^{2-} is covered by the sea spray component PC1.

26 The results of the PCA are in good correspondence with those of a correlation analysis of the
27 10-year-averaged records of the ionic species and the melt percent for the pre-industrial
28 period (Table 3). Strong correlation is observed for the sea spray related ions Na^+ , K^+ , Mg^{2+} ,
29 and Cl^- ($0.59 < r^2 < 0.98$). Furthermore, MSA and NO_3^- are highly correlated and share 60% of
30 data variability. NH_4^+ , Ca^{2+} , melt percent and SO_4^{2-} are not significantly correlated with any
31 other species.

3.1 Nitrate and methane-sulfonate (NO₃⁻ and MSA)

In the Arctic MSA is well known to originate from marine biogenic sources (Legrand, 1997). However, major pre-industrial NO₃⁻ sources in this region are still not fully understood (e.g., Wolff et al., 2008).

Varying atmospheric MSA concentrations have been related to changing sea ice conditions. Studies from Arctic and Antarctic ice cores found positive (Becagli et al., 2009; Legrand et al., 1997), but also negative correlations of MSA and sea ice extent (Rhodes et al., 2009; Sharma et al., 2012), see also the review on MSA and sea ice in Antarctica by Abram et al. (2013). After 1920 the Lomo97 core MSA correlates negatively with summer (August) sea-ice extent and sea surface temperature in the Barents Sea (O'Dwyer et al., 2000) and positively with the instrumental summer temperature record from Svalbard (Isaksson et al., 2005). During the period 1600-1920 Isaksson et al. (2005) detected a positive correlation of the Lomo97 MSA and winter (April) sea ice extent in the Barents Sea (Divine and Dick, 2006; Vinje, 2001). The Lomo97 MSA record reveals a pattern with twice as high values prior to about 1920 compared to those of the later 20th century (Isaksson et al., 2005). They suggest that it results from a change of source and/or more favourable growing conditions for the DMS-producing phytoplankton in a more extensive sea ice environment before 1920.

In the MSA record of the Lomo09 core we find a similar pattern as in the Lomo97 core with higher concentrations prior to the 20th century and a decreasing trend since around 1900 (Figs. 4 and 6). Hence, we investigate if a coupling of MSA with sea ice conditions around Svalbard exists, using three long-term reconstructions of sea ice extent. These reconstructions include the winter (April) ice extent in the Western Nordic Seas covering the last 800 years (Macias Fauria et al., 2010), the summer (August) location of the sea ice edge in the Barents Sea (BS) that covers the last 400 years (Kinnard et al., 2011), and the summer sea ice extent in the Arctic Seas extending back to the year 563 (Kinnard et al., 2011). The best agreement was observed between the 40-year-lowpass-filtered records of Lomo09 MSA and winter (April) Western Nordic Seas ice extent (Macias Fauria et al., 2010) (Fig. 6; $r=0.56$, $p<0.001$). The most striking feature in both records is the pronounced decrease starting around 1890 which is not seen in any of the summer (August) ice records before around 1910 (Fig. 6). Furthermore, the pronounced minimum around 1710 and the peak around 1640 in the BS ice record are not reflected in the Lomo09 MSA record. Thus, our data do not support the connection of MSA at Lomonosovfonna and the BS ice extent stated in O'Dwyer et al. (2000) for the period 1920-

1 1997, nor the assumption of Isaksson et al. (2005) that the MSA sources prior and after 1920
2 were the same, i.e. the BS. We explain the positive correlation of Lomo09 MSA and Western
3 Nordic Sea ice extent as follows. The marginal ice zone is known to be the area of highest
4 DMS production (Perrette et al., 2011). The larger the sea ice area, the more ice edge area is
5 available for phytoplankton growth and thus DMS production. Furthermore, more ice leads to
6 higher freshwater inflow by melting ice. This results in a stronger stratification of the ocean
7 water (Perrette et al., 2011) which keeps the phytoplankton in the euphotic zone. The good
8 correspondence of the Lomo09 MSA record with the Western Nordic Sea ice extent but not
9 with that of the BS is well supported by the findings of Beaudon et al. (2013) pointing to the
10 Greenland Sea as the main source for biogenic related MSA in Svalbard.

11 In the Lomo09 ice core MSA is highly correlated with NO_3^- before around 1900. The records
12 (Figs. 4 and 6) are similar with shared peaks around 1395, 1475, 1560, 1645, 1695, and 1795.
13 The only exception is the period at about 1350 when the correlation breaks down. After
14 around 1900 there is a decoupling of both species with enhanced NO_3^- concentrations from
15 anthropogenic Eurasian NO_x emissions (see above) and strongly decreased MSA
16 concentrations.

17 In previous studies NO_3^- was found to correlate with non-sea-salt- Ca^{2+} (nss- Ca^{2+}) (Legrand et
18 al., 1999; Röthlisberger et al., 2000, 2002), suggesting that nss- Ca^{2+} prevents NO_3^- from being
19 re-emitted from the snowpack. However, those studies are from Greenland, consider glacial
20 timescales, and include e.g. the last glacial maximum (LGM) with much higher nss- Ca^{2+}
21 concentrations. Other studies observed a correlation of NO_3^- and Ca^{2+} in summer and with sea
22 salt in winter but they considered only industrial times (Beine et al., 2003; Geng et al., 2010;
23 Teinilä et al., 2003). The empirical orthogonal function (EOF) analysis performed on the ion
24 data of the Lomo97 core suggests in general no correlation between Ca^{2+} and NO_3^- , but in
25 some parts of the last 200 years the two species are clearly associated (Kekonen et al., 2002).
26 Kekonen et al. (2002) found NO_3^- and NH_4^+ to covariate during the last 100 years. However,
27 the EOF of the whole core did not show a clear association of NH_4^+ and NO_3^- . Nevertheless,
28 they suggested that before 1920 and after 1960 ammonium nitrate (NH_4NO_3) has been
29 common at Lomonosovfonna. They explain this in recent years to be due to Arctic Haze and
30 significant natural sources of NH_4NO_3 during the earlier period. At Holtedahlfonna, Svalbard,
31 NH_4^+ was also associated with NO_3^- before 1880 which Beaudon et al. (2013) interpreted as
32 evidence for NH_4NO_3 to be present. Teinilä et al. (2003) also discovered a correlation of NO_3^-

1 and NH_4^+ in recent times which they concluded to result from anthropogenic emissions. Our
2 data neither support a correlation of NO_3^- and Ca^{2+} , nor of NO_3^- and the sea salt species Na^+ ,
3 nor of NO_3^- and NH_4^+ in pre-industrial times. Instead, they clearly suggest an association of
4 NO_3^- with MSA. Three hypotheses for the high correlation are discussed: (1) post-
5 depositional processes caused by melt water percolation affecting NO_3^- and MSA in the same
6 way, (2) a common source of NO_3^- and MSA, and (3) NO_3^- fertilisation of the ocean which
7 triggers phytoplankton growth and thus DMS and MSA formation.

- 8 1. The pre-industrial record of the melt percent does share some features with NO_3^- and
9 MSA but there is no significant correlation with NO_3^- or MSA ($r^2=0.1$ with either
10 NO_3^- or MSA) (Table 3, Fig. 4). This is also seen in the PCA where the melt percent
11 and the two ionic species have their highest loadings in different PCs (Table 2). Thus,
12 the correlation of NO_3^- and MSA is not a result of similar relocation during melt
13 events on the decadal time scales considered here.
- 14 2. If both species have a common source this would have to be the ocean because MSA
15 results only from marine DMS production and its oxidation in the atmosphere. NO_3^- is
16 only a minor component in sea water with concentrations in the micro-molar range
17 (Chester and Jickells, 2012; Codispoti et al., 2013). The ice core $\text{NO}_3^-/\text{Na}^+$ ratio of
18 ~ 0.066 in the Lomo09 core is up to a factor of ten higher than the sea water ratio of
19 0.006 to 0.038 (Keene et al., 1986). Additionally, we can exclude NO_3^- to be derived
20 from sea spray because NO_3^- and the major sea spray components Na^+ , K^+ , Mg^{2+} , and
21 Cl^- (PC1) do not correlate as seen in the PCA and the correlation analysis (Tables 2
22 and 3). Because of the high solubility of nitrate in water, outgassing of HNO_3 from the
23 ocean is unlikely. Thus, the major NO_3^- source is not the ocean which excludes a
24 common source to cause the strong correlation of NO_3^- and MSA.
- 25 3. Elevated atmospheric NO_3^- concentrations due to high NO_x emissions and/or
26 enhanced transport to the Arctic in the end of winter lead to an increased amount of
27 NO_3^- dissolved in the ocean surface water. Nutrient supply in the Arctic is known to
28 be limited and nitrate depletion is common during the vegetative season (Codispoti et
29 al., 2013). Hence, an increased nitrogen input by dissolved NO_3^- leads to a fertilisation
30 of the phytoplankton (Duce et al., 2008). As soon as light becomes available this
31 results in an enhanced production of DMS and finally higher MSA concentrations in
32 the atmosphere. This process takes weeks to months (Codispoti et al., 2013; Sharma et
33 al., 2012). However, such a potential short time lag cannot be resolved from our data.

1 We therefore propose the fertilising effect to be the dominant cause for the high correlation of
2 NO_3^- and MSA in pre-industrial times. In industrial times the records of NO_3^- and MSA
3 diverge with increasing NO_3^- and decreasing MSA concentrations. This reveals that during
4 the 20th century the effect of decreasing MSA concentrations following reduction in ice extent
5 in the Western Nordic Seas predominates compared to an expected MSA increase caused by
6 enhanced anthropogenic NO_3^- levels.

7 The major NO_3^- source region for the industrial time is Eurasia indicated by the similarity of
8 the NO_3^- records observed in the last 30-40 years in the ice cores from Lomo09, the Siberian
9 Altai, and Severnaya Zemlya (Eichler et al., 2009; Weiler et al., 2005) (Fig. 5). We assume
10 that the source region has not changed from pre-industrial to industrial times. In the period
11 1250-1940 NO_3^- in the Siberian Altai ice core was ascribed to forest fires and mineral dust as
12 main pre-industrial sources (Eichler et al., 2011). That NO_3^- record shows a maximum
13 between 1540 and 1680 (see Fig. 5), attributed to an increased mineral dust input from
14 Central Asian deserts (1540-1600) and enhanced fire activity from Siberian boreal forests
15 (1600-1680). This distinct peak in the 16th and 17th century is not observed in the Lomo09
16 NO_3^- record and also the general pre-industrial records do not correspond well. We cannot
17 exclude that other regional scale NO_3^- sources in Eurasia had a significant impact on the low
18 pre-industrial concentration level. From our data we can therefore not identify major pre-
19 industrial NO_3^- sources for the Lomo09 core.

20 **3.2 Ammonium (NH_4^+)**

21 The Lomo09 NH_4^+ record shows very low concentrations between the 13th and 18th century
22 and an increasing trend from around 1750 onwards (Fig. 7). The values are on the same order
23 of magnitude as those from other Arctic sites and the Lomo97 ice core (Beaudon et al., 2013;
24 Fuhrer et al., 1996; Kehrwald et al., 2012; Kekonen et al., 2005; Legrand and De Angelis,
25 1996; Legrand et al., 1992; Whitlow et al., 1994; Zennaro et al., 2014). The NH_4^+ record of
26 another Svalbard core from Holtedahlfonna, spanning the last 300 years, shows a differing
27 pattern prior to 1800, but a similar strong increasing trend as the Lomo09 record from the 18th
28 century on (Beaudon et al., 2013) (Fig. 4). The authors interpret the rising concentrations
29 from 1880 as result of anthropogenic mid-latitude pollution reaching the Arctic. However, the
30 earlier increase in NH_4^+ concentrations in the Lomo09 and Holte05 ice core from the 18th
31 century on cannot be related to anthropogenic emissions. As discussed above, anthropogenic

1 NH₃ emissions from Eurasia influence precipitation chemistry in Svalbard only after around
2 1940.

3 Pre-industrial NH₄⁺ was not studied in details in the Lomo97 core but Kekonen et al. (2002)
4 suggested NH₄NO₃ to have been common at Lomonosovfonna before 1920. Similarly,
5 Beaudon et al. (2013) postulated that at Hortedahlfonna natural NH₄NO₃ was a common
6 aerosol. Our data do not support this hypothesis since NH₄⁺ and NO₃⁻ are not significantly
7 correlated in pre-industrial times (Tables 2 and 3). In other studies pre-industrial NH₄⁺ was
8 attributed mainly to biomass burning (e.g., Fuhrer et al., 1996; Kehrwald et al., 2010; Legrand
9 et al., 1992; Whitlow et al., 1994). North America and Canada were identified as major
10 sources for NH₄⁺ in Greenland ice (Fuhrer et al., 1996), whereas Legrand and De Angelis
11 (1996) and Zennaro et al. (2014) suggest an additional Eurasian source. A period of
12 exceptional high fire activity around 1600-1680 in Siberian boreal forests of Eurasia was
13 detected in the ice core fire tracer records from the Siberian Altai and Greenland (Eichler et
14 al., 2011; Zennaro et al., 2014). This unique period did not lead to a maximum in the Lomo09
15 NH₄⁺ record. Therefore, we conclude that biomass burning is not a major source for NH₄⁺
16 arriving at Svalbard.

17 The trend in the Lomo09 NH₄⁺ record is similar to that in the ice core from Belukha glacier in
18 the Siberian Altai with increasing concentrations already from around 1750 and very low
19 concentrations between 1680 and 1750 (Eichler et al., 2009) (Fig. 7). Before ~1500 NH₄⁺
20 concentration records agree less, which could be explained by increasing dating uncertainty.
21 At the Belukha site long-term NH₄⁺ variations were related to temperature-induced changes of
22 biogenic NH₃ emissions from extended Siberian boreal forests (Eichler et al., 2009). The
23 strong increase after the 18th century was caused by a rise of Siberian temperatures since that
24 time. Hence, from the similarity in the Lomo09 and Siberian Altai NH₄⁺ concentration records
25 we conclude that biogenic NH₃ emissions from Siberian boreal forests are the dominant
26 source for NH₄⁺ at Lomonosovfonna. Due to the larger distance to the emission sources the
27 NH₄⁺ concentrations in the Lomo09 core are about one order of magnitude lower than in the
28 core from Belukha glacier. The NH₄⁺ concentrations in a Greenland ice core (NEEM, for
29 location see Fig. 1) do not show the increase after the 18th century (Zennaro et al., 2014) (Fig.
30 7), implying that biogenic emission trends in Northern America and Eurasia differ.

31

4 Summary

We presented the 800 year records of the two nitrogen species NO_3^- and NH_4^+ analysed in a new ice core collected from Lomonosovfonna, Svalbard, in 2009. In general, the NO_3^- and NH_4^+ records of the 2009 ice core reasonably agree with published data from two previous Svalbard ice cores, Lomonosovfonna 1997 (Kekonen et al., 2005) and Høltedahlfonna 2005 (Beaudon et al., 2013). On the decadal time scale considered here melt related effects did not significantly alter the concentrations of the nitrogen compounds. Both species show a clear impact of anthropogenic pollution in the 20th century, with peak concentrations in the 1970s/1980s. This temporal trend points to source regions in Eurasia and the Siberian Arctic, since emissions in Northern America and Western Europe kept rising into the 21st century. In pre-industrial times, i.e. prior to the 20th century, the dominant source of NH_4^+ was biogenic NH_3 emissions from Siberian boreal forests. During the same period NO_3^- was highly correlated to MSA on a decadal time scale. We explained this by a fertilising mechanism where higher atmospheric NO_3^- concentrations yield higher nitrogen input to the ocean, triggering the growth of DMS-producing phytoplankton. Elevated DMS concentrations then result in enhanced concentrations of MSA in the atmosphere. Based on our data it was not possible to resolve major pre-industrial NO_3^- sources for Svalbard.

Supplementary material

Here, an example for the annual layer counting (ALC) for the core section between 0 and 20 m weq (Fig. S1) is displayed, followed by the raw data of the ionic species and the ion balance along depth in m weq (Fig. S2) and the annual melt percent (Fig. S3) of the Lomo09 ice core versus age.

Acknowledgements

The drilling was funded by the Paul Scherrer Institut and the Norwegian Polar Institute with help from Serla.

We would like to thank Beat Rufibach[†], and Dieter Stampfli for drilling; NPI field logistics personnel, Mats Björkman, Gerit Rotschky, and Carmen Vega for their help during the expedition; Dmitry Divine for the help on the sea ice data, Richard Petterson for providing the ice depth data, John Moore for providing the Lomo97 ion data, Emilie Beaudon for providing

1 the Holte05 NO_3^- and NH_4^+ data, Leonhard Tobler for analysing ^{210}Pb , and Max Rüthi and
2 Jost Eikenberg for analysing tritium, and Audun Igesund for the help on the map in Fig. 1A.
3 This is a contribution to cryosphere-atmosphere interactions in a changing Arctic climate
4 (CRAICC), a top-level research initiative (TRI).
5

1 **References**

- 2 Abram, N. J., Wolff, E. W. and Curran, M. A. J.: A review of sea ice proxy information from
3 polar ice cores, *Quat. Sci. Rev.*, 79, 168-183, doi:10.1016/j.quascirev.2013.01.011, 2013.
- 4 Adams, P. J., Seinfeld, J. H. and Koch, D. M.: Global concentrations of tropospheric sulfate,
5 nitrate, and ammonium aerosol simulated in a general circulation model, *J. Geophys. Res.*
6 *Atmospheres*, 104, 13791–13823, doi:10.1029/1999JD900083, 1999.
- 7 Baltensperger, U., Schwikowski, M., Gäggeler, H. W., Jost, D. T., Beer, J., Siegenthaler, U.,
8 Wagenbach, D., Hofmann, H. J. and Synal, H. A.: Transfer of atmospheric constituents into
9 an alpine snow field, *Atmos. Environ. A Gen.*, 27, 1881–1890, doi:10.1016/0960-
10 1686(93)90293-8, 1993.
- 11 Beaudon, E., Moore, J. C., Martma, T., Pohjola, V. A., Van De Wal, R. S. W., Kohler, J. and
12 Isaksson, E.: Lomonosovfonna and Holtedahlfonna ice cores reveal east–west disparities of
13 the Spitsbergen environment since AD 1700, *J. Glaciol.*, 59, 1069–1083,
14 doi:10.3189/2013JoG12J203, 2013.
- 15 Becagli, S., Castellano, E., Cerri, O., Curran, M., Frezzotti, M., Marino, F., Morganti, A.,
16 Proposito, M., Severi, M. and Traversi, R.: Methanesulphonic acid (MSA) stratigraphy from a
17 Talos Dome ice core as a tool in depicting sea ice changes and southern atmospheric
18 circulation over the previous 140 years, *Atmos. Environ.*, 43, 1051–1058, 2009.
- 19 Beine, H. J., Dominé, F., Ianniello, A., Nardino, M., Allegrini, I., Teinilä, K. and Hillamo, R.:
20 Fluxes of nitrates between snow surfaces and the atmosphere in the European high Arctic,
21 *Atmos. Chem. Phys.*, 3, 335–346, doi:10.5194/acp-3-335-2003, 2003.
- 22 Bergin, M. H., Jaffrezo, J.-L., Davidson, C. I., Dibb, J. E., Pandis, S. N., Hillamo, R.,
23 Maenhaut, W., Kuhns, H. D. and Makela, T.: The contributions of snow, fog, and dry
24 deposition to the summer flux of anions and cations at Summit, Greenland, *J. Geophys. Res.-*
25 *Atmos.*, 100, 16275–16288, doi:10.1029/95JD01267, 1995.
- 26 Björkman, M. P., Kühnel, R., Partridge, D. G., Roberts, T. J., Aas, W., Mazzola, M., Viola,
27 A., Hodson, A., Ström, J. and Isaksson, E.: Nitrate dry deposition in Svalbard, *Tellus B*, 65,
28 19071, doi:10.3402/tellusb.v65i0.19071, 2013.

1 Brimblecombe, P., Tranter, M., Abrahams, P. W., Blackwood, I., Davies, T. D. and Vincent,
2 C. E.: Relocation and preferential elution of acidic solute through the snowpack of a small,
3 remote, high-altitude Scottish catchment, *Ann. Glaciol.*, 7, 141–147, 1985.

4 Chester, R. and Jickells, T. D.: *Marine Geochemistry*, John Wiley & Sons, Chichester, West
5 Sussex, UK, 2012.

6 Codispoti, L. A., Kelly, V., Thessen, A., Matrai, P., Suttles, S., Hill, V., Steele, M. and Light,
7 B.: Synthesis of primary production in the Arctic Ocean: III. Nitrate and phosphate based
8 estimates of net community production, *Prog. Oceanogr.*, 110, 126–150,
9 doi:10.1016/j.pocean.2012.11.006, 2013.

10 Dickerson, R. R.: Reactive nitrogen compounds in the Arctic, *J. Geophys. Res.-Atmos.*, 90,
11 10739–10743, doi:10.1029/JD090iD06p10739, 1985.

12 Divine, D. V. and Dick, C.: Historical variability of sea ice edge position in the Nordic Seas,
13 *J. Geophys. Res.-Oceans*, 111, C01001, doi:10.1029/2004JC002851, 2006.

14 Divine, D., Isaksson, E., Martma, T., Meijer, H. A. ., Moore, J., Pohjola, V., van de Wal, R. S.
15 W., and Godtliessen, F.: Thousand years of winter surface air temperature variations in
16 Svalbard and northern Norway reconstructed from ice core data, *Polar Res.*, 30, 7379,
17 doi:10.3402/polar.v30i0.7379, 2011.

18 Duce, R. A., LaRoche, J., Altieri, K., Arrigo, K. R., Baker, A. R., Capone, D. G., Cornell, S.,
19 Dentener, F., Galloway, J., Ganeshram, R. S., Geider, R. J., Jickells, T., Kuypers, M. M.,
20 Langlois, R., Liss, P. S., Liu, S. M., Middelburg, J. J., Moore, C. M., Nickovic, S., Oschlies,
21 A., Pedersen, T., Prospero, J., Schlitzer, R., Seitzinger, S., Sorensen, L. L., Uematsu, M.,
22 Ulloa, O., Voss, M., Ward, B. and Zamora, L.: Impacts of atmospheric anthropogenic
23 nitrogen on the open ocean, *Science*, 320, 893–897, doi:10.1126/science.1150369, 2008.

24 Eichler, A., Schwikowski, M., Gäggeler, H. W., Furrer, V., Synal, H.-A., Beer, J., Saurer, M.
25 and Funk, M.: Glaciochemical dating of an ice core from upper Grenzgletscher (4200 m
26 a.s.l.), *J. Glaciol.*, 46, 507–515, doi:10.3189/172756500781833098, 2000.

27 Eichler, A., Schwikowski, M. and Gäggeler, H. W.: Meltwater-induced relocation of chemical
28 species in Alpine firn, *Tellus B*, 53, 192–203, 2001.

- 1 Eichler, A., Brütsch, S., Olivier, S., Papina, T., and Schwikowski, M.: A 750 year ice core
2 record of past biogenic emissions from Siberian boreal forests, *Geophys. Res. Lett.*, 36, ,
3 L18813, doi:10.1029/2009GL038807, 2009.
- 4 Eichler, A., Tinner, W., Brütsch, S., Olivier, S., Papina, T. and Schwikowski, M.: An ice-core
5 based history of Siberian forest fires since AD 1250, *Quat. Sci. Rev.*, 30, 1027–1034,
6 doi:10.1016/j.quascirev.2011.02.007, 2011.
- 7 Eleftheriadis, K., Vratolis, S. and Nyeki, S.: Aerosol black carbon in the European Arctic:
8 measurements at Zeppelin station, Ny-Ålesund, Svalbard from 1998–2007, *Geophys. Res.*
9 *Lett.*, 36, L02809, doi:10.1029/2008GL035741, 2009.
- 10 Eneroth, K., Kjellström, E. and Holmén, K.: A trajectory climatology for Svalbard;
11 investigating how atmospheric flow patterns influence observed tracer concentrations, *Phys.*
12 *Chem. Earth*, 28, 1191–1203, 2003.
- 13 Feng, Y. and Penner, J. E.: Global modeling of nitrate and ammonium: Interaction of aerosols
14 and tropospheric chemistry, *J. Geophys. Res.-Atmos.*, 112, D01304,
15 doi:10.1029/2005JD006404, 2007.
- 16 Fibiger, D. L., Hastings, M. G., Dibb, J. E. and Huey, L. G.: The preservation of atmospheric
17 nitrate in snow at Summit, Greenland, *Geophys. Res. Lett.*, 40, 3484–3489,
18 doi:10.1002/grl.50659, 2013.
- 19 Fischer, H., Wagenbach, D. and Kipfstuhl, J.: Sulfate and nitrate firn concentrations on the
20 Greenland ice sheet: 2. Temporal anthropogenic deposition changes, *J. Geophys. Res.-*
21 *Atmos.*, 103, 21935–21942, doi:10.1029/98JD01886, 1998.
- 22 Forsström, S., Ström, J., Pedersen, C. A., Isaksson, E. and Gerland, S.: Elemental carbon
23 distribution in Svalbard snow, *J. Geophys. Res.-Atmos.*, 114, D19112,
24 doi:10.1029/2008JD011480, 2009.
- 25 Fuhrer, K., Neftel, A., Anklin, M., Staffelbach, T. and Legrand, M.: High-resolution
26 ammonium ice core record covering a complete glacial-interglacial cycle, *J. Geophys. Res.-*
27 *Atmos.*, 101, 4147–4164, doi:10.1029/95JD02903, 1996.

1 Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S.
2 P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F.,
3 Porter, J. H., Townsend, A. R. and Vöosmarty, C. J.: Nitrogen cycles: Past, present, and
4 future, *Biogeochemistry*, 70, 153–226, doi:10.1007/s10533-004-0370-0, 2004.

5 Geng, H., Ryu, J. Y., Jung, H. J., Chung, H., Ahn, K. H. and Ro, C. U.: Single-particle
6 characterization of summertime Arctic aerosols collected at Ny-Ålesund, Svalbard, *Environ.*
7 *Sci. Technol.*, 44, 2348–2353, 2010.

8 Geng, L., Alexander, B., Cole-Dai, J., Steig, E. J., Savarino, J., Sofen, E. D. and Schauer, A.
9 J.: Nitrogen isotopes in ice core nitrate linked to anthropogenic atmospheric acidity change, *P.*
10 *Natl. Acad. Sci. USA*, 111, 5808–5812, doi:10.1073/pnas.1319441111, 2014.

11 Ginot, P., Stampfli, F., Stampfli, D., Schwikowski, M. and Gäggeler, H. W.: FELICS, a new
12 ice core drilling system for high-altitude glaciers, *Mem. Natl. Inst. Polar Res. Spec. Issue*, 56,
13 38–48, 2002.

14 Ginot, P., Schotterer, U., Stichler, W., Godoi, M. A., Francou, B. and Schwikowski, M.:
15 Influence of the Tungurahua eruption on the ice core records of Chimborazo, Ecuador, *The*
16 *Cryosphere*, 4, 561–568, doi:10.5194/tc-4-561-2010, 2010.

17 Goto-Azuma, K. and Koerner, R. M.: Ice core studies of anthropogenic sulfate and nitrate
18 trends in the Arctic, *J. Geophys. Res.*, 106, 4959–4969, 2001.

19 Greenaway, K. R.: Experience with Arctic flying weather, Royal Meteorological Society,
20 Canadian Branch, Toronto, Canada, 1950.

21 Hastings, M. G., Steig, E. J. and Sigman, D. M.: Seasonal variations in N and O isotopes of
22 nitrate in snow at Summit, Greenland: Implications for the study of nitrate in snow and ice
23 cores, *J. Geophys. Res.-Atmos.*, 109, 1984–2012., 2004.

24 Henderson, K., Laube, A., Gäggeler, H. W., Olivier, S., Papina, T. and Schwikowski, M.:
25 Temporal variations of accumulation and temperature during the past two centuries from
26 Belukha ice core, Siberian Altai, *J. Geophys. Res.*, 111, D03104, doi:10.1029/2005JD005819,
27 2006.

1 Hicks, S. and Isaksson, E.: Assessing source areas of pollutants from studies of fly ash,
2 charcoal, and pollen from Svalbard snow and ice, *J. Geophys. Res.-Atmos.*, 111, D02113, ,
3 doi:10.1029/2005JD006167, 2006.

4 Hirdman, D., Sodemann, H., Eckhardt, S., Burkhardt, J. F., Jefferson, A., Mefford, T., Quinn,
5 P. K., Sharma, S., Ström, J. and Stohl, A.: Source identification of short-lived air pollutants in
6 the Arctic using statistical analysis of measurement data and particle dispersion model output,
7 *Atmos. Chem. Phys.*, 10, 669–693, 2010a.

8 Hirdman, D., Burkhardt, J. F., Sodemann, H., Eckhardt, S., Jefferson, A., Quinn, P. K.,
9 Sharma, S., Ström, J. and Stohl, A.: Long-term trends of black carbon and sulphate aerosol in
10 the Arctic: changes in atmospheric transport and source region emissions, *Atmos. Chem.*
11 *Phys.*, 10, 9351–9368, doi:10.5194/acp-10-9351-2010, 2010b.

12 Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B. and Campbell, B.:
13 Evidence of NO_x production within or upon ice particles in the Greenland snowpack,
14 *Geophys. Res. Lett.*, 26, 695–698, doi:10.1029/1999GL900077, 1999.

15 Isaksson, E., Pohjola, V., Jauhiainen, T., Moore, J., Pinglot, J. F., Vaikmaae, R., van de Wal,
16 R. S. W., Hagen, J. O., Ivask, J., Karlöf, L., Martma, T., Meijer, H. A., Mulvaney, R.,
17 Thomassen, M. and van den Broeke, M.: A new ice-core record from Lomonosovfonna,
18 Svalbard: viewing the 1920-97 data in relation to present climate and environmental
19 conditions, *J. Glaciol.*, 47, 335–345, 2001.

20 Isaksson, E., Kekonen, T., Moore, J. and Mulvaney, R.: The methanesulfonic acid (MSA)
21 record in a Svalbard ice core, *Ann. Glaciol.*, 42, 345–351, 2005.

22 Jauhiainen, T., Moore, J., Perämäki, P., Derome, J. and Derome, K.: Simple procedure for ion
23 chromatographic determination of anions and cations at trace levels in ice core samples, *Anal.*
24 *Chim. Acta*, 389, 21–29, 1999.

25 Kaufmann, P., Fundel, F., Fischer, H., Bigler, M., Ruth, U., Udisti, R., Hansson, M., de
26 Angelis, M., Barbante, C., Wolff, E. W., Hutterli, M. and Wagenbach, D.: Ammonium and
27 non-sea salt sulfate in the EPICA ice cores as indicator of biological activity in the Southern
28 Ocean, *Quat. Sci. Rev.*, 29, 313–323, doi:10.1016/j.quascirev.2009.11.009, 2010.

- 1 Keene, W. C., Pszenny, A. A. P., Galloway, J. N. and Hawley, M. E.: Sea-salt corrections and
2 interpretation of constituent ratios in marine precipitation, *J. Geophys. Res.-Atmos.*, 91,
3 6647–6658, doi:10.1029/JD091iD06p06647, 1986.
- 4 Kehrwald, N., Zangrando, R., Gambaro, A. and Barbante, C.: Fire and climate: Biomass
5 burning recorded in ice and lake cores, *EPJ Web Conf.*, 9, 105–114,
6 doi:10.1051/epjconf/201009008, 2010.
- 7 Kehrwald, N., Zangrando, R., Gabrielli, P., Jaffrezo, J.-L., Boutron, C., Barbante, C. and
8 Gambaro, A.: Levoglucosan as a specific marker of fire events in Greenland snow, *Tellus B*,
9 64, 18196, doi:10.3402/tellusb.v64i0.18196, 2012.
- 10 Kekonen, T., Moore, J. C., Mulvaney, R., Isaksson, E., Pohjola, V. and Van De Wal, R. S.
11 W.: A 800 year record of nitrate from the Lomonosovfonna ice core, Svalbard, *Ann. Glaciol.*,
12 35, 261–265, 2002.
- 13 Kekonen, T., Moore, J., Perämäki, P., Mulvaney, R., Isaksson, E., Pohjola, V. and van de
14 Wal, R. S. W.: The 800 year long ion record from the Lomonosovfonna (Svalbard) ice core, *J.*
15 *Geophys. Res.*, 110, D07304, doi:10.1029/2004JD005223, 2005.
- 16 Kellerhals, T., Brütsch, S., Sigl, M., Knüsel, S., Gäggeler, H. W. and Schwikowski, M.:
17 Ammonium concentration in ice cores: A new proxy for regional temperature reconstruction?,
18 *J. Geophys. Res.-Atmos.*, 115, D16123, doi:10.1029/2009JD012603, 2010.
- 19 Kinnard, C., Zdanowicz, C. M., Fisher, D. A., Isaksson, E., de Vernal, A. and Thompson, L.
20 G.: Reconstructed changes in Arctic sea ice over the past 1450 years, *Nature*, 479, 509–512,
21 2011.
- 22 Kühnel, R., Roberts, T. J., Björkman, M. P., Isaksson, E., Aas, W., Holmén, K. and Ström, J.:
23 20-year climatology of NO_3^- and NH_4^+ wet deposition at Ny-Ålesund, Svalbard, *Adv.*
24 *Meteorol.*, 406508, doi:10.1155/2011/406508, 2011.
- 25 Lavigne, F., Degeai, J.-P., Komorowski, J.-C., Guillet, S., Robert, V., Lahitte, P.,
26 Oppenheimer, C., Stoffel, M., Vidal, C. M., Surono, Pratomo, I., Wassmer, P., Hajdas, I.,
27 Hadmoko, D. S. and Belizal, E. de: Source of the great A.D. 1257 mystery eruption unveiled,

- 1 Samalas volcano, Rinjani Volcanic Complex, Indonesia, P. Natl. Acad. Sci. USA, 110,
2 16742–16747, doi:10.1073/pnas.1307520110, 2013.
- 3 Law, K. S. and Stohl, A.: Arctic air pollution: Origins and impacts, Science, 315, 1537–1540,
4 2007.
- 5 Legrand, M.: Ice-core records of atmospheric sulphur, Philos. T. Roy. Soc.. B., 352, 241–
6 250, 1997.
- 7 Legrand, M. and De Angelis, M.: Light carboxylic acids in Greenland ice: A record of past
8 forest fires and vegetation emissions from the boreal zone, J. Geophys. Res.-Atmos., 101,
9 4129–4145, 1996.
- 10 Legrand, M. and Mayewski, P.: Glaciochemistry of polar ice cores: a review, Rev. Geophys.,
11 35, 219–244, 1997.
- 12 Legrand, M., de Angelis, M. and Delmas, R. J.: Ion chromatographic determination of
13 common ions at ultratrace levels in Antarctic snow and ice, Anal. Chim. Acta, 156, 181–192,
14 doi:10.1016/S0003-2670(00)85549-X, 1984.
- 15 Legrand, M., De Angelis, M., Staffelbach, T., Nefel, A. and Stauffer, B.: Large perturbations
16 of ammonium and organic acids content in the summit-Greenland Ice Core. Fingerprint from
17 forest fires?, Geophys. Res. Lett., 19, 473–475, doi:10.1029/91GL03121, 1992.
- 18 Legrand, M., De Angelis, M. and Maupetit, F.: Field investigation of major and minor ions
19 along Summit (Central Greenland) ice cores by ion chromatography, J. Chromatogr. A, 640,
20 251–258, doi:10.1016/0021-9673(93)80188-E, 1993.
- 21 Legrand, M., Hammer, C., Angelis, M. D., Savarino, J., Delmas, R., Clausen, H. and Johnsen,
22 S. J.: Sulfur-containing species (methanesulfonate and SO₄) over the last climatic cycle in the
23 Greenland Ice Core Project (central Greenland) ice core, J. Geophys. Res., 102, 26663–
24 26,679, doi:10.1029/97JC01436, 1997.
- 25 Legrand, M., Wolff, E. and Wagenbach, D.: Antarctic aerosol and snowfall chemistry:
26 implications for deep Antarctic ice-core chemistry, Ann. Glaciol., 29, 66–72,
27 doi:10.3189/172756499781821094, 1999.

- 1 Macias Fauria, M., Grinsted, A., Helama, S., Moore, J., Timonen, M., Martma, T., Isaksson,
2 E. and Eronen, M.: Unprecedented low twentieth century winter sea ice extent in the Western
3 Nordic Seas since AD 1200, *Clim. Dyn.*, 34, 781–795, 2010.
- 4 Matoba, S., Narita, H., Motoyama, H., Kamiyama, K. and Watanabe, O.: Ice core chemistry
5 of Vestfonna ice cap in Svalbard, Norway, *J. Geophys. Res.*, 107, 4721,
6 doi:10.1029/2002JD002205, 2002.
- 7 Moore, J. C. and Grinsted, A.: Ion Fractionation and Percolation in Ice Cores with Seasonal
8 Melting, in *Physics of Ice Core Records II*, vol. 68, edited by: Institute of Low Temperature
9 Science, Hokkaido University, Hokkaido University Press, Sapporo, Japan, 287–298, 2009.
- 10 Moore, J. C., Grinsted, A., Kekonen, T. and Pohjola, V.: Separation of melting and
11 environmental signals in an ice core with seasonal melt, *Geophys. Res. Lett.*, 32, L10501,
12 doi:10.1029/2005GL023039, 2005.
- 13 Moore, J., Kekonen, T., Grinsted, A. and Isaksson, E.: Sulfate source inventories from a
14 Svalbard ice core record spanning the Industrial Revolution, *J. Geophys. Res.*, 111, D15307,
15 doi:10.1029/2005JD006453, 2006.
- 16 Moore, J. C., Beaudon, E., Kang, S., Divine, D., Isaksson, E., Pohjola, V. A. and van de Wal,
17 R. S. W.: Statistical extraction of volcanic sulphate from nonpolar ice cores, *J. Geophys. Res.*,
18 117, D03306, doi:10.1029/2011JD016592, 2012.
- 19 O’Dwyer, J., Isaksson, E., Vinje, T., Jauhiainen, T., Moore, J., Pohjola, V., Vaikmae, R. and
20 van de Wal, R. S. W.: Methanesulfonic acid in a Svalbard ice core as an indicator of ocean
21 climate, *Geophys. Res. Lett.*, 27, 1159–1162, 2000.
- 22 Opel, T., Fritzsche, D. and Meyer, H.: Eurasian Arctic climate over the past millennium as
23 recorded in the Akademii Nauk ice core (Severnaya Zemlya), *Clim. Past*, 9, 2379–2389,
24 doi:10.5194/cp-9-2379-2013, 2013.
- 25 Perrette, M., Yool, A., Quartly, G. D. and Popova, E. E.: Near-ubiquity of ice-edge blooms in
26 the Arctic, *Biogeosciences*, 8, 515–524, doi:10.5194/bg-8-515-2011, 2011.
- 27 Pohjola, V. A., Moore, J. C., Isaksson, E., Jauhiainen, T., Van de Wal, R. S. W., Martma, T.,
28 Meijer, H. A. J. and Vaikmäe, R.: Effect of periodic melting on geochemical and isotopic

1 signals in an ice core from Lomonosovfonna, Svalbard, *J. Geophys. Res.*, 107, ACL 1-1–ACL
2 1-14, doi:10.1029/2000JD000149, 2002.

3 Quinn, P. K., Shaw, G., Andrews, E., Dutton, E. G., Ruoho-Airola, T. and Gong, S. L.: Arctic
4 haze: current trends and knowledge gaps, *Tellus B*, 59, 99–114, 2007.

5 Rhodes, R. H., Bertler, N. A. ., Baker, J. A., Sneed, S. B., Oerter, H. and Arrigo, K. R.: Sea
6 ice variability and primary productivity in the Ross Sea, Antarctica, from methylsulphonate
7 snow record, *Geophys. Res. Lett.*, 36, L10704, doi:10.1029/2009GL037311, 2009.

8 Röthlisberger, R., Hutterli, M. A., Sommer, S., Wolff, E. W. and Mulvaney, R.: Factors
9 controlling nitrate in ice cores: Evidence from the Dome C deep ice core, *J. Geophys. Res.-*
10 *Atmos.*, 105, 20565–20572, doi:10.1029/2000JD900264, 2000.

11 Röthlisberger, R., Hutterli, M. A., Wolff, E. W., Mulvaney, R., Fischer, H., Bigler, M., Goto-
12 Azuma, K., Hansson, M. E., Ruth, U., Siggaard-Andersen, M.-L. and Steffensen, J. P.: Nitrate
13 in Greenland and Antarctic ice cores: a detailed description of post-depositional processes,
14 *Ann. Glaciol.*, 35, 209–216, doi:10.3189/172756402781817220, 2002.

15 Samuelsson, H., Distribution of melt layers on the ice field Lomonosovfonna, Spitsbergen,
16 Master thesis, Uppsala University, 2001.

17 Sharma, S., Chan, E., Ishizawa, M., Toom-Sauntry, D., Gong, S. L., Li, S. M., Tarasick, D.
18 W., Leaitch, W. R., Norman, A., Quinn, P. K., Bates, T. S., Levasseur, M., Barrie, L. A. and
19 Maenhaut, W.: Influence of transport and ocean ice extent on biogenic aerosol sulfur in the
20 Arctic atmosphere, *J. Geophys. Res.*, 117, D12209, doi:10.1029/2011JD017074, 2012.

21 Shaw, G. E.: The arctic haze phenomenon, *B. Am. Meteorol. Soc.*, 76, 2403–2414, 1995.

22 Sigl, M.: Ice core based reconstruction of past climate conditions from Colle Gnifetti, Swiss
23 Alps, Ph.D. thesis, University of Bern, Bern, Switzerland, 2009.

24 Sigl, M., McConnell, J. R., Layman, L., Maselli, O., McGwire, K., Pasteris, D., Dahl-Jensen,
25 D., Steffensen, J. P., Vinther, B., Edwards, R., Mulvaney, R. and Kipfstuhl, S.: A new bipolar
26 ice core record of volcanism from WAIS Divide and NEEM and implications for climate
27 forcing of the last 2000 years, *J. Geophys. Res.-Atmos.*, 118, 1151–1169,
28 doi:10.1029/2012JD018603, 2013.

- 1 Simões, J. C. and Zagorodnov, V. S.: The record of anthropogenic pollution in snow and ice
2 in Svalbard, Norway, *Atmos. Environ.*, 35, 403–413, 2001.
- 3 Stefels, J., Steinke, M., Turner, S., Malin, G. and Belviso, S.: Environmental constraints on
4 the production and removal of the climatically active gas dimethylsulphide (DMS) and
5 implications for ecosystem modelling, *Biogeochemistry*, 83, 245–275, doi:10.1007/s10533-
6 007-9091-5, 2007.
- 7 Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys.*
8 *Res.*, 111, D11306, doi:10.1029/2005JD006888, 2006.
- 9 Stohl, A., Berg, T., Burkhardt, J. F., Fjæraa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder,
10 C., Mcmillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström,
11 J., Tørseth, K., Treffeisen, R., Virkkunen, K., and Yttri, K. E.: Arctic smoke? Record high air
12 pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring
13 2006, *Atmos. Chem. Phys.*, 7, 511–534, doi:10.5194/acp-7-511-2007, 2007.
- 14 Teinilä, K., Hillamo, R., Kerminen, V.-M. and Beine, H. J.: Aerosol chemistry during the
15 NICE dark and light campaigns, *Atmos. Environ.*, 37, 563–575, doi:10.1016/S1352-
16 2310(02)00826-9, 2003.
- 17 Thompson, L. G., Davis, M. E., Mosley-Thompson, E., Sowers, T. A., Henderson, K. A.,
18 Zagorodnov, V. S., Lin, P.-N., Mikhailenko, V. N., Campen, R. K., Bolzan, J. F., Cole-Dai, J.
19 and Francou, B.: A 25 000-year tropical climate history from Bolivian ice cores, *Science*, 282,
20 1858–1864, doi:10.1126/science.282.5395.1858, 1998.
- 21 Tunved, P., Ström, J. and Krejci, R.: Arctic aerosol life cycle: linking aerosol size
22 distributions observed between 2000 and 2010 with air mass transport and precipitation at
23 Zeppelin station, Ny-Ålesund, Svalbard, *Atmos. Chem. Phys.*, 13, 3643–3660,
24 doi:10.5194/acp-13-3643-2013, 2013.
- 25 Udisti, R., Bellandi, S. and Piccardi, G.: Analysis of snow from Antarctica: a critical approach
26 to ion-chromatographic methods, *Fresen. J. Anal. Chem.*, 349, 289–293,
27 doi:10.1007/BF00323205, 1994.

- 1 Van De Wal, R. S. W., Mulvaney, R., Isaksson, E., Moore, J. C., Pinglot, J. F., Pohjola, V. A.,
2 Thomassen, M. P. A.: Reconstruction of the historical temperature trend from measurements
3 in a medium-length borehole on the Lomonosovfonna plateau, Svalbard, *Ann. Glaciol.*, 35,
4 371-378, 2002.
- 5 Vinje, T.: Anomalies and trends of sea-ice extent and atmospheric circulation in the Nordic
6 seas during the period 1864-1998, *J. Climate*, 14, 255–267, 2001.
- 7 Vogt, M. and Liss, P. S.: Dimethylsulfide and Climate, in *Surface Ocean–Lower Atmosphere*
8 *Processes*, edited by: Quéré, C. L. and Saltzman, E. S., American Geophysical Union,
9 Washington, D. C., 197–232, doi:10.1029/2008GM000790, 2009.
- 10 Weiler, K., Fischer, H., Fritzsche, D., Ruth, U., Wilhelms, F. and Miller, H.: Glaciochemical
11 reconnaissance of a new ice core from Severnaya Zemlya, Eurasian Arctic, *J. Glaciol.*, 51,
12 64–74, doi:10.3189/172756505781829629, 2005.
- 13 Whitlow, S., Mayewski, P., Dibb, J., Holdsworth, G. and Twickler, M.: An ice-core-based
14 record of biomass burning in the Arctic and Subarctic, 1750–1980, *Tellus B*, 46, 234–242,
15 1994.
- 16 Wolff, E. W.: Ice sheets and nitrogen, *Philos. T. Roy. Soc. B*, 368, 20130127,
17 doi:10.1098/rstb.2013.0127, 2013.
- 18 Wolff, E. W., Jones, A. E., Bauguitte, S. J.-B. and Salmon, R. A.: The interpretation of spikes
19 and trends in concentration of nitrate in polar ice cores, based on evidence from snow and
20 atmospheric measurements, *Atmos. Chem. Phys.*, 8, 5627–5634, doi:10.5194/acp-8-5627-
21 2008, 2008.
- 22 Zennaro, P., Kehrwald, N., McConnell, J. R., Schüpbach, S., Maselli, O. J., Marlon, J.,
23 Vallelonga, P., Leuenberger, D., Zangrando, R., Spolaor, A., Borrotti, M., Barbaro, E.,
24 Gambaro, A., and Barbante, C.: Fire in ice: two millennia of boreal forest fire history from the
25 Greenland NEEM ice core, *Clim. Past*, 10, 1905-1924, doi:10.5194/cp-10-1905-2014, 2014.

- 1 Table 1 Detection limits and median values [$\mu\text{eq/L}$] for the ions analysed with the
 2 Metrohm 850 Professional IC. Pre-ind. = pre-industrial time from 1222-1859;
 3 Ind. = industrial time from 1860-2009; MSA = CH_3SO_3^- .

Anions	Detection limit	Median		Cations	Detection limit	Median	
		Pre-ind.	Ind.			Pre-ind.	Ind.
MSA	0.005	0.09	0.05	Na⁺	0.02	8.77	7.18
Cl⁻	0.02	10.48	8.92	NH₄⁺	0.02	0.50	0.74
NO₃⁻	0.01	0.54	0.65	K⁺	0.02	0.25	0.19
SO₄²⁻	0.02	2.08	2.63	Mg²⁺	0.03	2.10	1.32
				Ca²⁺	0.04	1.43	1.02

- 4
5

- 1 Table 2 Results of the principal component analysis (PCA) after VARIMAX rotation. Time
 2 period: 1222-1859; data: 10-year averages; MSA = CH_3SO_3^- ; melt% = melt percent.
 3 Values >0.8 marked in bold.

	PC1	PC2	PC3	PC4	PC5	PC6
Na⁺	0.97	0.06	0.05	0.11	-0.03	0.08
K⁺	0.88	0.18	0.00	-0.04	-0.07	0.16
Mg²⁺	0.82	0.37	0.02	0.27	0.07	0.19
Cl⁻	0.97	0.08	0.06	0.12	0.01	0.08
MSA	0.33	0.80	0.13	0.22	0.23	0.11
NO₃⁻	0.11	0.89	0.22	0.16	0.09	0.22
NH₄⁺	0.06	0.23	0.96	-0.02	0.17	0.02
Ca²⁺	0.18	0.27	-0.02	0.92	0.07	0.19
Melt%	-0.05	0.19	0.16	0.07	0.96	0.07
SO₄²⁻	0.29	0.28	0.02	0.21	0.08	0.88
Variance explained [%]	38	19	11	11	11	10

4


5

1 Table 3 R² values of the correlation analysis of the ionic species and the melt percent
 2 (Melt%). Time period: 1222-1859; data: 10-year averages; MSA = CH₃SO₃⁻; 0.5<r²<1
 3 marked in bold.

r ²	Na ⁺	K ⁺	Mg ²⁺	Cl ⁻	MSA	NO ₃ ⁻	NH ₄ ⁺	Ca ²⁺	Melt%	SO ₄ ²⁻
Na ⁺	1									
K ⁺	0.71	1								
Mg ²⁺	0.71	0.59	1							
Cl ⁻	0.98	0.67	0.78	1						
MSA	0.17	0.16	0.41	0.20	1					
NO ₃ ⁻	0.04	0.08	0.27	0.06	0.60	1				
NH ₄ ⁺	0.01	0.01	0.03	0.02	0.14	0.19	1			
Ca ²⁺	0.09	0.06	0.27	0.10	0.26	0.21	0.00	1		
Melt%	0.00	0.00	0.02	0.00	0.15	0.11	0.13	0.04	1	
SO ₄ ²⁻	0.16	0.18	0.33	0.17	0.26	0.26	0.02	0.24	0.04	1

4


5



1

2 Figure 1 (A) Map of Svalbard with the locations of Lomonosovfonna and Holtedahlfonna. (B)
 3 Map with all ice core locations discussed in the text: Lomo = Lomonosovfonna (red triangle);
 4 NEEM, Sum = Summit, SZ = Severnaya Zemlya, Bel = Belukha, and CG = Colle Gnifetti
 5 (black triangles). Satellite image in (B) © PlanetObserver, extracted from DVD-ROM "Der
 6 Große 3D-Globus 4.0 Premium,"#2008 United Soft Media Verlag GmbH, Munich.

7



1


2 Figure 2 Depth-age relationship of the Lomo09 ice core showing all dating methods applied.

3 Depth is given in m weq to account for density variation.

4

5

1




2

3 Figure 3 Temporal records of annual average concentrations of MSA, Na⁺, NO₃⁻ and NH₄⁺

4


and annual melt percent of the Lomo09 ice core.



1

2 Figure 4 Records of NH_4^+ (green), NO_3^- (red), melt percent (light blue), Na^+ (yellow), and
 3 MSA (dark blue) of the Lomo09 ice core. Bold lines are 40-year-lowpass-filtered (40-yr-LP);
 4 dashed lines are 10-year averages (10-yr-avg). Raw data are available in the supplementary
 5 material (Figs. S2 and S3). Grey dashed lines are 40-year-lowpass-filtered records of NH_4^+ ,
 6 NO_3^- , Na^+ , and MSA of the Lomo97 ice core (Kekonen et al., 2005) calculated with the
 7 updated chronology of Divine et al. (2011). Black dashed lines are 40-year-lowpass-filtered
 8 records of NH_4^+ and NO_3^- of the Holte05 ice core (Beaudon et al., 2013).


1



2


3 Figure 5 NO₃⁻ records from Lomo09 (red), Belukha (dark red; Eichler et al., 2009), Summit,
4 Greenland (grey; Geng et al., 2014), and Colle Gnifetti, Swiss Alps (black; Sigl, 2009). Bold
5 lines are 100-year-lowpass-filtered (100-yr-LP); dashed lines are 10-year averages (10-yr-
6 avg).

7



1

2 Figure 6 Records of Lomo09 MSA (dark blue), pre-industrial NO₃⁻ (red), Western Nordic
 3 Seas winter (April) ice extent (turquoise; Macias Fauria et al., 2010), Arctic summer (August)
 4 sea ice extent (light purple; Kinnard et al., 2011), and August sea ice edge position in the
 5 Barents Sea (BS; dark purple; Kinnard et al., 2011). Bold lines are 100-year-lowpass-filtered
 6 (100-yr-LP); dashed lines are 40-year-lowpass-filtered (40-yr-LP).



1

2 Figure 7 NH₄⁺ records of the Lomo09 (green), Belukha (light green; Eichler et al., 2009), and
 3 the NEEM (grey; Zennaro et al., 2014) ice cores. Bold lines are 100-year-lowpass-filtered
 4 (100-yr-LP); dashed lines are the 10-year-averages (10-yr-avg).

5