Investigation of secondary formation of formic acid: urban environment vs. oil and gas producing region

3 Bin Yuan^{1,2}, Patrick R. Veres^{1,2}, Carsten Warneke^{1,2}, James M. Roberts¹, Jessica B. Gilman^{1,2},

4 Abigail Koss^{1,2}, Peter M. Edwards^{1,2,#}, Martin Graus^{1,2,∥}, William C. Kuster^{1,2}, Shao-Meng Li³,

5 Robert J. Wild^{1,2}, Steven S. Brown¹, William P. Dubé^{1,2}, Brian M. Lerner^{1,2}, Eric J. Williams¹,

6 James E. Johnson^{4,5}, Patricia K. Quinn⁵, Timothy S. Bates^{4,5}, Barry Lefer⁶, Patrick L. Hayes^{2,7,¶},

- 7 Jose L. Jimenez^{2,7}, Rodney J. Weber⁸, Robert Zamora¹, Barbara Ervens^{1,2}, Dylan B. Millet⁹,
- 8 Bernhard Rappenglück⁶, and Joost A. de Gouw^{1,2}
- 9 1. Chemical Sciences Division, Earth System Research Laboratory, National Oceanic and
- 10 Atmospheric Administration, Boulder, CO, USA

11 2. Cooperative Institute for Research in Environmental Sciences, University of Colorado at

12 Boulder, Boulder, CO, USA

13 3. Environment Canada, Science and Technology Branch, Toronto, ON, Canada

4. Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle,WA, USA

16 5. NOAA Pacific Marine Environmental Laboratory (PMEL), Seattle, WA, USA

17 6. Department of Earth and Atmospheric Sciences, University of Houston, Houston, TX, USA

- 7. Department of Chemistry and Biochemistry, University of Colorado at Boulder, Boulder, CO,USA
- 8. School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA,
- 21 USA
- 22 9. Department of Soil, Water and Climate, University of Minnesota, St. Paul, Minnesota, USA
- 23 #: Now at: Department of Chemistry, University of York, York, UK
- 24 ||: Now at: Institute of Meteorology and Geophysics, University of Innsbruck, Innsbruck, Austria
- 25 ¶: Now at: Université de Montréal, Department of Chemistry, Montreal, QC, Canada

26

27 Correspondence to: Bin Yuan (bin.yuan@noaa.gov)

28 Abstract:

29 Formic acid (HCOOH) is one of the most abundant carboxylic acids in the atmosphere. However, current photochemical models cannot fully explain observed concentrations and in 30 particular secondary formation of formic acid across various environments. In this work, formic 31 32 acid measurements made at an urban receptor site (Pasadena) in June-July of 2010 during 33 CalNex and a site in an oil and gas producing region (Uintah Basin) in Jan.-Feb. of 2013 during 34 UBWOS 2013 will be discussed. Although the VOC compositions differed dramatically at the two sites, measured formic acid concentrations were comparable: 2.3 ± 1.3 ppb in UBWOS 2013 35 36 and 2.0±1.0 ppb in CalNex. We determine that concentrations of formic acid at both sites were 37 dominated by secondary formation (>99%). A constrained box model using the Master Chemical Mechanism (MCM v3.2) underestimates the measured formic acid concentrations drastically at 38 39 both sites (by a factor of >10). Compared to the original MCM model that includes only ozonolysis of unsaturated organic compounds and OH oxidation of acetylene, when we updated 40 41 yields of ozonolysis of alkenes and included OH oxidation of isoprene, vinyl alcohol chemistry, reaction of formaldehyde with HO₂, oxidation of aromatics, and reaction of CH₃O₂ with OH, the 42 model predictions for formic acid were improved by a factor of 6.4 in UBWOS 2013 and 4.5 in 43 CalNex, respectively. A comparison of measured and modeled HCOOH/acetone ratios is used to 44 45 evaluate the model performance for formic acid. We conclude that the modified chemical 46 mechanism can explain 19% and 45% of secondary formation of formic acid in UBWOS 2013 47 and CalNex, respectively. The contributions from aqueous reactions in aerosol and 48 heterogeneous reactions on aerosol surface to formic acid are estimated to be 0-6% and 0-5% in 49 UBWOS 2013 and CalNex, respectively. We observe that air-snow exchange processes and 50 morning fog events may also contribute to ambient formic acid concentrations during UBWOS 2013 (~20% in total). In total, 53%-59% in UBWOS 2013 and 50%-55% in CalNex of 51 52 secondary formation of formic acid remains unexplained. More work on formic acid formation 53 pathways is needed to reduce the uncertainties in the sources and budget of formic acid and to narrow the gaps between measurements and model results. 54

55 Keywords: formic acid, secondary formation, box model

58 1. Introduction

59 Carboxylic acids are present in the gaseous phase, aqueous phase and in particles (Chebbi and Carlier, 1996). They are significant contributors to rain acidity in remote environments (16-60 65%) and they regulate aqueous reactions with pH-dependence in cloud (Khare et al., 1999;Vet 61 62 et al., 2014). Some higher carboxylic acids are proposed to enhance new particle formation in the 63 atmosphere (Zhang et al., 2004). These organic acids also play important roles in secondary organic aerosol (SOA) formation (Carlton et al., 2006). Research on the sources and sinks of 64 65 carboxylic acids is needed to understand the processes in acid rain, new particle formation and SOA formation, all of which are integral to our understanding of regional air quality and climate 66 change. 67

68 Formic acid (HCOOH) is the simplest organic acid, and is one of the most abundant 69 carboxylic acids detected in the atmosphere (Khare et al., 1999). The sources of formic acid are emissions from vehicle exhausts (Kawamura et al., 2000), biomass burning (Akagi et al., 2011), 70 biogenic activities (Jardine et al., 2011), and secondary formation from the oxidation of volatile 71 organic compounds (VOCs) (Khare et al., 1999), e.g. oxidation of alkenes by ozone (Neeb et al., 72 73 1997). Aqueous reactions of formaldehyde (Chameides and Davis, 1983), glyoxal (Carlton et al., 74 2007) and other species also produce formic acid. Previous studies have proposed that organic 75 aerosol aging by heterogeneous reactions with OH radical is also an important source of formic 76 acid (Molina et al., 2004; Paulot et al., 2011). The global sources of formic acid are thought to be 77 dominated by photochemical oxidation of biogenic VOCs (Paulot et al., 2011). Recent work also 78 indicated that secondary formation was the largest contributor to formic acid in polluted air in 79 the summertime (de Gouw et al., 2005; Veres et al., 2011), even though primary emissions may 80 account for a larger fraction in the wintertime (Bannan et al., 2014). Thus, studies focused on 81 secondary formation of formic acid will be helpful to better understand the oxidation chemistry 82 of anthropogenic and biogenic VOCs (Paulot et al., 2011). While dominantly present in the gas phase, formic acid appears to be present in aerosols at higher than expected concentrations (Liu 83 et al., 2012; Yatavelli et al., 2014), although instrument artifacts might play a role in those 84 85 measurements.

The diversity of emission sources, formation pathways and precursors of formic acid makes it challenging to fully understand its primary sources and secondary formation in the atmosphere.

88 Modeling studies showed that observed formic acid concentrations in both urban plumes (Le 89 Breton et al., 2014) and biogenically dominated areas (Paulot et al., 2011) could not be explained 90 by current chemical mechanisms (Table S1). Comparisons between satellite measurements and global three-dimensional modeling results indicate that formic acid is underpredicted in many 91 92 regions of the world, especially in tropical and boreal forests (Stavrakou et al., 2012), and in the 93 summertime and over biomass burning regions (Cady-Pereira et al., 2014). To address this 94 underestimation, many new formation pathways for formic acid have been proposed in recent 95 years, such as OH oxidation of isoprene (Paulot et al., 2009b) and formation from vinyl alcohol 96 (Andrews et al., 2012).

97 In this study, we show that formic acid concentrations are at comparable levels in two 98 different environments: (1) an urban downwind site and (2) a site in an oil and gas producing 99 region, even though the VOC composition is completely different. A box model constrained by 100 measurements will be used to simulate the secondary formation of formic acid and to evaluate 101 the recently proposed formation pathways of formic acid. Contributions from condensed phase 102 sources of formic acid will also be investigated.

103 2. Measurements and methods

104 **2.1 UBWOS campaigns**

Datasets collected from ground sites in three different campaigns are used in this study:
Horse Pool during the Uintah Basin Winter Ozone Studies (UBWOS) in 2012 and 2013
(Edwards et al., 2013) and Pasadena ground site during the California Research at the Nexus of
Air Quality and Climate Change (CalNex) campaign in 2010 (Ryerson et al., 2013).

Two campaigns at the Horse Pool site (40.1428 N, 109.4680 W) in the Uintah Basin, Utah were conducted in January - February of 2012 and of 2013, respectively. The Uintah Basin, where over 10000 active oil and gas wells are located, has started to experience severe ozone problems during wintertime in recent years. Measurements in 2012 occurred from Jan. 15 to Feb. 29, but no ozone episode was encountered, due to unusually warm conditions and a lack of ground snow cover (Edwards et al., 2013). The second campaign was performed from Jan. 25 to Feb. 22, 2013 and very high ozone concentrations were observed during this campaign (Edwards

et al., 2014). This work will focus on the dataset collected during the 2013 study, since
secondary formation of formic acid was more prominent than during the 2012 study.

118 In the two UBWOS campaigns, formic acid in the ambient air was measured using a negative-ion proton-transfer chemical ionization mass spectrometry (NI-PT-CIMS) using acetate 119 (CH₃COO⁻) as the reagent ion (Veres et al., 2008). Calibrations of formic acid were performed in 120 the field using diluted gas standards generated from permeation tubes. Formic acid 121 122 concentrations of these sources were determined by catalytically converting to CO_2 and 123 subsequently measuring using a CO₂ detector (Veres et al., 2010). Instrument backgrounds were measured every 2-3 hours by passing ambient air through a platinum (Pt) catalytic converter 124 125 maintained at 350 °C. Measurement accuracy of formic acid using NI-PT-CIMS is estimated to 126 be better than 25%, propagated from the uncertainties of permeation tube concentrations, 127 calibration and variations of background signals. Nitric acid (HNO₃) was also measured by NI-128 PT-CIMS during UBWOS 2013.

129 During UBWOS 2013, C2-C7 alkanes, C2-C3 alkenes, acetylene and benzene were measured by a gas chromatograph with flame ionization detection (GC-FID) (Bon et al., 2011). 130 Aromatics and selected oxygenates were measured by a proton transfer reaction mass 131 132 spectrometer (PTR-MS). A custom-built four-channel cavity ring down spectrometry instrument (NOx-CaRD) was used to measure ozone (O₃) and nitrogen oxides (NOx, including NO and NO₂) 133 (Wild et al., 2014). PAN and nitryl chloride (ClNO₂) were measured by a CIMS with iodide (I⁻) 134 135 as the reagent ion (Slusher et al., 2004). A cavity ring-down spectroscopy system was used to detect nighttime NO₃ and N₂O₅ in the atmosphere (Dubé et al., 2006). Methane was measured by 136 137 a commercial cavity ring-down spectrometry instrument (Picarro G2301). A Scanning Mobility 138 Particle Sizer (SMPS, TSI model 3081) and an Aerodynamic Particle Size (APS, TSI model 3321) were used to measure the number size distribution of aerosols. Filter samples were 139 collected and analyzed by a Sunset Laboratory thermal/optical analyzer for organic carbon (OC) 140 and by ion chromatography (IC) for nitrate, sulfate, ammonium and chloride. Measurement of 141 142 meteorological parameters, including temperature, relative humidity, wind direction, wind speed, precipitation, downwelling and upwelling solar radiation were made at the Horse Pool site by 143 144 NOAA ESRL PSD. During UBWOS 2013, the Uintah Basin was covered by snow with an

average depth of 14±4 cm. Snow samples were collected (top 5 cm) and the chemical
composition in snow was measured by ion chromatography from melted snow water.

To assist data interpretation in this study, some measurements from UBWOS 2012 will also be used. During UBWOS 2012, C2-C10 hydrocarbons and many oxygenates were measured by an online two-channel gas chromatograph mass spectrometer (GC-MS) (Gilman et al., 2013). Additionally, photolysis frequencies of O_3 and NO_2 were measured by a filter radiometer only during the 2012 campaign. Therefore, the aforementioned 2012 data will be used to estimate unmeasured concentrations of some hydrocarbons and photolysis frequencies during the 2013 study.

154 **2.2 CalNex 2010 campaign**

Measurements at the Pasadena site during CalNex were conducted from May 15 - June 15, 2010 on the campus of the California Institute of Technology (34.1406 N, 118.1225 W). The measurements at this site sampled outflow from Los Angeles (LA). A suite of state-of-the-art instruments was deployed at the Pasadena site (Ryerson et al., 2013).

159 During CalNex, the same NI-PT-CIMS instrument as in the UBWOS campaigns was used to measure formic acid; the CalNex dataset has been previously reported by Veres et al. (2011). 160 161 Measurements of hydrocarbons and oxygenates were performed by online GC-MS (Gilman et al., 2013) and proton-transfer-reaction ion-trap mass spectrometry (PIT-MS) (Warneke et al., 2005). 162 163 An analyzer based on the Hantzsch reaction was used to measure formaldehyde (HCHO) (Warneke et al., 2011). NOx (NO and NO₂) and ozone were measured by commercial 164 165 chemiluminescence gas analyzers (Thermo 42i-TL and 42i-TL with blue light converter) and a UV absorbance analyzer (Thermo 49c), respectively. Photolysis frequencies of NO₂ were 166 167 derived from filter radiometer measurements. A particle into liquid sampler (PILS) coupled with a total organic carbon (TOC) analyzer was used to measure water-soluble organic carbon 168 169 (WSOC) (Zhang et al., 2012). Aerosol size distributions were measured by an SMPS (TSI model 3936). 170

171 **2.3 Description of box model**

172 The Dynamically Simple Model of Atmospheric Chemical Complexity (DSMACC) 173 (Emmerson and Evans, 2009) is used to simulate secondary formation of formic acid in this 174 study. Hydrocarbons, NOx (including NO₃ and N₂O₅), ozone, methane, and formaldehyde are constrained to their average measured diurnal profiles throughout the simulation period of the 175 176 zero-dimensional box model for each campaign, and the box model is run toward a diurnal steady state (DSS). Unmeasured VOC species during UBWOS 2013 are calculated from VOCs 177 178 measured in 2013 and their respective enhancement ratio measured in 2012. The VOC pairs used 179 for the calculations are generally chosen to have similar reactivity and/or similar structures. For 180 example, 2-methylpentane in 2013 is calculated from n-hexane concentrations measured in 2013 and the 2-methylpentane/n-hexane ratios measured in 2012. Photolysis frequencies are also 181 182 scaled from the 2012 measurements and the inferred surface albedo from measurements of 183 downwelling and upwelling solar radiations (see (Edwards et al., 2014) for details).

184 The model is typically run for ten days, after which the simulated diurnal profiles of formic acid and other photochemical products (e.g. acetaldehyde and acetone) change little compared to 185 186 the previous day (Edwards et al., 2013) (Fig. S2). Modeled diurnal profiles of formic acid and 187 other related species (mainly acetone) in the last day will be shown in this study. We note that 188 primary emissions of formic acid and other photochemical products (e.g. acetone) are not 189 prescribed in the box model. The box model output will be compared to the calculated secondary 190 concentrations by subtracting the primary part. Chemical mechanisms for measured VOC species 191 and other inorganic species are extracted from the Master Chemical Mechanism (MCM) v3.2 192 website (http://mcm.leeds.ac.uk/MCM) (Jenkin et al., 2012). ClNO₂ chemistry is included in 193 MCM v3.2 and measured cycloalkanes are lumped into cyclohexane (the only cycloalkane in 194 MCM v3.2), following previous work (Edwards et al., 2013). A first-order physical loss term is used in the box model to characterize the processes of dilution due to mixing with background 195 air and/or deposition. A physical loss rate of 1.15×10^{-5} s⁻¹, corresponding to a lifetime of 24 196 hours, is applied in the model runs for both campaigns. This lifetime due to physical losses is 197 198 consistent with the setup used in simulations of other similar box models, e.g. (Edwards et al., 199 2013;Li et al., 2014;Fried et al., 2008;Fried et al., 2003). A sensitivity study for physical loss rates in the box model will be performed (Section 3.4). 200

201 **3. Results and discussions**

202 **3.1** Comparisons of formic acid in two different campaigns

Figure 1 shows the measured concentrations of formic acid from the CalNex and UBWOS campaigns. The average (arithmetic mean) concentrations of formic acid over the entire campaigns were 2.0±1.0 ppb and 2.3±1.3 ppb in CalNex and UBWOS 2013, respectively. Similar concentration ranges (from sub-ppb level to 8-10 ppb) were observed during the two campaigns. Diurnal variations of formic acid during the two different campaigns are also shown in Fig. 1. Higher formic acid concentrations are observed during the daytime in both of the campaigns.

210 In a previous paper, Veres et al. (2011) conducted diurnal profile analysis and correlations 211 of formic acid with other compounds and the authors concluded that formic acid at the Pasadena site in CalNex was dominated by secondary formation. This finding is also valid for the UBWOS 212 213 2013 campaign. The evidence includes: (1) There were very few concentration spikes in the 214 measured time series of formic acid that would indicate a local, primary source of formic acid. In 215 contrast, concentration spikes of hydrocarbons (e.g. benzene) were detected frequently at Horse 216 Pool site due to primary emissions from nearby oil and natural gas wells (Fig. S1) (Warneke et al., 2014). Measurements by the PTR-MS in a mobile laboratory sampling downwind oil and gas 217 wells also showed little enhancement of the formic acid signal (m/z 47, HCOOH•H⁺) (Warneke 218 219 et al., 2014). (2) Multi-day accumulation patterns of formic acid during stagnation events (e.g. 220 Jan. 29 - Feb. 9, 2013) are most similar to species with predominantly secondary sources (e.g. 221 acetone and ozone), but are different from species with primary emissions (e.g. benzene) (Fig. 2). 222 (3) Formic acid during UBWOS 2013 increased by a factor of 4 compared to measurements in UBWOS 2012 when photochemistry was weak (Edwards et al., 2013). Most hydrocarbons 223 224 showed enhancements in 2013 from 2012 by a factor of 2-3, due to shallower boundary layer 225 heights in 2013 and more stagnant meteorological conditions. The different enhancements 226 observed between primary species and secondary products from 2012 to 2013 reflect the fact that 227 primary compounds are affected linearly by mixing and dilution processes in the boundary layer, whereas photochemical formation of secondary products is non-linear. 228

The dominance of secondary formation for formic acid makes it hard to accurately estimate the contribution from primary sources. The potential primary sources of formic acid in CalNex are mainly vehicular emissions. In addition to vehicular emissions, other combustion sources

232 related to oil and gas extractions, e.g. compressors, dehydrators and pump jacks, can also 233 contribute to primary emissions of formic acid in UBWOS campaigns. As shown in Table 1, 234 large ranges of emission ratios of formic acid to combustion tracers are reported in the literature. Bannan et al. (2014) reported a HCOOH/CO emission ratio of 1.2 ppb/ppm based on wintertime 235 236 observations in London, whereas no direct emissions of formic acid were detected in the Northeastern U.S. (de Gouw et al., 2005). The most comprehensive and latest study showed that 237 HCOOH/CO emission ratios from gasoline vehicles are $42.2\pm30.0\times10^{-3}$ ppb/ppm during hot 238 running and $10.2\pm6.5\times10^{-3}$ ppb/ppm during cold start (Crisp et al., 2014). Here, the HCOOH/CO 239 emission ratios ($42.2\pm30.0\times10^{-3}$ ppb/ppm) obtained in Crisp et al. (2014) for gasoline vehicles 240 during hot running are used to determine primary contributions to formic acid in the two 241 campaigns. For UBWOS, we note the caveat that HCOOH/CO emission ratios related to oil and 242 gas extractions may be different from those in urban regions. Because CO was not measured 243 during UBWOS 2013, acetylene (C₂H₂), another common combustion tracer, is used instead for 244 the analysis. Utilizing the emission ratio of 5.78 ppb/ppm for C_2H_2/CO at the Pasadena site 245 (Borbon et al., 2013), the emission ratios of $HCOOH/C_2H_2$ from combustion source are 246 247 calculated. Formic acid concentrations from combustion sources are determined from the HCOOH/C₂H₂ emission ratio and the measured acetylene concentrations. The calculations show 248 249 that emissions from combustion sources only accounted for $0.46\pm0.32\%$ and $0.63\pm0.45\%$ of formic acid in CalNex and UBWOS 2013, respectively. This shows that primary emissions only 250 251 contributed very minor parts to formic acid concentrations in both CalNex and UBWOS 2013. Using the same procedure, we determine that primary emissions from combustion sources 252 253 accounted for 1.0±0.7% of formic acid in UBWOS 2012, although photochemistry was weaker in 2012 compared to 2013. 254

255 Ozonolysis of unsaturated species and OH oxidation of acetylene are included as the only two formation pathways for formic acid in most previous modeling studies (Le Breton et al., 256 2014; Ito et al., 2007; Le Breton et al., 2012), as well as in MCM v3.2. Acetylene is a ubiquitous 257 258 species in the atmosphere, but the reaction of acetylene with OH radicals is rather slow $(k_{OH}=7.8\times10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}, 298 \text{ K} \text{ and } 1 \text{ bar})$ (Atkinson et al., 2006). Thus, unsaturated 259 species are the most important precursors of formic acid in MCM v3.2 in polluted environments. 260 Figure 3 shows the measured concentration ratios of various VOCs in the UBWOS campaigns 261 (2012 and 2013) relative to those measured in CalNex. The concentrations of alkanes were much 262

263 higher (5-60 times) in the UBWOS campaigns than in CalNex, mainly due to large emissions from oil and gas production in the Uintah Basin (Helmig et al., 2014). In contrast, levels of 264 265 alkene and other unsaturated species were much lower in UBWOS than those in CalNex, especially for biogenic species (e.g. isoprene and its oxidation products). Aromatics are generally 266 267 higher for UBWOS compared to CalNex. Thus, the much lower concentrations of alkenes and other unsaturated species in UBWOS 2013 compared to CalNex would imply a lower formic 268 269 acid concentration in UBWOS 2013 if alkene ozonolysis was the main secondary formation 270 pathway. However, formic acid levels are similar at the two different locations. This disagreement between measurements and expectation from known chemistry will be investigated 271 using the box model described in Section 2.3. 272

273 **3.2 Base box model run**

274 MCM v3.2 was extracted from the official website (http://mcm.leeds.ac.uk/MCM) and used in the box model. This run is referred as the base case. The modeled formic acid diurnal steady 275 276 state concentrations for UBWOS 2013 and CalNex are shown in Fig. 4 (also Fig. S2). Modeled 277 average formic acid concentrations are 0.05±0.003 ppb and 0.18±0.02 ppb for UBWOS 2013 and CalNex, respectively. The higher modeled formic acid concentrations in CalNex are 278 279 consistent with higher levels of alkenes that react with ozone to produce formic acid. However, 280 the modeled formic acid concentrations are 40 and 13 times lower than the measurements for UBWOS 2013 and CalNex, respectively. Modeled formic acid concentrations are higher in the 281 282 daytime and lower at night, as expected. But, the modeled formic acid concentrations are highest in late afternoon (around 6 pm, local time) for both campaigns, in contrast to the measurements 283 284 that show broad afternoon peaks. This is mainly due to the constant physical loss rates that are 285 used to represent the processes of dilution and deposition. The transport of air masses from downtown of Los Angeles to Pasadena site during noontime (10 am -2 pm) (Veres et al., 2011) 286 could be another reason for the different peak time between measurements and model results in 287 CalNex. This issue about diurnal profile patterns in the box model will be discussed again in 288 Section 3.4. 289

3.3 Modifications to MCM mechanisms

To investigate the large underestimation of formic acid concentration in the base model run, a thorough examination of MCM v3.2 and a literature review for formation pathways of formic acid has been conducted. Based on these results, some recent findings incorporated in the box model are:

295 (1) Formic acid yields of ozonolysis of alkenes and other unsaturated species in MCM v3.2 296 are compared with literature values (Table 2). Even though production of formic acid from 297 ozonolysis of these unsaturated compounds is represented in MCM v3.2, the yields in MCM v3.2 298 are lower than literature values by various factors (as high as 77% for methyl vinyl ketone, MVK), with the exception of β -pinene. Formation of formic acid from O₃ oxidation occurs via 299 300 reaction of Criegee intermediate (CI) biradicals with H₂O. The CH₂OO radical also reacts with 301 CO, SO₂, NO and NO₂, which compete with the formation of formic acid. CH₂OO is formed from seven excited biradicals (CHOOA, CHOOB, CHOOC, CHOOD, CHOOE, CHOOF and 302 303 CHOOG), which originate from different alkenes and unsaturated compounds based on the degree of alkyl substitution (Saunders et al., 2003). These excited biradicals undergo 304 305 decomposition (producing CO, HO₂ and OH), isomerization (producing CO and H₂O) or stabilization (producing CH₂OO). The branching ratios among decomposition, isomerization and 306 307 stabilization determine the yields of formic acid from the seven different groups of species 308 (Table S2). Branching ratios of the three pathways from seven excited biradicals in MCM v3.2 309 are modified either using values reported in literature when available (such as for ethene (Alam et al., 2011)), or by matching yields in the modified MCM with the reported yields in the 310 311 literature (Table S2). The yield of methacrolein (MACR) is not modified, since the difference 312 (10%) between the yields in the literature and in MCM v3.2 is small. The yields of the two 313 monoterpenes (α -pinene and β -pinene) also remain unchanged, since the two compounds were at 314 low levels in both campaigns (45±40 ppt in CalNex, below detection limit in UBWOS). The 315 overestimated yield for β -pinene and underestimated yield for α -pinene also partially cancel out 316 the differences. It should be noted that the modifications of branching ratios here also affect the 317 yields of other products (e.g. formaldehyde) and use of these numbers determined here in other studies should be done with caution. It is not an issue in this study, as formaldehyde has been 318 319 constrained using measurements.

320 (2) OH oxidation of isoprene and the subsequent products can lead to formation of formic 321 acid, but this is not included in MCM v3.2 (Table 2). A recent chamber study showed that OH 322 oxidation of isoprene forms formic acid with a yield of 10%, with a significant share of the yield attributed to the oxidation of glycolaldehyde and hydroxyacetone (Paulot et al., 2009a). Earlier 323 324 studies also showed that formic acid is formed from photo-oxidation of glycolaldehyde (Butkovskaya et al., 2006a) and hydroxyacetone (Butkovskaya et al., 2006b) with yields of 18% 325 326 and 8%, respectively. However, the findings of glycolaldehyde and hydroxyacetone as precursors of formic acid are questioned by another study (Orlando et al., 2012). Other second-327 generation reactions may also contribute to formic acid formation, including δ -hydroxy isoprene 328 nitrates (Paulot et al., 2009a), hydroperoxy methylbutenals (HPALDs) (Stavrakou et al., 2012) 329 330 and epoxides (IEPOX) (Bates et al., 2014). Considering the complexity of isoprene chemistry, a 331 detailed update of isoprene chemistry that includes all secondary reactions producing formic acid was beyond the scope of this study. Alternatively, the reported effective yield of formic acid 332 333 from isoprene photooxidation in Paulot et al. (2009a) is used as the benchmark. After including the formation of formic acid from OH oxidation of glycolaldehyde and hydroxyacetone in MCM 334 335 v3.2 to match the reported yields in Butkovskaya et al. (2006a;2006b), the effective yield of formic acid from isoprene oxidation in the modified MCM v3.2 is 8%, which is slightly lower 336 337 than the reported value (10%) in Paulot et al. (2009a). OH oxidation of β -pinene and acetylene show only small differences between the literature values and MCM v3.2, and therefore no 338 339 change is made for these two species. The modifications of O₃ and OH oxidation of alkenes and other unsaturated species discussed above will be referred as the "modified alkenes" case. 340

341 (3) Vinyl alcohol (CH₂=CHOH) has been suggested to be a precursor of formic acid when it 342 is oxidized by OH radicals (Archibald et al., 2007;So et al., 2014). Vinyl alcohol is formed through tautomerization of acetaldehyde by photolysis (Andrews et al., 2012). Organic acids (da 343 Silva, 2010) and inorganic acids (Karton, 2014) can catalyze the tautomerization processes 344 345 between acetaldehyde and vinyl alcohol. Here, both photo-induced and organic-acids catalyzed tautomerization is incorporated in the box model (Table S3). The tautomerization catalyzed by 346 inorganic acids is not included since the rate coefficients are not available. In MCM v3.2, vinyl 347 348 alcohol is produced from the photolysis of 3-hydroxy-cyclohexanone and it further reacts with OH to form glycolaldehyde and an HO₂ radical. In this study, the oxidation mechanisms of vinyl 349 alcohol proposed by Archibald et al. (2007) and So et al. (2014) are used to in place of the MCM 350

v3.2 default with the two cases referred as "*VINOH from Archibald*" and "*VINOH from So*",
respectively.

(4) Reactions of HOCH₂OO, a product from the reaction of formaldehyde (HCHO) with HO₂ radicals, also contribute to formic acid formation (Jenkin et al., 2007;Atkinson et al., 2006). The equilibrium constant between HOCH₂OO and HCHO+HO₂ (5.3×10^{-16} cm³ molecule⁻¹ at 298K and 1.6×10^{-14} cm³ molecule⁻¹ at 263K) is much larger at low temperature (Atkinson et al., 2006). As a result, the reactions of HOCH₂OO are more important during UBWOS 2013 due to the low ambient temperatures (-8.0±4.0 °C). This modification is referred as the "*HCHO/HO*₂" case.

360 (5) Many studies have shown that formic acid is formed from OH oxidation of aromatics (Berndt et al., 1999;Berndt and Böge, 2001;Baltensperger et al., 2005;Wyche et al., 2009) (Table 361 S4). The reported yields of formic acid range from 2% to 13% for various aromatics. The yields 362 found in the literature are highly variable, not only among different species, but also for a single 363 364 species (e.g. 1,3,5-trimethylbenzene). Formic acid is not treated as a product from oxidation of aromatics in MCM v3.2. Here, a yield of 10% is applied to all of the aromatics included in MCM 365 v3.2. We note that the yields used here for aromatics should be near an upper limit under real 366 atmospheric conditions. This modification will be referred as the "modified aromatics" case. 367

(6) Several studies have proposed that the reaction of CH₃O₂ with OH might be an important 368 source of formic acid (Archibald et al., 2009;Fittschen et al., 2014). A recent measurement 369 confirms that this reaction can occur with a relatively high rate constant $(2.8\pm1.4\times10^{-10} \text{ cm}^3)$ 370 molecule⁻¹ s⁻¹) (Bossolasco et al., 2014). Reaction of CH₃O₂ with OH radicals may proceed in 371 three pathways with different products: CH₂O₂+H₂O, CH₃O+HO₂ and CH₃OH+O₂. Formic acid 372 production from CH₂O₂ radicals only occurs via the first of those three pathways. The branching 373 ratio to the first pathway ranges between 49% (Maricq et al., 1994) and 91% (Daele and Poulet, 374 1996), both based on branching ratio measurements for the reaction of CH₃O₂ with chlorine 375 376 radicals (Cl) as reference for respective OH reaction. Here, a unity branching ratio was used to 377 simulate the upper limit of the contribution from this reaction to formic acid. This modification will be referred to as the " CH_3O_2 " case. 378

The only chemical sink of formic acid in MCM v3.2 is reaction with the OH radical (k_{OH} = 4.5×10⁻¹³ cm³ molecule⁻¹ s⁻¹, independent of temperature). A recent study proposed that formic acid react with Criegee radicals with rate coefficients in excess of 1×10⁻¹⁰ cm³ molecule⁻¹ s⁻¹ (Welz et al., 2014). All of the cases examined here include reaction with OH radicals as the sole chemical loss pathway. A sensitivity test to the newly proposed sink due to Criegee radicals will be conducted separately.

Simulated results from the six modified cases are shown in Fig. 4 (for magnified lower ranges of the plots, refer to Fig. S3). For UBWOS 2013, the biggest improvement to the modeled formic acid concentration comes from the inclusion of aromatics as precursors of formic acid. Other modified cases in UBWOS 2013 somewhat increase formic acid concentrations, but to a much smaller extent. Unlike UBWOS 2013, modifications to the alkenes mechanisms, inclusion of aromatics as precursors and one variant of vinyl alcohol chemistry significantly enhance the modeled formic acid concentrations during the CalNex study.

392 The different responses of modeled formic acid to various formation pathways in UBWOS and CalNex are due to the different environments and VOC emission patterns in the two 393 campaigns. The vinyl alcohol oxidation mechanism proposed by So et al. (2014) results in larger 394 formic acid production than when using mechanism proposed by Archibald et al. (2007), because 395 396 So et al. (2014) estimated a much higher rate constant for the reaction of vinyl alcohol with OH radical (k_{OH} =6.8×10⁻¹¹ cm³ molecule⁻¹ s⁻¹ at 298 K versus k_{OH} = 6.0×10⁻¹² cm³ molecule⁻¹ s⁻¹ by 397 Archibald et al. (2007)). The contribution of vinyl alcohol chemistry to formic acid formation in 398 399 CalNex is noticeably larger than for UBWOS 2013, since modeled OH concentrations during CalNex are higher $(1.5 \times 10^6 \text{ molecule cm}^{-3} \text{ versus } 2.9 \times 10^5 \text{ molecule cm}^{-3} \text{ in UBWOS } 2013, 24 \text{ -h}$ 400 average) and therefore OH oxidation of vinyl alcohol is more competitive with the 401 tautomerization processes back to acetaldehyde. The modest contribution of vinyl alcohol to 402 formic acid formation in the two campaigns is consistent with a simple calculation from a global 403 scale perspective (Muller and Peeters, 2014). The HCHO+HO₂ reaction slightly increases the 404 405 modeled formic acid concentration during UBWOS 2013, but only contributes a very small amount in CalNex because of the higher ambient temperatures in Los Angeles (18.4±4.6 °C). 406 The reaction of CH₃O₂ with OH is not an important contributor to formic acid in either 407 campaigns, because the dominant sink of CH₃O₂ is through reaction with NO and NO₂ at the 408

observed NOx levels (4.3 ± 4.1 ppb in UBWOS 2013 and 14.8 ± 8.6 ppb in CalNex) during the two campaigns. However, this reaction could provide a persistent source for formic acid, as CH₃O₂ is produced through reaction of methane with OH radicals.

All of the modified cases for formation pathways of formic acid are combined together and 412 the results are also shown in Fig. 4. Note that the oxidation mechanism of vinyl alcohol from So 413 et al. (2014) is used here, as it is the latest one and the results are based on quantum chemical 414 415 calculations. Combining all of the modifications in the mechanisms, the modeled formic acid concentrations for UBWOS 2013 and CalNex increase to 0.32±0.05 ppb and 0.81±0.12 ppb, 416 respectively. Modeled formic acid concentrations are enhanced by a factor of 6.4 in UBWOS 417 418 2013 and a factor of 4.5 in CalNex, compared to the base model case in Section 3.2. However, 419 despite the large enhancements of modeled formic acid concentrations, these concentrations are still significantly lower than the measurements. A sensitivity model run with the reaction of 420 421 formic acid with Criegee radicals was conducted in the box model. Inclusion of reaction with Criegee radicals reduces modeled concentrations of formic acid by 20.0% in UBWOS 2013 and 422 423 17.4% in CalNex, which is the combined effect of higher sink and lower formation rates from 424 Criegee radicals. It implies that the discrepancy between measurement and model may be even 425 higher, if the reaction of formic acid with Criegee radicals occurs as proposed.

426 **3.4 Quantification of box model performance**

For all of the simulations in Section 3.3, all of the non-chemical losses due to physical 427 dilution and/or deposition are parameterized using a first-order physical loss rate of 1.15×10^{-5} s⁻¹ 428 for formic acid and other species (corresponding to a lifetime of 24 hours due to physical losses). 429 430 However, atmospheric processes of both turbulent/entrainment mixing with background air and dry deposition are difficult to parameterize and the physical loss rate due to such processes can 431 432 exhibit large day-to-day and diurnal variations (e.g. due to diurnal changes of boundary layer height). It is also acknowledged that meteorological conditions during UBWOS 2013 and 433 434 CalNex were quite different, which can result in different physical loss rates.

In order to evaluate the model sensitivity to the physical loss rate in our box model, larger and smaller physical loss rates $(2.3 \times 10^{-5} \text{ s}^{-1} \text{ or a lifetime of } 12 \text{ hours}; 5.75 \times 10^{-6} \text{ s}^{-1} \text{ or a lifetime}$ of 48 hours) were applied in the box model to investigate the model sensitivity to the physical

loss rates. As shown in Fig. 5, a longer lifetime due to the physical losses results in larger 438 439 modeled formic acid, and vice versa. The variation of physical loss rate by a factor of 2 would 440 change the modeled concentrations of formic acid by a factor of 2.0-2.1 in UBWOS 2013 and 2.2-2.3 in CalNex. This phenomenon is consistent with the fact that the sinks of formic acid 441 442 during both campaigns are dominated by physical losses and that the chemical losses of formic acid are slow. The lifetime of formic acid with respect to reaction with the OH radical is 87 days 443 444 and 18 days in UBWOS 2013 and CalNex, respectively. The sensitivity tests show that including the reaction with Criegee radicals reduces the lifetime of formic acid with respect to chemical 445 losses to 5.3 days in UBWOS 2013 and 5.4 days in CalNex. In either case, chemical losses of 446 447 formic acid are slow compared with the physical losses (dilution and deposition) in both 448 campaigns.

449 As a test of the diurnal steady state (DSS) method using a constant physical loss rate in the 450 box model, an emission-based box model simulation that utilizes emission rate terms to reproduce concentrations of primary species was conducted for UBWOS 2013 (for details refer 451 452 to (Edwards et al., 2014)). A bi-modal physical loss rate (higher in daytime and lower at night) is 453 applied to reflect variations in mixing rates with background air at different times of the day. The 454 results of the emission-based box model associated with the modified MCM v3.2 for formic acid 455 in UBWOS 2013 are shown in Fig. S4. Besides the emission-based box model, a bi-modal 456 physical loss rate is applied to the simulation of the DSS method in UBWOS 2013 (Fig. 5). It is clear that the bi-modal physical loss rate is able to simulate diurnal variations of formic acid 457 458 better than constant physical loss rates.

459 In summary, physical loss rate in the model do affect both the modeled absolute 460 concentrations and the diurnal profile of formic acid significantly, as the chemical loss of formic acid is very slow. Physical loss rates in the box model not only influence formic acid, but also 461 other secondary products (e.g. acetone, Fig. 5). We are able to obtain reasonable agreements 462 463 between measurements and model results for formic acid in both magnitudes and diurnal profiles 464 by "tuning" the physical loss rates in the box model during the two campaigns. However, 465 simulation results for other compounds (e.g. acetone, Fig. 5) from the box model would then be 466 much higher than the measurements. To account for the effects of physical loss processes, scatter plots of formic acid versus acetone from box model simulations are shown in Fig. 6. Acetone is 467

468 selected, since (1) acetone was measured in both campaigns; (2) photochemical degradation of acetone, including reactions with OH radical (k_{OH} =1.7 x 10⁻¹³ cm³ molecule⁻¹ s⁻¹ at 298 K) 469 (Atkinson and Arey, 2003) and photolysis in MCM v3.2, is slow; (3) acetone can be modeled 470 well using MCM (or other similar chemical mechanisms) for the UBWOS 2013 data (Edwards et 471 472 al., 2014) and in urban emission outflows (Sommariva et al., 2011; Apel et al., 2010). Good linear correlations between modeled formic acid and acetone are found for all tests of physical loss and 473 474 the slopes are independent of the physical loss rates. The slopes in UBWOS 2013 and CalNex are 0.05 ppb/ppb and 0.21 ppb/ppb, respectively. 475

476 We acknowledge that the treatment of dilution and deposition in the box model by combining the two terms and assuming the same physical loss rate for different species may affect the 477 478 modeled slopes of formic acid to acetone. (1) The background air that dilutes modeled air parcels in the box model contains no formic acid and acetone. A test simulation that assumes the 479 480 modeled air parcel is diluted by background air with 0.1 ppb of formic acid (Paulot et al., 2011) and 0.5 ppb of acetone (Hu et al., 2013) for UBWOS 2013 is shown in Fig. S5. Very small 481 482 changes of the simulated slope of formic acid versus acetone are observed (~4%), compared with the simulation assuming background air without formic acid and acetone. (2) Deposition 483 484 velocities for various species can be different. Based on the parameterization of deposition 485 velocity (Wesely, 1989), the more soluble formic acid likely has a faster deposition rate than 486 acetone. Accounting for this difference in deposition velocities, the modeled slope of formic acid versus acetone would be even lower than those shown in Fig. 6. 487

Scatter plots of measured formic acid versus acetone are compared with model results in Fig. 6, where diurnal average data are shown. The measured enhancement ratios are 0.25 ppb/ppb in UBWOS 2013 and 0.42 ppb/ppb in CalNex, which are much larger than the modeled slopes. It should be noted that morning data points (7:30 am - 12:00 pm) are excluded from the fit for UBWOS 2013, because the data at this time of day may have been influenced by fog events, which will be discussed in Section 3.5. Linear regressions using all ambient data points rather than diurnal averages produce only slightly different enhancement ratios (Fig. S6).

Using the measured and modeled enhancement ratios of formic acid to acetone (Fig. 6), we estimate that the modified MCM v3.2 can explain 20% and 50% of formic acid concentrations in UBWOS 2013 and CalNex, respectively. The box-model simulations do not account for primary

498 emissions for both formic acid and acetone, thus concentrations due to secondary production for 499 the two species are calculated by subtracting the primary concentrations (see discussions in 500 Section 3.1) from the measured concentrations. This slightly changes the enhancement ratios in Fig. 6 and the explainable fractions of formic acid by modified MCM v3.2 changes to 19% in 501 502 UBWOS 2013 and 45% in CalNex, respectively. As emission compositions of hydrocarbons 503 were not found to be different between UBWOS 2012 and UBWOS 2013, the performance of 504 the emission-base box model can also represent the conditions in UBWOS 2012. As shown in 505 Fig. 6, we observe a similar explainable fraction of formic acid in UBWOS 2012 as that in 506 UBWOS 2013 by the box model.

507 Based on box model simulations with the modified MCM v3.2, the formation rates of 508 formic acid are 3.2 ppb/day in UBWOS 2013 and 8.2 ppb/day in CalNex. We determine that 509 additional formation rates of formic acid of 13.6 ppb/day and 10.0 ppb/day are required to 510 achieve agreement between box model simulations and measurements in UBWOS 2013 and CalNex, respectively. The additional formation rates of formic acid are equivalent to additional 511 512 ethene concentrations of 30.3 ppb in UBWOS 2013 and 19.1 ppb in CalNex (daily-average), 513 which are 14.4 and 10.6 times the measured ethene concentrations in the two campaigns 514 (UBWOS 2013: 2.1±0.2 ppb; CalNex: 1.8±0.2 ppb). Alternatively, the additional formation rate 515 of formic acid is equivalent to 1.3 ppb of isoprene for the CalNex study, which is again much 516 larger than the measured isoprene concentration $(0.33\pm0.32 \text{ ppb})$. Studies have shown that not all species are measured in the atmosphere by GC-MS, based on either OH reactivity measurements 517 518 (Di Carlo et al., 2004) or more advanced measurement techniques (Lewis et al., 2000). However, 519 the additional concentrations needed for alkenes to reproduce the observed formic acid 520 concentrations are much larger than the fractions of potentially unmeasured species in the 521 atmosphere.

We evaluated the chemical mechanisms of both alkenes and aromatics and included them in MCM v3.2. However, alkanes (including cycloalkanes), the most abundant compounds in UBWOS 2013 due to the emissions of oil and gas extraction (e.g. ethane: 305 ± 30 ppb, propane: 141 ± 14 ppb), are not included. Laboratory and field measurements showed that photooxidation of >C5-alkanes produces substituted dihydrofurans, which can react with ozone to form carboxylic acids (Zhang et al., 2014;Russell et al., 2011). However, it is unclear whether formic

acid is produced. Assuming that alkanes were precursors of formic acid, yields of 6.9% and 21.3%

from alkane oxidation could explain the missing formic acid source in UBWOS 2013 and

530 CalNex, respectively. It is hard to determine the relevance of the calculated yields to real

atmospheric conditions due to lack of information. The calculated yields seem to be high

532 compared with other compounds considered in this study (e.g. aromatics, 0-10%).

533 **3.5** Contributions from aqueous and heterogeneous reactions

534 The box model simulations shown in Section 3.3 only account for formation pathways of 535 formic acid in the gas phase. Previous studies showed that formic acid can also be formed from 536 aqueous-phase oxidation of formaldehyde (Chameides and Davis, 1983), glyoxal (Carlton et al., 537 2007), methylglyoxal (Tan et al., 2010) and glycolaldehyde (Perri et al., 2009). Formic acid formed in the aqueous phase can enter the gas phase through gas-particle partitioning or 538 539 evaporation of water in cloud/aerosol. Formic acid can also be produced through heterogeneous 540 reactions of aerosol with OH radicals (Vlasenko et al., 2008; Eliason et al., 2004) and ozone 541 (Thomas et al., 2001;Dubowski et al., 2004).

The rate of aqueous reactions between OH radicals and dissolved organic carbon (DOC)can be expressed as:

$Rate = k_{c,OH} \times [OH] \times [DOC]$

where $k_{c,OH}$ is the rate constant for the reaction, and [OH] and [DOC] are the concentrations of 544 OH radicals and DOC in the aqueous phase. The rate constant $k_{c,OH}$ is adopted as $3.8\pm1.9\times10^8$ L 545 $(mol C)^{-1} s^{-1}$ and [OH] is taken as 10^{-15} M for atmospheric particles (Arakaki et al., 2013). 546 547 Measured organic carbon (OC) in UBWOS 2013 and water-soluble organic carbon (WSOC) in 548 CalNex are used as surrogates for DOC concentration in particles, respectively. Reaction rates 549 are calculated to be 0.11±0.05 ppb C/day for UBWOS 2013 and 0.06±0.04 ppb C/day for CalNex. Assuming all of the carbon fluxes from DOC oxidation converts to formic acid and 550 551 transfers to gas phase, formation rates of formic acid from aqueous reactions are only 0.11±0.05 ppb/day for UBWOS 2013 and 0.06±0.04 ppb/day for CalNex. It should be noted that formic 552 acid is also consumed rapidly by OH radicals in the aqueous phase $(3 \times 10^9 \text{ M}^{-1} \text{ s}^{-1})$ and formic 553 acid (pKa=3.74 at 298 K) will only transfer efficiently to the gas phase when pH values in the 554 aqueous phase are low. By comparing the formation rates from aqueous reactions and those 555

needed to reproduce formic acid concentrations (Table 3), we conclude that aqueous reactions

557 contribute little to formic acid concentration in both UBWOS 2013 (0-0.7%) and CalNex (0-

558 0.3%).

559 The heterogeneous reaction rate (*Rate*) of oxidants on aerosol surface is determined from 560 the collision flux of oxidants with aerosol and the uptake coefficient (γ):

$$Rate = \frac{1}{4} \times v \times S \times [O_x] \times \gamma$$

where v is the molecular velocity (m/s), calculated as $(8RT/\pi M)^{0.5}$ (R is the universal gas 561 constant, T is the temperature, and M is the molecular weight of the oxidant) (Kwan et al., 2006). 562 S is the ambient aerosol surface area $(cm^2 cm^{-3})$. The dry surface area of aerosol is calculated 563 from measurements of aerosol size distributions. Wet ambient aerosol surface area is determined 564 by taking account for hygroscopic diameter growth, using the reported hygroscopicity parameter 565 for CalNex (κ =0.37) (Hersey et al., 2013) and a derived hygroscopicity parameter in UBWOS 566 2013 (κ =0.31) based on the measured chemical compositions of aerosol. [Ox] is the oxidant 567 concentration. Here we consider two different oxidants: OH radical and ozone. The value of γ for 568 OH radical is taken as unity (Bertram et al., 2001; Abbatt et al., 2012). The uptake coefficient γ 569 for ozone is estimated from the reported dependence of γ on ozone concentrations (McCabe and 570 Abbatt, 2008) (4.1 \pm 0.5 \times 10⁻⁵ in UBWOS 2013 and 1.1 \pm 0.5 \times 10⁻⁵ in CalNex). Here, diurnal 571 averaged data for aerosol surface areas and concentrations of oxidant are used for the 572 573 calculations. Heterogeneous reaction rates of OH radicals with aerosol are calculated to be 0.06 ppb/day in UBWOS 2013 and 0.33 ppb/day in CalNex, whereas reaction rates of ozone with 574 575 aerosol are 12±2 ppb/day (UBWOS 2013) and 13±4 ppb/day (CalNex) (Table 3). If yields of formic acid from heterogeneous reactions (e.g. <5% for alkenes with ozone (Thomas et al., 2001)) 576 577 are considered, formation rates of formic acid would then be less than 0.6±0.1 ppb/day in UBWOS 2013 and less than 0.7±0.2 ppb/day in CalNex. Thus, our best estimates for formation 578 579 rates of formic acid from heterogeneous reactions should be <1 ppb/day, which are not large 580 contributions to the formic acid formation for either UBWOS 2013 (0-6%) or CalNex (0-5%). It 581 should be noted that the calculations in the estimates of heterogeneous reactions are associated 582 with large uncertainties, from both uptake coefficients and the yields of formic acid from the reactions. Future studies are warranted to provide better constraints on the contributions from 583

584 heterogeneous reactions.

585 Besides aerosols, several morning fog events (e.g. Feb. 2, Feb. 3 and Feb. 7) occurred during UBWOS 2013 that could potentially serve as reaction media for aqueous and heterogeneous 586 reactions contributing to formic acid formation. Formic acid (measured as formate) has been 587 identified as one of the main components of organic matter in fog droplets (Herckes et al., 2013). 588 589 Figure 7 shows the time series of formic acid and other VOCs during a fog event on the morning 590 of Feb. 7, 2013. During this fog event, maximum concentrations of formic acid (up to 10.3 ppb) in UBWOS 2013 were recorded between 7:30-9:00 am. Concentrations of other VOCs, including 591 benzene, acetaldehyde, acetone and acetic acid, also increased during this period, indicating a 592 593 more polluted air mass was encountered. However, enhancements in concentrations of both 594 acetaldehyde and acetone were lower than those of formic acid (and acetic acid recorded by 595 PTR-MS). Larger enhancements of formic acid relative to acetone during fog events on Feb. 2 596 and Feb. 3 were also observed (not shown). It is also clear that formic acid and acetic acid 597 decreased more than other VOCs in the early morning (6:00-7:30 am). Time variations of 598 formaldehyde are similar with two carboxylic acids, but different from acetaldehyde and acetone. 599 This may reflect the dynamic absorption and release processes for these highly soluble species 600 (formic acid, acetic acid and formaldehyde) to and from fog droplets in the fog event. Due to the 601 lack of chemical measurements of fog water, we are not able to conclude whether formaldehyde 602 contributes to formic acid enhancement during the fog events (Chameides and Davis, 1983;Keene et al., 1995). The different behavior of formic acid and acetone in the fog events 603 604 lead to deviation from the otherwise high correlations between the two compounds in the 605 morning (Fig. 6 and Fig. S6). Here, linear fit results from non-morning data points in Fig. 6 are 606 used to estimate formic acid concentration without the influence from fog events, and the 607 difference between measured and estimated formic acid concentrations in the morning can be 608 attributed to fog events (Fig. S7). Using this approach, fog events in the morning in UBWOS 609 2013 are determined to enhance the campaign-averaged concentration of formic acid by $4\pm7\%$. 610 This contribution is not a large source for formic acid for the four-week campaign, but fog 611 formation accounted for significant formic acid concentrations in certain periods (e.g. the 612 morning of Feb. 7).

613 **3.6 Contribution of Air-snow exchange**

614 As mentioned earlier, the Uintah Basin was covered by snow during UBWOS 2013. The 615 processes of air-snow exchange may provide another pathway for secondary formation of formic 616 acid in the atmosphere (Jacobi et al., 2004; Dibb and Arsenault, 2002). Measured formic acid concentrations in pore spaces of the snowpacks (firn air) are much higher than ambient air and 617 618 formic acid may be formed from oxidation of carbonyls and alkenes within snowpack (Dibb and 619 Arsenault, 2002). During UBWOS 2013, sampling inlets of both NI-PT-CIMS and PTR-MS 620 were mounted on a small tower that can move inlet heights between 1.0 m and 7.5 m. The inlet height was changed every 20 min from Feb. 7 to Feb. 16, 2013 to obtain concentration gradients 621 622 of formic acid, acetone, and other measured compounds.

623 Here, the concentration gradient is defined as the averaged concentration of each 20 min 624 interval measured at 1.0 m minus the averaged concentration measured at 7.5 m in the preceding 625 and subsequent cycles. Thus, positive (negative) vertical gradients indicate upward (downward) 626 flux from (to) the snow surface. The calculated concentration gradients of formic acid and acetone are shown in Fig. 8. We did not observe a clearly gradient direction for acetone during 627 628 Feb. 7-16. The averaged concentration gradient observed during the time was 0.07±0.37 ppb (average \pm standard deviation). However, two different periods showed strong gradients of 629 630 formic acid: Feb. 7-12 with a negative $(-0.21\pm0.29 \text{ ppb})$ and Feb. 12-16 with a positive gradient $(0.22\pm0.25 \text{ ppb})$, both of the values are significantly different from zero (p<0.01). Statistical tests 631 632 also indicate that the gradient differences between the two periods are significant for formic acid (p<0.01), but not for acetone. The different formic acid gradients in the two periods might be due 633 634 to varying chemical compositions in the snow, possibly as the results of the snow event on Feb. 9. Unfortunately, formate data in the snow was not available during UBWOS 2013. Measured ions 635 636 in the snow during UBWOS 2013 include oxalate and many other inorganic ions (e.g. nitrate). As the only measured organic ion in the snow, oxalate may be used as a proxy for formate, since 637 formate and oxalate in the snow strongly correlate with each other (Norton, 1985). As shown in 638 639 Fig. 8, neither time variations of oxalate in the snow nor nitric acid concentrations in ambient air 640 correlated well with the concentration gradients of formic acid (R=0.20 and R=0.21, respectively), whereas the concentration products of oxalate in the snow and nitric acid in 641 642 ambient air show a reasonable correlation with formic acid gradients (R=0.58). This suggests that deposition of nitric acid to the snow surface and the acid displacement reactions due to nitric 643 acid, which is thermodynamically favored (HNO₃ + HCOO⁻ = NO₃⁻ + HCOOH, $\Delta_r G^\circ$ =-41 kJ/mol) 644

(Lide, 2005), may play important roles in the air-snow exchange of formic acid (and other organic
acids). However, we cannot rule out other chemical processes and physical mechanisms that may
account for air-snow exchange of formic acid in UBWOS 2013. Concurrent measurements of
formate in snow, formic acid in ambient air and in firn air, and other chemical compositions in
snow would be needed to answer this question.

650 Differences in vertical gradients for formic acid and acetone could be used to investigate the 651 importance of air-snow exchange to formic acid concentrations in UBWOS 2013. The downward 652 flux before Feb. 12 and upward flux after Feb. 12 of formic acid should cause different 653 enhancement ratios of formic acid to acetone in the two periods. The measurements are in 654 support of this statement (Fig. S8): enhancement ratios of 0.285 ppb/ppb before Feb. 12 and 655 0.337 ppb/ppb after Feb. 12 were observed. Thus, the difference in enhancement ratios between 656 the two periods $(18\pm1\%)$ can be considered an upper limit for the contribution of air-snow 657 exchange to formic acid.

658 **3.7 Summary for both gas phase and non-gas phase processes**

659 Figure 9 shows the fractional contributions of various formation pathways to secondary 660 formation of formic acid, including both gas phase reactions and other non-gas phase processes. Note that the upper limits of contributions from aerosol (aqueous reactions and heterogeneous 661 662 reactions) are used in Fig. 9. Combining all of the processes and considering the lower and upper limits of contributions from aerosol-related reactions, current knowledge could explain 41%-47% 663 664 and 45%-50% of the secondary formation of formic acid in UBWOS 2013 and CalNex, respectively. Inclusion of the non-gas phase processes helped narrow the gap of formic acid 665 666 sources significantly, especially for UBWOS 2013 (from 81% to as low as 53%). Even though the explained fractions of formic acid in UBWOS 2013 and CalNex are comparable, the 667 668 processes producing formic acid are quite different. In CalNex, gas phase reactions (according to box model results) are much more important than aerosol-related production. This is in contrast 669 670 to UBWOS 2013, where aerosol processes, air-snow exchange and fog events may be account for significant contributions. In the gas phase, ozonolysis of alkenes, OH oxidation of isoprene 671 672 (13%) and OH oxidation of aromatics (12%) are all important in CalNex, whereas OH oxidation 673 of aromatics (12%) dominates gas-phase contribution to formic acid formation in UBWOS 2013.

674 Note that Fig. 9 should be viewed as the most "optimistic" case for formic acid formation in 675 both UBWOS 2013 and CalNex. The upper limits are used to determine the fractions from many 676 formation pathways in the production of formic acid, e.g. yields of formic acid from oxidation of aromatics. The newly proposed reaction with Criegee radicals may provide an additional sink of 677 678 formic acid and the sink can reduce the modeled formic acid by $\sim 20\%$ in the two campaigns, 679 which is also not reflected in Fig. 9. Our treatment of deposition by assuming formic acid has 680 same deposition velocities as acetone may overestimate the modeled ratio of formic acid to acetone and consequently, the percentages explained by the box model. One exception is that the 681 formic acid yield from OH oxidation of isoprene used in this study is 20% lower than the 682 literature value, which would only increase the fraction of isoprene reaction with OH radicals in 683 684 Fig. 9 to 16% in CalNex. The reason of providing the most "optimistic" case in Fig. 9 is that many of the formation pathways included in the gas phase box model or non-gas phase processes 685 686 are associated with large uncertainties. For example, all of the knowledge on vinyl alcohol oxidation producing formic acid comes from theoretical calculations without any evidence from 687 688 direct measurements. Additional work on essentially all of the processes discussed in this study 689 would be helpful to reduce the uncertainties in our understanding of secondary formic acid sources. In addition, half of observed secondary formation of formic acid are still unexplained in 690 691 both campaigns, thus explorations of new formation pathways for formic acid are needed to 692 accurately reproduce the observed formic acid concentrations.

693 **4. Conclusions**

Formic acid was measured at an urban receptor site during CalNex and a site in an oil and gas producing region during UBWOS 2013. Secondary formation was the main source of formic acid during the two campaigns. Formic acid concentrations were comparable at both sites, even though the VOC composition was very different from each other.

A box model is used to simulate secondary formation of formic acid for the two campaigns. The original chemical mechanisms derived from MCM v3.2 gave very large discrepancies between measured and modeled formic acid (lower by a factor of 40 in UBWOS 2013 and a factor of 13 in CalNex). Chemical mechanisms for formation pathways of formic acid reported in many recent studies are incorporated into MCM v3.2 for the box model, including updated yields of ozonolysis of alkenes, OH oxidation of isoprene, vinyl alcohol chemistry, reaction of

formaldehyde with HO₂, oxidation of aromatics and reaction of CH_3O_2 with OH. The updated chemical mechanisms increase the modeled formic acid concentrations significantly, by a factor of 6.4 for the UBWOS 2013 case and a factor of 4.5 for the CalNex case. Based on correlations of formic acid with acetone from both measurement and model results, the influences from physical losses that are hard to represent in a box model are taken into account. We determine that the box model using an updated chemical mechanism can explain 19% and 45% of secondary formation of formic acid in UBWOS 2013 and CalNex, respectively.

711 Besides gas phase reactions, contributions from aerosol-related reactions, fog events and airsnow exchange are also evaluated. Aerosol related reactions (including aqueous and 712 heterogeneous reactions) may account for 0-6% and 0-5% of formic acid concentration in 713 714 UBWOS 2013 and CalNex, respectively. Fog events and air-snow exchange in UBWOS 2013 contribute additional small fractions (~20% in total) to formic acid concentrations. Adding up all 715 716 of the pathways, 41%-47% and 45%-50% of secondary formation of formic acid can be 717 explained in UBWOS 2013 and CalNex, respectively. However, the dominant formic acid 718 sources in UBWOS 2013 and CalNex are completely different, which is a result of the different 719 environments and atmospheric compositions for the two locations.

The Pasadena site investigated in this study during the summertime CalNex campaign is 720 721 downwind of a major urban area. Considering the similar VOC compositions across different mega-cities (Borbon et al., 2013), secondary formation of formic acid in the urban plumes of 722 other cities should demonstrate similar results shown in Fig. 9. Formic acid in urban plumes in 723 724 winter may be more influenced by primary emissions, as discussed in a recent paper (Bannan et al., 2014). However, the wintertime UBWOS 2013 campaign at the Horse Pool site provides a 725 726 good opportunity to investigate secondary formation of formic acid without either significant 727 primary emissions or biogenic influence. The UBWOS 2013 case is unique, since 728 unconventional photochemistry at Horse Pool in winter helped to promote formation of secondary products (Edwards et al., 2014), including formic acid. Comparisons between the two 729 730 different sites have helped to better understand formic acid secondary formation. Nevertheless, 731 more studies on formation pathways of formic acid, including those discussed in this study and 732 new possible routes, are needed to narrow the gap between measurement and model results.

734 Acknowledgement

- 735 This work was supported in part by the Western Energy Alliance. This work was also supported
- by the NOAA Health of the Atmosphere Program and by the NOAA Climate Program Office –
- 737 Atmospheric Composition and Climate Program. We thank the contribution from Colm Sweeney
- (NOAA ESRL GMD) to methane data in UBWOS 2013. Bin Yuan acknowledges support from
- 739 National Research Council (NRC) Research Associateship Programs (RAP). Patrick Hayes and
- 740 Jose-Luis Jimenez thank a CIRES Visiting Fellowship and funding from CARB (11-305) and
- 741 NOAA NA13OAR4310063. Dylan Millet acknowledges support from NSF Grant #AGS-
- 742 1148951.
- 743
- 744
- 745

746 **References:**

- Abbatt, J. P. D., Lee, A. K. Y., and Thornton, J. A.: Quantifying trace gas uptake to tropospheric aerosol:
- recent advances and remaining challenges, Chem Soc Rev, 41, 6555-6581, 2012.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and
- Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models,
- 751 Atmos. Chem. Phys., 11, 4039-4072, 10.5194/acp-11-4039-2011, 2011.
- Alam, M. S., Camredon, M., Rickard, A. R., Carr, T., Wyche, K. P., Hornsby, K. E., Monks, P. S., and
- Bloss, W. J.: Total radical yields from tropospheric ethene ozonolysis, Phys Chem Chem Phys, 13,
 11002-11015, 10.1039/c0cp02342f, 2011.
- Andrews, D. U., Heazlewood, B. R., Maccarone, A. T., Conroy, T., Payne, R. J., Jordan, M. J. T., and
- 756 Kable, S. H.: Photo-tautomerization of Acetaldehyde to Vinyl Alcohol: A Potential Route to
- 757 Tropospheric Acids, Science, 10.1126/science.1220712, 2012.
- Apel, E. C., Emmons, L. K., Karl, T., Flocke, F., Hills, A. J., Madronich, S., Lee-Taylor, J., Fried, A.,
- 759 Weibring, P., Walega, J., Richter, D., Tie, X., Mauldin, L., Campos, T., Weinheimer, A., Knapp, D., Sive,
- 760 B., Kleinman, L., Springston, S., Zaveri, R., Ortega, J., Voss, P., Blake, D., Baker, A., Warneke, C.,
- Welsh-Bon, D., de Gouw, J., Zheng, J., Zhang, R., Rudolph, J., Junkermann, W., and Riemer, D. D.:
- 762 Chemical evolution of volatile organic compounds in the outflow of the Mexico City Metropolitan area,763 Atmos. Chem. Phys., 10, 2353-2375, 2010.
- Adinos. Chem. Phys., 10, 2555-2575, 2010.
 Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S.,
- Arakaki, T., Anastasio, C., Kuroki, T., Nakajina, H., Okada, K., Kotani, T., Handa, D., Azechi, S.,
 Kimura, T., Tsuhako, A., and Miyagi, Y.: A General Scavenging Rate Constant for Reaction of Hydroxyl
- Radical with Organic Carbon in Atmospheric Waters, Environmental Science & Technology, 47, 8196-
- 767 8203, 10.1021/es401927b, 2013.
- 768 Archibald, A. T., McGillen, M. R., Taatjes, C. A., Percival, C. J., and Shallcross, D. E.: Atmospheric
- transformation of enols: A potential secondary source of carboxylic acids in the urban troposphere,
- 770 Geophys. Res. Lett., 34, L21801, 10.1029/2007gl031032, 2007.

- Archibald, A. T., Petit, A. S., Percival, C. J., Harvey, J. N., and Shallcross, D. E.: On the importance of
- the reaction between OH and RO2 radicals, Atmospheric Science Letters, 10, 102-108, 10.1002/asl.216,
 2009.
- Atkinson, R., and Arey, J.: Atmospheric degradation of volatile organic compounds, Chemical Reviews,
- 775 103, 4605-4638, Doi 10.1021/Cr0206420, 2003.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E.,
- 777 Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume
- II gas phase reactions of organic species, Atmospheric Chemistry and Physics, 6, 3625-4055, 2006.
- Baltensperger, U., Kalberer, M., Dommen, J., Paulsen, D., Alfarra, M., Coe, H., Fisseha, R., Gascho, A.,
- 780 Gysel, M., and Nyeki, S.: Secondary organic aerosols from anthropogenic and biogenic precursors,
- 781 Faraday Discuss, 130, 265-278, 2005.
- 782 Bannan, T. J., Bacak, A., Muller, J. B. A., Booth, A. M., Jones, B., Le Breton, M., Leather, K. E.,
- Ghalaieny, M., Xiao, P., Shallcross, D. E., and Percival, C. J.: Importance of direct anthropogenic
- emissions of formic acid measured by a chemical ionisation mass spectrometer (CIMS) during the Winter
- 785 ClearfLo Campaign in London, January 2012, Atmospheric Environment, 83, 301-310, Doi
- 786 10.1016/J.Atmosenv.2013.10.029, 2014.
- 787 Bates, K. H., Crounse, J. D., St. Clair, J. M., Bennett, N. B., Nguyen, T. B., Seinfeld, J. H., Stoltz, B. M.,
- and Wennberg, P. O.: Gas Phase Production and Loss of Isoprene Epoxydiols, The Journal of Physical
- 789 Chemistry A, 118, 1237-1246, 10.1021/jp4107958, 2014.
- Berndt, T., Böge, O., and Herrmann, H.: On the formation of benzene oxide/oxepin in the gas-phase
- reaction of OH radicals with benzene, Chemical Physics Letters, 314, 435-442, 10.1016/S0009-
- 792 2614(99)01041-6, 1999.
- Berndt, T., and Böge, O.: Gas-phase reaction of OH radicals with benzene: products and mechanism,
- 794 Phys Chem Chem Phys, 3, 4946-4956, 10.1039/b106667f, 2001.
- 795 Bertram, A. K., Ivanov, A. V., Hunter, M., Molina, L. T., and Molina, M. J.: The Reaction Probability of
- OH on Organic Surfaces of Tropospheric Interest, The Journal of Physical Chemistry A, 105, 9415-9421,
 10.1021/jp0114034, 2001.
- Bon, D. M., Ulbrich, I. M., de Gouw, J. A., Warneke, C., Kuster, W. C., Alexander, M. L., Baker, A.,
- 799 Beyersdorf, A. J., Blake, D., Fall, R., Jimenez, J. L., Herndon, S. C., Huey, L. G., Knighton, W. B.,
- 800 Ortega, J., Springston, S., and Vargas, O.: Measurements of volatile organic compounds at a suburban
- ground site (T1) in Mexico City during the MILAGRO 2006 campaign: measurement comparison,
- 802 emission ratios, and source attribution, Atmos. Chem. Phys., 11, 2399-2421, 10.5194/acp-11-2399-2011, 2011.
- 804 Borbon, A., Gilman, J. B., Kuster, W. C., Grand, N., Chevaillier, S., Colomb, A., Dolgorouky, C., Gros,
- 805 V., Lopez, M., Sarda-Esteve, R., Holloway, J., Stutz, J., Petetin, H., McKeen, S., Beekmann, M.,
- 806 Warneke, C., Parrish, D. D., and de Gouw, J. A.: Emission ratios of anthropogenic volatile organic
- 807 compounds in northern mid-latitude megacities: Observations versus emission inventories in Los Angeles
- and Paris, Journal of Geophysical Research: Atmospheres, 118, 1-17, 10.1002/jgrd.50059, 2013.
- 809 Bossolasco, A., Faragó, E. P., Schoemaecker, C., and Fittschen, C.: Rate constant of the reaction between
- 810 CH3O2 and OH radicals, Chemical Physics Letters, 593, 7-13, 10.1016/j.cplett.2013.12.052, 2014.
- 811 Butkovskaya, N. I., Pouvesle, N., Kukui, A., and Le Bras, G.: Mechanism of the OH-Initiated Oxidation
- of Glycolaldehyde over the Temperature Range 233–296 K, The Journal of Physical Chemistry A, 110,
- 813 13492-13499, 10.1021/jp064993k, 2006a.
- 814 Butkovskaya, N. I., Pouvesle, N., Kukui, A., Mu, Y., and Le Bras, G.: Mechanism of the OH-Initiated
- 815 Oxidation of Hydroxyacetone over the Temperature Range 236–298 K⁺, The Journal of Physical
- 816 Chemistry A, 110, 6833-6843, 10.1021/jp056345r, 2006b.
- 817 Cady-Pereira, K. E., Chaliyakunnel, S., Shephard, M. W., Millet, D. B., Luo, M., and Wells, K. C.:
- 818 HCOOH measurements from space: TES retrieval algorithm and observed global distribution, Atmos.
- 819 Meas. Tech., 7, 2297-2311, 10.5194/amt-7-2297-2014, 2014.

- 820 Carlton, A. G., Turpin, B. J., Lim, H. J., Altieri, K. E., and Seitzinger, S.: Link between isoprene and
- 821 secondary organic aerosol (SOA): Pyruvic acid oxidation yields low volatility organic acids in clouds, 822
- Geophysical Research Letters, 33, Doi 10.1029/2005gl025374, 2006.
- 823 Carlton, A. G., Turpin, B. J., Altieri, K. E., Seitzinger, S., Reff, A., Lim, H.-J., and Ervens, B.:
- Atmospheric oxalic acid and SOA production from glyoxal: Results of aqueous photooxidation 824
- 825 experiments, Atmospheric Environment, 41, 7588-7602, 10.1016/j.atmosenv.2007.05.035, 2007.
- 826 Chameides, W. L., and Davis, D. D.: Aqueous-phase source of formic acid in clouds, Nature, 304, 427-827 429, 1983.
- 828 Chebbi, A., and Carlier, P.: Carboxylic acids in the troposphere, occurrence, sources, and sinks: A review, 829 Atmospheric Environment, 30, 4233-4249, 1996.
- 830 Crisp, T. A., Brady, J. M., Cappa, C. D., Collier, S., Forestieri, S. D., Kleeman, M. J., Kuwayama, T.,
- 831 Lerner, B. M., Williams, E. J., Zhang, Q., and Bertram, T. H.: On the primary emission of formic acid
- 832 from light duty gasoline vehicles and ocean-going vessels, Atmospheric Environment, 98, 426-433,
- 833 10.1016/j.atmosenv.2014.08.070, 2014.
- 834 da Silva, G.: Carboxylic Acid Catalyzed Keto-Enol Tautomerizations in the Gas Phase, Angewandte
- 835 Chemie International Edition, 49, 7523-7525, 10.1002/anie.201003530, 2010.
- 836 Daele, V., and Poulet, G.: Kinetics and products of the reactions of CH3O2 with Cl and ClO, Journal De
- 837 Chimie Physique Et De Physico-Chimie Biologique, 93, 1081-1099, 1996.
- de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M., 838
- 839 Fehsenfeld, F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C., Marchewka, M.,
- 840 Bertman, S. B., and Bates, T. S.: Budget of organic carbon in a polluted atmosphere: Results from the
- 841 New England Air Quality Study in 2002, Journal of Geophysical Research-Atmospheres, 110, D16305, 842 10.1029/2004jd005623, 2005.
- Di Carlo, P., Brune, W. H., Martinez, M., Harder, H., Lesher, R., Ren, X., Thornberry, T., Carroll, M. A., 843
- Young, V., Shepson, P. B., Riemer, D., Apel, E., and Campbell, C.: Missing OH Reactivity in a Forest: 844
- 845 Evidence for Unknown Reactive Biogenic VOCs, Science, 304, 722-725, 10.1126/science.1094392, 2004.
- 846 Dibb, J. E., and Arsenault, M.: Shouldn't snowpacks be sources of monocarboxylic acids?, Atmospheric
- 847 Environment, 36, 2513-2522, 10.1016/S1352-2310(02)00131-0, 2002.
- Dubé, W. P., Brown, S. S., Osthoff, H. D., Nunley, M. R., Ciciora, S. J., Paris, M. W., McLaughlin, R. J., 848
- 849 and Ravishankara, A. R.: Aircraft instrument for simultaneous, in situ measurement of NO3 and N2O5
- 850 via pulsed cavity ring-down spectroscopy, Review of Scientific Instruments, 77, -,
- 851 doi:http://dx.doi.org/10.1063/1.2176058, 2006.
- Dubowski, Y., Vieceli, J., Tobias, D. J., Gomez, A., Lin, A., Nizkorodov, S. A., McIntire, T. M., and 852
- 853 Finlayson-Pitts, B. J.: Interaction of Gas-Phase Ozone at 296 K with Unsaturated Self-Assembled
- 854 Monolayers: A New Look at an Old System, The Journal of Physical Chemistry A, 108, 10473-10485,
- 855 10.1021/jp046604x, 2004.
- Edwards, P. M., Young, C. J., Aikin, K., deGouw, J., Dubé, W. P., Geiger, F., Gilman, J., Helmig, D., 856
- Holloway, J. S., Kercher, J., Lerner, B., Martin, R., McLaren, R., Parrish, D. D., Peischl, J., Roberts, J. M., 857
- 858 Ryerson, T. B., Thornton, J., Warneke, C., Williams, E. J., and Brown, S. S.: Ozone photochemistry in an
- 859 oil and natural gas extraction region during winter: simulations of a snow-free season in the Uintah Basin,
- Utah, Atmos. Chem. Phys., 13, 8955-8971, 10.5194/acp-13-8955-2013, 2013. 860
- Edwards, P. M., Brown, S. S., Roberts, J. M., Ahmadov, R., Banta, R. M., deGouw, J. A., Dube, W. P., 861
- 862 Field, R. A., Flynn, J. H., Gilman, J. B., Graus, M., Helmig, D., Koss, A., Langford, A. O., Lefer, B. L.,
- Lerner, B. M., Li, R., Li, S. M., McKeen, S. A., Murphy, S. M., Parrish, D. D., Senff, C. J., Soltis, J., 863
- Stutz, J., Sweeney, C., Thompson, C. R., Trainer, M. K., Tsai, C., Veres, P. R., Washenfelder, R. A., 864
- Warneke, C., Wild, R. J., Young, C. J., Yuan, B., and Zamora, R.: High winter ozone pollution from 865
- 866 carbonyl photolysis in an oil and gas basin, Nature, 514, 351-354, 10.1038/nature13767, 2014.
- Eliason, T. L., Gilman, J. B., and Vaida, V.: Oxidation of organic films relevant to atmospheric aerosols, 867
- Atmospheric Environment, 38, 1367-1378, 10.1016/j.atmosenv.2003.11.025, 2004. 868
- 869 Emmerson, K. M., and Evans, M. J.: Comparison of tropospheric gas-phase chemistry schemes for use
- 870 within global models, Atmos. Chem. Phys., 9, 1831-1845, 10.5194/acp-9-1831-2009, 2009.

- 871 Fittschen, C., Whalley, L. K., and Heard, D. E.: The reaction of CH3O2 radicals with OH radicals: a
- neglected sink for CH3O2 in the remote atmosphere, Environ Sci Technol, 48, 7700-7701,
- 873 10.1021/es502481q, 2014.
- Fried, A., Crawford, J., Olson, J., Walega, J., Potter, W., Wert, B., Jordan, C., Anderson, B., Shetter, R.,
- Efer, B., Blake, D., Blake, N., Meinardi, S., Heikes, B., O'Sullivan, D., Snow, J., Fuelberg, H., Kiley, C.
- M., Sandholm, S., Tan, D., Sachse, G., Singh, H., Faloona, I., Harward, C. N., and Carmichael, G. R.:
- Airborne tunable diode laser measurements of formaldehyde during TRACE-P: Distributions and box
- model comparisons, Journal of Geophysical Research-Atmospheres, 108, NO. D20, 8798,
- 879 10.1029/2003jd003451, 2003.
- 880 Fried, A., Walega, J. G., Olson, J. R., Crawford, J. H., Chen, G., Weibring, P., Richter, D., Roller, C.,
- Tittel, F. K., Heikes, B. G., Snow, J. A., Shen, H., O'Sullivan, D. W., Porter, M., Fuelberg, H., Halland, J.,
- and Millet, D. B.: Formaldehyde over North America and the North Atlantic during the summer 2004
- 883 INTEX campaign: Methods, observed distributions, and measurement-model comparisons, J. Geophys.
- 884 Res., 113, D10302, 10.1029/2007jd009185, 2008.
- Gilman, J. B., Lerner, B. M., Kuster, W. C., and de Gouw, J. A.: Source signature of volatile organic
- compounds from oil and natural gas operations in northeastern Colorado, Environ Sci Technol, 47, 1297-
- 887 1305, 10.1021/es304119a, 2013.
- 888 Grosjean, D., Williams, E. L., and Grosjean, E.: Atmospheric Chemistry of Isoprene and of Its Carbonyl
- 889 Products, Environmental Science & Technology, 27, 830-840, 1993.
- 890 Hatakeyama, S., Washida, N., and Akimoto, H.: Rate constants and mechanisms for the reaction of
- 891 hydroxyl (OD) radicals with acetylene, propyne, and 2-butyne in air at 297 ± 2 K, The Journal of Physical
- 892 Chemistry, 90, 173-178, 10.1021/j100273a039, 1986.
- 893 Helmig, D., Thompson, C. R., Evans, J., Boylan, P., Hueber, J., and Park, J. H.: Highly elevated
- atmospheric levels of volatile organic compounds in the Uintah Basin, Utah, Environ Sci Technol, 48,
- 895 4707-4715, 10.1021/es405046r, 2014.
- 896 Herckes, P., Valsaraj, K. T., and Collett Jr, J. L.: A review of observations of organic matter in fogs and
- 897 clouds: Origin, processing and fate, Atmospheric Research, 132–133, 434-449,
- 898 10.1016/j.atmosres.2013.06.005, 2013.
- Hersey, S. P., Craven, J. S., Metcalf, A. R., Lin, J., Lathem, T., Suski, K. J., Cahill, J. F., Duong, H. T.,
- 900 Sorooshian, A., Jonsson, H. H., Shiraiwa, M., Zuend, A., Nenes, A., Prather, K. A., Flagan, R. C., and
- 901 Seinfeld, J. H.: Composition and hygroscopicity of the Los Angeles Aerosol: CalNex, Journal of
- 902 Geophysical Research: Atmospheres, 118, 3016-3036, 10.1002/jgrd.50307, 2013.
- Hu, L., Millet, D. B., Kim, S. Y., Wells, K. C., Griffis, T. J., Fischer, E. V., Helmig, D., Hueber, J., and
- 904 Curtis, A. J.: North American acetone sources determined from tall tower measurements and inverse
- 905 modeling, Atmos. Chem. Phys., 13, 3379-3392, 10.5194/acp-13-3379-2013, 2013.
- 906 Ito, A., Sillman, S., and Penner, J. E.: Effects of additional nonmethane volatile organic compounds,
- 907 organic nitrates, and direct emissions of oxygenated organic species on global tropospheric chemistry,
- Journal of Geophysical Research-Atmospheres, 112, D06309, Doi 10.1029/2005jd006556, 2007.
- Jacobi, H.-W., Bales, R. C., Honrath, R. E., Peterson, M. C., Dibb, J. E., Swanson, A. L., and Albert, M.
- 910 R.: Reactive trace gases measured in the interstitial air of surface snow at Summit, Greenland,
- 911 Atmospheric Environment, 38, 1687-1697, 10.1016/j.atmosenv.2004.01.004, 2004.
- 912 Jardine, K., Yañez Serrano, A., Arneth, A., Abrell, L., Jardine, A., Artaxo, P., Alves, E., Kesselmeier, J.,
- 913 Taylor, T., Saleska, S., and Huxman, T.: Ecosystem-scale compensation points of formic and acetic acid
- 914 in the central Amazon, Biogeosciences, 8, 3709-3720, 10.5194/bg-8-3709-2011, 2011.
- 915 Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel of the
- 916 CH3C(O)O2 + HO2 reaction in the gas phase, Phys Chem Chem Phys, 9, 3149-3162, 10.1039/b702757e, 917 2007.
- Jenkin, M. E., Wyche, K. P., Evans, C. J., Carr, T., Monks, P. S., Alfarra, M. R., Barley, M. H.,
- 919 McFiggans, G. B., Young, J. C., and Rickard, A. R.: Development and chamber evaluation of the MCM
- 920 v3.2 degradation scheme for β -caryophyllene, Atmos. Chem. Phys., 12, 5275-5308, 10.5194/acp-12-
- 921 5275-2012, 2012.

- 922 Karton, A.: Inorganic acid-catalyzed tautomerization of vinyl alcohol to acetaldehyde, Chemical Physics
- 923 Letters, 592, 330-333, 10.1016/j.cplett.2013.12.062, 2014.
- Kawamura, K., Steinberg, S., and Kaplan, I. R.: Homologous series of C1-C10 monocarboxylic acids and
- 925 C1-C6 carbonyls in Los Angeles air and motor vehicle exhausts, Atmospheric Environment, 34, 4175-
- 926 4191, 2000.
- 927 Keene, W. C., Mosher, B. W., Jacob, D. J., Munger, J. W., Talbot, R. W., Artz, R. S., Maben, J. R.,
- Daube, B. C., and Galloway, J. N.: Carboxylic acids in clouds at a high-elevation forested site in central
- Virginia, Journal of Geophysical Research: Atmospheres, 100, 9345-9357, 10.1029/94JD01247, 1995.
- Khare, P., Kumar, N., Kumari, K. M., and Srivastava, S. S.: Atmospheric formic and acetic acids: An
- 931 overview, Rev Geophys, 37, 227-248, 1999.
- 932 Kwan, A. J., Crounse, J. D., Clarke, A. D., Shinozuka, Y., Anderson, B. E., Crawford, J. H., Avery, M. A.,
- 933 McNaughton, C. S., Brune, W. H., Singh, H. B., and Wennberg, P. O.: On the flux of oxygenated volatile
- 934 organic compounds from organic aerosol oxidation, Geophysical Research Letters, 33,
- 935 doi:10.1029/2006GL026144, Doi 10.1029/2006gl026144, 2006.
- 936 Le Breton, M., McGillen, M. R., Muller, J. B. A., Bacak, A., Shallcross, D. E., Xiao, P., Huey, L. G.,
- 937 Tanner, D., Coe, H., and Percival, C. J.: Airborne observations of formic acid using a chemical ionization
- mass spectrometer, Atmos. Meas. Tech., 5, 3029-3039, 10.5194/amt-5-3029-2012, 2012.
- Le Breton, M., Bacak, A., Muller, J. B. A., Xiao, P., Shallcross, B. M. A., Batt, R., Cooke, M. C.,
- 940 Shallcross, D. E., Bauguitte, S. J. B., and Percival, C. J.: Simultaneous airborne nitric acid and formic
- acid measurements using a chemical ionization mass spectrometer around the UK: Analysis of primary
- and secondary production pathways, Atmospheric Environment, 83, 166-175,
- 943 10.1016/j.atmosenv.2013.10.008, 2014.
- Leather, K. E., McGillen, M. R., Cooke, M. C., Utembe, S. R., Archibald, A. T., Jenkin, M. E., Derwent,
- R. G., Shallcross, D. E., and Percival, C. J.: Acid-yield measurements of the gas-phase ozonolysis of
- ethene as a function of humidity using Chemical Ionisation Mass Spectrometry (CIMS), Atmos. Chem.
- 947 Phys., 12, 469-479, 10.5194/acp-12-469-2012, 2012.
- Lee, A., Goldstein, A. H., Keywood, M. D., Gao, S., Varutbangkul, V., Bahreini, R., Ng, N. L., Flagan, R.
- 949 C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the ozonolysis of ten
- different terpenes, Journal of Geophysical Research: Atmospheres, 111, D07302, 10.1029/2005jd006437,
 2006.
- 952 Lewis, A. C., Carslaw, N., Marriott, P. J., Kinghorn, R. M., Morrison, P., Lee, A. L., Bartle, K. D., and
- Pilling, M. J.: A larger pool of ozone-forming carbon compounds in urban atmospheres, Nature, 405, 778781, 2000.
- Li, X., Rohrer, F., Brauers, T., Hofzumahaus, A., Lu, K., Shao, M., Zhang, Y. H., and Wahner, A.:
- 956 Modeling of HCHO and CHOCHO at a semi-rural site in southern China during the PRIDE-PRD2006
- 957 campaign, Atmospheric Chemistry and Physics, 14, 12291-12305, Doi 10.5194/Acp-14-12291-2014,
- 958 2014.
- Lide, D. R.: CRC Handbook of Chemistry and Physics, in, 2005.
- 960 Liu, J., Zhang, X., Parker, E. T., Veres, P. R., Roberts, J. M., de Gouw, J. A., Hayes, P. L., Jimenez, J. L.,
- 961 Murphy, J. G., Ellis, R. A., Huey, L. G., and Weber, R. J.: On the gas-particle partitioning of soluble
- organic aerosol in two urban atmospheres with contrasting emissions: 2. Gas and particle phase formic
 acid, J. Geophys. Res., 117, D00V21, 10.1029/2012jd017912, 2012.
- 964 Maricq, M. M., Szente, J. J., Kaiser, E. W., and Shi, J.: Reaction of Chlorine Atoms with Methylperoxy
- and Ethylperoxy Radicals, The Journal of Physical Chemistry, 98, 2083-2089, 10.1021/j100059a017,
 1994.
- 967 McCabe, J., and Abbatt, J. P. D.: Heterogeneous Loss of Gas-Phase Ozone on n-Hexane Soot Surfaces:
- 968 Similar Kinetics to Loss on Other Chemically Unsaturated Solid Surfaces[†], The Journal of Physical
- 969 Chemistry C, 113, 2120-2127, 10.1021/jp806771q, 2008.
- 970 Molina, M. J., Ivanov, A. V., Trakhtenberg, S., and Molina, L. T.: Atmospheric evolution of organic
- 971 aerosol, Geophysical Research Letters, 31, L22104, 10.1029/2004gl020910, 2004.

- 972 Muller, J. F., and Peeters, J.: Interactive comment on "HCOOH measurements from space: TES retrieval
- algorithm and observed global distribution" by K. E. Cady-Pereira et al., Atmos. Meas. Tech. Discuss., 7,
 C312-C314, 2014.
- 975 Neeb, P., Sauer, F., Horie, O., and Moortgat, G. K.: Formation of hydroxymethyl hydroperoxide and
- 976 formic acid in alkene ozonolysis in the presence of water vapour, Atmospheric Environment, 31, 1417-
- 977 1423, 10.1016/S1352-2310(96)00322-6, 1997.
- 978 Norton, R. B.: Measurements of formate and acetate in precipitation at Niwot Ridge and Boulder,
- 979 Colorado, Geophysical Research Letters, 12, 769-772, 10.1029/GL012i011p00769, 1985.
- 980 Orlando, J., Tyndall, G., and Taraborrelli, D.: Atmospheric Oxidation of Two Isoprene By-Products,
- Hydroxyacetone and Glycolaldehyde, AGU Fall Meeting 2012, San Francisco, 3-7 December, 2012.
- 982 Orlando, J. J., Nozière, B., Tyndall, G. S., Orzechowska, G. E., Paulson, S. E., and Rudich, Y.: Product
- studies of the OH- and ozone-initiated oxidation of some monoterpenes, Journal of Geophysical Research:
 Atmospheres, 105, 11561-11572, 10.1029/2000jd900005, 2000.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kroll, J. H., Seinfeld, J. H., and Wennberg, P. O.: Isoprene
- 986 photooxidation: new insights into the production of acids and organic nitrates, Atmospheric Chemistry
- 987 and Physics, 9, 1479-1501, 2009a.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kurten, A., St Clair, J. M., Seinfeld, J. H., and Wennberg, P.
- O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, Science, 325, 730-733,
 DOI 10.1126/science.1172910, 2009b.
- 991 Paulot, F., Wunch, D., Crounse, J. D., Toon, G. C., Millet, D. B., DeCarlo, P. F., Vigouroux, C.,
- 992 Deutscher, N. M., González Abad, G., Notholt, J., Warneke, T., Hannigan, J. W., Warneke, C., de Gouw,
- J. A., Dunlea, E. J., De Mazière, M., Griffith, D. W. T., Bernath, P., Jimenez, J. L., and Wennberg, P. O.:
- 994 Importance of secondary sources in the atmospheric budgets of formic and acetic acids, Atmos. Chem.
- 995 Phys., 11, 1989-2013, 10.5194/acp-11-1989-2011, 2011.
- 996 Perri, M. J., Seitzinger, S., and Turpin, B. J.: Secondary organic aerosol production from aqueous
- 997 photooxidation of glycolaldehyde: Laboratory experiments, Atmospheric Environment, 43, 1487-1497,
- 998 10.1016/j.atmosenv.2008.11.037, 2009.
- Russell, L. M., Bahadur, R., and Ziemann, P. J.: Identifying organic aerosol sources by comparing
- 1000 functional group composition in chamber and atmospheric particles, Proc Natl Acad Sci U S A, 108,
- 1001 3516-3521, 10.1073/pnas.1006461108, 2011.
- 1002 Ryerson, T. B., Andrews, A. E., Angevine, W. M., Bates, T. S., Brock, C. A., Cairns, B., Cohen, R. C.,
- 1003 Cooper, O. R., de Gouw, J. A., Fehsenfeld, F. C., Ferrare, R. A., Fischer, M. L., Flagan, R. C., Goldstein,
- 1004 A. H., Hair, J. W., Hardesty, R. M., Hostetler, C. A., Jimenez, J. L., Langford, A. O., McCauley, E.,
- 1005 McKeen, S. A., Molina, L. T., Nenes, A., Oltmans, S. J., Parrish, D. D., Pederson, J. R., Pierce, R. B.,
- 1006 Prather, K., Quinn, P. K., Seinfeld, J. H., Senff, C. J., Sorooshian, A., Stutz, J., Surratt, J. D., Trainer, M.,
- 1007 Volkamer, R., Williams, E. J., and Wofsy, S. C.: The 2010 California Research at the Nexus of Air
- 1008 Quality and Climate Change (CalNex) field study, Journal of Geophysical Research: Atmospheres, 118, 5830-5866, 10 1002/jard 50331, 2013
- 1009 5830-5866, 10.1002/jgrd.50331, 2013.
- 1010 Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the
- 1011 Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile
- 1012 organic compounds, Atmos. Chem. Phys., 3, 161-180, 10.5194/acp-3-161-2003, 2003.
- 1013 Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F. M., and Roberts, J. M.: A thermal dissociation
- 1014 chemical ionization mass spectrometry (TD-CIMS) technique for the simultaneous measurement of
- 1015 peroxyacyl nitrates and dinitrogen pentoxide, J. Geophys. Res., 109, D19315, 10.1029/2004jd004670,
- 1016 2004.
- 1017 So, S., da Silva, G., and Wille, U.: The Atmospheric Chemistry of Enols: A Theoretical Study of the
- 1018 Vinyl Alcohol + OH + O2 Reaction Mechanism, Environ Sci Technol, 10.1021/es500319q, 2014.
- 1019 Sommariva, R., de Gouw, J. A., Trainer, M., Atlas, E., Goldan, P. D., Kuster, W. C., Warneke, C., and
- 1020 Fehsenfeld, F. C.: Emissions and photochemistry of oxygenated VOCs in urban plumes in the
- 1021 Northeastern United States, Atmos. Chem. Phys., 11, 7081-7096, 10.5194/acp-11-7081-2011, 2011.

- 1022 Stavrakou, T., Muller, J. F., Peeters, J., Razavi, A., Clarisse, L., Clerbaux, C., Coheur, P. F., Hurtmans, D.,
- 1023 De Maziere, M., Vigouroux, C., Deutscher, N. M., Griffith, D. W. T., Jones, N., and Paton-Walsh, C.:
- Satellite evidence for a large source of formic acid from boreal and tropical forests, Nature Geosci, 5, 26-30, 2012.
- 1026 Talbot, R. W., Beecher, K. M., Harriss, R. C., and Cofer, W. R., III: Atmospheric Geochemistry of
- 1027 Formic and Acetic Acids at a Mid-Latitude Temperate Site, J. Geophys. Res., 93, 1638-1652,
- 1028 10.1029/JD093iD02p01638, 1988.
- 1029 Tan, Y., Carlton, A. G., Seitzinger, S. P., and Turpin, B. J.: SOA from methylglyoxal in clouds and wet
- aerosols: Measurement and prediction of key products, Atmospheric Environment, 44, 5218-5226,
- 1031 10.1016/j.atmosenv.2010.08.045, 2010.
- 1032 Thomas, E. R., Frost, G. J., and Rudich, Y.: Reactive uptake of ozone by proxies for organic aerosols:
- Surface-bound and gas-phase products, Journal of Geophysical Research: Atmospheres, 106, 3045-3056,
 1034 10.1029/2000jd900595, 2001.
- 1035 Veres, P., Roberts, J. M., Warneke, C., Welsh-Bon, D., Zahniser, M., Herndon, S., Fall, R., and de Gouw,
- 1036 J.: Development of negative-ion proton-transfer chemical-ionization mass spectrometry (NI-PT-CIMS)
- 1037 for the measurement of gas-phase organic acids in the atmosphere, International Journal of Mass
- 1038 Spectrometry, 274, 48-55, DOI 10.1016/j.ijms.2008.04.032, 2008.
- 1039 Veres, P., Gilman, J. B., Roberts, J. M., Kuster, W. C., Warneke, C., Burling, I. R., and de Gouw, J.:
- 1040 Development and validation of a portable gas phase standard generation and calibration system for
- 1041 volatile organic compounds, Atmos. Meas. Tech., 3, 683-691, 10.5194/amt-3-683-2010, 2010.
- 1042 Veres, P. R., Roberts, J. M., Cochran, A. K., Gilman, J. B., Kuster, W. C., Holloway, J. S., Graus, M.,
- Flynn, J., Lefer, B., Warneke, C., and de Gouw, J.: Evidence of rapid production of organic acids in an urban air mass, Geophys. Res. Lett., 38, L17807, 10.1029/2011gl048420, 2011.
- 1045 Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C.-U., Aas, W., Baker, A., Bowersox, V. C., Dentener, F.,
- 1046 Galy-Lacaux, C., Hou, A., Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher, T.,
- 1047 Mahowald, N. M., Nickovic, S., Rao, P. S. P., and Reid, N. W.: A global assessment of precipitation
- 1048 chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and
- 1049 phosphorus, Atmospheric Environment, 93, 3-100, 10.1016/j.atmosenv.2013.10.060, 2014.
- 1050 Vlasenko, A., George, I. J., and Abbatt, J. P. D.: Formation of Volatile Organic Compounds in the
- Heterogeneous Oxidation of Condensed-Phase Organic Films by Gas-Phase OH, The Journal of Physical
 Chemistry A, 112, 1552-1560, 10.1021/jp0772979, 2008.
- 1053 Warneke, C., de Gouw, J. A., Lovejoy, E. R., Murphy, P. C., Kuster, W. C., and Fall, R.: Development of
- 1054 proton-transfer ion trap-mass spectrometry: On-line detection and identification of volatile organic
- 1055 compounds in air, Journal of the American Society for Mass Spectrometry, 16, 1316-1324,
- 1056 10.1016/j.jasms.2005.03.025, 2005.
- 1057 Warneke, C., Veres, P., Holloway, J. S., Stutz, J., Tsai, C., Alvarez, S., Rappenglueck, B., Fehsenfeld, F.
- 1058 C., Graus, M., Gilman, J. B., and de Gouw, J. A.: Airborne formaldehyde measurements using PTR-MS:
- calibration, humidity dependence, inter-comparison and initial results, Atmos. Meas. Tech., 4, 2345-2358,
 1060 10.5194/amt-4-2345-2011, 2011.
- 1061 Warneke, C., Geiger, F., Edwards, P. M., Dube, W., Pétron, G., Kofler, J., Zahn, A., Brown, S. S., Graus,
- 1062 M., Gilman, J., Lerner, B., Peischl, J., Ryerson, T. B., de Gouw, J. A., and Roberts, J. M.: Volatile
- 1063 organic compound emissions from the oil and natural gas industry in the Uinta Basin, Utah: point sources
- 1064 compared to ambient air composition, Atmos. Chem. Phys., 14, 10977-10988, 10.5194/acpd-14-11895-
- 1065 2014, 2014.
- 1066 Welz, O., Eskola, A. J., Sheps, L., Rotavera, B., Savee, J. D., Scheer, A. M., Osborn, D. L., Lowe, D.,
- 1067 Murray Booth, A., Xiao, P., Anwar H. Khan, M., Percival, C. J., Shallcross, D. E., and Taatjes, C. A.:
- 1068 Rate Coefficients of C1 and C2 Criegee Intermediate Reactions with Formic and Acetic Acid Near the
- 1069 Collision Limit: Direct Kinetics Measurements and Atmospheric Implications, Angewandte Chemie
- 1070 International Edition, 126, 4635-4638, 10.1002/ange.201400964, 2014.

- 1071 Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale
- 1072 numerical models, Atmospheric Environment (1967), 23, 1293-1304, <u>http://dx.doi.org/10.1016/0004-</u>
 1073 6981(89)90153-4, 1989.
- 1074 Wild, R. J., Edwards, P. M., Dube, W. P., Baumann, K., Edgerton, E. S., Quinn, P. K., Roberts, J. M.,
- 1075 Rollins, A. W., Veres, P. R., Warneke, C., Williams, E. J., Yuan, B., and Brown, S. S.: A Measurement of
- 1076 Total Reactive Nitrogen, NOy, together with NO2, NO, and O3 via Cavity Ring-down Spectroscopy,
- 1077 Environ Sci Technol, 48, 9609-9615, 10.1021/es501896w, 2014.
- 1078 Wyche, K. P., Monks, P. S., Ellis, A. M., Cordell, R. L., Parker, A. E., Whyte, C., Metzger, A., Dommen,
- 1079 J., Duplissy, J., Prevot, A. S. H., Baltensperger, U., Rickard, A. R., and Wulfert, F.: Gas phase precursors
- 1080 to anthropogenic secondary organic aerosol: detailed observations of 1,3,5-trimethylbenzene
- 1081 photooxidation, Atmos. Chem. Phys., 9, 635-665, 10.5194/acp-9-635-2009, 2009.
- 1082 Yatavelli, R. L. N., Stark, H., Thompson, S. L., Kimmel, J. R., Cubison, M. J., Day, D. A., Campuzano-
- 1083 Jost, P., Palm, B. B., Hodzic, A., Thornton, J. A., Jayne, J. T., Worsnop, D. R., and Jimenez, J. L.:
- 1084 Semicontinuous measurements of gas–particle partitioning of organic acids in a ponderosa pine forest
- 1085 using a MOVI-HRToF-CIMS, Atmos. Chem. Phys., 14, 1527-1546, 10.5194/acp-14-1527-2014, 2014.
- 1086 Zhang, R., Suh, I., Zhao, J., Zhang, D., Fortner, E. C., Tie, X., Molina, L. T., and Molina, M. J.:
- 1087 Atmospheric New Particle Formation Enhanced by Organic Acids, Science, 304, 1487-1490,
- 1088 10.1126/science.1095139, 2004.
- 1089 Zhang, X., Liu, J., Parker, E. T., Hayes, P. L., Jimenez, J. L., de Gouw, J. A., Flynn, J. H., Grossberg, N.,
- 1090 Lefer, B. L., and Weber, R. J.: On the gas-particle partitioning of soluble organic aerosol in two urban
- atmospheres with contrasting emissions: 1. Bulk water-soluble organic carbon, J. Geophys. Res., 117,
 D00V16, 10.1029/2012jd017908, 2012.
- 1093 Zhang, X., Schwantes, R. H., Coggon, M. M., Loza, C. L., Schilling, K. A., Flagan, R. C., and Seinfeld, J.
- H.: Role of ozone in SOA formation from alkane photooxidation, Atmos. Chem. Phys., 14, 1733-1753,
- 1095 10.5194/acp-14-1733-2014, 2014.
- 1096

1098 Tables

Pairs	Emission ratios, ppb/ppm	Location	References
HCOOH/CO	0.21	Virginia, US	(Talbot et al., 1988)
HCOOH/CO	1.22	London, UK	(Bannan et al., 2014)
HCOOH/NO _x	8.35	London, UK	(Bannan et al., 2014)
HCOOH/C ₂ H ₂	0	Northeastern US	(de Gouw et al., 2005)
HCOOH/CO	42.2±20.0×00 ⁻³	Laboratory study in California (hot running)	(Crisp et al., 2014)
HCOOH/CO	10.2±6.5×10 ⁻³	Laboratory study in California (cold start)	(Crisp et al., 2014)

1099 Table 1. Emission ratios (ERs) of formic acid to anthropogenic tracers from combustion

1009 1 able 1. Emission ratios (EKS) of re 1100 emissions reported in the literature

1101

Species	Literature values	MCM v3.2	Modified MCM v3.2		
	Reaction with oz	one			
Ethene	0.41^{1}	0.23	0.34		
Propene	0.14^{2}	0.074	0.14		
Isobutene	0.13^{2}	0.056	0.13		
Isoprene	0.30^{2}	0.15	0.31		
MVK	0.32^{3}	0.074	0.31		
MACR	0.22^{3}	0.20			
α-pinene	0.075^4	0			
<u>β-pinene</u>	0.04^{4}	0.09			
Reaction with OH					
Isoprene	0.10^{5}	0	0.08		
Glycolaldehyde	0.18^{6}	0	0.18		
Hydroxyacetone	0.08^{7}	0	0.08		
β-pinene	0.02^{8}	0			
Acetylene	0.40^{9}	0.36			

Table 2. Yields of formic acid from reactions of alkenes and other unsaturated compounds withozone and OH radical

1105 '--' indicates there is no modification to chemical mechanism of the species.

1106 References in the table: 1. (Leather et al., 2012); 2. (Neeb et al., 1997); 3. (Grosjean et al., 1993);

1107 4. (Lee et al., 2006); 5. (Paulot et al., 2009a); 6. (Butkovskaya et al., 2006a); 7. (Butkovskaya et

1108 al., 2006b); 8. (Orlando et al., 2000); 9. (Hatakeyama et al., 1986).

1109

1110

1113	Table 3. Production rates of formic acid in the modified MCM v3.2 and the gaps between the
------	--

box model simulations and measurements, and the reaction rates of aerosol-related reactions.

Campaign	UBWOS 2013	CalNex
HCOOH production rate in modified MCM v3.2 (ppb/day)	3.2	8.2
Additional HCOOH production rate needed (ppb/day)	13.6	10.0
Additional C ₂ H ₄ (ppb)	30.3	19.1
Additional C_5H_8 (ppb)		1.3
Yield of alkanes needed	6.9%	21.3%
DOC loss rate due to aqueous reactions (ppb C/day)	0.11±0.05	0.058 ± 0.044
Reaction rate of OH with aerosol (ppb/day)	0.06	0.33
Reactions rate of O ₃ with aerosol (ppb/day)	12±2	13±4

1115 "--" indicates no calculation.

1116 Figures



1117

Fig. 1. (A) Box-whisker plots of formic acid concentrations in UBWOS 2012, UBWOS 2013
and CalNex. The boxes denote the central 50% of the data (25-75 percentile), and the bars within
the box indicate the median value. The ends of the whiskers show the maximum and minimum
of the data. The cross markers show arithmetic mean concentrations of formic acid. (B) Diurnal
variations (arithmetic mean) of formic acid in UBWOS 2012, UBWOS 2013 and CalNex.



Fig. 2. Buildup patterns of formic acid, acetaldehyde, benzene and ozone in Jan. 29 -Feb 9 in
 UBWOS 2013. Daily averages of various species are shown.



Fig. 3. Ratios of average concentrations of various VOCs in UBWOS 2012 and UBWOS 2013
relative to CalNex. The dash line indicates a ratio of unity.



1133 Fig. 4. Comparison of measured and modeled diurnal profiles of formic acid for UBWOS 2013

(left) and CalNex (right). A zoom of the figure is shown in Fig. S3. Note that the scales of y-axis
are different.



Fig. 5. Comparison of measured and modeled diurnal profiles of formic acid and acetone for UBWOS 2013 (A and B) and CalNex (C and D) using different lifetimes due to physical losses.



Fig. 6. Scatter plots of formic acid versus acetone in UBWOS 2013 (A) and CalNex (B). The 1143 measured values are 30 min diurnal-averaged data. The measured values in UBWOS 2012 are 1144 1145 also shown in (A). The last-day diurnal results from the diurnal steady state (DSS) box model simulations are shown. Model results from the emission-based model and the DSS simulations 1146 using bi-modal physical loss rates in UBWOS 2013 are also shown. Dashed and dash-dotted 1147 lines show fitted results from measured concentrations and estimated secondary concentrations, 1148 1149 respectively. The fits in UBWOS 2013 exclude data in 7:30 am - 12:00 pm (empty symbols, see 1150 text for details). Solid lines indicate the fitted results from model simulations with modified MCM in the two campaigns. Numbers in the graphs indicate enhancement ratios of formic acid 1151 to acetone from various datasets. 1152

- 1154
- 1155





Fig. 7. Time series of formic acid and other VOCs (benzene, acetone, formaldehyde,
acetaldehyde and acetic acid) during the fog event on the morning of Feb. 7, 2013 in UBWOS
2013. Measured ambient temperature and relative humidity (RH) at different heights (2 m, 9 m
and 18 m) and downwelling and upwelling solar radiation are also shown.



1163 Fig. 8. (A) Time series of concentration gradients of formic acid and acetone during UBWOS 1164 2013. Time series of oxalate measured in melted snow water and the product of oxalate in the 1165 snow and nitric acid (HNO₃) in ambient air are also shown. The vertical black bars indicate periods with snow fall. (B) Scatter plot of concentration gradient of formic acid versus oxalate in 1166 the snow. (C) Scatter plot of concentration gradient of formic acid versus nitric acid 1167 concentration in ambient air. (D) Scatter plot of concentration gradients of formic acid versus the 1168 1169 products of oxalate in the snow and nitric acid in ambient air. The blue line is the linear regression to the data points. 1170

- 1171
- 1172



1174 Fig. 9. Contributions from various formation pathways to secondary production of formic acid in

1175	UBWOS 2013 (upper) and CalNex (bottom). Wedges with black outlines indicate the
1176	contributions from gas phase reactions.