




Supplement of

A new model of the global biogeochemical cycle of carbonyl sulfide – Part 2: Use of ocs to constrain gross primary productivity of current vegetation models

T. Launois et al.

Correspondence to: T. Launois (thomas.launois@lsce.ipsl.fr)

1



2

3 Figure A1: Upper row: annual mean indirect oceanic emissions of OCS from DMS (resulting from the
 4 oxidation of DMS in the atmosphere), for January (left column) and July (right column). Second and third
 5 rows: monthly means of oceanic indirect emissions of OCS from DMS provided by large latitudinal bands
 6 and at the global scale.

7

8


9

10


11

12

13




1
2 Figure A2: Sensitivity of monthly exchanges of OCS between the atmosphere and the soils (over the period
3 2001-2002) to the choice of (1) the deposition velocities of H₂ to soils (Morfopoulos et al., 2012 vs.
4 Bousquet et al., 2011), and (2) the ratio between OCS and H₂ deposition velocities (0.5:1 (H. Chen pers.
5 com.) vs. 1:1 (Belviso et al., 2013)). Results are provided for the global scale and by large latitudinal bands.
6 Note that the OCS emissions by anoxic soils were kept unchanged between the simulations, following the
7 mean flux values proposed by Whelan et al. (2013). The Kettle et al. (2002) soil fluxes are shown in black
8 soil fluxes are shown in black
9
10



1
2 Figure A3: Monthly uptakes of OCS by vegetation (for 2001) deduced from CO₂ gross fluxes (GPP)
3 calculated by three different vegetation models (ORC, LPJ and CLM4CN). Results are provided for the
4 global scale and by large latitudinal bands. The Kettle et al. (2002) vegetation fluxes are shown in black
5 solid line.

6
7
8
9
10
11
12
13
14



1
2 Figure A4: Annual variations of monthly mean GPPs (solid lines) and monthly mean respiration fluxes
3 (dotted lines) simulated by the ORC, CLM4CN and LPJ vegetation models. Data are for the following three
4 latitudinal bands: 50°N-90°N (boreal vegetation, upper panel), 30°N-50°N (vegetation of temperate regions
5 of the northern hemisphere, central panel), and 30°S-30°N (tropical vegetation, lower panel).
6
7