

Large-eddy simulation of contrail evolution in the vortex phase and its interaction with atmospheric turbulence

J. Picot¹, R. Paoli¹, O. Thouron¹, and D. Cariolle^{1,2}

¹CNRS/CERFACS, URA 1875, Sciences de l'Univers au CERFACS, Toulouse, France

²Météo France, Toulouse, France

Correspondence to: R. Paoli (paoli@cerfacs.fr)

Abstract. In this work, the evolution of contrails in the vortex and dissipation regimes is studied by means of fully three-dimensional large-eddy simulation (LES) coupled to a Lagrangian particle tracking method to treat the ice phase. This is the first paper where fine-scale atmospheric turbulence is generated and sustained by means of a stochastic forcing that mimics the properties of stably stratified turbulent flows as those occurring in the upper troposphere lower stratosphere. The initial flow-field is composed by the turbulent background flow and a wake flow obtained from separate LES of the jet regime. Atmospheric turbulence is the main driver of the wake instability and the structure of the resulting wake is sensitive to the intensity of the perturbations, primarily in the vertical direction. A stronger turbulence accelerates the onset of the instability, which results in shorter contrail decent and more effective mixing in the interior of the plume. However, the self-induced turbulence that is produced in the wake after the vortex break-up dominates over background turbulence at the end of the vortex regime and dominates the mixing with ambient air. This results in mean microphysical characteristics such as ice mass and optical depth that are be slightly affected by the intensity of atmospheric turbulence. On the other hand, the background humidity and temperature have a first order effect on the survival of ice crystals and particle size distribution, which is in line with recent studies in the literature.

1 Introduction

Aircraft-induced cloudiness in the form of condensation trails (contrails) and contrail cirrus is among the most uncertain contributors of aviation to the Earth radiative forcing (Lee et al., 2009). One of the main reasons of this uncertainty is the large disparity of scales that exists between the

near-field of aircraft engines where emissions are produced, and the grid-boxes of global models that are used to evaluate their global atmospheric impact (Burkhardt et al., 2010). In order to bridge this gap and develop more refined parameterizations of unresolved plume processes into global models, one has to gain better knowledge of the dynamical and the microphysical properties of contrails and their precursors. For that purpose, high-fidelity numerical simulations such as large-eddy simulations (LES) are now becoming a reliable instrument of research in the atmospheric science community as they allow for accurate description of the physicochemical processes that occur in aircraft wakes (Unterstrasser and Gierens, 2010a; Paugam et al., 2010; Naiman et al., 2011).

According to the classification given by (Gerz et al., 1998), the wake of an aircraft can be split into four successive regimes: during the first few seconds after emission (jet regime), ice forms by condensation of water vapor onto particle nucleation sites like soot emitted by the engines (Schumann, 1996; Kärcher et al., 1996), while the vorticity shed by the aircraft wings rolls up into a pair of counter-rotating vortices where the ice and vapor exhausts are trapped (primary wake). In the following minutes (vortex regime), the primary wake descends because of the downward motion induced by the vortices on each other; then it interacts with the surrounding atmosphere via a baroclinic instability, which eventually leads to the formation of an upward-moving secondary wake (Spalart, 1996) where part of the exhausts and ice particles are detrained (Gerz and Ehret, 1997). The vortices also interact with each other via a kinematic process of mutual instability (Crow, 1970; Widnall et al., 1974) until they break up and release the exhaust in the atmosphere (dissipation regime). For high ambient humidity, ice particles size can increase dramatically by condensation of ambient water vapor (Lewellen and Lewellen, 2001; Paugam et al., 2010; Naiman et al., 2011). During the final diffusion

regime, the contrail evolution is controlled by atmospheric motion via shear and turbulence (Gerz *et al.*, 1998), and by radiative transfer processes and sedimentation (Unterstrasser and Gierens, 2010a,b) until complete mixing usually within a few hours.

From a computational perspective, the vortex regime is particularly challenging because of the intricate transformations that occur in the wake and changes in the flow topology, which manifest as abrupt expansions of the contrail in both horizontal and vertical directions (Sussmann, 1999). These transformations necessarily affect the ice microphysical properties. Most of numerical simulations of contrails in the vortex regime in the literature rely on Eulerian formulations for the treatment of ice particles. For example Lewellen and Lewellen (2001), Huebsch and Lewellen (2006) and Paugam *et al.* (2010) used three-dimensional LES with different levels of sophistication of ice microphysics, different set-ups of the dynamics and different resolution to cover timescales beyond the end of the vortex regime. The results of these studies showed that ambient humidity, stratification and turbulence are among the factors controlling the survival of ice crystals and their spatial distribution at the end of the vortex regime. These findings have been confirmed by LES coupled to a Lagrangian Particle Tracking (LPT) method in three-dimensional (Naiman *et al.*, 2011) and two-dimensional frameworks (Unterstrasser and Sölch, 2010), respectively. The latter authors also compared the results obtained with Eulerian and Lagrangian based methods on the same computational grid and observed that the LPT allow for a more accurate representation of ice microphysics. This aspect was recently investigated by Unterstrasser *et al.* (2014) in the context of tracer dispersion in aircraft wake vortices. They observed that LPT appears to be less dissipative than Eulerian formulations in reproducing particle dispersion in aircraft wake vortices although this conclusion has to be taken with caution because it is quite sensitive to the accuracy of the numerical scheme and the computational grid.

The dynamics of aircraft wake vortices has been studied traditionally in the wake vortex community for wake hazard applications. (The reader is invited to consult the reviews by Spalart (1998); Gerz *et al.* (2002); Holzäpfel *et al.* (2003) and references therein for further details on this subject). The interaction between wake vortices, turbulence and tracer dispersion has been studied in the past by Gerz and Holzäpfel (1999) and more recently by Misaka *et al.* (2012) who analyzed the fine-scale dynamics and the mixing of exhaust tracers with ambient air.

An aspect that has not been fully investigated so far is the role of atmospheric turbulence in determining the three-dimensional structure of contrails and the ice microphysical properties. For example, it is likely that turbulence enhances the mixing of ice crystals with atmospheric water vapor, especially in the secondary wake, and it influences the detrainment of exhausts at the vortex boundaries. In addition, vortex instabilities such as Crow instability (Crow,

1970) or elliptical instability (Widnall *et al.*, 1974) can be triggered by atmospheric turbulence, which ultimately lead to their break-up and decay. Unterstrasser *et al.* (2008) used 2D LES with modified diffusion of the vortex core to mimic the effects of three-dimensional vortex decay in a turbulent environment, a procedure that was also used by Unterstrasser and Gierens (2010a) and Unterstrasser and Sölch (2010). Lewellen *et al.* (2014) recently proposed a ‘quasi-3D’ approach which consists in resolving a certain narrow axial range of 3D eddies that are imposed a priori rather than emerging naturally from the turbulent cascade. Comparisons of mean contrail parameters with those obtained using full 3D LES were reported to be satisfactory for selected case studies. While the procedures described above are fast and suitable for sensitivity analysis of mean contrail properties to background turbulence, the treatment of vortex dynamics and the representation of atmospheric turbulence remain strongly parametrized. In fully 3D LES of contrail realized to date, turbulence models have been mainly used to trigger vortex instabilities rather than investigate the effects of background turbulence on the wake structure and the contrail properties. Paugam *et al.* (2010) introduced rather idealized perturbations in the form of sine waves, while Naiman *et al.* (2011), Lewellen *et al.* (2014), and Unterstrasser (2014) used divergence-free random fields with imposed spectrum which are evolved in order to develop coherent turbulent eddies that constitute the initial condition for the contrail simulations. However, during this transient time, turbulence decays, which makes difficult to control the level of the atmospheric turbulence that effectively interacts with the wake vortices.

This work is part of a larger project that aims at evaluating the atmospheric impact of contrail cirrus using a systematic chain of models and LES computations that cover the contrail lifetime from the formation to its demise in the atmosphere. While this strategy is similar to that employed recently by other authors who explored a large set of parameter space (Lewellen *et al.*, 2014; Unterstrasser, 2014), the present paper focuses on a specific point that is the fine-scale description of atmospheric turbulence and its effects on the mechanics of the wake and the contrail physics during the vortex phase -preliminary results of this work were presented by Picot *et al.* (2012). To that end, turbulence is generated by means of a spectral stochastic forcing technique that allows the nonlinear turbulent cascade to develop at smaller scales and was specially devised for strongly stratified turbulent flows as those occurring in the upper troposphere lower stratosphere (UTLS) (Paoli and Shariff, 2009; Paoli *et al.*, 2014). To the authors knowledge, this is the first time such sophisticated treatment of atmospheric turbulence is employed for contrail simulations. This point is crucial to have a correct representation of turbulence that is strongly anisotropic and hence substantially different and more complex than classical Kolmogorov turbulence. The application

of the forcing technique to the UTLS turbulence is described in the LES by Paoli *et al.* (2014) where the statistical properties were analyzed and compared with theoretical analysis of strongly stratified flows (Brethouwer *et al.*, 2007). The study showed that a substantial amount of turbulent kinetic energy is present at sub-kilometer scales relevant to the wake flow dynamics, which have then to be properly resolved. In particular, the LES data revealed the typical flow structures characterized by horizontally layered eddies or ‘pancakes’ (Riley and Lelong, 2000) that are a signature of the background field used for the present contrail simulations.

Although full unsteady spatial simulations of wake formation including the exhaust jets would help understanding the details of ice nucleation in rapidly changing thermodynamic conditions, they have not been performed so far as they would require far higher resolution and computational resources that are hardly available on present supercomputers. Hence, for the present study, the initial wake-flow field was reconstructed using data from temporal LES of the jet regime at a wake age of 10 s that takes into account the jet/vortex interaction process (Paoli *et al.*, 2013). In addition, the numerical model uses a fully 3D Lagrangian approach to track the crystals. Since turbulence can be fully controlled, this methodology allows for a sensitivity analysis of the wake and contrail properties in the vortex regime as a function of turbulence intensity. The sensitivity to background humidity and temperature is also discussed and compared to the sensitivity to turbulence. The paper is organized as follows: Sect. 2 summarizes the model equations, Sect. 3 describes computational setup of the simulations, and Sect. 4 presents the results of the simulations. Conclusions are drawn in Sect. 5.

2 Model equations

The computational model used in this work is based on an Eulerian-Lagrangian approach: the large-scale eddies of the gaseous atmosphere are solved with compressible Navier-Stokes equations, while ice crystals are treated by a Lagrangian tracking method and mass transfer between the two phases. As this model has already been presented elsewhere (Paoli *et al.*, 2008, 2013), only a summary is presented here. The density ρ , velocity \mathbf{u} , total energy $E = C_v T + \frac{1}{2} \mathbf{u}^2$ (where T is the temperature) and water vapor mass fraction Y_v are solution of the Favre-filtered Navier-Stokes equations:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = \omega_v, \quad (1a)$$

$$\frac{\partial \rho \mathbf{u}}{\partial t} + \nabla \cdot (\rho \mathbf{u} \otimes \mathbf{u}) + \nabla p = \nabla \cdot \mathbf{T} + \rho \mathbf{g}, \quad (1b)$$

$$\frac{\partial \rho E}{\partial t} + \nabla \cdot ((\rho E + p) \mathbf{u}) = \nabla \cdot (\mathbf{T} \mathbf{u}) + \rho \mathbf{g} \cdot \mathbf{u} - \nabla \cdot \mathbf{q}, \quad (1c)$$

$$\frac{\partial \rho Y_v}{\partial t} + \nabla \cdot (\rho Y_v \mathbf{u}) = -\nabla \cdot \boldsymbol{\xi} + \omega_v, \quad (1d)$$

where the source term ω_v in (1a) and (1d) represents the mass transfer between vapor and ice phases detailed in Sect. 2.1; the tensor $\mathbf{T} = 2\mu \left(\mathbf{S} - \frac{1}{3} S_{kk} \mathbf{1} \right)$ is the shear stress with

$\mathbf{S} = \frac{1}{2} (\nabla \mathbf{u} + \nabla \mathbf{u}^T)$, $\mu = \mu_{\text{mol}} + \mu_{\text{sgs}}$ where μ_{mol} and μ_{sgs} are, respectively, the molecular viscosity of air (mol) and the sub-grid scale (sgs) viscosity, the latter obtained from the filtered structure function model by (Ducros *et al.*, 1996); the vector \mathbf{g} is the gravitational acceleration; the pressure p is obtained from the perfect gas law $p = \rho R T$ with R the gas constant for air; the vectors $\mathbf{q} = \mathbf{q}_{\text{mol}} + \mathbf{q}_{\text{sgs}}$ and $\boldsymbol{\xi} = \boldsymbol{\xi}_{\text{mol}} + \boldsymbol{\xi}_{\text{sgs}}$ are the heat and vapor diffusion fluxes, respectively, each one composed of the molecular and sgs fluxes. Given the high Reynolds number flows considered in this study, the molecular transport are small compared to the turbulent transport so that in practice $\mu \simeq \mu_{\text{sgs}}$, $\mathbf{q} \simeq \mathbf{q}_{\text{sgs}}$ and $\boldsymbol{\xi} \simeq \boldsymbol{\xi}_{\text{sgs}}$. These are related to the gradient of temperature and vapor mass fraction by means of turbulent Prandtl and Schmidt numbers:

$\mathbf{q}_{\text{sgs}} = -C_p \frac{\mu^{\text{sgs}}}{Pr_t} \nabla T$ and $\boldsymbol{\xi}_{\text{sgs}} = -\frac{\mu^{\text{sgs}}}{Sc_t} \nabla Y_v$. For typical atmospheric flows Pr and Sc depends on Richardson number and other atmospheric parameters (see e.g. Schumann and Gerz (1995)), however at the resolution of $\mathcal{O}(1\text{ m})$ considered of here (see Tab. 1) the subgrid-scale turbulence can be considered isotropic (the Ozmidov scale in the UTLS is one order of magnitude larger, see e.g. Brethouwer *et al.* (2007)) and they are taken constant with values $Pr_t = Sc_t = 0.419$ as in Gerz and Holzäpfel (1999).

Equations (1) are discretized on collocated, Cartesian meshes with non-uniform grid spacing. The spatial derivatives are computed with a sixth-order compact scheme reformulated for stretched grids (Lele, 1992; Gamet *et al.*, 1999). A high-order selective artificial dissipation (Tam and Webb, 1993; Tam *et al.*, 1993) was used to enforce numerical stability while retaining the sixth-order spectral-like resolution at the resolved scales (see Barone (2003) for details). Finally, time integration is performed by a third-order Runge-Kutta method with a Courant-Friedrichs-Lewy number fixed to $CFL = 0.5$.

2.1 Ice model

The model uses a Lagrangian approach to track the trajectories of particles: ice crystals and nucleation sites (soot particles emitted by engines in the present study) onto which ice forms by deposition of water vapor. Given the large soot emission index $EI_s = \mathcal{O}(10^{15}) \text{ kg}^{-1}$ (Kärcher and Yu, 2009), it would be prohibitive to track all soot particles in the wake, so only a reduced number of numerical particles are tracked by the model. A numerical particle represents a cluster of $n_c = n/n_p$ (identical) physical particles that have mass m_p , center of mass \mathbf{x}_p and velocity \mathbf{u}_p . As particles remain small compared to flows structures, the point-force approximation (Boivin *et al.*, 1998) can be used for particle-flow interac-

tions, which means that all fluid properties at particle position are obtained through an interpolation of the values at neighborhood grid points:

$$\frac{d\mathbf{x}_p}{dt} = \mathbf{u}_p, \quad p = 1, \dots, n_p, \quad (2a)$$

$$\frac{d\mathbf{u}_p}{dt} = -\frac{\mathbf{u}_p - \mathbf{u}(\mathbf{x}_p)}{\tau_p} + \mathbf{g}, \quad p = 1, \dots, n_p \quad (2b)$$

where the relaxation time τ_p takes into account the aerodynamic drag of the numerical particle

$$\tau_p = \frac{\rho_{\text{ice}} d_p^2}{18\mu} \frac{1}{1 + 0.15 Re_p^{0.687}}, \quad Re_p = \frac{\rho d_p}{\mu} \|\mathbf{u}_p - \mathbf{u}(\mathbf{x}_p)\| \quad (3)$$

where $\rho_{\text{ice}} = 917 \text{ kg m}^{-3}$ is the density of ice and Re_p is the Reynolds number of the particle. The mass of vapor removed through deposition is

$$-\omega_v(\mathbf{x}) = \sum_p n_c \frac{dm_p}{dt} \delta(\mathbf{x} - \mathbf{x}_p) \quad (4)$$

where δ is the Dirac delta function. In numerical applications, the vapor mass must be removed from the nodes surrounding a numerical particles in order to conserve the total mass, which is done by replacing δ in Eq. (4) by a weighting function that depends on the inverse distance between the node and the particle (Boivin et al., 1998). Similar budgets can be established with drag momentum and thermal exchanges, but they are neglected in Eq. (1) because the mass ratio between ice and gaseous phases remain small.

To close the problem a model for the deposition rate dm_p/dt is needed in Eq. (4). The formation of nucleation sites is the result of complex microphysical processes that are beyond the scope of the present study (Kärcher et al., 1998). In a simplified picture, for a nucleation site to be activated, the air surrounding the particle needs to be saturated with respect to water to form a supercooled droplet that freeze afterwards. The time needed for this freezing depends on the particle size among other parameters but this is in general much smaller than the time step used for fluid dynamics simulations. Hence, as done in previous LES we assume instantaneous freezing (Unterstrasser and Sölch, 2010; Naiman et al., 2011; Paoli et al., 2013). This newly formed crystal subsequently grows by deposition of water vapor. Hence, in the present model, activation is represented by the flag

$$H_p^{\text{act}} = \begin{cases} 1 & \text{if } m_p > 0 \text{ or } Y_v(\mathbf{x}_p) \geq Y_v^{s,w}, \\ 0 & \text{else} \end{cases} \quad (5)$$

where $Y_v^{s,w} = Y_v^{s,w}(T(\mathbf{x}_p))$ and $Y_v^{s,w}$ is the equilibrium specific humidity with respect to water. This picture is consistent with recent studies (Kärcher and Yu, 2009) showing that in

the soot-rich regimes as those characterizing modern turbofans, the activation is a thermodynamically-controlled process. The deposition process is obtained by assuming spherical crystals of diameter d_p , i. e.

$$m_p = \frac{\pi}{6} \rho_{\text{ice}} d_p^3 \quad (6)$$

where the soot core size (of the order of nanometers) can be neglected compared to the mass of deposited ice. The deposition rate is (Kärcher et al., 1996)

$$\frac{dm_p}{dt} = H_p^{\text{act}} \frac{\pi}{2} d_p G(Kn_p) D_{v,p} \rho(\mathbf{x}_p) (Y_v(\mathbf{x}_p) - Y_v^{s,i}) \quad (7)$$

where $Kn_p = 2\lambda_{v,p}/d_p$ is the Knudsen number of the particle; $Y_v^{s,i} = Y_v^{s,i}(T(\mathbf{x}_p))$ and $Y_v^{s,i}$ is the equilibrium specific humidity with respect to ice; the collisional factor G accounts for the transition from gas kinetic to continuum regime (Kärcher et al., 1996)

$$G(Kn_p) = \left(\frac{1}{1 + Kn_p} + \frac{4}{3} \frac{Kn_p}{\alpha} \right)^{-1} \quad (8)$$

where the accommodation factor is taken $\alpha = 0.5$, the upper limit of the analysis by Kärcher et al. (1996) (we did not explore the sensitivity to this parameter in this study); the mean free path of vapor molecules in air $\lambda_{v,p} = \lambda_v(p(\mathbf{x}_p), T(\mathbf{x}_p))$ and the diffusion coefficient of vapor in air $D_{v,p} = D_v(p(\mathbf{x}_p), T(\mathbf{x}_p))$ are computed with relations that explicitly account for the dependence on pressure and temperature (Pruppacher and Klett, 1997)

$$\lambda_v(p, T) = \lambda_v^0 \left(\frac{p^0}{p} \right) \left(\frac{T}{T^0} \right), \quad (9)$$

$$D_v(p, T) = D_v^0 \left(\frac{p^0}{p} \right) \left(\frac{T}{T^0} \right)^{1.94} \quad (10)$$

with $p^0 = 1 \text{ atm}$, $T^0 = 0 \text{ °C}$, $\lambda_v^0 = 61.5 \text{ nm}$, and $D_v^0 = 0.211 \text{ cm}^2 \text{ s}^{-1}$. The equilibrium specific humidity are obtained from corresponding equilibrium vapor pressure through $Y_v^{s,\chi} = \varepsilon p_v^{s,\chi} / (\varepsilon p_v^{s,\chi} + (p - p_v^{s,\chi}))$ where $\chi = w, i$ denotes either water or ice and $\varepsilon \approx 0.6219$ is the ratio between vapor and dry air molar masses. The equilibrium vapor pressure in the temperature of interest, i. e. between 200 K to 300 K, is taken from the fit by Sonntag (1994). Following Kärcher et al. (1996) and unlike the latest work of Unterstrasser (2014) and Lewellen et al. (2014) the Kelvin effect is not considered. The impact of this assumption on the results relative to other potential effects is discussed in Sect. 4.3. It is readily seen from Eq. (7) that the variation of crystal size depends on the local ice saturation ratio

$$s(\mathbf{x}_p) = \frac{\rho_v(\mathbf{x}_p)}{\rho_{v,p}^{s,i}} = \frac{Y_v(\mathbf{x}_p)}{Y_v^{s,i}} \approx \frac{p_v(\mathbf{x}_p)}{p_{v,p}^{s,i}} \quad (11)$$

with respect to 1. Thus, the model can also deal with evaporation ($dm_p/dt < 0$) that occurs when $s < 1$. The lower limit for crystal shrinking is the soot core diameter $d_n = 20 \text{ nm}$.




Fig. 1. Vertical cross-sections of the computational domain and along the contrail longitudinal axis (left panel) and transverse axis (right panel). The mesh is regular in the subdomain $[0 \text{ m}, 400 \text{ m}] \times [-200 \text{ m}, 200 \text{ m}] \times [-350 \text{ m}, +100 \text{ m}]$. For the sake of clarity 1 out of 5 grid points are shown.




Fig. 2. Vertical (left) and horizontal (right) cross-sections of potential temperature difference $\theta - \theta_b$ in K for the atmospheric flow-field used in cases 2 and 4. The data from Paoli et al. (2014) are extracted in the regions delimited by the black boxes and used to initialize the present simulations.

Table 1. Reference values common to all cases. Aircraft data refer to a B747 (4-engines) aircraft. Exhaust jet values are expressed in meter of flight.

| Quantity | unit | Value(s) |
|---|--------------|-------------------------------|
| <i>Computational Domain</i> | | |
| Grid cells | | $519 \times 619 \times 100$ |
| Grid spacing $\Delta x \times \Delta y \times \Delta z$ | m | $1 \times 1 \times 4$ |
| Domain $L_x \times L_y \times L_z$ | m | $1000 \times 1000 \times 400$ |
| <i>Atmosphere</i> | | |
| Brunt-Väisälä frequency N | s^{-1} | 0.012 |
| Reference altitude z_0 | m | 11 000 |
| Reference pressure p_0 | Pa | 24 286 |
| <i>Wake vortices (B747 data)</i> | | |
| Initial circulation Γ | $m^2 s^{-1}$ | 565 |
| Core radius r_c | m | 4 |
| Initial separation b | m | 47 |
| <i>Exhaust jets (B747 data)</i> | | |
| Emitted vapor mass $4 \times M_{v,e}$ | $g m^{-1}$ | 4×3.75 |
| Emitted nucleation sites n_e | m^{-1} | 10^{12} |

3 Numerical Set-Up and Initial Condition

The numerical simulations are based on a temporal approach, which is commonly used in the large-eddy simulation (LES) of wake vortex systems. The computational domain is a box of lengths $L_x = 400$ m, $L_y = 1$ km, and $L_z = 1$ km in the axial, transverse, and vertical directions, respectively. The axial domain is chosen to capture the long-wave instability of the vortex system (Crow instability). The reference transverse and vertical coordinates $y_0 = 0$, $z_0 = 0$ for the computational domain corresponds to the position of the aircraft, which cruises at an altitude of 11 km. The mesh is regular with uniform grid spacing $\Delta x = 4$ m in the axial direction and $\Delta y = \Delta z = 1$ m in the region of the cross-sectional domain containing the wake, while it is stretched far away from the wake (see Fig. 1). Periodic and open boundary conditions are imposed in the axial and transverse direction respectively. In the vertical direction, all flow variables are relaxed toward a thermodynamic state representative of a stratified atmosphere with a uniform Brunt-Väisälä frequency (Paoli et al., 2013). The initial condition consists of an atmospheric flow-field onto which the wake of the aircraft is inserted.

The background atmosphere is representative of an upper troposphere lower stratosphere region, i. e. stably stratified. The background thermodynamic variables T_b , p_b and $\rho_b = p_b/(RT_b)$ depend on z and satisfy the relations

$$\frac{N^2}{g} = \frac{1}{\theta_b} \frac{d\theta_b}{dz}, \quad (12a)$$

$$\frac{dp_b}{dz} = -\rho_b g \quad (12b)$$

Table 2. Table of simulations. The parameter ϵ is the eddy dissipation rate of the background turbulence, η_0 is the relative turbulence intensity, T_0 is the temperature at flight level, and s_0 is the saturation ratio (in the whole domain). Across the following figures, each simulation is represented by a unique symbol or line pattern, which are indicated in this table.

| case | ϵ ($m^2 s^{-3}$) | η_0 | T_0 (K) | s_0 | Symbol & line |
|------|-----------------------------|----------|-----------|-------|------------------|
| 1 | 12.5×10^{-5} | 0.095 | 218 | 1.30 | \times ---- |
| 2 | 1.6×10^{-5} | 0.048 | 218 | 1.30 | \diamond ---- |
| 3 | 0.57×10^{-5} | 0.034 | 218 | 1.30 | $+$ ---- |
| 4 | 1.6×10^{-5} | 0.048 | 218 | 1.10 | \triangle ---- |
| 5 | 1.6×10^{-5} | 0.048 | 218 | 0.95 | ∇ ---- |
| 6 | 1.6×10^{-5} | 0.048 | 215 | 1.30 | \square ---- |

where $N = 0.012 s^{-1}$ is the Brunt-Väisälä frequency while θ_b is the background potential temperature defined by

$$\frac{\theta_b}{p_b^{(\gamma-1)/\gamma}} = \frac{T_b}{p_b^{(\gamma-1)/\gamma}}, \quad (13)$$

where $p_\theta = 10^5$ Pa and $\gamma = 1.4$ is the ratio of specific heats for air. Pressure and density at flight level are kept fixed at $p_b(z_0) = p_0 = 242.86$ hPa and $\rho_b(z_0) = \rho_0 = 0.3881 m^3 s^{-1}$ while temperature at flight level $T_b(z_0) = T_0$ is varied to analyze the effects of temperature on contrail properties (see Tab. 2). The background specific humidity $Y_{v,b}$ is chosen to keep vapor saturation with respect to ice s_0 constant in the computational domain: $Y_{v,b}(z) = s_0 Y_v^{s,i}(T_b(z))$. Vapor saturation s_0 is also varied to analyze the effects of humidity on contrail properties. Atmospheric (or background) turbulence flow-fields were provided by separate LES of such stratified flows with specific eddy dissipation rate ϵ . Because of atmospheric stratification, turbulence is organized in horizontally layered structures, reflecting the anisotropy of turbulent fluctuations in the horizontal and vertical directions as shown in Fig. 2. In the work by Paoli et al. (2014)), it was shown that a resolution of 4 m in the mid and low turbulence cases and 10 m in the strong turbulence case, are adequate to resolve the Ozmidov and buoyancy scales. Further details of the turbulent field are given in the Appendix.

The initial turbulent fields for the present simulations were extracted from the larger computational domain as shown by the black box in Fig. 2.

The initial wake of the aircraft is reconstructed using data from simulations of the jet regime at a wake age of $t = 10$ s (Paoli et al., 2013). These simulations are representative of a B747 aircraft in cruise conditions, i. e. composed of a pair of counter-rotating vortices and engine emissions including vapor and nucleation sites. Each vortex has a circulation of $\Gamma = 565 m^2 s^{-1}$, a core radius $r_c = 4$ m, and the vortices are spaced by $b = 47$ m. The mass of vapor emitted per unit flight distance is $M_{v,e} = 3.75 g m^{-1}$ per engine (the total emitted vapor is denoted by $M_{v,0} = 15 g m^{-1}$) and the




Fig. 3. Top views of iso-contours $\lambda_2 = \lambda_{2,0}/10$ (black) and $\lambda_2 = \lambda_{2,0}/400\,000$ (colored) in a horizontal plane xy at different instants for case 2. Black contours show the cores of the trailing vortices whereas the colored contours show the vortical structures of atmospheric turbulence. The development of instabilities is dominated by the long-wave (Crow) instability. The short-wave (elliptical) instability is also visible after $t = 2$ min as well as secondary vorticity structures around each trailing vortex. The displacement of the wake along the y direction is due to the oscillation of a large-scale energetic mode in the background turbulent field.

Table 3. Summary of main contrail characteristics: lifespan t_b of wake vortices; fraction of surviving crystals at the end of the vortex regime; and mean optical thickness $\bar{\delta}$ at the end of the vortex regime.

| case | Lifespan t_b s | Fraction of surviving crystals at 4 min % | Optical thickness $\bar{\delta}$ at 4 min — |
|------|---------------------|---|---|
| 1 | 95 | 99.89 | 0.22 |
| 2 | 132 | 99.74 | 0.22 |
| 3 | 111 | 99.66 | 0.22 |
| 4 | 132 | 74.71 | 0.06 |
| 5 | 132 | 2.78 | < 0.03 |
| 6 | 118 | 99.71 | 0.18 |

number of nucleation sites emitted per unit flight distance is $n_e = 10^{12} \text{ m}^{-1}$. Thus, the number of nucleation sites in the computational domain is $n = 4n_e L_z$, although the number of numerical particles is $n_p = 2 \times 10^6$. At this wake age all particles are already activated. In order to characterize the interaction between background turbulence and vortex dynamics, it is useful to introduce the relative turbulence intensity η_0 that represents the velocity ratio between turbulent fluctuations and vortex descent velocity (Crow and Bate, 1976):

$$\eta_0 = \frac{v_b}{w_0} = \frac{2\pi}{\Gamma} \sqrt[3]{\epsilon b^4}. \quad (14)$$

The effective Reynolds number based on circulation $\text{Re}_{\text{eff}} = \Gamma/\nu_{\text{tot}}$ where $\nu_{\text{tot}} = \nu_{\text{mol}} + \nu_{\text{sgs}}$, ranges between 10^3 and 10^4 based on the maximum turbulent viscosity in the

flow-field, and between 10^6 and 10^7 based on the mean turbulent viscosity. The simulations have been organized as follows: the reference simulation, labeled case 2, is based on a supersaturated atmosphere at $s_0 = 1.30$, $T_0 = 218$ K, and mild' turbulence intensity. The background turbulence field has been changed in cases 1 and 3 with 'strong' and weak' turbulence intensity, respectively (this nomenclature is the same used by Paoli *et al.* (2014)). Saturation has been reduced in case 4 and 5 to $s_0 = 1.10$ and $s_0 = 0.95$, respectively. Finally, temperature has been reduced in case 6 to $T_0 = 215$ K. All simulations parameters are summarized in Tab. 1 and Tab. 2.

4 Results

4.1 Dynamics of wake vortices

Wake vortices are detected using the λ_2 field, which is defined as the second eigenvalue of the symmetric tensor $\mathbf{S}^2 + \mathbf{\Omega}^2$ with \mathbf{S} and $\mathbf{\Omega}$ being the symmetric and the anti-symmetric parts of the velocity gradient tensor, respectively (Jeong and Hussain, 1995). The vortex cores are characterized by negative values of λ_2 with lower values corresponding to stronger vortices. The initial minimum value of λ_2 is found in the flow region inside the wake vortices, $\lambda_{2,0} = \min_{x,y,z} \{\lambda_2(x,y,z;0)\} = -49.67 \text{ s}^{-2}$, which is much smaller than the value corresponding to the vortex tubes of atmospheric turbulence, $\lambda_2^{\text{atm}} = -0.001 \text{ s}^{-2}$. The evolution of λ_2 iso-surfaces reported in Fig. 3 illustrates the mechanism of vortex instability for the reference case. It is known from stability analysis that counter-rotating vortices are prone to both long-wave (Crow) instability with wavelength $\lambda_{\text{LW}} = 8.6b$, and short-wave (elliptical) instability with wavelength $\lambda_{\text{SW}} = 0.37b$ (Crow, 1970; Widnall *et al.*, 1970). For the present study, these values are $\lambda_{\text{LW}} = 405 \text{ m}$ and $\lambda_{\text{SW}} = 17.4 \text{ m}$, respectively. The vortical structures identifying the wake vortices and the background turbulence are characterized by the values $\lambda_2 = \lambda_{2,0}/10$ and $\lambda_2 = \lambda_{2,0}/400000 \approx \lambda_2^{\text{atm}}/10$, respectively. The figure shows that turbulence triggers perturbations of the wake vortex system that develop until the two vortex tubes collide and break up. The largest flow structure corresponds to the size of the axial domain, and can then be identified with λ_{LW} . Starting at $t = 2.0 \text{ min}$, the vortex tubes start to get closer and eventually collide at $t = 2.2 \text{ min}$ at two axial locations, $x = 250 \text{ m}$ and $x = 350 \text{ m}$. In the other simulations, a single collision between the vortex tubes is observed, see Fig. 4. It is also interesting to observe the emergence of another flow structure with wavelength of about 25 m starting at $t = 1 \text{ min}$, which can be identified with a short-wave instability. Although this has a higher growth rate (Widnall *et al.*, 1970) than the long-wave instability, in the present case it is not sufficiently strong to overwhelm it and break the vortices before the latter develops. Elliptical instability in combination with Crow

instability have been studied theoretically (e.g. Fabre *et al.*, 2000), experimentally (e.g. Laporte and Leweke, 2000), and numerically (e.g. Paugam *et al.*, 2010). Their emergency depends on characteristics of the vortex system (vortex core radius and spacing) and on the initial perturbation of the vortex system. Typical perturbations include sinusoidal waves corresponding to the excited instability modes (Laporte and Corjon, 2000; Le Dizès and Laporte, 2002; Paugam *et al.*, 2010), broadband white noise and spectral perturbations satisfying model spectra of isotropic or anisotropic turbulence. The latter approach has been used in the wake vortex literature and particularly in the studies that attempted to analyze the interaction of the wake with the background turbulence (Lewellen and Lewellen, 1996; Gerz and Holzäpfel, 1999; Han *et al.*, 2000; Holzäpfel *et al.*, 2001) and more recently by Hennemann and Holzäpfel (2011) who focused on the identification of vortex flow topology, and by Misaka *et al.* (2012) who analyzed passive scalar transport in addition to the fine-scale structures of the wake. The main difference with the present study is that here the background turbulence is sustained, yet the 3D snapshots shown in these studies as well as the various pictures of wake instabilities reported in the literature Crow (1970); Chevalier (1973); Sussmann and Gierens (1999) corroborate the structures visualized in Fig. 3. Isometric views of iso-surfaces $\lambda_2 = \lambda_{2,0}/10$ and $\lambda_2 = \lambda_{2,0}/1000$ are drawn in Fig. 4 for cases 1, 2, 3, and 6 taken immediately before and after the vortex tubes collide. The snapshots are similar for the three cases, except for the double collision appearing in case 2, which insures the reproducibility of these flow structures by the present LES. In general, the vortex break-up always starts in the lowermost part of the wake because the wake descent is inversely proportional to the vortex separation (which goes to zero at the collision point).

In order to measure the lifespan of wake vortices, their intensities are evaluated as a function of time: the minimum value of λ_2 of each vortex is measured along the longitudinal axis. The condition $\lambda_2 < \lambda_{2,0}/10$ is used to discriminate wake vortices from secondary vortices. Noticing on Fig. 4 that this condition may not be reached in some parts of the longitudinal axis, we define the vortex length x_v as the length of the longitudinal axis range covering all axial sections where this condition is verified. The length ratio x_v/L_x and the average vortex intensity $\bar{\lambda}_2$ are plotted in Fig. 5 for each vortex and for cases 1,2,3, and 6. The magnitude of $\bar{\lambda}_2$ quickly decreases at the beginning, then its rate of variation stabilizes at a constant value of $-0.1|\lambda_{2,0}|$ per minute. The ratio x_v/L_x is initially equal to 1, meaning that coherent vortices can be detectable over all the axial domain. After a couple of minutes, the magnitude of $\bar{\lambda}_2$ decreases abruptly and falls below $-|\lambda_{2,0}|/10$ within a few seconds. In the meantime, x_v decreases and reaches zero within the same time span. The abrupt slope change is the mark of the vortex break-up as shown by snapshots of Fig. 4, which are taken at the times indicated by the vertical dashed lines in Fig. 5. The wake lifespan t_b is defined by the abrupt slope




Fig. 4. Iso-contours $\lambda_2 = \lambda_{2,0}/10$ (black) and $\lambda_2 = \lambda_{2,0}/1000$ (colored) taken immediately before (left panels) and after (right panels) the vortex tubes collide. Black iso-contours represent the wake vortices while colored iso-contours represent the secondary vortices.




Fig. 5. Time histories of vortex length ratios x_v/L_x and $\lambda_2/\lambda_{2,0}$, for cases 1,2,3 and 6. The horizontal dashed lines for each case indicate the threshold values $\lambda_2 = \lambda_{2,0}/10$ that is used to discriminate the wake vortex tubes from the background turbulence. The two vertical dashed lines correspond to the times selected in the iso-contours of Fig. 4 (i.e. immediately before and after the vortex break-up) and define the lifespans of the wake vortices.




Fig. 6. Lifespans of wake vortices as a function of relative turbulence intensity. The solid line is the analytical estimation by Crow and Bate (1976), the dashed line outlines lifespans obtained by numerical simulations of (Spalart and Wray, 1996) and black crosses are *in-situ* measurements (Crow and Bate, 1976). Simulations: \times case 1, \diamond case 2, $+$ case 3, and \square case 6.

change and values are given in Tab. 3. The results obtained for all cases studied are compared in Fig. 6 to the analytical estimation of Crow and Bate (1976) and contrasted with a limited set of measurements and numerical data found in the literature (Crow and Bate, 1976; Spalart and Wray, 1996). In spite of the large scatter of data (even with similar turbulence level), the numerical values of t_b are in the range of experimental values. All collected data show a similar trend: weaker relative turbulence leads to longer lifespan (as observed for example by Holzäpfel (2003)). Similar trends were recently reported using the normalized Brunt-Väisälä frequency $N^* = N t_0$ and the vortex sinking distance $z_b = w_0 \times t_b$ instead of the vortex lifespan t_b (Schumann, 2012; Jeßberger et al., 2013; Schumann et al., 2013) and Fig. 7 shows a general good agreement with those data.

In order to evaluate the vertical displacement of the wake vortices, vertical profiles $\tilde{\lambda}_2(z, t) = \min_{x,y} \{\lambda_2(x, y, z; t)\}$ are taken at different times and shown in Fig. 8. They descend roughly 100 m until $t = 1$ min, in agreement with previous observational studies (Sussmann and Gierens, 1999). The minimum values of λ_2 are coherent with values found in Fig. 5 and indicate the region containing the vortices. The vertical spread of this region increases at a rate of 20 m min^{-1} , which is due to the development of instabilities discussed in Fig. 3. We observed in Fig. 4 that the vertical displacement increases in the regions where the vortex separation goes to zero, this phenomenon is observed again in Fig. 8 as the vertical spreading has accelerated at $t = 2$ min in cases 1, 3, 6. Besides, the minimum value of $\tilde{\lambda}_2$ has in-




Fig. 7. Non-dimensional maximum descent z_b/b of wake vortices as a function of non-dimensional Brunt-Väisälä frequency $N \times t_0$ for the present simulations and a collection of experimental and numerical data (adapted from Schumann (2012)). Simulations: \times case 1, \diamond case 2, $+$ case 3, and \square case 6. Note that $z_b/b = w_0 \times t_b/b = t_b/t_0$ with $t_0 = 2\pi b_0^2/\Gamma$. In all cases considered in this study, $t_0 \simeq 25 \text{ s}$ and $N \times t_0 \simeq 0.3$ (see Tab. 1).

creased towards zero in cases 1, 3 in accordance with Fig. 5. At the end of the vortex regime, the primary wake is found between 200 m and 300 m below flight level.

4.2 Contrail microphysics

As explained in Sect. 2.1, vapor deposits on ice crystals at a rate that depends on the local ice supersaturation. This process is represented in Fig. 9, which shows the spatial distribution of ice crystals colored with diameter d_p for different levels of atmospheric turbulence. In the early stages of the vortex regime (until $t = 0.85$ min), the crystals are distributed in the primary wake with diameters ranging between $2 \mu\text{m}$ and $6 \mu\text{m}$. The vortex cores are visible and resemble pictures of vortex tubes (see for example Scorer and Davenport, 1970). Note that regions of low crystal density (i. e. far from the cores) contain larger crystals because, for given ambient supersaturation, the same amount of ambient water vapor has to be shared among less crystals. At the end of the vortex regime ($t = 1.5$ min), the vortex tubes are still visible and the secondary wake has started to form. Crystals remain relatively small in the primary wake while they grow mostly in the secondary wake where they are exposed to a supersaturated environment. The presence of atmospheric turbulence folds the structure of the secondary wake and the peripheral regions of the primary wake. At $t = 3$ min, the primary wake is rearranged into a puff as a consequence of the turbulent dissipation that results from the break-up (named induced turbulence hereafter). The figure shows that turbulence increases the mixing of the contrail with ambient air and favors the development of Crow instability. The vertical




Fig. 8. Vertical profiles of $\tilde{\lambda}_2/\lambda_{2,0}$ that is used to track the wake vortices as they move downwards and break up. The four profiles shown for each case correspond to ---- $t=0.5$ min, --- $t=1$ min, ... $t=1.5$ min, — $t=2$ min.

extension reduces with the turbulence intensity because the vortices tend to destabilize and break up earlier in the case of strong turbulence compared to weak turbulence. This is further confirmed by the vertical profiles of ice crystal number and mass reported in Fig. 10 taken at the same time as in Fig. 4.

Contrail diffusion is analyzed in Fig. 11 that shows the evolution of the contrail volume per unit flight distance V_p normalized by the initial volume $V_{p,0}$. The contrail volume is computed as the sum of volume of the grid-cells containing at least one crystal and defined over a regular mesh with 1 m of resolution. Up to $t = 1$ min, the volume expansion is similar for the three turbulence levels. Approximately at $t = t_b$, the volume expands faster with stronger turbulence, as also observed in snapshots of Fig. 9. It is interesting to note that the expansion rate is the same for all cases at the end of the simulation, even for case 6 ($T_0 = 215$ K). This means that the wake turbulence is the main contributor to contrail diffusion in the dissipation regime (even though atmospheric turbulence is expected to become predominant as wake turbulence dissipates).

Figure 12 shows the evolution of ice mass per meter of flight M_i normalized by the mass of vapor emitted by the engines $M_{v,0} = 15 \text{ g m}^{-1}$. The mass M_i is obtained using Eq. (6)

$$M_i = \frac{1}{L_z} \sum_p n_c m_p = \frac{1}{L_z} \sum_p n_c \frac{\pi}{6} d_p^3 \rho_{\text{ice}}. \quad (15)$$

At the beginning of the vortex regime, the emitted water vapor $M_{v,0}$ has been completely deposited on ice crystals. The ambient vapor entrained in the plume during the first 10 seconds (see e.g. Paoli et al. (2013)) also contribute to the overall ice mass so that initially $M_i > M_{v,0}$. When ambient air is subsaturated (case 5), the ice mass decreases exponentially, consistently with observations of non-persistent contrails. When ambient air is supersaturated (all but case 5), the ice mass reaches a maximum at $t = 25$ s. This maximum indicates that all available vapor (exhaust vapor plus atmospheric vapor trapped in the wake) has been converted into ice so that $s = 1$ inside the contrail. As vapor density in the atmosphere scales with s_0 , this maximum is higher when the atmosphere is more supersaturated. In cases 2, 3, 4, and 6, the ice mass decreases from $t = 1$ min to the end of the vortex regime. This decrease indicates the sublimation of crystals, explained by the process of adiabatic compression occurring in the primary wake (Unterstrasser et al., 2008). On the other hand, the mixing of the secondary wake with ambient air partly compensates sublimation so that the ice mass reduction is less pronounced when increasing the turbulence intensity, and even completely compensates sublimation in case 1. In the dissipation regime, the ice mass grows again as the consequence of induced turbulence. At $t = 4$ min, M_i reaches $2 M_{v,0}$ when $s_0 = 1.10$ (case 4), $4.5 M_{v,0}$ when $T_0 = 215$ K (case 6), and $8.5 M_{v,0}$ when $s_0 = 1.30$, $T_0 = 218$ K and regardless of the turbulence level. The less pronounced increase of ice mass for lower ambient temperature can be understood by observing that ambient saturation is kept constant between cases 2 and 6. Hence, the density of water vapor $\rho_v = s \rho_v^{s,i}(T)$ decreases when decreasing T (since $\rho_v^{s,i}$ is a monotonic function of temperature). In case 6, the mass of vapor entraining in the contrail and condensing into ice is then reduced compared to case 2 as shown in Fig. 12.

The vertical structure of the contrail is further analyzed in Fig. 13 in the dissipation regime. While the ‘curtain’ connecting the primary and secondary wakes forms as soon as the vortex pair start the descent, ice tends to substantially accumulate in the secondary wake after the break-up until, at $t = 4.5$, the two peaks are approximately the same for case 2 ($s_0 = 1.3$). For case 4 ($s_0 = 1.1$) most of ice crystals in the primary wake sublimated so that only the peak in the secondary wake is apparent at the end of the dissipation regime. (Note these results closely resemble those obtained recently by Unterstrasser (2014)).

The number of particles surviving the adiabatic compression is an important parameter to consider when evaluating




Fig. 9. Snapshots of ice crystals spatial distribution for cases 1, 2, and 3. Crystals are colored with diameters. The figure shows the development of the secondary wake at $t = 1.5$ min and $t = 3$ min (center and right panels) and the formation of puffs at $t = 3$ min (right panels). It can be observed that the size of crystals slightly increases during the instability process of the vortex regime ($t \leq 1.5$ min) whereas it increases considerably in the dissipation regime ($t = 3$ min) with larger crystals found in the secondary wake. Crystals appear more mixed in the strong atmospheric turbulence case (top panels).




Fig. 10. Vertical profiles of normalized number $\tilde{n}(z)/n_{\text{tot}}$ (left panels) and mass $\tilde{M}_i(z)/M_{v,0}$ (right panels) of ice crystals. The four profiles shown for each case correspond to ---- $t=0.5$ min, -.- $t=1$ min, $t=1.5$ min, — $t=2$ min.




Fig. 11. Normalized contrail volume per unit flight distance. Note the increased mixing following the vortex the break-up where the exhaust material is released into the atmosphere.




Fig. 12. Ice mass per unit flight distance normalized by the mass of emitted water vapor. Adiabatic compression reduces \tilde{M}_i at the end of the vortex regime, and this is particularly effective when the atmosphere is weakly supersaturated (case 4). The break-up of the vortices causes \tilde{M}_i to increase at a rate that depends on atmospheric temperature, saturation, and, to a much lesser extent, on the atmospheric turbulence.




Fig. 13. Vertical profiles of normalized mass $\tilde{M}_i(z)/M_{v,0}$. Left panel: profiles at different wakes ages in the dissipation regime for case 2. Right panel: comparison at $t = 4.5$ min between case 2 ($s_0 = 1.3$) and case 4 ($s_0 = 1.1$).




Fig. 14. Normalized number of activated particles (fraction of surviving crystals). Adiabatic compression is strong enough in weakly supersaturated atmosphere (case 4) to completely sublimate 30 % of crystals, but it is not able to evaporate any crystals in strongly supersaturated atmosphere.




Fig. 15. Ratio of deposited mass. In the vortex regime, the contrail is close to equilibrium ($M_i \approx M_c$) only when the atmosphere is strongly supersaturated. In the dissipation regime, the strong mixing between the contrail and ambient air causes a depart from the equilibrium state ($M_i < M_c$).

the global and climate impact of contrails (e.g. Burkhardt et al., 2010; Burkhardt and Kärcher, 2011). Figure 14 shows the fraction of surviving crystals, and their values at the end of the simulation are summarized in Tab. 3 for all considered cases. These data indicate that almost all crystals survive when $s_0 = 1.30$ and 75 % survive when $s_0 = 1.10$, which is higher than the results obtained by Unterstrasser and Sölch (2010) and recently by Unterstrasser (2014) and Lewellen et al. (2014). As discussed in Sect. 4.3, this can be caused

by the the different microphysical set-up (cf. inclusion of Kelvin effect) and/or the mixing efficiency predicted by the two models, especially in the peripheral region of the primary wake in Fig 9.

It is interesting to evaluate the mass of ice that would be formed by a model that would enforce equilibrium between ice and vapor phase at each time step. This kind of models may be attractive as they are less computationally expensive.

The equilibrium ice mass M_e is defined by

$$M_e = M_i + M_{v,a} \quad \text{with} \quad M_{v,a} = \int_{V_p} (\rho_v - \rho_v^{s,i}) dV_p. \quad (16)$$

and represents the ice mass of ice if all available vapor were instantaneously deposited onto crystals (note that $M_{v,a}$ is not the available mass of vapor in the ‘true’ situation). Figure 15 shows the ratio of deposited mass M_i/M_e . At $t = 25$ s, the ratio is close to one consistently with the equilibrium state suggested earlier in this section. The competition between ice sublimation due to the adiabatic compression and ice deposition by mixing of the secondary wake is seen again here at the end of the vortex regime. Turbulence favors mixing and entrainment of supersaturated ambient air into the plume. This in turn increases the ice deposition rate, which scales with the local supersaturation or $Y_v(\mathbf{x}_p)$ (the amount of vapor available at particle position). When turbulence is strong (case 1) deposition is stronger than sublimation as $M_i/M_e \lesssim 1$, whereas when turbulence is weak (case 3) sublimation is stronger than deposition as $M_i/M_e \gtrsim 1$. When supersaturation is reduced to $s_0 = 1.10$ (case 4), sublimation is much stronger than deposition and the equilibrium mass M_e approaches zero (while M_i/M_e diverges so the assumption of equilibrium is not valid in for $s_0 = 1.10$). In the dissipation regime, the ratio M_i/M_e reduces due to the mixing produced by the induced turbulence, and reaches a constant value of 0.7 when $s_0 = 1.30$ (regardless turbulence intensity or background temperature) and 0.45 when $s_0 = 1.10$. This result can be explained with an estimation of the rate of change of vapor in the contrail. On one hand, the quantity of vapor is increased by mixing at a rate of $Q_m = s_0 \rho_v^{s,i}(T_0) dV_p/dt$, neglecting temperature variations in the contrail vicinity. On the other hand, the quantity of vapor is decreased by deposition at a rate of $Q_d = n dm_p/dt = C n (s - 1) \rho_v^{s,i}(T_0)$ where C depends on the mean particles radius and can be assumed constant between 1 and 2 min as the ice mass (see Fig. 12). The temperature dependence of $G(Kn_p) D_{v,p}$ in Eq.(7) can reasonably be neglected in this context. Figure 15 shows that these rates are balanced at the end of the simulation. When T_0 is reduced, both Q_m and Q_d are decreased by the same amount, which does not change the balance (although, the time needed to reach this balance increases). When s_0 is reduced by 15 % from 1.30 to 1.10, Q_m is reduced by 15 % while Q_d is reduced (through n) by 25 %, the balance is then changed towards higher amounts of vapor in the contrail. A potential drawback in the definition of $M_{v,a}$ in Eq. 16 is that it depends on the definition of V_p , which may not be suitable to evaluate the non-equilibrium in the actual contrail. To that end, we calculated the mean saturation ratio \bar{s} by means of an ensemble average $s(\mathbf{x}_p)$ over all ice particles. Its evolution is shown in Fig. 16. In the middle of the vortex phase between 1 and 2 minutes, thermodynamic conditions are very close to equilibrium (relative humidity is slightly less than 100 %) because of the sublimation due to adiabatic heating balance-




Fig. 16. Mean saturation ratio computed as an ensemble average over all ice particles. The contrails is slightly sub-saturated during the initial vortex descent between 1 and 2 min and supersaturated in the dissipation phase after 2 min.




Fig. 17. Mean optical thickness. The horizontal black line represent the visibility criterion $\bar{\delta} > 0.03$ (Kärcher et al., 1996). The mean optical thickness decreases during the vortex regime and stabilizes during the dissipation regime. Its evolution depends mainly on atmospheric temperature and saturation and, to a lesser extent, on atmospheric turbulence. The symbols on the right of the figure shows the values of the optical thickness for different aircraft and atmospheric situations measured in the CONCERT campaign (Voigt et al., 2011) as reported by Jeßberger et al. (2013).

ing the deposition due to the entrainment of fresh ambient vapor (similar levels of relative humidity were observed for example by Naiman et al. (2011), their Fig. 8). In the dissipation phase humidity is greater than 100 % as the ice crystals originally trapped in the vortices are fully exposed to ambient vapor although it does not exceed 105 % for the conditions of this study.

The optical thickness of the contrail $\bar{\delta}$ is shown in Fig. 17. It is evaluated by first computing the optical thickness $\delta(\mathcal{S}_{xy})$

over each column $\mathcal{S}_{xy} = [x, x + \Delta x] \times [y, y + \Delta y] \times L_z$, and averaging δ over the regions where $\delta(\mathcal{S}_{xy}) > 0$. The sunlight is assimilated to a monochromatic wave of wavelength $\lambda_0 = 550$ nm, the refractive index of water is $\mu_0 = 1.31$, and the extinction coefficient Q is approximated by the anomalous diffraction theory (Van De Hulst, 1957):

$$Q(\rho) = 2 - \frac{4}{\rho} \sin \rho + \frac{4}{\rho^2} (1 - \cos \rho). \quad (17)$$

where $\rho \equiv 2\pi(\mu_0 - 1)d_p/\lambda_0$. Over a column \mathcal{S}_{xy} , the optical thickness is computed as follows

$$\delta(\mathcal{S}_{xy}) = \frac{1}{\Delta x \Delta y} \sum_{\mathbf{x}_p \in \mathcal{S}_{xy}} n_c \frac{\pi}{4} d_p^2 Q(\rho). \quad (18)$$

In Fig. 17, when ambient air is not saturated (case 5), $\bar{\delta}$ decreases exponentially and, by 2.5 min, it falls below the threshold $\bar{\delta} < 0.03$ used by Kärcher et al. (1996) as a visibility criterion. Although the visibility of a contrail does depend not only on the optical depth but on many other parameters (angle between observation and sun, contrast against background, aerosols between observer and contrail etc.), the simplistic treatment used here is in line with those employed in previous numerical simulations of contrails (Unterstrasser and Gierens, 2010a; Naiman et al., 2011). When ambient air is supersaturated, $\bar{\delta}$ increases by 25 % of the initial value by the time the contrail reaches the equilibrium state. Afterwards, $\bar{\delta}$ decreases by the end of the vortex regime, which is due to the microphysical processes mentioned above combined with the dilution of the contrail that reduces the number density of ice crystals and thus $\bar{\delta}$ in every cases. In case 4 ($s_0 = 1.10$), the more pronounced sublimation results in a stronger reduction of $\bar{\delta}$. In the early stages of the dissipation regime, the mean optical thickness has larger fluctuations that are possibly due to the induced wake turbulence and to the form of the extinction coefficient that is an oscillating function of the argument (in particular, the ice crystal diameter d_p). As the contrail expands and diffuses, these oscillations are damped and the optical thickness attains a value of around 0.06 when $s_0 = 1.10$ (case 4), 0.18 when $T_0 = 215$ K (case 6), and 0.22 when $s_0 = 1.30$, $T_0 = 218$ K, regardless of the turbulence level. Mean contrail optical thickness of the order of 0.2 to 0.3 were reported by Voigt et al. (2011) and Jeßberger et al. (2013) for the CONCERT campaign and similar aircraft. Jeßberger et al. (2013) also reported qualitatively similar results with the EULAG-LCM LES model (Sölch and Kärcher, 2010) and the CoCiP model (Schumann, 2012), i. e. $0.05 < \bar{\delta} < 0.1$ in weakly supersaturated atmospheres and $0.1 < \bar{\delta} < 1$ in strongly supersaturated atmospheres. Optical thickness is lower when s_0 or T_0 is reduced as they both reduce density of water in the atmosphere.

Figure 18 compares the particle size distributions with those obtained by *in situ* measurements (Schröder et al., 2000). Note that a computed size distribution represents an average over the whole contrail whereas measurements are




Fig. 18. Crystal diameter distributions at 5 s, 1 min, 3 min for the three levels of saturation and measurements from (Schröder et al., 2000) (bottom).

done locally. This may lead to significantly different values where the contrail is highly inhomogeneous such as in the peripheral regions of the contrail with low densities and large crystals. Nevertheless, Fig. 18 shows many common properties between simulations and measurements: the distributions have log-normal shapes which broadens and reduces in density as time advances. When $s_0 = 1.30$ (case 2) the distribution readily shifts towards larger sizes, whereas when $s_0 = 1.10$ (case 4) the distribution does not shift. At $t = 3$ min ‘sublimation tail’ appears in the size distribution containing small ice crystals (with diameter less than $0.1 \mu\text{m}$) that are prone to sublimate as the result of adiabatic compression. When $s_0 = 0.95$ the sublimation tail appears as soon as $t = 1$ min, the distribution decreases more quickly and shifts towards smaller sizes showing that the contrail sublimates as a whole. Measurements have smaller sizes and larger densities compared to simulations. This difference may be due to the different conditions encountered in the measurements and those used in the present computations, but also from the uncertainty in the number of nucleation sites as the same mass of water has to be shared on a different number of crystals. These data were obtained from different campaigns or independent flight measurements where the ambient conditions and the characteristics of the aircraft generating the contrail could be substantially different from those considered in this study. In more recent campaigns, Jeßberger et al. (2013) carried out time-averaged measurements to reduce effects of contrail heterogeneity and obtained much broader distributions. Compared to the present results, the number of large crystals ($d_p > 2 \mu\text{m}$) is equivalent, but they measured an important number of smaller crystals ($d_p < 2 \mu\text{m}$) than the present model. This could be due to the limited number of numerical particles and should lead to an underestimation of $\bar{\delta}$. However, the smaller coefficient $d_p^2 Q (2\pi(\mu_0 - 1) \frac{d_p}{\lambda_0})$ in Eq. (18) reduces this underestimation. Despite these differences, Fig. 18 shows a reasonable agreement in terms of order of magnitude and shape of the crystal diameter distribution.

4.3 Impact of Kelvin effect on contrail properties

For the high supersaturated cases, $s_0 = 1.3$, the present study consistently shows a reduced crystal loss with more than 99% crystals surviving whereas for example Unterstrasser (2014) predicts that 90% crystals survive at $s_0 = 1.4$. As pointed by Lewellen et al. (2014) and confirmed by Unterstrasser (2014), the onset of crystal loss is delayed when the Kelvin effect in the ice growth rate is switched off. For the sake of comparison with these latest studies we ran two additional LES corresponding to cases 2 and 4 with Kelvin effect activated. This was done by replacing the ice saturation pressure $p_{v,p}^{s,i}$ by $p_{v,p}^{s,i} \times \exp(a_k/d_p)$ where a_k is a parameter that depends (among other variables) on the ice-vapor surface tension, in analogy to liquid water surface tension (Lewellen, 2012). Given the large uncertainty of ice-vapor tension, the




Fig. 19. Normalized number of activated particles (fraction of surviving crystals).

parameter was taken constant, $a_k = 10^{-9}$ m as in Lewellen et al. (2014). Figure 19 shows that the Kelvin effect increases the ice crystals loss although its impact is less pronounced compared to the aforementioned study especially in the highest supersaturation case. Besides the different ice microphysical treatment, other factors that can potentially affect this difference are: the different computational set-up and the numerical scheme (high-order vs low-order numerical schemes, upwind vs centered schemes) used for the discretization of the deposition rate. Finally, Fig. 19 also indicates that the contrail optical thickness (which scales with $\langle d_p^2 \rangle$) and mass (which scales with $\langle d_p^3 \rangle$, not shown) are insensitive to Kelvin effect, which is also consistent with Lewellen et al. (2014) results (their Fig. 8). This can be explained by the fact that the small particles are the most prone to sublimate and hence are most likely affected by the increase in the ice saturation pressure due to Kelvin effect, but they are also those that contribute the less to the moments of the size distribution. This is an important point to keep in mind when carrying on radiative calculations, particularly in the diffusion regime when the contrail transitions into a cirrus.

5 Discussion and conclusions

This study presented the results of three-dimensional large-eddy simulations of contrail evolution in the vortex and dissipation regimes of an aircraft wake immersed in a turbulent atmospheric flow-field. The computational model is based on an Eulerian-Lagrangian two-phase flow formulation where clusters of ice crystals are tracked as they move in the wake. The background turbulence and the initial condition for the contrail at the end of the jet regime were both generated from appropriate simulations. The focus of the study is to evaluate the effects of atmospheric turbulence on the wake dynamics and the contrail properties, and to compare with the effects of ambient humidity and temperature. The results showed a

good agreement with numerical and experimental literature work, in terms of descent and lifespan of wake vortices, overall mass of ice produced, and optical depth. Visual patterns of the primary and secondary wakes, Crow instability and the formation of ‘puffs’ also resembled qualitatively those found in observational analysis. The agreement of particle diameter distribution was also acceptable given the large data scatter and uncertainty of both ambient conditions and soot particle emissions in the experimental campaigns compared to those considered in the present study.

The main effect of atmospheric turbulence in the vortex regime is to trigger instabilities in the wake vortices and to accelerate their descent. Stronger turbulence accelerates the onset of the instability, leading to shorter contrail descent and more effective mixing in the interior of the plume. These results are in line with those published in recent wake vortex literature (Holzäpfel *et al.*, 2001; Hennemann and Holzäpfel, 2011; Misaka *et al.*, 2012) –although the latter authors analyzed weaker turbulence scenarios than ours– and are then a solid basis to investigate their impact on contrail microphysics. These effects influence the lifetime of the vortices, the vertical extent of the contrail and the number of surviving crystals. Atmospheric turbulence is then an important driver of the contrail evolution in the vortex and dissipation regime, which cannot be captured when the vortex instability is triggered by explicitly forcing one or more specific modes of the vortex system (Paugam *et al.*, 2010). However, it is found that the self-induced turbulence that is produced in the wake after the vortex break-up dominates over the background turbulence and effectively controls the mixing of the wake with ambient air. As a result, the intensity of the latter has small impact on the microphysical and optical properties at time scales of 4 min after emission. On the other hand, contrail microphysics is strongly influenced by atmospheric temperature and saturation which control the mass of water vapor in contrail surroundings, and the number of surviving crystals. In addition, the mixing induced by the wake turbulence generated during the vortex break-up is able to break the equilibrium between water vapor and ice phases inside the contrail, reducing the mass of ice by more than 30 % compared to model that would enforce equilibrium. Note that, once the wake turbulence has decayed, one can expect that background turbulence would be again the main driver of contrail dynamics in the early diffusion regime. This is the object of current investigation.

The mean contrail properties are in line with recent results obtained by numerical simulations that explored a larger set of parameter space (Lewellen *et al.*, 2014; Unterstrasser, 2014). However, for the high supersaturated cases, $s_0 = 1.3$, the present study consistently shows a reduced crystal loss with more than 99% crystals surviving whereas for example Unterstrasser (2014) predicts that 90% crystals survive at $s_0 = 1.4$. As pointed by Lewellen *et al.* (2014) and confirmed by Unterstrasser (2014), the onset of crystal loss is delayed when the Kelvin effect in the ice growth rate is switched off,

which can explain the difference with the present study in which Kelvin effect is not considered. Another factor that can potentially impact the prediction of crystal loss is the treatment of vapor mixing. Indeed, the deposition rate, Eq. 7, depends on the local vapor field which is sensitive to the numerical scheme adopted for scalar advection (Unterstrasser *et al.*, 2014), especially in the zones of high gradients as those occurring in the dissipation regime where the exhausts trapped in the wake is released and mix efficiently with ambient air. Whether it is the microphysical set-up or the order and resolution proprieties of the scalar advection schemes (e.g. high-order vs low-order numerical schemes, upwind vs centered schemes) that is mainly responsible for the difference in the deposition rate and the crystal loss in this phase is a point that deserves further investigation in follow-up studies.

Future work includes a sensitivity analysis of contrail characteristics to the initial number of nucleation sites n . The contrail properties are expected to change throughout its evolution: with increasing n , smaller crystals are expected to be found in the vortex regime as the same mass has to be shared with more crystals, and the contrail is expected to be closer to equilibrium in the dissipation regime as the rate of vapor deposition increases with n in Eq. (4). Finally, the data obtained in this work are being used to reconstruct the initial conditions for the simulations of the diffusion regime and the contrail-to-cirrus transition in an ongoing study. The long term objective is to contribute to the development of subgrid parameterizations for global models that can be used to investigate and evaluate the global radiative effects of contrail cirrus.

6 Appendix

6.1 Details of the atmospheric turbulence field

In the large domain used to generate the ambient turbulence ($L_{\text{box}} = 4 \text{ km}$, $\Delta = 4 \text{ m}$) the root mean squares (rms) of horizontal velocity fluctuations were $\sigma_u \simeq \sigma_v \simeq 0.4 \text{ ms}^{-1}$, while the anisotropy ratio between the rms of horizontal and vertical fluctuations was $\sigma_u/\sigma_w \simeq 4$ (see Table 1 in Paoli *et al.* (2014)). The integral length scale (estimated as

$$L_t = \frac{3\pi}{4\mathcal{K}} \int_0^\infty \frac{E(k)}{k} dk \text{ with } \mathcal{K} = 1/2(\sigma_u^2 + \sigma_v^2 + \sigma_w^2) \text{ the turbulent kinetic energy, } k \text{ the horizontal wavenumber, } E(k) \text{ the kinetic energy horizontal spectrum), ranged between 700 m and 1 km. Because the ratio } L_t/b \gg 1, \text{ the integral length scale does not significantly affect the vortex decay as observed in previous studies (e.g. Crow and Bate, 1976; Hennemann and Holzäpfel, 2011).}$$

When the atmospheric turbulent flow-field is truncated within the smaller domain denoted by the black box in Fig. 1, it will adapt to the new boundary conditions with a time scale that can be estimated as $\tau = \sigma_{u_{\text{vor}}}^2/\epsilon$ where $\sigma_{u_{\text{vor}}}^2$ is the rms

Table 4. Horizontal and vertical rms velocities measured in the regular subdomain.

| η_0 | $\sigma_u^{\text{vor}} (\text{m s}^{-1})$ | $\sigma_w^{\text{vor}} (\text{m s}^{-1})$ | $\tau (\text{s})$ |
|-----------------------|---|---|-------------------|
| 0.095 (case 1) | 0.41 | 0.15 | 47 |
| 0.048 (cases 2,4,5,6) | 0.23 | 0.08 | 15 |
| 0.095 (case 3) | 0.17 | 0.05 | 8 |

velocity calculated in the regular portion of the contrail domain (and similar expression for the vertical direction). We reported these values in Tab. 4, which shows that in all cases $\tau > 5 \text{ min}$. Hence, we can reasonably assume that ambient turbulence remains frozen (and the condition $L_t/b \gg 1$) remains valid during the vortex phase.

6.2 Details of sgs turbulent viscosity

The subgrid scale model in this study is based on filtered structure function by Ducros *et al.* (1996) that provides at each grid node the turbulent viscosity reconstructed locally using the nine closest grid points surrounding the grid node. Figure 20 below shows the evolution of the min, max and mean values of turbulent viscosity in the field, normalized by the molecular viscosity ν_{mol} . We can see for the mean turbulent viscosity reaches a decent factor of 10 times the molecular viscosity at the break-up time (where the gradients are the largest). Peak values of this ratio are of course larger although they are local in nature as they correspond to specific points in the flow-field. We computed the effective Reynolds number defined based on the circulation, $Re_{\text{eff}} = \Gamma/\nu_{\text{tot}}$ with $\nu_{\text{tot}} = \nu_{\text{mol}} + \nu_{\text{sgs}}$. The values based on the maximum turbulent viscosity (worst situation) are around 10^4 expect a short interval at the break-up time and in line with previous LES of the vortex phase (e.g. Gerz and Holzäpfel (1999))

Acknowledgements. Financial support from the Direction Générale de l'Aviation Civile through the project TC2 (Traînés de Condensation et Climat) is gratefully acknowledged. Computational resources were provided by CINES supercomputing center.

References

- Barone, M. F.: Receptivity of compressible mixing layers, Ph.D. thesis, Stanford University, 2003.
- Boivin, M., Simonin, O., and Squires, K. D.: Direct numerical simulation of turbulence modulation by particles in isotropic turbulence, *J. Fluid Mech.*, 375, 235–263, doi:10.1017/S0022112098002821, 1998.
- Brethouwer, G., Billant, P., Lindborg, E., and Chomaz, J.-M.: Scaling analysis and simulation of strongly stratified turbulent flows, *J. Fluid Mech.*, 585, 343, doi:10.1017/S0022112007006854, 2007.
- Burkhardt, U. and Kärcher, B.: Global radiative forcing from contrail cirrus, *Nat. Clim. Chang.*, 1, 54–58, doi:10.1038/nclimate1068, 2011.
- Burkhardt, U., Kärcher, B., and Schumann, U.: Global Modeling of the Contrail and Contrail Cirrus Climate Impact, *B. Am. Meteorol. Soc.*, 91, 479–484, doi:10.1175/2009BAMS2656.1, 2010.
- Chevalier, H.: Flight Test Studies of the Formation and Dissipation of Trailing Vortices, Tech. Rep. 730295, SAE International, Warrendale, PA, 1973.
- Crow, S. C.: Stability Theory for a Pair of Trailing Vortices, *AIAA Journal*, 8, 2172–2179, doi:10.2514/3.6083, 1970.
- Crow, S. C. and Bate, E. R., J.: Lifespan of trailing vortices in a turbulent atmosphere, *J. Aircraft*, 13, 476–482, doi:10.2514/3.44537, 1976.
- Ducros, F., Comte, P., and Lesieur, M.: Large-eddy simulation of transition to turbulence in a boundary layer developing spatially over a flat plate, *J. Fluid Mech.*, 326, 1–36, doi:10.1017/S0022112096008221, 1996.
- Fabre, D., Cossu, C., and Jacquin, L.: Spatio-temporal development of the long and short-wave vortex-pair instabilities, *Phys. Fluids*, 12, 1247–1250, doi:10.1063/1.870375, 2000.
- Gamet, L., Ducros, F., Nicoud, F., and Poinso, T. J.: Compact finite difference schemes on non-uniform meshes. Application to direct numerical simulations of compressible flows, *Int. J. Numer. Meth. Fl.*, 29, 159–191, doi:10.1002/(SICI)1097-0363(19990130)29:2<159::AID-FLD781>3.0.CO;2-9, 1999.
- Gerz, T. and Ehret, T.: Wingtip vortices and exhaust jets during the jet regime of aircraft wakes, *Aerosp. Sci. Technol.*, 1, 463–474, doi:10.1016/S1270-9638(97)90008-0, 1997.
- Gerz, T. and Holzäpfel, F.: Wing-Tip Vortices, Turbulence, and the Distribution of Emissions, *AIAA Journal*, 37, 1270–1276, doi:10.2514/2.595, 1999.




Fig. 20. Top panel: Evolution of the ratio of turbulent-to-molecular viscosity (minimum, maximum and mean values in the flow-field). Bottom panel: Evolution of the effective Reynolds number abased on circulation and the total viscosity: $Re_{\text{eff}} = \Gamma/\nu_{\text{tot}}$ with $\nu_{\text{tot}} = \nu_{\text{mol}} + \nu_{\text{sgs}}$. The Reynolds number based on the molecular viscosity is also plotted for reference.

- Gerz, T., Dürbeck, T., and Konopka, P.: Transport and effective diffusion of aircraft emissions, *J. Geophys. Res.*, 103, 25 905–25 913, doi:10.1029/98JD02282, 1998.
- Gerz, T., Holzäpfel, F., and Darracq, D.: Commercial aircraft wake vortices, *Prog. Aerosp. Sci.*, 38, 181–208, doi:10.1016/S0376-0421(02)00004-0, 2002.
- Han, J., Lin, Y.-L., Schowalter, D. G., Pal Arya, S., and Proctor, F. H.: Large Eddy Simulation of Aircraft Wake Vortices Within Homogeneous Turbulence: Crow Instability, *AIAA Journal*, 38, 292–300, doi:10.2514/2.956, 2000.
- Hennemann, I. and Holzäpfel, F.: Large-eddy simulations of aircraft wake vortex deformation and topology, *J. Aerospace Eng.*, 25, 1336–1350, doi:10.1177/0954410011402257, 2011.
- Holzäpfel, F.: Probabilistic two-phase wake vortex decay and transport model, *J. Aircraft*, 40, 323–331, doi:10.2514/2.3096, 2003.
- Holzäpfel, F., Gerz, T., and Baumann, R.: The turbulent decay of trailing vortex pairs in stably stratified environments, *Aerosp. Sci. Technol.*, 5, 95–108, doi:10.1016/S1270-9638(00)01090-7, 2001.
- Holzäpfel, F., Hofbauer, T., Darracq, D., Moet, H., Garnier, F., and Gago, C. F.: Analysis of wake vortex decay mechanisms in the atmosphere, *Aerosp. Sci. Technol.*, 7, 263–275, doi:10.1016/S1270-9638(03)00026-9, 2003.
- Huebsch, W. W. and Lewellen, D. C.: Sensitivity studies on contrail evolution, in: 36th AIAA Fluid Dynamics Conference and Exhibit, San Francisco, California, doi:10.2514/MFDC06, 2006.
- Jeong, J. and Hussain, F.: On the identification of a vortex, *J. Fluid Mech.*, 285, 69–94, doi:10.1017/s0022112095000462, 1995.
- Jeßberger, P., Voigt, C., Schumann, U., Sölch, I., Kaufmann, S., Petzold, A., Schäuble, D., and Gayet, J.: Aircraft type influence on contrail properties, *Atmos. Chem. Phys.*, 38, 11 965–11 984, doi:10.5194/acp-13-11965-2013, 2013.
- Kärcher, B. and Yu, F.: Role of aircraft soot emissions in contrail formation, *Geophys. Res. Lett.*, 36, L01 804, doi:10.1029/2008GL036649, 2009.
- Kärcher, B., Peter, T., Biermann, U. M., and Schumann, U.: The Initial Composition of Jet Condensation Trails, *J. Atmos. Sci.*, 53, 3066–3083, doi:10.1175/1520-0469(1996)053<3066:TICOJC>2.0.CO;2, 1996.
- Kärcher, B., Busen, R., Petzold, A., Schröder, F. P., Schumann, U., and Jensen, E. J.: Physicochemistry of aircraft-generated liquid aerosols, soot, and ice particles 2. Comparison with observations and sensitivity studies, *J. Geophys. Res.-Atmos.*, 103, 17,129–17,147, doi:10.1029/98JD01045, 1998.
- Laporte, F. and Corjon, A.: Direct numerical simulations of the elliptic instability of a vortex pair, *Phys. Fluids*, 12, 1016–1031, doi:10.1063/1.870357, 2000.
- Laporte, F. and Leweke, T.: Elliptic Instability of Counter-Rotating Vortices: Experiment and Direct Numerical Simulation, *AIAA J.*, 40, 2483–2494, doi:10.2514/2.1592, 2000.
- Le Dizès, S. and Laporte, F.: Theoretical predictions for the elliptical instability in a two-vortex flow, *J. Fluid Mech.*, 471, 169–201, doi:10.1017/S0022112002002185, 2002.
- Lee, D. S., Fahey, D., Forster, P., Newton, P., Wit, R., Lim, L., Owen, B., and Sausen, R.: Aviation and global climate change in the 21st century, *Atmos. Environ.*, 43, 3520–3537, doi:10.1016/j.atmosenv.2009.04.024, 2009.
- Lele, S. K.: Compact finite difference schemes with spectral-like resolution, *J. Comput. Phys.*, 103, 16–42, doi:10.1016/0021-9991(92)90324-R, 1992.
- Lewellen, D. C.: Analytic Solutions for Evolving Size Distributions of Spherical Crystals or Droplets Undergoing Diffusional Growth in Different Regime, *J. Atmos. Sci.*, 69, 417–434, doi:10.1175/JAS-D-11-029.1, 2012.
- Lewellen, D. C. and Lewellen, W. S.: Large-eddy simulations of the vortex-pair breakup in aircraft wakes, *AIAA J.*, 34, 2337–2345, 1996.
- Lewellen, D. C. and Lewellen, W. S.: The effects of aircraft wake dynamics on contrail development, *J. Atmos. Sci.*, 58, 390–406, doi:10.1175/1520-0469(2001)058<0390:TEOAWD>2.0.CO;2, CCC:000166475500005, 2001.
- Lewellen, D. C., Meza, O., and Huebsch, W. W.: Persistent contrails and contrail cirrus. Part 1: Large-eddy simulations from inception to demise, *J. Atmos. Sci.*, 71, 4399–4419, doi:10.1175/JAS-D-13-0316.1, 2014.
- Misaka, T., Holzäpfel, F., Hennemann, I., Gerz, T., and Manhart, M.: Vortex bursting and tracer transport of a counter-rotating vortex pair, *Phys. Fluids*, 24, 025 104, doi:10.1063/1.3684990, 2012.
- Naiman, A. D., Lele, S. K., and Jacobson, M. Z.: Large eddy simulations of contrail development: Sensitivity to initial and ambient conditions over first twenty minutes, *J. Geophys. Res.*, 116, D21 208, doi:10.1029/2011JD015806, 2011.
- Paoli, R. and Shariff, K.: Turbulent Condensation of Droplets: Direct Simulation and a Stochastic Model, *J. Atmos. Sci.*, 66, 723–740, doi:10.1175/2008JAS2734.1, 2009.
- Paoli, R., Vancassel, X., Garnier, F., and Mirabel, P.: Large-eddy simulation of a turbulent jet and a vortex sheet interaction: particle formation and evolution in the near field of an aircraft wake, *Meteorol. Z.*, 17, 131–144, doi:10.1127/0941-2948/2008/0278, 2008.
- Paoli, R., Nybelen, L., Picot, J., and Cariolle, D.: Effects of jet/vortex interaction on contrail formation in supersaturated conditions, *Phys. Fluids*, 25, 053 305–053 305–28, doi:10.1063/1.4807063, 2013.
- Paoli, R., Thouron, O., Escobar, J., Picot, J., and Cariolle, D.: High-resolution large-eddy simulations of stably stratified flows: application to subkilometer-scale turbulence in the upper troposphere–lower stratosphere, *Atmos. Chem. Phys.*, 14, 5037–5055, doi:10.5194/acp-14-5037-2014, 2014.
- Paugam, R., Paoli, R., and Cariolle, D.: Influence of vortex dynamics and atmospheric turbulence on the early evolution of a contrail, *Atmos. Chem. Phys.*, 10, 3933–3952, doi:10.5194/acp-10-3933-2010, 2010.
- Picot, J., Paoli, R., Thouron, O., and Cariolle, D.: Effects of atmospheric turbulence and humidity on the structure of a contrail in the vortex phase, in: TAC-3 Proceedings, pp. 110–113, Prien am Chiemsee, doi:10.2514/MFDC06, 2012.
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, vol. 18 of *Atmospheric and Oceanographic Sciences Library*, Kluwer Academic Publishers, Dordrecht, the Netherlands, 2nd edn., 1997.
- Riley, J. J. and Lelong, M.-P.: Fluid Motions in the Presence of Strong Stable Stratification, *Annu. Rev. Fluid Mech.*, 32, 613–657, doi:10.1146/annurev.fluid.32.1.613, 2000.
- Schröder, F., Kärcher, B., Durour, C., Ström, J., Petzold, A., Gayet, J., Strauss, B., Wendling, P., and Borrmann, S.: On the transition of contrails into cirrus clouds, *J. Atmos. Sci.*, 57, 464–480, doi:10.1175/1520-0469(2000)057<0464:OTTOCI>2.0.CO;2, 1999.

- 2000.
- 1190 Schumann, U.: On Conditions for Contrail Formation from Aircraft Exhausts, *Meteorol. Z.*, 5, 4–23, 1996. ¹²⁵⁰
- Schumann, U.: A contrail cirrus prediction model, *Geosci. Model Dev.*, 5, 543–580, doi:10.5194/gmd-5-543-2012, 2012.
- 1195 Schumann, U. and Gerz, T.: Turbulent mixing in stably stratified flows, *J. Appl. Meteorol.*, 34, 33–48, doi:10.1175/1520-0450-34.1.33, 1995. ¹²⁵⁵
- Schumann, U., Jeßberger, P., and Voigt, C.: Contrail ice particles in aircraft wakes and their climatic importance, *Geophys. Res. Lett.*, 25, 1336–1350, doi:10.1002/grl.50539, 2013.
- 1200 Scorer, R. S. and Davenport, L. J.: Contrails and aircraft downwash, *J. Fluid Mech.*, 43, 451–464, doi:10.1017/S0022112070002501, ¹²⁶⁰ 1970.
- Sonntag, D.: Advancements in the field of hygrometry, *Meteorol. Z.*, 3, 51–66, 1994.
- 1205 Spalart, P. R.: On the motion of laminar wing wakes in a stratified fluid, *J. Fluid Mech.*, 327, 139–160, ¹²⁶⁵ doi:10.1017/S002211209600849X, 1996.
- Spalart, P. R.: Airplane Trailing Vortices, *Annu. Rev. Fluid Mesh.*, 30, 107–138, doi:10.1146/annurev.fluid.30.1.107, 1998.
- 1210 Spalart, P. R. and Wray, A. A.: Initiation of the Crow Instability by Atmospheric Turbulence, in: *The characterisation & modification of wakes from lifting vehicles in fluids*, vol. 584, AGARD, Trondheim, 1996.
- Sussmann, R.: Vertical dispersion of an aircraft wake: Aerosol-lidar analysis of entrainment and detrainment in the vortex regime, *J. Geophys. Res.*, 104, 2117–2129, doi:10.1029/1998JD200033, ¹²¹⁵ 1999.
- Sussmann, R. and Gierens, K. M.: Lidar and numerical studies on the different evolution of vortex pair and secondary wake in young contrails, *J. Geophys. Res.*, 104, PP. 2131–2142, doi:10.1029/1998JD200034, 1999.
- 1220 Sölch, I. and Kärcher, B.: A large-eddy model for cirrus clouds with explicit aerosol and ice microphysics and Lagrangian ice particle tracking, *Q. J. Roy. Meteor. Soc.*, 136, 2074–2093, ¹²²⁵ doi:10.1002/qj.689, 2010.
- Tam, C. K. W. and Webb, J. C.: Dispersion-Relation-Preserving Finite Difference Schemes for Computational Acoustics, *J. Comput. Phys.*, 107, 262–281, doi:10.1006/jcph.1993.1142, 1993.
- 1230 Tam, C. K. W., Webb, J. C., and Dong, Z.: A study of the short wave components in computational acoustics, *J. Comput. Acoust.*, 1, 1–30, doi:10.1142/S0218396X93000020, 1993.
- Unterstrasser, S.: Large-eddy simulation study of contrail microphysics and geometry during the vortex phase and consequences on contrail-to-cirrus transition, *J. Geophys. Res.*, 119, 7537–7555, doi:10.1002/2013JD021418, 2014. ¹²³⁵
- Unterstrasser, S. and Gierens, K.: Numerical simulations of contrail-to-cirrus transition – Part 1: An extensive parametric study, *Atmos. Chem. Phys.*, 10, 2017–2036, doi:10.5194/acp-10-2017-2010, 2010a.
- 1240 Unterstrasser, S. and Gierens, K.: Numerical simulations of contrail-to-cirrus transition – Part 2: Impact of initial ice crystal number, radiation, stratification, secondary nucleation and layer depth, *Atmos. Chem. Phys.*, 10, 2037–2051, doi:10.5194/acp-10-2037-2010, 2010b.
- 1245 Unterstrasser, S. and Sölch, I.: Study of contrail microphysics in the vortex phase with a Lagrangian particle tracking model, *Atmos. Chem. Phys.*, 10, 10003–10015, doi:10.5194/acp-10-10003-2010, 2010.
- Unterstrasser, S., Gierens, K., and Spichtinger, P.: The evolution of contrail microphysics in the vortex phase, *Meteorol. Z.*, 17, 145–156, doi:10.1127/0941-2948/2008/0273, 2008.
- Unterstrasser, S., Paoli, R., Sölch, I., Kühnlein, C., and Gerz, T.: Dimension of aircraft exhaust plumes at cruise conditions: effect of wake vortices, *Atmos. Chem. Phys.*, 14, 2713–2733, doi:10.5194/acp-14-2713-2014, 2014.
- Van De Hulst, H. C.: *Light scattering by small particles*, Courier Dover Publications, 1957.
- Voigt, C., Schumann, U., Jeßberger, P., Jurkat, T., Petzold, A., Gayet, J., Krämer, M., Thornberry, T., and Fahey, W.: Extinction and optical depth of contrails, *Geophys. Res. Lett.*, 38, L111806, doi:10.1029/2011GL047189, 2011.
- Widnall, S. E., Bliss, D., and Zalay, A.: Theoretical and Experimental Study of The Stability of a Vortex Pair, in: *Aircraft Wake Turbulence and Its Detection*, pp. 305–338, Plenum Press, Seattle, Washington, 1970.
- Widnall, S. E., Bliss, D. B., and Tsai, C.-Y.: The instability of short waves on a vortex ring, *J. Fluid Mech.*, 66, 35–47, doi:10.1017/S0022112074000048, 1974.