



**Hygroscopic
properties of NaCl
and NaNO₃ mixture
particles**

D. Gupta et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Hygroscopic properties of NaCl and NaNO₃ mixture particles as reacted inorganic sea-salt aerosol surrogates

D. Gupta, H. Kim, G. Park, X. Li, H.-J. Eom, and C.-U. Ro

Department of Chemistry, Inha University, Incheon, 402-751, Korea

Received: 24 November 2014 – Accepted: 29 November 2014 – Published:
23 December 2014

Correspondence to: C.-U. Ro (curo@inha.ac.kr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

NaCl in fresh sea-salt aerosol (SSA) particles can partially or fully react with atmospheric NO_x/HNO_3 , so internally mixed NaCl and NaNO_3 aerosol particles can co-exist over a wide range of mixing ratios. Laboratory-generated, micrometer-sized NaCl and NaNO_3 mixture particles at ten mixing ratios (mole fractions of NaCl (X_{NaCl}) = 0.1 to 0.9) were examined systematically to observe their hygroscopic behavior, derive experimental phase diagrams for deliquescence and efflorescence, and understand the efflorescence mechanism. During the humidifying process, aerosol particles with the eutonic composition ($X_{\text{NaCl}} = 0.38$) showed only one phase transition at their mutual deliquescence relative humidity (MDRH) of $67.9(\pm 0.5)\%$. On the other hand, particles with other mixing ratios showed two distinct deliquescence transitions, i.e., the eutonic component dissolved at MDRH and the remainder in the solid phase dissolved completely at their DRHs depending on the mixing ratios, resulting in a phase diagram composed of four different phases, as predicted thermodynamically. During the dehydration process, NaCl-rich particles ($X_{\text{NaCl}} > 0.38$) showed two-stage efflorescence transitions: the first stage was purely driven by the homogeneous nucleation of NaCl and the second stage at the mutual efflorescence RH (MERH) of the eutonic components, with values in the range of 30.0–35.5%. Interestingly, aerosol particles with the eutonic composition ($X_{\text{NaCl}} = 0.38$) also showed two-stage efflorescence with NaCl crystallizing first followed by heterogeneous nucleation of the remaining NaNO_3 on the NaCl seeds. NaNO_3 -rich particles ($X_{\text{NaCl}} \leq 0.3$) underwent single-stage efflorescence transitions at ERHs progressively lower than the MERH, because of the homogeneous nucleation of NaCl and the almost simultaneous heterogeneous nucleation of NaNO_3 on the NaCl seeds. SEM/EDX elemental mapping indicated that the effloresced NaCl- NaNO_3 particles at all mixing ratios were composed of a homogeneously crystallized NaCl moiety in the center, surrounded either by the eutonic component (for $X_{\text{NaCl}} > 0.38$) or NaNO_3 (for $X_{\text{NaCl}} \leq 0.38$). During the humidifying or dehydration process, the amount of eutonic composed part drives particle/droplet growth or shrinkage at the MDRH or MERH

Hygroscopic properties of NaCl and NaNO_3 mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(second ERH), respectively, and the amount of remnant pure salts (NaCl or NaNO₃ in NaCl- or NaNO₃-rich particles, respectively) drives the second DRHs or first ERHs, respectively. Therefore, their behavior can be a precursor to the optical properties and direct radiative forcing for these atmospherically relevant mixture particles representing the coarse, reacted inorganic SSAs. In addition, the NaCl-NaNO₃ mixture aerosol particles can maintain an aqueous phase over a wider RH range than the genuine SSA surrogate (i.e., pure NaCl particles), making their heterogeneous chemistry more probable.

1 Introduction

Atmospheric aerosols play important roles in global climate change, directly by scattering or absorbing incoming solar radiation and indirectly by serving as cloud condensation nuclei (Pandis et al., 1995; Satheesh and Moorthy, 2005). The radiative effects depend on the chemical composition and sizes of the atmospheric aerosol particles. The optical properties and the chemical reactivity of the atmospheric aerosols also depend on their mixing states and different aerosol phases (Martin, 2000). Studies of the hygroscopic properties of inorganic salt particles as aerosol surrogates can provide important insights to several of these important aerosol properties, such as (i) alteration of aerodynamic properties, (ii) cloud-droplet nucleation efficiency, (iii) optical properties; and (iv) physicochemical changes, through complex heterogeneous chemical reactions with atmospheric gas-phase species (Wang and Martin, 2007; Haywood et al., 2000; ten Brink, 1998; Krueger et al., 2003).

Sea-salt or sea-spray aerosols (SSAs) comprise a large proportion of the atmospheric particulate mass (25–50 %) (Finlayson-Pitts and Pitts, 2000). Thus far, many studies have examined the hygroscopic behavior of both airborne and laboratory-generated SSAs (Tang et al., 1997; Wise et al., 2007, 2009; Prather et al., 2013), but the hygroscopicity of the SSAs is not completely understood (Meskhidze et al., 2013). NaCl in the nascent SSAs can react quickly (within a few minutes to a hours of

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



have been developed to predict the deliquescence behavior or the ionic activity coefficients of two-component aerosol particles (Tang, 1976; Ansari and Pandis, 1999; Clegg et al., 1998; Wexler and Clegg, 2002; Zuend et al., 2008, 2011). The Extended Atmospheric Inorganic Model (E-AIM) predicts the physical state and chemical compositions of aerosols containing several atmospherically relevant inorganic ionic species and/or organic species (<http://www.aim.env.uea.ac.uk/aim/aim.php>). The Aerosol Inorganic–Organic Mixtures Functional group Activity Co-efficient (AIOMFAC) model allows calculations of the activity coefficients in organic and/or inorganic mixtures from simple binary solutions to complex multicomponent systems (<http://www.aiomfac.caltech.edu>). On the other hand, there have been few systematic, experimental hygroscopic studies to support the theoretical models for mixed salt particles.

During the dehydration process, where the RH is decreased from high to low, the concentration of single salts in the aqueous droplets becomes dense and the inorganic single salts can be finally crystallized at their efflorescence RH (ERH). The ERH is sometimes significantly lower than the DRH. For example, pure NaCl particles have a DRH of ~ 75 % and ERH of ~ 45–47 % (Martin, 2000). From a thermodynamic point of view, or as observed in bulk ternary systems, aqueous droplets with double salts should show step-wise efflorescence transitions: a component in the aqueous droplets precipitates first at their ERH and the remnant aqueous phase of the eutonic composition effloresces at their mutual ERH (MERH), which should be lower than either ERHs of the pure salts. Therefore, effloresced mixed particles may form a heterogeneous, core-shell crystal structure owing to the step-wise crystallization process (Ge et al., 1996). On the other hand, aqueous droplets with a eutonic composition are expected to crystallize simultaneously, resulting in a homogeneous crystal structure. Efflorescence, however, is a kinetic or rate-driven process that requires a sufficient activation energy to overcome the kinetic barrier (Martin, 2000). This kinetic or critical-nucleation barrier in turn depends on a range of factors, such as the mixing states of the chemical components, micro-physical states, supersaturation levels, vapor pressure, interfacial tension, viscosity, inter-ionic forces, and solute-water and solute-solute interactions

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Cohen et al., 1987b). Therefore, the ERHs of single or multi-component salts are difficult to predict theoretically (Seinfeld and Pandis, 2006). A theoretical model for the efflorescence behavior of the NaCl-Na₂SO₄ mixed system was reported, where the efflorescence was considered to be driven primarily by the homogeneous nucleation of the more supersaturated salt, resulting in a sufficiently large seed for the subsequent heterogeneous nucleation of the other salt (Gao et al., 2007). However, there is no general model that covers the efflorescence of multi-component particles because it depends on many complicated parameters and varies with the salt characteristics. Therefore, the best way to understand the efflorescence behavior of aerosols is through experimental measurements (Seinfeld and Pandis, 2006). For example, a recent experimental study of the two component NaCl-KCl mixture particles showed that during the dehydration process, the aqueous droplets of various mixing ratios underwent single step efflorescence (Li et al., 2014). Based on the experimentally-obtained efflorescence phase diagram and X-ray elemental maps of the effloresced NaCl-KCl mixture particles at various mixing ratios, it was suggested that the more supersaturated salt nucleated homogeneously to crystallize in the center and the other salt underwent heterogeneous crystallization on the former almost simultaneously in the time-scale of the measurements. Full phase diagrams, covering the entire range of mixing ratios, are needed to fully understand the hygroscopic behavior of multi-component aerosol particles (Martin, 2000).

Up until now, there have been only a few studies on the hygroscopic properties of mixed NaCl-NaNO₃ aerosol system. Tang and Munkelwitz (1994a) examined the temperature dependent deliquescence behavior of equi-molar mixed NaCl-NaNO₃ particles using a single-particle levitation technique. Ge et al. (1998) also studied the deliquescence behavior of mixed NaCl-NaNO₃ particles with a NaCl mole fraction of 0.2, 0.378 (eutonic), and 0.8 using rapid single-particle mass spectrometry (RSMS), where only MDRH was measured experimentally and it was claimed that the second-stage DRHs agreed with the thermodynamic predictions of AIM. Although the crystallization process was not studied experimentally, a core-shell type of heterogeneous morphol-

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ogy in the NaCl-NaNO₃ particles was proposed based on the two-step deliquescence phase transitions observed during the humidifying process (Ge et al., 1998; Hoffmann et al., 2004) and by measuring the secondary electron yields from the nebulized mixture particles (Ziemann and McMurry, 1997). On the other hand, as observed for the NaCl-KCl mixture particles, particles that exhibit two-step deliquescence phase transitions do not always have a core-shell type (Li et al., 2014). Moreover, it was reported that pure NaNO₃ droplets do not crystallize easily during the dehydration process, and at very low RHs they appear to exist in an amorphous form (Hoffmann et al., 2004; Gibson et al., 2006; Kim et al., 2012). Therefore, the mixing states in NaCl-NaNO₃ particles can be understood only when the efflorescence phenomena for these mixture particles are elucidated. In this study, the hygroscopic properties and microstructure of mixed NaCl-NaNO₃ particles at various mixing ratios were examined extensively by optical microscopy and scanning electron microscopy/energy dispersive X-ray spectroscopy (SEM/EDX). The phase transitions of the mixed NaCl-NaNO₃ aerosol particles were observed by monitoring the size changes of the particles on the optical images as a function of the RH. SEM/EDX mapping was used to investigate the compositional distribution in the effloresced particles. This paper describes the full hygroscopic behavior of the NaCl-NaNO₃ binary aerosol particles as reacted SSA surrogates at ten different mixing ratios for the first time.

2 Experimental section

2.1 Preparation of mixed NaCl-NaNO₃ particles

Mixed NaCl-NaNO₃ particles were generated by the nebulization of mixed aqueous solutions. Pure solutions (1.0 M) of NaCl and NaNO₃ (NaCl, > 99.9% purity, Aldrich; NaNO₃, 99.9% purity, Aldrich) were prepared and the desired solution was made by mixing the two solutions volumetrically. A single jet atomizer (HCT4810) was used to generate aerosol particles to be deposited on TEM grids (200 mesh Cu coated with For-

mvar stabilized with carbon, Ted Pella, Inc.), which behave as hydrophobic substrates (Eom et al., 2014). The aqueous aerosol particles were dried by passing through a silica packed diffusion dryer (HCT4920) with a residence time of ~ 2 s. The size of the dry particles ranged from 1 to 10 μm .

In this study, NaCl-NaNO₃ particles with 10 different mixing ratios were investigated; i.e., 9 compositions with NaCl mole fractions of 0.1–0.9 ($X_{\text{NaCl}} = 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8,$ and 0.9 , where X_{NaCl} represents the mole fraction of NaCl.) and a eutonic composition ($X_{\text{NaCl}} = 0.378 \approx 0.38$, which was calculated from the ionic activity products predicted by the AIOMFAC model). Based on the mixing ratio of two salts, the mixed NaCl-NaNO₃ particles were divided into three categories: (1) eutonic particles ($X_{\text{NaCl}} = 0.38$), (2) NaCl-rich particles containing larger NaCl fraction than the eutonic composition ($X_{\text{NaCl}} > 0.38$), and (3) NaNO₃-rich particles containing larger NaNO₃ fraction than the eutonic composition ($X_{\text{NaCl}} < 0.38$).

2.2 Hygroscopic property measurement

The hygroscopic properties of the particles were investigated using a “see-through” inertia impactor apparatus equipped with an optical microscope. The experimental set-up is described in detail elsewhere (Ahn et al., 2010). Briefly, the apparatus is composed of three parts: (A) see-through impactor, (B) optical microscope, and (C) humidity controlling system. A TEM grid on which aerosol particles were deposited was mounted on the impaction plate in the see-through impactor. The RH inside the impactor was controlled by mixing dry and wet (saturated with water vapor) N₂ gases. The wet N₂ gas was obtained by bubbling through deionized water reservoirs. The flow rates of the dry and wet N₂ gases were controlled by mass flow controllers to obtain the desired RH in the range of ~ 3 –93 %, which was monitored using a digital hygrometer (Testo 645). The digital hygrometer was calibrated using a dew-point hygrometer (M2 Plus-RH, GE), providing RH readings with ± 0.5 % reproducibility. To achieve a steady state for condensing or evaporating water, each humidity condition was sustained for at least two minutes. The particles on the impaction plate were observed through a nozzle

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



throat using an optical microscope (Olympus, BX51M). Images of the particles were recorded continuously using a digital camera (Canon EOS 5D, full frame, Canon EF f/3.5 L macro USM lens) during the humidifying (by increasing RH from ~ 3 to $\sim 93\%$) and dehydration (by decreasing RH from ~ 93 to $\sim 3\%$) experiments. The image size was 4368×2912 pixels and the image recording condition was set to ISO200. The exposure time was 0.4 s, and the DOF was F/3.5. All hygroscopic experiments were conducted at room temperature ($T = 22 \pm 1^\circ\text{C}$).

The change in particle size with the variation of RH was monitored by measuring the particle areas in the optical images. The particle images were processed using image analysis software (Matrox, Inspector v9.0). The size of the imaging pixel was calibrated using $10\ \mu\text{m}$ Olympus scale bars. Particles with $D_p > 0.5\ \mu\text{m}$ could be analyzed using the present system.

2.3 SEM/EDX measurement

After the hygroscopicity measurements of the individual particles, SEM/EDX was performed for the effloresced particles to determine the morphology and spatial distribution of the chemical elements. The measurements were carried out using a Jeol JSM-6390 SEM equipped with an Oxford Link SATW ultrathin window EDX detector. The resolution of the detector was 133 eV for the Mn K_α X-rays. The X-ray spectra and elemental maps were recorded under the control of Oxford INCA Energy software. A 10 kV accelerating voltage and 0.2 nA beam current was used and the typical measuring times were 10 min. for elemental mapping.

3 Results and discussion

3.1 Hygroscopic behavior of pure NaCl and NaNO₃ particles

Aerosol particles generated from a pure NaCl aqueous solution showed typical hysteresis curves with DRH = $75.5 (\pm 0.5)\%$ and ERH = $47.6\text{--}46.3\%$, and these values were

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sition. For example, a droplet of $X_{\text{NaCl}} = 0.2$ decreased continuously in size with decreasing RH until it showed an efflorescence transition at $\text{RH} = 29.4\text{--}29.2\%$ (Fig. 2b). Similar to the NaCl-rich case, the NaNO₃-rich particles would be expected to exhibit two-stage transitions during the dehydration process, i.e., the first transition accompanying the precipitation of solid NaNO₃, and the second one due to the efflorescence of the eutonic component. However, all the NaNO₃-rich particles with $X_{\text{NaCl}} = 0.1, 0.2,$ and 0.3 showed single-stage efflorescence transitions during the dehydration process (Figs. 2a and b, 4), suggesting that all the components in the NaNO₃-rich particles crystallized (almost) simultaneously. The supersaturated NaNO₃ in the droplets did not appear to crystallize until the NaCl crystallized and acted as heterogeneous nuclei for the almost simultaneous solidification of NaNO₃. To confirm this assumption, particles with small fractions of NaCl, such as particles with $X_{\text{NaCl}} = 0.01, 0.03,$ and $0.05,$ were investigated. The particles with $X_{\text{NaCl}} = 0.05$ showed two-stage and single-stage transitions during the humidifying and dehydration processes, respectively, which is similar to that observed with the NaNO₃-rich particles with a composition of $X_{\text{NaCl}} \geq 0.1$. On the other hand, the particles with $X_{\text{NaCl}} = 0.01$ and 0.03 underwent continuous hygroscopic growth and shrinkage without phase transitions during the humidifying and dehydration processes, respectively, which is similar to that of aerosol particles generated from an aqueous single-component NaNO₃ solution. Pure nebulized NaNO₃ did not show clear efflorescence transitions apparently because of its amorphous nature (Hoffman et al., 2004; Gibson et al., 2006; Kim et al., 2012). Therefore, a sufficient quantity of heterogeneous nuclei (NaCl in this case) is needed to induce the crystallization of NaNO₃. A similar observation was reported for the crystallization of NH₄NO₃ and NH₄HSO₄ aerosol particles (Schlenker and Martin, 2005). Both the pure one-component salts did not effloresce, even at $\text{RH} = 1\%$ and their crystallization could be promoted with the addition of some fraction of inclusions that could serve as good heterogeneous nuclei.

3.3 Deliquescence phase diagram of mixed NaCl-NaNO₃ particles

Figure 3 presents the measured MDRHs (the first DRHs) and second DRHs of the NaCl-NaNO₃ mixture particles with different mole fractions along with the measured DRHs of the pure NaCl and NaNO₃ particles. As shown in Fig. 3, a clearly demarked phase diagram depicting their deliquescence behavior was obtained experimentally; i.e.,

1. NaCl(s) + NaNO₃(s) phase in Fig. 3: both NaCl and NaNO₃ are mixed as solids below the MDRH at all mole fractions;
2. NaCl(s) + eutonic(aq) phase: a mixed phase of solid NaCl and aqueous eutonic components between the MDRH and second DRHs for $X_{\text{NaCl}} > 0.38$;
3. NaNO₃(s) + eutonic(aq) phase: a mixed phase of solid NaNO₃ and aqueous eutonic components between the MDRH and second DRHs for $X_{\text{NaCl}} < 0.38$; and
4. NaCl(aq) + NaNO₃(aq) phase: both NaCl and NaNO₃ are mixed in the aqueous phase above the second DRHs at all mole fractions.

This MDRH and second DRHs obtained experimentally agrees well with the values calculated from the ionic activity products of constituents predicted by the AIOMFAC model, as shown in Fig. 3 (dotted line for MDRH and dash-dotted curve for the second DRHs). Although many theoretical models have been developed to predict the hygroscopic behavior of a mixed sodium chloride and nitrate system (Tang, 1976; Ansari and Pendis, 1999; Clegg et al., 1998; Wexler and Clegg, 2002), only a few experimental results have been reported. Tang and Munkelwitz (1994a) reported an MDRH = 68.0 (±0.4) % at 25 °C. Ge et al. (1998) examined the deliquescence behavior of mixed NaCl-NaNO₃ particles with a NaCl mole fraction of 0.2, 0.38(eutonic), and 0.8, using RSMS. They claimed that the DRHs were generally consistent with the AIM thermodynamic model predictions. On the other hand, the measured DRH values were not reported except for the MDRH of 67.0 %, which was lower than the 67.9 % obtained in

the present study, because the RSMS appeared to detect only the start of the transition similar to the present system where mutual deliquescence for $X_{\text{NaCl}} = 0.38$ began at 67.1 % (Fig. 2b), whereas the actual MDRH was at the end of the transition at 67.9 % when the eutonic particle dissolved completely.

All the mixed NaCl-NaNO₃ particles showed the first phase transition at the MDRH regardless of the mixing ratio of the two salts. Thermodynamically, as the phase transition of mixed-salts is governed by the water activity at the eutonic point, the MDRH of the mixed-salt particles is independent of the initial composition of the mixture. Other inorganic mixed particles, such as NaCl-KCl, Na₂SO₄-NaNO₃ and NH₄Cl-NaCl particles, also exhibited a mutual deliquescence transition at RHs, which were independent of the initial dry-salt compositions, and the phase diagram followed the typical pattern for these two-component inorganic salt + water ternary systems (Tang and Munkelwitz, 1994a; Li et al., 2014; Kelly et al., 2008). For the NaCl-rich particles of $X_{\text{NaCl}} > 0.38$, which contain more NaCl than the eutonic composition, the second DRH value approached the DRH of the pure NaCl salt as the NaCl concentration was increased. For the NaNO₃-rich particles of $X_{\text{NaCl}} < 0.38$, the DRH approached that of pure NaNO₃ as the NaNO₃ mole fraction was increased (Fig. 3). This suggests that the second-stage deliquescence is purely driven by the solid salt remaining after the first deliquescence of the eutonic composition.

3.4 Efflorescence phase diagram of mixed NaCl-NaNO₃ particles

Figure 4 shows the measured ERHs and MERHs for the mixed NaCl-NaNO₃ particles with various mixing ratios as a function of the NaCl mole fraction. Unlike the deliquescence phase diagram, which showed four systematic phases, the efflorescence phase diagram is composed of three distinct phases; i.e.,

1. NaCl(aq) + NaNO₃(aq) phase: both NaCl and NaNO₃ are mixed in the aqueous phase above the first ERHs at all mixing ratios;

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2. NaCl(s) + eutonic(aq) phase: a mixed phase of solid NaCl and aqueous eutonic components between the first ERH and second ERH (MERH) for $X_{\text{NaCl}} \geq 0.38$; and

3. NaCl(s) + NaNO₃(s) phase: both NaCl and NaNO₃ are mixed as solids below the second ERH (MERH) for $X_{\text{NaCl}} \geq 0.38$ and below the first ERHs for $X_{\text{NaCl}} < 0.38$.

This experimental phase diagram for efflorescence is reported for the first time, for which no theoretical predictions or other experimental reports exist to the best of the authors' knowledge.

The ERH of NaCl-rich droplets ($X_{\text{NaCl}} > 0.38$) shifted toward the pure NaCl limit (RH = 47.6–46.3 %) when the NaCl content was increased (see Fig. 4). The measured MERH was observed at a relatively wide range of RH = 30.0–35.5 %. The droplets with the eutonic composition ($X_{\text{NaCl}} = 0.38$) showed the first and second ERHs over a range of 37.7–35.7 and 35.4–33.4 %, respectively. The NaNO₃-rich droplets ($X_{\text{NaCl}} < 0.38$) had only one stage efflorescence transition and the decreasing trend of their ERHs as the NaCl mole fraction was decreased clearly follows that for the NaCl-rich and eutonic droplets (Fig. 4). This suggests that the first efflorescence of the NaCl-NaNO₃ mixture droplets at all mixing ratios is driven by the homogeneous nucleation of NaCl. For the NaCl-rich droplets, crystallized NaCl acts as a seed for further precipitation of the remaining metastable eutonic aqueous part. For the eutonic droplets, NaCl is crystallized homogeneously at the first ERH and the remaining NaNO₃ solidify heterogeneously on the NaCl seeds at the second ERH. For NaNO₃-rich droplets, due to the low NaCl content, the homogeneous nucleation rate of NaCl decreases with decreasing NaCl mole fraction, and the NaNO₃ appears to undergo almost simultaneous heterogeneous crystallization (precipitation) on the NaCl seeds under the time scale of measurements (i.e., within 2 min of the equilibrating time for recording images during the first observed efflorescence transition).

3.5 Spatial distribution of effloresced NaCl-NaNO₃ solid particles

To examine the morphology and spatial distribution of the chemical components in NaCl-NaNO₃ particles at various mixing ratios, SEM/EDX were performed for the effloresced solid particles formed after the humidifying and dehydration cycles. Figure 5a shows the secondary electron image (SEI) and elemental X-ray mapping images of a NaCl-rich ($X_{\text{NaCl}} = 0.8$) particle. The elemental X-ray maps suggest that Cl (from NaCl) is concentrated in the central part, whereas O (from NaNO₃) is more concentrated at the edges. This suggests that NaCl nucleates homogeneously to crystallize in the center at the first ERH, whereas NaCl and NaNO₃ from the eutonic phase crystallized on these central NaCl seeds and precipitated on the edges at the second ERH (MERH), respectively.

For a typical particle with a eutonic composition ($X_{\text{NaCl}} = 0.38$), the elemental map of Cl suggests that NaCl is again more concentrated at the central part like NaCl-rich particles and O from NaNO₃ is concentrated around this NaCl core (Fig. 5b), suggesting that NaCl is nucleated homogeneously to crystallize in the center at the first ERH, whereas the remaining NaNO₃ precipitated at the second ERH. The thermodynamically-expected, homogeneously-mixed eutonic particles were not observed, and the spatial distribution was rather ruled by the two stage efflorescence transitions.

In the case of NaNO₃-rich particles, as shown in Fig. 5c, the Cl component was localized in the core region, and Na and O are distributed over the entire particle. This suggests that even for NaNO₃-rich particles, NaCl is nucleated homogeneously to crystallize in the core at the single observed ERHs, whereas the NaNO₃ precipitate simultaneously on this core within the time scale of the measurements. Although the NaNO₃-rich droplet was supersaturated with NaNO₃ with decreasing RH during the dehydration process, NaNO₃ could not crystallize easily even at the MERH of the NaCl-rich mixtures. Therefore, RHs lower than MERH (see Fig. 4 for the lower ERHs in case of $X_{\text{NaCl}} \leq 0.3$ than MERH) were necessary for the homogeneous crystallization

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of NaCl followed by the almost simultaneous, induced heterogeneous crystallization of NaNO₃.

Up until now, the binary inorganic salt aerosols, except the eutonic composition, are generally believed to form a core-shell type heterogeneous solid of a pure salt core surrounded by the eutonic component. The formation of the core-shell type had been reported for a range of binary mixed aerosol particles, such as NaCl-KCl, KCl-KI, (NH₄)₂SO₄-NH₄NO₃ system (Ge et al., 1996). Based on the secondary electron yield measurements of the NaCl-NaNO₃ mixture particles, it was claimed that individual particles exist in core-shell form with the richer salt (NaCl in NaCl-rich or NaNO₃ in NaNO₃-rich), and the eutonic components occupy the core and shell, respectively (Ziemann and McMurry, 1997). However, the study was conducted on particles nebulized from aqueous solutions that had not gone through the proper dehydration process. Hoffmann et al. (2004) predicted a core-shell type structure based on their observations of NaCl inclusions in the partially deliquesced particles only. On the other hand, a previous report on NaCl-KCl mixture particles showed that NaCl and KCl were crystallized as separate phases and not necessarily in the core-shell configuration (Li et al., 2014). In addition, a eutonic solid shell in the dry particle was found to be unnecessary for the exhibition of the two stage deliquescence transitions during the humidifying process at all mixing ratios. The efflorescence phase diagram (Fig. 4) and the X-ray maps (Fig. 5) showed that effloresced NaCl-NaNO₃ particles of all mixing ratios had NaCl crystallized homogeneously in the center, surrounded either by the eutonic component for $X_{\text{NaCl}} > 0.38$ or NaNO₃ for $X_{\text{NaCl}} \leq 0.38$. These micro-structures and spatial distributions of chemical components have obvious atmospheric implications (Ziemann and McMurry, 1997).

4 Atmospheric implications

The particle/droplet size variations with RH for the different mixing states (Fig. 2), the four and three distinct phases observed during the humidifying (Fig. 3) and dehydra-

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion (Fig. 4) processes, respectively, have important atmospheric implications in terms of radiative forcing (Baynard et al., 2006; Ma et al., 2008), cloud nucleation efficiency (Petters et al., 2007; Wex et al., 2008), atmospheric chemistry (Ault et al., 2013a; Ryder et al., 2014; Wang and Laskin, 2014), and gas adsorption/desorption (gas-particle partitioning) (Woods et al., 2012). Some of these implications related to the observed experimental results are discussed.

4.1 Particle size change at different phase transitions

In this study, the measured NaCl-NaNO₃ mixture particles, 1–10 μm in size and their size variations according to the RH change, are atmospherically relevant because inorganics are dominant in the super-micron or coarse size fraction of SSAs (Keene et al., 2007; Ault et al., 2013b; Prather et al., 2013). In Fig. 6a, 2-D diameter ratios ($d/d_x = \sqrt{A/A_x}$, where x is the particle state at RHs before the transitions), which represent the size increase due to mutual deliquescence of the eutonic part at MDRH (compared to the rearranged solid particle) as well as to deliquescence of the remnant salts at the second DRHs (compared to the partially deliquesced particle at MDRH), were plotted as a function of the NaCl mole fractions. As shown in Figs. 2 and 6a, the particle size variations during the humidifying process depends on the mixing ratio of the two-component NaCl-NaNO₃ particles as the extent of the increase in particle size due to the water uptake is associated with the chemical composition of the particles. For single-component NaCl and NaNO₃ particles, their 2-D diameters increase ~ 2.0 and ~ 1.5 fold, respectively, when they deliquesced at their DRHs (Fig. 6a). The difference was attributed to the different solubility of the salts. Because NaCl is less soluble than NaNO₃ (their solubility is 36.0 g/100 g and 91.2 g/100 g, respectively) (Lide, 2002), NaCl requires more water than NaNO₃ to form a saturated solution, resulting in a larger increase in size when solid NaCl particles deliquesce. Therefore, for the mixed NaCl-NaNO₃ particles, the extent of the particle size increases at the MDRH or the second DRH is associated with the mixing ratios of the two salts. At the MDRH, where only the

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



eutonic component deliquesces, the extent of the size increase depends on the fraction of the eutonic component in the particles. Therefore, at the MDRH, particles with a eutonic composition ($X_{\text{NaCl}} = 0.38$) show the largest increase in size ($d/d_x \approx 1.7$ as shown in Fig. 6a). For NaCl-rich particles ($X_{\text{NaCl}} > 0.38$), the size increase at the MDRH decreases with increasing NaCl fraction (Figs. 2d–f and 6a) because the eutonic fraction, which depends on the NaNO₃ content in the particle, decreases with increasing NaCl mole fraction. In contrast, for NaNO₃-rich particles ($X_{\text{NaCl}} < 0.38$), the size at the MDRH increases with increasing NaCl fraction (Figs. 2a and b, 6a), because the eutonic fraction, which depends on the NaCl content, increases with increasing NaCl mole fraction. The particle size change at the RHs between MDRH and the second DRH during the humidifying process depends on the residual solid components after the mutual deliquescence. For NaCl-rich particles, the particles with a larger NaCl fraction grow more at the RHs between MDRH and their second DRHs (Figs. 2d–f and 6a) because the partially deliquesced particles have a larger residual NaCl solid fraction. In contrast, for the NaNO₃-rich particles, the particles with the larger NaCl fraction grow less (Figs. 2a and b, 6a), because they have a smaller residual NaCl solid fraction.

At the first ERH, where efflorescence occurs by the homogeneous nucleation of NaCl, the NaCl-rich and eutonic droplets ($X_{\text{NaCl}} \geq 0.38$) show a larger decrease in size for droplets of a larger NaCl fraction (Fig. 6b). For the NaNO₃-rich droplets, the simultaneous heterogeneous precipitation of the NaNO₃ on the homogeneously-crystallized NaCl seeds occurs at their ERH, and the droplets of higher NaNO₃ content show a larger decrease in size. The decrease in size at the second efflorescence transition for aerosol particles of $X_{\text{NaCl}} \geq 0.38$ should be governed by the eutonic content in the aerosol particles; i.e., the more eutonic content, the greater the size decrease (Fig. 6b).

Aerosol particles with mixing ratios of $X_{\text{NaCl}} \geq 0.38$ did not show a noticeable change in the 2-D area when the RH was decreased from their first ERH to the start of the second ERH (MERH) (see Figs. 1g–i and 2c–f). At this RH range, the aerosol particles are expected to gradually decrease in size due to the evaporation of water in the eu-

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tonic solution. As discussed above, however, less soluble NaCl solid particles require more water than NaNO₃ to form a saturated solution, resulting in a larger increase in size when NaCl particles deliquesce (Fig. 6a). On the other hand, when aqueous NaCl droplets crystallize, more water in the droplets evaporates than when NaNO₃ is present in the droplets. For NaCl-rich droplets, because a large portion of water in the droplet might be removed when NaCl crystallizes at their first ERH, only a small amount of water remains in the eutonic solution, which evaporates gradually until complete efflorescence occurs, even though shrinkage was not observed well on their optical images. However, in this RH range, the particle boundary appeared to be clearer when it became more concentrated with decreasing RH. Indeed, the change was clear when viewed directly through the optical microscope (unlike the digital images) during the dehydration measurements. The size variations of aerosol particles with different mixing ratios according to the RH change can help predict their aerodynamic properties and hence their residence time in ambient air. These variations, however, also depend on the original size of the particles, possibly more for sub-micron particles (Hu et al., 2010).

4.2 Hygroscopic growth and cloud droplet nucleation

Cloud droplet nucleation can begin when the air is supersaturated with water vapor (i.e., RH is above 100 %) and the hygroscopic growth at high RHs are correlated with the cloud condensation nuclei (CCN) activity (Petters et al., 2007; Wex et al., 2008). Under these experimental conditions, the highest working RH was ~ 93 %. On the other hand, as the DRHs of pure NaCl and NaNO₃ are 75.5 (±0.5) % and 74.0 (±0.5) %, respectively, the hygroscopic growth begins at less than ~ 76 % RH for particles of all mixing ratios. In Fig. 7, the growth factors in terms of the 2-D diameter ratios (d_{95}/d_0 , where $d_{95} \equiv$ the diameter at RH = 95 %, determined by extrapolation, and $d_0 \equiv$ dry diameter at ~ 3 % RH) are plotted as a function of the mole fraction of NaCl for the NaCl-NaNO₃ mixture particles to indicate the cloud droplet nucleation efficiency of these particles at various mixing ratios. NaCl, eutonic, and NaNO₃ particles, which have sin-

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gle deliquescence transitions (one DRH), appear to show relatively higher hygroscopic growth, in the order, NaCl > eutonic > NaNO₃, than particles with other mixing ratios, having two-stage deliquescence (MDRH and the second DRH). The hygroscopic growth first decreases with decreasing eutonic content for NaCl-rich particles with 0.38 < $X_{\text{NaCl}} \leq 0.6$ and then begins to increase with increasing NaCl content for particles with $0.6 < X_{\text{NaCl}} \leq 1.0$. Similar behavior was observed for NaNO₃-rich particles. Therefore, for NaCl-NaNO₃ mixture particles, the CCN efficiency should also follow the same trend. This can serve as a precursor to the hygroscopic growth and CCN activity of more complex, reacted SSAs containing other salts and organics (King et al., 2012). However, an estimation of the full CCN activity and cloud droplet number concentration require further experiments on a wider dry particle size range under both sub- and super-saturated conditions (Fuentes et al., 2011; Mochida et al., 2011).

4.3 Atmospheric chemistry

The knowledge of mixing states, phases, and spatial distribution of chemicals in the NaCl-NaNO₃ mixture particles, as reacted SSA surrogates, at various RHs is expected to help better understand the complexity of real ambient SSAs, their hygroscopic properties, aqueous phase chemistry, etc. For SSAs comprising NaCl-NaNO₃ mixture particles generated from a partial or full reaction with NO_x/HNO₃, their aqueous surface regions are crucial for atmospheric heterogeneous chemistry because the aqueous phases of NaCl, NaNO₃, or the eutonic composed part at different RHs are expected to be available for further reactions with gas phase species, such as N₂O₅ (Ault et al., 2013a; Ryder et al., 2014) or organics (Wang and Laskin, 2014), and/or for facile gas-particle partitioning (Woods et al., 2012).

SSAs are generated by sea-spray action, so that they are initially aqueous droplets when they become airborne. Although real ambient SSAs are complex mixtures containing NaCl, other inorganic salts, and organics (O'Dowd and de Leeuw, 2007; Keene et al., 2007; Prather et al., 2013; Beardsley et al., 2013; Ault et al., 2013b), NaCl particles have been studied as genuine SSA surrogates in many laboratories (Martin,

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2000; Krieger et al., 2012). Based on the experimental phase diagrams of deliquescence and efflorescence for the NaCl-NaNO₃ system (Figs. 3 and 4), once airborne as droplets, the genuine SSA surrogate would remain in the aqueous phase unless the droplets experience an ambient RH below their ERH of 47.6–46.3 % (pure NaCl).

5 Their aqueous phase can facilitate heterogeneous reactions with gaseous species, such as NO_x/HNO₃, to become NaCl-NaNO₃ mixture droplets. When mixture droplets are formed, their full efflorescence occurs at lower RHs (i.e., at MERH of ~ 33.1 % for the mixture droplets of $X_{\text{NaCl}} \geq 0.38$ or at ERH of ~ 35–21 % for those of $X_{\text{NaCl}} < 0.38$; see Fig. 4) than ERH for pure NaCl, so that their chance for a further gas-particle interaction would be larger with less probability to fully effloresce. Even when the mixed particles become solids due to the full efflorescence below either their MDRH or ERH, they can deliquesce at a lower DRH (i.e., the NaCl-NaNO₃ mixed particles with various mixing ratios partially deliquesce at a MDRH of ~ 67.9 %, whereas pure NaCl deliquesces at a DRH of ~ 75.5 %; see Fig. 3). In other words, NaCl-NaNO₃ mixture aerosol particles can maintain an aqueous phase over a wider RH range than the genuine SSA surrogate, making their heterogeneous chemistry more probable. Therefore, NaCl-rich particles with higher Cl⁻/NO₃⁻ can show higher uptake of N₂O₅ at ambient RHs (usually > 60 %) in the marine boundary layer (Ryder et al., 2014). On the other hand, a high uptake of HNO₃ due to surface adsorbed moisture on crystalline NaCl particles has also been reported (Saul et al., 2006; Liu et al., 2007). The ionic activities of aqueous Na⁺/Cl⁻/NO₃⁻ at high to very low water activities during hydration and dehydration, indicating the chemical reactivity of these ions, can be calculated using the AIOMFAC model.

5 Conclusion

25 The hygroscopic behavior and microstructures of mixed NaCl-NaNO₃ particles at various mixing ratios collected on TEM grid substrates were investigated by the use of optical microscopy and SEM/EDX. The DRHs and ERHs of the mixed NaCl-NaNO₃ parti-

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cles in the micrometer size range at room temperature were determined by monitoring the change in the particle area in 2-D optical images with the RH variation of $\sim 3\text{--}93\%$. During the humidifying process, the particles with a eutonic composition ($X_{\text{NaCl}} = 0.38$) exhibited single-stage deliquescence behavior at MDRH = $67.9 (\pm 0.5)\%$. The mixed NaCl-NaNO₃ particles, except for the eutonic composition, showed two-stage phase transitions: the first transition at MDRH, which is independent of the chemical compositions, and the second transition at the DRHs, which depend on the mixing ratios of the two salts. For the NaCl-rich particles ($X_{\text{NaCl}} > 0.38$), the increase in the mole fraction of NaCl shifted the DRH toward the pure NaCl limit (DRH = $75.5 (\pm 0.5)\%$). For NaNO₃-rich particles ($X_{\text{NaCl}} < 0.38$), the increase in the mole fraction of NaNO₃ shifted the DRH toward the pure NaNO₃ limit (DRH = $74.0 (\pm 0.5)\%$). The measured DRH values agreed well with those calculated from the thermodynamic AIOMFAC model.

The dehydration behavior of the mixed NaCl-NaNO₃ particles depend on the mixing ratios of the two salts. NaCl-rich particles ($X_{\text{NaCl}} > 0.38$) showed two-stage efflorescence transitions: the first stage, which is driven purely by the homogeneous nucleation of NaCl, and the second stage at MERH. Interestingly, the eutonic composed particles ($X_{\text{NaCl}} = 0.38$) also showed two stage efflorescence with NaCl crystallizing first followed by the heterogeneous nucleation of NaNO₃ on NaCl seeds. NaNO₃-rich particles ($X_{\text{NaCl}} \leq 0.3$) showed single-stage efflorescence transitions at ERHs progressively lower than MERH due to the slower homogeneous nucleation of NaCl and the simultaneous heterogeneous efflorescence of NaNO₃ on the NaCl seeds. SEM/EDX elemental mapping shows that effloresced NaCl-NaNO₃ particles of all mixing ratios had NaCl crystallized homogeneously in the center, surrounded either by the eutonic component for $X_{\text{NaCl}} > 0.38$ or NaNO₃ for $X_{\text{NaCl}} \leq 0.38$. A pure salt core and eutonic composed solid shell in the dry particle was not necessary for exhibiting the two stage deliquescence transitions in all mixing ratios.

During humidifying or dehydration processes, the amount of the eutonic composed part drives the particle/droplet growth or shrinkage at the MDRH or MERH (second ERH), respectively, and the amount of remnant pure salts (NaCl or NaNO₃ in NaCl-

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



or NaNO₃-rich particles, respectively) does at the second DRHs or first ERHs, respectively. Therefore, their behavior can be a precursor to the optical properties and direct radiative forcing for these atmospherically relevant mixture particles representing the coarse, reacted inorganic SSAs. On the other hand, the hygroscopic growth at a RH of 95 % for the various mixing ratios of these mixture particles can be a precursor to the cloud droplet nucleation efficiency and indirect radiative forcing. In addition, NaCl-NaNO₃ mixture aerosol particles can maintain an aqueous phase over a wider RH range than the genuine SSA surrogate (i.e., pure NaCl particles), making their heterogeneous chemistry more likely.

Acknowledgements. This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012R1A2A1A05026329) and by Metrology Research Center funded by Korea research Institute of Standards and Science (KRISS – 2013 – 13011055).

References

- Ahn, K.-H., Kim, S.-M., Jung, H.-J., Lee, M.-J., Eom, H.-J., Maskey, S., and Ro, C.-U.: Combined use of optical and electron microscopic techniques for the measurement of hygroscopic property, chemical composition, and morphology of individual aerosol particles, *Anal. Chem.*, 82, 7999–8009, 2010.
- Ansari, A. S. and Pandis, S. N.: Prediction of multicomponent inorganic atmospheric aerosol behavior, *Atmos. Environ.*, 33, 745–757, 1999.
- Ault, A. P., Guasco, T. L., Ryder, O. S., Baltrusaitis, J., Cuadra-Rodriguez, L. A., Collins, D. B., Ruppel, M. J., Bertram, T. H., Prather, K. A., and Grassian, V. H.: Inside versus outside: ion redistribution in nitric acid reacted sea spray aerosol particles as determined by single particle analysis, *J. Am. Chem. Soc.*, 135, 14528–14531, doi:10.1021/ja407117x, 2013a.
- Ault, A. P., Moffet, R. C., Baltrusaitis, J., Collins, D. B., Ruppel, M. J., Cuadra-Rodriguez, L. A., Zhao, D., Guasco, T. L., Ebben, C. J., Geiger, F. M., Bertram, T. H., Prather, K. A., and Grassian, V. H.: Size-dependent changes in sea spray aerosol composition and properties with different seawater conditions, *Environ. Sci. Technol.*, 47, 5603–5612, doi:10.1021/es400416g, 2013b.

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ault, A. P., Guasco, T. L., Baltrusaitis, J., Ryder, O. S., Trueblood, J. V., Collins, D. B., Ruppel, M. J., Cuadra-Rodriguez, L. A., Prather, K. A., and Grassian, V. H.: Heterogeneous reactivity of nitric acid with nascent sea spray aerosol: large differences observed between and within individual particles, *J. Phys. Chem. Lett.*, 5, 2493–2500, doi:10.1021/jz5008802, 2014.

Baynard, T., Garland, R. M., Ravishankara, A. R., Tolbert, M. A., and Lovejoy, E. R.: Key factors influencing the relative humidity dependence of aerosol light scattering, *Geophys. Res. Lett.*, 33, L06813, doi:10.1029/2005gl024898, 2006.

Beardsley, R., Jang, M., Ori, B., Im, Y., Delcomyn, C. A., and Witherspoon, N.: Role of sea salt aerosols in the formation of aromatic secondary organic aerosol: yields and hygroscopic properties, *Environ. Chem.*, 10, 167–177, doi:10.1071/EN13016, 2013.

Chang, S.-Y. and Lee, C.-T.: Applying GC-TCD to investigate the hygroscopic characteristics of mixed aerosols, *Atmos. Environ.*, 36, 1521–1530, 2002.

Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: A thermodynamic model of the system $H^+ - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O$ at 298.15 K, *J. Phys. Chem. A*, 102, 2155–2171, 1998.

Cohen, M. D., Flagan, R. C., and Seinfeld, J. H.: Studies of concentrated electrolyte solutions using the electrodynamic balance. 2. Water activities for mixed-electrolyte solutions, *J. Phys. Chem.*, 91, 4575–4582, doi:10.1021/j100301a030, 1987a.

Cohen, M. D., Flagan, R. C., and Seinfeld, J. H.: Studies of concentrated electrolyte solutions using the electrodynamic balance. 3. Solute nucleation, *J. Phys. Chem.*, 91, 4583–4590, doi:10.1021/j100301a031, 1987b.

Eom, H.-J., Gupta, D., Li, X., Jung, H.-J., Kim, H., and Ro, C.-U.: Influence of collecting substrates on the characterization of hygroscopic properties of inorganic aerosol particles, *Anal. Chem.*, 86, 2648–2656, doi:10.1021/ac4042075, 2014.

Finlayson-Pitts, B. J. and Pitts, J. N. J.: *Chemistry of the Upper and Lower Atmosphere Theory, Experiments, and Applications*, Academic Press, San Diego, 2000.

Frey, E. J., Martin, S. T., and Buseck, P. R.: Deliquescence and efflorescence of potassium salts relevant to biomass-burning aerosol particles, *Aerosol Sci. Tech.*, 43, 799–807, 2009.

Frey, E. J., Adachi, K., and Buseck, P. R.: Internally mixed atmospheric aerosol particles: hygroscopic growth and light scattering, *J. Geophys. Res.*, 115, D19210, doi:10.1029/2009JD013558, 2010.

Fuentes, E., Coe, H., Green, D., and McFiggans, G.: On the impacts of phytoplankton-derived organic matter on the properties of the primary marine aerosol – Part 2: Compo-

**Hygroscopic
properties of NaCl
and NaNO₃ mixture
particles**

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

sition, hygroscopicity and cloud condensation activity, *Atmos. Chem. Phys.*, 11, 2585–2602, doi:10.5194/acp-11-2585-2011, 2011.

Gao, Y., Yu, L. E., and Chen, S. B.: Efflorescence relative humidity of mixed sodium chloride and sodium sulfate particles, *J. Phys. Chem. A*, 111, 10660–10666, doi:10.1021/jp073186y, 2007.

Gard, E. E., Kleeman, M. J., Gross, D. S., Hughes, L. S., Allen, J. O., Morrical, B. D., Ferguson, D. P., Dienes, T., Galli, M. E., Johnson, R. J., Cass, G. R., and Prather, K. A.: Direct observation of heterogeneous chemistry in the atmosphere science, *Science*, 279, 1184–1187, 1998.

Ge, Z., Wexler, A. S., and Johnston, M. V.: Multicomponent aerosol crystallization, *J. Colloid Interf. Sci.*, 183, 68–77, 1996.

Ge, Z., Wexler, A. S., and Johnston, M. V.: Deliquescence behavior of multicomponent aerosols, *J. Phys. Chem. A*, 102, 173–180, 1998.

Gibson, E. R., Hudson, P. K., and Grassian, V. H.: Physicochemical properties of nitrate aerosols: implications for the atmosphere, *J. Phys. Chem. A*, 110, 11785–11799, doi:10.1021/jp063821k, 2006.

Gysel, M., Weingartner, E., and Baltensperger, U.: Hygroscopicity of aerosol particles at low temperatures. 2. Theoretical and experimental hygroscopic properties of laboratory generated aerosols, *Environ. Sci. Technol.*, 36, 63–68, doi:10.1021/es010055g, 2002.

Haywood, J. and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: a review, *Rev. Geophys.*, 38, 513–543, doi:10.1029/1999rg000078, 2000.

Hoffman, R. C., Laskin, A., and Finlayson-Pitts, B. J.: Sodium nitrate particles: physical and chemical properties during hydration and dehydration, and implications for aged sea salt aerosols, *J. Aerosol Sci.*, 35, 869–887, 2004.

Hu, D., Qiao, L., Chen, J., Ye, X., Yang, X., Cheng, T., and Fang, W.: Hygroscopicity of inorganic aerosols: size and relative humidity effects on the growth factor, *Aerosol Air Qual. Res.*, 10, 255–264, doi:10.4209/aaqr.2009.12.0076, 2010.

Keene, W. C., Maring, H., Maben, J. R., Kieber, D. J., Pszenny, A. A. P., Dahl, E. E., Iza-guirre, M. A., Davis, A. J., Long, M. S., Zhou, X., Smoydzin, L., and Sander, R.: Chemical and physical characteristics of nascent aerosols produced by bursting bubbles at a model air–sea interface, *J. Geophys. Res.-Atmos.*, 112, D21202, doi:10.1029/2007jd008464, 2007.

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kelly, J. T., Wexler, A. S. Chan, C. K., and Chan, M. N.: Aerosol thermodynamics of potassium salts, double salts, and water content near the eutectic, *Atmos. Environ.* 42, 3717–3728, 2008.
- Kim, H.-K., Lee, M.-J., Jung, H.-J., Eom, H.-J. Maskey, S., Ahn, K.-H., and Ro, C.-U.: Hygroscopic behavior of wet dispersed and dry deposited NaNO₃ particles, *Atmos. Environ.*, 60, 68–75, 2012.
- King, S. M., Butcher, A. C., Rosenoern, T., Coz, E., Lieke, K. I., de Leeuw, G., Nilsson, E. D., and Bilde, M.: Investigating primary marine aerosol properties: CCN activity of sea salt and mixed inorganic–organic particles, *Environ. Sci. Technol.*, 46, 10405–10412, doi:10.1021/es300574u, 2012.
- Krieger, U. K., Marcolli, C., and Reid, J. P.: Exploring the complexity of aerosol particle properties and processes using single particle techniques, *Chem. Soc. Rev.*, 41, 6631–6662, doi:10.1039/c2cs35082c, 2012.
- Krueger, B. J., Grassian, V. H., Iedema, M. J., Cowin, J. P., and Laskin, A.: Probing heterogeneous chemistry of individual atmospheric particles using scanning electron microscopy and energy-dispersive X-ray analysis, *Anal. Chem.*, 75, 5170–5179, doi:10.1021/ac034445t, 2003.
- Laskin, A., Gaspar, D. J., Wang, W., Hunt, S. W., Cowin, J. P., Colson, S. D., and Finlayson-Pitts, B. J.: Reactions at interfaces as a source of sulfate formation in sea-salt particles, *Science*, 301, 340–344, doi:10.1126/science.1085374, 2003.
- Lee, C.-T. and Hsu, W.-C.: The measurement of liquid water mass associated with collected hygroscopic particles, *J. Aerosol Sci.*, 31, 189–197, 2000.
- Li, X., Gupta, D., Eom, H.-J., Kim, H., and Ro, C.-U.: Deliquescence and efflorescence behavior of individual NaCl and KCl mixture aerosol particles, *Atmos. Environ.*, 82, 36–43, doi:10.1016/j.atmosenv.2013.10.011, 2014.
- Lide, D. R.: *Handbook of Chemistry and Physics*, 83rd edn., edited by: Lide, D. R., CRC Press, Boca Raton, Florida, 2002.
- Liu, Y., Cain, J. P., Wang, H., and Laskin, A.: Kinetic study of heterogeneous reaction of deliquesced NaCl particles with gaseous HNO₃ using particle-on-substrate stagnation flow reactor approach, *J. Phys. Chem. A*, 111, 10026–10043, doi:10.1021/jp072005p, 2007.
- Ma, X., von Salzen, K., and Li, J.: Modelling sea salt aerosol and its direct and indirect effects on climate, *Atmos. Chem. Phys.*, 8, 1311–1327, doi:10.5194/acp-8-1311-2008, 2008.

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Martin, S. T.: Phase transitions of aqueous atmospheric particles, *Chem. Rev.*, 100, 3403–3453, 2000.
- McInnes, L. M., Quinn, P. K., Covert, D. S., and Anderson, T. L.: Gravimetric analysis, ionic composition, and associated water mass of the marine aerosol, *Atmos. Environ.*, 30, 869–884, 1996.
- Meskhidze, N., Petters, M. D., Tsigaridis, K., Bates, T., O'Dowd, C., Reid, J., Lewis, E. R., Gantt, B., Anguelova, M. D., Bhave, P. V., Bird, J., Callaghan, A. H., Ceburnis, D., Chang, R., Clarke, A., de Leeuw, G., Deane, G., DeMott, P. J., Elliot, S., Facchini, M. C., Fairall, C. W., Hawkins, L., Hu, Y., Hudson, J. G., Johnson, M. S., Kaku, K. C., Keene, W. C., Kieber, D. J., Long, M. S., Mårtensson, M., Modini, R. L., Osburn, C. L., Prather, K. A., Pszenny, A., Rinaldi, M., Russell, L. M., Salter, M., Sayer, A. M., Smirnov, A., Suda, S. R., Toth, T. D., Worsnop, D. R., Wozniak, A., and Zorn, S. R.: Production mechanisms, number concentration, size distribution, chemical composition, and optical properties of sea spray aerosols, *Atmos. Sci. Lett.*, 14, 207–213, doi:10.1002/asl2.441, 2013.
- Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations, *Atmos. Chem. Phys.*, 9, 9491–9522, doi:10.5194/acp-9-9491-2009, 2009.
- Mochida, M., Nishita-Hara, C., Furutani, H., Miyazaki, Y., Jung, J., Kawamura, K., and Uematsu, M.: Hygroscopicity and cloud condensation nucleus activity of marine aerosol particles over the western North Pacific, *J. Geophys. Res.-Atmos.*, 116, D06204, doi:10.1029/2010jd014759, 2011.
- O'Dowd, C. D. and de Leeuw, G.: Marine aerosol production: a review of the current knowledge, *Philos. T. R. Soc. A*, 365, 1753–1774, doi:10.1098/rsta.2007.2043, 2007.
- Pandis, S. N., Wexler, A. S., and Seinfeld, J. H.: Dynamics of tropospheric aerosols (review), *J. Phys. Chem.*, 99, 9646–9659, 1995.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.
- Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M., Roberts, G. C., Russell, L. M., Ault, A. P., Baltrusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L.,

**Hygroscopic
properties of NaCl
and NaNO₃ mixture
particles**

D. Gupta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hersey, S. P., Kim, M. J., Lambert, W. F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C., and Zhao, D.: Bringing the ocean into the laboratory to probe the chemical complexity of sea spray aerosol, *P. Natl. Acad. Sci. USA*, 110, 7550–7555, doi:10.1073/pnas.1300262110, 2013.

5 Ro, C.-U., Oh, K.-Y., Kim, H., Kim, Y. P., Lee, C. B., Kim, K.-H., Osan, J., de Hoog, J., Worobiec, A., and Van Grieken, R.: Single particle analysis of aerosols at Cheju Island, Korea, using Low-Z electron probe X-ray microanalysis: a direct proof of nitrate formation from sea salts, *Environ. Sci. Technol.*, 35, 4487–4494, 2001.

Ryder, O. S., Ault, A. P., Cahill, J. F., Guasco, T. L., Riedel, T. P., Cuadra-Rodriguez, L. A.,
10 Gaston, C. J., Fitzgerald, E., Lee, C., Prather, K. A., and Bertram, T. H.: On the role of particle inorganic mixing state in the reactive uptake of N₂O₅ to ambient aerosol particles, *Environ. Sci. Technol.*, 48, 1618–1627, doi:10.1021/es4042622, 2014.

Satheesh, S. K. and Moorthy, K. K.: Radiative effects of natural aerosols: a review, *Atmos. Environ.*, 39, 2089–2110, 2005.

15 Saul, T. D., Tolocka, M. P., and Johnston, M. V.: Reactive uptake of nitric acid onto sodium chloride aerosols across a wide range of relative humidities, *J. Phys. Chem. A*, 110, 7614–7620, doi:10.1021/jp060639a, 2006.

Schlenker, J. C. and Martin, S. T.: Crystallization pathways of sulfate-nitrate-ammonium aerosol particles, *J. Phys. Chem. A*, 109, 9980–9985, 2005.

20 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2nd edn., J. Wiley & Sons Inc., Hoboken, N. J., 1203 pp., 2006.

Tang, I. N.: Phase transformation and growth of aerosol particles composed of mixed salts, *J. Aerosol Sci.*, 7, 361–371, 1976.

Tang, I. N. and Munkelwitz, H. R.: Composition and temperature dependence of the deliquescence properties of hygroscopic aerosols, *Atmos. Environ.*, 27, 467–473, 1993.

25 Tang, I. N. and Munkelwitz, H. R.: Aerosol phase transformation and growth in the atmosphere, *J. Appl. Meteorol.*, 33, 791–796, 1994a.

Tang, I. N. and Munkelwitz, H. R.: Water activities, densities, and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance, *J. Geophys. Res.*, 99, 18801–18808, 1994b.

30 Tang, I. N., Munkelwitz, H. R., and Davis, J. G.: Aerosol growth studies – IV phase transformation of mixed salt aerosols in a moist atmosphere, *J. Aerosol Sci.*, 9, 505–511, 1978.

Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559–4593, doi:10.5194/acp-8-4559-2008, 2008.

5 Zuend, A., Marcolli, C., Booth, A. M., Lienhard, D. M., Soonsin, V., Krieger, U. K., Topping, D. O., McFiggans, G., Peter, T., and Seinfeld, J. H.: New and extended parameterization of the thermodynamic model AIOMFAC: calculation of activity coefficients for organic-inorganic mixtures containing carboxyl, hydroxyl, carbonyl, ether, ester, alkenyl, alkyl, and aromatic functional groups, *Atmos. Chem. Phys.*, 11, 9155–9206, doi:10.5194/acp-11-9155-2011, 2011.

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 1. Optical images of NaCl-NaNO₃ particles with a mole fraction of $X_{\text{NaCl}} = 0.8$, obtained during humidifying (\uparrow) and dehydration (\downarrow) processes for the same image field.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 2. Plots of 2-D projected area ratio as a function of the relative humidity (humidifying process: closed circles; dehydration process: open triangles) for different mole fractions of NaCl-NaNO₃, expressed in X_{NaCl} values of **(a)** 0.1 (NaNO₃-rich), **(b)** 0.2 (NaNO₃-rich), **(c)** 0.38 (eutonic), **(d)** 0.5 (equi-molar/NaCl-rich), **(e)** 0.8 (NaCl-rich), and **(f)** 0.9 (NaCl-rich). The transition relative humidity in both humidifying and dehydration processes are marked with arrows.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 3. Measured first DRH or MDRH (open triangles), second DRH (closed circles) values, calculated MDRH (dotted line), and the second DRHs (dash-dotted curve) from the AIOMFAC, plotted as a function of the mole fraction of NaCl in NaCl-NaNO₃ mixture particles. The phase notations shown in brackets are s \equiv solid; and aq \equiv aqueous.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 4. Measured first ERH values (open circles) and second ERH values (open triangles) as a function of the mole fraction of NaCl in NaCl-NaNO₃ mixture particles as well as ERH (closed circle) for wet deposited NaNO₃ particles containing seeds. The phase notations shown in brackets are s ≡ solid; and aq ≡ aqueous.

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 5. Secondary electron images (SEIs) and elemental X-Ray maps for Cl (from NaCl), O (from NaNO₃), and Na of the effloresced NaCl-NaNO₃ mixture particles with compositions of **(a)** $X_{\text{NaCl}} = 0.8$ (NaCl-rich); **(b)** $X_{\text{NaCl}} = 0.38$ (eutonic); and **(c)** $X_{\text{NaCl}} = 0.2$ (NaNO₃-rich).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 6. (a) 2-D diameter ratios (d/d_x) of the particles/aerosols plotted as a function of the NaCl mole fractions (X_{NaCl}), which represent the size increase due to the mutual deliquescence of the eutonic part at MDRH (d_x) compared to that of the solid particle at the start of the first transition (d_x) (eutonic: open squares); and due to the complete deliquescence of the remnant salts (in the figure, NaNO₃: closed circles and NaCl: open circles) at their second DRHs (d_x) compared to that of the partially deliquesced particle at MDRH (d_x) and (b) 2-D diameter ratios (d/d_x) of droplets/aerosols plotted as a function of the NaCl mole fractions (X_{NaCl}), which represent the size decrease due to the efflorescence of NaCl (d_x) compared to that of the liquid droplets at the start of the first efflorescence (d_x) (first ERH: closed circles); and due to the mutual efflorescence of the remnant eutonic compositions (second ERH: open circles) (d_x) compared to that of the partially effloresced particles (d_x).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Hygroscopic properties of NaCl and NaNO₃ mixture particles

D. Gupta et al.




Figure 7. Hygroscopic growth factors in terms of the 2-D diameter ratios (d_{95}/d_0 , where $d_{95} \equiv$ the diameter at $\sim 95\%$, determined by extrapolation, and $d_0 \equiv$ dry diameter at lowest or $\sim 3\%$ RH) are plotted as a function of the mole fraction of NaCl for the NaCl-NaNO₃ mixture particles.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)