



A study of aerosol liquid water content

Y. X. Bian et al.

A study of aerosol liquid water content based on hygroscopicity measurements at high relative humidity in the North China Plain

Y. X. Bian¹, C. S. Zhao¹, N. Ma^{1,2}, J. Chen¹, and W. Y. Xu¹

¹Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing 100871, China

²Leibniz Institute for Tropospheric Research, Permoserstr. 15, 04318 Leipzig, Germany

Received: 28 November 2013 – Accepted: 29 January 2014 – Published: 14 February 2014

Correspondence to: C. S. Zhao (zcs@pku.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Water can be a major component of an aerosol particle. Also water serves as a medium for aqueous-phase reactions in aerosols. In this study, a novel method is presented to calculate the aerosol liquid water content at high relative humidity based on measurements of aerosol hygroscopic growth factor, particle number size distribution and relative humidity in the Haze in China (HaChi) summer field campaign (July–August 2009) in the North China Plain. The aerosol liquid water content calculated using this method agreed well with that calculated using a thermodynamic equilibrium model (ISORROPIA II) at high relative humidity (>60 %) with a correlation coefficient of 0.9658. At low relative humidity (<60 %), an underestimation was found in the calculated aerosol liquid water content by the thermodynamic equilibrium model. This discrepancy mainly resulted from the ISORROPIA II model, which only considered limited aerosol chemical compositions. The mean and maximum values of aerosol liquid water content during the HaChi campaign reached $1.69 \times 10^{-4} \text{ g m}^{-3}$ and $9.71 \times 10^{-4} \text{ g m}^{-3}$, respectively. A distinct diurnal variation of the aerosol liquid water content was found, with lower values during daytime and higher ones at night. The aerosol liquid water content depended strongly on the relative humidity. The aerosol liquid water content in the accumulation mode dominated the total aerosol liquid water content.

1 Introduction

Water uptake by a dry particle composed of inorganic salts, takes place at a critical relative humidity, namely, deliquescence relative humidity (DRH) (Seinfeld and Pandis, 2006). As the relative humidity (RH) increases, the hygroscopic growth of aerosols will continue and become even much faster when RH exceeds 80 % (Engelhart et al., 2011). Once the ambient RH reaches the critical super saturation ratio of aerosol particles, aerosols are activated into cloud droplets. Water uptake by aerosol particles exerts an influence on aerosol optical properties, leading to increased extinction co-

ACPD

14, 4089–4118, 2014

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



efficient and impaired atmospheric visibility, changing the planetary albedo, aerosol optical depth (AOD), and the direct climatic effect of aerosols (Dougle et al., 1996; Seinfeld and Pandis, 2006). Aerosol liquid water content (ALWC) can also speed up the atmospheric chemical reaction rates and serve as the medium for air-particle heterogeneous reactions (Arellanes et al., 2006).

ALWC mostly depends on aerosol hygroscopicity and the ambient RH. Under ambient RH higher than 90 %, liquid water makes up a great part of the aerosol volume (Kitamori et al., 2009). Although the mass concentration of ALWC under sub-saturated conditions is much lower than that the liquid water contents in clouds or fogs, ALWC plays an important role in the tropospheric chemistry (Gelencsér and Varga, 2005). ALWC has a significant effect on the phase partitioning of trace substances and the chemical reactions that takes place inside or on the surface of particulate matter (Kreidenweis et al., 2008).

ALWC can be calculated with detailed aerosol chemical information. However, simulations of aerosol hygroscopicity and phase state could still be quite complicated even under the thermodynamic equilibrium hypothesis. On one hand, ions could be combined in various ways into a variety of solid components. On the other hand, the behaviour of the liquid-phase substances in the aerosol are highly non-ideal due to the high concentrations of solutes under low RH (Wexler and Clegg, 2002). Various aerosols can exist in a metastable liquid state under the condition that dissolved salts are supersaturated, which makes it even more complicated for the simulation process (Martin, 2000). Many aerosol thermodynamic equilibrium models were developed to simulate the process of aerosols dissolving in water and forming ions, including EQUIL, MARS, AIM, E-AIM, SCAPE, EQUISOLV, ISORROPIA etc. (Bassett and Seinfeld, 1983; Saxena et al., 1986; Wexler and Seinfeld, 1991; Kim et al., 1993; Jacobson et al., 1996; Nenes et al., 1998; Wexler and Clegg, 2002). Zhang et al. (2000) compared the simulation results of some of the above mentioned models. Most models only consider how inorganic solutes influence the saturation vapour pressure as solute (Raoult effect). Topping et al. (2005a, b) developed a model called inorganic Aerosol Diam-

A study of aerosol liquid water content

Y. X. Bian et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

eter Dependent Equilibrium Model (ADDEM), which considers both the Raoult effect and the curvature effect (Kelvin effect) of the inorganic/organic aerosol components. McFiggans et al. (2006) suggested to simplify the organic components as primary and secondary components while evaluating their contribution, because it takes a lot of time to process different organic components in the ADDEM.

Another way to obtain the ALWC is based on using size-resolved aerosol hygroscopic growth factors and particle number size distributions (PNSD). Aklilu et al. (2006) used the Quadrupole Aerosol Mass Spectrometer (Q-AMS) to measure size-resolved aerosol chemical component information and assumed constant growth factors for organic compositions. A simplified model that is based on the Zdanovskii-Stokes-Robinson (ZSR; Stokes and Robinson, 1966) mixing rule (the hygroscopicity of mixed aerosols is computed by weighting hygroscopicity of each component according to their dry volume fractions) is then applied to simulate the aerosol growth factors. The comparison between the simulated growth factors and those measured by the Hygroscopicity Tandem Differential Mobility Analyzer (H-TDMA) shows that, for aerosols with sulphate as main inorganic component, the simulated result is similar to the measured one. For aerosols with large proportions of nitrate in the inorganic compositions, the simulated result is higher than the measurement. Similar results have been reported by Gysel et al. (2007). With aerosol chemical components from size segregated filter sampling, Meier et al. (2009) calculated the aerosol growth factors at different particle sizes with a model and then compared it respectively with the H-TDMA and Humidifying Differential Mobility Particle Sizer (HDMPS) measurements. The modelling results based on measured chemical components could only reproduce the hygroscopic growth factors' variation pattern with size as was observed by the other two methods.

Combining the aerosol PNSD with the aerosol hygroscopic growth characteristics, real time ALWC can be calculated. Engelhart et al. (2011) used the Dry-Ambient Aerosol Size Spectrometer (DAASS) to observe the volume growth factor (VGF) after aerosol water uptake, and then combined it with the PNSD measured by a Scanning Mobility Particle Sizer (SMPS) to calculate the ALWC. Comparison between the results

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calculated with VGF and PNSD and the ALWC measured by a Q-AMS that directly collected ambient aerosol in the same field campaign was made. Comparison results displayed a similar variation pattern of growth factors with size, however, the absolute values showed great discrepancies. For ambient RH greater than 70 %, the RH measured by the DAASS and Q-AMS showed outstanding differences, which is why the data observed under RH > 70 % were discarded in their research.

With increasing RH, aerosols will grow in size due to hygroscopic growth processes (Cheng et al., 2008). Especially under high RH (RH > 90 %), a rapid increase of aerosol hygroscopic growth factors (Liu et al., 2011) and aerosol extinction coefficients (Chen et al., 2012) were found, which will lead to more significant impacts of aerosols on climate. Therefore, the study of ALWC under high RH conditions is particularly important. However, only limited researches were performed on this subject. Jung et al. (2009) did a comparative analysis about aerosol mass scattering efficiency and single scattering albedo at both ambient and dry RH in the Pearl River Delta (PRD), China, and then used the result to represent the effect of ALWC on visibility and radiative forcing. The highest RH value was beyond 90 % during the campaign. Instead of calculating the ALWC to quantify the effect of ALWC on visibility and radiative forcing, they only used the ambient RH as a proxy for ALWC to make qualitative descriptions. Kreidenweis et al. (2008) measured the hygroscopic properties of several compounds (sodium chloride (NaCl), ammonium sulphate ((NH₄)₂SO₄), oxalic acid ((HOOC)–(COOH)), etc.) at RH ~ 90 % to calculate the ALWC at different RHs (10 ~ 99 %) in a laboratory, and then compared it with the ALWC simulated from the AIM model. The results retrieved from these two methods showed a significant difference for RH below 85 %. Aggarwal et al. (2007) used an H-TDMA to measure the ALWC at the RH of 5 % to 95 % generated from the water extracts of ambient aerosol filter samples that were collected by a high volume air sampler.

In this paper, High Humidity Tandem Differential Mobility Analyzer (HH-TDMA, Hennig et al., 2005) measurements under high RH (90 ~ 98.5 %) conditions are utilized to acquire aerosol hygroscopicity information using the κ -Köhler theory (Petters and

Kreidenweis, 2007). The result is used to estimate the size-resolved aerosol growth factors at RH from 20 % to 99 %, and then combined with the measured PNSD to calculate the ALWC in July–August 2009 in Wuqing area. A comparison is made between the ALWC obtained by the above described method and the results simulated by the ISORROPIA II model with size-segregated aerosol chemical component information as input. In addition, the influences of both RH and PNSD on the ALWC are analysed in detail.

2 Data and measurements

2.1 Site

The North China Plain (NCP) has experienced rapid economic developments during the past few decades. Meanwhile, the air pollution in this area has become a very serious problem (Xu et al., 2011). The HaChi project was jointly sponsored by Peking University and the Leibniz Institute for Tropospheric Research (TROPOS) during July–August 2009 at the Wuqing Meteorological Station (39°23' N, 117°01' E, 7.4 m.a.s.l.). The description of the site and measurements refer to Liu et al. (2011) and Ran et al. (2011). The time zone that contains this site is in UTC+8, so all of the time is reported as UTC+8 in this paper. Figure 1 shows the location of Wuqing site, the major neighbouring cities and the average geographical distribution of MODIS (Moderate Resolution Imaging Spectroradiometer) AOD at 550 nm wavelength in 2009. As shown by the AOD distribution, AOD values are high over the entire NCP (AOD > 0.6) and, in addition, the AOD distribution is relatively homogeneous over the intercity area. The AOD value is about 0.7 at Wuqing site, which is close to the average AOD of the NCP. Therefore, this site is highly representative of the general situation of aerosol pollution in the NCP.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Instrumentation

The aerosol hygroscopic growth properties at high humidity, the aerosol PNSD at dry state, the aerosol chemical components and the ambient temperature and RH were sampled from 15 July to 15 August 2009. Here the aerosol PNSD at dry state is referred to the PNSD measurements which were conducted under RH below 40%. During the campaign, the HH-TDMA and the Twin Differential Mobility Particle Sizer (TDMPS)/Aerodynamic Particle Sizer (APS) for the aerosol hygroscopicity and PNSD measurements were housed in a mobile laboratory consisting of a refitted shipping container. The sample flow was dried to RH less than 40% before flowing into the laboratory (Tuch et al., 2009). The room temperature of the laboratory was kept at a steady 22 °C. The Berner low-pressure Impactor (BLPI) for the aerosol filter sampling was placed near the Automatic Weather Station (AWS) in an open field. The stage films of the BLPI were changed twice a day. Measurements were made from 11:00 to 17:00 LT and from 20:00 to 02:00 LT. The samples were carefully stored at about -20 °C for further analysis.

The HH-TDMA is an important instrument in this study, designed to measure the size-resolved aerosol hygroscopic growth characteristics at high RH (90%–98.5%) (Hennig et al., 2005). The basic principle is similar to that of the H-TDMA (Rader and McMurry, 1986). The widely used H-TDMA are not able to hold the RH of aerosols and sheath flow stable, for RH > 90%. To measure the aerosol hygroscopic growth at high RH, TROPOS used the technique of temperature controlled water bath and re-designed the HH-TDMA based on the H-TDMA.

The TDMPS (Birmili et al., 1999) includes two Hauke-type DMAs that have different effective centre rod lengths which measure aerosol particles in size range of 20–800 nm and 3–20 nm, respectively. The two Condensation Particle Counters (CPCs) are used to count the aerosol particles downstream of the two DMAs. Combining the counts from the two CPCs, the TDMPS can measure the PNSD of aerosol particles from 3–800 nm (electrical mobility diameter). It took 10 min for a complete scan for the TDMPS. The

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



APS (Model 3320, TSI, Inc., Shoreview, MN USA) was used to measure the aerosol size distribution from 0.5 to 10 μm aerodynamic particle diameter (Armendariz and Leith, 2002).

Both TDMPs and APS take 10 min to complete a whole scan. Detailed information on the pre-processing of the TDMPs data can be found in Ma et al. (2011). Using the algorithm described by DeCarlo et al. (2004), electrical mobility equivalent diameter measured by TDMPs and aerodynamic equivalent diameter measured by APS were transformed to volume equivalent diameter. The density was assumed as 1.7 g cm^{-3} directly. Then the PNSD were interpolated into a logarithmically uniformly spaced diameter from 3 nm to 10 μm . Moreover, the PNSD data was adjusted for impaction losses, diffusion losses and gravitational losses. The temporal resolution of the PNSD data was 10 min.

The BLPI segregates aerosols into 10 stages. The sampler segregates diameters by different sized orifices which lead to different flow rates. Every stage is made up of two parts: air flows across the jet hold on the first part and particles with diameters larger than the cut-off diameter of that stage will be collected by the sampling filter on the second part. When the air flow passes through the instrument, particulates with different diameters will impact on different stages with filters collecting them (Hillamo and Kauppinen, 1991). Excluding stage 10, the measuring range is from 15 nm to 8 μm , segregated into 9 stages. The lower and upper bounds for each stage are shown in Table 1. The detail information about the particle size ranges and mean diameters for the individual impactor stages is refer to Liu et al. (2013). By preweighing and then weighing and analysing the films afterwards in the laboratory, the stage segregated aerosol chemistry information was determined.

A study of aerosol liquid water content

Y. X. Bian et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3 Methods

3.1 Size-resolved hygroscopic growth factors ($g(D_d, RH)$) based on HH-TDMA observation

Particles water uptake can be represented by the Köhler theory (Köhler, 1936). Recently, some single parameter representations of the Köhler equation have been proposed in previous studies (Rissler et al., 2006; Petters and Kreidenweis, 2007; Wex et al., 2007). In this study, a widely used hygroscopicity parameter κ is used to analyse the hygroscopic growth of particles in the subsaturated regime. Based on the Köhler equation, the physical quantities about aerosol hygroscopicity are simplified to κ to obtain a new equation, named “ κ -Köhler theory” (Petters and Kreidenweis, 2007). According to this theory, $S(D)$, the saturation ratio over the hydrated particulate solution surface can be calculated from,

$$S(D) = \frac{D^3 - D_d^3}{D^3 - D_d^3(1 - \kappa)} \exp\left(\frac{4\sigma_{s/a}M_w}{RT\rho_w D}\right), \quad (1)$$

where $\sigma_{s/a}$ is the surface tension coefficient, M_w is the mole mass of water, R is the universal gas constant, T is the temperature, ρ_w is the density of water, D_d is the diameter of dry particle and D is the diameter of wet particle. For a known T and S , the relationship between κ and $g(D_d, RH)$, the hygroscopic growth factor at a certain diameter D_d and a certain RH, can be calculated by,

$$\kappa = \left(g(D_d, RH)^3 - 1\right) \left[\frac{1}{S(D)} \exp\left(\frac{4\sigma_{s/a}M_w}{RT\rho_w D_d \cdot g(D_d, RH)}\right) - 1 \right], \quad (2)$$

where $g(D_d, RH) = D/D_d$. In this study, T used to calculate κ is set to 293 K, $\sigma_{s/a}$ is assumed to be the surface tension coefficient between water and air (about 0.0728 N m^{-1} at 293 K).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Liu et al. (2011) found that, for a certain particle diameter, the according growth factors varies with RH. The κ value, however, does not vary with RH. The HH-TDMA was used to measure the hygroscopic growth factors at four dry aerosol diameters (50 nm/100 nm/200 nm/250 nm) in this campaign. Calculations need to be made to derive a size-resolved hygroscopic growth factors. First, a 4 mode lognormal distribution is applied to fit the measured PNSD, yielding fitting parameters for each mode (Tareq et al., 2005). Afterwards, assuming that κ is constant within the same mode, the size-resolved κ is calculated by combining the 4-mode fitting results with the measured κ values at certain particle diameters (Fig. 2). The detail of the fitting process is described by Chen et al. (2012). Finally, the size-resolved κ is substituted into Eq. (2) to calculate the size-resolved growth factors at 20 ~ 99 % RH.

3.2 Calculation of ALWC from $g(D_d, RH)$

With the assumption of the conservation of volume during aerosol hygroscopic process, the volume of ALWC is equal to subtract the volume of dry aerosol particles from that of wet particles. Using the size-resolved hygroscopic growth factors and the PNSD measurement, the ALWC can be calculated:

$$ALWC = \left[\frac{\pi}{6} \sum_i N_i D_{d,i}^3 \left(g(D_d, RH)^3 - 1 \right) \right] \cdot \rho_w, \quad (3)$$

Where, $N_i(D_{d,i})$ represents the number concentration of dry particles of the i th bin, $D_{d,i}$ is the particle diameter of that bin, ρ_w is 1 g cm^{-3} in this study. ALWC is given in the unit of g m^{-3} of air.

3.3 Simulation of ALWC from ISORROPIA II model

ISORROPIA II is a computationally efficient thermodynamic equilibrium model developed recently by Fountoukis and Nenes (2007). All chemical components the BLPI filter

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sample analysis yielded are considered in this model except for the organic species. Liu et al. (2013) shows that the overall particle hygroscopicity is mostly contributed by inorganic compounds in the NCP and the water soluble organic compounds play a critical role only for ultrafine particles smaller than 100 nm. The thermodynamics of the crustal species including Ca^{2+} , K^+ and Mg^{2+} have been added to the model's treatment components along with NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- and H_2O , but organic species are not taken into account. ISORROPIA II is used to model the ALWC (the temperature was set to 293 K). ISORROPIA II exhibits robust and rapid convergence under all conditions, with especially prominent performance on computational speed. Based on its computational rigor and performance on computational speed, ISORROPIA II appears to be an extremely desirable choice for large scale atmospheric transport and air quality models (Fountoukis and Nenes, 2007). The ISORROPIA II model can simulate the ALWC with the input of aerosol chemical composition measured by BLPI sampling (Liu et al., 2013).

ISORROPIA II model assumes that the aerosol curvature effect can be ignored and the water uptake of aerosols has no effect to the ambient vapour pressure. Consequently, the water activity a_w is equal to the ambient RH caused by the phase equilibrium between gas and aerosols (Seinfeld and Pandis, 2006).

$$a_w = \text{RH}. \quad (4)$$

The water uptake of aerosols is calculated with the ZSR hypothesis,

$$\text{ALWC} = \sum_i \frac{M_i}{m_{oi}(a_w)}. \quad (5)$$

Where, M_i is the mole concentration of the electrolyte i (mol m^{-3} air), and $m_{oi}(a_w)$ is the corresponding molality of the binary solution of the electrolyte i under the same a_w with complex solution.

4 Results and discussion

4.1 Comparison between the two methods to estimate the ALWC

The aerosol liquid water content calculated from the measured hygroscopic growth factors is compared with the results modelled with ISORROPIA II. As can be seen in Fig. 3, at low RH, the ISORROPIA II ALWC values are much smaller than those calculated from the growth factors. With increasing RH, the two results gradually converge. For RH over 60 %, the results from the two methods agree well with a correlation coefficient R of 0.96. The possible explanations for these results can be summarized as follows.

First, the more complicated the aerosol chemical compositions are, the lower the DRH of aerosols becomes (Seinfeld and Pandis, 2006). Thus, due to the limited types of inorganic species considered in the ISORROPIA II model, the modelled DRH is much higher than that in the real atmosphere. For RH below the modelled DRH (around 60 %), aerosols are predicted to be still in a dry state, not significantly hydrated, and with small ALWC, whereas in the real atmosphere it might have already crossed over to a hydrated state. The modelled ALWC are therefore much lower than those calculated with the growth factors at RH below the simulated DRH.

Furthermore, ISORROPIA II assumes that the aerosol is completely dry at the beginning of the simulation, then dissolves each of its salt component and simulates water uptake as the ambient RH increases. This solution dynamics is different from that of some other models, like SCAPE2, which leads to the fact that the ISORROPIA II modelled ALWC results under low RH are not only lower than the growth factors calculated result, but also lower than that modelled by SCAPE2 (Fountoukis and Nenes, 2007).

The modelled ALWC is closer to the calculated one at high RH. The shape and mixing state of aerosols is not considered in the ISORROPIA II and the effect of non-spheroid is also ignored. Thus, the “inverse Kelvin effect” is disregarded in the ISORROPIA II. This effect enhances the water uptake of real aerosols compared with the flat surface that is assumed in ISORROPIA II and also leads to a decrease in the DRH

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Sjogren et al., 2007). On the other hand, the κ values used to calculate the size-resolved hygroscopic growth factors are measured under high RH conditions ($> 90\%$), so the deliquescence process is not considered in the calculation. For RH below the DRH, a discrepancy may exist between the calculated ALWC and the true value.

4.2 Time series and diurnal variation of ALWC

Figure 4a shows the time series of ALWC calculated from the growth factors at ambient RH during HaChi summer campaign. During the campaign, the maximum value of ALWC is $9.71 \times 10^{-4} \text{ g m}^{-3}$, with an average value of $(1.69 \pm 1.64) \times 10^{-4} \text{ g m}^{-3}$. Although these values are much lower than the ALWC in clouds measured by aircraft over this region (Deng et al., 2009), the significance of ALWC cannot be ignored. The life time of a cloud is about several hours, but the life of aerosols may last for several days in the air (Seinfeld and Pandis, 2006). The ALWC also plays an important role in atmospheric chemistry and radiation in the NCP due to high aerosol loading and long lifetime of aerosols.

The calculated ALWC shows an obvious diurnal variation (Fig. 4b). During the daytime, the ALWC stays at low level. After 18:00 LT, the ALWC begins to rise to relative high values, since particle emissions are accumulated in the narrowed surface layer with RH increasing simultaneously. The maximum ALWC appears at 6:00 LT. Afterwards it drops slightly as a result of increased surface temperature that leads to the growth of the boundary layer and reduces the relative humidity (Stull, 1988). Due to the vehicle combustion resulting from the rush hours, the ALWC does not decrease very quickly during 6:00–9:00 LT. After that period, a rapid reduction in the ALWC can be observed. The growth factors of most particles varies little in a day, hence has slight influence on the diurnal variation of the ALWC. During the campaign, no regular variation patterns of the daily averaged ALWC could be found (Fig. 4c).

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Factors influencing the ALWC

4.3.1 Impacts of RH

According to Eq. (3), the absolute value of ALWC depends to a great extent on the dry aerosol volume. To evaluate how RH impact the ALWC, the influence of the dry volume of aerosols has to be removed. Consequently, a physical quantity called relative ALWC is defined as:

$$\text{Relative ALWC} = \text{ALWC}/C_{\text{volume}} \quad (6)$$

where C_{volume} represents the volume concentration of dry aerosol particles.

The physical meaning of the relative ALWC is the mass of water absorbed by unit volume of dry aerosols. A significant positive correlation can be found between the relative ALWC and RH using all of the calculated ALWC and the according aerosol volume concentration and ambient RH values during the HaChi campaign (Fig. 5). Under low RH, the relative ALWC rises slowly with RH. When RH exceeds 80 %, the relative ALWC increases rapidly with RH, which means that RH is an important influencing factor in the ALWC variation.

4.3.2 Impacts of PNSD

PNSD is another important factor influencing the aerosol liquid water content. The aerosol particles are separated into four modes according to their different origins (Whitby, 1978; Birmili et al., 2001): nucleation mode (< 30 nm), Aitken mode (30–110 nm), accumulation mode (110 nm–1 μm) and coarse mode (> 1 μm). The relationship between ALWC and the number concentration of each mode has been analysed to evaluate the impacts of PNSD on the ALWC. These relationships under five different RH conditions are discussed based on the discussion of RH impact above. Then the ALWC is linearly fitted with the number concentrations of particles in each mode under varying RH conditions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2 shows the coefficient of determination of the linear regressions of ALWC and particulate number concentrations. The R^2 values of the nucleation and the Aitken mode particles are mostly less than 0.2 under all the RH conditions, which suggests that the particle number concentrations of those two modes have little influence on the aerosol ALWC due to their small volume concentration and low hygroscopicity. The hygroscopicity of aerosol particles in these two modes is relatively low because of their origins (Swietlicki et al., 1999). Freshly emitted hydrophobic particles such as externally mixed black carbon particles are mostly in the fine modes. Besides, the significant Kelvin effect also leads to higher hygroscopic growth factors for particles in the coarse mode than for those in fine modes (Väkevää et al., 2002). For the coarse mode, the R^2 are between 0.3 and 0.6 under all the RH conditions. It presents an existent but not remarkable correlation between the ALWC and the number concentration. Although some aerosols in the coarse mode are water-soluble and have certain impacts on the ALWC, nearly-hydrophobic aerosols, such as dust particles contribute a large number fraction to the coarse mode (Hussein et al., 2004; Liu et al., 2008, 2013).

The R^2 for the accumulation mode is significantly higher than those for the other three modes (Fig. 6). All the results are above 0.6 at $RH > 60\%$. This indicates that the number concentration of the accumulation mode does play a significant role in the ALWC variation. Accumulation mode particles are mostly aged, secondary aerosol and are transported within boundary layer and subjected to cloud processing, which leads to higher hygroscopicity than those of the nucleation and Aitken mode aerosols (Swietlicki et al., 1999; Vignati et al., 2004). Accumulation mode particles also have a large volume fraction of aerosols.

5 Summary

ALWC is an important physical quantity that can significantly influence aerosol optical and chemical properties. However, direct measurement of ALWC is hard to achieve, especially under high RH conditions. In this paper, a novel method to calculate the ALWC

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from aerosol hygroscopicity measurements is proposed based on the size-resolved aerosol hygroscopic growth factors, PNSD, and RH. The size-resolved aerosol growth factors is obtained using the κ -Köhler theory with the HH-TDMA measurements under high RH (90 ~ 98.5 %).

5 The ALWC is also modelled using ISORROPIA II model with the input of measured aerosol chemistry data during the summertime field campaign in 2009. Comparisons between the ALWC results calculated by hygroscopic growth factors and modelled by ISORROPIA II show that under low RH, the modelled results are much smaller than the calculated ones. With the increase of RH, the results from these two methods
10 gradually converge together. They agree well with each other at RH higher than 60 %, with a correlation coefficient of 0.96.

The maximum value of ALWC calculated with aerosol measurements in the NCP between July and August 2009 reaches $9.71 \times 10^{-4} \text{ gm}^{-3}$, with an average value of $1.69 \times 10^{-4} \text{ gm}^{-3}$. The ALWC experiences significant diurnal variations at Wuqing site,
15 with lower values during daytime and higher ones during night time. Aerosol liquid water content is highly related to the relative humidity. The contribution to the aerosol liquid water content from the accumulation mode is dominating among all the aerosol particle modes.

Acknowledgements. This work is supported by the National 973 project of China (2011CB403402), the National Natural Science Foundation of China under Grant No. 41375134, the Beijing Natural Science Foundation (8131003), the Beijing municipal science and technology plan project No. Z131100006113013, and the Tianjin municipal science and technology support program (13ZCZDSF02100).
20

References

25 Aggarwal, S. G., Mochida, M., Kitamori, Y., and Kawamura, K.: Chemical closure study on hygroscopic properties of urban aerosol particles in Sapporo, Japan, Environ. Sci. Technol., 41, 6920–6925, 2007.

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aklilu, Y., Mozurkewich, M., Prenni, A. J., Kreidenweis, S. M., Alfarra, M. R., Allan, J. D., Anlauf, K., Brook, J., Leaitch, W. R., Sharma, S., Boudries, H., and Worsnop, D. R.: Hygroscopicity of particles at two rural, urban influenced sites during Pacific 2001: comparison with estimates of water uptake from particle composition, *Atmos. Environ.*, 40, 2650–2661, 2006.

Arellanes, C., Paulson, S. E., Fine, P. M., and Sioutas, C.: Exceeding of Henry's Law by hydrogen peroxide associated with urban aerosols, *Environ. Sci. Technol.*, 40, 4859–4866, 2006.

Armendariz, A. J. and Leith, D.: Concentration measurement and counting efficiency for the aerodynamic particle sizer 3320, *J. Aerosol Sci.*, 33, 133–148, 2002.

Bassett, M. and Seinfeld, J. H.: Atmospheric equilibrium model of sulfate and nitrate aerosols, *Atmos. Environ.*, 17, 2237–2252, 1983.

Birmili, W., Stratmann, F., Wiedensohler, A.: Design of a DMA-based size spectrometer for a large particle size range and stable operation, *J. Aerosol Sci.*, 30, 549–553, 1999.

Birmili, W., Wiedensohler, A., Heintzenberg, J., and Lehmann, K.: Atmospheric particle number size distribution in central Europe: statistical relations to air masses and meteorology, *J. Geophys. Res.-Atmos.*, 106, 32005–32018, 2001.

Chen, J., Zhao, C. S., Ma, N., Liu, P. F., Göbel, T., Hallbauer, E., Deng, Z. Z., Ran, L., Xu, W. Y., Liang, Z., Liu, H. J., Yan, P., Zhou, X. J., and Wiedensohler, A.: A parameterization of low visibilities for hazy days in the North China Plain, *Atmos. Chem. Phys.*, 12, 4935–4950, doi:10.5194/acp-12-4935-2012, 2012.

Cheng, Y. F., Wiedensohler, A., Eichler, H., Heintzenberg, J., Tesche, M., Ansmann, A., Wendisch, M., Su, H., Althausen, D., Herrmann, H., Gnauk, T., Brüggemann, E., Hu, M., and Zhang, Y. H.: Relative humidity dependence of aerosol optical properties and direct radiative forcing in the surface boundary layer at Xinken in Pearl River Delta of China: an observation based numerical study, *Atmos. Environ.*, 42, 6373–6397, 2008.

DeCarlo, P. F., Slowik, J. G., Worsnop, D. R., Davidovits, P., and Jimenez, J. L.: Particle morphology and density characterization by combined mobility and aerodynamic diameter measurements – Part 1: Theory, *Aerosol Sci. Tech.*, 38, 1185–1205, 2004.

Deng, Z., Zhao, C., Zhang, Q., Huang, M., and Ma, X.: Statistical analysis of microphysical properties and the parameterization of effective radius of warm clouds in Beijing area, *Atmos. Res.*, 93, 888–896, 2009.

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dougle, P. G., Vlasenko, A. L., Veefkind, J. P., and Brink, H. M. T.: Humidity dependence of the light scattering by mixtures of ammonium nitrate, ammonium sulfate and soot, *J. Aerosol Sci.*, 27, 513–514, 1996.
- Engelhart, G. J., Hildebrandt, L., Kostenidou, E., Mihalopoulos, N., Donahue, N. M., and Pandis, S. N.: Water content of aged aerosol, *Atmos. Chem. Phys.*, 11, 911–920, doi:10.5194/acp-11-911-2011, 2011.
- Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for $\text{K}^+ - \text{Ca}^{2+} - \text{Mg}^{2+} - \text{NH}_4^+ - \text{Na}^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{Cl}^- - \text{H}_2\text{O}$ aerosols, *Atmos. Chem. Phys.*, 7, 4639–4659, doi:10.5194/acp-7-4639-2007, 2007.
- Gelencsér, A. and Varga, Z.: Evaluation of the atmospheric significance of multiphase reactions in atmospheric secondary organic aerosol formation, *Atmos. Chem. Phys.*, 5, 2823–2831, doi:10.5194/acp-5-2823-2005, 2005.
- Gysel, M., Crosier, J., Topping, D. O., Whitehead, J. D., Bower, K. N., Cubison, M. J., Williams, P. I., Flynn, M. J., McFiggans, G. B., and Coe, H.: Closure study between chemical composition and hygroscopic growth of aerosol particles during TORCH2, *Atmos. Chem. Phys.*, 7, 6131–6144, doi:10.5194/acp-7-6131-2007, 2007.
- Hennig, T., Massling, A., Brechtel, F. J., and Wiedensohler, A.: A tandem DMA for highly temperature-stabilized hygroscopic particle growth measurements between 90 % and 98 % relative humidity, *J. Aerosol Sci.*, 36, 1210–1223, 2005.
- Hillamo, R. E. and Kauppinen, E. I.: On the performance of the Berner low pressure impactor, *Aerosol Sci. Tech.*, 14, 33–47, 1991.
- Hussein, T., Puustinen, A., Aalto, P. P., Mäkelä, J. M., Hämeri, K., and Kulmala, M.: Urban aerosol number size distributions, *Atmos. Chem. Phys.*, 4, 391–411, doi:10.5194/acp-4-391-2004, 2004.
- Hussein, T., Dal Maso, M., Petäjä, T., Koponen, I. K., Paatero, P., Aalto, P. P., Hämeri, K., and Kulmala, M.: Evaluation of an automatic algorithm for fitting the particle number size distributions, *Boreal Environ. Res.*, 10, 337–355, 2005.
- Jacobson, M. Z., Tabazadeh, A., and Turco, R. P.: Simulating equilibrium within aerosols and nonequilibrium between gases and aerosols, *J. Geophys. Res.-Atmos.*, 101, 9079–9091, 1996.
- Kim, Y. P., Seinfeld, J. H., and Saxena, P.: Atmospheric gas-aerosol equilibrium I. Thermodynamic model, *Aerosol Sci. Tech.*, 19, 157–181, 1993.

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kitamori, Y., Mochida, M., and Kawamura, K.: Assessment of the aerosol water content in urban atmospheric particles by the hygroscopic growth measurements in Sapporo, Japan, *Atmos. Environ.*, 43, 3416–3423, 2009.
- Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *T. Faraday Soc.*, 32, 1152–1161, 1936.
- Kreidenweis, S. M., Petters, M. D., and DeMott, P. J.: Single-parameter estimates of aerosol water content, *Environ. Res. Lett.*, 3, 035002, doi:10.1088/1748-9326/3/3/035002, 2008.
- Jung, J., Lee, H., Kim, Y. J., Liu, X., Zhang, Y., Gu, J., and Fan, S.: Aerosol chemistry and the effect of aerosol water content on visibility impairment and radiative forcing in Guangzhou during the 2006 Pearl River Delta campaign, *J. Environ. Manage.*, 90, 3231–3244, 2009.
- Liu, H. J., Zhao, C. S., Nekat, B., Ma, N., Wiedensohler, A., van Pinxteren, D., Spindler, G., Müller, K., and Herrmann, H.: Aerosol hygroscopicity derived from size-segregated chemical composition and its parameterization in the North China Plain, *Atmos. Chem. Phys. Discuss.*, 13, 20885–20922, doi:10.5194/acpd-13-20885-2013, 2013.
- Liu, P. F., Zhao, C. S., Göbel, T., Hallbauer, E., Nowak, A., Ran, L., Xu, W. Y., Deng, Z. Z., Ma, N., Mildenerger, K., Henning, S., Stratmann, F., and Wiedensohler, A.: Hygroscopic properties of aerosol particles at high relative humidity and their diurnal variations in the North China Plain, *Atmos. Chem. Phys.*, 11, 3479–3494, doi:10.5194/acp-11-3479-2011, 2011.
- Liu, S., Hu, M., Wu, Z., Wehner, B., Wiedensohler, A., and Cheng, Y.: Aerosol number size distribution and new particle formation at a rural/coastal site in Pearl River Delta (PRD) of China, *Atmos. Environ.*, 42, 6275–6283, 2008.
- Ma, N., Zhao, C. S., Nowak, A., Müller, T., Pfeifer, S., Cheng, Y. F., Deng, Z. Z., Liu, P. F., Xu, W. Y., Ran, L., Yan, P., Göbel, T., Hallbauer, E., Mildenerger, K., Henning, S., Yu, J., Chen, L. L., Zhou, X. J., Stratmann, F., and Wiedensohler, A.: Aerosol optical properties in the North China Plain during HaChi campaign: an in-situ optical closure study, *Atmos. Chem. Phys.*, 11, 5959–5973, doi:10.5194/acp-11-5959-2011, 2011.
- Martin, S. T.: Phase transitions of aqueous atmospheric particles, *Chem. Rev.*, 100, 3403–3454, 2000.
- McFiggans, G., Artaxo, P., Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S., Gysel, M., Laaksonen, A., Lohmann, U., Mentel, T. F., Murphy, D. M., O'Dowd, C. D., Snider, J. R., and Weingartner, E.: The effect of physical and chemical aerosol properties on warm cloud droplet activation, *Atmos. Chem. Phys.*, 6, 2593–2649, doi:10.5194/acp-6-2593-2006, 2006.

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Meier, J., Wehner, B., Massling, A., Birmili, W., Nowak, A., Gnauk, T., Brüggemann, E., Herrmann, H., Min, H., and Wiedensohler, A.: Hygroscopic growth of urban aerosol particles in Beijing (China) during wintertime: a comparison of three experimental methods, *Atmos. Chem. Phys.*, 9, 6865–6880, doi:10.5194/acp-9-6865-2009, 2009.
- 5 Nenes, A., Pandis, S., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols, *Aquat. Geochem.*, 4, 123–152, 1998.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.
- 10 Rader, D. J. and McMurry, P. H.: Application of the tandem differential mobility analyzer to studies of droplet growth or evaporation, *J. Aerosol Sci.*, 17, 771–787, 1986.
- Ran, L., Zhao, C. S., Xu, W. Y., Lu, X. Q., Han, M., Lin, W. L., Yan, P., Xu, X. B., Deng, Z. Z., Ma, N., Liu, P. F., Yu, J., Liang, W. D., and Chen, L. L.: VOC reactivity and its effect on ozone production during the HaChi summer campaign, *Atmos. Chem. Phys.*, 11, 4657–4667, doi:10.5194/acp-11-4657-2011, 2011.
- 15 Rissler, J., Vestin, A., Swietlicki, E., Fisch, G., Zhou, J., Artaxo, P., and Andreae, M. O.: Size distribution and hygroscopic properties of aerosol particles from dry-season biomass burning in Amazonia, *Atmos. Chem. Phys.*, 6, 471–491, doi:10.5194/acp-6-471-2006, 2006.
- Saxena, P., Belle Hudischewskyj, A., Seigneur, C., and Seinfeld, J. H.: A comparative study of equilibrium approaches to the chemical characterization of secondary aerosols, *Atmos. Environ.*, 20, 1471–1483, 1986.
- 20 Seinfeld, J. and Pandis, S.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley, Inc., New Jersey, USA, 450 pp., 2006.
- Sjogren, S., Gysel, M., Weingartner, E., Baltensperger, U., Cubison, M. J., Coe, H., Zardini, A. A., Marcolli, C., Krieger, U. K., and Peter, T.: Hygroscopic growth and water uptake kinetics of two-phase aerosol particles consisting of ammonium sulfate, adipic and humic acid mixtures, *J. Aerosol Sci.*, 38, 157–171, 2007.
- 25 Stokes, R. H. and Robinson, R. A.: *Interactions in aqueous nonelectrolyte solutions. I. Solute-solvent equilibria*, *J. Phys. Chem.*, 70, 2126–2131, 1966.
- 30 Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Springer, New York, USA, 670 pp., 1988.

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Swietlicki, E., Zhou, J., Berg, O. H., Martinsson, B. G., Frank, G., Cederfelt, S. I., Dusek, U., Berner, A., Birmili, W., Wiedensohler, A., Yuskiewicz, B., and Bower, K. N.: A closure study of sub-micrometer aerosol particle hygroscopic behaviour, *Atmos. Res.*, 50, 205–240, 1999.
- 5 Topping, D. O., McFiggans, G. B., and Coe, H.: A curved multi-component aerosol hygroscopicity model framework: Part 1 – Inorganic compounds, *Atmos. Chem. Phys.*, 5, 1205–1222, doi:10.5194/acp-5-1205-2005, 2005a.
- Topping, D. O., McFiggans, G. B., and Coe, H.: A curved multi-component aerosol hygroscopicity model framework: Part 2 – Including organic compounds, *Atmos. Chem. Phys.*, 5, 1223–1242, doi:10.5194/acp-5-1223-2005, 2005b.
- 10 Tuch, T. M., Haudek, A., Müller, T., Nowak, A., Wex, H., and Wiedensohler, A.: Design and performance of an automatic regenerating adsorption aerosol dryer for continuous operation at monitoring sites, *Atmos. Meas. Tech.*, 2, 417–422, doi:10.5194/amt-2-417-2009, 2009.
- Väkevä, M., Kulmala, M., Stratmann, F., and Hämeri, K.: Field measurements of hygroscopic properties and state of mixing of nucleation mode particles, *Atmos. Chem. Phys.*, 2, 55–66, doi:10.5194/acp-2-55-2002, 2002.
- 15 Vignati, E., Wilson, J., and Stier, P.: M7: an efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res.-Atmos.*, 109, D22202, doi:10.1029/2003JD004485, 2004.
- Wex, H., Hennig, T., Salma, I., Ocskay, R., Kiselev, A., Henning, S., Massling, A., Wiedensohler, A., and Stratmann, F.: Hygroscopic growth and measured and modeled critical super-saturations of an atmospheric HULIS sample, *Geophys. Res. Lett.*, 34, L02818, doi:10.1029/2006GL028260, 2007.
- 20 Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *J. Geophys. Res.-Atmos.*, 107, ACH 14-11–ACH 14-14, 2002.
- 25 Wexler, A. S. and Seinfeld, J. H.: Second-generation inorganic aerosol model, *Atmos. Environ. A-Gen.*, 25, 2731–2748, 1991.
- Whitby, K. T.: The physical characteristics of sulfur aerosols, *Atmos. Environ.*, 12, 135–159, 1978.
- 30 Xu, W. Y., Zhao, C. S., Ran, L., Deng, Z. Z., Liu, P. F., Ma, N., Lin, W. L., Xu, X. B., Yan, P., He, X., Yu, J., Liang, W. D., and Chen, L. L.: Characteristics of pollutants and their correlation to meteorological conditions at a suburban site in the North China Plain, *Atmos. Chem. Phys.*, 11, 4353–4369, doi:10.5194/acp-11-4353-2011, 2011.

Zhang, Y., Seigneur, C., Seinfeld, J. H., Jacobson, M., Clegg, S. L., and Binkowski, F. S.: A comparative review of inorganic aerosol thermodynamic equilibrium modules: similarities, differences, and their likely causes, *Atmos. Environ.*, 34, 117–137, 2000.

ACPD

14, 4089–4118, 2014

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. The lower and upper bounds for each stage of the BLPI. They represent the 50 % cut-off aerodynamic diameters.

Impactor Stage	1	2	3	4	5	6	7	8	9	10
Lower Bound (μm)	0.015	0.03	0.06	0.125	0.25	0.5	1	2	4	8
Upper Bound (μm)	0.03	0.06	0.125	0.25	0.5	1	2	4	8	16

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. The coefficient of determination of each aerosol mode under different RH conditions.

RH (%)	Coefficient of determination (R^2)			
	Nucleation Mode	Aitken Mode	Accumulation Mode	Coarse Mode
≤ 60	0.16	0.02	0.54	0.35
60 ~ 70	0.11	0.01	0.76	0.59
70 ~ 80	0.10	0.01	0.74	0.58
80 ~ 90	0.11	0.10	0.62	0.48
> 90	0.04	0.32	0.78	0.35




Fig. 1. MODIS AOD distribution and the location of the measurement site (red star) and regional mega cities (circle).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A study of aerosol liquid water content

Y. X. Bian et al.




Fig. 2. Size-resolved κ (blue line) and the corresponding mean values ± 1 standard deviation of observed κ (green circles and error bars) of particles with dry diameters of 50 nm, 100 nm, 200 nm and 250 nm during the HaChi summer campaign.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Comparison between ALWC calculated from measured growth factors (ALWC-GF) and modelled with ISORROPIA II (ALWC-ISO II); the colour of the dots present their corresponding RHs; the solid line shows the 1 : 1 line.

A study of aerosol liquid water content

Y. X. Bian et al.




Fig. 4. (a) Time series of the estimated ALWC during the HaChi summer campaign (denoted by the colour). (b) Diurnal variation of the estimated ALWC. (c) Daily averaged time series of the estimated ALWC. The blue region of the time and diurnal series denotes ± 1 standard deviation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 5. The relationship between relative ALWC and RH. Relative ALWC is defined as the mass of water absorbed by unit volume of dry aerosols.

A study of aerosol liquid water content

Y. X. Bian et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



A study of aerosol liquid water content

Y. X. Bian et al.




Fig. 6. The coloured dots show the variation of the ALWC with corresponding number concentrations of accumulation mode particles, with the colour segregated by the RH stages. The dashed lines represent the fitting results in accumulation mode under varying RH conditions.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)