Modelling non-equilibrium secondary organic aerosol
 formation and evaporation with the aerosol dynamics, gas and particle-phase chemistry kinetic multi-layer model
 ADCHAM

5

```
    P. Roldin<sup>1,3</sup>, A. C. Eriksson<sup>1</sup>, E. Z. Nordin<sup>2</sup>, E. Hermansson<sup>1</sup>, D. Mogensen<sup>3</sup>, A.
    Rusanen<sup>3</sup>, M. Boy<sup>3</sup>, E. Swietlicki<sup>1</sup>, B. Svenningsson<sup>1</sup>, A. Zelenyuk<sup>4</sup>, J. Pagels<sup>2</sup>
```

9 [1]{Division of Nuclear Physics, Lund University, P.O. Box 118 SE-221 00 Lund, Sweden}

[2]{Ergonomics and Aerosol Technology, Lund University, P.O. Box 118 SE-221 00 Lund,
Sweden}

12 [3]{Department of Physics, P.O. Box 48, University of Helsinki, 00014, Finland}

[4]{Pacific Northwest National Laboratory, P.O. Box 999, MSIN K8-88, Richland, WA 99354
USA }

15 Correspondence to: P. Roldin (pontus.roldin@nuclear.lu.se)

16

17 Abstract

We have developed the novel Aerosol Dynamics, gas- and particle- phase chemistry model for 18 19 laboratory CHAMber studies (ADCHAM). The model combines the detailed gas phase Master Chemical Mechanism version 3.2, an aerosol dynamics and particle phase chemistry module 20 21 (which considers acid catalysed oligomerization, heterogeneous oxidation reactions in the particle phase and non-ideal interactions between organic compounds, water and inorganic ions) and a 22 kinetic multilayer module for diffusion limited transport of compounds between the gas phase, 23 24 particle surface and particle bulk phase. In this article we describe and use ADCHAM to study: (1) the evaporation of liquid dioctyl phthalate (DOP) particles, (2) the slow and almost particle 25 size independent evaporation of α -pinene secondary organic aerosol (SOA) particles, (3) the mass 26 27 transfer limited uptake of ammonia (NH₃) and formation of organic salts between ammonium (NH_4^+) and carboxylic acids (RCOOH), and (4) the influence of chamber wall effects on the 28 observed SOA formation in smog chambers. 29

1 ADCHAM is able to capture the observed α -pinene SOA mass increase in the presence of 2 NH₃(g). Organic salts of ammonium and carboxylic acids predominantly form during the early 3 stage of SOA formation. These salts contribute substantially to the initial growth of the 4 homogeneously nucleated particles.

5 The model simulations of evaporating α -pinene SOA particles support the recent experimental 6 findings that these particles have a semi-solid tar like amorphous phase state. ADCHAM is able 7 to reproduce the main features of the observed slow evaporation rates if the concentration of low-8 volatility and viscous oligomerized SOA material at the particle surface increases upon 9 evaporation. The evaporation rate is mainly governed by the reversible decomposition of 10 oligomers back to monomers.

Finally, we demonstrate that the mass transfer limited uptake of condensable organic compounds onto wall deposited particles or directly onto the Teflon chamber walls of smog chambers can have profound influence on the observed SOA formation. During the early stage of the SOA formation the wall deposited particles and walls themselves serve as a SOA sink from the air to the walls. However, at the end of smog chamber experiments the semi-volatile SOA material may start to evaporate from the chamber walls.

With these three model applications, we demonstrate that several poorly quantified processes, i.e. mass transport limitations within the particle phase, oligomerization, heterogeneous oxidation, organic salt formation, and chamber wall effects can have substantial influence on the SOA formation, lifetime, chemical and physical particle properties, and their evolution. In order to constrain the uncertainties related to these processes, future experiments are needed where as many of the influential variables as possible are varied. ADCHAM can be a valuable model tool in the design and analysis of such experiments.

24

25 1 Introduction

Aerosol particles in the atmosphere have substantial impact on the global climate, air quality, and public health. Measurements around the world have demonstrated that a large fraction of the submicron aerosol particles are composed of organic compounds (Jimenez et al., 2009). Today many important biogenic and anthropogenic secondary organic aerosol (SOA) precursors have been identified. However, the scientific knowledge about the SOA formation mechanisms, the
SOA composition and properties is still very uncertain (Kroll and Seinfeld, 2008 and Hallquist et
al., 2009).

Traditionally, the important SOA formation mechanisms are modelled as pure gas phase 4 oxidation processes followed by equilibrium partitioning between the gas and a liquid organic 5 particle phase (e.g. Pankow, 1994 and Donahue et al., 2011). However, during the last ~10 years 6 other processes occurring in the particle phase have also been identified as important mechanisms 7 8 for the formation and properties of SOA. These include acid catalysed oligomerization (e.g. Gao et al., 2004, Iinuma et al., 2004, Kalberer et al., 2004, and Tolocka et al., 2004), heterogeneous 9 oxidation reactions (e.g. Knopf et al., 2005, Nash et al., 2006, Rudich et al., 2007, Maksymiuk et 10 al., 2009), organic salt formation (e.g. Na et al., 2007, Smith et al., 2010, Kuwata and Martin, 11 2012 and Yli-Juuti et al., 2013), organosulphate formation (e.g. Liggio and Li, 2006, Surratt et 12 al., 2007) and salting-out effects (e.g. Smith et al., 2011, Bertram et al., 2011). The term salting-13 out refers to the process in which interactions with dissolved ions (generally inorganic) drive 14 15 nonpolar organic compounds out of the mixed phase, either into a different organic-rich (liquid) phase or out to the gas phase (Zuend et al., 2011). 16

Several independent laboratory experiments have also shown that secondary organic aerosol 17 particles can form a solid or semi-solid amorphous phase (e.g. Virtanen et al., 2010, Vaden et al., 18 2010, Vaden et al., 2011, Kuwata and Martin, 2012, Zelenyuk et al., 2012, Abramson et al., 2013 19 and Zhou et al., 2013), at least for relative humidities (RH) below 65 % (Saukko et al., 2012). 20 21 Recently, Abramson et al. (2013) measured the evaporation rates of pyrene that was imbedded inside SOA particles formed from α -pinene ozonolysis in the presence of pyrene vapour, based 22 on which the authors estimated a diffusion coefficient of 2.5×10^{-17} cm² s⁻¹ for pyrene in the fresh 23 SOA, at dry conditions. For particles aged for ~24 hours the diffusivity was an additional ~3 24 times slower. Using the Stokes-Einstein relation for the binary diffusion coefficients gives a 25 SOA viscosity of $\sim 10^8$ Pa s for fresh SOA and $\sim 3 \times 10^8$ Pa s for the aged particles. These high 26 viscosity values are typical for tar or pitch like substances (Koop et al., 2011). For a particle with 27 a diameter of 100 nm, this gives a characteristic time of mass-transport (e-folding time of 28 equilibration) of ~28 hours for fresh SOA and ~84 hours for the aged SOA particles (Seinfeld and 29 Pandis, 2006). A similar study has been performed by Zhou et al. (2013), which observed mass 30

1 transport limited degradation of benzo[a]pyrene (BaP) with ozone, when coated with α -pinene SOA. Based on kinetic double-layer model simulations the authors estimate that the BaP 2 diffusion coefficients (in cm² s⁻¹) are 2 x 10^{-14} , 8 x 10^{-14} and > 10^{-12} for dry (RH < 5 %), 50 % RH 3 and 70 % RH, respectively. The estimated diffusion coefficient of BaP in α-pinene SOA (at dry 4 conditions) from Zhou et al. (2013) is ~3 orders of magnitude larger than the diffusion coefficient 5 of pyrene, estimated by Abramson et al. (2013). One reason for this could be that the α -pinene 6 SOA in Zhou et al. (2013) is very fresh (~1 minute) while in Abramson et al. (2013) the SOA 7 particles are aged for at least 1 hour, thus having time to form a substantial fraction of viscous 8 9 oligomers.

10 If a viscous phase is formed, the mixing within the particle bulk will be kinetically limited and the gas to particle partitioning cannot be well represented by an equilibrium process (Pöschl, 11 2011 and Shiraiwa and Seinfeld, 2012), which the traditional partitioning theory assumes 12 (Pankow, 1994). This may not be evident from pure SOA mass formation experiments where the 13 condensable organic compounds are continuously formed by gas phase oxidation of different 14 precursor compounds (see e.g. Odum et al., 1996, Hoffmann et al., 1997, Griffin et al., 1999, Ng 15 et al., 2007, Pathak et al., 2007). However, in the atmosphere the aerosol particles are present 16 with a broad size range and are exposed to more variable concentration, temperature and 17 18 humidity conditions. Hence, atmospheric aerosol particles will never be entirely in equilibrium 19 with the gas phase. Dzepina et al. (2009) showed that their equilibrium partitioning model substantially overestimate the evaporation of SOA in the Mexico City metropolitan area. 20

In well controlled laboratory experiments Grieshop et al. (2007) and Vaden et al. (2011) have 21 illustrated that the evaporation of SOA particles formed from α -pinene ozonolysis is a slow 22 23 process (hours to days). Vaden et al. (2011) showed that this is orders of magnitude slower than expected from the 7-product VBS parameterization from Pathak et al., 2007. This 24 parameterization is based on a large number of smog chamber experiments of α -pinene 25 ozonolysis, which resemble the experiments by Vaden et al. (2011). This VBS lack substantial 26 fraction of low-volatile material. The slow evaporation of SOA can be due to presence of low-27 volatile oligomers in combination with mass transfer limitations and mixing effects (Grieshop et 28 al., 2007 and Vaden et al., 2011). Saleh et al., (2013) did not observe a strong evaporation 29 30 inhibition because of diffusion limitations in the particle phase of α -pinene SOA particles.

However, in Saleh et al., 2013 only ~20 % of the SOA particle mass (corresponding to a few monolayers) was evaporated in their thermodenuder. This early stage evaporation of freshly formed SOA particles may indeed not be strongly diffusion limited because low-volatile bulk-phase oligomers may not have formed a complete monolayer thick surface coverage, which inhibit further evaporation of monomers.

6 Vaden et al. (2011) illustrated that the evaporation of ambient SOA particles are even slower than 7 for the pure α-pinene SOA and better resembles the evaporation of aged α-pinene SOA particles 8 in the presence of different hydrophobic organic compounds. If the ambient SOA particles 9 studied by Vaden et al. (2011) are representative of typical atmospheric SOA particles, the 10 evaporation due to dilution in the atmosphere (e.g. in urban plumes downwind the source) will be 11 almost negligible. This can increase the lifetime and concentrations of SOA (and e.g. NH₄NO₃) in 12 the atmosphere (Vaden et al., 2011).

For the past decade, large discrepancies between field-measured and model-predicted SOA 13 14 loadings stimulated an intense research that was mostly focused on the search for additional SOA precursors. However, these models have all treated SOA assuming it to be semi-volatile 15 equilibrated solution. A recent study (Shrivastava et al. 2013) shows that it is possible to improve 16 17 agreement between measured and modelled SOA loadings by treating SOA, in accord with experimental data as a non-volatile, semi-solid. Similarly, considering the highly viscous, non-18 volatile nature of SOA offers a simple explanation for the observed long-range transport of 19 persistent organic pollutants by atmospheric particles (Zelenyuk et al. 2012). 20

A number of model studies have been performed to explore detailed gas phase reaction 21 22 mechanisms which can be responsible for the SOA formation of known biogenic and 23 anthropogenic SOA precursors (e.g. Bloss et al., 2005a-b, Johnson et al., 2005 and 2006, Li et al., 2007, Hu et al., 2007, Metzger et al., 2008, Rickard et al., 2010, Camredon et al., 2010 and 24 25 Valorso et al., 2011). However, relatively few attempts have been made to perform detailed process-based modelling on the influence of phase state (Shiraiwa et al., 2010, 2011 and 2012, 26 27 Pfrang et al., 2011), oligomerization (Vesterinen et al. 2007, Pun and Seigneur, 2007, Li et al., 2007, Hu et al., 2007, Ervens and Volkamer, 2010), heterogeneous oxidation mechanisms 28 (Shiraiwa et al., 2010, 2011 and 2012, Pfrang et al., 2011), organic-inorganic interactions (e.g. 29 salting-out effects, acidity effects) (Zuend et al. 2010, and Zuend and Seinfeld, 2012), organic 30

salt formation (Barsanti et al., 2009), and non-equilibrium gas-particle partitioning and aerosol
dynamics (e.g. Korhonen, et al. 2004, Vesterinen et al. 2007, Boy et al., 2006, and Roldin et al.,
2011a-b,) on the SOA formation and properties, and to our knowledge no one has previously
included all these processes in the same model.

In this article we describe and apply a newly developed aerosol dynamics and gas- and particle-5 phase chemistry model for chamber studies (ADCHAM). As the name implies the model is 6 primarily aimed to be used as a flexible tool for evaluation and design of controlled experiments 7 in smog chambers (e.g. Nordin et al., 2013), thermo-denuders (e.g. Riipinen et al., 2010), 8 evaporation chambers (e.g. Vaden et al., 2011), flow-tube reactors (e.g. Jonsson et al., 2008) or 9 hygroscopicity measurements set-ups (e.g. Svenningsson et al., 2006). However, the overall aim 10 is to gain better understanding of which processes (e.g. gas phase chemistry, particle phase 11 12 reactions, particle phase state, aerosol water uptake, cloud droplet activation, and aerosol dynamics) are relevant for the aerosol properties and formation in the atmosphere. 13

14 In ADCHAM the secondary aerosol formation is modelled by combining the Master Chemical Mechanism version 3.2 (MCMv3.2) (Jenkin et al., 1997, Jenkin et al., 2003, Saunders et al., 15 2003) and an updated version of the aerosol dynamics and particle phase chemistry module from 16 17 ADCHEM (2D-Lagrangian model for Aerosol Dynamics, gas phase CHEMistry and radiative 18 transfer) (Roldin et al., 2011a), which now considers acid catalysed oligomerization, oxidation reactions in the particle phase (e.g. secondary ozonide formation) and the diffusion limited 19 20 transport of compounds between the gas phase, particle surface and particle bulk phase. In this work we test the capability of ADCHAM to simulate: (1) the particle size dependent mass 21 22 evaporation loss rates of liquid DOP particles, (2) the slow and almost particle size-independent evaporation of α -pinene SOA particles (Vaden et al., 2011), (3) the mass transfer limited uptake 23 24 of NH₃ and formation of organic salts between ammonium and carboxylic acids (Na et al., 2007 and Kuwata and Martin, 2012), and (4) the influence of heterogeneous reactions and chamber 25 26 wall effects on the SOA formation and properties.

27 Regional and global scale chemistry transport models (e.g. the EMEP model (Bergström et al., 28 2012) rely on semi-empirical parameterizations for the SOA formation (e.g. VBS) derived from 29 smog chamber experiments. This is one of many reasons why it is important to constrain the 30 uncertainties related to specific chamber effects. Hence, as a final application, we illustrate how ADCHAM can be used to study the influence of chamber wall effects on the SOA mass
 formation, particle number size distribution and gas phase chemistry during a *m*-xylene oxidation
 experiment from Nordin et al. (2013).

4

5 2 Model description

6 ADCHAM consists of:

- 7 1) a detailed gas phase kinetic code (in this work with reactions from MCMv3.2),
- 8 2) an aerosol dynamics code (Roldin et al., 2011a) which include Brownian coagulation,
 9 homogeneous nucleation, dry deposition to chamber walls and a detailed
 10 condensation/evaporation algorithm,
- 11

12

- a novel particle phase chemistry module which is closely connected to the condensation/ evaporation algorithm and,
- 4) a kinetic multi-layer model which treats the diffusion of compounds between the particle
 surface and several bulk layers, analogous to Shiraiwa et al. (2010).
- Figure 1 shows a schematic picture of the ADCHAM model structure. The model explicitly treats 15 the bulk diffusion of all compounds (including oxidation agents (Z_{ox}) such as OH, O₃ and NO₂) 16 17 between different particle layers and bulk reactions. For all compounds except Z_{0x} the gas-surface partitioning is by default modelled as an absorption (dissolution) process with the 18 condensation/evaporation equation (Eq. 1, Jacobson, 2005a). Equation 1 considers the gas-19 20 surface diffusion limitations and potentially non-unity probability of adsorption (sticking) and dissolution into the particle surface-bulk layer (surface-bulk accommodation). The surface-bulk 21 layer we define as the monolayer thick particle surface layer where the condensing compounds 22 dissolve (absorb). In each particle layer the model considers acid catalysed oligomerization, 23 equilibrium reactions between inorganic and organic salts and their dissolved ions, and 24 25 heterogeneous oxidation of SOA.
- 26 In ADCHAM the different processes are solved with separate modules using operator splitting.
- 27 For each main model time step (in this work 10 s) ADCHAM considers homogeneous nucleation,
- followed by dry deposition of particles (Sect. 2.2.3) and potentially gases, emissions of gases and

particles, gas phase chemistry (Sect. 2.1) and coagulation (Sect. 2.2.2). After this ADCHAM 1 handles the condensation and evaporation of all organic and inorganic compounds (Sect. 2.2.1) 2 and the reversible adsorption, diffusion and reactions of Z_{ox} in the different particle layers (Sect. 3 2.4.2). In-between these processes ADCHAM uses operator splitting with a much shorter time 4 step (in this work 1-10 ms)." For the reversible adsorption, diffusion and reactions of the 5 different oxidation agents in the particle-phase, a kinetic-multilayer model (Sect. 2.4.2) is used. 6 This model consists of a coupled ordinary differential equation system which is solved using the 7 MATLAB ode15s solver with adaptive and error tolerance controlled internal time steps. The 8 gas-particle partitioning relies upon updated activity coefficients (Sect. 2.3.1), hydrogen ion 9 10 concentrations (Sect. 2.3.2), water content, concentrations of inorganic and organic salts (Sect. 2.3.3) and their corresponding anion and cations. Therefore, the gas-particle partitioning is 11 usually the most time demanding process in ADCHAM. Finally, the model considers the 12 diffusion of organic and inorganic compounds between all particle layers (Sect. 2.4.1) and acid 13 catalysed oligomerization (Sect. 2.3.4). 14

15 2.1 Gas phase chemistry

To be able to implement the detailed MCMv3.2 gas phase chemistry together with user specified 16 reactions and reaction rates (e.g. chamber wall effects) in a computationally efficient way in 17 MATLAB, we constructed a program which automatically creates a system of equations which 18 19 can be used to calculate the concentrations of the user specified compounds. The only required 20 input to the program is the MCMv3.2 names of the compounds which can be downloaded at http://mcm.leeds.ac.uk/MCM. The output from the program is a set of coupled ordinary 21 differential equations (one for each compound) and the Jacobian matrix which is used by the 22 23 ode15s solver in MATLAB. The constructed code can either be used as a standalone code for separate gas phase chemistry simulations, or used as a module in the ADCHEM or ADCHAM 24 25 model. The ode15s solver in MATLAB is intended to be used for stiff ordinary differential 26 equation systems. The solver uses an adaptive and error tolerance controlled internal time step in order to solve the gas phase chemistry. 27

In Sect. 3.2 and 3.3 we simulate the SOA formation from ozonolysis of α-pinene in the presence
of CO or cyclohexane as OH scavenger. We constructed an equation system consisting of all
MCMv3.2 reactions involving inorganic gas phase chemistry and all oxidation products of α-

pinene and cyclohexane (in total 668 compounds and 2093 reactions). In Sect. 3.4 we also model
the SOA formation oxidation of *m*-xylene with the MCMv3.2 gas phase chemistry (273
compounds and 878 reactions).

4 2.2 Aerosol dynamics

The aerosol dynamics module in ADCHAM is based on the aerosol dynamics code from the
ADCHEM model (Roldin et al., 2011a). A shorter description with focus on the important
updates is given below.

8 2.2.1 Condensation and evaporation

9 In ADCHAM the gas-particle partitioning depends on the chemical composition in the particle 10 surface-bulk layer. Analogous to Jacobson (2005b) the dissolution of ammonia into the particle 11 surface-bulk layer water- and/or organic phase is treated as an equilibrium process, considered 12 after the diffusion limited condensation/evaporation of HNO₃, H₂SO₄ and organic compounds 13 (Eq. 1, Seinfeld and Pandis, 2006) (of which carboxylic acids influence the particle acidity and 14 hence the ammonia dissolution).

15
$$I_{i} = 2D_{i}D_{p}f_{i}(Kn_{i},\alpha_{s,i})(C_{i,\infty} - C_{i,s}), \quad f_{i}(Kn_{i},\alpha_{s,i}) = \frac{0.75\alpha_{s,i}(1 - Kn_{i})}{Kn_{i}^{2} + Kn_{i} + 0.283Kn_{i}\alpha_{s,i} + 0.75\alpha_{s,i}}$$
(1)

In Eq. (1) I_i is the contributions of species *i* to the particle molar growth rates, f_i is the Fuchs-Sutugin correction factor in the transition region, $C_{i,\infty}$ is the gas phase concentration of species *i* far from the particle surface (mol m⁻³ air), $C_{i,s}$ is the saturation gas phase concentration at the particle surface (mol m⁻³ air), D_i is the gas phase diffusion coefficient (m² s⁻¹), D_p is the particle diameter (m), Kn_i is the non-dimensional Knudsen number and $\alpha_{s,i}$ is the surface-bulk accommodation coefficient.

In this work we estimate the pure-liquid saturation vapour pressures (p_0) of the MCMv3.2 oxidation products using either the group contribution method SIMPOL (Pankow and Asher, 2008) or the method by Nannoolal et al. (2008) (here referred to as the Nannoolal method). The corresponding equilibrium vapour pressures $(p_{s,i,j})$ for each particle size bin (j) are derived with Raoult's law, using the mole fractions of each organic compound $(x_{i,j})$, the activity coefficients 1 $(\gamma_{i,i})$ calculated with the AIOMFAC thermodynamic model (Zuend et al., 2008 and 2011) (Sect.

2 2.3.1), and the Kelvin effect ($C_{ki,j}$) (Eq. 2). The surface tension (σ_i) of the organic compounds 3 were assumed to be 0.05 Nm⁻¹ following Riipinen et al. (2010).

4
$$p_{s,i,j} = p_{0,i} x_{i,j} \gamma_{i,j} C_{ki,j}, \quad C_{ki,j} = e^{\left(\frac{4M_i \sigma_i}{RT \rho_{pD_{p,j}}}\right)}$$
 (2)

5 *T* is the temperature in Kelvin, *R* is the universal gas constant (J mol⁻¹ K⁻¹), M_i is the molar mass 6 of compound *i* and ρ_p is the density of the phase which the compound partition to.

The mole fraction for compound i in Eq. (2) is the mole fraction of the organic compound in the 7 surface-bulk layer organic phase which compound *i* partitions into (dissolves). In this work we 8 9 either treat all SOA (monomers + dimers + organic salts) as one phase or as two completely separate phases, with monomers as one phase, and the dimers and organic salts as a second phase. 10 This phase separation is not modelled explicitly (as in Zuend and Seinfeld, 2012). Instead we 11 simply assume that either the phase separation does occur or it does not. In future model 12 application, we intend to implement a simple approach to calculate liquid-liquid phase separation 13 (e.g. Topping et al., 2013). If the described phase separation occurs, then the monomers will not 14 dissolve in the phase made up of dimers and/or organic salts. Thus, their saturation vapour 15 pressures are not lowered by dimerization or organic salt formation, which result in less SOA 16 mass in the model compared to simulations without phase separation (Topping et al., 2013). 17

In this work the condensation and evaporation mechanism includes all organic compounds with modelled pure-liquid saturation vapour pressures less than 1 Pa. For the α -pinene oxidation experiments which we model in Sect. 3.2 and 3.3 this involves 154 non-radical MCMv3.2 organic compounds, while for the *m*-xylene SOA formation experiment modelled in Sect. 3.4 we consider 112 condensable organic MCMv3.2 compounds.

23 2.2.2 Coagulation

ADCHAM also includes a Brownian coagulation algorithm (Roldin et al., 2011a). The particle mass and number concentrations of the formed particles are split between the existing particle size bins using a full-stationary method. It still remains a challenge to combine the coagulation

algorithm with the kinetic multilayer model, when the number of particle layers depends on the 1 2 particle size. In this first version of ADCHAM it is only possible to treat coagulation between particles composed of maximum three layers (e.g. a surface monolayer layer, a bulk layer and a 3 seed aerosol core). When two particles composed of such a layer structure collide the layers are 4 5 simply assumed to merge together forming a new spherical particle with a surface-bulk layer, a bulk layer and a seed aerosol core. Because the surface area of the formed particle is always less 6 7 than the sum of the surface areas of the two original particles the width of the surface-bulk layer increases. Hence, in order to keep the width of the surface-bulk layer at approximately the 8 9 thickness of one monolayer, part of the surface-bulk layer material is transferred to the particle 10 bulk.

11 **2.2.3 Dry deposition and chamber wall effects**

It is well known that dry deposition losses of particles onto the chamber walls have large 12 influence on many chamber experiments (see e.g. Pierce et al., 2008). A commonly used method 13 (see e.g. Hildebrandt et al., 2009 and Loza et al., 2012) is to scale the measured SOA mass with 14 15 the measured relative seed aerosol (typically ammonium sulphate) loss rate. With this method it is assumed that the particles deposited on the chamber walls continue to take up condensable gas-16 phase compounds as if they were still present in the gas phase. A second method which was also 17 used by Hildebrandt et al. (2009) and Loza et al. (2012) is to assume that once the particles have 18 19 deposited on the chamber walls the gas-particle partitioning stops. These two correction methods 20 can be considered to be two extremes, where the first method gives an upper bound of the SOA 21 mass formed during the experiments while the second method gives a lower bound of the SOA 22 formed during the experiments (at least if the SOA particles on the chamber walls are not 23 evaporating and the gas phase losses directly to the chamber walls are negligible).

ADCHAM considers the dry deposition of particles onto chamber walls and also keeps track of the particles deposited on the walls. The model also treats the mass transfer limited gas to particle partitioning between the gas phase and the wall deposited particles. Hence, ADCHAM can be used to study the influence of chamber wall effects on the SOA mass formation and help constraining the uncertainties of the formed SOA mass (SOA mass yield).

For non-charged particles, ADCHAM uses the indoor dry deposition loss rate model from Lai 1 and Nazaroff (2000) which accounts for different dry deposition loss rates on upward, downward 2 and vertical facing surfaces. However, if a considerable fraction of the particles are charged (e.g. 3 at Boltzmann charge equilibrium) the effective dry deposition loss rate of particles can be 4 5 considerably enhanced (Pierce et al., 2008). Hence, in order to be able to model realistic dry deposition loss rates of charged particles, ADCHAM keeps track of the fraction of particles 6 suspended in the air with zero, one, two or three elemental charges in each particle size bin. The 7 first order deposition loss rate (s⁻¹) due to charge (k_{charge}) is calculated with Eq. (3) where v_e is the 8 characteristic average deposition velocity due to electrostatic forces (m/s) (McMurry and Rader, 9 10 1985). The dry deposition loss rates depend on the friction velocity and for charged particles also on the mean electrical field strength within the chamber (\overline{E}). Unfortunately both of these 11 parameters are usually poorly known and need to be constrained with model simulations of seed 12 aerosol deposition experiments (see Section 3.4). 13

McMurry and Rader, (1985) found that \overline{E} was ~45 V cm⁻¹ in an almost spherical ~0.25 m³ Teflon chamber. On the chamber surfaces they measured a negative electrical field strength of - 300 ± 150 V cm⁻¹. They attributed the lower empirically derived electric field within the chamber to the fact that the particles in the bag will be influenced by a net electrical field, which has contributions from all points on the chamber surfaces. Hence, the shape and size of the chamber will also influence the mean electrical field.

20
$$k_{charge} = \frac{A_{chamber} v_e}{V_{chamber}}, \qquad v_e = \frac{neC_c E}{3\pi\mu D_p}$$
 (3)

21 $A_{chamber}$ is the chamber surface area, $V_{chamber}$ is the chamber volume, *n* is the number of elemental 22 charges of the particle, *e* is the elementary charge, C_c is the Cunningham slip correction factor 23 and μ is the dynamic viscosity of air.

The mass transfer limited uptake of gases to and from the chamber walls need to be considered in order to take into account the potential uptake (dissolution) of organic compounds in the SOA particles deposited on the chamber walls (Hildebrandt et al., 2009), as well as direct uptake of gas phase molecules onto the Teflon chamber surfaces (Matsunaga and Ziemann, 2010). For the condensation uptake or evaporation of SOA from the particles deposited on the walls, we assume

that the particles deposited on the walls behave as if they were still suspended in (direct contact 1 with) a thin (by default 1 mm thick) air layer adjacent to the chamber walls. As more particles get 2 deposited on the walls, the SOA concentration on the chamber wall will increase. The 3 condensable organic compounds in the thin air layer next to the chamber walls then have an 4 increased probability to dissolve into the organic particle phase on the walls. However, semi-5 volatile organic compounds may also evaporate from the particles on the walls, when the gas-6 phase concentrations in the chamber are reduced. The gas-particle partitioning between the wall-7 deposited particles and the thin air layer next to the chamber walls is modelled with the 8 9 condensation and evaporation module described in Sect. 2.2.1.

ADCHAM also considers the adsorption and desorption of condensable organic compounds onto 10 the Teflon surface film. This is modelled as a reversible process in accordance with Matsunaga 11 12 and Ziemann, 2010. The adsorption of gas phase organic compounds onto the chamber walls is represented by a first order loss rate from the near wall gas phase to the walls $(k_{g,w})$. The 13 desorption rate from the Teflon surfaces out to the thin layer next to the chamber walls $(k_{w,g,i})$ 14 depends on the pure-liquid saturation vapour pressures $(p_{0,i})$ of the adsorbed compounds (Eq. 4) 15 (Matsunaga and Ziemann, 2010). Equation (5) and (6) describe the rate of change of the organic 16 17 compound (X_i) (due to adsorption and desorption) on the chamber walls and in the air layer adjacent to the wall, respectively. $[X_{i,g,w}]$ is the concentrations of compound X_i in the thin layer 18 adjacent to the chamber walls. The concentration at the chamber wall $([X_{i,w}])$ is given as an 19 effective chamber volume concentration (total number of X_i molecules on the walls divided by 20 the total chamber volume $(V_{chamber})$). V_{wall} is the air volume of the thin (1 mm) layer adjacent to 21 the chamber walls, C_w is an effective wall equivalent mass concentration which the organic 22 compounds can dissolve into, M_w is the average molar mass of the Teflon film, and $\gamma_{w,i}$ is the 23 24 activity coefficient of compound *i* in the Teflon film.

Because ADCHAM calculates the gas-wall uptake from the thin layer (volume) adjacent to the Teflon walls and not from the total chamber volume, the $k_{g,w}$ values used in the model need to be substantially larger than the (experimentally quantifiable) effective total chamber volume loss rate $k_{g,w}^*$ (e.g. from Matsunaga and Ziemann, 2010). If the mass transport across the laminar layer adjacent to the chamber walls do not pose a strong limitation on the gas-wall uptake ($\Delta x \approx 1$ mm), the $k_{g,w}$ value used in ADCHAM can be derived from the experimentally determined 1 effective loss rate, by scaling $k_{g,w}^*$ with the ratio between the total chamber volume and the air 2 volume adjacent to the chamber walls ($V_{chamber}/V_{wall}$). However, if Δx is relatively large (e.g. 10 3 mm), $k_{g,w}$ need to be increases in order to match the experimentally derived gas-wall losses. In the 4 supplementary material we illustrate this with a few examples. Here we also illustrate that the 5 modelled gas-wall uptake is not sensitive to the absolute width of the thin air layer adjacent to the 6 chamber walls, as long as $k_{g,w}$ is scaled with $V_{chamber}/V_{wall}$ (see Fig. S1).

7 $k_{g,w}^*$ and $C_w / (M_w \gamma_{w,i})$ in Eq. (4) was experimentally determined by Matsunaga and Ziemann 8 (2010) for a 5.9 m³ Teflon chamber. For *n*-alkanes, 1-alkenes, 2-alcohols and 2-ketones 9 $C_w / (M_w \gamma_{w,i})$ was 9, 20, 50 and 120 µmol m⁻³, respectively. $k_{g,w}^*$ varied between 1/3600 s⁻¹ and 10 1/480 s⁻¹ depending the type of compound.

11
$$k_{w,g,i} = \frac{k_{g,w}}{(RT / p_{0,i}C_w / (M_w \gamma_{w,i}))}$$
 (4)

12
$$\frac{d \left[X_{i,w} \right]}{dt} = \left(k_{g,w} \left[X_{i,g,w} \right] - k_{w,g,i} \left[X_{i,w} \right] \right) \frac{V_{wall}}{V_{chamber}}$$
(5)

13
$$\frac{d\left\lfloor X_{i,g,w}\right\rfloor}{dt} = -k_{g,w}\left[X_{i,g,w}\right] + k_{w,g,i}\left[X_{i,w}\right]$$
(6)

According to Eq. (4) a compound with $p_0=2.5 \times 10^{-2}$ Pa and $C_w / (M_w \gamma_{w,i}) = 10 \text{ }\mu\text{mol m}^{-3}$, partitions 14 ~50 % to the gas phase and ~50 % to the chamber walls, at equilibrium and room temperature. At 15 equilibrium, compounds with a vapour pressure $<10^{-3}$ Pa and $C_w / (M_w \gamma_{w,i}) > 10 \mu \text{mol m}^{-3}$ will 16 almost exclusively be found at the walls, if they are not able to form SOA rapidly enough. Hence, 17 the SOA formation in the smog chamber will depend on: (1) the formation rate of condensable 18 organic compounds, (2) the particle deposition losses, (3) the magnitude of the condensation sink 19 to the particles in the air and onto the chamber walls, and (4) the diffusion limited uptake onto the 20 chamber walls and particles on the walls. 21

The concentration gradient in the laminar layer adjacent to the chamber walls generally drives condensable gas phase components from the well mixed chamber volume to the chamber walls (thin model layer next to the wall). We explicitly model this mass transfer with Fick's first law of diffusion, assuming a linear concentration gradient across the laminar layer next to the chamber
wall (see Fig. 2).

In Sect. 3.4 we study how different values of the laminar layer width influence model results. The 3 4 gas phase chemistry and the gas to particle mass transfer (condensation) in the well mixed chamber volume and in the thin layer adjacent to the chamber wall were solved using operator 5 splitting with a model time step of 10 ms. The mass transfer between the well-mixed chamber 6 volume and the thin layer next to the chamber wall was modelled with a time step of 0.1 ms. The 7 8 model needs to take short time steps because of the large condensation sink (or evaporation source) of the wall deposited particles and the Teflon surfaces which may rapidly alter the 9 concentrations in the thin air layer next to the chamber walls. 10

In Sect. 3.2 we evaluate the potential influence of the reversible uptake of organic compounds to
the smog chamber Teflon walls on the volatility and evaporation rates of SOA particles. In Sect.
3.4 we test the capability of ADCHAM to simulate the losses of organic compounds from the
gas-phase to the Teflon walls and wall deposited particles.

15 2.2.4 Size distribution structures

Analogous to ADCHEM (Roldin et al., 2011a) ADCHAM include several methods (full-16 stationary, full-moving and moving-centre) in order to treat the changes in the particle number 17 18 size distribution upon condensation/evaporation or coagulation. These methods are all mass and 19 number conserving and have different advantages and disadvantages (Korhonen, et al. 2004, 20 Jacobson, 2005a and Roldin et al., 2011a). For all simulations performed in this article, we have used the full-moving method for condensation and evaporation. With this method the diameter 21 22 grid moves with the particles. Hence, this method has no numerical diffusion problems when particles grow by condensation or evaporate. Homogeneous nucleation is considered by adding 23 new particle size bins when new particles are formed (Sects. 3.2 and 3.3). For coagulation we use 24 a full-stationary method where the formed particles mass and number concentrations are split 25 between the existing diameter bins (Sect. 2.2.2). 26

27 2.3 Particle phase chemistry

28 2.3.1 Activity coefficients and organic-inorganic interactions

The AIOMFAC model is based on the UNIFAC model for organic mixtures but also considers 1 2 organic-inorganic interaction which allows us to study salt-effects on the SOA formation. AIOMFAC considers interactions between 12 different ions (including NH₄⁺, NO₃⁻, H⁺, SO₄⁻² 3 and HSO₄) and alkyls, hydroxyls, carboxyls, ketones, aldehydes, ethers, esters, alkenyls, 4 5 aromatic carbon-alcohols and aromatic hydrocarbons (Zuend et al., 2008 and 2011). For other important functional groups i.e. nitrates, nitros, PANs and peroxides we only consider organic-6 7 organic functional group interactions. In total the model considers 52 different UNIFAC functional subgroups, with interaction parameters from Hansen et al. (1991), except for alcohols 8 (Marcolli and Peter, 2005) and nitrates, PANs and peroxides for which we use the 9 10 parameterization from Compernolle et al. (2009). In ADCHAM, the activity coefficients are calculated before the condensation algorithm is used and when updating the hydrogen ion 11 concentration ([H⁺]) for the acid catalysed oligomerization. 12

13 **2.3.2** Acidity and dissociation of inorganic compounds in organic rich phases

The hydrogen ion concentration is calculated in the condensation algorithm and when considering acid catalysed oligomerization. Analogous to the procedure in ADCHEM (Roldin et al., 2011a) [H⁺] in the particle water or particle water + organics phase is calculated by solving the ion balance equation (Eq. 7). In ADCHAM we have extended the ion balance equation with dissociation products of carboxylic acids (RCOO⁻). In this work we assume that all carboxylic acids have identical dissociation constants (see Sect. 3.2). Hence, [RCOO⁻] in Eq. (7) represent the total concentration (mol/kg solvent) of dissociated carboxylic acids.

21
$$\begin{bmatrix} H^{+} \end{bmatrix} + \begin{bmatrix} NH_{4}^{+} \end{bmatrix} + \begin{bmatrix} Na^{+} \end{bmatrix} = \begin{bmatrix} NO_{3}^{-} \end{bmatrix} + 2\begin{bmatrix} SO_{4}^{2-} \end{bmatrix} + \begin{bmatrix} HSO_{4}^{-} \end{bmatrix} + \begin{bmatrix} CI^{-} \end{bmatrix} + \begin{bmatrix} OH^{-} \end{bmatrix} + \begin{bmatrix} HCO_{3}^{-} \end{bmatrix} + 2\begin{bmatrix} CO_{3}^{2-} \end{bmatrix} + \begin{bmatrix} RCOO^{-} \end{bmatrix}$$
(7)

In order to calculate $[H^+]$, all concentrations except the hydrogen ion concentration in Eq. (7) are replaced with known equilibrium coefficients, activity coefficients from AIOMFAC, and the total concentration of dissolved dissociated and non-dissociated compounds, ([RCOOH]+[RCOO⁻]), ([NH₃(aq)]+[NH₄⁺]), ([SO₄²⁻] +[HSO₄⁻]), ([HNO₃]+ [NO₃⁻]) and ([HCl(aq)]+ [Cl⁻]). In this work the uptake of CO₂ in the particles was treated as an equilibrium process. The HCO₃⁻ and CO₃²⁻ concentrations depend on the hydrogen ion concentration and the CO₂ partial pressure (390 ppbv). When all unknown ion concentrations have been replaced with the known parameters, Eq. (7) becomes an 8th order polynomial with [H⁺] as the only unknown variable. The hydrogen ion
 concentration is given by the maximum real root of this polynomial.

To treat the CO₂ uptake as an equilibrium process may not be realistic if the particles are very viscous (see Sect. 1). However, the estimated diffusion coefficients of other small "guest" molecules (e.g. O₃, OH and H₂O) in an amorphous glassy organic matrix is in the order of 10⁻¹⁰-10⁻¹² cm² s⁻¹ at room temperature (Koop et al., 2011, Zobrist et al., 2011). This gives corresponding e-folding times of equilibration for submicron particles in the range of seconds.

All experiments which we model in this paper were performed at dry conditions (RH \leq 5 %). For 8 9 the α -pinene SOA experiments (Sect. 3.2-3.3), the modelled particle water mass content is only ~ 0.4 % at a RH of 5 %. For these particles the solvent will therefore mainly be the organic 10 compounds and not water. Hence, in this work the concentrations of the inorganic ions (including 11 12 H⁺) is not given for the aqueous but for the combined organics and water phase. Henry's law coefficients (K_H) of inorganic compounds and dissociation rates (K_a) of inorganic compounds and 13 carboxylic acids, are (if at all) usually only available for aqueous solutions. However, there is 14 often a relationship between the ΔpK_a ($-\log_{10}(K_{a,base}) + \log_{10}(K_{a,acid})$) and the proton transfer 15 between the Brønsted acid and the Brønsted base, in protic ionic liquids (Greaves and 16 17 Drummond, 2008). Thus, for most of the simulations we will use the aqueous dissociation rates and Henry's law coefficients for the organic amorphous SOA and water mixtures, and take into 18 account the non-ideal interactions between the ions, organic solvents and water using AIOMFAC 19 (Sect. 2.3.1). However, we will also test to model the ammonium uptake with 10 times lower 20 Henry's law coefficients, which may be more appropriate for organic solvents. 21

22 With these assumptions in mind, the modelled absolute values of $[H^+]$ should be interpreted with caution. However, we believe that the model can give a realistic representation of the relative 23 influence of different types of dissolved compounds on the particle acidity. For instance, 24 25 carboxylic acids will most likely increase [H⁺] also in an organic rich phase, while dissolved ammonia will decrease [H⁺]. For all other organic compounds except the carboxylic acids, the 26 dissociation rates were assumed to be equal to that of pure water ($pK_a = 14$). Hence, equivalent to 27 aqueous solutions the acidity will mainly be governed by the carboxylic acids and inorganic 28 compounds. 29

1 **2.3.3** Inorganic and organic salt formation

In ADCHAM the inorganic salts (NH₄)₂SO₄, NH₄HSO₄ and NH₄NO₃ and the organic salts of 2 3 ammonium and different carboxylic acids (NH₄RCOO) can be considered to form. All these salts contain NH_4^+ and which of these salts that will be formed depend on the solubility constants, the 4 ammonium concentration, the concentration of the different anions and the activity coefficients. 5 6 Because all these salts contain ammonium the salt which forms first will limit the formation of other salts. In this work, we only simulate experiments performed on pure organic particles or 7 organic particles which take up NH₃(g). Hence, NH₄RCOO(s) was the only (solid) salt which was 8 9 considered to be formed in the particle organics-water phase. The solid salt concentrations are updated iteratively every time step which the condensation/evaporation algorithm is used. 10

11 When updating the NH₄RCOO(s) concentration, ADCHAM starts by estimating the activity coefficients and the hydrogen ion concentration (Eq. 7). After this non-equilibrium NH_4^+ and 12 RCOO⁻ concentrations ($[NH_4^+]^*$ and $[RCOO^-]^*$) can be derived, and the total concentrations of 13 NH_4 ([NH_4 tot]=[NH_4^+]^{*}+[NH_4RCOO]_{t-1}) and RCOO ([$RCOO_{tot}$]=[$RCOO^-$]^{*}+[NH_4RCOO]_{t-1}) are 14 estimated. These values are then inserted into the solubility product equation (Eq. 8). Rearranging 15 Eq. (8) gives a second order polynomial where the new NH₄RCOO concentration ([NH₄RCOO]_t) 16 is given by the smallest positive real root. Finally the NH_4^+ and $RCOO^-$ concentrations are 17 updated and the iteration starts from the beginning by deriving the hydrogen ion concentration 18 again. The iteration proceeds until the relative change in the NH₄⁺, RCOO⁻ and H⁺ concentrations 19 all are less than 10⁻³ between one iteration step. The activity coefficient of the dissociated 20 carboxylic acids (γ_{RCOO}) was assumed to be equal to the activity coefficient of undissociated 21 pinonic acid. 22

23
$$\frac{K_{NH_4RCOO} = \gamma_{NH_4} \gamma_{RCOO} \left[NH_4^+ \right] \left[RCOO^- \right] =}{\gamma_{NH_4} \gamma_{RCOO} \left(\left[NH_{4,tot} \right] - \left[NH_4RCOO \right] \right) \left(RCOO_{tot} - \left[NH_4RCOO \right] \right) \right) (RCOO_{tot} - \left[NH_4RCOO \right] \right)$$
(8)

24 2.3.4 Acid catalysed oligomerization

Any oligomerization mechanisms in the gas phase (g) or particle phase (p) (including different functional groups, ozonolysis, acid catalysed reactions, and radicals), can easily be implemented in ADCHAM. For the applications in this work, we only consider the reactions between

monomers which form dimers and not the possible reactions between dimers and dimers with 1 monomers. The acid catalysed formation rate of a dimer in the particle phase can generally be 2 considered to be proportional to the hydrogen ion concentration (see e.g. schemes by Tolocka et 3 al., 2004 and Iinuma et al., 2004). In contrast to thermodynamic equilibrium models (e.g. 4 5 Barsanti and Pankow, 2004), ADCHAM explicitly treats the kinetics of reversible dimerization with separate reactions (R1-R2) for the formation and the degradation of dimers back to 6 7 monomers. Based on observations several different particle phase oligomerization mechanism have been suggested. This includes: (i) esters formed from reactions between carboxylic acids 8 and alcohols (e.g. Surratt et al., 2006), (ii) hemiacetal formation from reactions between 9 10 carbonyls and alcohols (e.g. linuma et al., 2004), (iii) aldol reaction products formed from carbonyl-carbonyl reactions (Casale et al., 2007) and (iv) peroxyhemiacetals formed from 11 reactions between hydroperoxides and carbonyls (e.g. Tobias and Ziemann, 2000). Dimers can 12 also form when carbonyls react with ozone in the particle phase and form secondary ozonides 13 (R3) which then rapidly react with other organic compounds and form dimers (Maksymiuk et al., 14 15 2009). Based on thermodynamic calculations of different organic mixtures, it has been suggested that ester formation (Barsanti and Pankow, 2006) and peroxyhemiacetal formation (DePalma et 16 al., 2013) can be thermodynamically favourable, while hemiacetal formation is not (Barsanti and 17 Pankow, 2004 and DePalma et al., 2013). 18

If peroxyhemiacetal and hemiacetal formation are thermodynamically favourable these 19 mechanism are probably rapid enough to form substantial dimer mass on short timescales 20 (minutes to hours). The second-order rate constants (k_f) for uncatalysed reactions of different 21 hydroperoxides and aldehydes to form peroxyhemiacetals range from 0.5 to 70 M⁻¹ h⁻¹ (2.3x10⁻²⁵-22 3.2x10⁻²³ molecules⁻¹ cm³ s⁻¹) Ziemann and Atkinson (2012). According to Ziemann and 23 Atkinson (2012) peroxyhemiacetal formation can probably be acid catalysed. Thus, for acid 24 particles, k_f may be larger than the values reported by Ziemann and Atkinson (2012). Shiraiwa et 25 al. (2013a) found that in order for the KM-GAP model to capture the temporal evolutions of the 26 SOA formation and shape of the particle number size distribution during an dodecane 27 photooxidation experiment, the peroxyhemiacetal formation rate need to be 12 M^{-1} s⁻¹ (2x10⁻²⁰ 28 molecules⁻¹ cm³ s⁻¹). This high formation rate can be attributed to the presence of carboxylic 29 30 acids (Shiraiwa et al., 2013a). The formation of hemiacetal from the reaction between acetaldehyde and methanol is acid catalysed with a third-order reaction specific formation rate 31

1 constant $(k_{f,H+})$ equal to $4.9 \times 10^6 \text{ M}^{-2} \text{ h}^{-1}$ (Ziemann and Atkinson, 2012). With a pH of 4, this 2 corresponds to a k_f of 2.3×10^{-22} molecules⁻¹ cm³ s⁻¹.

3 monomer_i + monomer_i
$$\xrightarrow{H^+}$$
 dimer_{ji} (R1)

4
$$\operatorname{dimer}_{j_i} \longrightarrow \operatorname{monomer}_j + \operatorname{monomer}_i$$
 (R2)

5
$$\operatorname{carbonyls}(p) + O_3(p) \longrightarrow \operatorname{secondary ozonides}(p)$$
 (R3)

6 In the gas-phase low-volatile ROOR-type peroxide dimers can form when two peroxy radicals 7 (RO₂) react (e.g. Kroll and Seinfeld, 2008 and Ng et al., 2008) (R4). Recently, Ehn et al., 2014 8 also showed that extremely low-volatile organic compounds (ELVOC) can form during 9 ozonolysis of α -pinene. The authors propose that ELVOC is formed from RO₂ which rapidly take 10 up O₂ after intramolecular hydrogen abstraction, in several steps.

11
$$\operatorname{RO}_{2}(g) + \operatorname{RO}_{2}(g) \longrightarrow \operatorname{ROOR-type peroxides}(g)$$
 (R4)

The acid catalysed dimer formation rates in the particle phase $(F_f(\mathbf{p}))$ between monomer (denoted with index *i* and *j*) depend on the monomer concentrations, k_{f,H^+} (molecules⁻² cm⁶ s⁻¹) and the hydrogen ion concentration (c_{H^+}) (Eq. 9). If the dimerization process is uncatalyzed, the formation rate depends on the monomer concentrations and a second-order rate constant $(k_f$ (molecules⁻¹ cm³ s⁻¹)). The dimer degradation rate $(F_d(\mathbf{p}))$ of a dimer (denoted with index *q*) simply depends on the dimer concentration and a dimer specific first-order degradation reaction rate constant $(k_d(s^{-1}))$ (Eq. 10).

19
$$F_{f,i,j}(\mathbf{p}) = k_{f,H^+}(\mathbf{p})c_{H^+}(\mathbf{p})c_{m,i}(\mathbf{p})c_{m,j}(\mathbf{p})$$
 (9)

20
$$F_{d,q}(\mathbf{p}) = k_d(\mathbf{p})c_{d,q}(\mathbf{p})$$
 (10)

The temporal evolution of the dimer and monomer concentrations (c_d and c_m) in the particle bulk layers and surface-bulk layer are derived with a kinetic model. This code solves a coupled ordinary differential equation system, consisting of one ordinary differential equation for each SOA monomer (Eq. 11), and one ordinary differential for each dimer (Eq. 12). The equations are given by the sum of all dimer degradation and formation rates for the individual reactions which each monomer compound is involved in.

$$1 \qquad \frac{dc_{m,i}}{dt} = \sum_{q=1}^{Z} (F_{d,q} x_{m,i,q}) - \sum_{j=1}^{N} F_{f,i,j}$$
(11)

2
$$\frac{dc_{d,q}}{dt} = -F_{d,q} + \sum_{i=1}^{M} (\sum_{j=1}^{N} F_{f,i,j}) x_d$$
 (12)

3
$$x_{m,i,q} = \sum_{j=1}^{N} F_{f,i,j} / \sum_{i=1}^{M} (\sum_{j=1}^{N} F_{f,i,j})$$
 (13)

In order to not end up with an enormous coupled ordinary differential equation system, the 4 5 different dimers are classified into different categories (types) depending on the dimerization 6 mechanism (e.g. esters or peroxyhemiacetal formation). Secondary ozonide formation (R3) is treated by the kinetic multilayer module for heterogeneous oxidation (see Sect. 2.4.2). In order to 7 8 be mass conserving the number of moles of dimer formed is corrected with the molar ratio (x_d) 9 between the molar mass of the product(s) and the sum of the molar masses of the reacting 10 compounds. In ADCHAM all dimers have by default a molar mass of 400 g/mol. When we lump 11 the dimers into different categories the information about their exact chemical composition and 12 origin are lost. This can be a problem when considering the reversible reactions back to 13 monomers. In this work we have assumed that a dimer is converted back to the monomers it is formed from, with fractions (x_m) corresponding to the (current time step) relative contribution of 14 15 each monomer to the dimer formation (Eq. 13). This can be a reasonable assumption if the 16 monomer SOA composition does not change substantially on a time scale longer than the lifetime of the dimer. However, if this is not the case it can distort the modelled particle composition. 17 18 With this method we do not take into account that specific compounds of the same type of dimer (e.g. esters) may have substantially different formation and degradation rates, e.g. depending on 19 other functional groups in the molecules (Ziemann and Atkinson, 2012). However, in principle 20 21 the method still enables unlimited number of different dimer types and dimer specific formation 22 and degradation rates.

The modelled relative amount and composition of oligomer SOA in each particle layer depends on: (1) the monomer SOA composition, (2) the hydrogen ion concentration, (3) the dimer formation rates, (4) the oligomer degradation reaction rates, (5) possible evaporation and condensation of monomers and dimers (vapour pressures), (6) the mixing between different particle layers (diffusion coefficients of monomers and dimers), (7) the ozone uptake at the particle surface, (8) the ozone diffusion rate within the particle bulk phase, (9) the reaction rates of ozone with unsaturated organic compounds in the particle phase, and (10) the time of aging. Points 1-9 all bear large uncertainties, which need to be constrained in order to represent oligomerization in an accurate way. For the model applications in this work we will not explicitly model the influence of particle acidity on the dimer formation rates because these values are anyhow very uncertain.

8 2.4 Kinetic multi-layer model

9 To be able to model the diffusion limited mass transfer of ozone from the gas-particle interface to the particle core, and the reaction between ozone and the organic compounds in the particle 10 11 phase, Shiraiwa et al. (2010) developed the kinetic multilayer model KM-SUB which is based on the PRA concept of gas-particle interactions (Pöschl-Rudich-Ammann, 2007 and Ammann and 12 Pöschl, 2007). This model divides the particles into a sorption layer, a quasi-static surface layer, 13 near-surface bulk, and multiple bulk layers and considers the gas-surface transport, reversible 14 adsorption, surface layer reactions, surface-bulk transport, bulk diffusion and bulk reactions. 15 Recently, Shiraiwa et al. extended the kinetic multilayer model to also include condensation, 16 evaporation and heat transfer (KM-GAP) (Shiraiwa et al., 2012), thermodynamics (Shiraiwa et 17 al., 2013b) and simplified gas- and particle-phase chemistry (Shiraiwa et al., 2013a). 18

19 For the oxidation agents we model the uptake to the sorption layer as a reversible adsorption processes, followed by diffusion to and from the particle surface-bulk and bulk-layers. In this 20 21 work, the partitioning of organic and inorganic compounds to and from the monolayer thick particle surface-bulk layer is modelled as a condensation/evaporation process (Eq. 1), taking into 22 23 account the possibility of non-unity surface-bulk accommodation (Sect. 2.2.1). However, in principle ADCHAM could also model the gas-particle partitioning of organic and inorganic 24 compounds as a reversible adsorption process. Analogous to KM-SUB and KM-GAP, ADCHAM 25 explicitly treats the bulk diffusion of all compounds between the different layers using first-order 26 27 mass transport rate equations.

The kinetic multilayer model in ADCHAM consists of two separate modules. The first module (Sect. 2.4.1) treats the diffusion of all organic and inorganic compounds (except Z_{ox}) between the different bulk layers. The second module (Sect. 2.4.2) considers the uptake, diffusion and reactions of Z_{ox} with the organic compounds in the particle phase. The main reason why these processes are treated by two separate modules is that the uptake, diffusion and reaction of O_3 and other oxidation agents generally occur on substantially shorter timescales than the diffusion of the organic compounds.

6 2.4.1 Diffusion of organic and inorganic compounds

7 The transport velocity of compound X_i between the bulk layers or the surface and first-bulk layer 8 is given by Eq. (14). D_{X_i} is the diffusion coefficient of compound X_i, and δ_k and δ_{k+1} represent 9 the width of the two adjacent layers (*k* and *k*+1) which X_i is transported between.

10
$$k_{k+1,k,X_i} = k_{k,k+1,X_i} = \frac{4D_{X_i}}{\pi \left(\delta_{k+1} + \delta_k\right)/2}$$
 (14)

11

12 The transport of compound X_i between the particle layers (including the exchange between the surface and first-bulk layer) is modelled with Eq. (15). A_k is the area of exchange between layer 13 k-1 and k. $\left[V_{X_i}\right]_k$ is the absolute volume concentration (m³) and $\left[X_i\right]_k$ is the relative volume 14 concentration (volume fraction) of compound X_i in layer k. The total volume of each particle 15 layer (V_k) is given by the sum of the absolute volume concentrations of all compounds. Equation 16 16 contains no terms for the chemical formation and degradation of X_i because this is considered 17 by separate modules for oligomerization (Sect. 2.3.4), heterogeneous oxidation (Sect. 2.4.2) and 18 organic salt formation (Sect. 2.3.3). 19

20
$$\frac{d\left[V_{\mathbf{X}_{i}}\right]_{k}}{dt} = (k_{k-1,k}\left[\mathbf{X}_{i}\right]_{k-1} - k_{k,k-1}\left[\mathbf{X}_{i}\right]_{k})A_{k} + \left(-k_{k,k+1}\left[\mathbf{X}_{i}\right]_{k} + k_{k+1,k}\left[\mathbf{X}_{i}\right]_{k+1}\right)A_{k+1}$$
(15)

The equations describing the concentration change of all compounds in all layers (Eq. 15) comprise a system of $N \ge N_L$ coupled ordinary differential equations (N_L =number of particle layers) which we solve with the ode15s solver in MATLAB.

Figure 3 shows a schematic picture of the kinetic multilayer module in ADCHAM. In contrast to the kinetic multilayer model by Shiraiwa et al. (2010, 2012) the number of particle layers increases when the particles grow. Hence, particles of different sizes are composed of different
 number of layers.

Once the depth of the surface-bulk layer becomes larger than 1.1 nm, material is moved from this layer to the first bulk-layer, leaving a 1 nm thick surface-bulk layer. If the first bulk-layer becomes larger than a certain value (by default 3 nm thick) it is split into a first and second bulk layer with identical compositions, 1 and 2 nm thick, respectively.

7 Upon evaporation material is lost from the surface-bulk layer and if the layer thickness becomes 8 less than 0.99 nm, material is moved from the first bulk layer to the surface-bulk layer, to keep 9 the surface-bulk layer width intact. If the first bulk layer width becomes less than a certain value 10 (by default 0.8 nm), this layer is merged together with the second bulk layer and together they 11 form a new first bulk layer. The rest of the particle bulk is divided into layers with variable width, 12 which depends on the net mass transport to the adjacent layers and chemical reactions (e.g. 13 between O_3 and unsaturated hydrocarbons).

In each particle layer the model considers different oligomerization reactions and the equilibrium 14 reactions between salts and their dissolved ions in the organic + water phase (Sect. 2.3.3 and 15 2.3.4). The formed oligomers and salts make up a particle volume fraction (f_p) with generally 16 substantially lower diffusivity than the rest of the compounds. Hence, oligomerization and solid 17 salt formation increases the viscosity which also limits the diffusion of the liquid compounds 18 19 according to the obstruction theory (Stroeve, 1975). The treatment of the oligomers as solid nondiffusing compounds which limits diffusion of the liquid compounds were adopted from Pfrang 20 21 et al. (2011) which used KM-SUB to model degradation of an organic 12-component mixture with ozone. 22

According to the obstruction theory the diffusivity of compound X_i (D_{Xi}) is a function of the fraction of solid or semi-solid material and the diffusivity ($D_{0,Xi}$) without any solid or semi-solid material (Eq. 16). The diffusivity of organic compounds can vary from ~10⁻⁵ cm² s⁻¹ in a liquid to ~ 10⁻²⁰ cm² s⁻¹ in a solid organic matrix (Shiraiwa et al., 2011).

27
$$D_{\rm Xi} = D_{0,\rm Xi} \left(2 - 2f_p\right) / \left(2 + f_p\right)$$
 (16)

24

1 2.4.2 Diffusion of oxidation agents and reactions with SOA

The diffusion of oxidation agents (Z_{ox}) between the particle bulk layers is similar to the treatment of other compounds (Eq. 14 and 15), except that we do not consider that the dissolved Z_{ox} in the particle phase take up a bulk volume of its own.

5 The uptake of Z_{ox} from the gas phase to the particle surface is treated as a reversible adsorption 6 process (Fig. 3). This approach was adopted from Shiraiwa et al. (2010). The surface 7 accommodation coefficient of Z_{ox} is given by Eq. (17). $\theta_{s,Z_{ox}}$ is the relative coverage of adsorbed 8 Z_{ox} on the particle surface. The adsorption of Z_{ox} from the near surface gas phase (*gs*) to the 9 sorption layer (*so*) and the desorption from the sorption layer to the near surface-gas phase is 10 given by Eq. (18) and (19), respectively. $\omega_{Z_{ox}}$ is the mean thermal velocity of Z_{ox} and $\tau_{d,Z_{ox}}$ is the 11 desorption lifetime of Z_{ox} .

12 The transport velocity of Z_{ox} from the surface-bulk layer to the sorption layer is given by Eq. 13 (20). δ_s is the width of the monolayer thick surface-bulk layer and $d_{Z_{ox}}$ is the width of the 14 sorption layer. Hence, $(\delta_s + d_{Z_{ox}})/2$ in Eq. (20) represents the average travel distance between 15 the sorption and surface-bulk layer. The transport velocity of Z_{ox} from the sorption layer to the 16 surface-bulk layer can then be calculated from Eq. (21). $K_{H,Z_{ox}}$ is the Henry's law coefficient of 17 Z_{ox} .

18
$$\alpha_{s,Z_{ox}} = \alpha_{s,0,Z_{ox}} \left(1 - \theta_{s,Z_{ox}} \right), \quad \theta_{s,Z_{ox}} = \frac{\left[Z_{ox} \right]_{so} d_{Z_{ox}}^2 \pi}{4}$$
 (17)

$$19 J_{ads,Z_{ox}} = [Z_{ox}]_{gs} \omega_{Z_{ox}} \alpha_{s,Z_{ox}} / 4 (18)$$

20
$$J_{des,Z_{ox}} = \tau_{d,Z_{ox}}^{-1} [Z_{ox}]_{so}$$
 (19)

21
$$k_{su,so,Z_{ox}} \approx \frac{4D_{Z_{ox}}}{\pi \left(\delta_s + d_{Z_{ox}}\right)/2}$$
(20)

$$k_{so,su,Z_{ox}} = \frac{4k_{su,so,Z_{ox}}K_{H,Z_{ox}}\tau_{d,Z_{ox}}^{-1}}{\alpha_{s,Z_{ox}}\omega_{Z_{ox}}}TR$$
(21)

Equations (22-24) form a differential equation system which describes the rate of change of the 1 Zox concentration in the particle sorption layer, particle surface-bulk layer, and particle bulk 2 layers. The chemical oxidation reactions between Z_{ox} and the organic compounds (X_i) are 3 represented by the last term in Eq. (23) and Eq. (24), where the summation is over all compounds 4 5 which react and consume Zox in the particle phase. The module also calculates the temporal evolution of the organic compounds (X_i) which are consumed by Z_{ox} and the organic compounds 6 7 which are formed from the oxidation reactions (Yi) (Eq. 25 and 26). The diffusion of these compounds is treated by the kinetic multilayer module described in Sect. 2.4.1. 8

9
$$\frac{d[Z_{ox}]_{so}}{dt} = J_{ads,Z_{ox}} - J_{des,Z_{ox}} - k_{so,su,Z_{ox}} [Z_{ox}]_{so} + k_{su,so,Z_{ox}} [Z_{ox}]_{su}$$
(22)

11

$$\frac{d[Z_{ox}]_{su}}{dt} = (k_{so,su}[Z_{ox}]_{so} - k_{su,so}[Z_{ox}]_{su})\frac{A_{su}}{V_{su}} +$$

$$(-k_{su,b1}[Z_{ox}]_{su} + k_{b1,su}[Z_{ox}]_{b1})\frac{A_{b1}}{V_{su}} - \sum_{i=1}^{N} k_{Ox,i}[X_{i}]_{su}[Z_{ox}]_{su}$$
(23)

$$\frac{d[Z_{ox}]_{bk}}{dt} = (k_{bk-1,bk}[Z_{ox}]_{bk-1} - k_{bk,bk-1}[Z_{ox}]_{bk})\frac{A_{bk}}{V_{bk}} +$$
(24)

$$(-k_{bk,bk+1}[Z_{ox}]_{bk} + k_{bk+1,bk}[Z_{ox}]_{bk+1}) \frac{A_{bk+1}}{V_{bk}} - \sum_{i=1}^{N} k_{Ox,i}[X_i]_{bk}[Z_{ox}]_{bk}$$

12
$$\frac{d\left[\mathbf{X}_{i}\right]_{bk}}{dt} = -k_{Ox,i}\left[\mathbf{X}_{i}\right]_{bk}\left[Z_{ox}\right]_{bk}$$
(25)

13
$$\frac{d\left[\mathbf{Y}_{i}\right]_{bk}}{dt} = k_{Ox,i} \left[\mathbf{X}_{i}\right]_{bk} \left[Z_{ox}\right]_{bk}$$
(26)

Table 1 gives the values of different parameters used in the multilayer module for ozone uptake,
diffusion and reactions within the particle phase. Most of the values were adopted from Table 1
in Pfrang et al. (2011).

17 The coupled ordinary differential equation system describing the temporal evolution of Z_{ox} and 18 the concentration of compounds which are consumed or formed from the Z_{ox} oxidation is solved 19 with the ode15s solver in MATLAB.

20

1 3 Model applications

In order to test and illustrate the capability of ADCHAM we apply the model to four types of 2 3 published experimental results. In Sect. 3.1 we model the evaporation experiments of liquid dioctyl phthalate (DOP) particles presented in Vaden et al. (2011), which have been modelled by 4 Shiraiwa et al. (2012) with the KM-GAP model. In Sect. 3.2 we model the evaporation 5 experiments of α -pinene SOA particles by Vaden et al. (2011). In Sect. 3.3 we model the SOA 6 formation, ammonia uptake, and organic salt (NH₄RCOO) formation in the α -pinene - NH₃ - O₃ 7 experiments by Na et al. (2007). Finally, we apply ADCHAM to a *m*-xylene oxidation 8 9 experiment from Nordin et al., 2013 (Sect. 3.4). These examples serve to illustrate the wide applicability of ADCHAM. 10

11 For the simulations in Sect. 3.2-3.3 we model the condensational growth of particles formed by homogeneous nucleation using the condensation module described in Sect. 2.2.1 using the full 12 moving method (see Sect. 2.2.4). We start with one particle size and add new particle size bins 13 14 during the early stage of particle formation. The new particles are assumed to be composed of non-volatile SOA material and are introduced into the model at an initial diameter of 5 nm. 15 Hence, in this work we do not treat the initial activation and growth of the formed molecular 16 clusters. The new particle formation rate (J_{5nm}) is assumed to be constant during the experiments. 17 A new size bin is added for the time step when the smallest particle size grows larger than 10 nm 18 19 in diameter. For the experiments which we simulate in this work the SOA mass (condensation 20 sink) increases rapidly during the early stage of SOA formation. This effectively prevents the newly formed particles from growing and thus generally keeps the number of model particle size 21 22 bins down to ~20 (see Fig. S2 in the supplementary material).

Table 2 summarizes the different processes and range of parameter values used for the different
simulations in Sect. 3.1-3.4.

25 3.1 Simulations of DOP particle evaporation

Before modelling complex multicomponent SOA particle formation, growth and evaporation we test ADCHAM on the evaporation experiments of single component, liquid DOP particles (Vaden et al., 2011). The particles, in that study, were evaporated in a 7 L chamber with 1 L of activated charcoal at the bottom of the chamber. The particle number concentration was kept low (~150 cm⁻³) in order to keep the gas phase concentration close to zero. Before the aerosol was
introduced into the chamber, it was passed through two charcoal denuders in order to remove
most of the gas phase DOP (Vaden et al., 2011).

Here we adopt the approach from Shiraiwa et al. (2012) who modelled the gas phase loss to the 4 charcoal denuder using Fick's first law, on a laminar layer (Δx) adjacent to the charcoal denuder, 5 on the bottom of the chamber. Since the layer thickness is poorly known, we modelled the 6 DOP(g) loss rate using different Δx . Coagulation and particle wall losses were not considered. In 7 this small chamber, the wall losses can be substantial, however particles deposited on the 8 chamber walls not coated with charcoal will likely continue to evaporate and contribute to the gas 9 phase DOP. Neglecting the particle wall losses has the same effect as assuming that the particles 10 deposited on the walls continue to take up vapours as if they were still suspended in the air (Sect. 11 2.2.3). 12

Vaden et al. (2011) and Shiraiwa et al. (2012) used a binary diffusion coefficient for DOP in air of 4.4 x 10^{-2} cm² s⁻¹ from Ray et al. (1988). This value was measured at a pressure of 98 Torr (0.13 atm), which is lower than the pressure used in experiments. We have therefore estimated the diffusion coefficient (D_{DOP}) with Eq. (27) (Jacobson, 2005a) and with Eq. (28) (Chapman and Cowling, 1970, in accordance with Zhang et al., 1993). Equation (27) gives a D_{DOP} of 1.5 x 10^{-2} cm² s⁻¹, while with the Chapman-Enskog theory, utilizing a value of 1.34 for the collision integral ($\Omega_{DOP,air}^{(1,1)}$) (Hirschfelder et al., 1954), yields 2.9 x 10^{-2} cm² s⁻¹, for D_{DOP} at 1 atm and 296 K.

$$20 D_i = \frac{5}{16N_a d_i^2 \rho_{air}} \sqrt{\frac{RTM_{air}}{2\pi} \left(\frac{M_i + M_{air}}{M_i}\right)} (27)$$

21
$$D_i = \frac{3}{8\pi \Omega_{i,air}^{(1,1)} p d_{i,air}^2} \sqrt{\frac{\pi k_b^3 T^3 (m_i + m_{air})}{2m_i m_{air}}}, \quad d_{i,air} = \frac{d_i + d_{air}}{2}$$
 (28)

22 N_a is Avogadro's number, ρ_{air} is the density of air, M_{air} is the molar mass of air, M_i is the molar 23 mass of compound *i* (M_{DOP} =390.56 g mol⁻¹), d_i is the collision diameter of compound *i* 24 (d_{DOP} =1.012 nm (Ray et al., 1979)), $d_{i,air}$ is the collision diameter for binary collisions between 25 compound *i* and air molecules (d_{air} =0.362 nm), m_{air} is the molecular mass of air, m_i is the 26 molecular mass of compound *i*, k_b is the Boltzmann constant and *p* is the total pressure. 1 When we use Eq. (27), a laminar layer of 0.1 cm adjacent to the charcoal denuder wall and unity 2 $\alpha_{s,DOP}$, the model is in good agreement with the observed evaporation rates for all particle sizes. 3 Similar results are also achieved when using Eq. (28), unity $\alpha_{s,DOP}$ and a laminar layer of 0.6 cm 4 adjacent to the charcoal denuder (Fig. 4).

In section 3.2 we compare the modelled and measured α -pinene SOA evaporation rates using the same evaporation chamber. Based on the DOP evaporation experiments the simulations of the α pinene SOA particle experiments were performed with a Δx of 0.1 cm, binary diffusion coefficients calculated with Eq. (27), and unity surface-bulk accommodation coefficients.

9 3.2 Evaporation of α-pinene SOA

Here we use ADCHAM to explore which processes are responsible for the slow and nearly size 10 independent evaporation loss rates of α -pinene SOA particles observed by Vaden et al. (2011). α -11 pinene SOA particles were produced by homogeneous nucleation in a 0.1 m³ Teflon chamber 12 under dark conditions with ~200 ppb α -pinene, ~250 ppm cyclohexane as OH-scavenger and 13 ~500 ppb O₃. Once SOA particles stopped growing (approximately after 1.5 hours, fresh 14 particles), monodisperse aerosol particles were selected with a differential mobility analyser 15 (DMA), passed through two charcoal denuders (residence time ~2 minutes), and introduced at 16 low concentration ($\sim 10-200$ cm⁻³) into the evaporation chamber described in Sect. 3.1 (Vaden et 17 al., 2011). Alternatively, the particles were aged for 10-15 hours (aged particles) in the Teflon 18 chamber before being transferred into the evaporation chamber. 19

Vaden et al. (2011) showed that the evaporation rate of the pure α -pinene SOA particles is more 20 than 100 times slower than expected from modelled evaporation rates of liquid-like monomer 21 SOA, and that it consists of two stages. ~50 % of the particle mass evaporates during the first 100 22 23 min at relatively slow rate, followed by a second stage with even slower mass loss rate, in which additional ~25% of the initial mass is lost in 24 hours. Another interesting finding is that the 24 fractional volume loss by evaporation is almost size-independent. Vaden et al. (2011) concluded 25 26 that the nearly size-independent evaporation loss rates indicate that these type of SOA particles are not liquid-like, which later was verified with measurements by Abramson et al. (2013). 27

Here we use ADCHAM to examine how the processes listed below influence the α -pinene SOA 1 evaporation rates. Note that while the model includes various specific mechanisms, the 2 conclusions should be taken in terms that are more general. 3

1) Vapour pressures of the condensable monomers (pure-liquid saturation vapour pressure 4 5

method).

2) Slow and imperfect mixing within semi-solid amorphous SOA particles. 6

3) Dimerization in the particle phase, and their reversible decomposition back to monomers. 7

8 4) Accumulation of low volatility dimers at the particle surface, creating a coating material 9 which prevents the more volatile SOA monomers from evaporating.

5) Wall deposition losses of the α -pinene oxidation products in the Teflon chamber. 10

11 For all simulations presented in this section, the monomer SOA surface-bulk accommodation coefficients were assumed to be unity. The simulations were conducted for 23 °C, RH of 5 %, 12 and a pressure of 1 atm. The laminar layer width adjacent to the charcoal denuder in the 13 14 evaporation chamber was assumed to be 0.1 cm (see motivation in Sect. 3.1). Pure-liquid saturation vapour pressures were estimated with the SIMPOL model, except where otherwise 15 noted. Particles of different sizes were formed by homogeneous nucleation and were allowed to 16 grow in the presence of each other. After 1.5 hours or 12 hours of aging (fresh or aged aerosol) 17 size-selected particles with concentration $\sim 100 \text{ cm}^{-3}$ were introduced into the modelled charcoal 18 denuder chamber and allowed to evaporate by continuous removal of the gas phase compounds. 19 The gas-wall partitioning to the Teflon chamber walls were modelled with an effective gas-wall 20 loss rate $(k_{g,w}^*)$ in the range of 0-1/1000 s⁻¹ and $C_w / (M_w \gamma_{w,i})$ equal to 100 µmol m⁻³ (see Sect. 21 2.2.3). For each model application in Sect. 3.2 we test how sensitive the model results are to the 22 value of $k_{g,w}^{*}$. Particle wall losses were not considered (see discussion in Sect. 3.1 and Sect. 3.4). 23 24 The dimer and monomer SOA compounds were assumed to form one organic phase (no phase separation). 25

3.2.1 Evaporation of pure monomer SOA particles 26

27 Cappa and Wilson, 2011 did not find any substantial differences in chemical composition of αpinene SOA particles upon evaporation in a thermodenuder. Hence, according to this study these 28 particles do not seem to obey absorptive partitioning theory upon evaporation. This could 29

possibly be explained by a diffusion-limited transport of the organic compounds within an 1 2 amorphous (glassy) particle phase (Cappa and Wilson, 2011). However, in a similar study Kuwata et al. (2011) observed a substantial change of the CCN properties of α -pinene SOA 3 particles after termodenuder treatment, which indicates a relative enrichment of low-volatile 4 oligomers after evaporation. In Vaden et al. (2011) they note that the mass spectral peak at 5 m/z=201 rapidly disappears on evaporation and that the only other change is a gradual increase in 6 7 relative intensity of peaks at higher m/z. Thus, this study also suggests an increase in the relative oligomer content, which could indicate that the smaller, higher vapour pressure molecules 8 9 evaporate and oligomerization continues at a slow rate during evaporation, consistent with the 10 observed SOA hardening (Abramson et al. 2013).

11 To set the stage, we start by calculating if the evaporation rates can be explained by the volatility distribution of the condensing monomers formed in the gas phase, in combination with non-12 perfect mixing within a semi-solid amorphous particle phase. The evaporation of the more 13 volatile organic compounds will then be controlled by the evaporation rate of the least volatile 14 organic compounds enriched in the particle surface-bulk layer, and not by their own species 15 16 specific saturation vapour pressures. The measured mass spectra and densities of small and large SOA particles formed by ozonolysis of α -pinene are undistinguishable (Zelenyuk et al., 2008). 17 18 Despite this fact, we use the model to evaluate whether it gives a relative enrichment of the least volatile monomer SOA compounds in the smaller particles during their formation and growth 19 (see e.g. Roldin et al., 2011b), and if this can explain the observed size-independent SOA 20 evaporation. 21

22 The pure-liquid saturation vapour pressures were calculated with the SIMPOL (Pankow and Asher, 2008), Nannoolal et al. (2008) vapour pressure methods or with the semi-empirical 7-23 24 product model (VBS) parameterization from Pathak et al. (2007), which was also used by Vaden et al. (2011). Here we evaluate its influence on the modelled evaporation rates of ~160 nm and 25 ~250 nm particles. The model results presented in Fig. 5 are from simulations with $k_{g,w}^* = 1/2000$ 26 s⁻¹ and liquid-like SOA ($D_{monomer}$ =10⁻¹⁰ cm² s⁻¹) or solid-like SOA particles with negligible mixing 27 $(D_{monomer}=0 \text{ cm}^2 \text{ s}^{-1})$. In Fig. 5a the results are from simulations with the VBS from Pathak et al. 28 (2007), Fig. 5b shows the results when we use SIMPOL and Fig. 5c results from simulations with 29 the Nannoolal method. In Fig. S3 we compare the modelled evaporation losses for simulations 30

with or without reversible gas-wall partitioning onto the smog chamber Teflon walls. The figure illustrates that the uptake of α -pinene oxidation products onto the smog chamber walls lower the volatility of the formed SOA particles. But this does not substantially improve the agreement between the modelled and measured evaporation rates.

In all model runs except with the Nannoolal method and solid-like amorphous particles, the evaporation rates are orders of magnitude faster than the observations. According to the curve fitted to the measurements only ~3 % of the SOA mass is lost during the first 2 minutes. In the model runs 7-80 % are lost, depending on vapour pressure method used, the particle size, the value of $k_{e,w}^*$ and if the SOA is treated as liquid (1) or solid (s) like.

Another difference is that the observed evaporation loss rate is almost linear for the first 30 minutes while in all model runs the loss rate is first very rapid and then gradually slows down. This is because in the model the SOA is composed of molecules with different volatility. Hence, the most volatile molecules are lost early and the remaining compounds that are less volatile evaporate later and slower, inconsistent with observations by Cappa and Wilson, (2011) and Vaden et al. (2011). Moreover, all calculated evaporation rates are size dependent, similarly inconsistent with the observed SOA evaporation (Vaden et al., 2011, Zelenyuk et al., 2012).

17 When the SOA is treated as a solid the evaporation rates are much slower with the Nannoolal method compared to the other two methods, even though most of the other SOA mass (without 18 wall losses) is somewhat more volatile than with the SIMPOL method (see Fig. S4). This is 19 20 mainly because of two low-volatile MCMv3.2 compounds called C922OOH and C813OOH, which before evaporation together make up 10±3 % and 7.5±2 % of the particle mass in the 160 21 nm and 250 nm particles, respectively (see modelled mass spectrum in Fig. S5). These 22 compounds have vapour pressures of 4.8×10^{-8} and 5.3×10^{-8} Pa (at 296 K) when calculated with 23 the Nannoolal method, while according to SIMPOL their vapour pressures are 1.7x10⁻⁶ and 24 1.8x10⁻⁶ Pa (at 296 K). Hence, if the SOA particles are considered to be solid or semi-solid, and 25 the Nannoolal method is used, these compounds accumulate in the particle surface-bulk layer 26 27 upon evaporation and limit the loss of other more volatile compounds.

From the discrepancies between the model and measurement results in Fig. 5 we can conclude that it is unlikely that the observed evaporation rates can be explained purely by incomplete mixing and the vapour pressure controlled evaporation of SOA monomers. We note however,
that when a nearly non-volatile component is introduced and the SOA is treated as solid like, the
evaporation rate significantly decreases.

4 3.2.2 Evaporation governed by mass transport limited mixing and dimer 5 degradation.

Here we are evaluating a hypothesis where dimers comprise a significant fraction (~50%) of the 6 7 particles' mass prior to the transfer of particles into the evaporation chamber (see e.g. Gao et al., 2004). In this case, monomer evaporation dominates the first evaporation stage, which leads to 8 increased dimer concentration in the particle surface-bulk layer (Widmann et al., 1998). The 9 dimers form a low volatile viscous barrier that slows evaporation (modelled with the obstruction 10 theory (Eq. 16)). The dimer SOA is partly mixed by diffusion with the less viscous monomer 11 SOA. The second, slow evaporation stage starts when nearly all monomers are lost and the 12 evaporation rate is determined by the dimer formation/decomposition rates and the transport of 13 the degradation products (monomers) to the surface-bulk layer. 14

In order to test this hypothesis we searched for a possible group of monomer compounds that 15 comprise ~50 % of the SOA mass if they dimerize. Most of the dimers should also form 16 relatively rapidly (within ~1 h), and be relatively long lived ($k_d < 1$ h⁻¹). Peroxyhemiacetal 17 formation has been shown to be thermodynamically favourable (DePalma et al., 2013), and it is 18 probably rapid enough to form substantial dimer mass in the relatively fresh SOA (~1.5 h), (see 19 Sect. 2.3.4). With an equilibrium constants (K_{eq} =[peroxyhemiacetal]/[aldehyde][hydroperoxide]) 20 in the range 0.16-120 M⁻¹ Ziemann and Atkinson (2012) and k_f equal to 10⁻²³ molecules⁻¹ cm³ s⁻¹ 21 the first order degradation rate should be in the range of $1/5 - 40 h^{-1}$. However with a dimer 22 formation rate of 1×10^{-23} molecules⁻¹ cm³ s⁻¹ and decomposition rate of <1 h⁻¹ peroxyhemiacetal 23 dimers contributes to ~80 % of the particle mass. Thus, instead we decided to only consider 24 dimerization between four monomers (C108OOH, C922OOH, C97OOH and C813OOH), which 25 all contain at least one carbonyl and one hydroperoxide functional group. With this assumption 26 the dimer particle content is ~50 %, for particles aged 1.5 h. The dimer mass fraction is nearly the 27 same for all particle sizes (see Fig. S6). Thus, for the results presented in this section we will 28 29 assume that only these four monomers contribute to the dimer formation.

For the diffusion coefficients of monomers and dimers we assume that D_{dimer} are two orders of magnitude smaller than $D_{0,monomer}$, and calculate $D_{monomer}$ with the obstruction theory. The dimers and monomers where assumed to be composed of one well-mixed organic phase.

In order to fit the model to the observed evaporation rates we varied $D_{0,Xi}$ for the monomers and dimers in the range of $1 \times 10^{-16} - 1 \times 10^{-13}$ cm² s⁻¹ and $1 \times 10^{-18} - 1 \times 10^{-15}$ cm² s⁻¹, respectively. With these values of $D_{0,Xi}$ the dimers are enriched in the particle surface-bulk layer upon evaporation, but mass transport limited monomer evaporation across the viscous surface-bulk layer is still possible. The dimer formation and degradation rate was varied in the range of $10^{-22} - 10^{-24}$ molecules⁻¹ cm³ s⁻¹ and 1/20 - 1 h⁻¹, respectively. We also tested to run the model with or without gas-wall partitioning to the smog chamber walls ($k_{g,w}^* = 1/2000$ s⁻¹ or $k_{g,w}^* = 0$ s⁻¹).

With a $D_{0,\text{monomer}}$ of $2 \times 10^{-14} \text{ cm}^2 \text{ s}^{-1}$ in agreement with Zhou et al. (2013), D_{dimer} of $2 \times 10^{-16} \text{ cm}^2 \text{ s}^{-1}$, 11 k_f of 10⁻²³ molecules⁻¹ cm³ s⁻¹, k_d of 1/10 h⁻¹, no phase separation and $k_{g,w}^*$ of 1/2000 s⁻¹ the model 12 reproduce the main features of the observed evaporation behaviour of fresh SOA particles (Fig. 13 6). However, other combination of values of these parameters reproduces the observations 14 equally well (e.g. with $D_{\text{monomer}} \approx 2 \times 10^{-15} \text{ cm}^2 \text{ s}^{-1}$, $D_{\text{dimer}} \approx 2 \times 10^{-15} \text{ cm}^2 \text{ s}^{-1}$, $k_f \approx 10^{-23} \text{ molecules}^{-1}$ 15 cm³ s⁻¹ and $k_d \approx 1/20$ h⁻¹). For aged particles, the model somewhat underestimates the evaporation 16 losses. This is because the relative dimer content in the particles increases with ageing. This 17 18 effect is most pronounced when considering chamber wall losses in the smog chamber (Fig. S6). Thus, when we run the model without reversible gas-wall partitioning to the smog chamber 19 20 Teflon walls the aging effect on the modelled evaporation rates is negligible (Fig. S7).

In the simulations the early evaporation rate is governed by the monomer diffusion rate to the 21 surface. The small particles have a shorter characteristic time of mass-transport than the large 22 23 particles (see Sect. 1). This is the reason why the loss rate during the first hour is somewhat larger for the ~160 nm particles than the ~250 nm particles. When most of the monomers have 24 evaporated (after ~3 hours for the modelled 160 nm particles and ~6 hours for the 250 nm 25 particles) (Fig. S8), the second slow evaporation-stage begins. This stage is determined by dimer 26 degradation, formation and by the diffusion of monomer to the particle surface-bulk layer. Again, 27 28 because of the shorter characteristic time of mass-transport for the small particles, the evaporation losses of the small particles are somewhat larger (steeper slope of the curves in Fig.
 6). This is not completely consistent with the measurements.

From these simulations we can conclude that the model can reproduce the main features of the observed evaporation rates for fresh and aged α -pinene SOA particles, if the reversible gas-wall partitioning in the smog chamber only has a small influence on the particle composition. However, the observed nearly size independent evaporation rates can probably not be explained by an particle phase mass transfer limited evaporation of the monomer SOA, followed by a slow decomposition of the remaining (~50 % by mass) oligomer SOA.

9 3.2.3 Evaporation controlled by the degradation of short- and long-lived dimers in 10 semi-solid tar like SOA particles

Here we examine whether the observed slow evaporation rate can be explained by nearly solid-11 12 like SOA in combination with two types of dimers; the first being relatively short-lived (lifetime of a few minutes) and a second long-lived (lifetime of more than a day). For this paradigm the 13 dimers will accumulate and stay in the particle surface-bulk layer upon evaporation. Thus, the 14 15 size independent evaporation rates will mainly be controlled by the decomposition rate of dimers back to monomers in the surface-bulk layer. We also test if gas-wall losses can contribute to an 16 enrichment of dimers in the particle surface-bulk layer already in the smog chamber, thus helping 17 to explaining the observed relatively slow and size independent first evaporation stage of α -18 pinene SOA. The evaporation is then first controlled by the degradation of the relatively short-19 lived dimers which gradually are replaced by long-lived but less numerous dimers from the 20 particle bulk. 21

In order to test the general mechanism principle, we consider that the dimers are 22 peroxyhemiacetals, which as in Sect. 3.2.2 are formed from the monomers C108OOH, 23 C922OOH, C97OOH and C813OOH. However, the long-lived dimer is only assumed to be 24 25 formed from the least volatile MCMv.3.2 oxidation products C922OOH. With this assumption and because of the Kelvin effect, the relative amount of long-lived dimer increases with 26 decreasing particle sizes (Fig. S9). For the short-lived dimers we varied the values of k_f and k_d in 27 the range of 1×10^{-22} - 1×10^{-24} molecules⁻¹ cm³ s⁻¹ and 30-6 h⁻¹, respectively. For the long-lived 28 dimer we used a k_f of 1×10^{-22} molecules⁻¹ cm³ s⁻¹ and varied the k_d values in the range of 1/20-1/4029

h⁻¹. The monomer SOA was treated as a semi-solid tar like mixture ($D_{0,\text{monomer}}=5 \times 10^{-17} \text{ cm}^2 \text{ s}^{-1}$) according to Abramson et al. (2013) and the dimer SOA as solid ($D_{dimer}=0 \text{ cm}^2 \text{ s}^{-1}$). The gas-wall partitioning was modelled with $k_{g,w}^*$ in the range of 0 - 1/500 s⁻¹ and $C_w / (M_w \gamma_{w,i})$ equal to 100 umol m⁻³.

In Fig. 7, we compare the modelled and observed evaporation rates of fresh and aged α -pinene 5 SOA particles for simulations with k_f and k_d values of 1×10^{-23} , 1×10^{-22} molecules⁻¹ cm³ s⁻¹ and 12, 6 $1/30 \text{ h}^{-1}$ for the short- and long-lived dimers, respectively. $k_{g,w}^*$ was set to $1/1000 \text{ s}^{-1}$. During the 7 first ~20 minutes of evaporation, before the surface-bulk layer has been entirely filled with a 8 9 mixture of short- and long-lived dimers, the modelled evaporation rates are size dependent. However, once the surface-bulk layer has been filled with dimers the evaporation is controlled by 10 the dimer degradation, and becomes nearly size independent. After ~2 hours of evaporation 11 almost all short-lived dimers in the surface-bulk layer are lost and replaced by the long-lived 12 13 dimer (see Fig. S9). This is when the second slow evaporation stage starts. If the long-lived dimer mass fraction would have been size independent, a substantially larger mass fraction of the small 14 particles would have needed to evaporate before they reach this stage. Thus, in-order for the 15 model to capture the observed nearly size independent evaporation, the long-lived dimers need to 16 be formed from the least volatile monomers, or formed in the gas-phase (e.g. by peroxy radical 17 termination reactions or hydrogen abstraction, see Sect. 2.3.4). 18

For the aged particles the model substantially underestimates the early stage evaporation losses. 19 This is because of the modelled gas-wall losses in the smog chamber. In Fig. S10 we compare the 20 modelled evaporation losses with or without chamber wall losses ($k_{e,w}^* = 0$ or 1/1000 s⁻¹) and with 21 or without ageing. Without chamber wall losses the effect of ageing in the smog chamber 22 becomes negligible, but at the same time model substantially overestimates the mass fraction loss 23 during the first evaporation stage. This is because (for this set-up) the bulk mass fraction of long-24 lived dimers is too small and ~65 mass % instead of the desired ~50 mass % need to evaporate 25 before the long-lived dimer has formed a monolayer thick surface-bulk layer coverage. With a 26 doubling of the long-lived dimer content this model and measurement discrepancy disappears. 27
From these simulations we can conclude that ADCHAM is able to reproduce the main features of
 the measured nearly size independent evaporation losses of SOA particles from Vaden et al.
 (2011) if:

- If relatively short-lived dimers are present in and near the particle surface-bulk layer,
 before the particles are introduced into the evaporation chamber.
- 6 7

2) A relatively small mass fraction of long-lived dimers, accumulate in the particle surface-bulk layer upon evaporation.

- 3) The long-lived dimer mass fraction is higher in the small particles compared to the
 large ones. As illustrated by the model simulations, this is possible (because of the
 Kelvin effect) if the dimer preferentially is formed from the least-volatile monomer
 compounds. But it could also be explained by ELVOC (e.g. dimers) formed in the gas
 phase.
- 13

14

4) The reversible gas-wall losses to the smog chamber Teflon walls have only small influences on the particle composition.

15 **3.3** Modelling of organic salt formation between carboxylic acids and ammonia

16 Here we model the SOA formation in the α -pinene – NH₃ – O₃ experiments by Na et al. (2007), in a dark indoor 18 m³ Teflon chamber. In the experiments CO (~200 ppm) was used as OH-17 18 scavenger. The chamber was operated at a temperature of 21±1 °C, and dry conditions. For the simulations we use a RH of 5 % and a temperature of 21 °C. Once the α-pinene and NH₃ initial 19 target concentrations were reached, the experiments started by injecting O₃ for approximately 20 20 21 minutes, to produce an O_3 concentration of 200 ± 5 ppb. In the model, emissions corresponding to 250 ppb unreacted O_3 were added during the first 20 minutes, in order to simulate the 22 experimental target O₃ concentrations. 23

In the experiments Na et al. (2007) observed a substantially higher SOA formation when $NH_3(g)$ was present. The authors also performed experiments on *cis*-pinonic acid (a common α -pinene oxidation product), and found a dramatic increase in particle number and volume concentration when NH_3 was added to the system. From these experiments they concluded that most of the observed SOA mass enhancement in the presence of NH_3 could be explained by acid-base reactions which drive the carboxylic acids into the particle phase. Similar organic salt formation in the presence of NH_3 was observed both at dry and humid conditions (RH=50 %). Several experiments were performed at initial $NH_3(g)$ concentration between 0-400 ppb and an α pinene concentration of ~220 ppb (see Table 1 in Na et al., 2007). The formed aerosol particle mass increases when more NH_3 is added. However, when the ammonia concentration exceeded 200 ppb no substantial additional mass formation was observed. The reason for this could be that in principle all gas phase carboxylic acids already have formed particle mass at 200 ppb NH_3 (Na et al., 2007).

Recently, Kuwata and Martin (2012) conducted experiments with an Aerosol Mass Spectrometry 7 8 (AMS) on SOA formed from ozonolysis of α -pinene at low and high relative humidity (RH<5% and RH>94%). In these experiments, the α -pinene SOA particles were formed at dry conditions 9 before they were exposed to varying degree of humidification and ammonia (see Fig. 1 in 10 Kuwata and Martin, 2012). An ~10 times greater uptake of ammonia was observed at high RH 11 12 compared to low RH, which was attributed to a more rapid diffusion uptake of ammonia in the less viscous humidified aerosol particles. Because the gas phase was not removed from the 13 aerosol between the generation and the exposure to ammonia, part of the ammonia uptake could 14 15 be attributed to reactive uptake of NH₃ and organic acids from the gas phase (Kuwata and Martin, 2012). 16

17 In this work, we model the organic salt formation between ammonium and carboxylic acids as a process occurring in the particle surface-bulk layer and particle bulk, analogous to inorganic salt 18 formation (e.g. NH₄NO₃). The partitioning of carboxylic acids and ammonia between the gas 19 phase and particle surface-bulk layer are modelled as separate pH dependent dissolution 20 processes using the condensation/evaporation module (Sect. 2.2.1). The amount of organic acids, 21 22 ammonia/ammonium and organic salts which exists in the particles depend on the pure-liquid saturation vapour pressures or Henry's law constant (K_H) , acid dissociation constants (K_a) , 23 24 activity coefficients, surface tension (Barsanti et al., 2009) and the solubility product of the formed salts (K_s) (R5-R9). The aerosol particle formation will be favoured by low pure-liquid 25 26 saturation vapour pressures of the carboxylic acids, large solubility (Henry's law coefficient) of NH₃, large difference between the carboxylic acids and NH_4^+ K_a values (Greaves and 27 Drummond, 2008) and low solubility of the formed salts (K_s). 28

29 $RCOOH(g) \leftrightarrow RCOOH(1)$

38

(R5)

1
$$\operatorname{RCOOH}(l) \leftrightarrow \operatorname{RCOO^{-}} + H^{+}, \quad K_{a} = \frac{\left[\operatorname{RCOO^{-}}\right]\left[H^{+}\right]\gamma_{H^{+}}\gamma_{\operatorname{RCOO^{-}}}}{\left[\operatorname{RCOOH}\right]\gamma_{\operatorname{RCOOH}}}$$
 (R6)

2
$$\operatorname{NH}_{3}(g) \leftrightarrow \operatorname{NH}_{3}(1), \quad K_{H} = \frac{\left[\operatorname{NH}_{3}(1)\right]\gamma_{\operatorname{NH}_{3}}}{P_{\operatorname{NH}_{3}}}$$
 (R7)

3
$$\operatorname{NH}_{3}(1) + \operatorname{H}^{+} \leftrightarrow \operatorname{NH}_{4}^{+}, \quad \frac{1}{K_{a,NH_{4}^{+}}} = K_{NH_{3}} = \frac{\left[\operatorname{NH}_{4}^{+}\right]\gamma_{NH_{4}^{+}}}{\left[\operatorname{NH}_{3}(1)\right]\left[\operatorname{H}^{+}\right]\gamma_{NH_{3}}\gamma_{H^{+}}}$$
(R8)

4
$$\operatorname{NH}_{4}^{+} + \operatorname{RCOO}^{-} \leftrightarrow \operatorname{NH}_{4}\operatorname{RCOO}(s), \quad K_{s} = \left[\operatorname{NH}_{4}^{+}\right]\left[\operatorname{RCOO}^{-}\right]\gamma_{\operatorname{NH}_{4}^{+}}\gamma_{\operatorname{RCOO}^{-}}$$
 (R9)

5 Table 3 lists different model parameter values used for the base case simulations in this section. The K_a values are unknown for most carboxylic acids, even in aqueous solutions. However, for 6 two major ozonolysis products (cis-pinic acid) and (cis-pinonic acid) (Hallquist et al., 2009), 7 aqueous pK_a values were found in the literature (see e.g. Hyder et al., 2012 and Barsanti et al., 8 2009). These acids have nearly the same pK_a values (~4.6). Hence, in this work we assume that 9 all carboxylic acids from α-pinene ozonolysis which partition into the particle organic rich phase 10 11 have a pK_a values equal to 4.6. The carboxylic acid and ammonia dissociation rate coefficients and the Henry's law coefficient of NH₃ are valid for dilute water solutions (see Sect. 2.3.2). Thus, 12 as a sensitivity test we also tested to model the NH₃ uptake with a 10 times lower Henry's law 13 coefficient, which may be more appropriate for organic solvents. 14

Unfortunately we could not find any values of solubility products between carboxylic acids and ammonium in the literature. Hence, we decided to define an effective solubility product (K_s^*) as the product between the ammonium concentration and the total deprotonated carboxylic acid concentration ([RCOO⁻]_{tot}) (Eq. 29). K_s^* was the only parameters which we systematically varied in order to find the best possible agreement between the model and measurements.

20
$$K_s^* = \left[\mathrm{NH}_4^+ \right] \left[\mathrm{RCOO}^- \right]_{\mathrm{tot}}$$
 (29)

If not otherwise specified, the pure-liquid saturation vapour pressures of the organic compounds were estimated with the SIMPOL method, K_s^* was set to 0.1 mol² m⁻⁶, and the NH₄RCOO salts were mixed with the other organic compounds (no separate phase). Because the interactions

between the NH₄RCOO and other organic compounds and inorganic ions are unknown (see Sect. 1 2 2.3.1), NH₄RCOO was not considered to influence the activity coefficients of the other compounds. However, as a second extreme condition we performed simulations where we treated 3 NH₄RCOO and the other organic compounds + inorganics as two completely separate phases 4 (liquid-liquid phase separated or NH₄RCOO as crystalline salts (see Sect. 1)). The diffusion 5 coefficients for monomer SOA and ammonia/ammonium were estimated with the Stokes-6 Einstein relationship using a viscosity of $\sim 10^8$ Pa s (Abramson et al., 2013). Because the viscosity 7 of the SOA is uncertain and depend on the experimental conditions and time of aging, we also 8 performed simulations with less viscous particles ($D_{0,monomer,SOA}=10^{-15}$ cm² s⁻¹, $D_{0,ammonium}=10^{-13}$ 9 $cm^2 s^{-1}$). 10

In Table 4 we have listed the measured and model initial concentrations, concentration change of 11 ozone $(\Delta[O_3] = [O_3]_{max} - [O_3]_{t=6h})$ and α -pinene $\Delta[\alpha$ -pin.], and SOA yields. Figure 8 shows the 12 modelled temporal evolution of the α -pinene, O₃, NH₃ and OH concentrations in the gas phase. 13 The O₃ concentration rises during the first 20 minutes while O₃ is continuously applied to the 14 chamber. The OH concentration reaches a maximum of $\sim 10^6$ molecules cm⁻³ at the same time as 15 the maximum O₃ concentration. Hence, according to the model the experiments with CO as OH 16 scavenger are not pure O_3 oxidation experiments, but a fraction of the α -pinene and the oxidation 17 products are also oxidized with OH. Figure S11 in the supplementary material shows the 18 cumulative fraction of reacted α -pinene which was oxidized by O₃ during the evolution of the 19 20 experiment. In the beginning of the experiment only 86 % of the consumed a-pinene was oxidized by O₃, while at the end of the experiment 92 % of the consumed α-pinene was oxidized 21 by O₃. 22

In Fig. 9 we compare the modelled and measured SOA yields from experiments conducted with 23 24 approximately 220 ppb α -pinene, 200 ppb O₃ and varying initial NH₃ concentrations. The model results in Fig. 9a are from the base case simulation set-up (Table 3). Figure 9b shows model 25 26 results from simulations performed with pure-liquid saturation vapour pressures from Nannoolal et al. (2008). The results in Fig. 9c are from model runs with unity activity coefficients (Raoult's 27 law for ideal solution) and Fig. 9d shows results from simulations with less viscous particles 28 $(D_{0,monomer,SOA}=10^{-15} \text{ cm}^2 \text{ s}^{-1}, D_{0,ammonium}=10^{-13} \text{ cm}^2 \text{ s}^{-1} \text{ and } D_{0,NH4RCOO}=0 \text{ cm}^2 \text{ s}^{-1})$. For a particle 29 with a diameter of 250 nm these values of the diffusion coefficients gives an expected e-folding 30

time of equilibration of 2.6 minutes for ammonium and 4.4 hours for SOA monomers (Seinfeld
and Pandis, 2006). However, since a substantial fraction of the ammonium can be bound into
NH₄RCOO, the actual e-folding time can be longer.

4 For the base case simulations the agreement between the modelled and measured SOA mass and SOA yields are surprisingly good, both with and without addition of NH₃. One reason for this is 5 that the organic salt effective solubility product (Eq. 29) was used as a model fitting parameter. 6 However, in order for the model to agree with the measurements the amount of semi-volatile 7 8 carboxylic acids formed from the α -pinene oxidation still needs to be reasonably well predicted, 9 which seems to be the case. It is also important to mention that for these model simulations we 10 did not consider any chamber wall losses. Figure S12 in the supplementary material shows the 11 modelled temporal evolution of the total carboxylic acid concentration (gas + particle phase).

We find the largest difference between the model runs, and between the model and measurements, when we use the pure-liquid saturation vapour pressure method from Nannoolal et al. (2008) instead of SIMPOL (Pankow and Asher, 2008) (Fig. 9b). The model then underestimates the SOA mass with ~200 μ g m⁻³ (~30 %), irrespectively of the amount of NH₃ added.

Figure S4 in the supplementary material shows a comparison of the volatility basis set (VBS) 17 parameterization from Pathak et al. (2007) and VBS parameterizations which we have derived 18 19 from the MCMv3.2 condensable α-pinene oxidation products using either the method from Nannoolal et al. (2008) or SIMPOL. The MCMv3.2 a-pinene oxidation product VBS 20 21 parameterizations are given both for CO and cyclohexane as OH-scavenger. The VBS parameterizations show large differences both between the vapour pressure methods and the type 22 of OH scavenger used. By comparing the VBS parameterizations we can conclude that SIMPOL 23 gives the largest SOA mass at high α -pinene concentrations (this work). However, at low 24 25 (atmospherically more realistic) α-pinene concentrations the Nannoolal method will give the least 26 volatile SOA and highest SOA mass.

Barley and McFiggans, 2010 have shown that the uncertainties of the calculated pure-liquid
saturation vapour pressures are large, especially for low-volatility compounds with several
functional groups. However, because of other large uncertainties, e.g. oligomerization processes

and gas phase chemistry mechanisms (see Sect. 1), we cannot predict which of the two liquid saturation vapour pressure methods that give the most realistic vapour pressures. In Sect. 3.3 we illustrate how the estimated volatility of the α -pinene gas phase oxidation products can have substantial effects on the particle evaporation loss rates.

- In contrast to the vapour pressures, the modelled activity coefficients have only small influence 5 on the simulated SOA mass formation (compare Fig. 9a and 9c). This is consistent with the 6 conclusions from McFiggans et al. (2010), and Zuend and Seinfeld (2012) for conditions without 7 8 dissolved inorganic ions and low relative humidity. The mass difference between the model runs ([OA_{ideal}]-[OA_{activity}]) is small without added NH₃, but increases when the free particle 9 ammonium concentration increases. The reason for this is that the dissolved ammonium ions 10 generally increase the organic molecule activity coefficients (salting-out effect). At atmospheric 11 more realistic relative humidities (>30 %), salt effects which either cause liquid-liquid phase 12 separation or drive the organic compounds out from the particles, may have large effects on the 13 SOA formation (see e.g. Zuend and Seinfeld, 2012). 14
- 15 If we assume that the SOA is less viscous (Fig. 9d), the mass yields are slightly larger (60.7 % 16 compared to 57.5% without NH_3 addition, and 69.1 % compared to 67.0 % when 200 ppb NH_3 is 17 added at the start of the experiments).
- Figure S13 in the supplementary material shows the total SOA mass and NH₄RCOO mass for varying initial NH₃ concentration, $K_s^* = 0.01$ or $0.1 \text{ mol}^2 \text{ m}^{-6}$ and semi-solid SOA particles. As expected the NH₄RCOO mass concentration, and the total particle mass increases when K_s^* is lowered. However, for 200 ppb NH₃ the difference becomes negligible since almost all carboxylic acids are anyhow found in the particle phase. The results also reveal a moderate salting-out effect of the ammonium on the SOA (see the decrease in the total particle mass with increasing NH₃ when $K_s^*=0.01 \text{ mol}^2 \text{ m}^{-6}$).

We also performed simulations with 10 times lower Henry's law coefficients and $K_s^* = 0.01$ or 0.1 mol² m⁻⁶ (Fig. S14 and S15). With $K_s^* = 0.1 \text{ mol}^2 \text{ m}^{-6}$ no NH₄RCOO is formed even if 200 ppb NH₃ is added. However, the added NH₃ still contributes to an enhanced dissociation and uptake of the carboxylic acids. When 200 ppb NH₃ is added and K_s^* is 0.1 mol² m⁻⁶ the model gives a SOA mass increase of 11 % and the measurements an increase of 22 %. If we decrease 1 K_s^* to 0.01 mol² m⁻⁶, substantial amount of NH₄RCOO is formed and the model are nearly able to 2 captures the observed SOA mass increase with increasing NH₃ concentrations (a 17 % increase in 3 mass when 200 ppb NH₃ is added) (Fig. S15).

In order to test which processes that are responsible for the observed NH₃ uptake in α -pinene SOA particles (Kuwata and Martin, 2012 and Na et al., 2007), we also performed simulations where the SOA particles were allowed to age for 6 hours before they were exposed to 200 ppb NH₃(g). To test the effect of mass transfer limited uptake of NH₃, the particles were either treated as glassy solids (no mixing) or semi-solid less viscous ($D_{0,monomer,SOA}=10^{-15}$ cm² s⁻¹, $D_{0,mmonium}=10^{-13}$ cm² s⁻¹).

In Fig. 10a the temporal evolution of the modelled SOA mass from these simulations is shown. 10 As a comparison, the results from simulations with 200 ppb NH₃(g) added at the start of the 11 12 experiments are also plotted. After the addition of NH₃, the SOA mass increases rapidly both with and without mass transfer limited diffusion uptake in the particles (semi-sold or solid 13 particles). This indicates that the rapid uptake of $NH_3(g)$, by the particles, mainly is caused by 14 reactive uptake of carboxylic acids(g) and $NH_3(g)$ and not by the diffusion of NH_3/NH_4^+ into the 15 particle bulk. However, the temporal evolution of the formed NH₄RCOO salts and dissociated 16 17 and non-dissociated carboxylic acids (Fig. 10b), reveal that the mass of NH₄RCOO salts formed in the semi-solid particles are twice as high, and the carboxylic acid mass concentration is 18 substantially lower, than if treating the SOA as solid. This difference is attributed to the mass 19 transfer limited uptake and reaction of NH₃/NH₄⁺ with the carboxylic acids found in the semi-20 solid particle bulk interior. 21

22 However, although the NH₄RCOO concentration becomes higher if the particles are semi-solid (less viscous), the total aerosol mass 3 hours after the addition of ammonium is lower than if the 23 particles are solid (compare simulation Nr. 4 and 5 in Fig. 10a). The reason for this is the salting-24 out effect of NH4⁺ which causes the nonpolar organic compounds to evaporate. For these 25 simulations, the salting-out effect is mainly important if both the NH₃/NH₄⁺ and the organic 26 compounds can be transported between the bulk and particle surface-bulk layer. In the laboratory 27 experiments (see Fig. 2a in Na et al., 2007) no SOA mass loss could be seen after the NH₄RCOO 28 formation. This experiment continued less than 1 hour after the addition of NH₃, but it at least 29 indicates that the mixing of organic compounds within the particle phase is mass transfer limited, 30

and/or that the NH₄RCOO salts form a separate phase, which limits the salting-out of other SOA
 compounds from the particles to the gas phase.

Figure 10a also shows the simulated SOA mass formation when we treat the NH₄RCOO salts as a 3 separate phase (e.g. crystalline salt) which other condensable organic compounds cannot dissolve 4 into. When $NH_3(g)$ is added during the start of the experiments the difference between the model 5 runs with and without a separate NH₄RCOO phase is relatively small. However, if the NH₃(g) is 6 added after the solid SOA particles have formed, only a moderate SOA mass increase is 7 8 accomplished (~9%). This is in sharp contrast to the results from the simulations with solid particles and only one organic phase (mass increase of ~39 %). The reason for this is that the 9 ammonium salts are enriched in the particle surface-bulk layer, and if no other compounds can 10 dissolve into this phase their uptake is limited. On the other hand if NH₄RCOO is part of a single 11 amorphous organic phase, it will lower the mole fractions of the other compounds and hence 12 increase (at least for ideal conditions) the uptake of them (see Eq. 2). This is the reason why the 13 total SOA mass increase is larger (~270 μ g m⁻³, ~39 %), than the increase explained purely by the 14 carboxylic acids and NH₄RCOO (46+84=130 µg m⁻³, ~19 %) (see simulation Nr. 4 in Fig. 10a 15 and Fig. 10b). Na et al. (2007) observed a mass increase of 15 % when 1000 ppb NH₃ was added 16 after the α -pinene SOA particle mass formation had ceased. This increase is larger than the 17 modelled increase when considering complete phase separation between NH₄RCOO and the 18 19 other condensable organic compounds, but substantially smaller than for the simulations with 20 only one organic phase. This may indicates that in reality, there will neither be perfect (ideal) mixing between NH₄RCOO and the other condensable organic compounds, nor a complete phase 21 separation. 22

Figure 11 shows a) the modelled pH, b) the total ammonium mass fraction (free and bonded in 23 ammonium salts), c) the NH₄RCOO mass fraction and d) the carboxylic acid mass fraction 24 ([RCOO⁻]+[RCOOH]) for a semi-solid SOA particle, at different distances from the particle core. 25 26 The figure includes results from simulations with initial NH₃(g) concentrations of 50, 100 and 200 ppb, respectively, and at 1 or 6 hours of aging. A large fraction of the SOA formed early 27 during the experiments are due to condensation of carboxylic acids. This explains the large mass 28 fractions of carboxylic acids and the lower pH in the particle cores (Fig. 11a-b). For the 29 30 simulations with 200 ppb NH₃, a large fraction of the carboxylic acids and ammonium form salts

1 (Fig. 11c), while when only 50 ppb NH₃ is added, ammonium salts are only present during the 2 early stage of particle formation, when the carboxylic acid mass fraction is large. Because of the 3 assumed relatively rapid mixing of ammonium (e-folding time of a few minutes), the free 4 ammonium concentration (not bound in organic salts) are almost constant in all particle layers. 5 Hence, the differences in the NH₄RCOO concentrations between different layers are largely 6 caused by differences in the carboxylic acid concentrations, which even after 6 hours of aging are 7 not uniformly mixed.

8 It has been suggested that organic salt formation between carboxylic acids and NH₃ or ammines could possibly be responsible for the early growth of nanometre sized particles in the atmosphere 9 (e.g. Smith et al. 2008, Barsanti et al, 2009 and Smith et al., 2010, Yli-Juuti et al., 2013). In order 10 to be able to draw any conclusions from our simulations concerning the potential effect of 11 12 NH₄RCOO formation in the atmosphere, we performed simulations where we decreased the α pinene concentration to 50 ppb and varied the NH₃ concentration in the range 0-2 ppb. We used 13 50 ppb α -pinene, because in the model ~30 ppb α -pinene needs to react before the particles with 14 an initial diameter of 5 nm start to grow. Furthermore, the model simulations do not consider 15 inorganic salt formation between NH₃ and the strong acids H₂SO₄ or HNO₃. Yli-Juuti et al. 16 (2013) have shown that for typical conditions over boreal forest, NH₃ will preferentially form 17 inorganic salts with H₂SO₄ and not with carboxylic acids. In our simulations, very little 18 NH₄RCOO is formed even if the NH₃ concentration is 2 ppb and the average growth rate between 19 20 5 and 20 nm is only amplified with ~7 % (see Fig. S16).

From the simulations in this section we can conclude that ADCHAM (with the pure-liquid saturation vapour pressures from SIMPOL and activity coefficients from AIOMFAC), are able to reproduce the observed SOA formation at different concentrations of $NH_3(g)$. With NH_3 present during the formation, reactive uptake of carboxylic acids contributes substantially to the modelled early growth of the particles formed by homogeneous nucleation. However, this is probably not the case for atmospheric more relevant NH_3 and α -pinene concentrations.

27 **3.4 SOA formation from oxidation of** *m***-xylene**

Here we model the SOA formation from an *m*-xylene oxidation experiment (Exp. P2 in Nordin et al. 2013). The experiment was conducted in a 6 m^3 Teflon chamber in the Aerosol Laboratory at Lund University. The experiment started with dark conditions by adding (NH₄)₂SO₄ seed aerosol
 into the chamber (~20 μg m⁻³), followed by ~40 ppb NO and ~240 ppb *m*-xylene. Approximately
 30 minutes before the UV-lights were turned on (~90 minutes after the start of the experiment),
 (NH₄)₂SO₄ particles were added a second time in order to achieve the target (NH₄)₂SO₄ mass of
 ~20 μg m⁻³.

The seed aerosol was formed by nebulizing an $(NH_4)_2SO_4$ – water solution and then drying the 6 7 droplets. Before the dry (NH₄)₂SO₄ particles were introduced into the chamber they were passed 8 through a bi-polar charger in order to achieve a well-defined nearly Boltzmann distributed charge distribution (Wiedensohler et al., 2012). The experiment was performed at a temperature of 22 °C 9 \pm 2 °C, dry conditions (RH of 3-5 %) and in the presence of UV-light with an experimentally 10 derived NO₂ photolysis rate of 0.2 min⁻¹. The experimental set-up has been described in detail by 11 Nordin et al. (2013). The measured UV-light spectrum (320-380 nm) is given in the 12 supplementary material to Nordin et al. (2013). 13

In the model we used a temperature of 21 °C and a RH of 5 %. The photolysis rates were calculated with the recommended cross sections and quantum yields from MCMv3.2 and the measured 1 nm resolution UV-spectrum from Nordin et al. 2013, with a total light intensity of 23 W/m² which gives a NO₂ photolysis rate of 0.20 min⁻¹.

18 **3.4.1 Particle deposition loss rates**

To be able to quantify the effect of dry deposition on the estimated SOA formation from chamber 19 experiments, the deposition losses of particles to the chamber walls needs to be evaluated. The 20 dry deposition depends both on the friction velocity (u^*) , the particle size and charge 21 distributions, the mean electrical field strength (\overline{E}) in the chamber, and the chamber surface area 22 to volume ratio (see Sect. 2.2.3). \overline{E} and u^* are commonly not known, but can be estimated by 23 fitting the model to particle number size distribution measurements. For this purpose an 24 25 experiment with (NH₄)₂SO₄ seed particles but without condensable organic compounds was performed. 26

As the experiments in the chamber proceed, the chamber surface area to volume ratio increasedbecause of instrument sampling and leakage out from the chamber due to a small over pressure

1 inside the chamber (see Nordin et al., 2013). We estimate the chamber volume loss rates 2 $(\Delta V / \Delta t)$ during the experiments to $0.8 \pm 0.2 \text{ m}^3 \text{ h}^{-1}$.

In Fig. S17 we compare the modelled and measured temporal evolution of the particle number-3 and particle volume concentration for simulations with different values of \overline{E} and u^* . With a 4 $\Delta V / \Delta t$ of 0.8 m³ h⁻¹, coagulation and deposition, and a \overline{E} of 50 V cm⁻¹ and a u^* of 0.05 m s⁻¹, 5 ADCHAM is able to nearly reproduce the measured $(NH_4)_2SO_4$ particle number size distributions 6 (Fig. 12a), the temporal evolution of the total particle number (Fig. 12c) and volume 7 8 concentrations (Fig. 12d). The coagulation has no direct influence on the particle volume 9 concentration but is important for the particle number concentration at the end of the experiment. In the beginning of the experiment the charged smallest particles are rapidly deposited to the 10 chamber walls resulting in a high effective wall deposition loss rate $(k_w (s^{-1}))$ (Fig. 12b). But, as 11 the experiment proceeds the fraction of charged particles (especially the small ones) decreases in 12 the air. At the same time the surface area to volume ratio increases in the chamber, which in turn 13 increases the deposition loss rates of all particle sizes (see the gradual upward displacement of the 14 curves in Fig. 12b). Recharging of particles by collision with air ions was not considered in the 15 16 model.

After the tuning of the dry deposition loss rates on the pure seed aerosol experiments we used ADCHAM to simulate the SOA formation experiment with *m*-xylene as precursor (Nordin et al. 2013). However, if we use the same $(\Delta V / \Delta t)$, \overline{E} and u^* as in the pure seed particle deposition experiment, the model underestimates the seed aerosol mass loss (especially during the first 2 hours after the UV-lights are turned on), but substantially overestimates the particle number concentration losses before the UV-lights are turned on.

The heating of the air by the UV-lights and the air condition units which blow on the outer chamber walls can produce an increased mixing within the chamber. Therefore, before the UVlights are turned on u^* may be smaller. By decreasing u^* to 0.01 m s⁻¹ before the UV-lights are turned on the model better captures the measured initial particle number concentration losses.

27 Another important difference between the pure seed particle experiment and the *m*-xylene 28 precursor experiment is that the latter experiment was performed during almost twice as long 29 time (\sim 6 h). Hence, the effect of particle recharging when colliding with air ions may be more 1 important to consider. Furthermore, the chamber volume during the end of the *m*-xylene 2 experiment was substantially smaller (1.5-2 m³). This might have increased the effective mean 3 electrical field strength within the chamber (see Sect. 2.2.3). In the model we try to account for 4 this by calculating $\overline{E_t}$ (at time t) as the quotient between the initial mean electrical field strength 5 $\overline{E_0}$ (50 V cm⁻¹) and the relative change of the approximate distance between the roof and ceiling 6 (*h*) of the chamber (h_t/h_0) (which is approximately equal to the relative chamber volume change 7 (V_t/V_0)) (Eq. 30).

8
$$\overline{E_t} = \frac{\overline{E_0}}{h_t / h_0} \approx \frac{\overline{E_0}}{V_t / V_0}$$
 (30)

9 Figure S18 in the supplementary material compares the modelled and measured (with AMS and scanning mobility particle sizer (SMPS)) temporal evolution of the sulphate seed particle mass 10 11 concentration, particle number concentration and the particle number size distribution, and the modelled initial and final effective dry deposition loss rates. The model results are from 12 simulations with, $\overline{E_t} = 50 \text{ V cm}^{-1}$ and $u^* = 0.05 \text{ m s}^{-1}$ or $\overline{E_t}$ calculated with Eq. (30) and with $u^* =$ 13 0.01 m s⁻¹ before the UV-lights are turned on. With the latter values the model shows 14 15 substantially better agreement with the measured temporal evolutions of the sulphate seed aerosol mass concentration. However, the model still overestimates the particle number concentration 16 loss rates (especially after the UV-light are turned on). For the model simulations presented 17 below we will use Eq. (30) to estimate $\overline{E_t}$, and $u^* = 0.01 \text{ m s}^{-1}$ before the UV-lights are turned on 18 and $u^* = 0.05 \text{ m s}^{-1}$ after the UV-lights are turned on. 19

20 3.4.2 Gas - particle partitioning and heterogeneous reactions

Since the *m*-xylene experiment was performed at dry conditions the $(NH_4)_2SO_4$ seed particles will initially be in a solid crystalline phase. Therefore, we assume that no material is mixed between the crystalline solid salt cores and the SOA coating (see e.g. Fig. 1a in Bertram et al., 2011). Hence, in the model there will be no salting-out effect (increase of the nonpolar organic compound activity coefficients caused by NH_4^+ , SO_4^{-2} and HSO_4^- from the seed aerosol particles) (see discussion in Sect. 3.2 on possible salting-out effects of NH_4^+).

In total we considered 112 potentially condensable ($p_0 < 1$ Pa) non-radical organic MCMv3.2 1 compounds. The pure-liquid saturation vapour pressures were calculated with either the SIMPOL 2 (Pankow and Asher, 2008) or the method from Nannoolal et al. (2008). We also used a third 3 (semi-empirical) method to model the SOA formation. This method considers in total three 4 oxidation products with vapour pressures and molar based stoichiometric yields (α_i) derived from 5 the parameterizations for low and high NO conditions from Ng et al. (2007). For this we assume 6 that the condensable organic compounds have a molar mass of 200 g mol⁻¹. The two most volatile 7 compounds ($p_{0,1} = 6.4 \times 10^{-6}$ Pa, $\alpha_1 = 0.021$ and $p_{0,2} = 1.7 \times 10^{-4}$ Pa, $\alpha_2 = 0.061$) represent the volatility 8 distribution of the condensable oxidation products formed through the $RO_2 + NO$ pathway. The 9 10 third non-volatile product ($p_{0,3} = 0$ Pa, $\alpha_3 = 0.245$) represent the generally less volatile organic compounds formed through the $RO_2 + HO_2$ pathway. The gas phase was still modelled with the 11 MCMv3.2. The fraction of condensable organic compounds which was formed through the RO₂ 12 + HO₂ pathway (product 3) was derived with the ratio $(k_{RO2+HO2}[HO_2]/(k_{RO2+NO}[NO]+$ 13 14 $k_{RO2+HO2}[HO_2]$) as proposed by Ng et al. (2007).

The partitioning of the condensable organic compounds to the wall deposited particles and the 15 Teflon walls were modelled according to the procedure described in Sect. 2.2.3. The uptake onto 16 the Teflon film and the particles deposited on the chamber walls depends on the laminar layer 17 width adjacent to the chamber walls (Δx). The uptake (adsorption) on the Teflon film also 18 depends on the first order loss rate from the near wall gas phase to the walls $(k_{g,w})$ and the 19 20 desorption rate from the Teflon surfaces out to the thin layer next to the chamber walls $(k_{w,g,i})$ (Eq. 4). In Sect. 3.4.4 we test different values of $\Delta x_i k_{g,w}$ and $k_{w,g,i}$ in order to find the best possible 21 agreement between the modelled and the measured SOA formation. 22

Because coagulation has a considerable influence on the modelled particle number size 23 distribution (see Fig. 12) we will consider this process as well. However, with the current version 24 of ADCHAM coagulation cannot be combined with the complete kinetic multi-layer model (see 25 Sect. 2.2.2). Hence, for the simulations presented in this section the particles were only divided 26 into a solid seed particle core, and a second (well-mixed) bulk layer and a surface monolayer 27 which are composed of the condensable organic compounds. Additionally (if specified) we also 28 consider the adsorption and desorption of O₃ and NO₂, the mass transfer limited diffusion of O₃ 29 30 and NO₂ from the sorption layer into the particle bulk, and the particle phase reactions between

O₃ and unsaturated organic compounds (see Sect. 2.4.2) or between NO₂ and oxidized aromatic
compounds (see Sect. 3.4.3).

Table 1 in Sect. 2.4.2 gives the model parameter values used for O₃ uptake. For the simulations 3 presented here the diffusion coefficient of ozone (D_{0,O_3}) was set to values between 10^{-7} and 10^{-8} 4 cm² s⁻¹ (semi-solid SOA (see e.g. Table 1 in Shiraiwa et al., 2011)), and the reaction rate 5 constants between ozone and the unsaturated (non-aromatic carbon-carbon double bond) organic 6 compounds (k_{0_2}) were varied between 10^{-16} and 10^{-17} molecules⁻¹ cm³ s⁻¹. This can be compared 7 with the measured k_{0_3} of 10⁻¹⁶ molecules⁻¹ cm³ s⁻¹ for the heterogeneous ozonolysis of oleic and 8 palmitoleic acid (Huff Hartz et al., 2007). Berkemeier et al. (2013) used the KM-SUB model to 9 constrain the kinetic parameter values which limits the ozonolysis of oleic acid. According to 10 their model simulations k_{0_3} should be somewhere in the range of 1.7×10^{-15} - 1.7×10^{-17} molecules⁻¹ 11 cm³ s⁻¹. The formed particle phase oxidation products were assumed to be non-volatile, which 12 likely is an acceptable assumption if the oxidation products rapidly react and form dimer SOA 13 (see e.g. Maksymiuk et al., 2009). Apart from increasing the SOA mass formation and changing 14 the chemical composition of the SOA these heterogeneous reactions may also serve as an 15 additional ozone sink (which is not accounted for by the MCMv3.2 gas phase chemistry 16 mechanism). 17

Additionally, we will also test peroxyhemiacetal and hemiacetal dimer formation in the particlephase.

3.4.3 Gas phase chemistry and influence from chamber walls and heterogeneous reactions

Bloss et al. (2005a-b) have previously shown that the MCMv3.1 (without particle SOA formation and particle phase chemistry) generally overestimates the ozone concentration and underestimates the OH concentration during oxidation of light aromatic compounds (e.g. xylene and toluene). Hence, for these systems MCM also tends to underestimate the NO and hydrocarbon oxidation (loss) rates. In order to account for the missing OH source Bloss et al. 2005b had to include an artificial OH source of $4x10^8$ molecules cm⁻³ s⁻¹ when modelling a toluene oxidation experiment from the EUPHORE chamber. Conversion of NO₂ to HONO on the organic particle surfaces may partly explain the discrepancy between the modelled and measured particle phase chemistry (Bloss et al., 2005b). These reactions have been observed on diesel exhaust particles (Gutzwiller et al., 2002) and on organic aerosol surfaces e.g. by George et al. (2005). Metzger et al. (2008) instead proposed that the NO₂ primarily is converted to HONO on the Teflon chamber walls.

In this work we will test the heterogeneous NO_2 to HONO conversion mechanism. Bloss et al. 6 (2005b) modelled this mechanism using a constant reaction probability (γ_{HONO}) of 0.025 for the 7 NO₂ molecules which collide with a particle. In this work we model this proposed mechanism in 8 a more detailed way by considering the adsorption, diffusion and reaction of NO₂ with specific 9 organic compounds in the particle phase. NO₂ has approximately the same Henry's law 10 coefficient for dissolution in water (Seinfeld and Pandis, 2006) and molecule size as O₃. Hence, 11 for these simulations, we will use the same parameter values for NO₂ (e.g. Henry's law 12 coefficient and diffusion coefficient) as specified for O₃ in Sect. 3.4.2 and in Table 1. 13

Gutzwiller et al. (2002) suggested that the organic compounds which react with NO₂ in the particle phase and form HONO are oxygenated aromatics (e.g. 2-methoxyphenol). Hence, we assume that it is only the compounds that contain an aromatic ring which will be oxidized by NO₂ and form HONO. The organic oxidation products formed from these heterogeneous reactions were assumed to be non-volatile.

The NO₂ to HONO conversion mechanism was considered both on the particles deposited on the chamber walls and in the air. We will also test an additional photo-enhanced background reactivity caused by HONO release from the chamber walls (Rohrer et al., 2005). The strength of the HONO emissions (from the walls to the near surface gas phase) in the Lund Teflon chamber (Nordin et al, 2013) was estimated to be 4.6×10^8 molecules cm⁻² s⁻¹. This value is based on the estimated HONO wall production rate of 9.1×10^6 molecules cm⁻³ s⁻¹ in Metzger et al. (2008) and their chamber volume to surface area characteristics (Paulsen et al., 2005).

Analogous to the MCM light aromatic model simulations by Bloss et al. (2005a-b) we underestimate the OH and overestimate the maximum O_3 concentration, without tuning the MCM gas phase chemistry (Fig. 13). MCMv3.2 also underestimate the initial O_3 formation rate, the amount of reacted *m*-xylene and the rapid NO to NO₂ conversion which starts approximately 20 minutes after the UV-lights were turned on. Therefore, analogous to Bloss et al. (2005b) we decided to include an artificial OH source, in our case with a rate of 10^8 cm⁻³ s⁻¹ from 20 minutes after the UV-light were turned on until the end of the experiment. This substantially improves the agreement between the modelled and measured NO, NO₂, O₃ and *m*-xylene concentrations. However, the model still substantially overestimates the maximum O₃ concentration.

6 The poor agreement between the modelled and measured NO_2 in the latter half of the experiment 7 (Fig. 13b) is because of the interference from peroxy acyl nitrates (PAN), HNO₃, HONO, N₂O₅ 8 and other nitrate containing compounds in the chemiluminescence instrument used (see Nordin et 9 al., 2013 and references there in).

Figure 13 also shows the results from a simulation where we additionally include HONO 10 emissions from the chamber walls. Because the surface area to volume ratio increases during the 11 12 experiments (~5 times) these emissions have an increasing influence on the modelled gas phase chemistry. With HONO emissions and the OH source, the OH concentration at the end of the 13 model run is 1.5×10^6 cm⁻³, while without these emissions but with the OH source the 14 concentration is $7x10^5$ cm⁻³. Hence, with HONO wall emissions more *m*-xylene reacts in the 15 simulation than what is indicated by the gas chromatography – mass spectrometry (GC-MS) 16 17 measurements. Additionally, the model O₃ concentration becomes even higher.

In order to be able to compare the modelled and measured SOA formation during the experiment 18 it is crucial that we are able to accurately simulate both the amount of m-xylene which is 19 consumed and the fraction of RO₂ which reacts with HO₂ and NO, respectively (see e.g. Ng et al., 20 2007 and Kroll and Seinfeld, 2008). Hence, if not otherwise specified we included the artificial 21 22 OH source but not any HONO emissions from the chamber walls. With this model set-up, the cumulative fraction of the *m*-xylene first generation RO₂ oxidation products which have reacted 23 with HO_2 at the end of the experiment is about ~65 %. When we also include HONO wall 24 emissions this value is ~50 % and with the non-tuned MCMv3.2 chemistry we get a value of ~35 25 % (see Fig. S19 in the supplementary material). 26

In order to test whether HONO formation from heterogeneous reactions between NO_2 and oxidized aromatic compounds can improve the agreement between the modelled and measured O_3 concentration, we performed a simulation with what we believe are upper estimates of the 1 reaction rates between NO₂ and the oxidized aromatic compounds and the NO₂ diffusion 2 coefficient ($k_{NO_2} = 10^{-15}$ molecules⁻¹ cm³ s⁻¹ and $D_{0,NO_2} = 10^{-7}$ cm² s⁻¹). With these values ~60 % of 3 the aromatic SOA was oxidized by NO₂. The formed (in the model non-volatile) oxidation 4 products comprise 20 % of the total SOA mass in the end of the simulation (Fig. S20a in the 5 supplementary material). Still, this has only a moderate influence on the HONO concentration 6 (Fig. S20b) and the NO₂ and O₃ decrease is equal or less than ~1 % (Fig. S20c-d).

Figure S20d also shows the modelled $O_3(g)$ concentration when including heterogeneous 7 reactions between O₃ and the unsaturated organic compounds ($k_{O_3} = 10^{-16}$ molecules⁻¹ cm³ s⁻¹ and 8 $D_{0.0} = 10^{-7}$ cm² s⁻¹). For this simulation ~98 % of the unsaturated organic compounds in the 9 particle phase were oxidized by O3 and the formed non-volatile SOA products comprise 37 % of 10 the total SOA mass. However, comparable to the heterogeneous NO₂ to HONO conversion this 11 12 has a very small influence on the modelled $O_3(g)$ concentration (~1% decrease). Hence, we can conclude that it seems unlikely that heterogeneous reactions between NO2 and oxidized aromatic 13 compounds and/or between O₃ and the unsaturated organic compounds can explain why 14 measurements generally gives much lower O₃(g) concentrations than MCM model. However, as 15 will be shown in Sect. 3.4.4 these heterogeneous reactions can still be important for the amount 16 17 and type of SOA which is formed.

3.4.4 SOA formation, properties and the potential influence from chamber wall effects and heterogeneous reactions

In Fig. 14 we compare the modelled and measured particle volume concentrations during the m-20 xylene experiment. The model results are from simulations with the SIMPOL vapour pressure 21 method. The desorption of condensable organic compounds from the chamber walls was 22 modelled with $C_w / (M_w \gamma_{w,i})$ in Eq. (4) equal to 100 µmol m⁻³. This value is between those 23 measured by Matsunaga and Ziemann (2010) for 2-alcoholes and 2-ketones (see Sect. 2.2.3). For 24 the model results in Fig. 14a we used a Δx of 0.1 cm and $k_{g,w}$ was set to 1/20 s⁻¹ while for the 25 results in Fig. 14b we used a Δx of 1.0 cm and $k_{g,w}$ was set to 1/6 s⁻¹. Hence, the model simulation 26 in Fig. 14a represent conditions with only relatively small mass transfer limitations for the gas 27 28 exchange between the air and the chamber walls and particles on the walls, and a relatively slow uptake of organic compounds directly onto the Teflon walls. The model simulation in Fig. 15b
instead represents conditions where the mass transfer limitations between the air and the chamber
walls and particles on the walls are substantial while the uptake of gases directly onto the Teflon
walls is relatively effective.

The simulations were performed both with and without heterogeneous oxidation of unsaturated organic compounds using k_{0_3} and $D_{0,0_3}$ as model fitting parameters. The oxidation products (ox. prod.) from these reactions were assumed to form one organic semi-solid phase together with the other organic compounds ($D_{0,monomer}=5x10^{-17}$ cm² s⁻¹ and $D_{ox,prod}=0$ cm² s⁻¹).

In the model simulations presented in Fig. 14 it is shown that the model is able to capture the 9 volume loss rates of the seed aerosol and the onset of the SOA formation in the experiment (~0.5 10 hours after UV-lights were turned on). With a k_{0_3} of 10^{-16} molecules⁻¹ cm³ s⁻¹ and a $D_{0,0_3}$ of 10^{-8} 11 cm² s⁻¹ the model shows the best agreement with the observed particle volume concentration 12 more than 1.5 hours after the UV-light are turned on. However, for all simulations in Fig. 14, 13 ADCHAM underestimates the observed rapid SOA formation between 0.5 and 1.25 hours for the 14 particles suspended in air. Additionally, ADCHAM overestimates the total particle volume loss 15 rates of the suspended particle at the end of the experiments, especially without heterogeneous 16 ozonolysis and relatively rapid uptake of organic compounds onto the Teflon walls (Fig. 15b). 17 18 According to this simulation the particle losses are not only caused by deposition but also evaporation. Heterogeneous ozonolysis or other particle phase reactions allows more gas phase 19 20 monomers to partition into the particle phase and delay the time when the evaporation and deposition losses dominates over the SOA formation (see Fig. S21 in the supplementary 21 material). Additionally, the SOA formed from these particle phase reactions is less volatile (in the 22 model non-volatile) and will therefore decrease the evaporation loss rates (see Sect. 3.2). 23

Opposite to the simulation results in Fig. 14b, the maximum particle volume is larger without heterogeneous ozonolysis in Fig. 14a. For these simulations the SOA formation onto the wall deposited particles is more efficient ($\Delta x = 0.1$ cm) and the gas uptake onto the Teflon walls smaller. The wall deposited particles may not always serve as a sink of SOA but can also become a source of condensable organic compounds from the walls to the air. This is especially the case if the formed SOA is relatively volatile. The more volatile the SOA is, the smaller the SOA fraction found on the wall deposited particles will be. Hence, while the formed total SOA mass
 (air + walls) is larger with heterogeneous reactions in Fig. 14a, the SOA mass formed on the
 particles in the air is smaller (see also Fig. S22 in the supplementary material).

Figure S23 in the supplementary material compares the modelled particle volume from 4 simulations with the SIMPOL and Nannoolal vapour pressure method or the semi-empirical two 5 product model parameterization (see Sect. 3.4.2). The model simulations were performed with a 6 Δx of 0.1 cm and $k_{g,w}=1/20$ s⁻¹. For the simulations with the SIMPOL and the Nannoolal method, 7 heterogeneous ozonolysis was also considered ($k_{0_3} = 10^{-16}$ molecules⁻¹ cm³ s⁻¹ and $D_{0,0_3} = 10^{-8}$ cm² 8 s⁻¹). From this figure it is evident that both methods give almost identical SOA mass formation at 9 the end of the experiment. However, with the Nannoolal method the onset of the SOA formation 10 is approximately 15 minutes too late. The reason for this is that the modelled early stage SOA 11 formation is dominated by two MCM oxidation products (MXNCATECH and MXYMUCNO3) 12 (formed through the high NO oxidation pathway (see Sect. 3.4.3)). Both of these compounds 13 have higher vapour pressures with the Nannoolal method $(3.1 \times 10^{-3} \text{ and } 1.31 \times 10^{-4} \text{ Pa})$ compared 14 to the SIMPOL method $(1.9 \times 10^{-4} \text{ and } 7.5 \times 10^{-5} \text{ Pa}).$ 15

With the semi-empirical parameterization, derived from experiments in a similar but larger 16 Teflon chamber (28 m³) (Ng et al., 2007), ADCHAM gives a too early onset of the SOA 17 formation and overestimate the SOA formation when $k_{g,w}=1/20 \text{ s}^{-1}$. The reason for this is that the 18 three model compounds of this method all have relatively low vapour pressures (see Sect. 3.4.2). 19 Hence, the gas phase is rapidly saturated with respect to all these three compounds and they are 20 effectively taken up by the particles before they are lost to the Teflon wall surfaces. In order to 21 not overestimate the final SOA mass, $k_{g,w}$ need to be much larger ~1 s⁻¹. However, then the model 22 substantially underestimates the early stage SOA formation rate. 23

We also modelled the SOA formation without losses of condensable organic compounds onto the Teflon wall (see Fig. S24 in the supplementary material). With a Δx of 0.1 cm ADCHAM is now able to capture the rapid early stage SOA formation in the chamber. However, the final particle volume concentration in the air is overestimated with ~40 %. If we instead assume that the gas particle partitioning onto the chamber wall deposited particles is identical to the uptake onto the particles suspended in the air ($\Delta x = 0$ cm) (see Sect. 2.2.3 and references there in), the model again substantially underestimates the early stage SOA formation rate, while it gives reasonable
 particle volume concentrations at the end of the simulation.

3 Finally we also tested if a relatively rapid oligomerization process in the particle phase could improve the agreement with the modelled and measured SOA formation. For these simulations 4 we again use the SIMPOL vapour pressure method and assume that peroxyhemiacetal and 5 hemiacetal dimers form in the particle phase. The best agreement between the modelled and 6 measured SOA formation we find when we use a k_f of 10⁻²² molecules⁻¹ cm³ s⁻¹. This value of k_f 7 corresponds well with previously reported values of k_f for hemiacetal and peroxyhemiacetal 8 formation at weekly acidic conditions (pH≈4) (see Sect. 2.3.4). In order to shift the equilibrium 9 toward the particle phase (which might explain the rapid early stage SOA formation seen in the 10 11 experiment) we assume that the oligomers and monomers form one mixed phase.

Figure 15 shows the modelled particle volume concentrations when considering peroxyhemiacetal and hemiacetal dimer formation and with $\Delta x = 0$ or 0.1 cm and $k_{g,w} = 0$ or 1/15 s⁻¹. Without gas phase losses onto the Teflon walls and ideal uptake onto wall deposited particles ($\Delta x = 0$ cm) the model is able to capture the rapid early stage SOA formation seen in the experiment. After this the modelled particle volume concentration in the air continues to increase slowly for additionally ~2 hours, while in the experiment the measured particle volume slowly decreases.

With mass transfer limited diffusion and losses of condensable organic compounds from the near wall gas phase to the Teflon walls ($\Delta x = 0.1$ cm and $k_{g,w} = 1/15$ s⁻¹) the model results are in better agreement with the measurements in the end of the experiment and can nearly reproduce the rapid SOA formation in the beginning of the experiment.

Hence, these simulations indicate that relatively rapid heterogeneous reactions (either oligomerization or oxidation) are required in order to explain the observed rapid SOA formation in the beginning of the *m*-xylene oxidation experiment. Still, the model cannot fully explain the sharp transition between the rapid SOA formation between 0.5 and 1.25 hours after the UV-light is turned on and the slow almost linear volume (mass) loss observed during the latter half of the experiment.

In Fig. 16 we compare the temporal evolution of the modelled SOA formation without wall 1 2 losses to the chamber walls (ideal chamber), with the SIMPOL, Nannoolal or the semi-empirical parameterization method from Ng et al. (2007). The figure also illustrates the influence from 3 heterogeneous ozonolysis (O₃ ox.) of unsaturated organic compounds ($k_{O_2} = 10^{-16}$ molecules⁻¹ cm³ 4 s⁻¹ and $D_{0,0_2} = 10^{-8}$ cm² s⁻¹) and peroxyhemiacetal and hemiacetal oligomer formation ($k_f = 10^{-22}$ 5 molecules⁻¹ cm³ s⁻¹). We have also included the measured wall loss corrected SOA mass (SOA 6 mass scaled with the measured relative sulphate loss rate from the time when the UV-lights are 7 8 turned on) (see Sect. 2.2.3).

9 The simulation with SIMPOL and no heterogeneous reactions gives best agreement with the measured final SOA mass formation (70 and 65 μ g m⁻³, respectively). However, this simulation 10 substantially underestimates the SOA formation during the start of the experiment. The best 11 agreement between the model and measurements in the beginning of the experiment is instead 12 reached when we include relatively rapid oligomerization in the particle phase. The results from 13 14 this simulation also show surprisingly good agreement with the model simulation using the semiempirical parameterizations from Ng et al. (2007). This again indicates that heterogeneous 15 reactions are likely to be important for the SOA formation. The larger SOA formation from these 16 17 model simulations compared to the measurements can likely be attributed to substantial gas phase losses directly onto the Teflon walls in the chamber. This effect will be especially pronounced in 18 the end of the experiment when the surface area to volume ratio is large (see Sect. 3.4.1). Hence 19 for this experiment, the model simulations indicate that the wall corrections (which assume 20 continued uptake of condensable organic compounds onto the wall deposited particles) do not 21 give an upper estimate of the actual (atmospheric relevant) SOA formation (see Sect. 2.2.3). 22

23

24 4 Summary and conclusions

We have developed a novel aerosol dynamics, gas- and particle- phase chemistry model for chamber studies (ADCHAM). ADCHAM combines the detailed gas phase chemistry from MCMv3.2, a kinetic multilayer module for diffusion limited transport of compounds between the gas phase, particle surface and particle bulk phase, and an aerosol dynamics and particle phase chemistry module which is based on the ADCHEM model (Roldin et al., 2011a) but with important updates, among others process-based algorithms for: non-ideal interactions (salt
 effects) between water, organic and inorganic compounds, acidity catalysed oligomerization, and
 oxidation of organic compounds in the particle phase.

In this work we have illustrated the usefulness of ADCHAM in studying potentially influential but poorly known processes, i.e. different dimerization mechanisms, organic salt formation, salting-out effects, heterogeneous oxidation reactions and mass transfer limitations between the gas-particle phase, between the particle surface and particle bulk phase, and within the particle bulk phase. All these processes influence the modelled SOA formation and chemical and physical properties (e.g. volatility, phase state, oxidation state and hygroscopicity).

10 Additionally, we have also shown how ADCHAM can be used to study the influence of the 11 chamber wall effects on the SOA mass formation, evaporation properties, particle number size 12 distribution and gas phase chemistry. These effects are important to constrain because current 13 knowledge concerning SOA formation in the atmosphere is to a large extent based on smog 14 chamber experiments, and global climate models and chemistry transport models rely on 15 simplified semi-empirical parameterizations of SOA formation derived from these experiments.

16 The most important findings from the model simulations performed in this article are:

1) Our simulations of the α -pinene SOA evaporation experiments from Vaden et al. (2011) 17 supports the recent experimental findings that these particles are very viscous (tar like 18 19 amorphous SOA) (Virtanen et al., 2010, Vaden et al., 2010, Vaden et al., 2011, Kuwata and Martin, 2012, Zelenyuk et al., 2012, Abramson et al., 2013 and Zhou et al., 2013). In 20 21 these particles low-volatile dimers can accumulate in the particle surface-bulk layer upon 22 evaporation. With this dimer coating, ADCHAM is able to reproduce the main features of the observed slow evaporation rates if it is controlled by the reversible degradation of 23 dimers back to monomers. The model simulations illustrates that the mass fraction of 24 long-lived dimers needs to increase with decreasing particle size to explain the nearly size 25 26 independent evaporation rates. Because of the Kelvin effect, this can be accomplished if a considerable fraction of the dimers are formed in the gas phase or if they are formed in the 27 particle phase from the least volatile monomer compounds. The model simulations also 28 reveal that the dimer particle content and thus the observed evaporation rates of α -pinene 29

1 2 SOA particles may not only depend on chemical ageing but can also depend on the wall losses in chamber where the particles are formed.

- 2) The effect of $NH_3(g)$ on the α -pinene SOA properties and formation depends on: 1) the 3 reactive uptake of carboxylic acids and $NH_3(g)$ from the gas phase, 2) the viscosity of the 4 5 SOA particles (ammonium and organic compound diffusion rates) and 3) the salting-out effects of NH₄⁺. In order to distinguish between these effects we recommend future 6 experiments with AMS, in which the SOA particles are exposed to NH₃ in the absence of 7 gas phase carboxylic acids. In the model simulations the organic salts between ammonium 8 9 and carboxylic acids are involved in the initial growth of the particles. However, for atmospheric more relevant NH₃(g) and α -pinene concentrations, NH₃ has only a minor 10 influence on the uptake of carboxylic acids to the particle phase. Thus analogous to Yli-11 Juuti et al. (2013), our simulations indicate that it is unlikely that NH_3 and carboxylic 12 acids from α -pinene oxidation are responsible for the initial growth of nanometre sized 13 14 particles over the boreal forest.
- 3) Mass transfer limitations between the smog chamber air volume and the chamber walls 15 16 because of a thin laminar layer adjacent to the walls have large influence on the uptake of gases onto the wall deposited particles or directly onto the walls. If the formed SOA 17 material is semi-volatile the SOA particles on the chamber walls may even start to 18 evaporate and hence become a source of SOA at the end of smog chamber experiments. 19 20 Paradoxically, heterogeneous reactions which give less volatile SOA and generally more 21 SOA mass can increase the fraction of SOA which is found on the chamber walls and can thus even decrease the detectable SOA mass suspended in the chamber air volume. 22
- 4) In order to capture the rapid SOA formation observed during the oxidation of *m*-xylene in 23 24 the Lund University smog chamber we need to consider relatively rapid dimerization and/or some other heterogeneous reactions (e.g. ozonolysis of unsaturated organic 25 26 compounds). When considering peroxyhemiacetal and hemiacetal dimer formation in the particle phase, ADCHAM is able to capture both the observed early stage rapid SOA 27 formation in our own *m*-xylene experiment and gives almost identical SOA mass 28 formation as the semi-empirical parameterizations from Ng et al. (2007). This indicates 29 that heterogeneous particle phase reactions are not only important for the SOA properties 30 (e.g. volatility) but also for the concentration and formation rates. 31

Another more general conclusion, which can be drawn from the simulations performed in this 1 2 work, is that many of the parameters (processes) with large uncertainties (e.g. SOA viscosity, oligomerization rates and mechanisms, pure-liquid saturation vapour pressures, surface tension 3 and chamber wall effects) have large influence on the SOA formation and/or the chemical and 4 5 physical properties of the SOA. To be able to constrain the uncertainties related to these parameters (processes), the experiments need to be designed where as many variables as possible 6 7 are varied (e.g. time of aging, temperature, RH, concentrations, dilution, oxidation agents and light intensities). Apart from evaluating experimental results, ADCHAM can be used as a 8 9 valuable model tool when planning, designing and selecting which experiments and 10 instrumentation are needed in order to be able to answer specific research questions. The mxylene experiment studied in Sect. 3.4 is part of a larger experiment campaign designed in order 11 to study aging of anthropogenic SOA precursors and gasoline car exhausts (Nordin et al., 2013). 12 In that paper an early version of ADCHAM was used to study chamber wall effects, gas phase 13 chemistry and SOA formation before the experiments were performed. Currently we are applying 14 15 ADCHAM to study the aging of gasoline car exhausts and ELVOC formation from α -pinene ozonolysis. We have also started to implement many of the detailed processes (e.g. the kinetic 16 17 multilayer model, different dimerization processes and the detailed MCMv3.2 gas phase chemistry) in the ADCHEM model (Roldin et al., 2011a) which we use for detailed atmospheric 18 19 process studies.

20

21 Appendix A

22 **Table A1.** Nomenclature.

Symbol	Description
α_s	Surface-bulk accommodation coefficient
$\alpha_{0,s}$	Surface accommodation coefficient of surface free from adsorbing material
γ	Activity coefficient
γ _{w,i}	Activity coefficient of compound <i>i</i> in a Teflon wall film
$\mathcal{\delta}_{_k}$	Width of particle layer k
Δx	Laminar layer width adjacent to chamber walls or charcoal denuder

θ_{s}	Relative surface coverage of the adsorbed species
μ	Dynamic viscosity of air
$ ho_{air}$	Density of air
$ ho_p$	Particle phase density
σ	Surface tension of organic compounds
$ au_{d,Z_{ox}}$	Desorption lifetime of Z _{ox}
$\omega_{Z_{ox}}$	Mean thermal velocity of Z _{ox}
Ve	Characteristic average deposition velocity due to electrostatic forces
$\mathbf{\Omega}_{\scriptscriptstyle AB}^{(1,1)}$	Collision integral between
A_k	Area of exchange between particle layer <i>k</i> -1 and <i>k</i>
$A_{chamber}$	Chamber surface area
C_d	Dimer particle phase concentration
$\mathcal{C}_{\mathrm{H}^+}$	Hydrogen ion concentration
C_m	Monomer particle phase concentration
C_c	Cunningham slip correction factor
C_k	Kelvin effect
C_∞	Gas phase concentration far from the particle surfaces
C_s	Saturation gas phase concentration at the particle surface
C_w	Effective wall equivalent mass concentration
$d_{Z_{ox}}$	Width of the Z _{ox} sorption layer
d_{i}	Collision diameter of compound <i>i</i>
$d_{_{air}}$	Collision diameter of compound air molecules
$d_{_{i,air}}$	Collision diameter for binary collisions between compound <i>i</i> and air molecules
D_p	Particle diameter
D_{va}	Vacuum aerodynamic diameter
$D_{0,\mathrm{Xi}}$	Diffusivity coefficient of compound Xi without obstructing material
$D_{ m Xi}$	Diffusion coefficient of compound X _i
е	Elementary charge of a single proton
\overline{E}	Mean electrical field strength

$\overline{E_0}$	Initial mean electrical field strength
F	Fuchs-Sutugin correction factor in the transition region
f_p	Particle volume fraction of solid or semi-solid obstructing material
$F_{f}(\mathbf{p})$	Dimer formation rates in the particle phase
h	Distance between the roof and ceiling of the chamber
Ι	Molar condensation growth rate
$J_{ads,Z_{ox}}$	Adsorption rate of Z_{ox} to the sorption layer
$\boldsymbol{J}_{des, Z_{ox}}$	Desorption rate of Z_{ox} from the sorption layer
k_b	The Boltzmann constant
k_{charge}	First order deposition loss rate due to charge
k_d	First-order dimer specific degradation reaction rate constant
k_{f}	Second-order dimer formation rate constant
$k_{f,H+}$	Acid catalyzed third-order dimer formation rate constant
$k_{k,k+1,\mathrm{X_i}}$	Transport velocity of compound X_i between the layers k and layer $k+1$.
k_{Ox}	Oxidation reaction rate constant in the particle phase
$k_{so,su,Z_{ox}}$	Transport velocity of Z_{ox} from the sorption layer to the surface-bulk layer
$k_{su,so,Z_{ox}}$	Transport velocity of Z_{ox} from the surface-bulk layer to the sorption layer
$k_{g,w}$	First order loss rate from the near wall gas phase to the walls
$k_{w,g}$	Desorption rate from the chamber wall Teflon surfaces
k_w	Effective wall deposition loss rate
K_a	Acid dissociation constant
K_{H}	Henry's law constant
K_{s}	Solubility product of salt
K_s^*	Effective solubility product of organic salts
Kn	Non-dimensional Knudsen number
m_i	Molecular mass of compound <i>i</i>
m _{air}	Average air molecular mass
M_{i}	Molar mass of compound <i>i</i>

M_w	Average molar mass of a Teflon wall film
Ν	Number of elemental charges of a particle
N_a	Avogadros's number
р	Total pressure
p_0	Pure-liquid saturation vapour pressure
P_s	Equilibrium vapour pressure
pH	Negative 10-logarithm of the hydrogen ion concentration
pK_a	Negative 10-logarithm of the acid dissociation constant
R	Universal gas constant (8.3145 J K ⁻¹ mol ⁻¹)
RH	Relative humidity in %
t	Time
Т	Temperature in Kelvin
u [*]	Friction velocity
V_k	Volume of particle layer <i>k</i>
$V_{chamber}$	Chamber volume
V_{wall}	Air volume of a thin layer adjacent to the chamber walls
$\begin{bmatrix} V_{\mathrm{X}_{\mathrm{i}}} \end{bmatrix}_k$	Absolute volume concentration of compound X_i in particle layer k .
X	Mole fractions
X_k	Ratio between the smaller and larger of the two volume fluxes across A_k
Х	Condensable organic compound
$\left[\mathbf{X}_{\mathrm{i}}\right]_{k}$	Relative volume concentration of compound X_i in particle layer k .
$[\mathbf{X}_{\mathbf{i},g,w}]$	Concentrations of compound X_i in the thin layer adjacent to the chamber walls
$[X_{i,w}]$	Concentration of compound X _i on the chamber wall
Y	Organic compound formed by particle phase oxidation reaction
Z _{ox}	Oxidation agent in the particle phase (e.g. OH, O ₃ . NO ₃ and NO ₂)

1 Acknowledgements

2 This work was supported by the strategic research area MERGE at Lund University, Swedish
3 Research Council for Environment, Agricultural Sciences and Spatial Planning FORMAS

through projects 2007-1205, 2008-1467, 2009-615 and 2010-1678, the Swedish Research
Council through project 2006-5940 and by Metalund, the centre for Medicine and Technology for
Working Life and Society, a competence centre at Lund University, Sweden, supported by FAS,
the Swedish Council for Working Life and Social Research. Support for A. Zelenyuk was
provided by US Department of Energy (DOE), Office of Science, Office of Basic Energy
Sciences (BES), Chemical Sciences, Geosciences, and Biosciences Division.

The authors would like to thank prof. Gordon McFiggans research group at the University of 7 Manchester, and especially Dr. David Topping, for helpful discussions and for providing the 8 9 Python public script (now a available function called CompSysProp: http://ratty.cas.manchester.ac.uk/informatics/) to calculate Nannoolal based sub-cooled liquid 10 11 saturation vapour pressures for all included organic compounds in this paper.

2 References

- Abramson, E., Imre, D., Beránek, J., Wilson, J., Zelenyuk, A.: Experimental determination of
 chemical diffusion within secondary organic aerosol particles, Phys. Chem. Chem. Phys.,15,
 2983-2991, 2013
- Ammann, M. and Pöschl, U.: Kinetic model framework for aerosol and cloud surface chemistry
 and gas-particle interactions Part 2: Exemplary practical applications and numerical
 simulations, Atmos. Chem. Phys., 7, 6025–6045, 2007.
- 9 Barley, M. H. and McFiggans, G.: The critical assessment of vapour pressure estimation methods
 10 for use in modelling the formation of atmospheric organic aerosol, Atmos. Chem. Phys., 10, 74911 767, doi:10.5194/acp-10-749-2010, 2010.
- Barsanti, K. C., and Pankow, J. F: Thermodynamics of the formation of atmospheric organic
 particulate matter by accretion reactions—Part 1: aldehydes and ketones, Atmos. Environ., 38,
 4371–4382, 2004.
- Barsanti, K. C., and Pankow, J. F: Thermodynamics of the formation of atmospheric organic
 particulate matter by accretion reactions—Part 3: Carboxylic and dicarboxylic acids, Atmos.
 Environ., 40, 6676–6686, 2006.
- Barsanti, K. C., McMurry, P. H., and Smith, J. N.: The potential contribution of organic salts to
 new particle growth, Atmos. Chem. Phys., 9, 2949-2957, doi:10.5194/acp-9-2949-2009, 2009.
- Bergström, R., Denier van der Gon, H. A. C., Prévôt, A. S. H., Yttri, K. E., and Simpson, D.:
 Modelling of organic aerosols over Europe (2002–2007) using a volatility basis set (VBS)
 framework: application of different assumptions regarding the formation of secondary organic
 aerosol, Atmos. Chem. Phys., 12, 8499-8527, doi:10.5194/acp-12-8499-2012, 2012.
- Berkemeier, T., Huisman, A. J., Ammann, M., Shiraiwa, M., Koop, T., and Pöschl, U.: Kinetic
 regimes and limiting cases of gas uptake and heterogeneous reactions in atmospheric aerosols and
 clouds: a general classification scheme, Atmos. Chem. Phys., 13, 6663-6686, doi:10.5194/acp13-6663-2013, 2013.

1

Bertram, A. K., Martin, S. T., Hanna, S. J., Smith, M. L., Bodsworth, A., Chen, Q., Kuwata, M.,
Liu, A., You, Y., and Zorn, S. R.: Predicting the relative humidities of liquid-liquid phase
separation, efflorescence, and deliquescence of mixed particles of ammonium sulphate, organic
material, and water using the organic-to-sulphate mass ratio of the particle and the oxygen-tocarbon elemental ratio of the organic component, Atmos. Chem. Phys., 11, 10995-11006,
doi:10.5194/acp-11-10995-2011, 2011.

- Bloss, C., Wagner, V., Bonzanini, A., Jenkin, M. E., Wirtz, K., Martin-Reviejo, M., and Pilling,
 M. J.: Evaluation of detailed aromatic mechanisms (MCMv3 and MCMv3.1) against
 environmental chamber data, Atmos. Chem. Phys., 5, 623-639, 2005a.
- Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E.,
 Wirtz, K., Martin-Reviejo, M., Rea, G., Wenger, J. C., and Pilling, M. J.: Development of a
 detailed chemical mechanism (MCMv3.1) for the atmospheric oxidation of aromatic
 hydrocarbons, Atmos. Chem. Phys., 5, 641-664, doi:10.5194/acp-5-641-2005, 2005b.
- Boy, M., Hellmuth, O., Korhonen, H., Nilsson, E. D., ReVelle, D., Turnipseed, A., Arnold, F.,
 and Kulmala, M.: MALTE model to predict new aerosol formation in the lower troposphere,
 Atmos. Chem. Phys., 6, 4499-4517, doi:10.5194/acp-6-4499-2006, 2006.
- Camredon, M., Hamilton, J. F., Alam, M. S., Wyche, K. P., Carr, T., White, I. R., Monks, P. S.,
 Rickard, A. R., and Bloss, W. J.: Distribution of gaseous and particulate organic composition
 during dark α-pinene ozonolysis, Atmos. Chem. Phys., 10, 2893-2917, doi:10.5194/acp-10-28932010, 2010.
- Cappa, C. D. and Wilson, K. R.: Evolution of organic aerosol mass spectra upon heating:
 implications for OA phase and partitioning behavior, Atmos. Chem. Phys., 11, 1895-1911,
 doi:10.5194/acp-11-1895-2011, 2011.
- Casale, M. T, Richman, A. R., Elrod, M. J., Garland, R. M., Beaver M. R., and Tolbert, M. A.:
 Kinetics of acid-catalyzed aldol condensation reactions of aliphatic aldehydes Atmos. Environ.,
 2007, 41, 6212–6224.
- 27 Chapman, S., and Cowling, T. G.: The Mathematical Theory of Nonuniform Gases, Cambridge
- 28 University Press, Cambridge, 1970.

- Compernolle, S., Ceulemans, K., and Müller, J.-F.: Influence of non-ideality on condensation to
 aerosol, Atmos. Chem. Phys., 9, 1325–1337, doi:10.5194/acp-9-1325-2009, 2009.
- 2 = 100000, 10000, 100
- 3 DePalma, J. W., Horan, A. J., Hall, W. A., and Johnston, M. V.: Thermodynamics of oligomer
- 4 formation: implications for secondary organic aerosol formation and reactivity, Phys. Chem.
- 5 Chem. Phys., 15, 6935, 2013
- Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility
 basis set: 1. organic-aerosol mixing thermodynamics, Atmos. Chem. Phys., 11, 3303-3318,
 doi:10.5194/acp-11-3303-2011, 2011.
- Dzepina, K., Volkamer, R. M., Madronich, S., Tulet, P., Ulbrich, I. M., Zhang, Q., Cappa, C. D.,
 Ziemann, P. J., and Jimenez, J. L.: Evaluation of recently-proposed secondary organic aerosol
 models for a case study in Mexico City, Atmos. Chem. Phys., 9, 5681-5709, doi:10.5194/acp-95681-2009, 2009.
- Ervens, B. and Volkamer, R.: Glyoxal processing by aerosol multiphase chemistry: towards a
 kinetic modelling framework of secondary organic aerosol formation in aqueous particles,
 Atmos. Chem. Phys., 10, 8219-8244, doi:10.5194/acp-10-8219-2010, 2010.
- 16 Fuchs, N. A., Sutugin, A. G.: In Topics in Current Aerosol Research, Pergamon Press, 1971.
- Gao, S., Keywood, M., Ng, N. L., Surratt, J., Varutbangkul, V., Bahreini, R., Flagan, R. C., and
 Seinfeld, J. H.: Low-Molecular-Weight and Oligomeric Components in Secondary Organic
 Aerosol from the Ozonolysis of Cycloalkenes and α-Pinene, J. Phys. Chem. A, 108, 1014710164, 2004.
- George, C., Strekowski, R. S., Kleffmann, J., Stemmler, K., and Ammann, M.: Photoenhanced
 uptake of gaseous NO2 on solid organic compounds: a photochemical source of HONO?,
 Faraday Discuss., 130, 195–210, 2005.
- Greaves, T. L. and Drummond, C. J.: Protic ionic liquids: Properties and applications, Chem.
 Rev., 108, 206–237, 2008.

- 1 Grieshop, A. P., Donahue, N. M., and Robinson, A. L.: Is the gas-particle partitioning in alpha-
- 2 pinene secondary organic aerosol reversible?, Geophys. Res. Lett., 34, L14810,
- 3 doi:10.1029/2007GL029987, 2007.
- Griffin, R. J., Cocker III, D. R., Flagan, R. C., and Seinfeld, J. H.: Organic aerosol formation
 from the oxidation of biogenic hydrocarbons, J. Geophys. Res., 104, 3555–3567, 1999.
- Gutzwiller, L., Arens, F., Baltensperger, U., G^{*}aggeler, H. W., and Ammann, M.: Significance of
 Semivolatile Diesel Exhaust Organics for Secondary HONO Formation, Environ. Sci. Technol.,
 36, 677–682, 2002.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen,
 J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T.,
 Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W.,
 McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D.,
 Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic
 aerosol: current and emerging issues, Atmos. Chem. Phys., 9, 5155–5236, doi:10.5194/acp-95155-2009, 2009.
- 16 Hansen, H. K., Rasmussen, P., Fredenslund, A., Schiller, M., and Gmehling, J.: Vapour–liquid
- equilibria by UNIFAC group contribution. 5. Revision and extension, Ind. Eng. Chem. Res., 30,
 2352–2355, 1991.
- 19 Hildebrandt, L., Donahue, N. M. and Pandis, S. N.: High formation of secondary organic aerosol
- 20 from the photo-oxidation of toluene. Atmos. Chem. Phys. 9, 2973-2986, 2009.
- Hirschfelder, J. O., Curtiss, C. F., and Bird, R. B.: Molecular Theory of Gases and Liquids. John
 Wiley, New York, 1954.
- 23 Hoffmann, T., Odum, J. R., Bowman, F., Collins, D., Klockow, D., Flagan, R. C., and Seinfeld, J.
- 24 H.: Formation of organic aerosols from the oxidation of biogenic hydrocarbons, J. Atmos. Chem.,
- 25 26, 189–222, 1997.

Hu, D., Tolocka, M., Li, Q., and Kamens, R. M.: A kinetic mechanism for predicting secondary
 organic aerosol formation from toluene oxidation in the presence of NOx and natural sunlight,
 Atmos. Environ., 41, 6478–6496, 2007.

4 Huff Hartz, K. E. H., Weitkamp, E. A., Sage, A. M., Donahue, N. M., and Robinson, A. L.:

5 Laboratory measurements of the oxidation kinetics of organic aerosol mixtures using a relative

6 rate constants approach, J. Geophys. Res.-Atmos., 112, D04204, doi:10.1029/2006jd007526,

7 2007.

8 Hyder, M., Genberg, J., Jönsson, J. Å.: Application of hollow fiber liquid phase microextraction

9 for pinic acid and pinonic acid analysis from organic aerosols, Analytica Chimica Acta, 713, 79–
10 85, 2012.

Iinuma, Y., Böge, O., Gnauk, T., and Herrmann, H.: Aerosolchamber study of the α-pinene/O3
reaction: Influence of particle acidity on aerosol yields and products, Atmos. Environ., 38, 761–
773, 2004.

Jacobson, M. Z.: Fundamentals of Atmospheric Modelling (2nd edition), Cambridge University
Press, Cambridge, United Kingdom and New York, NY, USA, ISBN: 0 521 54865 9, 2005a.

16 Jacobson, M. Z.: A Solution to the Problem of Nonequilibrium Acid/Base Gas-Particle Transfer

17 at Long Time Step, Aerosol Sci. Technol., 39, 92–103, 2005b.

Jenkin, M. E., Saunders, S. M, and Pilling, M. J.: The tropospheric degradation of volatile
organic compounds: A protocol for mechanism development, Atmos. Environ., 31, 81-104, 1997.

Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of
the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic
volatile organic compounds, Atmos. Chem. Phys., 3, 181-193, doi:10.5194/acp-3-181-2003,
2003.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H.,
DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M.,
Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin,

27 C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M.,

- 1 Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A.,
- 2 Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F.,
- 3 Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A.,
- 4 Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y, Zhang, Y. M., Dzepina, K., Kimmel, J. R.,
- 5 Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C.,
- 6 Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic
- 7 Aerosols in the Atmosphere, Science, 326, 1525-1529, 2009.
- Johnson, D., Jenkin, M.E., Wirtz, K., Martin-Reviejo, M.: Simulating the formation of secondary
 organic aerosol from the photooxidation of aromatic hydrocarbons. Environmental Chemistry 2,
 35–48, 2005.
- 11 Johnson, D., Utembe, S. R., and Jenkin, M. E.: Simulating the detailed chemical composition of

12 secondary organic aerosol formed on a regional scale during the TORCH 2003 campaign in the

- 13 southern UK, Atmos. Chem. Phys., 6, 419–431, 2006.
- Jonsson, Å. M., Hallquist, M., and Ljungström, E.: The effect of temperature and water on
 secondary organic aerosol formation from ozonolysis of limonene, Δ³-carene and α-pinene,
 Atmos. Chem. Phys., 8, 6541-6549, doi:10.5194/acp-8-6541-2008, 2008.
- 17 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prévôt, A. S. H., Fisseha, R.,
- 18 Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as
- 19 major components of atmospheric organic aerosols, Science, 303, 1659–1662, 2004.
- Knopf, D. A., Anthony, L. M., and Bertram, A. K.: Reactive uptake of O3 by multicomponent
 and multiphase mixtures containing oleic acid, J. Phys. Chem. A, 109, 5579–5589, 2005.
- 22 Koop, T., Bookhold, J., Shiraiwa, M., and Pöschl, U.: Glass transition and phase state of organic
- 23 compounds: dependency on molecular properties and implications for secondary organic aerosols
- 24 in the atmosphere, Phys. Chem. Chem. Phys., 13, 19238–19255, 2011.
- 25 Korhonen, H., Lehtinen, K. E. J., and Kulmala, M.: Multicomponent aerosol dynamics model
- 26 UHMA: modle development and validation, Atmos. Chem. Phys., 4, 471–506, 2004.

- Kroll, J., and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of
 low-volatility organics in the atmosphere, Atmos. Env., 42, 3593–3624, 2008.
- Kuwata, M., and Martin, S., T.: Phase of atmospheric secondary organic material affects its
 reactivity, PNAS, 109, 17354-17359, doi/10.1073/pnas.1209071109, 2012.
- Lai, A. and Nazaroff, W.W.: Modelling indoor particle deposition from turbulent flow onto
 smooth surfaces, J. Aerosol Sci., 31, 463-476, 2000.
- Li, Q., Hu, D., Leungsakul, S., and Kamens, R. M.: Large outdoor chamber experiments and
 computer simulations: (I) Secondary organic aerosol formation from the oxidation of a mixture of
 d- limonene and α-pinene, Atmos. Environ., 41(40), 9341–9352, 2007.
- Lide, D. R. (Ed.): CRC Handbook of chemistry and physics, Taylor and Francis Group, Boca
 Raton, FL, USA, 88th Edition, 2008.
- Liggio, J. and Li, S.-M.: Organosulfate formation during the uptake of pinonaldehyde in acidic
 sulphate aerosols, Geophys. Res. Lett., 33, L13808, doi:10.1029/2006GL026079, 2006.
- Loza, C. L., Chhabra, P. S., Yee, L. D., Craven, J. S., Flagan, R. C., and Seinfeld, J. H.: Chemical
 aging of *m*-xylene secondary organic aerosol: laboratory chamber study, Atmos. Chem. Phys.,
 12, 151-167, doi:10.5194/acp-12-151-2012, 2012.
- Maksymiuk, C. S., Gayahtri, C., Gil, R. R., and Donahue, N. M.: Secondary organic aerosol
 formation from multiphase oxidation of limonene by ozone: mechanistic constraints via twodimensional heteronuclear NMR spectroscopy. Phys. Chem. Chem. Phys., 2009, 11, 7810–7818,
 2009.
- 21 Marcolli, C. and Peter, T.: Water activity in polyol/water systems: new UNIFAC
- 22 parameterization, Atmos. Chem. Phys., 5, 1545–1555, 2005.
- 23 Matsunaga, A., and Ziemann, P.J.: Gas-Wall Partitioning of Organic Compounds in a Teflon
- 24 Film Chamber and Potential Effects on Reaction Product and Aerosol Yield Measurements,
- 25 Aerosol Sci. Technol., 44, 881-892, 2010.

- McFiggans, G., Topping, D. O., and Barley, M. H.: The sensitivity of secondary organic aerosol
 component partitioning to the predictions of component properties Part 1: A systematic
 evaluation of some available estimation techniques, Atmos. Chem. Phys., 10, 10255–10272,
 2010.
- McMurry, P. H. and Rader, D. J.: Aerosol wall losses in electrically charged chambers. Aerosol
 Sci. Technol., 4, 249-268, 1985.
- Metzger, A., Dommen, J., Gaeggeler, K., Duplissy, J., Prévôt, A. S. H., Kleffmann, J.,
 Elshorbany, Y., Wisthaler, A., and Baltensperger, U.: Evaluation of 1,3,5 trimethylbenzene
 degradation in the detailed tropospheric chemistry mechanism, MCMv3.1, using environmental
 chamber data, Atmos. Chem. Phys., 8, 6453–6468, 2008.
- Na, K., Song, C., Switzer, C., and Cocker, D.: Effect of Ammonia on Secondary Organic Aerosol
 Formation from α-Pinene Ozonolysis in Dry and Humid Conditions. Environ. Sci. Technol., 41,
 6096-6102, 2007.
- Nannoolal, J., Rarey, J., Ramjugernath, D.: Estimation of pure component properties Part 3.
 Estimation of the vapour pressure of non-electrolyte organic compounds via group contributions
 and group interactions Fuild Phase Equilibria, 269, 117-133, 2008.
- Nash, D. G., Tolocka, M. P., and Baer, T.: The uptake of O₃ by myristic acid-oleic acid mixed
 particles: evidence for solid surface layers, Phys. Chem. Chem. Phys., 8, 4468–4475,
 doi:10.1039/b609855j, 2006.
- Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.:
 Secondary organic aerosol formation from *m*-xylene, toluene, and benzene, Atmos. Chem. Phys.,
 7, 3909–3922, 2007.
- 23 Ng, N. L., Kwan, A. J., Surratt, J. D., Chan, A. W. H., Chhabra, P. S., Sorooshian, A., Pye, H. O.
- 24 T., Crounse, J. D., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol
- (SOA) formation from reaction of isoprene with nitrate radicals (NO3), Atmos. Chem. Phys., 8,
 4117–4140, 2008.
- Nordin, E. Z., Eriksson, A. C., Roldin, P., Nilsson, P. T., Carlsson, J. E., Kajos, M. K.,
 Hellén, H., Wittbom, C., Rissler, J., Löndahl, J., Swietlicki, E., Svenningsson, B., Bohgard, M.,
 Kulmala, M., Hallquist, M., and Pagels, J. H.: Secondary organic aerosol formation from idling
 gasoline passenger vehicle emissions investigated in a smog chamber, Atmos. Chem. Phys., 13,
- 5 6101-6116, doi:10.5194/acp-13-6101-2013, 2013.
- Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H.:
 Gas/particle partitioning and secondary organic aerosol yields, Environ. Sci. Technol., 30, 2580–
 2585, 1996.
- 9 Pankow, J. F.: An absorption-model of the gas aerosol partitioning involved in the formation of
 10 secondary organic aerosol, Atmos. Environ., 28, 189–193, 1994.
- 11 Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting

vapour pressures and enthalpies of vaporization of multifunctional organic compounds, Atmos.
Chem. Phys., 8, 2773–2796, 2008.

- Pathak, R. K., Presto, A. A., Lane, T. E., Stanier, C. O., Donahue, N. M., and Pandis, S. N.:
 Ozonolysis of α-pinene: parameterization of secondary organic aerosol mass fraction, Atmos.
 Chem. Phys., 7, 3811-3821, doi:10.5194/acp-7-3811-2007, 2007.
- Paulsen, D., Dommen, J., Kalberer, M., Prevot, A. S. H., Richter, R., Sax, M., Steinbacher,
 M.,Weingartner, E., and Baltensperger, U.: Secondary organic aerosol formation by irradiation of
 1,3,5-trimethylbenzene-NOx-H2O in a new reaction chamber for atmospheric chemistry and
 physics, Environ. Sci. Technol., 39, 2668–2678, 2005.
- Pfrang, C., Shiraiwa, M., and Pöschl, U.: Chemical ageing and transformation of diffusivity in
 semi-solid multi-component organic aerosol particles, Atmos. Chem. Phys., 11, 7343-7354,
 doi:10.5194/acp-11-7343-2011, 2011.
- Pierce, J. R., Engelhart, G. J., Hildebrandt, L., Weitkamp, E. A., Pathak, R. K., Donahue, N. M.,
 Robinson, A. L., Adams, P. J., and Pandis, S. N.: Constraining Particle Evolution from Wall
 Losses, Coagulation, and Condensation-Evaporation in Smog-Chamber Experiments: Optimal
 Estimation Based on Size Distribution Measurements, Aerosol Sci. Technol., 42:12, 1001-1015,
 2008.

Pun, B. K. and Seigneur, C.: Investigative modelling of new pathways for secondary organic
 aerosol formation, Atmos. Chem. Phys., 7, 2199-2216, doi:10.5194/acp-7-2199-2007, 2007.

Pöschl, U., Rudich, Y., and Ammann, M.: Kinetic model framework for aerosol and cloud
surface chemistry and gas-particle interactions – Part 1: General equations, parameters, and
terminology, Atmos. Chem. Phys., 7, 5989–6023, 2007.

- Pöschl, U.: Gas-particle interactions of tropospheric aerosols: Kinetic and thermodynamic
 perspectives of multiphase chemical reactions, amorphous organic substances, and the activation
 of cloud condensation nuclei, Atmos. Res. 101, 562–573, 2011.
- 9 Rader, D. J., McMurry, P. H. and Smith, S.: Evaporation Rates of Monodisperse Organic
- 10 Aerosols in the 0.02- to 0.2-µm-Diameter Range, Aerosol Sci. Technol., 6,247-260, 1987.
- 11 Ray, A. K., Davis, E. J., and Ravindran, P.: Determination of ultralow vapour-pressures by sub-
- 12 micron droplet evaporation, J. Chem. Phys. 71, 582-587, 1979.
- Ray, A. K, Lee, J., and Tilley, H. L.: Direct Measurements of Evaporation Rates of Single
 Droplets at Large Knudsen Numbers, Langmuir, 4, 631-637, 1988.
- 15 Rickard, A. R., Wyche, K.P., Metzger, A., Monks, P.S., Ellis, A.M., Dommenc, J., Baltensperger,
- 16 U., Jenkin, M.E., Pilling, M.J.: Gas phase precursors to anthropogenic secondary organic aerosol:
- 17 Using the Master Chemical Mechanism to probe detailed observations of 1,3,5-trimethylbenzene
- 18 photo-oxidation, Atmos. Env. 44, 5423–5433, 2010.
- Riipinen, I., Pierce, J. R., Donahue, N. M, Pandis, S. N.: Equilibration time scales of organic
 aerosol inside thermodenuders: Evaporation kinetics versus thermodynamics. Atmos Environ
 44:597–607, 2010.
- 22 Rohrer, F., Bohn, B., Brauers, T., Bruning, D., Johnen, F. J., Wahner, A., and Kleffmann, J.:
- 23 Characterisation of the photolytic HONO-source in the atmosphere simulation chamber SAPHIR,
- 24 Atmos. Chem. Phys., 5, 2189–2201, 2005.

- Roldin, P., Swietlicki, E., Schurgers, G., Arneth, A., Lehtinen, K. E. J., Boy, M., and Kulmala,
 M.: Development and evaluation of the aerosol dynamics and gas phase chemistry model
 ADCHEM, Atmos. Chem. Phys., 11, 5867-5896, 2011a.
- Roldin, P., Swietlicki, E., Massling, A., Kristensson, A., Löndahl, J., Eriksson, A., Pagels, J., and
 Gustafsson, S.: Aerosol ageing in an urban plume implication for climate, Atmos. Chem. Phys.,
- 6 11, 5897–5915, 2011b.
- Rudich, Y., Donahue, N. M., and Mentel, T. F.: Aging of organic aerosol: Bridging the gap
 between laboratory and field studies, Annu. Rev. Phys. Chem., 58, 321–352, 2007.
- 9 Saukko, E., Lambe, A. T., Massoli, P., Koop, T., Wright, J. P., Croasdale, D. R., 10 Pedernera, D. A., Onasch, T. B., Laaksonen, A., Davidovits, P., Worsnop, D. R., and Virtanen, A.: Humidity-dependent phase state of SOA particles from biogenic and anthropogenic 11 precursors, Atmos. Chem. Phys., 12, 7517-7529, doi:10.5194/acp-12-7517-2012, 2012. 12
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development
 of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic
 volatile organic compounds, Atmos. Chem. Phys., 3, 161-180, doi:10.5194/acp-3-161-2003,
 2003.
- Seinfeld, J. H., and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to
 Climate Change, (2nd edition), Wiley, New Jersey, ISBN: 0-471-72018-6, 2006.
- Shiraiwa, M., Pfrang, C., and Pöschl, U.: Kinetic multi-layer model of aerosol surface and bulk
 chemistry (KM-SUB): the influence of interfacial transport and bulk diffusion on the oxidation of
 oleic acid by ozone, Atmos. Chem. Phys., 10, 3673-3691, doi:10.5194/acp-10-3673-2010, 2010.
- Shiraiwa, M., Ammann, M., Koop, T., Pöschl, U.: Gas uptake and chemical aging of semisolid
 organic aerosol particles, Proc. Natl Acad. Sci., 108, 1100311008 doi:10.1073/pnas.1103045108, 2011.
- Shiraiwa, M., Pfrang, C., Koop, T., and Pöschl, U.: Kinetic multi-layer model of gas-particle
 interactions in aerosols and clouds (KM-GAP): linking condensation, evaporation and chemical
 reactions of organics, oxidants and water, Atmos. Chem. Phys., 12, 2777–2794, 2012.

- Shiraiwa, M., and Seinfeld, J. H.: Equilibration timescale of atmospheric secondary organic
 aerosol partitioning, Geophys. Res. Lett., 39, L24801, 2012.
- Shiraiwa, M., Yee, L. D., Schilling, K. A., Loza, C. L., Craven, J. S., Zuend, A., Ziemann, P. J.,
 and Seinfeld, J. H.: Size distribution dynamics reveal particle-phase chemistry in organic aerosol
- 5 formation, Proc. Natl Acad. Sci., 110, 11746-11750, 2013a.
- 6 Shiraiwa, M., Zuend, A., Bertram, A. K., and Seinfeld, J. H.: Gas-particle partitioning of
 7 atmospheric aerosols: interplay of physical state, non-ideal mixing and morphology, Phys. Chem.
 8 Chem. Phys., 15, 11441-11453, 2013b.
- 9 Smith, J. N., Dunn, M. J., VanReken, T. M., Iida, K., Stolzenburg, M. R., McMurry, P. H., and
 10 Huey, L. G.: Chemical composition of atmospheric nanoparticles formed from nucleation in
 11 Tecamac, Mexico: Evidence for an important role for organic species in nanoparticle growth,
- 12 Geophys. Res. Lett., 35, L04808, doi:10.1029/2007gl032523, 2008.
- Smith, J. N., Barsanti, K. C, Friedli, H. R., Ehn, M., Kulmala, M., Collins, D. R., Scheckman, J.
 H., Williams, B. J., and McMurry, P. H.: Observations of aminium salts in atmospheric
 nanoparticles and possible climatic implications, Proc. Natl Acad. Sci. 107, 6634–6639, 2010.
- Smith, M. L., Kuwata, M., and Martin, S. T.: Secondary Organic Material Produced by the Dark
 Ozonolysis of α-Pinene Minimally Affects the Deliquescence and Efflorescence of Ammonium
 Sulphate, Aerosol Sci. Technol., 45, 244–261, doi:10.1080/02786826.2010.532178,
 http://dx.doi.org/10.1080/02786826.2010.532178, 2011.
- Stroeve, P.: On the Diffusion of Gases in Protein Solutions, Ind. Eng. Chem. Fund., 14, 140-141,
 1975.
- Surratt, J. D, Murphy, S. M, Kroll, J. H, Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski,
 R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, F. C., and Seinfeld, J. H.: Chemical
 composition of secondary organic aerosol formed from the photooxidation of isoprene, J. Phys.
 Chem. A, 110, 9665-9690, 2006

- 1 Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L.,
- 2 Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for
- 3 organosulfates in secondary organic aerosol, Environ. Sci. Technol., 41, 517–527, 2007.
- Svenningsson, B., Rissler, J., Swietlicki, E., Mircea, M., Bilde, M., Facchini, M. C., Decesari, S.,
 Fuzzi, S., Zhou, J., Mønster, J., and Rosenørn, T.: Hygroscopic growth and critical
 supersaturations for mixed aerosol particles of inorganic and organic compounds of atmospheric
 relevance, Atmos. Chem. Phys., 6, 1937-1952, doi:10.5194/acp-6-1937-2006, 2006.
- 8 Tobias, H. J, and Ziemann, P. J.: Thermal desorption mass spectrometric analysis of organic
 9 aerosol formed from reactions of 1-tetradecene and O-3 in the presence of alcohols and
 10 carboxylic acids, Environ. Sci. Technol., 34, 2105–2115, 2000.
- Tolocka, M. P., Jang, M., Ginter, J. M., Cox, F. J., Kamens R. M., and Johnston, M. V.:
 Formation of Oligomers in Secondary Organic Aerosol, Environ. Sci. Technol., 38, 1428-1434,
 2004.
- Topping, D., Barley, M., and McFiggans, G.: Including phase separation in a unified model to
 calculate partitioning of vapours to mixed inorganic–organic aerosol particles, Faraday Discuss.,
 165, 273, 2013.
- 17 Vaden, T. D., Song, C., Zaveri, R. A., Imre, D., Zelenyuk, A.: Morphology of mixed primary
- and secondary organic particles and the adsorption of spectator organic gases during aerosol
 formation, Proc. Natl. Acad. Sci., 107, 6658–6663, 2010.
- Vaden, T. D., Imreb, D., Beráneka, J., Shrivastavaa, M., and Zelenyuka, A.: Evaporation
 kinetics and phase of laboratory and ambient secondary organic aerosol, Proc. Natl. Acad. Sci.,
 108, 2190-2195, 2011.
- 23 Valorso, R., Aumont, B., Camredon, M., Raventos-Duran, T., Mouchel-Vallon, C., Ng, N. L.,
- 24 Seinfeld, J. H., Lee-Taylor, J., and Madronich, S.: Explicit modelling of SOA formation from α-
- pinene photooxidation: sensitivity to vapour pressure estimation. Atmos. Chem. Phys., 11, 6895–
 6910, 2011.

- 1 Vesterinen, M., Lehtinen, K. E. J., Kulmala, M., and Laaksonen, A.: Effect of particle phase
- 2 oligomer formation on aerosol growth, Atmos. Environ., 41, 1768–1776, 2007.
- 3 Virtanen, A., Joutsensaari, J., Koop, T., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J.
- 4 K., Pöschl, U., Kulmala, M., Worsnop, D. R. and Laaksonen, A.: An amorphous solid state of
- 5 biogenic secondary organic aerosol particles, Nature, 467, 824-827, 2010.
- Widmann, J, Heusmann, C. M. and Davis, E. J..: The effect of a polymeric additive on the
 evaporation of organic aerocolloidal droplets, Colloid and Polymer Science, 276, 197-205, 1998.
- 8 Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., 9 Tuch, T., Pfeifer, S., Fiebig, M., Fjäraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., 10 Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., 11 12 Grüning, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G., 13 O'Dowd, C. D., Marinoni, A., Horn, H.-G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., 14 Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Löschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to 15 facilitate high quality long-term observations of atmospheric particle number size distributions, 16 Atmos. Meas. Tech., 5, 657-685, doi:10.5194/amt-5-657-2012, 2012. 17
- Yli-Juuti, T., Barsanti, K., Hildebrandt Ruiz, L., Kieloaho, A.-J., Makkonen, U., Petäjä, T.,
 Ruuskanen, T., Kulmala, M. and Riipinen, I.: Model for acid-base chemistry in nanoparticle
 growth (MABNAG), Atms. Chem. Phys. Discuss., 13, 7175-7222, 2013.
- Zelenyuk, A., Yang, J., Song, C., Zaveri, R. A. and Imre, D.: A New Real-Time Method for
 Determining Particles' Sphericity and Density: Application to Secondary Organic Aerosol
 Formed by Ozonolysis of α-Pinene, Environ. Sci. Technol., 42, 8033-8038, 2008.
- Zelenyuk, A., Imre, D., Beránek, J., Abramson, E., Wilson, J. and Shrivastava, M.: Synergy
 between Secondary Organic Aerosols and Long-Range Transport of Polycyclic Aromatic
 Hydrocarbons, Environ. Sci. Technol., 46, 12459-12466, 2012.
- Zhang, S-H., Seifeld, J. H., and Flagan, R. C.: Determination of Particle Vapour Pressures Using
 the Tandem Differential Mobility Analyzer, Aerosol Sci. Technol., 19,3-14, 1993.

- Zhou, S., Shiraiwa, M., McWhinney, R., Pöschl, U., and Abbatt, J. P. D.: Kinetic limitations in
 gas-particle reactions arising from slow diffusion in secondary organic aerosol, Faraday Discuss.,
 165, 391-406, 2013.
- Zobrist, B., Soonsin, V., Luo, B. P., Krieger, U. K., Marcolli, C., Peter, T., and Koop, T.: Ultraslow water diffusion in aqueous sucrose glasses, Phys. Chem. Chem. Phys., 13, 3514–3526, doi:10.1039/c0cp01273d, 2011.
- Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organicinorganic aerosols to predict activity coefficients, Atmos. Chem. Phys., 8, 4559–4593,
 doi:10.5194/acp-8-4559-2008, 2008.
- Zuend, A., Marcolli, C., Peter, T., and Seinfeld, J. H.: Computation of liquid-liquid equilibria and
 phase stabilities: implications for RH-dependent gas/particle partitioning of organic-inorganic
 aerosols, Atmos. Chem. Phys., 10, 7795-7820, doi:10.5194/acp-10-7795-2010, 2010.
- Zuend, A., Marcolli C., Booth , A. M., Lienhard, D. M., Soonsin, V., Krieger, U. K., Topping, D.
 O., McFiggans G., Peter, T., and Seinfeld, J. H.: New and extended parameterization of the
 thermodynamic model AIOMFAC: calculation of activity coefficients for organic-inorganic
 mixtures containing carboxyl, hydroxyl, carbonyl, ether, ester, alkenyl, alkyl, and aromatic
 functional groups., Atmos. Chem. Phys., 11, 9155–9206, 2011.
- Zuend, A. and Seinfeld, J. H.: Modelling the gas-particle partitioning of secondary organic
 aerosol: the importance of liquid-liquid phase separation, Atmos. Chem. Phys., 12, 3857-3882,
 doi:10.5194/acp-12-3857-2012, 2012.
- 21

Parameter	Definition	Value
$\alpha_{s,0,0_3}$	Surface accommodation coefficient of O_3 on a free substrate	1 ^a
$ au_{d,\mathrm{O}_3}(\mathrm{s})$	O ₃ desorption lifetime	10 ^{-9 a}
$K_{\rm H,O_3} (\rm mol \ m^3 \ Pa^{-1})$	Henry's law coefficient of O ₃	$4.7 \times 10^{-3a^*}$
$D_{0,0_3}$ (cm ² s ⁻¹)	Bulk diffusion coefficient O ₃ without obstruction	Variable
ω_{0_3} (cm s ⁻¹)	Mean thermal velocity O ₃	3.6x10 ^{4a}
<i>d</i> ₀₃ (nm)	Effective diameter cross section O ₃	0.4 ^a
k_{0_3} (molec ⁻¹ cm ³ s ⁻¹)	Reaction rate constant between O_3 and organic comp.	Variable

1 Table 1. Model parameters used in the multilayer module for O_3 uptake, diffusion and reactions 2 in the particle phase.

^a Values from Pfrang et al. (2011), ^{*}Different unit than in Pfrang et al. (2011)

4

1	Table 2. Summary of which processes and parameter values that were used for the simulations
2	presented in Sect. 3.1 to 3.4.

Parameter (unit)	Sect. 3.1	Sect. 3.2	Sect. 3.3	Sect. 3.4	
Gas-wall losses	Yes	Yes	No	Yes	
Δx (cm)	^a 0.1	^b 0.1	-	0.1 or 1	
$V_{chamber}$ (m ³)	-	0.1	-	^c 6	
V_{wall} (m ³)	-	-	-	^d 0.02	
$k_{g,w}(s^{-1})$	-	-	-	0-1/6	
$k_{g,w}^{*}(s^{-1})$	-	0-1/500	-	-	
$C_{w}/(M_{w}\gamma_{w,i}) \;(\mu \mathrm{mol}\;\mathrm{m}^{-3})$	-	100	-	100	
Particle-wall deposition	No	No	No	Yes	
\overline{E} (V cm ⁻¹)	-	-	-	50 or Eq. 30	
u^{*} (m s ⁻¹)	-	-	-	0.01 or 0.05	
Coagulation	No	No	No	Yes	
Homogeneous nucleation	No	^e Yes	^e Yes	No	
Organic salt formation	No	No	Yes (Table 3)	No	
Condensation/Evaporation	Yes	Yes	Yes	Yes	
α_s	1	1	1	1	
σ (N m ⁻¹)	0.05	0.05	0.05	0.05	
Kinetic multilayer model	Yes	Yes	Yes	3-layer model	
$D_{0,monomer}$ (cm ² s ⁻¹)	0 or 10 ⁻¹⁰	$5 \times 10^{-17} - 10^{-13}$	0-10 ⁻¹⁵	5×10^{-17}	
$D_{0,dimer} ({\rm cm}^2{ m s}^{-1})$	-	0-10 ⁻¹⁵	-	0	
$D_{0,ammonium} (\mathrm{cm}^2 \mathrm{s}^{-1})$	-	-	0-10 ⁻¹³	-	
$D_{0,\rm NH4RCOO}~(\rm cm^2~s^{-1})$	-	-	0	-	

$D_{0,O_3} \ (\mathrm{cm}^2 \ \mathrm{s}^{-1})$	-	-	-	10 ⁻⁷ or 10 ⁻⁸
$D_{0,\rm NO_2}~(\rm cm^2~s^{-1})$	-	-	-	10 ⁻⁷
Heterogeneous oxidation	No	No	No	Yes
k_{0_3} (molec ⁻¹ cm ³ s ⁻¹)	-	-	-	0-10 ⁻¹⁶
k_{NO_2} (molec ⁻¹ cm ³ s ⁻¹)	-	-	-	0 or 10 ⁻¹⁵
Particle phase dimerization	No	Yes	No	Yes
k_f peroxyhemiacetals (molec ⁻¹ cm ³ s ⁻¹)	-	10 ⁻²⁴ -10 ⁻²¹	-	0-10 ⁻²¹
k_f hemiacetal (molec ⁻¹ cm ³ s ⁻¹)	-	-	-	0-10 ⁻²¹
k_d peroxyhemiacetals (h ⁻¹)	-	1/40-30	-	0
k_d hemiacetal (h ⁻¹)	-	-	-	0

^a Value used for the gas uptake onto the charcoal denuder.

2 ^b Same value used for the charcoal denuder and the Teflon chamber walls.

3 ^c Initial value. During the experiments $V_{chamber}$ gradually decreases.

4 ^d Derived with the assumption that the width of the thin air layer adjacent to the chamber walls is 1 mm.

⁶ Represented by adding new size bins with an initial particle diameter of 5 nm (see Sect. 3).

Daramatar	Definition	Value ^a
I diameter	Definition	v aluc
рК _{а,СООН}	Logarithm of carboxylic acid dissociation constant	4.6
pK _{a,NH3}	Logarithm of NH_4^+ dissociation constant	9.25 ^b
$K_s^* (\mathrm{mol}^2 \mathrm{m}^{-6})$	Effective solubility product (see Eq. 32)	0.1
$K_H (\mathrm{mol} \mathrm{m}^{-3} \mathrm{atm}^{-1})$	Henry's law coefficient for NH ₃	57.6 ^c
$p_{0,i}$	Pure-liquid saturation vapour pressure comp. i	SIMPOL
$\gamma_{\rm i}$	Activity coefficient for compound <i>i</i>	AIOMFAC
$D_{0,monomer,SOA} (\mathrm{cm}^2 \mathrm{s}^{-1})$	Diffusion coefficient for SOA monomers	5x10 ^{-17d}
$D_{0,ammonium} (\mathrm{cm}^2 \mathrm{s}^{-1})$	Diffusion coefficient for NH ₃ /NH ₄ ⁺	1.3×10^{-16d}
$D_{NH4RCOO} (\mathrm{cm}^2 \mathrm{s}^{-1})$	Diffusion coefficient of organic salts	0

Table 3. Base case model set-up values for the simulation of organic salt formation between
 carboxylic acids and dissolved ammonium ions.

^aBase case simulation value. ^bLide, 2008 (CRC Handbook of Chemistry and Physics) at 298 K.

⁴ ^cJacobson, 2005a. ^dBased on the Stokes-Einstein relationship and a SOA viscosity of 10⁸ Pa s

5 (Abramson et al., 2013).

Date	Initial	Initial	$\Delta[O_3]$	Δ[α-	Yield	Initial	$\Delta[O_3]$	Δ[α-	Yield
	[α-pin.]	NH_3	exp.	pin.]	exp.	[α-pin.]	model	pin.]	model.
	exp.	(nnh)	(ppb) exp.	ovn	(%)	model (ppb)	(ppb)	model	(%) nodel
	(ppb)	(ppu)		ехр.				moder	
								(ppb)	
01/25/05	221		130	218	54.3	222	149	216	57.5
01/11/05	221	50	150	203	60.3	222	149	216	63.6
01/10/05	223	100	150	206	64.0	222	149	216	65.4
01/06/05	224	200	151	220	65.3	222	149	216	67.0

1 Table 4. Initial conditions and results from the α -pinene – O_3 – NH_3 – CO experiments (Na et al., 2007) and base case model simulations.



2 Figure 1. Schematic picture of the ADCHAM model structure.



2 Figure 2. Schematic figure which illustrates how ADCHAM treat the diffusion limited mass

3 transfer of gas phase compounds across a laminar layer next to the chamber walls. The thin (1

4 mm thick) air layer next to the chamber walls is treated as a separate volume which exchange gas

5 phase compounds with the well mixed chamber and the walls + wall deposited particles.



Figure 3. Schematic picture which illustrates the model structure and processes included in the

- 3 kinetic multilayer model in ADCHAM. The double arrows represent the mass transport between
- 4 the layers.
- 5



Figure 4. Modelled and measured (Vaden et al., 2011) evaporation losses of DOP particles of different initial diameters. The model results are from simulations with (1) Eq. 30 and a laminar layer width of 0.1 cm adjacent to the charcoal denuder (solid lines), and (2) Eq. 31 and a laminar layer of 0.6 cm (dashed lines). The measurements are given by the solid circles. The DOP mass accommodation coefficient was assumed one.



2 Figure 5. Modelled and measured (Vaden et al., 2011) evaporation rates of fresh α -pinene SOA. 3 (a) Model results for SOA particles with volatility according to the VBS parameterization from Pathak et al. (2007). (b) Model results are from simulations using vapour pressures of the 4 MCMv3.2 compounds were estimated with SIMPOL. (c) Model results using vapour pressures 5 6 estimated with the Nannoolal method. The evaporation loss rates are given for particles with a diameter of ~160 and ~250 nm, treated as liquid (1) or solid (s) (no diffusion between the particle 7 layers). The reversible gas-wall partitioning of all α -pinene oxidation products onto the smog 8 chamber Teflon walls was modelled with $k_{g,w}^* = 1/2000 \text{ s}^{-1}$ and $C_w / (M_w \gamma_{w,i}) = 100 \text{ }\mu\text{mol m}^{-3}$. 9



1

Figure 6. Measured (Vaden et al., 2011) and modelled evaporation losses for particles composed of approximately equal amount of dimer and monomer SOA, prior to the introduction in the charcoal denuder chamber. The saturation vapour pressures were calculated with SIMPOL, $D_{0,\text{monomer}} = 2x10^{-14} \text{ cm}^2 \text{ s}^{-1}$, $D_{\text{dimer}} = 2x10^{-16} \text{ cm}^2 \text{ s}^{-1}$, $k_f = 10^{-23} \text{ molecules}^{-1} \text{ cm}^3 \text{ s}^{-1}$, $k_d = 1/10 \text{ h}^{-1}$, no phase separation and a $k_{g,w}^* = 1/2000 \text{ s}^{-1}$. The results are given both for fresh and aged particles with a diameter of ~160 nm and ~250 nm, respectively.



1

Figure 7. Measured (Vaden et al., 2011) and modelled evaporation loss rates for semi-solid tar like particles ($D_{0,monomer}=5x10^{-17}$ cm² s⁻¹ and $D_{oligomer}=0$ cm² s⁻¹). The fresh SOA particles are composed of short lived dimers (~20 mass %) and long lived dimers (2.19 and 1.44 mass % for 160 nm and 250 nm particles, respectively) (see Fig. S9), before they are introduced in the charcoal evaporation chamber. The saturation vapour pressures were calculated with SIMPOL and $k_{g,w}^* = 1/1000$ s⁻¹.



Figure 8. Modelled NH₃(g), O₃(g), α-pinene(g) and OH(g) concentrations for the α-pinene
oxidation experiments by Na et al. (2007).





2 Figure 9. Comparison of modelled and measured SOA mass and mass yields at different initial levels of NH₃(g). In Fig. a) the model results are from simulations with vapour pressures from 3 4 SIMPOL, activity coefficients from AIOMFAC and very slow mixing between the particle layers (base case), in Fig. b) the results are from simulations with vapour pressures from Nannoolal et 5 al. (2008), activity coefficients from AIOMFAC and very slow mixing between particle layers, in 6 Fig. c) the results are from simulations with vapour pressures from SIMPOL, unity activity 7 8 coefficients (ideal solution) and very slow mixing between the particle layers, and in Fig. d) the model results are from simulations with vapour pressures from SIMPOL, activity coefficients 9 from AIOMFAC and semi-solid less viscous particles with $D_{0,monomer,SOA}=10^{-15}$ cm² s⁻¹, 10 $D_{0,ammonium} = 10^{-13} \text{ cm}^2 \text{ s}^{-1}$ and $D_{0,\text{NH4RCOO}} = 0 \text{ cm}^2 \text{ s}^{-1}$. 11



1

Figure 10. Modelled SOA mass formation for α -pinene – O₃ – NH₃ – CO experiments with 200 ppb NH₃ added at the start or after 6 hours of aging. For all simulations SIMPOL was used to estimate the vapour pressures of the organic compounds. The SOA particles were either treated as completely solid (no mixing between particle layers) (simulation 1-4) or semi-solid with $D_{0,monomer,SOA}=10^{-15}$ cm² s⁻¹, $D_{0,ammonium}=10^{-13}$ cm² s⁻¹ and $D_{0,NH4RCOO}=0$ cm² s⁻¹ (simulation 5). For simulation 2 and 3 we assume that the NH₄RCOO salts form a separate phase which other organic compounds cannot dissolved into.



Figure 11. Modelled a) pH, b) Ammonium (NH₄⁺ + NH₃(1) + NH₄ in NH₄RCOO) mass fractions, c) NH₄RCOO mass fractions and d) Carboxylic acid (COOH + COO⁻) mass fractions at different distances from the particle core, for α -pinene SOA particles with a diameter of approximately 240 nm after 1 hour and 280 nm after 6 hours of aging. The model results are from three different simulations with an initial [NH₃(g)] of 50, 100 or 200 ppb. The SOA particles were assumed to be semi-solid with $D_{0,monomer,SOA}=10^{-15}$ cm² s⁻¹, $D_{0,ammonium}=10^{-13}$ cm² s⁻¹ and $D_{0,NH4RCOO}=0$ cm² s⁻¹ 1.



Figure 12. Modelled dry deposition and coagulation losses of $(NH_4)_2SO_4$ seed aerosol particles in the Lund University 6 m³ Teflon chamber. We used a friction velocity of 0.05 m/s and a mean electrical field strength of 50 V cm⁻¹. The chamber volume loss rate was set to 0.8 m³ h⁻¹. The model results gives both the particle concentrations in the air (with or without coagulation) and on the particle walls. Figure a) shows the modelled and measured particle number size distributions, b) effective wall loss rates (modelled), c) number concentration and d) volume concentration.



Figure 13. Modelled and measured gas phase concentrations from the *m*-xylene oxidation experiment from Nordin et al. (2013). Figure a-d gives the modelled and measured NO, NO₂, O₃ and *m*-xylene concentration, respectively. Figure e-f shows the modelled OH and HONO concentrations. The model results are from simulations with: 1) the original MCMv3.2 gas phase chemistry, 2) with MCMv3.2 gas phase chemistry and an artificial OH source of 10⁸ cm³ s⁻¹, and 3) with MCMv3.2 gas phase chemistry, the artificial OH source and wall emissions of HONO.



Figure 14. Modelled and measured volume concentrations of (seed aerosol + SOA coating) 2 during the *m*-xylene oxidation experiment by Nordin et al. (2013). The model results are given 3 4 both for the particles in the air and for those that have deposited on the chamber walls. The results in (a) are from simulations with a laminar layer width (Δx) of 0.1 cm adjacent to the 5 6 chamber walls and a first order loss rate from the near wall gas phase to the walls $(k_{g,w})$ of 1/20 s⁻ ¹. The results in (b) are from simulations with a Δx of 1.0 cm and a $k_{g,w}$ of 1/6 s⁻¹. The figures 7 show both the results from simulations without or with heterogeneous reactions between O3 and 8 unsaturated organic compounds ($k_{0_3} = 10^{-16}$ molecules⁻¹ cm³ s⁻¹ and $D_{0,0_3} = 10^{-8}$ cm² s⁻¹). 9





Figure 15. Modelled and measured volume concentrations (seed aerosol + SOA coating) during the *m*-xylene oxidation experiment from Nordin et al. (2013). The model results are from simulations with relative rapid ($k_f = 10^{-22}$ molecules⁻¹ cm³ s⁻¹) peroxyhemiacetal and hemiacetal formation, $\Delta x = 0$ or 0.1 cm and $k_{g,w} = 0$ or 1/15 s⁻¹. The model results are given both for the particles in the air and the particles deposited on the chamber walls.





