

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Methane and carbon dioxide fluxes and their regional scalability for the European Arctic wetlands during the MAMM project in summer 2012

S. J. O'Shea¹, G. Allen¹, M. W. Gallagher¹, K. Bower¹, S. M. Illingworth¹, J. B. A. Muller¹, B. Jones¹, C. J. Percival¹, S. J-B. Bauguitte², M. Cain³, N. Warwick³, A. Quiquet³, U. Skiba⁴, J. Drewer⁴, K. Dinsmore⁴, E. G. Nisbet⁵, D. Lowry⁵, R. E. Fisher⁵, J. L. France⁵, M. Aurela⁶, A. Lohila⁶, G. Hayman⁷, C. George⁷, D. Clark⁷, A. J. Manning⁸, A. D. Friend⁹, and J. Pyle¹⁰

¹School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Oxford Road, Manchester, M13 9PL, UK

²Facility for Airborne Atmospheric Measurements (FAAM), Building 125, Cranfield University, Cranfield, Bedford, MK43 0AL, UK

³Centre for Atmospheric Science, University of Cambridge, Cambridge CB2 1EW, UK

⁴Centre for Ecology and Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

⁵Department of Earth Sciences, Royal Holloway, University of London, Egham TW20 0EX, UK

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁶Climate Change Research, Finnish Meteorological Institute, P.O. Box 503, 00101, Helsinki, Finland

⁷Centre for Ecology and Hydrology, Maclean Building, Crowmarsh Gifford, Wallingford, OX10 8BB, UK

⁸UK Meteorological Office, Fitzroy Rd, Exeter, UK

⁹Department of Geography, Downing Place, Cambridge CB2 3EN, UK

¹⁰National Centre for Atmospheric Science, UK

Received: 7 February 2014 – Accepted: 12 March 2014 – Published: 31 March 2014

Correspondence to: S. J. O'Shea (sebastian.oshea@manchester.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Airborne and ground-based measurements of methane (CH_4), carbon dioxide (CO_2) and boundary layer thermodynamics were recorded over the Fennoscandian wetlands in July 2012 as part of the MAMM (Methane and other greenhouse gases in the Arctic – Measurements, process studies and Modelling) field campaign. Employing these airborne measurements and a simple boundary layer box model, net regional scale (~ 100 km) fluxes were calculated to be $1.2 \pm 0.5 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$ and $-350 \pm 143 \text{ mg CO}_2 \text{ h}^{-1} \text{ m}^{-2}$. These airborne fluxes were found to be relatively consistent with seasonally-averaged surface chamber ($1.3 \pm 1.0 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$) and eddy covariance ($1.3 \pm 0.3 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$ and $-309 \pm 306 \text{ mg CO}_2 \text{ h}^{-1} \text{ m}^{-2}$) flux measurements in the local area. The internal consistency of the aircraft-derived fluxes across a wide swath of Fennoscandia coupled with an excellent statistical comparison with local seasonally-averaged ground-based measurements demonstrates the potential scalability of such localised measurements to regional scale representivity. Comparisons were also made to longer term regional CH_4 climatologies from the JULES (Joint UK Land Environment Simulator) and Hybrid8 land surface models within the area of the MAMM campaign. The average hourly emission flux output for the summer period (July–August) across the years 1980 to 2010 was $0.054 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$ (minimum 0.0 and maximum $0.38 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$) for the JULES model and $0.073 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$ (minimum -0.0018 and maximum $4.62 \text{ mg CH}_4 \text{ h}^{-1} \text{ m}^{-2}$) for Hybrid8. Based on these observations both models were found to significantly underestimate the CH_4 emission flux in this region, which was linked to the under prediction of the wetland extents generated by the models.

1 Introduction

Temperatures at high northern latitudes have been observed to be increasing at a rate of twice the global average over the past two decades (Forster and Ramaswamy, 2007).

Methane and carbon dioxide fluxes

S. J. O’Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



It has been suggested that this rise will continue (Parmentier et al., 2013). This is likely to have significant consequences for natural greenhouse gas emissions in the region, which contains potentially large sources that are known to be highly sensitive to changes in temperature; such as the boreal wetlands and the reservoirs of carbon that are sequestered in permafrost and as methane-hydrates (Smith et al., 2004; Zimov et al., 2006a, b; Ping et al., 2008). As well as contributing to radiative forcing, such emissions have the potential to significantly perturb atmospheric chemistry, including oxidant capacity (Isaksen et al., 2011). Palaeo-records indicate that strong positive feedbacks exist between climate and greenhouse gas emissions in the region (Walter et al., 2007; Nisbet and Chappellaz, 2009) and recent studies have already reported newly identified or growing CH₄ emissions from some of these reservoirs (Westbrook et al., 2009; Shakhova et al., 2010; Kort et al., 2012; Anthony et al., 2012).

Wetland regions are the single largest source of atmospheric CH₄, accounting for approximately a third of total global CH₄ emissions equivalent to 142–208 Tgyr⁻¹ (Kirschke et al., 2013), for which Boreal and Arctic regions make a significant contribution (approximately 25%; Smith et al., 2004; Zhuang et al., 2006). Much of the remainder is currently suggested, (by potentially poorly constrained inversion models), to originate from tropical wetlands (Bridgham et al., 2013). Biogenic CH₄ is produced in anoxic soils through the decomposition of organic matter by methanogenic bacteria (Bridgham et al., 2013). Emission rates by this process are dependent on soil moisture, temperature and the availability of organic matter (Pelletier et al., 2007; Strom and Christensen, 2007). Much of this CH₄ does not reach the atmosphere due to consumption that occurs in oxic soil regions by methanotrophic bacteria (O'Connor et al., 2010). As a result of these competing environment-dependant factors, emissions show a large degree of spatial and temporal variability (Zhuang et al., 2006; Pickett-Heaps et al., 2010).

CO₂ exchange between the surface and the atmosphere in these regions displays a similar degree of complexity. It is governed by the interplay between release of CO₂ through respiration and uptake by photosynthesis. At high latitudes, as temperatures

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

rise and the ground thaw reaches greater soil depths, more organic carbon becomes available for decomposition, potentially liberating large carbon reservoirs to the atmosphere (Oelke et al., 2004). However, a simultaneous increase in plant production and biomass may also occur during the growing season. Rapid warming at high latitudes is increasing both plant growth and soil decomposition, making it difficult to determine the overall impact a warmer climate has on the total net carbon budget of Arctic and boreal regions (Zhuang et al., 2006; Davidson and Janssens, 2006; Schuur et al., 2009; Sitch et al., 2007).

Previously, Arctic wetland emissions have been determined by up-scaling surface chamber and eddy covariance flux measurements (Pelletier et al., 2007) or by process-based and inverse models (Petrescu et al., 2010; Pickett-Heaps et al., 2010; Wania et al., 2010; Bousquet et al., 2011). However, due to the heterogeneous nature of wetlands, uncertainties exist when multiple studies are synthesized to determine net emissions for large areas (Christensen et al., 2007). Currently, there is a lack of flux measurements at the same spatial scale as the resolution of land surface models (typically 0.5°), which has been identified as a key reason why models are not able to confidently simulate the global wetland CH_4 flux (Melton et al., 2013). Airborne measurements have been shown to be a powerful tool in reducing these uncertainties (Desjardins et al., 1997; Miller et al., 2007; Peischl et al., 2012), where the greater spatial coverage afforded may be an advantage over ground-based measurements under appropriate conditions, especially when testing the scalability of fluxes derived for local scales across wider areas.

Ground-based CH_4 flux measurements have now been made for multiple years in several wetland locations within northern Fennoscandia, these include: the Stordalen wetlands, in sub-arctic Sweden (68.33°N , 19.05°E , Christensen et al., 2012) and both Kaamanen (69.1°N , 27.2°E , Maanavilja et al., 2011) and Lompolojännkä (68.0°N , 24.2°E , Aurela et al., 2009) in Finland. Stordalen summer CH_4 emissions have been reported as $4.7 \text{ mgCH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (2004 to 2006) and $6.2 \pm 2.6 \text{ mgCH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (2006 and 2007) (Petrescu et al., 2008; Jackowicz-Korczynski et al., 2010). Mean July CO_2

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fluxes are $-1152 \text{ mgCO}_2 \text{ m}^{-2} \text{ h}^{-1}$, $-576 \text{ mgCO}_2 \text{ m}^{-2} \text{ h}^{-1}$ and $-504 \text{ mgCO}_2 \text{ m}^{-2} \text{ h}^{-1}$ for Lompolojännkä, Kaamanen and Siikaneva, respectively (Aurela et al., 2009).

This paper uses in-situ measurements collected on board the UK's Facility for Airborne Atmospheric Measurements (FAAM) BAe-146 research aircraft to quantify greenhouse gas net fluxes from the Fennoscandian wetlands during a dedicated case study. A simple boundary layer mass budget approach (described in Sect. 3.1) is employed to derive regional fluxes using the aircraft observations under pseudo-stationary boundary layer flow assumptions. This estimate is then compared to smaller footprint ground-based eddy covariance and chamber measurements within the aircraft's sampling footprint that were made over much of Summer 2012 to address scalability and spatiotemporal heterogeneity (Sect. 3.3). Finally, the regional scale aircraft derived flux is used to assess the skill of two land surface models (Sect. 3.4).

2 Methods

The measurements reported in this paper were collected as part of the MAMM (Methane and other greenhouse gases in the Arctic – measurements, process studies and modelling, <http://arp.arctic.ac.uk/projects/methane-and-other-greenhouse-gases-arctic-measrem/>) project. The aim of the MAMM project is to quantify greenhouse gas fluxes at high northern latitudes using a combination of measurement, process and modelling studies. As part of this project, sorties were performed from Kiruna, Sweden by the FAAM BAe-146 research aircraft during July 2012 (6 flights), August 2013 (9 flights) and September 2013 (7 flights). This paper focuses on one flight conducted on the 22 July 2012 (flight number B720) and simultaneous ground based flux measurements within the operational area. This flight has been chosen for this case study due to the favourable meteorological and flight conditions for applying a mass budget approach to derive fluxes (Sect. 3.1). The MAMM campaign is ongoing at the time of writing and we anticipate that a seasonal analysis will be addressed in the future.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 FAAM BAe-146 research aircraft

CO₂ and CH₄ dry air mole fractions were determined through cavity-enhanced absorption spectroscopy on board the FAAM BAe-146 (CEAS, Model RMT-200, Los Gatos Research Inc, USA). In-flight CO₂ uncertainty was calculated as ± 0.17 ppm; typical 1 Hz precision is ± 0.70 ppm (all precisions are 1σ). CH₄ uncertainty is calculated at ± 1.31 ppb; 1 Hz precision is ± 2.37 ppb (for a detailed description of this system see O'Shea et al., 2013b). Separate measurements of CO₂ and CH₄ were made by analysing whole-air-samples (WAS). These were collected in stainless steel flasks (for a description see Lewis et al., 2013), and analysed post-flight in the laboratory using cavity-ring down spectroscopy (Model G1301, Picarro Inc, USA). Uncertainty is estimated at ± 0.5 ppb and ± 0.1 ppm for CH₄ and CO₂, respectively. During the MAMM flights the mean bias of the whole air samples (400 samples) relative to the in situ measurements was -0.16 (± 0.46 at 1σ) ppm for CO₂ and -0.5 (± 4.6 at 1σ) ppb for CH₄. Flask samples were also analysed for $\delta^{13}\text{C}$ isotopic ratios of CO₂ and CH₄, using continuous-flow gas chromatography/isotope-ratio mass spectrometry (CF-GC-IRMS), with a precision of 0.1 ‰ (Fisher et al., 2006).

A range of other chemical, tracer and thermodynamic parameters were measured simultaneously on board the FAAM BAe-146; these include pressure, temperature and the 3-D wind vector with an estimated uncertainty of 0.3 hPa, 0.1 K and 0.2 ms^{-1} , respectively (Allen et al., 2011). Measurements of carbon monoxide (CO) and hydrogen cyanide (HCN) are used here to identify air masses that have been strongly influenced by either biomass burning or anthropogenic activity using an enhancement-over-background-threshold technique described by O'Shea et al. (2013a), as such air masses would bias the calculation of the biogenic flux. Mole fractions of CO were determined through VUV (vacuum ultraviolet) fast-fluorescence spectrometry, with an uncertainty of 2 % (AL5002, Aerolaser GmbH, Germany; Gerbig et al., 1999). In situ HCN measurements were made using a chemical ionisation mass spectrometer, with an uncertainty of 10 % (Le Breton et al., 2013).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Surface measurements

CH₄ and CO₂ eddy covariance and chamber flux measurements were made in So-
dankylä, Finland from 1 July 2012 to 15 August 2012. The eddy covariance sys-
tem used included a USA-1 (METEK GmbH, Germany) three-axis sonic anemome-
ter/thermometer, a RMT-200 (Los Gatos Research, Inc., USA) CH₄ analyzer and a LI-
7200 (Li-Cor, Inc., USA) CO₂/H₂O gas analyzer. The measurement height was 6 m
(a.g.l.). The length of the inlet tubes for both gases was 8 m for CH₄ and 1 m for CO₂,
with flow rates of 15 and 20 L min⁻¹, respectively. For more details of the eddy covari-
ance measurement system, see Aurela et al. (2009).

Half-hour flux values were calculated using standard eddy covariance methods. The
original 10 Hz data was block-averaged, and a double rotation of the coordinate sys-
tem was performed (McMillen, 1988). The time lag between the anemometer and gas
analyzer signals, resulting from the transport through the inlet tube, was taken into ac-
count in the on-line calculations. An air density correction related to the sensible and
latent heat fluxes was conducted according to Webb et al. (1980). Corrections for the
systematic high-frequency flux loss owing to the imperfect properties and setup of the
sensors (i.e. insufficient response time, sensor separation, damping of the signal in
the tubing and averaging over the measurement paths) were carried out off-line using
transfer functions with empirically-determined time constants (Aubinet et al., 2000). All
data with wind directions from sector 240–290° were discarded due to insufficient fetch.
Some data were also discarded due to instrument failures during weak turbulence (fric-
tion velocity < 0.1 ms⁻¹). CO₂ fluxes during the period 14 July 2012 to 1 August 2012
are missing due to instrumental problems.

Fluxes of CH₄ were also measured using the static chamber method, as follows.
These were positioned to cover a range of vegetation types and water saturations that
can be broadly classified into either those situated in wetlands (39 chambers) and those
in the forest (21 chambers). Shallow frames were installed the day before first sampling
to a depth of ~ 10 cm, and remained in situ for the duration of the study period; fluxes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calculated from the first sampling were not significantly different from subsequent sampling occasions suggesting that the short settling period after frame installation had no effect. Fluxes were measured at ~ 2 day intervals between 12 July and 2 August. For measurements, chamber lids were attached to the frames and internal air samples were collected into vials 4 times over a 45 min incubation period. Samples were analyzed by gas chromatography and fluxes calculated using GCFlux, version 2 (Levy et al., 2012). Reported CH_4 fluxes correlate to the best-fit model for individual chambers (either linear or asymptotic). Fluxes of N_2O and CO_2 were also measured by the static chamber method. However, since static-chamber-measured CO_2 fluxes are only a measure of soil respiration and do not include uptake by plants, they cannot be directly compared with the aircraft-derived flux estimates; this will be presented in a separate study.

2.3 Flight sampling and study area

On the 22 July 2012 the FAAM BAe-146 surveyed the Northern Fennoscandian landscape. Four large transects (~ 340 km) were performed within the planetary boundary layer (PBL) at an approximate altitude range of 100 m to 2000 m: two east–west (east to west transect 10:42 to 11:46 GMT; west to east 15:26 to 16:04 GMT) and two north–south. Figure 1a shows the geographic coverage of this flight along with the location of waypoints: Kiruna (67.9° N, 20.2° E), Sodankylä (67.4° N, 26.6° E) and Kaamanen (69.1° N, 27.2° E). To show the prevalent vegetation and land use types within the region, the flight track is also shown overlaying the land classification (Fig. 1b, Corine land cover 2006, <http://www.eea.europa.eu/data-and-maps/data/corine-land-cover-2006-raster>). As seen, the sampling domain is largely characterised by coniferous forests (33%, dark green Fig. 1b), peat bogs (23%, blue Fig. 1b) and mixed forests (16%, green Fig. 1b).

During the FAAM flight, in situ measurements showed winds to be consistently easterly ($6 \pm 2 \text{ ms}^{-1}$). HYSPLIT (Hybrid Single Particle Lagrangian Integrated Trajectory Model, Draxler and Rolph, 2003) back trajectories, calculated along the FAAM BAe-

Methane and carbon dioxide fluxes

S. J. O’Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



146's flight track, also show a predominant easterly flow in the region. The majority of air masses sampled by the FAAM BAe-146 spent the previous 5 days within the Arctic region.

2.4 Methane emission models

In Sect. 3.4, we assess the skill of two land surface models: the Joint UK Land Earth Simulator (JULES: Best et al., 2011; Clark et al., 2011) and Hybrid8 (Friend et al., 2010). The JULES model contains a CH₄ wetland emission parameterization, developed and tested by Gedney et al. (2004) for use at large spatial scales. The wetland parameterization is coupled to the large-scale hydrology scheme of Gedney and Cox (2003), which predicts the distribution of sub-grid scale water table depth and wetland fraction (f_w) from the overall soil moisture content and the sub-grid scale topography using the approach of Beven and Kirby (1979). The CH₄ flux from wetlands, $F_w(\text{CH}_4)$, is parameterized as a function of temperature, water table height and substrate availability, as follows:

$$F_w(\text{CH}_4) = f_w k(\text{CH}_4) C_s Q_{10}(T_{\text{soil}})^{(T_{\text{soil}} - T_0)/10} \quad (1)$$

where T_{soil} is the soil temperature (in K) averaged over the top 10 cm and $k(\text{CH}_4)$ is a global constant which is calibrated to give the required global CH₄ flux. The Q_{10} is a temperature coefficient to account for the temperature dependency of the flux. Soil carbon content (C_s) was used for substrate availability. The default parameter values chosen were $k(\text{CH}_4) = 7.4 \times 10^{-12} \text{ kg m}^{-2} \text{ s}^{-1}$, $T_0 = 273.15 \text{ K}$ and $Q_{10}(T_0) = 3.7$ (as per Clark et al., 2011).

The surface physics of the Hybrid8 model are based on the NASA-GISS ModelE land surface component (Schmidt et al., 2006). This model contains a sophisticated canopy representation (Friend and Kiang, 2005) allowing for a realistic canopy conductance to various environmental factors (light, temperature, humidity, CO₂ and canopy height). Recently, we have implemented a TOPMODEL approach to model the hydrology following Niu et al. (2005). Very similar to the implementation in the JULES land

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

surface model, the TOPMODEL hydrological module in Hybrid8 uses a topographical index and interactively computes the wetland fraction in each grid box (f_w). In our approach only saturated soils (determined by f_w) contribute to CH_4 emissions. The fluxes of CH_4 are parameterised in a very similar way in Hybrid8 as in JULES. The governing equation for CH_4 production at depth z , is,

$$P_w(\text{CH}_4) = k(\text{CH}_4)F_{ps}(z)C_{\text{som}}(z)Q_{10}^{T(z)-T_0} \quad (2)$$

Where $k(\text{CH}_4)$ is the baseline production rate; $F_{ps}(z)$ is the total pore space fraction in a specific layer (a function of soil texture); $C_{\text{som}}(z)$ is the soil organic matter at the depth z ; and $T(z)$ is the soil temperature. For this study, the following representative parameters were chosen: $k(\text{CH}_4) = 1.3 \times 10^{-11} \text{ kg m}^{-2} \text{ s}^{-1}$, $Q_{10} = 3$ and $T_0 = 22^\circ\text{C}$. The CH_4 produced is then transported to adjacent layers via diffusivity, eventually reaching the atmosphere.

3 Results and discussion

On the 22 July 2012 consistent linear gradients were observed in both CH_4 and CO_2 along the longitudinal transects (Fig. 1). CH_4 was found to be approximately 20 ppb higher at the eastern boundary compared to the western, while CO_2 decreased by several ppm over the same interval. No clear latitudinal trends were observed in either species. However, a region of significantly enhanced CH_4 (up to 20 ppb) was observed to the north of Sodankylä (Fig. 1), a region with a slightly higher proportion of wetlands (29%).

With a mean PBL mole fraction of 89 ppb for CO and 26 ppt for HCN, both species remained at mole fractions throughout the flight that are representative of a typical background for the summer at these latitudes (Vay et al., 2011; O'Shea et al., 2013a). This indicates that any biomass burning and anthropogenic emissions within the region were small and well-mixed when sampled by the FAAM BAe-146. To identify the source of the observed enhancements we use the measured $\delta^{13}\text{C}$ isotopic ratios and a Keeling

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

plot methodology. Figure 2 shows a Keeling plot for all PBL measurements of $\delta^{13}\text{C}-\text{CH}_4$ during the flight on the 22 July 2012 (B720). The vertical intercept represents the isotopic ratio of the source of the enhancements. A source of $-72.0 \pm 4.4\%$ as seen here is consistent with wetland CH_4 emissions (-71 to -59% ; Fisher et al., 2011; Sriskantharajah et al., 2012).

3.1 Regional scale fluxes derived using aircraft observations

Mass budget approaches have been employed on several occasions to derive regional scale (> 1 km) fluxes of trace species (White et al., 1976; Gallagher et al., 1994; Choularton et al., 1995; Wratt et al., 2001; Mays et al., 2009). Observations are typically made in a background location and then down-wind of a source region to determine the net enhancement due to this region. In this study, we use the fact that the East–West transect performed during the 22 July 2012 flight was aligned nearly parallel with the prevailing wind bearing, which was 258° . If it can then be assumed that the non-reactive tracer species, S , is well mixed from the surface up to the top of the PBL, Z_1 , and that entrainment into (and detrainment from) the PBL can be neglected, then the net flux of S can be determined by

$$\text{flux} = \frac{\bar{U}}{\cos\theta} \frac{\Delta S}{\Delta x} \int_0^{Z_1} n dz, \quad (3)$$

where \bar{U} (m s^{-1}) is the mean wind speed, and n (molecules m^{-3}) is the atmospheric number density, which is integrated from the surface to the top of the boundary layer (m). The ΔS ($\text{molecules molecules}^{-1}$) term is the enhancement in species S along the transect x of increment Δx (m) parallel to the prevailing wind. the angle θ is between the mean wind vector and transect x . This gradient ($\Delta S/\Delta x$) is determined here by first averaging the data to 500 m intervals (equivalent to around 4 s of sampling time) along x , before performing an orthogonal distance regression (Fig. 3). The regression slope

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is weighted by the quadrature addition of the analytical uncertainty and the vertical variability of S throughout the PBL (Fig. 4).

Several assumptions are implicit in this model; firstly, that emissions are immediately mixed throughout the PBL column. This is expected to be reasonable given that the time taken for air to advect across the transect (up to ~ 16 h) is much larger than the average turnover time of the PBL (~ 80 min, calculated using thermodynamic parameters measured here and following Stull et al., 1988, and Karion et al., 2013). Secondly, the transect, x , should ideally be aligned parallel to the wind vector. However, we note that there was a 12° offset between the mean wind vector and transect x (θ , Eq. 3), while the wind also showed some variation about the mean ($1\sigma = 24^\circ$). It then has to be assumed that mole fractions perpendicular to the wind vector are constant. The 1σ of the wind direction is used in the flux uncertainty propagation.

Vertical profiles of potential temperature, from the FAAM BAe-146, performed over Sodankylä (Fig. 1) and from the two dropsondes released, show a clear capping inversion was present over the area during the flight (Fig. 4). Following Mays et al. (2009), a PBL height of 1740 m (a.g.l.) is determined here from both ascending and descending vertical profiles, which show strong mixing (constant potential temperature profile) between the ground and 1740 m. Uncertainty (based on the observed changing vertical gradient in potential temperature between the PBL beneath and the free troposphere above) is estimated at ± 200 m around this mixing height (see Fig. 4) between the start and end of the sampling period. In addition, above the boundary layer, both CO_2 and CH_4 show abrupt changes in their mole fraction and the vertical wind speed becomes less variable (variance in the wind speed above the boundary layer is typically less than $0.2 \text{ m}^2 \text{ s}^{-2}$), supporting the assumption that entrainment into and out of the boundary layer is relatively small and so can be neglected for this exercise. However, within the boundary layer some structure exists in the altitude profile. The CH_4 standard deviation was 4.5 ppb for the ascending profile and 1.7 ppb for the descending profile, while for CO_2 this was 1 ppm for both the ascending and descending profiles. Some of this variability is likely to be due to the fact that these profiles are recorded slant-wise in

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the horizontal and therefore reflect both variability in vertical mixing and the existing horizontal gradient. This variability is included in the error propagation described below to derive a representative and comprehensive uncertainty on the calculated flux.

The calculated fluxes are found to be $1.2 \pm 0.5 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ and $-350 \pm 143 \text{ mgCO}_2 \text{ h}^{-1} \text{ m}^{-2}$ (Table 2 and Fig. 5). The uncertainty in the total flux is determined by propagating the uncertainties associated with the individual terms in Eq. (3); these include the uncertainty in the observed (fitted) spatial mole fraction gradient, known variability in the wind, and boundary layer mixing height, as identified above. Similar to previous studies (e.g. Ryerson et al., 1998), the largest known source of uncertainty was found to be the assumption of a single wind vector for the whole of the transect x . Within the uncertainties, the fluxes are in agreement whether separately-derived eastward, westward, or combined transects are used in the calculation. The repeatability of this measured gradient indicates that both species were vertically well-mixed since the transects were performed at slightly different altitudes (eastward mean = 507 m, range = 70 to 1287 m; westward mean = 717 m, range = 103 to 1382 m). The fluxes calculated using the 11 WAS measurements, collected along the East–West transect, are also in excellent agreement (see Table 2) with that from the continuous in situ measurements. However, in the case of CO_2 this is with a large uncertainty.

3.2 Dispersion modelling

The flux derived from the aircraft measurements has also been tested using forward model runs with the UK Met Office's Numerical Atmospheric-dispersion Modelling environment (NAME) to diagnose whether the calculated ground flux might be expected to translate into the observed enhancements seen in measurements observed aloft when advected. NAME is a 3-D Lagrangian particle dispersion model (Ryall and Maryon, 1998; Ryall et al., 1998), which is run here using the UK Met Office's Unified Model meteorological fields (Cullen, 1993). A flux of $1.2 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ was emitted from the ground in the region bounded by 20° E to 28° E and 67° N to 69.5° N continuously for the period from 00:00 GMT on 20 July 2012 until 17:00 GMT on 22 July 2012, and the

model was run forwards to disperse the CH₄ through the modelled atmosphere. The particle motions are calculated based on the large-scale winds, wind meander and sub-grid scale stochastic turbulence.

Figure 6 shows a cross section of the atmosphere that is co-incident with flight B720.

The contours show the 1 h-average mixing ratio of CH₄ averaged over 67.75 to 68.00° N (upper panel for 11:00 GMT and lower panel for 16:00 GMT). This shows the modelled increment of CH₄ that comes from the local region, based on the flux calculated by the aircraft observations. At 11:00 GMT (the time of the eastward transect), the increment in CH₄ at the eastern end of the flight is approximately 15 to 20 ppb higher than the western part of the transect. By 16:00 GMT, the difference in the model has reduced to 12 to 15 ppb. This is because the model PBL is well mixed, and so gradients within it decline as the day progresses and the PBL top rises. It can be seen in Fig. 6a that the model PBL height is about 2200 m at 11:00 GMT (corresponding to our eastward transect) and has increased to about 3000 m by 16:00 GMT (the time of the westward transect). The higher late afternoon modelled PBL would act to dilute the CH₄, which can be seen in the lower modelled mixing ratio enhancements at 16:00 GMT (Fig. 6b). However, this dilution was not observed in the late afternoon aircraft measurements, which also showed a much lower PBL height of 1740 m (Fig. 3), similar to that observed earlier in the day.

Despite this, the increment to CH₄ is comparable for the 11:00 case (approximately 20 ppb in the observations, and approximately 15–20 ppb in the dispersion model). The reason for the difference in PBL height between the model and measurement cannot currently be explained and is beyond the scope of this study: however, these results confirm that observed enhancements can be reasonably represented by dispersion modelling when treating the land as a constant source equal to that derived here, for a PBL mixing height of ~ 2200 m (as modelled for the 11:00 GMT transect).

Methane and carbon dioxide fluxes

S. J. O’Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Ground-based flux measurements

In this section, we compare the aircraft-derived flux with seasonally averaged surface measurements to examine scalability and potential sources of bias (e.g. spatial heterogeneity). The ground based-measurements during the MAMM campaign comprised both chamber and eddy covariance flux measurements, as described in Sect. 2.2. A comparison between these two techniques and the aircraft-determined flux is complicated by the differences in their respective footprints. Chambers are the smallest scale (< 1 m) and are specific to a single land type. While eddy covariance fluxes are typically representative of 100 to 1000 m and as a result may average the flux across several land types. The aircraft represents a regional flux, in this case > 300 km, which encompasses several ecosystems with air mixed over all.

During the MAMM field campaign, 60 chambers were used to determine CH₄ fluxes. Fluxes for the entire measurement period, as well as those for just 22 July 2012 are given in Table 2 and Fig. 5. Forested regions are found to have negligible net flux, varying between a small source or sink (Ridgwell et al., 1999), while the wetlands show a wide range of net emissions, which could be expected since the chambers covered a wide range of soil moisture saturations.

The aircraft-derived CH₄ flux is within the wide range spanned by the forest and wetland chamber measurements (−0.09 to 11.6 mgCH₄ h^{−1} m^{−2}). This might be expected as both ecosystems are present within the aircraft's footprint (Fig. 1). For a more direct comparison we perform a weighted average of the two classes of chamber fluxes. This was done by first determining the aircraft's surface footprint using the NAME model. The Corine land cover map was then used to identify the prevalence of the each land classification within this footprint (Table 1). Each Corine classification was grouped as either a forest (coniferous forest, mixed forest, transitional woodland, broad leafed forest) or a wetland (peat bog, moor and heathland) land type. Using this methodology, during the 22 July flight's East–West transect, 28 % of the land footprint was classified as wetland and 65 % was classified as forest. These proportions were then

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



used to weight the averaging of the two chamber flux categories. The result of this is $1.3 \pm 1.0 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ (mean \pm standard deviation) using the summer mean chambers and $1.5 \pm 1.6 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ if just the 22 July 2012 measurements are used. Though poorly constrained, these are both in good agreement with the aircraft derived flux, which is only $0.1 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ and $0.3 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ lower, respectively (Fig. 5). Uncertainties exist in this comparison since the partitioning is quite broad and in the assumption of a zero flux for 7% of the land area. A more sophisticated comparison would assign measured fluxes for each 2006 Corine land cover. Nevertheless this simple approach provides a useful validation of the airborne calculation.

The CH_4 and CO_2 eddy covariance flux measurements were calculated for the Sodankylä wetland from 1 July 2012 to 15 August 2012 (Table 2 and Figs. 7 and 8). CH_4 chamber fluxes show a wider range than the eddy covariance fluxes, which could be expected since they covered the dryer and wetter parts of the wetland, while the eddy covariance method spatially integrates these regions and as a consequence is within this range. CH_4 fluxes do not show large variation over diurnal (Fig. 7) or weekly timescales (Fig. 8). However, CO_2 was emitted for several hours around midnight, while uptake occurred during the day. The mean daytime (06:00 to 18:00 GMT) eddy covariance CO_2 measurement of -309 ($1\sigma = 306$) $\text{mgCO}_2 \text{ h}^{-1} \text{ m}^{-2}$ is only $41 \text{ mgCO}_2 \text{ h}^{-1} \text{ m}^{-2}$ higher than the aircraft derived flux, well within the measurement uncertainty.

However, the mean daytime eddy covariance CH_4 flux of $4.5 \pm 1.2 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ for the summer period is a factor of 4 larger than the aircraft. This is comparable with some other previous studies in wetlands such as $4.7 \text{ mgCH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (2004 to 2006) and $6.2 \pm 2.6 \text{ mgCH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (2006 and 2007) for Stordalen (Petrescu et al., 2008; Jackowicz-Korczynski et al., 2010). Similar to the chamber measurements, this may be because the eddy covariance footprint is more specific to a single land type than the aircraft in this instance. To test this, the same scaling was repeated using the Corine land classification but this time using the Sodankylä wetland eddy covariance flux instead of that from the wetland chambers, which resulted in a flux of $1.3 \pm 0.3 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$. This then displays similarly good agreement with the aircraft derived flux.

3.4 Comparison against modelled wetland emission estimates

In this section, we compare our measurement-derived CH₄ emission fluxes with those predicted from wetlands in Fennoscandia by two land surface models: JULES and Hybrid8. The purpose of this comparison is to investigate how representative the regional snapshot we discuss above is, in the context of predicted seasonal and interannual variability, and to discuss potential sources of systematic bias. To this end, we have chosen a 31 year model period prior to the MAMM campaign for statistical comparison.

For this comparison, runs of the JULES and Hybrid8 models were to a 0.5° × 0.5° terrestrial grid covering Scandinavia, using the CRU-NCEP meteorological dataset (Viovy and Ciais, 2009). Hourly CH₄ emission fluxes from wetlands were derived between January 1980 and December 2010 (the last year currently available in the CRU-NCEP driving meteorological dataset). Table 3 summarises the statistics derived from the modelled hourly CH₄ emission for the domain covered by the aircraft (20.0–29.0° E, 67.5–68.5° N). As the JULES run does not cover the actual period of the MAMM campaign due to a lack of forcing data, we present climatological statistics for every July and August between 1980 and 2010.

Even accounting for the fact that the model dataset does not extend to 2012, it is evident that the two models significantly underestimate (a factor ~ 14 for Hybrid8 and ~ 22 for JULES in the mean) the CH₄ emission flux in this region for the 31 year climatology modelled here, when compared to our regionally representative case study. Furthermore, even the upper quartile maximum monthly-averaged flux in the 31 year climatology (0.085 mgCH₄ h⁻¹ m⁻² for JULES and 0.095 mgCH₄ h⁻¹ m⁻² for Hybrid8) does not approach the measured aircraft and ground-based results in this case study. This is possibly because of an under prediction of wetland extent by both models in this region, which could be linked to the topographical dataset used and/or the absence of an organic soil type related to peatlands. Such soils would have very different hydraulic properties to the mineral soil types currently used in JULES and Hybrid8. Water would be retained at or close to the surface increasing the area of wetlands. Model emission

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

fluxes were derived assuming that each grid cell is all wetland. These results were found to be much closer to the aircraft values for both JULES (1980–2010: median $1.5 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$, inter-quartile range 1.2 to $1.8 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$) and Hybrid8 (1980–2010: median $2.2 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$, inter-quartile range 1.7 to $3.0 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$). This suggests that underestimation of the area of wetlands in both models is probably the major reason for the under-prediction of the wetland emission fluxes in this region. Petrescu et al. (2010) investigated the sensitivity to the wetland area and found a wide variation in methane emission fluxes (37.7 to $157.3 \text{ TgCH}_4 \text{ yr}^{-1}$) from wetlands and floodplains above 30° N for the years 2001 to 2006 for different estimates of wetland extent. The wetland model intercomparison (Melton et al., 2013) has further highlighted the major challenges and uncertainties that exist in modelling wetlands and the associated CH_4 emissions.

Both the JULES and Hybrid8 models have been used to simulate the response of future emissions to climate change (Gedney et al., 2004; Friend et al., 2010). The results from this comparison suggest that there are significant uncertainties when emissions are simulated at regional scales and/or at specific times. Although our snapshot of a regionally representative flux on a single day should not be directly extrapolated to demonstrate a systematic under-bias in the climatological Arctic wetland CH_4 flux as predicted by JULES and Hybrid8, these results do point to the important need for further such case studies from which to build diagnostic statistics to validate such models. Given that this study suggests a minimum factor 3 under-bias in modelled fluxes, this uncertainty is potentially very important for climate studies that model CH_4 emissions scenarios.

4 Conclusions

As part of the MAMM field project, airborne measurements of CH_4 and CO_2 were collected in the European Arctic in summer 2012. An airborne mass balance approach was used to derive regional scale fluxes for the northern Scandinavian wet-

lands from one flight on the 22 July 2012. These were established to be $1.2 \pm 0.5 \text{ mgCH}_4 \text{ h}^{-1} \text{ m}^{-2}$ and $-350 \pm 143 \text{ mgCO}_2 \text{ h}^{-1} \text{ m}^{-2}$, which were comparable with simultaneous seasonally-averaged chamber and eddy covariance flux measurements made in Sodankylä (within 11% for CO_2 and 8% for CH_4 if the fluxes were scaled using the land type). The internal consistency of the aircraft-derived fluxes across a wide swath of Fennoscandia coupled with an excellent statistical comparison with local seasonally-averaged ground-based measurements demonstrates the potential scalability of such localised measurements to regional scale representivity.

Though the fluxes calculated here do not provide information about the wider temporal variability of fluxes, they do provide a snapshot that can be compared with the statistical climatology for model fluxes in the region; which is representative of a spatial scale that is comparable with the resolution of regional chemical transport and land surface models. This together with a well characterised uncertainty mean that these fluxes can provide a useful constraint for “bottom-up” regional flux calculations. To this end, a comparison with both the Hybrid8 and JULES land surface model suggests that they both significantly underestimate the net CH_4 flux from these regions (a factor ~ 16 for Hybrid8 and ~ 22 for JULES in the mean).

Although our snapshot of a regionally representative flux on a single day should not be directly extrapolated to demonstrate a systematic under-bias in the modelled climatological Arctic wetland methane flux, the results presented here do point to the important need for further such case studies from which to build diagnostic statistics to validate such models as this uncertainty is potentially very important for climate studies that model CH_4 emissions scenarios. Future field campaigns and studies are planned to exploit the MAMM airborne dataset from the 2013 flights to derive additional regional scale fluxes of key greenhouse gases either through mass balance approaches, as illustrated here, or inverse modelling. These may provide additional information on the consistency of the disagreement between observations and the JULES/Hybrid8 models at high northern latitudes.

Methane and carbon dioxide fluxes

S. J. O’Shea et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. The MAMM project was funded by the UK Natural Environment Research Council (Grant NE/I029293/1). S. J. O'Shea is in receipt of a NERC studentship and G. Allen is a NERC Fellow (#NE/I021276/1). Airborne data were obtained using the FAAM BAe-146 Atmospheric Research Aircraft (ARA) operated by Directflight Ltd (DFL) and managed by the Facility for Airborne Atmospheric Measurements (FAAM), which is a joint entity of the Natural Environment Research Council (NERC) and the UK Meteorological Office. The Corine land cover map was obtained from <http://www.eea.europa.eu/data-and-maps/data/corine-land-cover-2006-raster>, the copyright holder is the European Environment Agency (EEA). A. Friend received funding from the European Community's Seventh Framework Programme (FP7 2007–2013) under grant agreement #238366.

References

- Allen, G., Coe, H., Clarke, A., Bretherton, C., Wood, R., Abel, S. J., Barrett, P., Brown, P., George, R., Freitag, S., McNaughton, C., Howell, S., Shank, L., Kapustin, V., Brekhovskikh, V., Kleinman, L., Lee, Y.-N., Springston, S., Toniazzo, T., Krejci, R., Fochesatto, J., Shaw, G., Krecl, P., Brooks, B., McMeeking, G., Bower, K. N., Williams, P. I., Crosier, J., Crawford, I., Connolly, P., Allan, J. D., Covert, D., Bandy, A. R., Russell, L. M., Trembath, J., Bart, M., McQuaid, J. B., Wang, J., and Chand, D.: South East Pacific atmospheric composition and variability sampled along 20° S during VOCALS-REx, *Atmos. Chem. Phys.*, 11, 5237–5262, doi:10.5194/acp-11-5237-2011, 2011.
- Anthony, K. M. W., Anthony, P., Grosse, G., and Chanton, J.: Geologic methane seeps along boundaries of Arctic permafrost thaw and melting glaciers, *Nat. Geosci.*, 5, 419–426, doi:10.1038/ngeo1480, 2012.
- Aubinet, M., Grelle, A., Ibrom, A., Rannik, U., Moncrieff, J., Foken, T., Kowalski, A. S., Martin, P. H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grunwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., and Vesala, T.: Estimates of the annual net carbon and water exchange of forests: the EUROFLUX methodology, *Adv. Ecol. Res.*, 30, 113–175, 2000.
- Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Riutta, T., and Laurila, T.: Carbon dioxide exchange on a northern boreal fen, *Boreal Environ. Res.*, 14, 699–710, 2009.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 4, 677–699, doi:10.5194/gmd-4-677-2011, 2011.
- 5 Beven, K. J. and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology, *Hydrol. Sci. Bull.*, 24, 43–69, 1979.
- Bousquet, P., Ringeval, B., Pison, I., Dlugokencky, E. J., Brunke, E.-G., Carouge, C., Chevalier, F., Fortems-Cheiney, A., Frankenberg, C., Hauglustaine, D. A., Krummel, P. B., Langenfelds, R. L., Ramonet, M., Schmidt, M., Steele, L. P., Szopa, S., Yver, C., Viovy, N., and Ciais, P.: Source attribution of the changes in atmospheric methane for 2006–2008, *Atmos. Chem. Phys.*, 11, 3689–3700, doi:10.5194/acp-11-3689-2011, 2011.
- 10 Bridgham, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q.: Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales, *Glob. Change Biol.*, 19, 1325–1346, 2013.
- Choularton, T. W., Gallagher, M. W., Bower, K. N., Fowler, D., Zahniser, M., and Kaye, A.: Trace gas flux measurements at the landscape scale using boundary-layer budgets, *Philos. T. Roy. Soc. A*, 351, 357–368, doi:10.1098/rsta.1995.0039, 1995.
- Christensen, T. R., Johansson, T., Olsrud, M., Strom, L., Lindroth, A., Mastepanov, M., Malmer, N., Friborg, T., Crill, P., and Callaghan, T. V.: A catchment-scale carbon and greenhouse gas budget of a subarctic landscape, *Philos. T. Roy. Soc. A*, 365, 1643–1656, doi:10.1098/rsta.2007.2035, 2007.
- 20 Christensen, T. R., Jackowicz-Korczynski, M., Aurela, M., Crill, P., Heliasz, M., Mastepanov, M., and Friborg, T.: Monitoring the multi-year carbon balance of a Subarctic Palsa Mire with micrometeorological techniques, *Ambio*, 41, 207–217, doi:10.1007/s13280-012-0302-5, 2012.
- Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722, doi:10.5194/gmd-4-701-2011, 2011.
- 30 Cullen, M. J. P.: The unified forecast/climate model, *Meteorol. Mag.*, 1449, 81–94, 1993.
- Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, doi:10.1038/nature04514, 2006.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Desjardins, R. L., MacPherson, J. I., Mahrt, L., Schuepp, P., Pattey, E., Neumann, H., Baldocchi, D., Wofsy, S., Fitzjarrald, D., McCaughey, H., and Joiner, D. W.: Scaling up flux measurements for the boreal forest using aircraft-tower combinations, *J. Geophys. Res.*, 102, 29125–29133, 1997.

5 Draxler, R. R. and Rolph, G. D.: HYSPLIT (Hybrid Single-Particle Lagrangian Integrated Trajectory) Model, NOAA Air Resources Laboratory, Silver Spring, MD, USA, 2003.

Gedney, N. and Cox, P. M.: The sensitivity of global climate model simulations to the representation of soil moisture heterogeneity, *J. Hydrometeorol.*, 4, 1265–1275, 2003.

Gedney, N., Cox, P. M., and Huntingford, C.: Climate feedback from wetland methane emissions. *Geophys. Res. Lett.*, 31, L20503, doi:10.1029/2004GL020919, 2004.

10 Fisher, R., Lowry, D., Wilkin, O., Sriskantharajah, S., and Nisbet, E. G.: High-precision, automated stable isotope analysis of atmospheric methane and carbon dioxide using continuous-flow isotope-ratio mass spectrometry, *Rapid Commun. Mass Sp.*, 20, 200–208, doi:10.1002/rcm.2300, 2006.

15 Fisher, R. E., Sriskantharajah, S., Lowry, D., Lanoisellé, M., Fowler, C. M. R., James, R. H., Hermansen, O., Lund Myhre, C., Stohl, A., Greinert, C., Nisbet-Jones, P. B. R., Mienert, J., Nisbet, E. G.: Arctic methane sources: isotopic evidence for atmospheric inputs, *Geophys. Res. Lett.*, 38, L21803, doi:10.1029/2011GL049319, 2011.

Forster, P. and Ramaswamy, V.: Changes in Atmospheric Constituents and in Radiative Forcing, *Climate Change 2007: the Physical Science Basis*, Cambridge University Press, Cambridge, 129–234, 2007.

20 Friend, A. D.: Terrestrial plant production and climate change, *J. Exp. Bot.*, 61, 1293–1309, 2010.

Friend, A. D. and Kiang, N. Y.: Land surface model development for the GISS GCM: effects of improved canopy physiology on simulated climate, *J. Climate*, 18, 2883–2902, doi:10.1175/JCLI3425.1, 2005.

25 Gallagher, M. W., Choularton, T. W., Bower, K. N., Stromberg, I. M., Beswick, K. M., Fowler, D., and Hargreaves, K. J.: Measurements of methane fluxes on the landscape scale from a wetland area in north Scotland, *Atmos. Environ.*, 28, 2421–2430, doi:10.1016/1352-2310(94)90394-8, 1994.

30 Gerbig, C., Schmitgen, S., Kley, D., Volz-Thomas, A., Dewey, K., and Haaks, D.: An improved fast-response vacuum-UV resonance fluorescence CO instrument, *J. Geophys. Res.-Atmos.*, 104, 1699–1704, doi:10.1029/1998jd100031, 1999.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Isaksen, I. S. A., Gauss, M., Myhre, G., Anthony, K. M. W., and Ruppel, C.: Strong atmospheric chemistry feedback to climate warming from Arctic methane emissions, *Global Biogeochem. Cy.*, 25, GB2002, doi:10.1029/2010gb003845, 2011.

Jackowicz-Korczynski, M., Christensen, T. R., Backstrand, K., Crill, P., Friborg, T., Maste-
 5 tepanov, M., and Strom, L.: Annual cycle of methane emission from a subarctic peatland, *J. Geophys. Res.-Biogeo.*, 115, G02009, doi:10.1029/2008jg000913, 2010.

Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Berga-
 10 maschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S.,
 Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B.,
 Fraser, P. J., Krummel, P. B., Lamarque, J.-F., Langenfelds, R. L., Le Quere, C., Naik, V.,
 O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M.,
 Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P.,
 Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M.,
 15 Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and
 sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/ngeo1955, 2013.

Kort, E. A., Wofsy, S. C., Daube, B. C., Diao, M., Elkins, J. W., Gao, R. S., Hints, E. J.,
 Hurst, D. F., Jimenez, R., Moore, F. L., Spackman, J. R., and Zondlo, M. A.: Atmospheric
 observations of Arctic Ocean methane emissions up to 82° north, *Nat. Geosci.*, 5, 318–321,
 doi:10.1038/ngeo1452, 2012.

20 Le Breton, M., Bacak, A., Muller, J. B. A., O'Shea, S. J., Xiao, P., Ashfold, M. N. R., Cooke, M. C.,
 Batt, R., Shallcross, D. E., Oram, D. E., Forster, G., Bauguitte, S. J.-B., Palmer, P. I., Parrington,
 M., Lewis, A. C., Lee, J. D., and Percival, C. J.: Airborne hydrogen cyanide measure-
 ments using a chemical ionisation mass spectrometer for the plume identification of biomass
 burning forest fires, *Atmos. Chem. Phys.*, 13, 9217–9232, doi:10.5194/acp-13-9217-2013,
 25 2013.

Levy, P. E., Burden, A., Cooper, M. D. A., Dinsmore, K. J., Drewer, J., Evans, C., Fowler, D.,
 Gaiawyn, J., Gray, A., Jones, S. K., Jones, T., McNamara, N. P., Mills, R., Ostle, N., Shep-
 30 pard, L. J., Skiba, U., Sowerby, A., Ward, S. E., and Zielinski, P.: Methane emissions from
 soils: synthesis and analysis of a large UK data set, *Glob. Change Biol.*, 18, 1657–1669,
 doi:10.1111/j.1365-2486.2011.02616.x, 2012.

Lewis, A. C., Evans, M. J., Hopkins, J. R., Punjabi, S., Read, K. A., Purvis, R. M., Andrews, S. J.,
 Moller, S. J., Carpenter, L. J., Lee, J. D., Rickard, A. R., Palmer, P. I., and Parrington, M.: The

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



influence of biomass burning on the global distribution of selected non-methane organic compounds, *Atmos. Chem. Phys.*, 13, 851–867, doi:10.5194/acp-13-851-2013, 2013.

Maanavilja, L., Riutta, T., Aurela, M., Pulkkinen, M., Laurila, T., and Tuittila, E.-S.: Spatial variation in CO₂ exchange at a northern aapa mire, *Biogeochemistry*, 104, 325–345, doi:10.1007/s10533-010-9505-7, 2011.

Mays, K. L., Shepson, P. B., Stirm, B. H., Karion, A., Sweeney, C., and Gurney, K. R.: Aircraft-based measurements of the carbon footprint of Indianapolis, *Environ. Sci. Technol.*, 43, 7816–7823, doi:10.1021/es901326b, 2009.

McMillen, R. T.: An eddy-correlation technique with extended applicability to non-simple terrain, *Bound.-Lay. Meteorol.*, 43, 231–245, doi:10.1007/bf00128405, 1988.

Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, doi:10.5194/bg-10-753-2013, 2013.

Miller, J. B., Gatti, L. V., d'Amelio, M. T. S., Crotwell, A. M., Dlugokencky, E. J., Bakwin, P., Artaxo, P., and Tans, P. P.: Airborne measurements indicate large methane emissions from the eastern Amazon basin, *Geophys. Res. Lett.*, 34, L10809, doi:10.1029/2006gl029213, 2007.

Nisbet, E. G. and Chappellaz, J.: Shifting gear, quickly, *Science*, 324, 477–478, doi:10.1126/science.1172001, 2009.

Niu, G.-Y., Yang, Z.-L., Dickinson, R. E., and Gulden, L. E.: A simple TOPMODEL-based runoff parameterization (SIMTOP) for use in GCMs, *J. Geophys. Res.*, 110, D21106, doi:10.1029/2005JD006111, 2005

O'Connor, F. M., Boucher, O., Gedney, N., Jones, C. D., Folberth, G. A., Coppel, R., Friedlingstein, P., Collins, W. J., Chappellaz, J., Ridley, J., and Johnson, C. E.: Possible role of wetlands, permafrost, and methane hydrates in the methane cycle under future climate change: a review, *Rev. Geophys.*, 48, RG4005, doi:10.1029/2010rg000326, 2010.

O'Shea, S. J., Allen, G., Gallagher, M. W., Bauguutte, S. J.-B., Illingworth, S. M., Le Breton, M., Muller, J. B. A., Percival, C. J., Archibald, A. T., Oram, D. E., Parrington, M., Palmer, P. I., and Lewis, A. C.: Airborne observations of trace gases over boreal Canada during BORTAS:

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

campaign climatology, air mass analysis and enhancement ratios, *Atmos. Chem. Phys.*, 13, 12451–12467, doi:10.5194/acp-13-12451-2013, 2013a.

O'Shea, S. J., Bauguitte, S. J.-B., Gallagher, M. W., Lowry, D., and Percival, C. J.: Development of a cavity-enhanced absorption spectrometer for airborne measurements of CH₄ and CO₂, *Atmos. Meas. Tech.*, 6, 1095–1109, doi:10.5194/amt-6-1095-2013, 2013b.

Oelke, C., Zhang, T. J., and Serreze, M. C.: Modeling evidence for recent warming of the Arctic soil thermal regime, *Geophys. Res. Lett.*, 31, L07208, doi:10.1029/2003gl019300, 2004.

Petrescu, A. M. R., van Huissteden, J., Jackowicz-Korczynski, M., Yurova, A., Christensen, T. R., Crill, P. M., Bäckstrand, K., and Maximov, T. C.: Modelling CH₄ emissions from arctic wetlands: effects of hydrological parameterization, *Biogeosciences*, 5, 111–121, doi:10.5194/bg-5-111-2008, 2008.

Petrescu, A. M. R., van Beek, E. J. R., van Huissteden, J., Prigent, C., Sachs, T., Corradi, C. A. R., Parmentier, F. J. W., and Dolman, A. J.: Modeling regional to global CH₄ emissions of boreal and arctic wetlands, *Global Biogeochem. Cy.*, 24, GB4009, doi:10.1029/2009GB003610, 2010.

Parmentier, F.-J. W., Christensen, T. R., Sorensen, L. L., Rysgaard, S., McGuire, A. D., Miller, P. A., and Walker, D. A.: The impact of lower sea-ice extent on Arctic greenhouse-gas exchange, *Nature Clim. Change*, 3, 195–202, 2013.

Peischl, J., Ryerson, T. B., Holloway, J. S., Trainer, M., Andrews, A. E., Atlas, E. L., Blake, D. R., Daube, B. C., Dlugokencky, E. J., Fischer, M. L., Goldstein, A. H., Guha, A., Karl, T., Kofler, J., Kosciuch, E., Misztal, P. K., Perring, A. E., Pollack, I. B., Santoni, G. W., Schwarz, J. P., Spackman, J. R., Wofsy, S. C., and Parrish, D. D.: Airborne observations of methane emissions from rice cultivation in the Sacramento Valley of California, *J. Geophys. Res.-Atmos.*, 117, D00V25, doi:10.1029/2012jd017994, 2012.

Pelletier, L., Moore, T. R., Roulet, N. T., Garneau, M., and Beaulieu-Audy, V.: Methane fluxes from three peatlands in the La Grande Riviere watershed, James Bay lowland, Canada, *J. Geophys. Res.-Biogeo.*, 112, G01018, doi:10.1029/2006jg000216, 2007.

Pickett-Heaps, C. A., Jacob, D. J., Wecht, K. J., Kort, E. A., Wofsy, S. C., Diskin, G. S., Worthy, D. E. J., Kaplan, J. O., Bey, I., and Drevet, J.: Magnitude and seasonality of wetland methane emissions from the Hudson Bay Lowlands (Canada), *Atmos. Chem. Phys.*, 11, 3773–3779, doi:10.5194/acp-11-3773-2011, 2011.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ping, C.-L., Michaelson, G. J., Jorgenson, M. T., Kimble, J. M., Epstein, H., Romanovsky, V. E., and Walker, D. A.: High stocks of soil organic carbon in the North American Arctic region, *Nat. Geosci.*, 1, 615–619, doi:10.1038/ngeo284, 2008.
- Ridgwell, A. J., Marshall, S. J., and Gregson, K.: Consumption of atmospheric methane by soils: a process-based model, *Global Biogeochem. Cy.*, 13, 59–70, doi:10.1029/1998gb900004, 1999.
- Ryerson, T. B., Buhr, M. P., Frost, G. J., Goldan, P. D., Holloway, J. S., Hubler, G., Jobson, B. T., Kuster, W. C., McKeen, S. A., Parrish, D. D., Roberts, J. M., Sueper, D. T., Trainer, M., Williams, J., and Fehsenfeld, F. C.: Emissions lifetimes and ozone formation in power plant plumes, *J. Geophys. Res.-Atmos.*, 103, 22569–22583, doi:10.1029/98jd01620, 1998.
- Ryall, D. B. and Maryon, R. H.: Validation of the UK Met Office's NAME model against the ETEX dataset, *Atmos. Environ.*, 32, 4265–4276, 1998.
- Ryall, D. B., Maryon, R. H., Derwent, R. G., and Simmonds, P. G.: Modelling long-range transport of CFCs to Mace Head, Ireland, *Q. J. Roy. Meteor. Soc.*, 124, 417–446, 1998.
- Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Del Genio, A., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner, J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D., Romanou, A., Russell, G. L., Sato, M., Shindell, D. T., Stone, P. H., Sun, S., Tausnev, N., Thresher, D., and Yao, M.-S.: Present day atmospheric simulations using giss model: comparison to in-situ, satellite and reanalysis data, *J. Climate*, 19, 153–192, 2006.
- Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, doi:10.1038/nature08031, 2009.
- Shakhova, N., Semiletov, I., Salyuk, A., Yusupov, V., Kosmach, D., and Gustafsson, O.: Extensive methane venting to the atmosphere from sediments of the East Siberian Arctic Shelf, *Science*, 327, 1246–1250, doi:10.1126/science.1182221, 2010.
- Sriskantharajah, S., Fisher, R., Lowry, D., Aalto, T., Hatakka, T., Aurela, M., Laurila, T., Lohila, A., Kuitunen, E., and Nisbet, E.: Stable carbon isotope signatures of methane from a Finnish subarctic wetland, *Tellus B*, 64, 18818 doi:10.3402/tellusb.v64i0.18818, 2012.
- Sitch, S., McGuire, A. D., Kimball, J., Gedney, N., Gamon, J., Engstrom, R., Wolf, A., Zhuang, Q., Clein, J., and McDonald, K. C.: Assessing the carbon balance of circumpo-

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lar Arctic tundra using remote sensing and process modeling, *Ecol. Appl.*, 17, 213–234, doi:10.1890/1051-0761(2007)017[0213:atcboc]2.0.co;2, 2007.

Smith, L. C., MacDonald, G. M., Velichko, A. A., Beilman, D. W., Borisova, O. K., Frey, K. E., Kremenetski, K. V., and Sheng, Y.: Siberian peatlands a net carbon sink and global methane source since the early Holocene, *Science*, 303, 353–356, doi:10.1126/science.1090553, 2004.

Strom, L. and Christensen, T. R.: Below ground carbon turnover and greenhouse gas exchanges in a sub-arctic wetland, *Soil Biol. Biochem.*, 39, 1689–1698, doi:10.1016/j.soilbio.2007.01.019, 2007.

Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Kluwer Academic Publishers, Dordrecht, 670 pp., 1988.

Vay, S. A., Choi, Y., Vadrevu, K. P., Blake, D. R., Tyler, S. C., Wisthaler, A., Hecobian, A., Kondo, Y., Diskin, G. S., Sachse, G. W., Woo, J. H., Weinheimer, A. J., Burkhardt, J. F., Stohl, A., and Wennberg, P. O.: Patterns of CO₂ and radiocarbon across high northern latitudes during International Polar Year 2008, *J. Geophys. Res.-Atmos.*, 116, D14301, doi:10.1029/2011jd015643, 2011.

Viovy, N. and Ciais, P.: A combined dataset for ecosystem modelling, available at: http://dods.extra.cea.fr/store/p529viov/cruncep/V4_1901_2012/ (last access: 10 July 2013), 2009.

Walter, K. M., Edwards, M. E., Grosse, G., Zimov, S. A., and Chapin III, F. S.: Thermokarst lakes as a source of atmospheric CH₄ during the last deglaciation, *Science*, 318, 633–636, doi:10.1126/science.1142924, 2007.

Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHyMe v1.3.1, *Geosci. Model Dev.*, 3, 565–584, doi:10.5194/gmd-3-565-2010, 2010.

Webb, E. K., Pearman, G. I., and Leuning, R.: Correction of flux measurements for density effects due to heat and water vapour-transfer, *Q. J. Roy. Meteor. Soc.*, 106, 85–100, doi:10.1002/qj.49710644707, 1980.

Westbrook, G. K., Thatcher, K. E., Rohling, E. J., Piotrowski, A. M., Paelike, H., Osborne, A. H., Nisbet, E. G., Minshull, T. A., Lanoiselle, M., James, R. H., Huehnerbach, V., Green, D., Fisher, R. E., Crocker, A. J., Chabert, A., Bolton, C., Beszczynska-Moeller, A., Berndt, C., and Aquilina, A.: Escape of methane gas from the seabed along the West Spitsbergen continental margin, *Geophys. Res. Lett.*, 36, L15608, doi:10.1029/2009gl039191, 2009.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



White, W. H., Anderson, J. A., Blumenthal, D. L., Husar, R. B., Gillani, N. V., Husar, J. D., and Wilson, W. E.: Formation and transport of secondary air-pollutants-ozone and aerosols in St-Louise Urban Plume, *Science*, 194, 187–189, doi:10.1126/science.959846, 1976.

5 Wratt, D. S., Gimson, N. R., Brailsford, G. W., Lassey, K. R., Bromley, A. M., and Bell, M. J.: Estimating regional methane emissions from agriculture using aircraft measurements of concentration profiles, *Atmos. Environ.*, 35, 497–508, doi:10.1016/s1352-2310(00)00336-8, 2001.

10 Zhuang, Q., Melillo, J. M., Sarofim, M. C., Kicklighter, D. W., McGuire, A. D., Felzer, B. S., Sokolov, A., Prinn, R. G., Steudler, P. A., and Hu, S.: CO₂ and CH₄ exchanges between land ecosystems and the atmosphere in northern high latitudes over the 21st century, *Geophys. Res. Lett.*, 33, L17403, doi:10.1029/2006gl026972, 2006.

Zimov, S. A., Davydov, S. P., Zimova, G. M., Davydova, A. I., Schuur, E. A. G., Dutta, K., and Chapin III, F. S.: Permafrost carbon: stock and decomposability of a globally significant carbon pool, *Geophys. Res. Lett.*, 33, L20502, doi:10.1029/2006gl027484, 2006a.

15 Zimov, S. A., Schuur, E. A. G., and Chapin, F. S.: Permafrost and the global carbon budget, *Science*, 312, 1612–1613, doi:10.1126/science.1128908, 2006b.

Table 1. Land classification key corresponding to Fig. 1 from Corine land cover 2006. Also included is the proportion of the aircrafts footprint that each classification accounted for during the B720 E–W transects.

Number	Land type	Proportion of footprint during E–W transect (%)
2	Discontinuous urban fabric	0.1
3	Industrial or commercial units	0.0
4	Road and rail networks and associated land	0.0
6	Airports	0.0
7	Mineral extraction sites	0.0
8	Dump sites	0.0
10	Green urban areas	0.0
11	Sport and leisure facilities	0.0
12	Non-irrigated arable land	0.0
18	Pastures	0.1
20	Complex cultivation patterns	0.0
21	Land principally occupied by agriculture with significant areas of natural vegetation	0.1
23	Broad-leaved forest	10.4
24	Coniferous forest	24.4
25	Mixed forest	16.3
26	Natural grasslands	0.1
27	Moors and heathland	8.3
29	Transitional woodland-shrub	13.8
31	Bare rocks	0.1
32	Sparsely vegetated areas	2.3
33	Burnt areas	0.0
34	Glaciers and perpetual snow	0.0
35	Inland marshes	0.1
36	Peat bogs	19.6
40	Water courses	0.5
41	Water bodies	3.7

Methane and carbon dioxide fluxes

S. J. O’Shea et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Table 2. Mean fluxes determined using the FAAM BAe-146, chamber and eddy covariance techniques. All uncertainties given are as one standard deviation (1σ). Chamber measurements are separated into the geometric mean of all seasonally averaged measurements and only those from the 22 July 2012. A weighted mean of the wetland and forest chamber fluxes is calculated using the number of occurrences of each land type within the east–west transect (Fig. 1b).

		Flux ($\text{mg h}^{-1} \text{m}^{-2}$)	
		CH ₄	CO ₂
FAAM BAe-146	Eastward transect	1.1 ± 0.6	-375 ± 202
	Westward transect	1.6 ± 0.5	-357 ± 135
	Both transects	1.2 ± 0.5	-350 ± 143
	Whole air samples	1.0 ± 0.6	-315 ± 368
Eddy covariance	Summer	4.5 ± 1.4	-135 ± 344
	Summer day	4.5 ± 1.2	-309 ± 306
	Summer night	4.4 ± 1.6	71 ± 264
	22 Jul 2012	4.5 ± 0.9	
	22 Jul 2012 day	4.9 ± 0.6	
	22 Jul 2012 night	4.4 ± 1.0	
Chamber	Wetland summer	4.5 ± 3.7	
	Wetland 22 Jul 2012	5.6 ± 5.6	
	Forest summer	0.05 ± 0.07	
	Forest 22 Jul 2012	-0.07 ± 0.05	
	Weighted average	1.3 ± 1.0	
	Weighted average 22 Jul 2012	1.5 ± 1.6	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Distribution of the modelled hourly wetland methane emission fluxes ($\text{mg CH}_4 \text{ m}^{-2} \text{ h}^{-1}$) for the domain ($20.0\text{--}29.0^\circ \text{ E}$, $67.5\text{--}68.5^\circ \text{ N}$) every July and August between 1980 and 2010.

Hourly emission flux ($\text{mg CH}_4 \text{ m}^{-2} \text{ h}^{-1}$)	JULES	Hybrid8
	Jul–Aug 1980–2010	Jul–Aug 1980–2010
Number of hours	1 660 608	1 660 608
Minimum	0.0	−0.0018
Lower quartile	0.0	0.013
Median	0.033	0.023
Upper quartile	0.085	0.095
Maximum	0.38	4.62
Mean	0.054	0.073

Methane and carbon dioxide fluxes

S. J. O'Shea et al.




Fig. 1. (a) FAAM BAe-146 flight track for flight B720 (22 July 2012). Observations of CH_4 in the PBL are coloured according to the legend. Black diamonds mark Kiruna, Sodankylä and Kaamanen. (b) Flight track where the surface is coloured using the land use type (Corine land cover 2006). Numbers correspond to land types given in Table 1.

Methane and carbon dioxide fluxes

S. J. O'Shea et al.




Fig. 2. Keeling plot showing PBL measurements of $\delta^{13}\text{C}\text{-CH}_4$ during flight on the 22 July 2012 (B720). The intercept of $-72 \pm 4.4 \text{ ‰}$ is representative of a wetland source of CH_4 .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Methane and carbon dioxide fluxes

S. J. O'Shea et al.




Fig. 3. (a) CH₄ and (b) CO₂ observations along a flight transect, which was aligned with the prevailing wind direction. The origin is 20° E, 68° N and the transect extends in an eastward direction. The gradients observed in both species were used to determine a net emission flux for the region using Eq. (3).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. (a) Ascending (01:00 GMT) and (b) descending (15:00 GMT) potential temperature profiles performed over Sodankylä during flight B720, used to determine the boundary layer height as described in the text.




Fig. 5. A comparison between different techniques used to determine fluxes. The box extents define the 25th and 75th percentiles, and whiskers are the 10th and 90th percentiles. Note: the Eddy covariance percentiles are for daytime (06:00 to 18:00 GMT) only. Forest and wetland chamber fluxes represent summer seasonal statistics for 60 chamber measurements (21 in forest regions and 39 in wetland regions). The scaled chamber (black circle) is determined by averaging the wetland and forest chamber fluxes as described in Sect. 3.3. The FAAM BAe-146 and scaled chamber error bar shows the 1σ uncertainty as described in Sect. 3.1.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Methane and carbon dioxide fluxes

S. J. O'Shea et al.



1h-average CH₄ increment (ppbv) averaged over 67.75–68.00N at 11Z on 22 July 20121h-average CH₄ increment (ppbv) averaged over 67.75–68.00N at 16Z on 22 July 2012

Fig. 6. (a) Dispersion model results from NAME for the mixing ratio of CH₄ originating from the local wetlands in a cross section of the atmosphere averaged over 67.75 to 68.00° N and 1 h surrounding 11:00 GMT on 22 July 2012. The local wetland CH₄ source was defined as a 1.2 mgCH₄ h⁻¹ m⁻² source emitted from the ground between 20 to 28° E and between 67 to 69.5° N. (b) shows the same but for a 1 h average surrounding 16:00 GMT on the same day.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Methane and carbon dioxide fluxes

S. J. O'Shea et al.




Fig. 7. CH₄ and CO₂ hourly fluxes at Sodankylä wetland site between 1 July and 15 August 2012 determined using the eddy covariance technique. CH₄ diurnal variation is noted to be small. Net CO₂ uptake occurs during the day, with net emission during the night.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Methane and carbon dioxide fluxes

S. J. O'Shea et al.




Fig. 8. Daytime (06:00–18:00) CH_4 and CO_2 fluxes at Sodankylä wetland site between 1 July and 15 August 2012 determined using the eddy covariance technique. Note: CO_2 fluxes are not shown for the period 14 July 2012 to 1 August 2012 as it was not possible to calibrate the LI-7200's CO_2 channel in that period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)