

# Origin, variability and age of biomass burning plumes intercepted during BORTAS-B

D. P. Finch<sup>1</sup>, P. I. Palmer<sup>1</sup>, and M. Parrington<sup>1,2</sup>

<sup>1</sup>School of GeoSciences, University of Edinburgh, UK

<sup>2</sup>Now at the European Centre for Medium-range Weather Forecasts, Reading, UK

*Correspondence to:* D. P. Finch  
(d.finch@ed.ac.uk)

## Abstract.

We use the GEOS-Chem atmospheric chemistry transport model to interpret aircraft measurements of carbon monoxide (CO) in biomass burning outflow taken during the 2011 BORTAS-B campaign over eastern Canada. The model has some skill reproducing the observed variability (Spearman's rank correlation  $r_s = 0.65$  but has a positive (negative) bias for observations  $< 100$  ppb ( $> 300$  ppb). We find that observed CO variations are largely due to NW North American biomass burning, as expected, with smaller and less variable contributions from fossil fuel combustion from eastern Asia and NE North America. To help interpret observed variations of CO we develop an Eulerian effective physical age of emissions ( $\bar{A}$ ) metric, accounting for mixing and chemical decay, which we apply to pyrogenic emissions of CO. We find that during BORTAS-B the age of emissions intercepted over Halifax, Nova Scotia is typically 4–11 days, and on occasion as young as two days. We show that  $\bar{A}$  is typically 1–5 days older than the associated photochemical ages inferred from colocated measurements of different hydrocarbons. We find that the median difference between the age measures ( $\Delta\tau$ ) in plumes ( $\text{CH}_3\text{CN} > 150$  ppt) peaks at 3 days. This corresponds to a chemical retardation of 50%. We find a strong relationship in biomass burning plumes between  $\bar{A}$  and  $\Delta\tau$  ( $r^2 = 0.80$ ), which is not present outwith these plumes ( $r^2 = 0.28$ ). We argue that these observed relationships, together with a robust observed relationship between CO and black carbon aerosol during BORTAS-B ( $r^2 > 0.7$ ), form the basis of indirect evidence that aerosols co-emitted with gases during pyrolysis markedly slowed down the plume photochemistry during BORTAS-B with respect to photochemistry at the same latitude and altitude in clear skies.



## 1 Introduction

The open burning of biomass is an inefficient combustion process, resulting in the release of a wide range of chemically reactive gases and particles that contribute to the production of ozone in the troposphere (Goode et al., 2000; Koppmann et al., 2005; Akagi et al., 2011), with implications for national surface air quality and air quality mitigation strategies. However, the rate and extent of photochemical ozone production in biomass burning outflow is still a matter of debate that largely reflects the sensitivity of results to environmental conditions (Jaffe and Wigder, 2012). In this paper we present an analysis of measurements of carbon monoxide (CO) from the BORTAS-B aircraft campaign during July 2011 (Palmer et al., 2013), in conjunction with a 3-D chemistry transport model to understand the processes that determine observed CO variability and relate the ages of emissions to the observed photochemical production of ozone.

Ozone production within biomass burning plumes intercepted during the BORTAS-B campaign (Palmer et al., 2013) has previously been studied using photochemical age (Parrington et al., 2013; Parrish et al., 2007). Estimating the photochemical age of a pyrogenic air mass relies on an accurate knowledge of the OH radical, and assumes constant trace gas emission factors from pyrogenic processes. However, despite these weaknesses the photochemical age is still a useful metric for helping to interpret observed trace gas variations. We introduce a complementary (weighted-mean) effective physical age metric  $\bar{A}$ . Using  $\bar{A}$  we can infer age distributions within an air mass and by comparing  $\bar{A}$  with the photochemical age we can estimate how the physical and chemical environment of the plume has modified the speed of the in situ plume chemistry.

The main source of CO is from the incomplete combustion of fossil fuel, biomass, and biofuel. There is also a source of CO from the oxidation from methane and non-methane volatile organic compounds (NMVOCs) (Duncan et al., 2007). The main sink is from the oxidation by the hydroxyl radical (OH), resulting in an atmospheric lifetime of weeks to months depending on latitude and season. We use airborne CO measurements from phase B of the Quantifying the impacts of Boreal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites (BORTAS-B) project, July 2011. The overall objective of BORTAS was to better understand the production of tropospheric ozone in respect to the chemical evolution of plumes from boreal forest fires, which was achieved by integrating aircraft (Lewis et al., 2013; Le Breton et al., 2013; O'Shea et al., 2013), surface (Gibson et al., 2013; Griffin et al., 2013), sonde (Parrington et al., 2012), and satellite measurements Tereszchuk et al. (2013) of atmospheric composition. Phase A of BORTAS was conducted without aircraft in July 2010 (Parrington et al., 2012).

In the next section we briefly describe the CO data we analyse. The GEOS-Chem chemistry transport model is described in section 3, including a description of a new age of emission calculation which we use to interpret the data. Our results are reported in section 4, including a statistical analysis of the data and a model interpretation of the data. We conclude in section 5.

## 2 Data and Methods

### 2.1 BORTAS-B Carbon Monoxide and CH<sub>3</sub>CN Data

Here we use data exclusively from the BORTAS-B aircraft campaign. The focus of the work shown here is the analysis of CO measurements, which are operated by the Facility for Airborne Atmospheric Measurements on the BAe-146 atmospheric research aircraft using a fast-response vacuum-UV resonance fluorescence instrument (Gerbig et al., 1999). The instrument has an averaging time of 1 s and a precision and accuracy of 1 ppb and 3%, respectively. We use measurements of acetonitrile (CH<sub>3</sub>CN, not shown), an additional tracer of biomass burning, measured by Proton Transfer Reaction Mass Spectrometer (Murphy et al., 2010) to isolate plumes within BORTAS-B. These measurements have a mean precision of 37 ppt during BORTAS-B (Palmer et al., 2013). We define plumes as CO measurements corresponding to CH<sub>3</sub>CN > 150 ppt.

### 2.2 The GEOS-Chem Atmospheric Chemistry Model

We use the GEOS-Chem atmospheric chemistry model ([www.geos-chem.org](http://www.geos-chem.org)) to interpret the BORTAS-B CO measurement. The model has been documented extensively (e.g., Bey et al. (2001); Duncan et al. (2007); Gonzi et al. (2011); Parrington et al. (2012)) and here we only include those details relevant to our study.

We use v9-01-03 of the model, driven by GEOS-5 assimilated meteorological data from the NASA Global Modelling and Assimilation Office (GMAO) Goddard Earth Observing System (GEOS). For global simulations we use a spatial resolution of 2° latitude by 2.5° longitude (a degradation of the native resolution of 0.5° × 0.667°) with 47 vertical levels with a temporal resolution of 30 minutes. For the model runs we use to compare against BORTAS-B aircraft data and the  $\bar{A}$  calculations we use the the native model resolution with 47 vertical levels. We use the Global Fire Emissions Database (GFED-3), describing biomass burning emissions (Giglio et al., 2010), which has a three-hour temporal resolution; fossil fuel emissions from the Emissions Database for Global Atmospheric Research (EDGAR, Olivier et al. (1999)); and biogenic emissions from the Model of Emissions of Gases and Aerosol from Nature (MEGAN, Guenther et al. (2006)). We report model calculations from the summers (JJA) of 2008–2011. We initialize the model in 2007, using previous model output, and run using a single total CO tracer for 9 months until our study period JJA 2008. During the summer periods we use “tagged” tracers (described below) and between the successive summer periods we collapse these tagged tracers back to the single tracer for computational expediency.

For the CO attribution calculations, we use a “tagged” version of the model (e.g., Jones et al. (2003); Palmer et al. (2003, 2006); Feng et al. (2009); Fisher et al. (2010)), which uses pre-calculated monthly 3-D OH fields. Using these fields allows us to linearly decompose the CO originating from specific processes and geographical regions. Figure 1 shows the geographical regions we use. For biomass burning in the northern hemisphere we split North America into four quadrants, consider

Europe as one region, and split Russia/Siberia into three regions (western, mid, and eastern). We show below that most of the observed CO over eastern Canada during JJA originates from these regions. For northern hemisphere fossil fuel we have combined some regions that do not play a significant role in the interpretation of the BORTAS-B data. The chemical source of CO from the oxidation of methane and NMVOCs is treated as one global tracer. In total, we have 28 tracers (including the background) that sum to the total atmospheric CO. Wherever we compare the model against data we sample the model at the time and location of the measurement.

### 2.3 Age of Emission Model Calculation

We use the same model structure for the “tagged” CO simulation to calculate  $\bar{A}$ . To calculate  $\bar{A}$ , we adapt the model to instead emit an arbitrary constant amount to a day-specific tracer wherever there is active burning during our study period (informed by GFED-3). Once emitted the tracer is left to disperse. We assume a constant chemical lifetime of 60 days from the oxidation by OH, corresponding to an OH concentration of  $1.9 \times 10^6$  molec/cm<sup>-3</sup>. At the end of a 31-day simulation for July, say, we have 31 tracers. The age of each tracer is simply the number of days since the start of the run minus the tracer number (not value).

To account for older air being subject to more dispersion than younger air we define an effective age of air  $\bar{A}$ :

$$\bar{A} = \frac{\sum_i A_i L_{M,i} L_{C,i}}{\sum_i L_{M,i} L_{C,i}}, \quad (1)$$

which represents a weighted mean of the age of each tracer  $A_i$ , the value of each tracer  $L_{M,i}$  that is a measure of atmospheric mixing processes, and the chemical lifetime of each tracer  $L_{C,i}$ . More generally, we can extend this formulation to include other sink terms such as, for example, dry and wet deposition. For  $L_{C,i}$  we assume a constant chemical lifetime of 60 days as described above. This method does not account for the magnitude of CO emitted from any fire, regardless of its size.

For this paper, calculating  $\bar{A}$  allows us to quantify the physical age of emissions intercepted during BORTAS-B, providing additional information to interpret the chemical signature of the sampled air masses. It also allows us to determine whether the air masses intercepted during BORTAS-B were representative of that summer and of a similar period from preceding years.

We compare our  $\bar{A}$  estimates against the associated photochemical ages of plumes intercepted in BORTAS-B (Parrington et al., 2013). The photochemical age is the time taken for a tracer to be photochemically removed from a biomass burning plume. It is based on ratios of NMVOCs close to the emission source (here, determined by flight B626 in the BORTAS-B campaign over Northwestern Ontario on the 26th July 2011) and the ratio of NMVOCs at the time of observations. Assuming a representative OH value, an age determined solely by photochemistry can be calculated for a plume.



### 3 Results

#### 3.1 Statistical Analysis of BORTAS-B CO Data

Figure 2 show the mean statistics of the nested model and observed CO concentrations during BORTAS-B. Mean and median values of observed CO suggests the model has a small positive bias; mean (median) model minus observed CO residual of 18.4 ppb (-0.1 ppb). We find the model has a positive bias below observations of 100 ppb and a negative bias for observations  $>300$  ppb. The largest discrepancies between the model and the observations generally occur at the largest values of CO. The 99th percentile value for model and observed CO concentrations are 670 ppb and 436 ppb, respectively. The interquartile range for BORTAS-B observed CO is 87–133 ppb with a mode of 87 ppb. This is consistent with background CO measurements during the NASA ARCTAS-B campaign (Arctic Research of the Composition of the Troposphere from Aircraft and Satellites, Jacob et al. (2010)). The largest observed CO concentrations during BORTAS-B are larger than those observed during ARCTAS-B (Liang et al., 2011). Figure 3 shows that the relative model error [model minus observation]/model is typically within  $\pm 0.5$  but has a range of  $\pm 1.0$ . The model overestimates observed CO concentrations between the surface and 800 hPa, reflecting the outflow of anthropogenic and biomass burning pollution, with the largest discrepancy around 800 hPa. The model generally has more skill in reproducing the variability of observed CO during BORTAS-B in the below 4 km (Spearman’s rank correlation  $r_s = 0.65$ ) than above 4 km ( $r_s = 0.19$ ). We use Spearman’s rank correlation because it is a more appropriate metric that can describe the model ability to reproduce qualitatively observed elevated values (plumes) above a background but not necessarily capture the value of these elevated values. The inability of the model to capture these elevated values may reflect errors in emissions or atmospheric transport.

#### 3.2 Tagged CO Model Output

Table 1 shows the tagged model analysis sampled at the times and locations of the BORTAS-B CO measurements. The largest source of CO and the largest source of CO variability during BORTAS was from NE North American biomass burning, as expected. There are also large but much less variable contributions from the background (air older than JJA) and from the oxidation of methane and NMVOCs. CO produced by  $\text{CH}_4$  oxidation typically contributes around 30% to global concentrations (Duncan et al., 2007).

Figure 4 shows the mean JJA model contributions of total surface CO from different geographically based sources 2008–2011, described using a horizontal resolution of  $2^\circ$  latitude by  $2.5^\circ$  longitude. The contribution from biomass burning over NW North America is broadly constant from year to year, although the distribution of the fires varies substantially, with Alaska playing a dominant role only in 2009 during our study period. The magnitude and the distribution of fossil fuel emissions from NE North America (predominately the NE USA) appear reasonably consistent over

the four years, with emissions generally travelling up the eastern seaboard with eventual outflow to the Atlantic ocean close to Halifax, Nova Scotia. Similar to North America, Siberian biomass burning has substantial spatial variability from year to year, with their location playing a key role in determining their eventual impact on North America and Europe. During 2010 the largest CO concentrations originated from East Siberia and spread across the northern hemisphere. In contrast, during BORTAS-B in 2011 most of the fire activity was further SW and had less of an impact over eastern Canada. Fires from mid-Siberia had a larger influence on total CO during 2008–2009, with very little activity during 2010–2011. There is a consistently small source from fossil fuel combustion from East Asia (not shown), peaking at around 400 ppb over China but quickly dropping off to around 10 ppb by the time it has crossed the Pacific. Differences between Figure 4 and Figure 9 from Palmer et al. (2013), also showing polar CO concentrations during BORTAS-B, are due to different biomass burning inventories. Here, we use GFED-3 (see above) and Palmer et al. (2013) used the Fire Locating and Modeling of Burning Emissions inventory (Reid et al., 2009).

Figure 5 shows that biomass burning from the NW North America is still a dominant factor in the variability of total CO in the free troposphere. During 2010, these fires contributed around 50 ppb of CO into the upper troposphere causing widespread pollution during July. Typically these emissions contribute about 10 ppb of CO over Europe. Contributions from east and mid Siberia appear to be consistent over the four years except during 2010 when there is very little material transported into the free troposphere. In general, the magnitude and distribution of the fossil fuel source is consistent across the four years with weather systems lifting up surface emission to the free troposphere.

The widespread and persistent source of CO (approximately 10 ppb) from Asian anthropogenic sources over the northern hemisphere (not shown) agrees with the ARCTAS-A study (Fisher et al., 2010) and ARCTAS-B (Bian et al., 2013). Relative to ARCTAS-A, BORTAS-B generally shows a much larger contribution to the total CO from boreal biomass burning, reflecting the timing of ARCTAS-A in April 2008 before the beginning of the main fire season. For JJA 2008–2011 we find boreal biomass burning represents a significant contribution to the total surface CO over the Western Arctic region defined as 50°N–90°N, 170°W–40°W, following Bian et al. (2013). Analysis of ARCTAS-B data showed that boreal biomass burning contributed approximately 25% of CO in this region during July 2008 (Bian et al., 2013). We find boreal biomass burning contributes 25%–45% of the total CO for the defined Western Arctic region for all years, peaking at >90% of the total CO over intense burning areas. During 2008 we find our results are broadly consistent with Bian et al. (2013) but at the lower end of their range. The discrepancy between these results is likely due to using different emission inventories, with Bian et al. (2013) using the Quick Fire Emissions Database.

### 195 3.3 Effective Physical Age of Air

Figure 6 shows the mean model statistics for  $\bar{A}$  during July 2008–2011 at 95°W and 50°W, averaged over 45°–60°N, representing the approximate western and eastern boundaries of the measurements sampled during BORTAS-B. We consider four altitude bins, corresponding to the boundary layer (0–2 km), lower and mid troposphere (2–4 km and 4–6 km, respectively), and upper troposphere  
 200 (>6 km). At the western boundary, emissions from 2008 have a median age of seven days, 2009 has a median age of 10 days, and 2010 and 2011 have a median age of five days. We find the older age of emissions during 2009 is due to fewer fires along the western boundary and consequently a larger influence from Alaska and further afield. Air sampled at the eastern boundary is older, as expected with the except of the upper troposphere (>6 km) which shows similar medians from the western  
 205 boundary to the eastern boundary. The eastern boundary for all years and at all altitudes have median values between 11 and 13 days and a range of 2–23 days. The difference in  $\bar{A}$  between the two boundaries decreases with altitude; above 4 km, ages are indistinguishable because air is more well-mixed and influenced by emissions outside of the domain. The age of emissions clearly get older moving towards the Atlantic, with an increase in median age of approximately three days for all  
 210 years except 2009, when the age difference between the west and east boundaries is approximately one day in the lower troposphere and insignificant at higher altitudes. We attribute this lack of ageing in 2009 to fewer Canadian fires during July of this year, as mentioned above. Boreal biomass burning in 2009 started earlier in the year and was more prominent in Alaska and Siberia. Emissions tend to be older at 50°W at lower altitudes (not shown). This may be a result from fresher emissions getting  
 215 lofted higher into the atmosphere as they travel. Figure 6 also shows that the median value of  $\bar{A}$  (six days) sampled during BORTAS-B falls within the range of model emission ages at 95°W, with the majority of measurements taken 65°–40°W. We find a similar observed median value of  $\bar{A}$  for all altitudes, which is typically lower than the model free troposphere. This bias towards younger ages reflects the sampling strategy of BORTAS-B that was to intercept fresh biomass biomass plumes.

220 Figure 7 shows an example of  $\bar{A}$  on 20th July 2011, which is used as a case study in other BORTAS studies (e.g., Griffin et al. (2013); Franklin et al. (2014)). The longitudinal cross section shows fresh emissions from the Thunder Bay region (50°N, 88°W), peaking at approximately a week old on the 20th of July (corresponding to the 14th–15th July), that eventually age as they are transported eastwards. When they were intercepted by the surface measurements over Toronto (43.70°N,  
 225 69.40°W) (Griffin et al., 2013) and Halifax (44.6°N, 63.59°W) (Franklin et al., 2014) emissions are typically 5–7 days old. Figure 7 shows the transported air, intercepted at 63°W, is composed of a young plume (4–5 days old) surrounded by older air (seven days old) over 47°–55°N. Previous analysis used the NOAA HYbrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model (Draxler and Hess, 1998), driven by meteorological reanalyses data from the National Center for  
 230 Environmental Prediction (NCEP) Global Data Assimilation Program (GDAS), to interpret ground-based remote-sensing data collected during BORTAS-B (Griffin et al., 2013; Franklin et al., 2014).

We find our Eulerian age of emission estimates are typically a day older than those determined by the the HYSPLIT model during the 19–21 July 2011 when ground-based observations reported elevated concentrations due to biomass burning (Gibson et al., 2013; Griffin et al., 2013; Franklin et al., 2014).

Figure 8 shows the mean statistics of  $\bar{A}$  intercepted during BORTAS-B are generally consistent with the associated median photochemical age, determined using NMVOC ratios, of six days (Partridge et al., 2013). The photochemical ages range 0–16 days while  $\bar{A}$  has a range of 2–14 days. There are substantial differences between the photochemical and  $\bar{A}$  for each flight. Figure 9 shows that  $\bar{A}$  minus photochemical age ( $\Delta\tau$ ) have distinct frequency distributions within and outwith plumes. Within plumes, the distribution peaks at  $\Delta\tau=2$ –4 days (median 1.8 days), while outwith plumes there is a relatively flat distribution. We find no significant difference to our results if we average the photochemical ages onto the model grid prior to the analysis. Figure 9 shows that high values of CO relate to  $\Delta\tau$  of 2–4 days, corresponding to a mean (median) fractional difference of 0.44 (0.40); and a weaker relationship between  $\Delta\tau$  outwith plumes, corresponding to a mean (median) fractional difference of 0.11 (0.10). Figure 9 also shows that  $\Delta\tau$  increases with  $\bar{A}$  within plumes ( $r^2=0.80$ ) from close to zero at 3–4 days to  $>5$  days for plumes older than 10 days. There is a much weaker relationship between these two age variables outwith plumes ( $r^2=0.28$ ).

#### 4 Concluding Remarks

We used the GEOS-Chem global atmospheric chemistry model to interpret observed variations of CO taken during the BORTAS-B aircraft campaign over eastern Canada in July-August 2011. We reported a mean difference between the observations and the model for the whole atmosphere of 18.4 ppb and a median of -0.1 ppb. The distribution of these residuals has a high kurtosis, with the majority of residuals within 50 ppb of the mean value. The model has a positive bias below observed values of 100 ppb and a negative bias above 300 ppb. We found that the larger differences between the model and the observations in the mid troposphere, where we found pyrogenic CO peaked. The model has some skill at reproducing the mean observed statistics in the lower troposphere (Spearman’s rank correlation  $r_s = 0.65$ ), and less skill in the higher troposphere ( $r_s = 0.19$ ). The difference in model skill at different altitudes is likely due to a) misdiagnosing sub-gridscale vertical mixing of pyrogenic material lofted by surface heating due to fires, and b) errors in biomass burning emission inventories.

Using a linearly-decomposed version of the model we found that most of the observed variability in CO concentration during BORTAS-B was due to Canadian biomass burning, as expected, with a smaller contribution from Siberian biomass burning and NE North American fossil fuel combustion.

We used the model to put BORTAS-B into the wider temporal context of 2008–2011. We found that North American biomass burning is broadly constant (45% of total CO) over this period although the

spatial distribution of fires varies substantially. The variation of Siberian biomass burning is more extreme with large contributions to total CO over North American during some years (2008–2009) and very little activity in other years (2010–2011), reflecting the spatial extent and geographical position of the fires. Based on our analysis of the source contributions to North American CO over during the (limited) four-year period we conclude that BORTAS-B (2011) was not anomalous.

Previous work has shown that ozone production within biomass burning plumes can be described using photochemical ageing (Parrington et al., 2013). In this paper we describe a new Eulerian method to quantify the effective physical age of emissions  $\bar{A}$ , taking into account mixing and chemical decay of emitted air masses of varying age. We found that during BORTAS-B, values of  $\bar{A}$  for airmasses intercepted over Halifax, Nova Scotia are typically 4–11 days old but occasionally as young as two days, corresponding well to analysis of concurrent ground-based observations at the Dalhousie ground station (DGS) in Halifax, NS (Gibson et al., 2013). We found our Eulerian age of emission estimates are typically a day older than those determined by the the HYSPLIT model during the 19–21 July 2011 when ground-based observations reported elevated concentrations due to biomass burning (Gibson et al., 2013; Griffin et al., 2013; Franklin et al., 2014). We attributed this difference in age to our method accounting for older air masses that are not explicitly considered by HYSPLIT. We compared the values of  $\bar{A}$  calculated using our method to the corresponding photochemical ages, using ratios of NMVOCs (Parrington et al., 2013), and found that values of  $\bar{A}$  are typically 1–5 days older. We found that the median difference between the age measures in plumes (defined as  $\text{CH}_3\text{CN} > 150$  ppt) peaks at 3–5 days, compared to a muted distribution for background CO concentrations, corresponding to a chemical retardation of 50%. We also found that in plumes  $\Delta\tau$  increased with physical age ( $r^2=0.80$ ), which was much less pronounced outwith plumes ( $r^2=0.23$ ). Based on  $\Delta\tau$  within and outwith plumes and on a strong relationship between CO and, for example, black carbon aerosol during BORTAS-B ( $r^2 > 0.70$ , Taylor et al. (2014)) we hypothesize that  $\Delta\tau$  variations provide evidence that pyrogenic aerosols slow down the plume photochemistry for many days downwind from the point of burning.

Previous work has showed using a photochemical model that the observed ozone tendency of Alaskan forest fire plumes observed over the North Atlantic during 2004 was consistent with a reduced photolysis rate of approximately 20% that could be due to aerosol loading within/above these plumes (Real et al., 2007). Our analysis of photochemical age versus  $\bar{A}$  suggests a larger retardation to the plume photochemistry. One important counter argument to our analysis of  $\Delta\tau$  that could reconcile  $\bar{A}$  and photochemical ages is that the photochemical age could have a negative bias. The method, described in detail by Parrington et al. (2013), relies on variation of NMVOC ratios that have different chemical lifetimes against oxidation by OH. The lifetime calculation is anchored by an assumption of a constant OH concentration. The BORTAS-B data analysis assumed an OH concentration of  $2 \times 10^6$  molec/cm<sup>3</sup> that was chosen to be representative of a northern midlatitude summertime OH concentration (Spivakovsky et al., 2000). Halving (doubling) the assumed OH con-

centration would increase (decrease) the photochemical age. For many flights the median value of  $\bar{A}$  is substantial higher than the photochemical age. As discussed in Yokelson et al. (2013), there are inherit limitations to using photochemistry as a proxy for the age of emissions. The photochemical age calculation also assumes a constant emission ratio of NMVOCs from the fires, and a constant background concentration. Our method also assumes constant OH concentrations, but it is much less sensitive to changes in OH concentration. All values of  $\bar{A}$  we report in our domain are well within the e-folding lifetime set by OH therefore diffusion is the dominant effect on  $\bar{A}$  for our time scale. By running our analysis with a doubled and halved lifetime of CO against OH oxidation we find that values of  $\bar{A}$  differ by less than one day.

Our method of calculating  $\bar{A}$  does not consider the size of the fire or the amount of CO that is emitted. While this will not affect the  $\bar{A}$  calculation it may complicate the interpretation of data. If, for example, old, high-CO air masses mix with young, low-CO air masses our method will assign more weight to the younger air mass and a stronger attribution to the observed CO variability. We tried to minimize this issue by using a fixed chemical decay but some residual of this issue will unavoidably remain. Other measures of age inferred from Lagrangian back trajectories, say, will suffer from similar problems and in some circumstances be more problematic if the mixing of different air masses is not considered.

Accounting for biomass burning in regional air quality budgets downwind of fires presents a number of challenges, not least due to the ability of models to predict where plume chemistry will result in net production of  $O_3$ . It is well established that this production is a function of the pyrogenic emissions (themselves a function of many environmental variables), the associated vertical mixing and transport pathways, and the photochemical environment. Using BORTAS-B we have only inferred that aerosols have slowed photochemical ageing of plumes but there is insufficient data to characterize directly how the aerosols have affected the photochemical environment within the plume as a function of time. Further studies of similar pyrogenic plumes should include a full suite of aerosol and radiation instruments in addition to gas-phase atmospheric chemistry instruments. This kind of integrative analysis will become progressively more important as we analyze more complex environments such as megacities where there is typically a mix of biogenic, anthropogenic, and pyrogenic material determining ozone photochemistry.

*Acknowledgements.* This research was supported by the Natural Environment Research Council under grant number NE/F017391/1. DF acknowledges NERC studentship NE/K500835/1. PIP acknowledges support from his Philip Leverhulme Prize and his Royal Society Wolfson Research Merit Award. DF and PIP designed the numerical calculations, performed the analysis, and wrote the paper; PIP was also the principal BORTAS mission scientist. MP was a mission scientist on BORTAS, and provided the photochemical ages and comments on the paper.

## References

- 340 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, doi:10.5194/acp-11-4039-2011, <http://www.atmos-chem-phys.net/11/4039/2011/>, 2011.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. a., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23 073, doi:10.1029/2001JD000807, <http://doi.wiley.com/10.1029/2001JD000807>, 2001.
- 345 Bian, H., Colarco, P. R., Chin, M., Chen, G., Rodriguez, J. M., Liang, Q., Blake, D., Chu, D. a., da Silva, A., Darmenov, a. S., Diskin, G., Fuelberg, H. E., Huey, G., Kondo, Y., Nielsen, J. E., Pan, X., and Wisthaler, A.: Source attributions of pollution to the Western Arctic during the NASA ARCTAS field campaign, *Atmos. Chem. Phys.*, 13, 4707–4721, doi:10.5194/acp-13-4707-2013, <http://www.atmos-chem-phys.net/13/4707/2013/>, 2013.
- 350 Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT 4 modeling system of trajectories, dispersion, and deposition, *Aust. Meteor. Mag.*, 47, 295–308, 1998.
- 355 Duncan, B. N., Logan, J. a., Bey, I., Megretskaia, I. a., Yantosca, R. M., Novelli, P. C., Jones, N. B., and Rinsland, C. P.: Global budget of CO, 1988–1997: Source estimates and validation with a global model, *J. Geophys. Res.*, 112, D22 301, doi:10.1029/2007JD008459, <http://doi.wiley.com/10.1029/2007JD008459>, 2007.
- Feng, L., Palmer, P. I., Bösch, H., and Dance, S.: Estimating surface CO<sub>2</sub> fluxes from space-borne CO<sub>2</sub> dry air mole fraction observations using an ensemble Kalman Filter, *Atmos. Chem. Phys.*, 9, 2619–2633, doi:10.5194/acp-9-2619-2009, <http://www.atmos-chem-phys.net/9/2619/2009/>, 2009.
- 360 Fisher, J. a., Jacob, D. J., Purdy, M. T., Kopacz, M., Le Sager, P., Carouge, C., Holmes, C. D., Yantosca, R. M., Batchelor, R. L., Strong, K., Diskin, G. S., Fuelberg, H. E., Holloway, J. S., Hyer, E. J., McMillan, W. W., Warner, J., Streets, D. G., Zhang, Q., Wang, Y., and Wu, S.: Source attribution and interannual variability of Arctic pollution in spring constrained by aircraft (ARCTAS, ARCPAC) and satellite (AIRS) observations of carbon monoxide, *Atmos. Chem. Phys.*, 10, 977–996, doi:10.5194/acp-10-977-2010, <http://www.atmos-chem-phys.net/10/977/2010/>, 2010.
- 365 Franklin, J. E., Drummond, J. R., Griffin, D., Pierce, J. R., Waugh, D. L., Palmer, P. I., Parrington, M., Lee, J. D., Lewis, A. C., Rickard, A. R., Taylor, J. W., Allan, J. D., Coe, H., Walker, K. A., Chisholm, L., Duck, T. J., Hopper, J. T., Blanchard, Y., Gibson, M. D., Curry, K. R., Sakamoto, K. M., Lesins, G., Dan, L., Kliever, J., and Saha, A.: A case study of aerosol depletion in a biomass burning plume over Eastern Canada during the 2011 BORTAS field experiment, *Atmospheric Chemistry and Physics Discussions*, 14, 3395–3426, doi:10.5194/acpd-14-3395-2014, <http://www.atmos-chem-phys-discuss.net/14/3395/2014/>, 2014.
- 370 Gerbig, C., Schmitgen, S., Kley, D., Volz-Thoms, A., Dewey, K., Haaks, D., and Volz-Thomas, A.: An improved fast-response vacuum-UV resonance fluorescence CO instrument, *J. Geophys. Res.*, 104, 1699–1704, doi:10.1029/1998JD100031, <http://doi.wiley.com/10.1029/1998JD100031>, 1999.
- Gibson, M. D., Pierce, J. R., Waugh, D., Kuchta, J. S., Chisholm, L., Duck, T. J., Hopper, J. T., Beauchamp, S., King, G. H., Franklin, J. E., Leaitch, W. R., Wheeler, A. J., Li, Z., Gagnon, G. A., and Palmer, P. I.:

Identifying the sources driving observed PM<sub>2.5</sub> temporal variability over Halifax, Nova Scotia, during  
380 BORTAS-B, *Atmos. Chem. Phys.*, 13, 7199–7213, 2013.

Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries,  
R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products,  
*Biogeosciences*, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, <http://www.biogeosciences.net/7/1171/2010/>,  
2010.

385 Gonzi, S., Feng, L., and Palmer, P. I.: Seasonal cycle of emissions of {CO} inferred from {MOPITT} profiles  
of {CO}: Sensitivity to pyroconvection and profile retrieval assumptions, *Geophys. Res. Lett.*, 38, 2011.

Goode, J. G., Yokelson, R. J., Ward, D. E., Susott, R. A., Babbitt, R. E., Davies, M. A., and Hao, W. M.:  
Measurements of excess O<sub>3</sub>, CO<sub>2</sub>, CO, CH<sub>4</sub>, C<sub>2</sub>H<sub>4</sub>, C<sub>2</sub>H<sub>2</sub>, HCN, NO, NH<sub>3</sub>, HCOOH, CH<sub>3</sub>COOH,  
HCHO, and CH<sub>3</sub>OH in 1997 Alaskan biomass burning plumes by airborne Fourier transform infrared  
390 spectroscopy (AFTIR), *J. Geophys. Res.*, 105, 22 147, doi:10.1029/2000JD900287, <http://doi.wiley.com/10.1029/2000JD900287>, 2000.

Griffin, D., Walker, K. a., Franklin, J. E., Parrington, M., Whaley, C., Hopper, J., Drummond, J. R., Palmer,  
P. I., Strong, K., Duck, T. J., Abboud, I., Bernath, P. F., Clerbaux, C., Coheur, P.-F., Curry, K. R., Dan, L.,  
Hyer, E., Kliever, J., Lesins, G., Maurice, M., Saha, a., Tereszchuk, K., and Weaver, D.: Investigation of  
395 CO, C<sub>2</sub>H<sub>6</sub> and aerosols in a boreal fire plume over eastern Canada during BOR-  
TAS 2011 using ground- and satellite-based observations and model simulations, *Atmos. Chem. Phys.*, 13,  
10 227–10 241, doi:10.5194/acp-13-10227-2013, <http://www.atmos-chem-phys.net/13/10227/2013/>, 2013.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terres-  
trial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos.*  
400 *Chem. Phys. Discuss.*, 6, 107–173, doi:10.5194/acpd-6-107-2006, <http://www.atmos-chem-phys-discuss.net/6/107/2006/>, 2006.

Jacob, D. J., Crawford, J. H., Maring, H., Clarke, a. D., Dibb, J. E., Emmons, L. K., Ferrare, R. a.,  
Hostetler, C. a., Russell, P. B., Singh, H. B., Thompson, a. M., Shaw, G. E., McCauley, E., Pederson,  
J. R., and Fisher, J. a.: The Arctic Research of the Composition of the Troposphere from Aircraft and  
405 Satellites (ARCTAS) mission: design, execution, and first results, *Atmos. Chem. Phys.*, 10, 5191–5212,  
doi:10.5194/acp-10-5191-2010, <http://www.atmos-chem-phys.net/10/5191/2010/>, 2010.

Jaffe, D. a. and Wigder, N. L.: Ozone production from wildfires: A critical review, *Atmos. Environ.*, 51, 1–10,  
doi:10.1016/j.atmosenv.2011.11.063, <http://linkinghub.elsevier.com/retrieve/pii/S1352231011012507>, 2012.

Jones, D. B. a., Bowman, K. W., Palmer, P. I., Worden, J. R., Jacob, D. J., Hoffman, R. N., Bey, I., and Yantosca,  
410 R. M.: Potential of observations from the Tropospheric Emission Spectrometer to constrain continental  
sources of carbon monoxide, *J. Geophys. Res. Atmos.*, 108, n/a—n/a, doi:10.1029/2003JD003702, <http://doi.wiley.com/10.1029/2003JD003702>, 2003.

Koppmann, R., von Czapiewski, K., and Reid, J. S.: A review of biomass burning emissions, part I:  
gaseous emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen contain-  
415 ing compounds, *Atmos. Chem. Phys. Discuss.*, 5, 10 455–10 516, doi:10.5194/acpd-5-10455-2005, <http://www.atmos-chem-phys-discuss.net/5/10455/2005/>, 2005.

Le Breton, M., Bacak, A., Muller, J. B. A., O'Shea, S. J., Xiao, P., Ashfold, M. N. R., Cooke, M. C., Batt,  
R., Shallcross, D. E., Oram, D. E., Forster, G., Bauguitte, S. J.-B., Palmer, P. I., Parrington, M., Lewis,



420 A. C., Lee, J. D., and Percival, C. J.: Airborne hydrogen cyanide measurements using a chemical ionisation  
mass spectrometer for the plume identification of biomass burning forest fires, *Atmos. Chem. Phys.*, 13,  
9217–9232, 2013.

Lewis, A. C., Evans, M. J., Hopkins, J. R., Punjabi, S., Read, K. A., Purvis, R. M., Andrews, S. J., Moller,  
S. J., Carpenter, L. J., Lee, J. D., Rickard, A. R., Palmer, P. I., and Parrington, M.: The influence of biomass  
burning on the global distribution of selected non-methane organic compounds, *Atmos. Chem. Phys.*, 13,  
425 851–867, 2013.

Liang, Q., Rodriguez, J. M., Douglass, A. R., Crawford, J. H., Olson, J. R., Apel, E., Bian, H., Blake, D. R.,  
Brune, W., Chin, M., Colarco, P. R., da Silva, A., Diskin, G. S., Duncan, B. N., Huey, L. G., Knapp, D. J.,  
Montzka, D. D., Nielsen, J. E., Pawson, S., Riemer, D. D., Weinheimer, A. J., and Wisthaler, A.: Reactive  
nitrogen, ozone and ozone production in the Arctic troposphere and the impact of stratosphere-troposphere  
430 exchange, *Atmospheric Chemistry and Physics*, 11, 13 181–13 199, doi:10.5194/acp-11-13181-2011, 2011.

Murphy, J. G., Oram, D. E., and Reeves, C. E.: Measurements of volatile organic compounds over West  
Africa, *Atmospheric Chemistry and Physics*, 10, 5281–5294, doi:10.5194/acp-10-5281-2010, [http://www.  
atmos-chem-phys.net/10/5281/2010/](http://www.atmos-chem-phys.net/10/5281/2010/), 2010.

Olivier, J. G. J., Bloos, J. P. J., Berdowski, J. J. M., Visschedijk, A. J. H., and Bouwman, A. F.: A 1990 global  
435 emission inventory of anthropogenic sources of carbon monoxide on 1<sup>st</sup> January developed in the framework  
of EDGAR/GEIA, *Chemosph. - Glob. Chang. Sci.*, 1, 1–17, doi:10.1016/S1465-9972(99)00019-7, [http:  
//linkinghub.elsevier.com/retrieve/pii/S1465997299000197](http://linkinghub.elsevier.com/retrieve/pii/S1465997299000197), 1999.

O’Shea, S. J., Allen, G., Gallagher, M. W., Bauguutte, S. J.-B., Illingworth, S. M., Le Breton, M., Muller, J.  
B. A., Percival, C. J., Archibald, A. T., Oram, D. E., Parrington, M., Palmer, P. I., and Lewis, A. C.: Airborne  
440 observations of trace gases over boreal Canada during BORTAS: campaign climatology, airmass analysis  
and enhancement ratios, *Atmos. Chem. Phys. Discuss.*, 13, 14 069–14 114, 2013.

Palmer, P. I., Jacob, D. J., Jones, D. B. A., Heald, C. L., Yantosca, R. M., and Logan, J. A.: Inverting for  
emissions of carbon monoxide from Asia using aircraft observations over the western Pacific, *J. Geophys.  
Res.*, 108, 8828, doi:10.1029/2003JD003397, <http://doi.wiley.com/10.1029/2003JD003397>, 2003.


445 Palmer, P. I., Suntharalingam, P., Jones, D. B. a., Jacob, D. J., Streets, D. G., Fu, Q., Vay, S. a., and Sachse,  
G. W.: Using CO<sub>2</sub>:CO correlations to improve inverse analyses of carbon fluxes, *J. Geophys. Res.*, 111,  
D12 318, doi:10.1029/2005JD006697, <http://doi.wiley.com/10.1029/2005JD006697>, 2006.

Palmer, P. I., Parrington, M., Lee, J. D., Lewis, a. C., Rickard, a. R., Bernath, P. F., Duck, T. J., Waugh, D. L.,  
Tarasick, D. W., Andrews, S., Aruffo, E., Bailey, L. J., Barrett, E., Bauguutte, S. J.-B., Curry, K. R., Di Carlo,  
450 P., Chisholm, L., Dan, L., Forster, G., Franklin, J. E., Gibson, M. D., Griffin, D., Helmig, D., Hopkins, J. R.,  
Hopper, J. T., Jenkin, M. E., Kindred, D., Kliever, J., Le Breton, M., Matthiesen, S., Maurice, M., Moller,  
S., Moore, D. P., Oram, D. E., O’Shea, S. J., Owen, R. C., Pagniello, C. M. L. S., Pawson, S., Percival, C. J.,  
Pierce, J. R., Punjabi, S., Purvis, R. M., Remedios, J. J., Rotermund, K. M., Sakamoto, K. M., da Silva,  
a. M., Strawbridge, K. B., Strong, K., Taylor, J., Trigwell, R., Tereszchuk, K. a., Walker, K. a., Weaver, D.,  
455 Whaley, C., and Young, J. C.: Quantifying the impact of BOREal forest fires on Tropospheric oxidants over  
the Atlantic using Aircraft and Satellites (BORTAS) experiment: design, execution and science overview,  
*Atmos. Chem. Phys.*, 13, 6239–6261, doi:10.5194/acp-13-6239-2013, [http://www.atmos-chem-phys.net/13/  
6239/2013/](http://www.atmos-chem-phys.net/13/6239/2013/), 2013.


- Parrington, M., Palmer, P. I., Henze, D. K., Tarasick, D. W., Hyer, E. J., Owen, R. C., Helmig, D., Clerbaux, C., Bowman, K. W., Deeter, M. N., Barratt, E. M., Coheur, P.-F., Hurtmans, D., Jiang, Z., George, M., and Worden, J. R.: The influence of boreal biomass burning emissions on the distribution of tropospheric ozone over North America and the North Atlantic during 2010, *Atmos. Chem. Phys.*, 12, 2077–2098, doi:10.5194/acp-12-2077-2012, <http://www.atmos-chem-phys.net/12/2077/2012/>, 2012.
- Parrington, M., Palmer, P. I., Lewis, a. C., Lee, J. D., Rickard, a. R., Di Carlo, P., Taylor, J. W., Hopkins, J. R., Punjabi, S., Oram, D. E., Forster, G., Aruffo, E., Moller, S. J., Bauguitte, S. J.-B., Allan, J. D., Coe, H., and Leigh, R. J.: Ozone photochemistry in boreal biomass burning plumes, *Atmos. Chem. Phys. Discuss.*, 13, 1795–1853, doi:10.5194/acpd-13-1795-2013, <http://www.atmos-chem-phys-discuss.net/13/1795/2013/>, 2013.
- Parrish, D. D., Stohl, A., Forster, C., Atlas, E. L., Blake, D. R., Goldan, P. D., Kuster, W. C., and de Gouw, J. a.: Effects of mixing on evolution of hydrocarbon ratios in the troposphere, *J. Geophys. Res.*, 112, D10S34, doi:10.1029/2006JD007583, <http://doi.wiley.com/10.1029/2006JD007583>, 2007.
- Real, E., Law, K. S., Weinzierl, B., Fiebig, M., Petzold, a., Wild, O., Methven, J., Arnold, S., Stohl, a., Huntrieser, H., Roiger, a., Schlager, H., Stewart, D., Avery, M., Sachse, G., Browell, E., Ferrare, R., and Blake, D.: Processes influencing ozone levels in Alaskan forest fire plumes during long-range transport over the North Atlantic, *J. Geophys. Res.*, 112, D10S41, doi:10.1029/2006JD007576, <http://doi.wiley.com/10.1029/2006JD007576>, 2007.
- Reid, J. S., Hyer, E. J., Prins, E. M., Westphal, D. L., Zhang, J., Wang, J., Christopher, S. a., Curtis, C. a., Schmidt, C. C., Eleuterio, D. P., Richardson, K. a., and Hoffman, J. P.: Global Monitoring and Forecasting of Biomass-Burning Smoke: Description of and Lessons From the Fire Locating and Modeling of Burning Emissions (FLAMBE) Program, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 2, 144–162, doi:10.1109/JSTARS.2009.2027443, <http://ieeexplore.ieee.org/lpdocs/epic03/wrapper.htm?arnumber=5208306>, 2009.
- Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B. A., Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A. M., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Three-dimensional climatological distribution of tropospheric OH: Update and evaluation, *Journal of Geophysical Research: Atmospheres*, 105, 8931–8980, doi:10.1029/1999JD901006, <http://dx.doi.org/10.1029/1999JD901006>, 2000.
- Taylor, J. W., Allan, J. D., Allen, G., Coe, H., Williams, P. I., Flynn, M. J., Le Breton, M., Muller, J. B. A., Percival, C. J., Oram, D., Forster, G., Lee, J. D., Rickard, A. R., and Palmer, P. I.: Size-dependent wet removal of black carbon in Canadian biomass burning plumes, *Atmospheric Chemistry and Physics Discussions*, 14, 19469–19513, doi:10.5194/acpd-14-19469-2014, <http://www.atmos-chem-phys-discuss.net/14/19469/2014/>, 2014.
- Tereszczuk, K. A., Moore, D. P., Harrison, J. J., Boone, C. D., Park, M., Remedios, J. J., Randel, W. J., and Bernath, P. F.: Observations of peroxyacetyl nitrate (PAN) in the upper troposphere by the Atmospheric Chemistry Experiment-Fourier Transform Spectrometer (ACE-FTS), *Atmos. Chem. Phys.*, 13, 5601–5613, 2013.
- Yokelson, R. J., Andreae, M. O., and Akagi, S. K.: Pitfalls with the use of enhancement ratios or normalized excess mixing ratios measured in plumes to characterize pollution sources and aging, *Atmospheric Measurement Techniques*, 6, 2155–2158, doi:10.5194/amt-6-2155-2013, 2013.

**Table 1.** Contribution of CO from geographical sources averaged over all BORTAS-B flights, lumping all other contributions <2 ppb into “Other”. Background refers to any residual CO before the beginning of the BORTAS-B period.


Tracer Source	Mean (ppb)	Std Dev (ppb)
NE USA and Canada Biomass Burning	55.4	134.6
East Asian Fossil Fuel	11.9	3.4
North East American Fossil Fuel	7.9	15.4
NW USA and Canada Biomass Burning	6.2	4.5
North West American Fossil Fuel	3.4	2.2
Mid Siberia Biomass Burning	2.0	0.7
Other	7.9	4.2
Methane & NMVOCS	45.5	12.4
Background	10.7	2.3




**Fig. 1.** Source regions for the tagged CO simulation. Regions outlined in red denote fossil fuel tagged tracers and regions outlined in green refer to biomass burning tagged tracers.




**Fig. 2.** Statistical comparison of model and observed CO from BORTAS-B. The observations have been averaged over the  $2^\circ \times 2.5^\circ$  model grid. Left panel shows the frequency distributions and the right panel show the frequency distribution of the model minus observed CO residuals. Mean and median values are shown inset of each panel.




**Fig. 3.** Relative model error in the GEOS-Chem simulation of CO during BORTAS-B as a function of altitude described by the box and whiskers approach. The red line and grey cross denotes the median and mean values, respectively.



**Fig. 4.** Mean June-August GEOS-Chem model surface CO concentrations (ppb) from the four largest sources over the northern hemisphere for 2008–2011. Individual contributions (Figure 1) are from NW North America biomass burning emissions (first row), NE North America fossil fuel (second row), East Siberia biomass burning (third row), and mid-Siberia biomass burning (bottom row).




**Fig. 5.** Same as Figure 4 but at 4 km altitude. Note the different upper limit to the colour bar.




**Fig. 6.** Box and whiskers plot showing the mean age of emissions for different altitudes (<2 km, 2–4 km, 4–6 km, and >6 km) at the longitudinal boundaries of the BORTAS-B domain (45°N–60° N, 95°W–50°W) during July 2008, 2009, 2010, 2011, and for the model sampled along the BORTAS-B flights. Within the box, the upper, middle, and lower horizontal bars denote the first quartile, the median, and the third quartile. The full range of data is shown by the whiskers.






**Fig. 7.** Age of emissions on 20th July 2011 (left) from 120°–40°W and 0–7 km, averaged over 45°–55°N; and (right) 40°–70°N and 0–7 km at 63°W, the same longitude as the Dalhousie University, Halifax, NS surface measurements (Palmer et al., 2013). Location of Dalhousie University is shown by the black arrow.



**Fig. 8.** Box and whiskers plot of the age of air observed during individual BORTAS-B aircraft flights using photochemical age using BORTAS-B data (red) and effective age  $\bar{A}$  using the GEOS-Chem model (blue). The box and whiskers plot for all flights are shown on the last two right columns. Within the box, the upper, middle, and lower horizontal bars denote the first quartile, the median, and the third quartile. The full range of data is shown by the whiskers.



**Fig. 9.** Top Panel: Frequency of effective age  $\bar{A}$  minus photochemical age (days, left axis) and CO concentration (ppb, right axis). The number of measurements  $n$  for each classification are shown inset. Bottom Panel: Scatterplot of  $\Delta\tau$  and  $\bar{A}$ . Red dots denote CO concentrations within a plume ( $\text{CH}_3\text{CN} > 150$  ppt) and the blue dots denote CO concentrations outwith a plume.