

Dear Dr. Kaiser,

Please find our revised manuscript, "On the origin of the occasional spring nitrate peak in Greenland snow" by Geng et al. In this file, we combine the "point-to-point" responses to the editor and the reviewer (**Pages 2-9**), followed by a "tracked-changes" manuscript (**Pages 10-60**).

As summarized by the editor, the reviewer has three points. The first point is regarding the sub-annual dating of the firn core. In the response below (also in the revised manuscript), we explicitly explain that due to the lack of a reliable summer layer indicator, we are unable to perform sub-annual dating of the firn core. Therefore, in the revised manuscript we have replaced the term "spring nitrate peak" with "years with double nitrate peaks" for the firn core.

The second point is regarding the role of pollution transport. In the analysis of the snowpit for the spring of 2005, we excluded pollution transport as a source of the spring nitrate peak using a combination of the negative NAO phase and the lack of a simultaneous sulfate peak. In the revised manuscript, we also compare nitrate and sulfate records in the firn core, and find that one (usually the secondary, or the smaller) of the double nitrate peaks in most years (14 out of 19 years) coincides with a sulfate peak. This suggests that long-range transport of nitrate is likely the source of the additional nitrate peak in these years. **According to this new result, , we have revised our conclusion as "the combination of firn core and snowpit results suggest that there are two mechanisms leading to the observed double nitrate peaks, 1) transport of nitrate from polluted areas and 2) enhanced local photochemical production of nitrate"**. This is significantly different from the conclusion in previous manuscripts, and we thank the reviewer for pointing this out.

Finally, the reviewer argued that "there is not a whole lot of sunlight at Summit in spring" and "1.3 times increase in snow nitrate photolysis" hints little increase in OH production. Using the TUV model, we have calculated that OH production may be increased by 90% due to enhanced surface ozone photolysis during the spring of 2005. Together with photolysis of snow-sourced  $\text{CH}_2\text{O}$ ,  $\text{H}_2\text{O}_2$ , HONO, and the effect of snow-sourced  $\text{NO}_x$  on OH production, it is plausible that OH production is enhanced by ~200% in the spring of 2005. As discussed in the manuscript, this alone is sufficient to cause the observed enhancement in nitrate production in the spring of 2005..

We are grateful to the editor for the helpful suggestions, and look forward to hearing from you at your earliest convenience.

Sincerely,  
Lei Geng, Ph.D  
Department of Atmospheric Sciences  
University of Washington  
ATG 408, Box 351640  
Seattle, WA 98195  
[leigeng@uw.edu](mailto:leigeng@uw.edu)

## Point-to-point response to editor and the reviewer:

We thank the editor and the reviewer very much for the suggestions and comments. Below we provide point-to-point responses to the comments. The suggestions/comments are in italics, followed by the response in normal font.

### Editor's Comment:

*the reviewer has suggested that the manuscript be rejected for three main reasons*

- *dating*
- *possibility of long-range transport during low NAO phases*
- *the proposed production mechanism*

*1) To address the first point, it would perhaps help if you showed a figure similar to Figure 2 for the years covered by satellite ozone column observations (1979 onwards), in which you identify the purported winter and summer months and calendar years as well as all nitrate peaks and classify them according to season. You do this to some extent in the text (p. 30), but these arguments are difficult to rationalise without further analysis of the data, which leaves unacceptable margins for interpretation.*

**Response:** We thank the editor very much for this suggestion. However, we are unable to perform sub-annual dating of the firn core in this study, mainly due to the lack of a reliable summer layer indicator. Previous studies use H<sub>2</sub>O<sub>2</sub> (Burkhart et al., 2006) or δ<sup>18</sup>O(H<sub>2</sub>O) (Whitlow et al., 1992) peak as a summer layer indicator. These parameters are independent of chemical ion (e.g., NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>) concentrations. However, we didn't measure such parameters in the snowpit or firn core, and don't have samples left for additional measurements. For the snowpit samples, we use Cl<sup>-</sup>/Na<sup>+</sup> peak as a summer layer indicator. However, Cl<sup>-</sup>/Na<sup>+</sup> is not independent of nitrate and sulfate concentrations. Especially, in the period of high atmospheric acidity (i.e., the industrial era when anthropogenic sulfate and nitrate increased the acidity of the atmosphere and snow (Geng et al., 2014)), the ability of Cl<sup>-</sup>/Na<sup>+</sup> peak as a summer layer indicator is reduced **for reasons detailed in our response below**. This applies to the firn core but not to the snowpit, because the snowpit covers 2004-2007 when sulfate concentrations are as low as they were prior to 1900 (Geng et al., 2014), minimizing its impact on the timing of the Cl<sup>-</sup>/Na<sup>+</sup> peak.

To avoid confusion on the use of "spring nitrate peak" in the firn core, in the revised manuscript, we have replaced the term "years with a spring nitrate peak" with "years with double nitrate peaks".

*2) To address the second point, you might be able draw on the work of Kramer et al. (2014) or extend it, and maybe include the NAO phase in Figure 3.*

**Response:** Please refer to our response to the reviewer regarding this point.

3) The reviewer also suggests more careful numerical photochemical calculations to address the third point and to check whether the suggested enhanced UV flux could account quantitatively for the nitrate peak.

**Response:** Please refer to our response to the reviewer regarding this point.

## Reviewer's comments:

*Geng et al. revision 2. The authors claim that this draft of the manuscript addresses the two main issues they feel I raised in regard to the previous version; dismissing long range transport of pollution as viable source of extra (outside of summer) nitrate peaks in the snow and firn records too quickly, and concern about dating of the pit and core. In my earlier review I also noted that I agreed with most of the comments raised by reviewer 2 on the initial submission, and that I felt these needed to be addressed in the manuscript (not just in the response to reviews).*

**Comment 1)** *This draft does directly address dating of the snowpit, and establishes the timing of the extra nitrate as a late winter/spring event. In fact, the authors claim that the peak is in snow they think accumulated in Feb (which is problematic for their hypothesis, more below). However, the authors choose not to attempt to date the firn core records, and continue to assume that any year with 2 nitrate peaks must have one in the spring and the other in the summer. They justify this decision on the basis of previous work by Cole-Dai et al who estimated that subannual dating of Summit firn cores was imprecise by up to +/- 4 months. I fully agree that it is not wise to assert that a peak in Na or Cl/Na can be dated to a specific week, or even month, but I am pretty certain that the winter peak in Na shows up in snow that accumulated in DJFM while the summer peak in Cl/Na occurs JJA. A "springtime" nitrate peak might show up in MAMJ in any given year, but it has to be between the winter and summer snow layers. As noted by reviewer 2, and myself, several of the extra nitrate peaks shown in Fig 2 (just 2 short sections of the firn core) fail this test and thus can not be springtime events. As a result, the comparison between O3 column variations and years with so-called springtime nitrate peaks (Fig 3 and discussion thereof) is not informative.*

**Response:** We agree partly with the reviewer on the statement that " Na shows up in snow that accumulated in DJFM while the summer peak in Cl/Na occurs JJA". The Cl/Na<sup>+</sup> peak is a good summer layer indicator in the preindustrial period when atmospheric acidity reaches the highest level in summer (due to the summer peaks of sulfate and nitrate). However, in the industrial era, the timing of the Cl/Na<sup>+</sup> peak differs due to seasonal changes in atmospheric acidity before and after the industrial revolution. Here we discuss why the Cl/Na<sup>+</sup> peak is affected by atmospheric


acidity. The  $\text{Cl}^-/\text{Na}^+$  calculation can be expressed by the following equation:

$$\text{Cl}^-/\text{Na}^+ = \frac{\text{ssCl}^- + \text{Cl}^-_{\text{ex}}}{\text{Na}^+} = \frac{\text{ssCl}^-}{\text{Na}^+} + \frac{\text{Cl}^-_{\text{ex}}}{\text{Na}^+} \quad \text{Equation 1}$$

where  $\text{ssCl}^-$  represents sea-salt chloride, and  $\text{Cl}^-_{\text{ex}}$  is excess chloride originating from acid-sea salt displacement (Laskin et al., 2012; Legrand and Delmas, 1988) and/or potential anthropogenic emissions in the Northern Hemisphere. Assuming  $\text{Na}^+$  in Summit snow is solely from sea salt, the first term in E1 is then equal to 1.79 (the mass ratio of  $\text{Cl}^-/\text{Na}^+$  in standard sea water). Therefore, variability in the value of  $\text{Cl}^-/\text{Na}^+$  in snow/ice cores is determined by the second term of E1,  $\text{Cl}^-_{\text{ex}}/\text{Na}^+$ . In the preindustrial period,  $\text{Na}^+$  reaches the lowest level in summer when the acidity is the highest (due to summer peaks of nitrate and sulfate), so that  $\text{Cl}^-/\text{Na}^+$  peak almost precisely falls in summer. However, in the industrial era, the sulfate (and acidity) peak becomes irregular, with a major peak in winter. This leads to the shift of annual  $\text{Cl}^-/\text{Na}^+$  peak toward the main sulfate (winter) peak. As shown in **Figure S2**, the affect of the winter sulfate peak (acidity effect) on the  $\text{Cl}^-/\text{Na}^+$  peak is clearly indicated in the industrial era (indicated as green arrows).

There are additional reasons why  $\text{Cl}^-/\text{Na}^+$  peak is not a good summer indicator, especially in the industrial period with relatively high atmospheric and snow acidity.  $\text{Cl}^-$  is subject to post-depositional process which is influenced by many factors including snow acidity (Röthlisberger et al., 2003). This process could induce further uncertainties when using  $\text{Cl}^-/\text{Na}^+$  peak to identify the summer snow layer.

Due to the above reasons, we don't attempt to sub-annually date the firn core (we have added this discussion in the revised manuscript) in this study. We understand the concern the reviewer had on the use of "spring nitrate peak" in the firn core. In the revised manuscript, we no longer use the term "spring nitrate peak" for the firn core, instead we use "years with double nitrate peaks".



**Figure S2.** Concentrations of  $\text{NO}_3^-$ ,  $\text{nss-SO}_4^{2-}$  and  $\text{Na}^+$ , and  $\text{Cl}^-/\text{Na}^+$  in the firn core. Top: the industrial period (depth interval of 10.75 to 12.5 m, covering the years of 1984 to 1987); Bottom: the preindustrial period (depth interval of 47.75 to 49.5 m, covering the years of 1879 to 1884)

**Comment 2):** the Authors dismiss the possibility that the nitrate peak in Feb/Mar 2005 resulted from long-range transport primarily due to the phase of the NAO, suggesting that meridional transport into the Arctic requires strong positive NAO index. It may be true that on average northward transport of mid latitude pollution is enhanced when the NAO is in positive phase, but advection of pollution plumes to Summit is a common event during winter and early spring. Kramer et al. (2004) identified 21 such events during DJFM in both 2008/09 and 2009/10 (42 winter/early spring transport events in just two years) using Flexpart and measurements of  $\text{NO}_x$ ,  $\text{NO}_y$ , PAN and NMHC in the air just above the snow. Similarly, Duderstadt et al., (2014) show numerous peaks of nitrate in the surface snow at Summit during this same season in 2000/01 and 2001/02, and indicate that the NCAR WACCM model does a reasonable job of simulating these via tropospheric transport of pollution from the mid latitudes. It should be noted that the NAO index was negative in 3 out of these 4 winters. (As an aside, I also point out that Kramer et al. (2014) clearly show that PAN is the dominant component of the  $\text{NO}_y$  that reaches Summit, accounting for a low of 50% in mid summer and up to 80% in April.)

**Response:** We used the NAO record to assess the inter-annual variability of meridional transport over the years covered by the snowpit, and showed that in 2005 winter/spring the transport is not particularly strong compared with other years without a spring nitrate peak. However, even in years with weak meridional transport (usually in the negative NAO phase), pollution transport events occur and may result in a nitrate peak. Thus, we also **compared the sulfate and nitrate data in the snowpit. In the snow layer of the 2005 spring nitrate peak, there is no increase in sulfate concentrations (Figure 1), suggesting that enhanced pollution transport is unlikely at this time.** These have been stated in the relevant part of the previous manuscripts (Section 4.1.2, Page 20).

In the revised manuscript, we repeat the comparison of the timing of the sulfate and nitrate double peaks in the firn core. We find that indeed in 14 out of 19 years, the additional nitrate peak is accompanied by a sulfate peak, suggesting that the additional nitrate peak in these years may arise from pollution transport. Therefore, we have revised Figure 3, indicating which years the double nitrate peaks coincide with a sulfate peak. We also have revised our conclusion and stated that " The results from the snowpit and firn core suggest that there are probably two mechanisms leading to the observed double nitrate peaks, 1) transport of nitrate from polluted areas and 2) enhanced local photochemical production of nitrate. We note that both of these mechanisms are related to pollution transport, as the additional nitrate from either direct transport or enhanced local photochemistry requires enhanced nitrogen sources from anthropogenic emissions. " We thank the reviewer for pointing this out.

***Comment 3):** Regarding the hypothesis that the spring 2005 nitrate is locally produced (recycled) from snow sourced NO<sub>x</sub> and OH, the Feb/Mar timing of this peak is kind of problematic. There is not a whole lot of sunlight at Summit at that time of year, for example the midday measured J NO<sub>3</sub><sup>-</sup> on 25 Mar 2004 was just  $0.6 \times 10^{-8}$ , and both NO and OH were close to detection limits in the air 1 m above the snow. In early Feb the sun is still below the horizon for more than 20 hours each day (sunrise approx. 27 Jan). Therefore it is hard to imagine that O<sub>3</sub> depletion could enhance the UV flux enough to stimulate significant release of NO<sub>x</sub> and OH for much of Feb and Mar. Of course, the solar elevation does increase quickly so far north (by 13 Apr 04 the midday J NO<sub>3</sub><sup>-</sup> was almost 4 times higher than on 25 Mar) but relaxation of the polar vortex and replenishment of the O<sub>3</sub> column is usually pretty well advanced by April. The TUV calculations presented for PAN and nitrate photolysis should have been a hint that photolysis of nitrate (and also hydrogen peroxide as additional source of OH) in snow at Summit in Feb would be expected to be quite weak (1.3 times a very small number is still very small). Even if I am mistaken and there are enough photons hitting the snow in March to recycle nitrate, why should it stop and preserve a spring peak? The actinic flux continues to increase to much higher levels through April, May and much of June, so why would locally formed nitrate deposited in Feb/Mar not continue to photolyze and get redeposited at the new surface, eventually merging with the usual summer peak? Fibiger et al. (2013) suggest that very little of*

*the initially deposited nitrate ever gets photolyzed, such that the recycled fraction is also a very minor component of preserved nitrate.*

**Response:** We think the reviewer misunderstood our hypothesis. We hypothesized that the spring 2005 nitrate peak is from enhanced local photochemistry, but we did not state it is entirely from "locally (recycled) from **snow-sourced** NO<sub>x</sub> and OH". We re-state our hypothesis here: under the situation of sufficient local NO<sub>x</sub> at Summit in the industrial era (probably due to winter accumulated NO<sub>x</sub> and its precursors), elevated local oxidant concentrations will increase local nitrate production.

By using the TUV model, we then calculated the enhancement in actinic flux in the spring of 2005 relative to other years with normal column ozone density, as well as the associated increases in photolysis of surface ozone and snow nitrate. We calculated that snow nitrate photolysis is enhanced by ~30% due to the enhanced actinic flux. We stated that this could contribute to the spring nitrate peak in 2005 by up to 30%, if snow nitrate photolysis is the dominant NO<sub>x</sub> source at Summit.

Based on the size of the spring 2005 nitrate peak, we calculated that a ~200% increase in OH production is required to account for the additional nitrate in the spring of 2005. This assumes sufficient local NO<sub>x</sub> due to accumulation of NO<sub>y</sub> species (as well as NO<sub>x</sub>) in the polar winter through a combination of pollution transport from the mid-latitudes and lack of sunlight. Using the TUV model, we calculated that under the situation of 2005 springtime column ozone density, elevated surface UV radiation can result in a ~90% increase in OH production due to enhanced surface ozone photolysis. This is about half of the desired amount. Further, we discussed that in addition to ozone photolysis, in snow covered regions, OH could be produced through photolysis of snow sourced CH<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, and HONO. Previous work suggested that the photolysis of these species in/from snow is able to produce more OH than ozone photolysis (Yang et al., 2002). Although these species are more sensitive to UV-A radiation, the production of HONO and CH<sub>2</sub>O from the snowpack is sensitive to UV-B radiation (Thomas et al., 2012). In addition, snow sourced NO will shift the balance between HO<sub>2</sub> and OH, leading to as much OH production in the boundary layer as that from ozone photolysis at Summit (Thomas et al., 2012). Bromine chemistry in the snowpack (sensitive to UV-B radiation) is also suggested to be able to increase boundary OH abundance by 10-18% (Thomas et al., 2012). Whether or not snow photochemistry can account for the additional OH enhancement will depend on their relative importance for OH production in the Greenland boundary layer. Model studies suggested that snow photochemistry at Summit can lead to OH production up to 2-3 times more than ozone photolysis (Yang et al., 2002; Thomas et al., 2012). We thus concluded that a 200% increase in OH production in 2005 spring is plausible.

The reviewer critiqued that " *The TUV calculations presented for PAN and nitrate photolysis should have been a hint that photolysis of nitrate (and also hydrogen peroxide as additional*

*source of OH) in snow at Summit in Feb would be expected to be quite weak (1.3 times a very small number is still very small)". However, a 30% increase in snow nitrate photolysis under the situation of low column zone density in 2005 spring does not mean photolysis of other species is also enhanced by only 30%. The peak wavelength of photolysis of different species is different. For example, the peak wavelength of snow nitrate photolysis is ~320 nm (Frey et al., 2009), while that of ozone photolysis is ~310 nm as inferred from the spectral actinic flux in Arctic spring and the respective spectral absorptivity of ozone at the earth surface. The TUV model calculations indicate that the degrees of enhancement in different bins of actinic flux are not equivalent. In particular, the largest enhancement occurs in the bin of 305-315 nm (by ~100%), while in the bin of 315-325nm the enhancement is less than 50%. This is consistent with the difference between the enhancement in ozone photolysis (by ~90%) and snow nitrate photolysis (~30%).*

Finally, we note that it is true that in 2005 spring the surface UV radiation is still low compared to the summer levels even with particular low column ozone density. But it is stronger than in other springs (e.g., ~100% increase in 305-315 nm). Boundary photochemistry in this spring is enhanced, with amplified OH production through the combined effects of ozone photolysis and snowpack photochemistry. More nitrate is produced, and then a nitrate peak in snow is formed and preserved. We note post-depositional processing of snow nitrate is active at Summit, but is small and not able to inhibit the preservation of seasonal cycles in nitrate concentrations and its isotopic composition, as suggested by the data of this study and others (Hastings et al., 2004). It is possible for snow photochemistry to have a minimal impact on the preservation of snow nitrate while at the same time have a significant impact on the chemistry of the overlying boundary layer. For example, in a 1-D box model, Thomas et al. (2012; 2011) set only 6% of total snow nitrate is able to be photolyzed, and this much photolysis is able to account for the observed NO<sub>x</sub> concentrations as well as double OH concentrations above the snowpack.

**Comment 4):** *As noted, reviewer 2 raises additional concerns with this hypothesis that I still feel have not been adequately addressed.*

**Response:** We would like specific comments on which parts the reviewer thought we didn't address. Otherwise, we find it is difficult to respond.

**Reference:**

Burkhart, J. F., Bales, R. C., McConnell, J. R., and Hutterli, M. A.: Influence of North Atlantic Oscillation on anthropogenic transport recorded in northwest Greenland ice cores, *J. Geophys. Res.*, 111, D22309, 10.1029/2005jd006771, 2006.

Frey, M. M., Savarino, J., Morin, S., Erbland, J., and Martins, J. M. F.: Photolysis imprint in the nitrate stable isotope signal in snow and atmosphere of East Antarctica and implications for reactive nitrogen cycling, *Atmos. Chem. Phys.*, 9, 8681-8696, 10.5194/acp-9-8681-2009, 2009.



Geng, L., Alexander, B., Cole-Dai, J., Steig, E. J., Savarino, J., Sofen, E. D., and Schauer, A. J.: Nitrogen isotopes in ice core nitrate linked to anthropogenic atmospheric acidity change, *Proceedings of the National Academy of Sciences*, 111, 5808-5812, 10.1073/pnas.1319441111, 2014.

Hastings, M. G., Steig, E. J., and Sigman, D. M.: Seasonal variations in N and O isotopes of nitrate in snow at Summit, Greenland: Implications for the study of nitrate in snow and ice cores, *J. Geophys. Res.*, 109, D20306, 10.1029/2004jd004991, 2004.

Laskin, A., Moffet, R. C., Gilles, M. K., Fast, J. D., Zaveri, R. A., Wang, B., Nigge, P., and Shutthanandan, J.: Tropospheric chemistry of internally mixed sea salt and organic particles: Surprising reactivity of NaCl with weak organic acids, *Journal of Geophysical Research: Atmospheres*, 117, D15302, 10.1029/2012jd017743, 2012.

Legrand, M. R., and Delmas, R. J.: Formation of HCl in the Antarctic atmosphere, *Journal of Geophysical Research: Atmospheres*, 93, 7153-7168, 10.1029/JD093iD06p07153, 1988.

Röthlisberger, R., Mulvaney, R., Wolff, E. W., Hutterli, M. A., Bigler, M., de Angelis, M., Hansson, M. E., Steffensen, J. P., and Udisti, R.: Limited dechlorination of sea-salt aerosols during the last glacial period: Evidence from the European Project for Ice Coring in Antarctica (EPICA) Dome C ice core, *Journal of Geophysical Research: Atmospheres*, 108, 4526, 10.1029/2003jd003604, 2003.

Thomas, J. L., Stutz, J., Lefer, B., Huey, L. G., Toyota, K., Dibb, J. E., and von Glasow, R.: Modeling chemistry in and above snow at Summit, Greenland – Part 1: Model description and results, *Atmos. Chem. Phys.*, 11, 4899-4914, 10.5194/acp-11-4899-2011, 2011.

Thomas, J. L., Dibb, J. E., Huey, L. G., Liao, J., Tanner, D., Lefer, B., von Glasow, R., and Stutz, J.: Modeling chemistry in and above snow at Summit, Greenland - Part 2: Impact of snowpack chemistry on the oxidation capacity of the boundary layer, *Atmos. Chem. Phys.*, 12, 6537-6554, 10.5194/acp-12-6537-2012, 2012.

Whitlow, S., Mayewski, P. A., and Dibb, J. E.: A Comparison of Major Chemical-Species Seasonal Concentration and Accumulation at the South-Pole and Summit, Greenland, *Atmos. Environ.*, 26, 2045-2054, 10.1016/0960-1686(92)90089-4, 1992.

Yang, J., Honrath, R. E., Peterson, M. C., Dibb, J. E., Sumner, A. L., Shepson, P. B., Frey, M., Jacobi, H. W., Swanson, A., and Blake, N.: Impacts of snowpack emissions on deduced levels of OH and peroxy radicals at Summit, Greenland, *Atmos. Environ.*, 36, 2523-2534, 10.1016/S1352-2310(02)00128-0, 2002.

1           **On the origin of the occasional spring nitrate peak in Greenland snow**

2   L. Geng<sup>1,\*</sup>, J. Cole-Dai<sup>1</sup>, B. Alexander<sup>2</sup>, J. Erbland<sup>3,4</sup>, J. Savarino<sup>3,4</sup>, A. J. Schauer<sup>5</sup>, E.  
3   J. Steig<sup>5</sup>, P. Lin<sup>2,\*\*</sup>, Q. Fu<sup>2</sup>, M. C. Zatzko<sup>2</sup>

4   <sup>1</sup> Department of Chemistry & Biochemistry, South Dakota State University,  
5   Brookings, SD, USA

6   <sup>2</sup> Department of Atmospheric Sciences, University of Washington, Seattle, WA, USA

7   <sup>3</sup> CNRS, LGGE (UMR5183), F-38041 Grenoble, France

8   <sup>4</sup> Univ. Grenoble Alpes, LGGE (UMR5183), F-38041 Grenoble, France.

9   <sup>5</sup> Department of Earth and Space Sciences, University of Washington, Seattle, WA,  
10   USA

11   \* Now at Department of Atmospheric Sciences, University of Washington, Seattle,  
12   WA, USA

13   \*\* Now at Program in Atmospheric and Oceanic Sciences / GFDL, Princeton  
14   University, Princeton, NJ, USA

15

16

17

18

19

## Abstract



Ice-core nitrate concentrations peak in the summer in both Greenland and Antarctica. Two nitrate concentration peaks in one annual layer have been ~~was~~ observed in some years in ice cores in Greenland from samples dating post-1900, with the additional nitrate peak occurring in the spring. The origin of the spring nitrate peak was hypothesized to be pollution transport from the mid-latitudes in the industrial era. We perform a case study on the origin of a spring nitrate peak in 2005 measured from a snowpit at Summit, Greenland covering three years of snow accumulation. The effect of long range transport of nitrate on this spring peak is excluded by using sulfate as a pollution tracer. The isotopic composition of nitrate ( $\delta^{15}\text{N}$ ,  $\delta^{18}\text{O}$  and  $\Delta^{17}\text{O}$ ) ~~was also measured. Isotopic data~~ combined with photochemical calculations suggest that the occurrence ~~presence~~ of this spring peak is linked to a significantly weakened stratospheric ozone ( $\text{O}_3$ ) layer. The weakened  $\text{O}_3$  layer resulted in elevated UV-B (Ultraviolet B) radiation on the snow surface, where the production of OH and  $\text{NO}_x$  from the photolysis of their precursors was ~~were~~ enhanced. Elevated ~~enhanced~~  $\text{NO}_x$  and OH concentrations resulted in enhanced ~~more~~ nitrate production mainly through ~~by~~ the  $\text{NO}_x + \text{OH}$  formation pathway, as indicated by decreases in  $\delta^{18}\text{O}$  and  $\Delta^{17}\text{O}$  of nitrate associated with the spring peak. We further examined the nitrate concentration record from a shallow ice core covering the period from 1772 to 2006 and ~~and compared this record to satellite observations of~~ springtime  $\text{O}_3$  column density data from 1979 to 2006. We found 19 years with double ~~two~~ nitrate peaks after the 1950s. Out of these 19 years, 14 of the secondary nitrate peaks are accompanied by sulfate peaks, suggesting long-range transport of nitrate as their source. In the other five years ~~After 1979, all years with two nitrate~~

1 ~~peaks are also years with low~~ springtime O<sub>3</sub> column density ~~near or below the 1979-~~  
2 ~~2006 average is observed, suggesting enhanced local production of nitrate as their~~  
3 ~~source. The results suggest that, in addition to direct transport of nitrate from polluted~~  
4 ~~regions, enhanced local photochemistry can also lead to a spring nitrate peak. The~~  
5 ~~enhanced local photochemistry is probably associated with~~ We thus hypothesize that  
6 ~~the presence of the spring nitrate peak is largely associated with and may be~~  
7 ~~determined by~~ the interannual variability of O<sub>3</sub> column density in the Arctic which  
8 leads to elevated surface UV radiation in some years. In this scenario, enhanced  
9 photochemistry causes increased local nitrate production, under the condition of  
10 elevated local NO<sub>x</sub> abundance in the industrial era. at Summit after the 1950s  
11 ~~resulting from transport of anthropogenic NO<sub>x</sub> precursors, though direct transport of~~  
12 ~~nitrate and other factor(s) may be important in some years. Isotopic data covering~~  
13 ~~additional years of low O<sub>3</sub> column density are needed to further examine this~~  
14 ~~hypothesis.~~



## 15 **1. Introduction**

16 Knowledge of the abundance and variability of reactive nitrogen oxides (NO<sub>x</sub> =  
17 NO + NO<sub>2</sub>) is valuable because of the critical role that NO<sub>x</sub> plays in determining the  
18 oxidative capacity of the atmosphere. The oxidative capacity of the atmosphere is  
19 determined by the tropospheric abundance of hydrogen oxide radicals (HO<sub>x</sub> = OH +  
20 HO<sub>2</sub>) and O<sub>3</sub> and largely controls the residence times of pollutants (e.g., CO) and  
21 greenhouse gases (e.g., CH<sub>4</sub>). NO<sub>x</sub> is emitted from a variety of sources including  
22 fossil fuel combustion, biomass burning, soil emissions, and lightning (Logan, 1983).  
23 In the atmosphere, NO<sub>x</sub> cycle rapidly between NO and NO<sub>2</sub> according to:





3 The main sink of  $\text{NO}_x$  is thought to be atmospheric oxidation to nitrate which  
4 distributes between the gas- and particulate phases. In general, the formation of nitrate  
5 involves reactions of  $\text{NO}_x$  with OH and  $\text{O}_3$ :



11 Reactive halogen species (e.g., BrO) may also play a role in  $\text{NO}_x$  cycling and  
12 nitrate formation, especially in the Arctic (Morin et al., 2007). R4 mainly occurs  
13 during daytime, as the result of diurnal variation in OH concentrations. The  
14 conversion to nitrate via  $\text{O}_3$  (R5-R8) is negligible during the day, since the  $\text{NO}_3$   
15 radical is rapidly photolyzed back to  $\text{NO}_x$  in sunlight. Globally, oxidation of  $\text{NO}_x$  by  
16 OH (R4) is thought to be the dominant nitrate formation pathway on an annual basis  
17 (Alexander et al., 2009). Once formed, nitrate is removed from the atmosphere mainly  
18 through wet and dry deposition to the surface. Efforts have been made to use ice core  
19 nitrate records to assess information about past changes in the global  $\text{NO}_x$   
20 environment (Fischer et al., 1998; Mayewski et al., 1990), and potentially about the  
21 past variations in the atmospheric oxidation capacity (Alexander et al., 2004).  
22 However, nitrate concentration in polar snow is also influenced by variations in snow  
23 accumulation rate and by post-depositional processes such as evaporation and

1 photolysis (Röthlisberger et al., 2002) of nitrate in near-surface snow. Post-  
2 depositional processing of snowpack nitrate has hampered the interpretation of ice  
3 core nitrate records in terms of assessing past atmospheric NO<sub>x</sub> variability.

4 Examination of detailed chronological nitrate records in snowpits and ice cores  
5 may provide clues on the primary factor(s) controlling nitrate concentration and its  
6 temporal variability in snow. For example, a clear seasonal pattern in nitrate  
7 concentration may be linked to a source with an annual cycle and/or modulated by  
8 seasonally varying atmospheric conditions. Many previous studies of snow samples  
9 and ice cores from central Greenland have found that nitrate concentration reaches a  
10 maximum in summer snow and a minimum in winter snow (Davidson et al., 1989;  
11 Finkel and Langway, 1986; Whitlow et al., 1992). This seasonal variation in nitrate  
12 concentration in general is due to the active photochemical reactions during polar  
13 summer (Yang et al., 1995), when peroxyacetyl nitrate (PAN), snowpack nitrate and  
14 other NO<sub>x</sub> reservoir species act as local NO<sub>x</sub> sources through recycling reactions  
15 (Moxim et al., 1996; Thomas et al., 2012). A few investigators (Burkhart et al., 2006;  
16 Yang et al., 1995; Whitlow et al., 1992) noticed that a single nitrate peak appears  
17 annually in summer snow/ice layer deposited prior to 1900, whereas two nitrate peaks  
18 are seen in some post-1900 annual layers. Although it seems the timing of the  
19 additional nitrate peak differs as observed in ice cores from different locations  
20 (Burkhart et al., 2006), Yang et al. (1995) stated that at Summit the additional peak  
21 occurs in late winter/early spring, which is consistent with the dating in Burkhart et al.  
22 (2006).

23 The occurrence of this occasional spring nitrate peak has been hypothesized to be  
24 associated with pollution transport from the mid-latitudes (Burkhart et al., 2006; Yang

1 et al., 1995), though this hypothesis and the occasional nature of the spring nitrate  
2 peak were not carefully examined. Anthropogenic NO<sub>x</sub> emissions from fossil fuel  
3 combustion have increased the concentration of atmospheric NO<sub>x</sub> and other nitrogen  
4 species (e.g., HNO<sub>3</sub>) in the northern hemisphere since 1900, especially in the last 50  
5 years (Galloway et al., 2003). A corresponding increase in nitrate concentrations in  
6 Greenland ice cores has been documented (Mayewski et al., 1990). Direct transport of  
7 NO<sub>x</sub> from the mid-latitudes to the Arctic is unlikely given the short lifetime of NO<sub>x</sub>  
8 (1-3 days (Levy et al., 1999)). Alternatively, long range transport of nitrate and NO<sub>x</sub>  
9 precursors (e.g., PAN) from anthropogenic NO<sub>x</sub> source regions could lead to increases  
10 in Greenland snow nitrate concentrations. However, the relative importance of nitrate  
11 and NO<sub>x</sub> precursors to Greenland snow nitrate is unclear. An earlier study suggested a  
12 significant contribution from PAN (e.g., (Ford et al., 2002)), whereas a recent study  
13 (Geng et al., 2014) suggests that transport to Greenland occurs mainly in the form of  
14 gaseous HNO<sub>3</sub>. In addition to PAN decomposition as a potential local source of NO<sub>x</sub>,  
15 the photolysis of nitrate in snowpack produces NO<sub>x</sub>, which is quickly transported to  
16 the overlying atmosphere by diffusion and wind pumping (Zatko et al., 2013). This is  
17 consistent with the elevated NO<sub>x</sub> concentrations observed in the ice sheets during  
18 polar summers (Honrath et al., 1999). In particular, a model study (Thomas et al.,  
19 2012) suggested that at Summit, NO<sub>x</sub> produced from snow nitrate photolysis in  
20 summer can account for all of the observed NO<sub>x</sub> concentrations in the overlying  
21 atmosphere. In addition, Kramer et al. (2014) implied that, after polar sunrise,  
22 atmospheric NO<sub>y</sub> (odd nitrogen species other than PAN and NO<sub>x</sub>) can release NO<sub>x</sub>  
23 through photochemistry. In summary, there are two ways that pollution transport can  
24 influence the springtime nitrate budget at Summit, 1) direct transport of nitrate, and 2)

1 transport of NO<sub>x</sub> precursors. In order to cause a spring nitrate peak, the latter requires  
2 local conditions favoring NO<sub>x</sub> release from its precursors and its subsequent  
3 conversion to nitrate. Elevated local NO<sub>x</sub> concentrations, via enhanced concentrations  
4 and/or decomposition rates of NO<sub>x</sub> precursors, could also lead to a nitrate  
5 concentration peak in surface snow, in addition to enhanced pollution transport.

6 The isotopic composition of nitrate can provide valuable information not  
7 available from concentration measurement alone, for example, regarding the  
8 pathways of NO<sub>x</sub> conversion to nitrate in the atmosphere (Michalski et al., 2003).  
9 Stable isotope ratios in nitrate are expressed as  $\delta^{15}\text{N}$ ,  $\delta^{18}\text{O}$  and  $\Delta^{17}\text{O}$ , where  $\Delta^{17}\text{O} =$   
10  $\delta^{17}\text{O} - 0.52 \times \delta^{18}\text{O}$  and  $\delta = R_{\text{sample}}/R_{\text{reference}} - 1$  with R denoting the  $^{15}\text{N}/^{14}\text{N}$ ,  $^{18}\text{O}/^{16}\text{O}$   
11 and  $^{17}\text{O}/^{16}\text{O}$  isotope ratios. The references are N<sub>2</sub>-AIR and VSMOW for N and O,  
12 respectively.  $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  of nitrate (hereafter denoted as  $\delta^{15}\text{N}(\text{NO}_3^-)$  and  
13  $\Delta^{17}\text{O}(\text{NO}_3^-)$ , respectively) have been used to investigate the origin and fate of NO<sub>x</sub> in  
14 the Arctic troposphere (Morin et al., 2008). In particular,  $\Delta^{17}\text{O}(\text{NO}_3^-)$  is related to  
15 oxidizing conditions and nitrate formation pathways (R4 vs. R5-R8) in the  
16 atmosphere (Alexander et al., 2004; Alexander et al., 2009; Kunasek et al., 2008;  
17 Michalski et al., 2003; Morin et al., 2011). For example, Michalski et al. (2003) found  
18 that nitrate formed during nighttime (i.e., via R5-R8) has higher  $\Delta^{17}\text{O}$  than nitrate  
19 formed during daytime (via R4) because of the high  $\Delta^{17}\text{O}$  of the dominant nighttime  
20 oxidant O<sub>3</sub> (Johnson et al., 2000; Lyons, 2001) which is transferred to nitrate. The  
21  $\Delta^{17}\text{O}(\text{NO}_3^-)$  in Greenland snow also reflects the seasonality of nitrate production, with  
22 O<sub>3</sub> oxidation (R1, R5) being more important in winter than in summer (Kunasek et al.,  
23 2008).



1 After atmospheric nitrate is deposited to the snow, UV photolysis will convert  
2 snow nitrate back to  $\text{NO}_x$ , which is then released to the atmosphere (Honrath et al.,  
3 1999; Thomas et al., 2012; Zatzko et al., 2013). This snow-sourced  $\text{NO}_x$  will be re-  
4 oxidized in the atmosphere to nitrate, which is subsequently re-deposited to surface  
5 snow or transported away. Reformation of nitrate in the condensed phase of snow  
6 grains can also occur if the nitrate being photolyzed is trapped inside the snow grain  
7 instead of on the surface (Meusinger et al., 2014). This is the so-called post-  
8 depositional processing of snow nitrate, which includes the steps of photolysis,  
9 recombination of photoproducts in the condensed phase and in the overlying  
10 atmosphere (i.e., recycling), and the re-deposition and/or export of snow-sourced  
11 nitrate. Post-depositional processing induces isotopic effects (Frey et al., 2009). In  
12 general, the photolysis of nitrate will enrich  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  of nitrate remaining in  
13 snow, but will not alter  $\Delta^{17}\text{O}$  as photolysis induces mass-dependent fractionation. If  
14 the photolytic products of nitrate are re-oxidized and re-deposited to the snow surface  
15 (i.e., no export/net loss), no measurable effect on  $\delta^{15}\text{N}(\text{NO}_3^-)$  will be observed  
16 assuming negligible nitrogen isotopic fractionation during the re-formation of nitrate.  
17 However, changes in  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  $\Delta^{17}\text{O}(\text{NO}_3^-)$  are expected even without net loss  
18 of nitrate. The reformation of nitrate in the condensed phase of snow grains will cause  
19 oxygen isotope exchange with water (Frey et al., 2009), which lowers  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  
20  $\Delta^{17}\text{O}(\text{NO}_3^-)$ . In addition, the re-oxidation of the snow-sourced  $\text{NO}_x$  to nitrate in the  
21 overlying atmosphere will occur mainly through the OH formation pathway. This is  
22 because active snow nitrate photolysis requires strong radiation, which is also when  
23 atmospheric nitrate is mainly formed through the daytime reaction channel ( $\text{NO}_x +$   
24 OH). Recycling of snow-sourced  $\text{NO}_x$  in the atmosphere will thus likely lower

1  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  $\Delta^{17}\text{O}(\text{NO}_3^-)$ , particularly if the initially deposited nitrate was formed  
2 through the  $\text{O}_3$  oxidation pathway (e.g., formed in polar winters when  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  
3  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values are the highest (Alexander et al., 2009; Kunasek et al., 2008)). The  
4 deposition of this reformed nitrate will then lower the bulk snow  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  
5  $\Delta^{17}\text{O}(\text{NO}_3^-)$ . At sites with snow accumulation rates greater than  $100 \text{ kg/m}^2/\text{yr}$ , the  
6 oxygen isotope exchange in the condensed phase is minimal (Erbland et al., 2013).  
7 This is consistent with the discussion in Fibiger et al. (2013) that the condensed phase  
8 process can't explain the observed relationship between  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  $\Delta^{17}\text{O}(\text{NO}_3^-)$  in  
9 surface snow, given the high snow accumulation rate at Summit ( $\sim 260 \text{ kg/m}^2/\text{yr}$ ,  
10 (Cole-Dai et al., 2013)). However, the Fibiger et al. (2013) analysis neglected  
11 atmospheric process, i.e., the reformation of nitrate from snow-sourced  $\text{NO}_x$  in the  
12 atmosphere and its subsequent deposition to surface snow. This process alters  
13  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  $\Delta^{17}\text{O}(\text{NO}_3^-)$  simultaneously and can explain the relationship between  
14  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  $\Delta^{17}\text{O}(\text{NO}_3^-)$  in surface snow observed by Fibiger et al. (2013). In  
15 addition, Fibiger et al. (2013) suggested that there is little to no post-depositional loss  
16 of nitrate at Summit, which might be true. But post-depositional loss is only one  
17 (possible) step of the post-depositional processing. Post-depositional processing of  
18 snow nitrate at Summit could be very active while little post-depositional loss occurs,  
19 given a fast recycling rate of nitrate in the air-snow interface.

20 In this study, the concentrations of major ions ( $\text{Na}^+$ ,  $\text{NH}_4^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  
21  $\text{NO}_3^-$ , and  $\text{SO}_4^{2-}$ ), and the isotopic composition of nitrate in a snowpit from central  
22 Greenland were measured. The concentration data were used to establish their  
23 temporal patterns and to identify any spring nitrate peak(s). The isotopic data were  
24 used to assess the chemistry of nitrate in any identified spring peaks and to discern the

1 origin of the peak. Further examinations were conducted on a shallow ice core to  
2 determine whether the mechanism leading to the spring nitrate peak observed in the  
3 snowpit is representative of the occasional nature of the spring nitrate peaks observed  
4 in Greenland snow since 1900.

## 5 **2. Methods**

### 6 **2.1. Snowpit and ice core sample collection**

7 In July 2007, six snow blocks (dimensions: 0.35 m long × 0.25 m wide × 0.35 m  
8 deep) were excavated from the surface down to a depth of 2.10 m at Summit,  
9 Greenland (72.5 °N, 38.5 °W; elevation: 3200 m). These snow blocks were wrapped  
10 with clean polyester film with their dimensions labeled and stored in a hard-shell box  
11 to avoid external compaction during delivery. Several ice cores including a 79 meter  
12 shallow core were drilled approximately 100 meters from the location of the snow  
13 blocks. These snow blocks and the ice cores were shipped frozen to the laboratory at  
14 South Dakota State University (SDSU) and stored in a freezer at -20 °C until analysis.  
15 At the same time, a set of snowpit samples were also collected in the field every 5 cm  
16 from the surface down to the depth of 2.10 m, at the same location of the collected  
17 snow blocks. This set of snowpit samples (referred to as SP-1) were double sealed in  
18 clean plastic bags and shipped frozen to Laboratoire de Glaciologie et Géophysique  
19 de l'Environnement (LGGE) for nitrate concentration and isotope analysis. All  
20 tools/containers directly touching the snowpit samples and/or the snow blocks were  
21 pre-cleaned with 18 MΩ water before use.

22 The density profile of the snow blocks were also measured in the field. In  
23 particular, in the field, a small snow cube was collected every 5 cm from the wall of a  
24 snowpit. The weight and the volume of the snow cube were measured and then the

1 density of the snow was calculated. The snow density profile from the surface to the  
2 depth of 2.1 m was included as supplemental data.

### 3 **2.2. Chemical and isotope analysis**

4 At SDSU, another set of snow samples (referred to as SP-2) in a vertical sequence  
5 was chiseled out from the snow blocks, after the removal of a surface layer of at least  
6 1 cm in thickness, and collected in clean plastic sample containers. All tools and  
7 sample containers were pre-cleaned with 18 MΩ water prior to use, and clean  
8 disposable plastic gloves were always worn during sample handling. In total, 71  
9 samples with a depth resolution of 3 cm were obtained. These samples were allowed  
10 to melt at room temperature and the meltwater samples were then analyzed by ion  
11 chromatography for concentrations of major ions in snow ( $\text{Na}^+$ ,  $\text{NH}_4^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  
12  $\text{Cl}^-$ ,  $\text{NO}_3^-$ , and  $\text{SO}_4^{2-}$ ). The ion chromatography instrumentation and methodology are  
13 similar to that described by Cole-Dai et al. (2006). Ice blanks were made from  
14 deionized water, processed and analyzed following the same procedure as that for the  
15 snow samples. High blank values of  $\text{Ca}^{2+}$  (up to  $5 \mu\text{g L}^{-1}$ ) indicated that the snowpit  
16 samples were probably contaminated with  $\text{Ca}^{2+}$  during sample preparation. No  
17 evidence of contamination was found for the other ions. Therefore, the snowpit  $\text{Ca}^{2+}$   
18 data were not used in interpretation in this study. The 79 meter shallow core was  
19 analyzed for the ionic species using the technique of continuous flow analysis with  
20 ion chromatography (CFA-IC) detection (Cole-Dai et al., 2006). One advantage of the  
21 CFA-IC technique is that it minimizes the potential contamination by eliminating  
22 sample preparation. Replicate analysis of blanks with CFA-IC showed no  
23 contamination of any of the ions including  $\text{Ca}^{2+}$ .

1           The concentration measurements of the SP-2 samples described above were used  
2 to estimate the volume of meltwater needed to provide sufficient nitrate mass (500  
3 nmoles, allowing for replicate measurements) for isotopic analysis. Based on that  
4 estimate, the snow blocks were carved vertically to yield large samples for isotope  
5 measurement. A total of 29 clean snow samples (referred to as SB) were obtained  
6 from the depth interval of 0.70 m to 1.75 m of the snow blocks (snow outside this  
7 depth interval had been consumed for other purposes). The depth resolution of these  
8 samples varies from 2 cm to 6 cm because the concentration of nitrate is different at  
9 different depths. These samples were melted at room temperature and concentrated,  
10 following the method described by Frey et al. (2009), to 10 mL solutions, collected in  
11 HDPE bottles, and stored frozen until isotopic analysis. Measurements of N and O  
12 isotope ratios in nitrate of the SB samples were performed in the stable isotope  
13 laboratory at University of Washington (UW) using the bacterial denitrifier method  
14 (Kaiser et al., 2007). Briefly, nitrate in the samples was first converted into N<sub>2</sub>O gas  
15 by the bacteria *Pseudomonas aureofaciens*; the N<sub>2</sub>O was carried on-line by helium  
16 gas into a heated gold tube where it was thermally decomposed to N<sub>2</sub> and O<sub>2</sub>. These  
17 were then separated by gas chromatography and the isotopic ratio(s) of each gas  
18 (<sup>15</sup>N/<sup>14</sup>N for N<sub>2</sub>, and <sup>18</sup>O/<sup>16</sup>O and <sup>17</sup>O/<sup>16</sup>O for O<sub>2</sub>) was measured with an isotope ratio  
19 mass spectrometer. The δ<sup>15</sup>N values were calculated with respect to N<sub>2</sub>-Air and  
20 calibrated against the two international reference materials IAEA-NO-3 (δ<sup>15</sup>N = 4.7  
21 ‰) and USGS34 ( δ<sup>15</sup>N = -1.8 ‰) (Kaiser et al., 2007). The δ<sup>17</sup>O and δ<sup>18</sup>O values  
22 were calculated with respect to VSMOW and calibrated against the two international  
23 reference materials USGS34 (δ<sup>17</sup>O = - 14.5 ‰, δ<sup>18</sup>O = - 27.9 ‰) and USGS35 (δ<sup>17</sup>O  
24 = 51.3 ‰, δ<sup>18</sup>O = 57.5 ‰) (Kaiser et al., 2007). The Δ<sup>17</sup>O values were then calculated

1 by using the linear equation  $\Delta^{17}\text{O} = \delta^{17}\text{O} - 0.52 \times \delta^{18}\text{O}$ . No replicate samples were  
2 possible due to the limited amount of snow available in the snow blocks, and no  
3 seasonally resolved isotopic measurements of the ice core samples were performed  
4 because of the limited amount of ice available. The analytical uncertainty of  
5  $\Delta^{17}\text{O}(\text{NO}_3^-)$  and  $\delta^{18}\text{O}(\text{NO}_3^-)$  measured at the UW laboratory was estimated to be 0.1  
6 ‰ and 0.5 ‰ ( $1\sigma$ ), and that of  $\delta^{15}\text{N}$  was 1.0 ‰ ( $1\sigma$ ), based on repeated measurements  
7 of the oxygen and nitrogen isotope ratios in the international reference materials  
8 USGS35 and IAEA-NO-3, respectively (Table 1 for details).

9       The SP-1 samples at LGGE were measured for nitrate concentration and  
10 isotopic composition ( $\delta^{15}\text{N}$ ,  $\delta^{18}\text{O}$  and  $\Delta^{17}\text{O}$ ). A small portion of each sample was  
11 taken for nitrate concentration measurement using the well-established Griess method  
12 in continuous flow analysis (Seal Analytical, Method No 1000223-1784-09). The rest  
13 of samples were processed following the similar procedure described above and the N  
14 and O isotope ratios were determined by the bacterial denitrifier method.  
15 Instrumentation details at the LGGE laboratory are similar to that described in  
16 Erbland et al. (2013). International reference materials (USGS34, USGS35 and  
17 IAEA-NO-3) were used for data reduction. The uncertainties of  $\Delta^{17}\text{O}(\text{NO}_3^-)$ ,  
18  $\delta^{18}\text{O}(\text{NO}_3^-)$  and  $\delta^{15}\text{N}(\text{NO}_3^-)$  measured at the LGGE laboratory were 0.3 ‰, 2.0 ‰ and  
19 0.3 ‰, respectively, estimated as the reduced standard deviation of the residuals from  
20 the linear regression between the measured reference materials and their accepted  
21 values.

## 22 **3. Results**

### 23 **3.1. Dating and identifying spring nitrate peak(s)**

1 The concentration data from the SP-1 and SP-2 samples, and the isotopic data  
2 from the SP-1 and SB samples, were plotted as a function of depth in Figure 1 (data  
3 are available in Supplemental Material). The concentration profiles of nitrate from  
4 SP-1 and SP-2 are nearly identical and each displays five peaks approximately at the  
5 same depths (labeled 1, 2, 3, 3' and 4 in Figure 1d). We used the sodium ( $\text{Na}^+$ ) peak  
6 as a winter snow layer indicator and the peak of the  $\text{Cl}^-/\text{Na}^+$  ratio as a summer snow  
7 layer indicator (Whitlow et al., 1992). Sodium concentration in Greenland snow peaks  
8 in winter/late winter due to winter Arctic aerosols (with high sea salt content) and/or  
9 late winter cyclonic storms, while  $\text{Cl}^-/\text{Na}^+$  peaks in summer likely due to long range  
10 transport of aged sea salt aerosols (Whitlow et al., 1992) which is depleted in  $\text{Na}^+$   
11 because of the preferential removal of  $\text{Na}^+$  in aerosols relative to gaseous HCl  
12 following sea salt-acid displacement (Legrand and Delmas, 1988). Since the snowpit  
13 and snow blocks were excavated in July of 2007, the partial peak of  $\text{Cl}^-/\text{Na}^+$  at the  
14 surface identifies the summer of 2007. The other  $\text{Cl}^-/\text{Na}^+$  peaks represent the summer  
15 of 2006, 2005 and 2004, respectively (Figure 1e) The nitrate peaks 1, 2, 3 and 4 are in  
16 almost identical layers as the  $\text{Cl}^-/\text{Na}^+$  peaks, providing independent evidence that they  
17 can be considered summer peaks. Nitrate Peak 3' is between a winter  $\text{Na}^+$  peak and a  
18 summer  $\text{Cl}^-/\text{Na}^+$  peak, clearly identifying it as a spring peak. This dating by  $\text{Cl}^-/\text{Na}^+$   
19 and  $\text{Na}^+$  peaks (referred to here as Method A) is consistent with the results of weekly  
20 recorded snow accumulation data determined by stake height measurements at  
21 Summit (data are available at  
22 [ftp://ftp.summitcamp.org/pub/data/GEOSummit/Bales\\_UCM/Bamboo%20Forest/Bamboo%20Forest%20Accumulation%20Log.xls](ftp://ftp.summitcamp.org/pub/data/GEOSummit/Bales_UCM/Bamboo%20Forest/Bamboo%20Forest%20Accumulation%20Log.xls)) (referred to here as Method B). The  
23  
24 approximate months of snowfall at depths of 0, 0.5, 1.0, 1.5 and 2.0 m identified by

1 Method B were indicated with vertical dashed lines of Figure 1. According to Method  
2 B, snow in the layer of the Peak 3' fell in February of 2005, suggesting it is a early  
3 spring peak, consistent with previous studies that the additional annual nitrate peak  
4 occurs in late winter/early spring (Burkhart et al., 2006; Yang et al., 1995). Some  
5 small discrepancies exist between the two dating methods. For example, the month of  
6 snowfall at the surface was identified as May of 2007 using Method B instead of July.  
7 This discrepancy is due to negligible snowfall (around 2 cm) from May to July 2007.  
8 The month of snowfall at the depth of 2 m was identified as March of 2004 using  
9 Method B, suggesting that Peak 4 is also a spring peak. However, according to the  
10 snow accumulation data, there was only about 3 cm of snow accumulation from  
11 March to June of 2004 (Supplemental Figure 1). Negligible snowfall during this time  
12 resulted in nitrate produced in summer dry depositing to the prior spring snow layers,  
13 making it appear as if the peak occurs in spring using Method B. In addition, the  
14 nitrate profiles in SP-1 and SP-2 are slightly different (i.e., nitrate peaks are not  
15 exactly at the same depths), which is likely due to the spatial variability of snow  
16 accumulation rates in the field due to snow drifting. But the overall temporal patterns  
17 of the nitrate profiles in SP-1 and SP-2 are similar, suggesting there was negligible  
18 disturbance of the stratigraphy of the snow blocks during delivery and storage.

19 The record of nitrate concentrations from the shallow ice core is shown in Figure  
20 2 (data are available in Supplemental Material). The annual layer dating of this core  
21 was performed by counting the annual spring peak of calcium (Cole-Dai et al., 2013).  
22 From this ice core, we found there are nineteen years with two nitrate peaks in the  
23 period of 1960 to 2006, obtained by subtracting total calcium peaks from total nitrate  
24 peaks in this period (Figure 2). We ~~didn't~~don't attempt to specifically identify the



1 seasonality of these peaks due to the lack of a reliable summer layer indicator in the  
2 firn core. The  $\text{Cl}^-/\text{Na}^+$  ratio is influenced by atmospheric acidity (details in  
3 Supplemental Material). In the preindustrial period, atmospheric acidity peaks in  
4 summer due to high summer production of non-sea-salt sulfate and nitrate (Whitlow  
5 et al., 1992). This leads to an enhancement in excess chloride in summer through  
6 acid-sea salt displacement (Legrand and Delmas, 1988). Given the summer low  $\text{Na}^+$   
7 concentrations, the  $\text{Cl}^-/\text{Na}^+$  peak falls almost precisely in summer in the preindustrial  
8 period, making it a good summer layer indicator. However, in much of the industrial  
9 era, the annual  $\text{Cl}^-/\text{Na}^+$  peak shifts toward the major annual non-sea-salt sulfate peak  
10 (usually in winter, Figure S2, top panel) inhibiting its ability to serve as a summer  
11 layer indicator. This makes it difficult to unambiguously identify the secondary nitrate  
12 peak in the years with double nitrate peaks as occurring in a specific season (e.g.,  
13 spring), though we note that previous studies have been able to identify the secondary  
14 nitrate peak as occurring in the spring by using summer layer indicators (e.g.,  $\text{H}_2\text{O}_2$ ,  
15 and  $\delta^{18}\text{O}(\text{H}_2\text{O})$ ) independent of major ion concentrations (Burkhart et al., 2006;  
16 Whitlow et al., 1992; Yang et al., 1995). We note that, in the snowpit, sulfate  
17 concentrations are as low as they were prior to 1900 (Geng et al., 2014), minimizing  
18 its impact on the timing of the  $\text{Cl}^-/\text{Na}^+$  peak. As discussed explicitly in sub-annual  
19 layer dating of the ice core samples by seasonal peaks of ion concentrations is  
20 imprecise and carries an estimated uncertainty of  $\pm 4$  months for ice cores drilled at  
21 Summit (sub-annual layer dating of the snowpit samples has a much lower uncertainty  
22 due to relatively high temporal resolution). However, we note that the practice of  
23 referring to the additional nitrate peak as a spring peak is both consistent with  
24 previous studies and with our sub-annual layer dating of the snowpit. The frequency

1 of ~~years with double nitrate peaks the spring peak~~ in each decade ~~in this period~~ is  
2 listed in Table 2. Prior to 1960, only a single nitrate peak is observed in each annual  
3 layer~~No such spring peak was seen before 1960, consistent with similar to~~ the finding  
4 by Finkel and Langway (1986) that the spring nitrate peak started appearing after the  
5 1950s, though others (Burkhart et al., 2006; Yang et al., 1995) stated in general that  
6 elevated spring nitrate concentrations are seen in snow layers after around 1900.

### 7 **3.2. $\delta^{18}\text{O}(\text{NO}_3^-)$ , $\delta^{15}\text{N}(\text{NO}_3^-)$ and $\Delta^{17}\text{O}(\text{NO}_3^-)$ variations**

8 The  $\delta^{18}\text{O}(\text{NO}_3^-)$ ,  $\delta^{15}\text{N}(\text{NO}_3^-)$  and  $\Delta^{17}\text{O}(\text{NO}_3^-)$  data from the SP-1 and SB samples  
9 are shown in Figure 1a, 1b and 1c, respectively. The depth range of the SP-1 samples  
10 (0 to 2.10 m) corresponds to the period of the 2007 summer to 2004 summer, while  
11 the depth range of the SB samples (0.7 to 1.75 m) corresponds to the time period of  
12 the 2006 spring/summer to the 2004/2005 winter. In general, all three isotopic  
13 signatures of nitrate display large seasonal variations.  $\delta^{18}\text{O}(\text{NO}_3^-)$  is low in summer  
14 and high in winter, while  $\delta^{15}\text{N}(\text{NO}_3^-)$  is high in spring/early summer and low in  
15 winter, consistent with previous measurements of Summit snow samples from  
16 Hastings et al. (2004). In the spring of 2005 (the time period corresponding to the  
17 depth of the spring nitrate peak), the means ( $\pm 1\sigma$ ) of  $\delta^{15}\text{N}(\text{NO}_3^-)$  from the SP-1 and  
18 SB samples were  $(5.3 \pm 3.3) \text{‰}$  and  $(6.4 \pm 2.1) \text{‰}$ , respectively, much higher than  
19 winter values  $(-9.2 \pm 3.3) \text{‰}$  from the SP-1 samples and  $(-7.4 \pm 4.4) \text{‰}$  from the SB  
20 samples. A local minimum in  $\delta^{18}\text{O}(\text{NO}_3^-)$  was also observed from both the SP-1 and  
21 the SB samples at the depth of the 2005 spring nitrate peak (Figure 1a).

22 The observed seasonality of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  (Figure 1c) is consistent with the  
23 expectation of high  $\Delta^{17}\text{O}(\text{NO}_3^-)$  during winter (polar night) due to the dominance of  
24 the  $\text{O}_3$  oxidation pathway of  $\text{NO}_x$  (R5-R8). The magnitude of the seasonality is also

1 consistent with the observations of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  at Summit by Kunasek et al.(2008).  
2 The seasonality of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  is regulated by the shift of the dominant nitrate  
3 formation pathway from OH oxidation (R4) in polar summer/day to that via  $\text{O}_3$  (R5-  
4 R8) in polar winter/night (low summer values and high winter values). Modeling by  
5 Kunasek et al. (2008) showed that, in early spring,  $\Delta^{17}\text{O}$  of locally produced nitrate at  
6 Summit should be close to that in winter snow, as oxidation by OH (R4) is very  
7 limited at this time due to the lack of sunlight. In other words, at Summit  $\Delta^{17}\text{O}(\text{NO}_3^-)$   
8 values in winter and early spring should be similar. However, in the early spring of  
9 2005, the  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values display apparent declines from the prior winter values  
10 (Figure 1c). To test whether the declines are significant, we calculate the mean  
11  $\Delta^{17}\text{O}(\text{NO}_3^-)$  value in the early spring of 2005 and compare it to the mean of measured  
12 winter  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values. The winter mean  $\Delta^{17}\text{O}(\text{NO}_3^-)$  was calculated from the  
13 2004/2005 to 2006/2007 winters from the SP-1 samples, and from the 2004/2005 to  
14 2005/2006 winters for the SB samples (samples used to calculate the winter and 2005  
15 early spring means are as marked in Figure 1c). The mean winter value was  $(32.9 \pm$   
16  $1.1) \text{‰}$  from the SP-1 samples and  $(32.4 \pm 0.6) \text{‰}$  from the SB samples. In contrast,  
17 the mean  $\Delta^{17}\text{O}(\text{NO}_3^-)$  value in the early spring of 2005 was  $(31.1 \pm 0.6) \text{‰}$  from SP-1  
18 and  $(30.8 \pm 0.8) \text{‰}$  from SB, approximately 1.7 ‰ lower than the winter means,  
19 which is statistically significant based on one-tailed t-test (for SP-1 samples:  $t =$   
20  $3.434$ ,  $\text{DOF} = 24$ ,  $P = 0.001$ ; for SB samples:  $t = 4.637$ ,  $\text{DOF} = 17$ ,  $P \leq 0.0005$ ). A  
21 previous study measuring  $\Delta^{17}\text{O}(\text{NO}_3^-)$  in Summit snow by Kunasek et al. (2008) also  
22 noted significantly low  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values (around 26 ‰) in the spring of 2005,  
23 compared to the prior winter of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  around 33 ‰.

#### 24 **4. Discussion**

1 The low  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values in the spring of 2005 suggest an increase in OH  
2 oxidation in the formation of nitrate associated with the peak. This is qualitatively  
3 consistent with the observed decrease in  $\delta^{18}\text{O}(\text{NO}_3^-)$  from a previous study (Hastings  
4 et al., 2004), as increases in the relative importance of OH oxidation will also result in  
5 decreases in  $\delta^{18}\text{O}(\text{NO}_3^-)$ . The extra nitrate deposited during the spring of 2005 was  
6 estimated to be  $5.7 \text{ nmol} \times \text{cm}^{-2}$ , which was calculated by subtracting the flux of  
7 nitrate ( $9.4 \text{ nmol} \times \text{cm}^{-2}$ ) in the spring of 2006 from the nitrate flux in the spring of  
8 2005 ( $15.1 \text{ nmol} \times \text{cm}^{-2}$ ). The 2006 spring was a normal spring without nitrate peak.  
9 This suggests a 60 % increase in nitrate deposited in the spring of 2005 compared to  
10 the spring of 2006. The additional nitrate in the spring of 2005 could either come from  
11 enhanced transport of nitrate produced elsewhere or enhanced local nitrate production  
12 ~~be produced locally~~ involving  $\text{NO}_x$  precursors. In the discussion to follow, we  
13 consider separately the possibility of enhanced transport and enhanced local  
14 production of nitrate in the Arctic being responsible for the spring 2005 nitrate peak.  
15 We further examine whether the explanation for the case of spring 2005 is  
16 representative of the occasional nature of the additional nitrate peaks observed in the  
17 firm shallow ice core.

#### 18 **4.1. Enhanced transport of nitrate**

##### 19 **4.1.1. Stratospheric denitrification**

20 Stratospheric denitrification refers to the sedimentation process of Polar  
21 Stratospheric Clouds (PSCs) containing nitric acid trihydrate. Research on nitrate in  
22 Antarctic snow (Mulvaney and Wolff, 1993) suggested that stratospheric  
23 denitrification associated with the winter polar vortex could result in a late  
24 winter/early spring nitrate peak. Although denitrification occurs less frequently and

1 less extensively in the Arctic than in Antarctica due to the warmer winter and the  
2 weaker and less persistent Arctic vortex (Waugh and Randel, 1999), significant  
3 denitrification has been observed in Arctic for some exceptionally cold winters,  
4 including the winter of 2004/2005 (Mann et al., 2003; Jin et al., 2006; Kleinbohl et  
5 al., 2005). Could stratospheric denitrification in the winter of 2004/2005 have resulted  
6 in the early spring nitrate peak?

7 Stratospheric nitrate is expected to possess high  $\Delta^{17}\text{O}$  since it is mainly formed  
8 via  $\text{O}_3$  oxidation (McCabe et al., 2007; Savarino et al., 2007) and stratospheric  
9  $\Delta^{17}\text{O}(\text{O}_3)$  is up to 5 ‰ higher than that in the troposphere (Liang et al., 2006; Lyons,  
10 2001; Mauersberger et al., 2001). In addition, stratospheric nitrate can also be formed  
11 via ClO oxidation (McCabe et al., 2007). Nitrate formed via ClO oxidation also  
12 possesses high  $\Delta^{17}\text{O}$  (McCabe et al., 2007; Savarino et al., 2007) due to the internal  
13 non-statistical distribution of isotopes in  $\text{O}_3$  which transfers its terminal oxygen atoms  
14 to ClO (Bhattacharya et al., 2008).

15 If the additional nitrate in the 2005 spring snow originated from stratospheric  
16 denitrification, the  $\Delta^{17}\text{O}(\text{NO}_3^-)$  would be expected to be anomalously high, not low as  
17 was observed. It is possible that the  $\Delta^{17}\text{O}$  value of nitrate deposited to the snow  
18 surface is reduced by post-depositional processing, for the photolytic recycling of  
19 snowpack nitrate (nitrate photolysis followed by re-formation and re-deposition of  
20 nitrate) can reduce the  $\Delta^{17}\text{O}$  signature of nitrate finally preserved in snow (Erbland et  
21 al., 2013). If all of the nitrate produced in the stratosphere undergoes photolytic  
22 recycling, the stratospheric  $\Delta^{17}\text{O}$  signature would be completely erased (i.e.,  $\Delta^{17}\text{O}$  of  
23 the nitrate would be the same as that of nitrate produced in the tropospheric boundary  
24 layer). Given the high snow accumulation rate at Summit (the degree of snowpack

1 nitrate photolysis decreases with increasing snow accumulation rate (R öthlisberger et  
2 al., 2002)), it is unlikely that all additional nitrate underwent photolytic recycling.  
3 Consequently,  $\Delta^{17}\text{O}$  of the preserved nitrate originally produced in the stratosphere  
4 should be no lower than that of the tropospheric nitrate deposited in a typical spring.  
5 Therefore, the observed  $\Delta^{17}\text{O}(\text{NO}_3^-)$  suggests stratospheric denitrification cannot  
6 account for the spring nitrate peak.

#### 7 **4.1.2. Transport from mid-latitudes**

8 Previous studies (Burkhart et al., 2006; Yang et al., 1995) hypothesized that the  
9 spring nitrate peak results from pollution transport in the industrial era, though the  
10 occasional nature of the peak was not explicitly discussed. In case of strong  
11 meridional transport of nitrate ~~and/or  $\text{NO}_x$  precursors~~, a spring nitrate peak may be  
12 expected. In addition, the  $\Delta^{17}\text{O}$  of nitrate formed in the mid-latitudes is likely lower  
13 than that of nitrate formed in the Arctic troposphere due to the latitudinal gradient in  
14  $\Delta^{17}\text{O}(\text{NO}_3^-)$  resulting from the latitudinal gradient in  $\text{O}_3/\text{HO}_x$  ratio (Alexander et al.,  
15 2009). Thus, enhanced long-range transport of mid-latitude nitrate to Greenland  
16 during the spring could elevate nitrate concentrations in snow with relatively low  
17  $\Delta^{17}\text{O}(\text{NO}_3^-)$ , consistent with the observations. Direct transport of nitrate from the mid-  
18 latitudes to Summit is also consistent the conclusions in Kunasek et al. (2008) and  
19 Geng et al. (2014).

20 Pollution from the Northern mid-latitudes is transported to the Arctic by  
21 poleward meridional circulation, which is strong when the North Atlantic Oscillation  
22 (NAO) is in its positive phase (Eckhardt et al., 2003). However, the NAO index data  
23 in the 2004/2005 winter and early spring of 2005 is not strong compared with other  
24 years during the period of 1997 to 2009 (Osborn, 2011), suggesting that meridional

1 ~~transport is not particularly strong no-enhanced transport from the mid-latitudes in the~~  
2 ~~spring of 2005 at this time compared to other springs. However, frequent pollution~~  
3 ~~transport events occur even in years with a negative NAO phase~~ (Kramer et al., 2014).  
4 ~~To examine the possibility of pollution transport as the direct cause of the 2005 spring~~  
5 ~~nitrate peak, we examine the profiles of  $\text{SO}_4^{2-}$  and  $\text{Mg}^{2+}$  from the snowpit. If the~~  
6 ~~additional nitrate in the spring of 2005 was from Eenhanced transport from the mid-~~  
7 ~~latitudes will also, elevated~~ concentrations of other species derived from  
8 anthropogenic and continental sources, such as  $\text{SO}_4^{2-}$  and  $\text{Mg}^{2+}$ , ~~would also be~~  
9 ~~expected, especially for~~  $\text{SO}_4^{2-}$  because it has a very similar atmospheric lifetime (4 to  
10 6 days) as nitrate (3 to 7 days) (Park et al., 2004; Pye et al., 2009). However, the data  
11 from the snowpit show that neither  $\text{SO}_4^{2-}$  nor  $\text{Mg}^{2+}$  concentrations were elevated  
12 (Figure 1f and 1g) in the spring of 2005 when the nitrate peak was present. Therefore,  
13 it is unlikely that the 2005 spring nitrate peak was caused by enhanced long-range  
14 transport of nitrate from the mid-latitudes during that time period.

15 In addition to pollution from industrial areas, biomass burning (e.g., wild fire)  
16 plumes from Boreal North America could also potentially increase snow nitrate  
17 concentrations in Greenland. However, the global fire emission data (Giglio et al.,  
18 2013) suggests that fire emissions in Boreal North America (~~n~~or in Temperate North  
19 America and Europe) are not strong in ~~the 2005-spring of 2005~~ compared to other  
20 years during the period from 1997 to 2011. In addition, chemical species transported  
21 from fire events will likely cause episodic, sharp spikes, such as the  $\text{NH}_4^+$  spikes  
22 frequently observed in Greenland ice cores (Savarino and Legrand, 1998). The  
23 ~~duration (> 1 month) morphology (shape)~~ of the 2005 spring nitrate peak ~~argues~~

1 | ~~against an episodic biomass-burning source seems to suggest a source different from~~  
2 | ~~fire events.~~

## 3 | **4.2. Enhanced local production**

### 4 | **4.2.1. PAN decomposition**

5 | Previous work has found that peroxyacetyl nitrate (PAN) is the dominant form  
6 | of reactive nitrogen in the Arctic troposphere during winter and spring (Beine and  
7 | Krognes, 2000; Bottenheim et al., 1993; Honrath and Jaffe, 1992; Jacobi et al., 1999)  
8 | and its decomposition yields NO<sub>x</sub> that is subsequently converted to nitrate. If the  
9 | decomposition rate of PAN is enhanced in the spring of 2005, more NO<sub>x</sub> would be  
10 | produced resulting in an increase in local nitrate production. The possibility of PAN  
11 | decomposition in spring leading to the spring nitrate peak has been mentioned in  
12 | Yang et al. (1995), but not explicitly examined. Here we provide a detailed  
13 | examination of the possible role of PAN decomposition in the spring 2005 nitrate  
14 | peak.

15 | PAN decomposes to NO<sub>x</sub> either thermally or via photolysis (Talukdar et al.,  
16 | 1995). At temperatures above 255 K, thermal decomposition dominates, while  
17 | photolysis becomes more important at lower temperatures (Talukdar et al., 1995). The  
18 | typical spring (February and March) air temperature at Summit is about 240 K (242 K  
19 | in 2005 and average of 236 K in the springs of 2006 to 2008; data are from  
20 | <http://www.summitcamp.org/resources/files>). At these low temperatures, the NO<sub>x</sub>  
21 | release from PAN is dominated by photolysis at wavelengths between 290-345 nm  
22 | (Talukdar et al., 1995; Flowers et al., 2005).

23 | The stratospheric O<sub>3</sub> layer filters out most of the UV-B ( $\lambda = 290\text{-}320\text{ nm}$ )  
24 | portion of the solar spectrum. A weakened stratospheric O<sub>3</sub> layer will allow more UV-



1 B penetration into the troposphere. In Figure 3, the spring (average of February and  
2 March values) O<sub>3</sub> column density at Summit for the years of 1979-2006 are shown  
3 (data are from NASA Total Ozone Mapping Spectrometer program). The 2005 spring  
4 level (290 Dobson Unit (DU)) was significantly lower than the average of spring  
5 values from 1979 to 2006 ( $390 \pm 50$  ( $1\sigma$ ) DU), suggesting a relatively high UV-B flux  
6 to the surface in the spring of 2005.

7 The photolysis of PAN in the spring of 2005 could have been enhanced due to  
8 the elevated UV-B radiation caused by stratospheric O<sub>3</sub> loss, leading to increased  
9 local NO<sub>x</sub> abundance, and subsequently resulting in the unusual spring nitrate peak in  
10 snow. The increase in PAN photolysis resulting from the decreased O<sub>3</sub> column  
11 density (290 DU) is estimated with the UCAR Tropospheric Ultraviolet & Visible  
12 (TUV) radiation model (available at <http://cprm.acd.ucar.edu/Models/TUV/>).  
13 Calculations with the TUV model showed a 24 % increase in the photolysis rate  
14 constant of PAN ( $j = (9.0 \pm 5.4) \times 10^{-8} \text{ s}^{-1}$  vs.  $(7.3 \pm 4.3) \times 10^{-8} \text{ s}^{-1}$ , both are daytime  
15 averages) in the spring of 2005 compared to that in a typical spring (column O<sub>3</sub>  
16 density of 390 DU). This would lead to an increase of only  $1.6 \times 10^5 \text{ cm}^{-3}$  in NO<sub>x</sub>  
17 number concentration, which is, on average, 0.04 % of the observed springtime NO<sub>x</sub>  
18 concentration of  $(3.9 \pm 3.1) \times 10^8 \text{ cm}^{-3}$  in the Arctic (Stroud et al., 2003). The above  
19 calculation assumes steady state of PAN with number concentration of  $(2.3 \pm 0.7) \times$   
20  $10^9 \text{ cm}^{-3}$  (Stroud et al., 2003) and  $[\text{NO}_2] = (([\text{PAN}] \times j_{\text{PAN}}) / k)^{1/2}$ , where  $[\text{NO}_2]$  and  
21  $[\text{PAN}]$  represent the number concentrations of NO<sub>2</sub> and PAN, respectively, and  $j_{\text{PAN}}$   
22 ( $\text{s}^{-1}$ ) is the photolysis rate constant of PAN calculated from the TUV model and  $k$  ( $\text{cm}^3$   
23  $\times \text{s}^{-1}$ ) is the reaction rate constant of PAN formation at  $T = 240 \text{ K}$  and  $P = 650 \text{ hPa}$   
24 (Summit springtime condition) calculated based on the equation from Atkinson et al.

1 (2006). The additional production of  $\text{NO}_x$  from enhanced PAN photolysis is  
2 negligible compared to the observed 60 % enhancement in nitrate deposition flux ( $5.7$   
3  $\text{nmol} \times \text{cm}^{-2}$ ) during the spring of 2005. This is consistent with the model prediction  
4 by Stroud et al. (2003) that, during spring, PAN is a net sink of  $\text{NO}_x$ , rather than a  
5 source. A similar conclusion was also reached by Singh et al. (1992) who found that  
6 the PAN reservoir is not a significant source of  $\text{NO}_x$  until summer.

#### 7 **4.2.2. Snowpack $\text{NO}_x$ emissions**

8 The photolysis of nitrate in the snowpack at snow depths of up to 20 cm or  
9 more (Zatko et al., 2013; France et al., 2011; Grannas et al., 2007) emits  $\text{NO}_x$  to the  
10 overlying atmosphere (Honrath et al., 1999; Frey et al., 2009), serving as a local  $\text{NO}_x$   
11 source.  $\text{NO}_x$  originating from the photolysis of nitrate in the snowpack can be re-  
12 oxidized in the atmosphere to nitrate and re-deposited to the surface. A recent model  
13 study (Thomas et al., 2012) suggested that at Summit, the photolysis of snowpack  
14 nitrate alone can sustain observed  $\text{NO}$  concentrations in the local atmospheric  
15 boundary layer. If the emission of  $\text{NO}_x$  from snowpack were enhanced, local  
16 atmospheric nitrate production would also be expected to be elevated. The reduction in  
17  $\text{O}_3$  column density in the spring of 2005 led to more UV radiation at the surface,  
18 which in turn may significantly enhance the photolysis of snow nitrate  
19 photolysis. Consequently, the oxidation of  $\text{NO}_x$  released from the snowpack by  
20 enhanced photolysis of snowpack nitrate at depths, and subsequent re-deposition of  
21 the nitrate to the surface in spring, would result in elevated nitrate concentration in  
22 surface snow during spring.

23 In order to estimate the impact of the reduced overhead  $\text{O}_3$  column density in the  
24 spring of 2005 on local  $\text{NO}_x$  concentration via the photolysis of snowpack nitrate, we


1 used the TUV model to calculate the surface actinic flux at Summit in the spring of  
2 2005 (290 DU) versus normal springtime with average O<sub>3</sub> column density (390 DU,  
3 which was close to the value (380 DU) in the spring of 2006). The parameterization  
4 from Zatzko et al. (2013) was then used to calculate the photolysis frequency (s<sup>-1</sup>) of  
5 snowpack nitrate at wavelengths from 298 to 345 nm (peak wavelength for nitrate  
6 photolysis is around 320 nm (Chu and Anastasio, 2003)). This calculation suggested a  
7 30 % enhancement of the snowpack nitrate photolysis frequency in the spring of 2005  
8 relative to 2006, and thus a similar enhancement of NO<sub>x</sub> emission from the snowpack,  
9 due to the reduced overhead O<sub>3</sub> column density alone. This likely contributed to the  
10 enhanced local nitrate production in the spring of 2005. However, the relative  
11 importance of snowpack nitrate photolysis to local NO<sub>x</sub> abundance in springtime is  
12 unknown, making it difficult to quantitatively assess the contribution from snowpack  
13 emissions to the observed spring nitrate peak. If snowpack nitrate photolysis nitrate  
14 photolysis is the dominant source of boundary layer NO<sub>x</sub> at Summit as suggested by  
15 Thomas et al. (2012), then this could account for up to half of the additional nitrate in  
16 the observed 2005 spring nitrate peak.


17       The relatively high  $\delta^{15}\text{N}(\text{NO}_3^-)$  values in the spring of 2005 (SP-1:  $(5.3 \pm 3.3)$   
18 ‰; SP-2  $(6.4 \pm 2.1)$  ‰) indicate some post-depositional loss of snowpack nitrate at  
19 that time. Since post-depositional loss is driven by photolysis (Berhanu et al., 2014;  
20 Frey et al., 2009), the observed high  $\delta^{15}\text{N}(\text{NO}_3^-)$  values suggest active snow nitrate  
21 photolysis during spring 2005. This is qualitatively consistent with the observations at  
22 Summit by Hastings et al. (2004) that  $\delta^{15}\text{N}(\text{NO}_3^-)$  in surface snow is significantly  
23 higher in daytime when snow nitrate photolysis is active compared to during the  
24 nighttime.

### 4.2.3. Enhanced conversion of NO<sub>x</sub> to nitrate



Field observations suggest that in present day, there is probably sufficient NO<sub>x</sub> at Summit to ensure local nitrate production even in the spring. The number concentration of NO<sub>x</sub> in the Arctic mid-troposphere (58-85° N, 3-6 km) during spring of 2002 has been reported to be  $(3.9 \pm 3.1) \times 10^8 \text{ cm}^{-3}$  (Stroud et al., 2003), which is similar to the summer NO<sub>x</sub> level at Summit (Jacobi et al., 2004). In addition, Kramer et al. (2014) reported that February NO<sub>x</sub> concentration at Summit is  $(1.6 \pm 3.2) \times 10^8 \text{ cm}^{-3}$  which overlaps with summer values due to the large uncertainty. Kramer et al. (2014) also suggested that pollution transport during polar winters leads to accumulation of NO<sub>y</sub> as well as NO<sub>x</sub> in the Arctic, and implied that part of the NO<sub>y</sub> will be converted to NO<sub>x</sub> after polar sunrise through photolysis. Under ~~this scenario~~ with sufficient local NO<sub>y</sub> and NO<sub>x</sub>, an increase in nitrate concentration would be expected if the oxidation rate of NO<sub>x</sub> to HNO<sub>3</sub> is also enhanced, ~~even in the absence of increased springtime NO<sub>x</sub> concentrations.~~ The NO<sub>x</sub> oxidation rate can be enhanced by increased oxidant (O<sub>3</sub>, OH, BrO) levels. Both O<sub>3</sub> and BrO have high  $\Delta^{17}\text{O}$  values (Lyons, 2001; Morin et al., 2007), thus an increase in either O<sub>3</sub> or BrO concentrations would result in additional nitrate with high  $\Delta^{17}\text{O}$ . The relatively low  $\Delta^{17}\text{O}(\text{NO}_3^-)$  in the spring 2005 snow (Figure 1c) suggests that increased tropospheric O<sub>3</sub> and/or BrO concentrations are unlikely to be the direct cause of enhanced nitrate production.

On the other hand, increased oxidation of NO<sub>x</sub> by OH would produce additional nitrate with low values of  $\Delta^{17}\text{O}$  as was observed. In general, the concentration of tropospheric OH is dependent on concentrations of tropospheric O<sub>3</sub> and water vapor and the available UV-B radiation through following reactions:





2 Either a substantial increase in UV-B in the troposphere or an increase in the  
3 water vapor content at elevated atmospheric temperatures can increase OH production  
4 and therefore enhance the conversion of  $\text{NO}_x$  to  $\text{HNO}_3$  via OH oxidation (R4). In  
5 addition, two other important oxidants involved in NO- $\text{NO}_2$  cycling (R2),  $\text{HO}_2$  and  
6  $\text{RO}_2$ , are formed by reactions of carbon monoxide (CO) and hydrocarbons (RH) with  
7 OH:



10 Increased OH will enhance NO- $\text{NO}_2$  cycling via  $\text{HO}_2$  and  $\text{RO}_2$  and reduce the relative  
11 importance of  $\text{O}_3$  in  $\text{NO}_x$  cycling, which also leads to lowered  $\Delta^{17}\text{O}$  in nitrate.

12 Using temperature and relative humidity data at Summit from the Greenland  
13 Climate Network Data (GCND,  
14 <http://cires.colorado.edu/science/groups/steffen/gcnet/order/admin/station.php>), we  
15 estimate that the spring (February and March average) atmospheric water vapor  
16 mixing ratio in 2005 was 40 % higher than that in 2006 due to high temperatures in  
17 the 2005 spring (-30 °C) relative to the 2006 spring (-34 °C). Assuming this high  
18 water vapor in spring 2005 will result in a maximum increase in OH production of 40  
19 %, the increase in water vapor is not enough to account for the 60 % enhancement in  
20 the nitrate deposition flux in the spring of 2005. In fact, as discussed later, OH must  
21 be increased by 200 % to explain the full magnitude of the spring nitrate maximum.  
22 Although the relatively high temperature in the spring of 2005 may have contributed  
23 around 20 % (40 % increase versus the required 200 % increase) to the enhanced local

1 nitrate production, it was likely not the dominant factor producing the spring nitrate  
2 peak in 2005.

3 To determine the effect of reduced column O<sub>3</sub> density on OH production at the  
4 surface, we used the TUV model to estimate the OH production rate (R9 and R10) at  
5 Summit due to enhanced photolysis of boundary layer O<sub>3</sub>. Our calculation showed  
6 that the OH production rate due to this mechanism alone was increased by 90 % in the  
7 spring of 2005 resulting from the decrease in O<sub>3</sub> column density from 390 to 290 DU.  
8 Assuming that the production of HNO<sub>3</sub> via OH oxidation was increased by the same  
9 factor, enhanced local OH production could easily account for the 60 % enhancement  
10 of nitrate flux in the spring of 2005, if all of the additional nitrate was formed via OH  
11 oxidation (R4). During a typical Arctic spring, O<sub>3</sub> is expected to be the major oxidant  
12 in the oxidation of NO<sub>x</sub> to HNO<sub>3</sub> (Alexander et al., 2009), so it is unlikely that R4 is  
13 solely responsible for the local oxidation of NO<sub>x</sub> to HNO<sub>3</sub>. Using a global chemical  
14 transport model (GEOS-Chem), Alexander et al. (2009) estimated that about 30 % of  
15 total nitrate in a normal spring (average in February and March) is produced via OH  
16 oxidation at Summit. With this more realistic value, a 200 % increase in the local OH  
17 production rate was needed to account for the additional nitrate flux in the spring of  
18 2005. Therefore, the enhanced local production rate of OH due to solely the enhanced  
19 photolysis of tropospheric O<sub>3</sub> in the spring of 2005 can explain about 50 % of the  
20 observed spring nitrate peak.

21 In addition to production via O<sub>3</sub> photolysis (R9 and R10), OH can be formed by  
22 the photolysis of snowpack emitted CH<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, and HONO (Dassau et al., 2002;  
23 Hutterli et al., 2001; Zhou et al., 2001; Sjostedt et al., 2007). Some Mm model studies  
24 suggested that at Summit photolysis of snow-sourced CH<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub> and HONO

1 contributes ~~to OH 2-3 times more than that a similar amount to local OH production~~  
2 ~~as that~~ from O<sub>3</sub> photolysis [e.g., (Yang et al., 2002)] ~~while the others seem to have~~  
3 ~~different results~~ [e.g., (Chen et al., 2007)]. ~~Although the photolysis of these species~~  
4 ~~are more sensitive to UV-A radiation (Grannas et al., 2007), the production/emission~~  
5 ~~of HONO and CH<sub>2</sub>O from the snowpack are influenced by snow nitrate photolysis~~  
6 ~~(Thomas et al., 2012) which is sensitive to UV-B radiation. In addition, a~~ recent  
7 model study ~~by Thomas et al.~~ (2012) suggested that at Summit, snow-sourced NO<sub>x</sub>  
8 ~~leads to as much OH production as that from ozone photolysis by photolysis of snow~~  
9 ~~nitrate also contributes to the local OH concentration (20-50%)~~ via shifting the local  
10 OH/HO<sub>2</sub> ratio in favor of OH, ~~with additional 10-18 % contribution from snowpack~~  
11 ~~bromine chemistry.~~ These processes are also sensitive to UV-B radiation. These  
12 ~~model studies suggested that snow photochemistry is more important than ozone~~  
13 ~~photolysis for OH production at Summit~~ (Thomas et al., 2012; Yang et al., 2002). ~~The~~  
14 ~~production of OH via CH<sub>2</sub>O, HONO and H<sub>2</sub>O<sub>2</sub> photolysis is also strongly influenced~~  
15 ~~by UV-B.~~ Therefore, ~~i~~ It is thus conceivable that under the condition of reduced O<sub>3</sub>  
16 column density in the spring of 2005, the OH production rate could be increased by  
17 200 % from the combined contribution of enhanced snow photochemistry OH  
18 ~~precursors from the snowpack and enhanced~~ tropospheric O<sub>3</sub> photolysis, even  
19 without considering the effect of elevated water vapor mixing ratio due to the  
20 temperature increase (around 3 °C, but this will enhance the emissions of H<sub>2</sub>O<sub>2</sub>, CH<sub>2</sub>O  
21 from snow (Grannas et al., 2007)).

22 Thus, we propose that the elevated tropospheric UV-B level due to a weakened  
23 stratospheric ozone layer in the spring of 2005 likely led to additional local nitrate  
24 production in sufficient quantities to account for the spring nitrate peak. The

1 additional nitrate was generated primarily through increased tropospheric OH  
2 production from the enhanced photolysis of tropospheric O<sub>3</sub> and snowpack  
3 photochemistry involving NO<sub>x</sub>, halogens, HONO, CH<sub>2</sub>O and H<sub>2</sub>O<sub>2</sub> and HONO  
4 emitted from the snowpack, and also possibly from enhanced Increased NO<sub>x</sub>  
5 abundance emission from the photolysis of snowpack nitrate and/or other NO<sub>x</sub>  
6 precursors that accumulated during the polar winter also contributes (Kramer et al.,  
7 2014). ~~The enhanced snowpack NO<sub>x</sub> emission, contributed to the spring nitrate peak~~  
8 ~~by enhancing the local NO<sub>x</sub> source and by increasing OH production rate.~~

#### 9 **4.3. Further examination Justification with additional snow and ice core data**

10 Hastings et al. (2004) reported seasonal variations in concentrations,  $\delta^{18}\text{O}$  and  
11  $\delta^{15}\text{N}$  of snow nitrate at Summit covering the years of 2000 and 2001. Qualitatively,  
12  $\delta^{18}\text{O}(\text{NO}_3^-)$  is similar to  $\Delta^{17}\text{O}(\text{NO}_3^-)$ , because the  $\delta^{18}\text{O}(\text{NO}_3^-)$  value is also determined  
13 by the relative importance of O<sub>3</sub> versus OH oxidation (higher  $\delta^{18}\text{O}(\text{NO}_3^-)$  values are  
14 associated with increased O<sub>3</sub> oxidation and lower  $\delta^{18}\text{O}(\text{NO}_3^-)$  values indicate  
15 increased HO<sub>x</sub> oxidation) (Hastings et al., 2004). Consequently,  $\delta^{18}\text{O}(\text{NO}_3^-)$  is high in  
16 winter snow and low in summer snow (as shown in Figure 1a, and also in Hastings et  
17 al. (2004)). The data shown in Hastings et al. (2004) indicated a spring nitrate peak in  
18 2000. The spring nitrate concentration peak in 2000 in Hastings et al. (2004) had  
19  $\delta^{18}\text{O}(\text{NO}_3^-)$  values ( $(69.8 \pm 2.1) \text{‰}$ ) similar to those during the summer of 2000 ( $(70.5$   
20  $\pm 2.4) \text{‰}$ ). In contrast, in the 2001 spring when no nitrate concentration peak was  
21 observed,  $\delta^{18}\text{O}(\text{NO}_3^-)$  was  $(77.5 \pm 2.4) \text{‰}$ , which is similar to the  $\delta^{18}\text{O}(\text{NO}_3^-)$  values  
22 observed during the prior winter ( $(77.4 \pm 1.9) \text{‰}$ ), and higher than those observed  
23 during the subsequent summer ( $(68.9 \pm 2.1) \text{‰}$ ). In addition, the mean  $\delta^{15}\text{N}(\text{NO}_3^-)$  in  
24 the spring of 2000 ( $(5.9 \pm 6.2) \text{‰}$ ) from Hastings et al. (2004) is similar to that



1 observed in the spring of 2005 ( $6.4 \pm 2.1$ ) ‰, and is higher than that observed in  
2 Hastings et al. (2004) during the spring of 2001 ( $-1.4 \pm 3.0$ ) ‰. In summary, the  
3 isotopic features of nitrate associated with the spring peaks observed in 2000  
4 (Hastings et al., 2004) and in 2005 are similar, each suggesting enhanced local  
5 photochemistry as a contributor to the observed spring nitrate peaks. This is consistent  
6 with the low O<sub>3</sub> column density in the spring of 2000 (337 DU, Figure 3) and 2005  
7 (294 DU, Figure 3). Therefore, the results of Hastings et al. (2004) support the  
8 explanation that the appearance of the spring nitrate peak in 2000 is caused by a  
9 weakened stratospheric ozone layer, consistent with the observations in the spring of  
10 2005.

11 We further examined the 19 years with double nitrate peaks found in the firn  
12 core. In order to assess the effect of long range transport of nitrate on the occurrence  
13 of the double nitrate peaks, we compared the nitrate sulfate records. This comparison  
14 indicates that in most (14 out of 19) years, the secondary (earlier and smaller) peak is  
15 accompanied by a sulfate peak. This suggests that direct transport of nitrate from  
16 polluted regions is likely responsible for the occurrence of the double nitrate peaks in  
17 these years. However, there are five years, including 2000 and 2005, in which the  
18 secondary peak does not appear with a rise in sulfate concentration. To determine if  
19 enhanced local photochemistry could be responsible for the occurrence of the double  
20 nitrate peaks in these years, we ~~further determine whether the above explanation is~~  
21 ~~representative of the occasional nature of the spring nitrate peak observed in modern~~  
22 ~~snow in Central Greenland, we ~~compared~~ examined the nitrate concentration record~~  
23 ~~from the shallow ice core with~~ O<sub>3</sub> column density data from 1979 to 2006, the time  
24 period when global O<sub>3</sub> data are available from satellite observations. The year-to-year

1 variability of polar stratospheric O<sub>3</sub> is largely controlled by the Brewer-Dobson  
2 circulation (BDC) through direct transport and indirect coupling between dynamics  
3 and chemistry [e.g., (Holton et al., 1995; Randel et al., 2002; Shepherd, 2008; Weber  
4 et al., 2011)]. Halogen-catalyzed chemical destruction leads to a decreasing trend in  
5 column O<sub>3</sub> density since 1980 (WMO, 2007) and causes sudden drops in O<sub>3</sub> column  
6 density in years when the winter temperatures are anomalously low [e.g., the winter of  
7 2004/2005 (Jin et al., 2006; Kleinbohl et al., 2005)]. The strength of the BDC is  
8 related to the wave force, which is represented by extratropical poleward eddy heat  
9 flux in the lower stratosphere (Fusco and Salby, 1999; Weber et al., 2011). From the  
10 NCEP/NCAR reanalysis data  
11 (<http://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.pressure.html>), we  
12 retrieved the average eddy heat flux in the lower stratosphere at 40-80 °N in January  
13 and February (due to the transport lag, January and February eddy heat flux  
14 determines the February and March polar stratospheric O<sub>3</sub> abundance) from 1979 to  
15 2006 (Figure 3). Reanalysis data before 1979 are also available, but less reliable due  
16 to lack of constraints by satellite observations. Therefore in this study, we only use the  
17 data after 1979.

18 — As shown in Figure 3, in the period from 1979 to 2006, there are 12 years with  
19 double nitrate peaks and five of them are years with the spring nitrate peak only  
20 appears when O<sub>3</sub> column density well below is near or below the average for that  
21 period. In other years, one of the double peaks is accompanied by a sulfate peak. The  
22 results from the firn core, combined with that from the snowpit, thus suggest that  
23 there are two mechanisms leading to occurrence of double nitrate peaks, 1) direct  
24 transport of nitrate from polluted regions, and 2) enhanced local photochemical

1 ~~production of nitrate. We note that although spring nitrate concentration peaks are~~  
2 ~~observed in all years with especially low (lower than the average) O<sub>3</sub> column density,~~  
3 ~~there are a few years (3 out of 12) with a spring nitrate peak when O<sub>3</sub> column density~~  
4 ~~is near the 1979–2006 average. In addition, there are other years with similarly~~  
5 ~~average O<sub>3</sub> column density when no spring nitrate peak is observed. We suggest that~~  
6 ~~this is because the presence of the spring nitrate peak also requires other conditions~~  
7 ~~(e.g., sufficient local NO<sub>x</sub> abundance), and that this is especially true when O<sub>3</sub> column~~  
8 ~~density is near the 1979–2006 average. Additional factors may also be important, such~~  
9 ~~as stratospheric input of nitrate and/or long range transport of nitrate, as discussed~~  
10 ~~previously. With isotopic data and other relevant information available, we were able~~  
11 ~~to exclude many possible sources of nitrate to the spring peak in 2005; however, it is~~  
12 ~~difficult to explicitly assess the dominant source(s) of nitrate contributing to the~~  
13 ~~spring peak in each individual year observed in the shallow ice core, mainly due to the~~  
14 ~~lack of isotopic data. It is possible that episodic events bring sufficient nitrate so that a~~  
15 ~~spring peak is detected, when local photochemistry is not significantly enhanced.~~

16 ~~——The second mechanism requires Possible local sources of NO<sub>x</sub> sources. Possible~~  
17 ~~local NO<sub>x</sub> sources~~ -at Summit include PAN decomposition and the photolysis of  
18 snowpack nitrate ~~photolysis, as well as photolysis of other winter accumulated NO<sub>y</sub>~~  
19 ~~species in the atmosphere~~ (Kramer et al., 2014). ~~For example, Aa~~ model study (Stroud  
20 et al., 2003) ~~also~~ suggested that HNO<sub>4</sub> is a source of NO<sub>x</sub> in remote regions through  
21 recycling reactions, but its importance is unclear due to a lack of field observations. ~~In~~  
22 ~~spring, PAN acts as a sink of NO<sub>x</sub>, which leaves the snowpack nitrate photolysis as~~  
23 ~~the most likely local source of NO<sub>x</sub>. As shown in Figure 2a, snow nitrate~~  
24 ~~concentrations at Summit began to increase around 1950 due to increasing~~

1 anthropogenic NO<sub>x</sub> emissions in the mid-latitudes (Fischer et al., 1998), and reached  
2 and maintained the highest level from 1970 to the present. Since snow nitrate can be  
3 photolyzed, releasing NO<sub>x</sub> to the boundary layer, the increase in snow nitrate  
4 concentrations represents an increase in a potentially important NO<sub>x</sub> reservoir in  
5 Greenland. These NO<sub>x</sub> precursors accumulate in the polar winter through a  
6 combination of pollution transport from the mid-latitudes and lack of sunlight, and  
7 Anthropogenic NO<sub>x</sub> emissions also increase other reservoir species that may  
8 contribute to springtime NO<sub>x</sub> abundance through photochemical decomposition after  
9 polar sunrise (Kramer et al., 2014). at Summit. This is consistent with the observation  
10 that there are no double nitrate peaks observed prior to the 1950s as dramatic  
11 increases in anthropogenic emissions of NO<sub>x</sub> did not occur until ~1950 (Lamarque et  
12 al., 2010). Prior to the 1950s, when local NO<sub>x</sub> abundance was not sufficiently high, no  
13 spring nitrate concentration peaks were detected.

## 14 5. Conclusion

15 Spring nitrate peaks have been observed in Greenland ice core records beginning  
16 ~1900 and was hypothesized to originate from pollution transport from the mid-  
17 latitudes in the industrial era (Burkhart et al., 2006; Whitlow et al., 1992; Yang et al.,  
18 1995). But the specific mechanism leading to the additional peak is unclear as it could  
19 be from direct transport of nitrate or transport of NO<sub>x</sub> precursors followed by  
20 enhanced local photochemistry. Here we report observations of nitrate concentrations  
21 and isotopes in a snowpit from Summit, Greenland. After identifying a spring nitrate  
22 peak in the year of 2005, we performed a case study regarding its origin. The isotopic  
23 composition of nitrate in the snowpit, combined with photochemical calculations,  
24 suggests the presence of the 2005 spring nitrate peak was caused by enhanced local

1 nitrate production, instead of long range pollution transport. An analysis of the  
2 possible causes suggests that this enhancement was primarily due to increased OH  
3 concentration mainly resulting from enhanced photolysis of OH precursors (O<sub>3</sub>,  
4 HONO, H<sub>2</sub>O<sub>2</sub>, etc.) in the troposphere caused by elevated UV-B radiation at the  
5 surface. Elevated UV-B radiation at the surface was the result of a significant  
6 reduction of the stratospheric ozone layer in the Arctic in spring 2005. Increased local  
7 NO<sub>x</sub> concentrations due to enhanced snowpack nitrate photolysis may also contribute  
8 to up to half of the additional nitrate in the spring peak, if snowpack nitrate photolysis  
9 is the dominant local NO<sub>x</sub> source. Either the increase in OH concentration or the  
10 increase in local NO<sub>x</sub> concentration relies on enhanced photochemical reactions  
11 initiated by the reduced O<sub>3</sub> column density. This mechanism is supported by the  
12 appearance of a spring nitrate peak in 2000 (Hastings et al., 2004), when a reduced O<sub>3</sub>  
13 column density also occurred.

14 Further examination on a firn core indicates that in most years direct transport of  
15 nitrate from polluted regions is likely responsible for the additional nitrate peak. This,  
16 combined with the snowpit results, suggests that there are two mechanisms for the  
17 occurrence of the additional annual nitrate peak in the industrial era, 1) direct  
18 transport of nitrate from polluted regions, and 2) enhanced local photochemical  
19 production of nitrate. We note that both of these mechanisms are related to pollution  
20 transport, as the additional nitrate from either direct transport or enhanced local  
21 photochemistry requires enhanced nitrogen sources from anthropogenic emissions. In  
22 addition, we examined the nitrate concentration record from a shallow ice core from  
23 Summit. Years with two nitrate peaks are frequently observed after 1950. Although  
24 we did not conduct sub-annual layer dating of this core, the practice of referring to the

1 additional nitrate peak as a spring peak is both consistent with previous studies  
2 (Burkhart et al., 2006; Whitlow et al., 1992; Yang et al., 1995), and with our sub-  
3 annual layer dating of the snowpit. Most of the years with two nitrate peaks observed  
4 in snow layers after 1979 are also years with significantly low spring ozone column  
5 density, consistent with our explanation of the nitrate peak in spring 2005.

6 ——— Based on the above evidence, we present an alternative hypothesis that the  
7 presence of the occasional spring nitrate peak in Greenland snow is largely associated  
8 with the interannual variability of  $O_3$  column density in the Arctic. The interannual  
9 variability of  $O_3$  column density is mainly controlled by the Brewer-Dobson  
10 circulation, while chemical destruction may also play a role beginning around 1980  
11 (WMO, 2007). The presence of the spring nitrate peak, however, also requires  
12 sufficient  $NO_x$  at Summit. Local  $NO_x$  sources likely increased dramatically after the  
13 1950s due to increases in anthropogenic  $NO_x$  emissions in the mid-latitudes, leading  
14 to a buildup of  $NO_x$  reservoir species (PAN and snow nitrate) at Summit. Seasonally  
15 resolved isotopic data ( $\delta^{17}O(NO_3^-)$  and  $\delta^{15}N(NO_3^-)$ ) covering additional years with  
16 spring nitrate concentration peaks are needed to further examine how robust the  
17 connection is between the spring nitrate peak and  $O_3$  column density. Although it is  
18 difficult to conduct such efforts with ice core measurements because of the large  
19 sample requirements ( $> 50$  g ice), measurements of snowpit samples can be  
20 performed. Snow samples from Summit covering the years of 2010 to 2012 may be  
21 ideal for this purpose because there is unprecedented  $O_3$  loss in the spring of 2011 in  
22 Arctic (Manney et al., 2011).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

## 6. Acknowledgments

Total column O<sub>3</sub> data are from <http://jwocky.gsfc.nasa.gov/eptoms/ep.html>, by Earth Probe Total Ozone Mapping Spectrometer, NASA. We thank Ice Drilling Design and Operations (formerly Ice Coring and Drilling Services), University of Wisconsin for field assistance in drilling the Summit ice cores. A. L. Lanciki and D. G. Ferris of South Dakota State University performed the continuous analysis of the shallow ice core. We also thank J. Bautista of University of Washington for assistance in isotopic measurements and acknowledge the support from the Quaternary Research Center at University of Washington. This work is financially supported by the NSF (Awards 0612461 and 0839066 to J. Cole-Dai, PLR 0944537 and PLR 1106317 to B. Alexander, and 0538049 and 0454803 to E. J. Steig). JS and JE has been supported by a grant from Labex OSUG@2020 (Investissements d’avenir – ANR10 LABX56). The French Polar Institute IPEV is acknowledged for its financial and logistical support to JS (programme 457). LEFE-IMAGO, a scientific program of the Institute National des Sciences de l’Univers (INSU/CNRS), has also provided partial funding for JS.

1

## 2 **References**

- 3 Alexander, B., Savarino, J., Kreutz, K. J., and Thiemens, M. H.: Impact of  
4 preindustrial biomass-burning emissions on the oxidation pathways of tropospheric  
5 sulfur and nitrogen, *J. Geophys. Res.*, 109, D08303, 10.1029/2003jd004218, 2004.
- 6 Alexander, B., Hastings, M. G., Allman, D. J., Dachs, J., Thornton, J. A., and  
7 Kunasek, S. A.: Quantifying atmospheric nitrate formation pathways based on a  
8 global model of the oxygen isotopic composition ( $\Delta^{17}\text{O}$ ) of atmospheric nitrate,  
9 *Atmos. Chem. Phys.*, 9, 5043-5056, 10.5194/acp-9-5043-2009, 2009.
- 10 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R.  
11 G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data  
12 for atmospheric chemistry: Volume II - gas phase reactions of organic species, *Atmos.*  
13 *Chem. Phys.*, 6, 3625-4055, 10.5194/acp-6-3625-2006, 2006.
- 14 Beine, H. J., and Krognes, T.: The seasonal cycle of peroxyacetyl nitrate (PAN) in the  
15 European Arctic, *Atmos. Environ.*, 34, 933-940, 10.1016/S1352-2310(99)00288-5,  
16 2000.
- 17 Berhanu, T. A., Meusinger, C., Erbland, J., Jost, R., Bhattacharya, S. K., Johnson, M.  
18 S., and Savarino, J.: Laboratory study of nitrate photolysis in Antarctic snow. II.  
19 Isotopic effects and wavelength dependence, *The Journal of Chemical Physics*, 140,  
20 244306, doi:<http://dx.doi.org/10.1063/1.4882899>, 2014.
- 21 Bhattacharya, S. K., Pandey, A., and Savarino, J.: Determination of intramolecular  
22 isotope distribution of ozone by oxidation reaction with silver metal, *J. Geophys.*  
23 *Res.*, 113, D03303, 10.1029/2006jd008309, 2008.
- 24 Bottenheim, J. W., Barrie, L. A., and Atlas, E.: The Partitioning of Nitrogen-Oxides  
25 in the Lower Arctic Troposphere during Spring 1988, *J. Atmos. Chem.*, 17, 15-27,  
26 10.1007/BF00699111, 1993.
- 27 Burkhart, J. F., Bales, R. C., McConnell, J. R., and Hutterli, M. A.: Influence of North  
28 Atlantic Oscillation on anthropogenic transport recorded in northwest Greenland ice  
29 cores, *J. Geophys. Res.*, 111, D22309, 10.1029/2005jd006771, 2006.
- 30 Chen, G., Huey, L. G., Crawford, J. H., Olson, J. R., Hutterli, M. A., Sjostedt, S.,  
31 Tanner, D., Dibb, J., Lefer, B., Blake, N., Davis, D., and Stohl, A.: An assessment of  
32 the polar HO<sub>x</sub> photochemical budget based on 2003 Summit Greenland field  
33 observations, *Atmos. Environ.*, 41, 7806-7820,  
34 <http://dx.doi.org/10.1016/j.atmosenv.2007.06.014>, 2007.
- 35 Chu, L., and Anastasio, C.: Quantum yields of hydroxyl radical and nitrogen dioxide  
36 from the photolysis of nitrate on ice, *J. Phys. Chem. A*, 107, 9594-9602,  
37 10.1021/Jp0349132, 2003.
- 38 Cole-Dai, J., Budner, D. M., and Ferris, D. G.: High speed, high resolution, and  
39 continuous chemical analysis of ice cores using a melter and ion chromatography,  
40 *Environ. Sci. Technol.*, 40, 6764-6769, 10.1021/Es061188a, 2006.
- 41 Cole-Dai, J., Ferris, D. G., Lanciki, A. L., Savarino, J., Thiemens, M. H., and  
42 McConnell, J. R.: Two likely stratospheric volcanic eruptions in the 1450s C.E. found



1 in a bipolar, subannually dated 800 year ice core record, *J. Geophys. Res.*, 118, 7459-  
2 7466, 10.1002/jgrd.50587, 2013.

3 Dassau, T. M., Sumner, A. L., Koeniger, S. L., Shepson, P. B., Yang, J., Honrath, R.  
4 E., Cullen, N. J., Steffen, K., Jacobi, H. W., Frey, M., and Bales, R. C.: Investigation  
5 of the role of the snowpack on atmospheric formaldehyde chemistry at Summit,  
6 Greenland, *J. Geophys. Res.*, 107, 4394, 10.1029/2002jd002182, 2002.

7 Davidson, C. I., Harrington, J. R., Stephenson, M. J., Small, M. J., Boscoe, F. P., and  
8 Gandley, R. E.: Seasonal variations in sulfate, nitrate and chloride in the Greenland  
9 ice sheet: relation to atmospheric concentrations, *Atmos. Environ.*, 23, 2483-2493,  
10 10.1016/0004-6981(89)90259-X, 1989.

11 Eckhardt, S., Stohl, A., Beirle, S., Spichtinger, N., James, P., Forster, C., Junker, C.,  
12 Wagner, T., Platt, U., and Jennings, S. G.: The North Atlantic Oscillation controls air  
13 pollution transport to the Arctic, *Atmos. Chem. Phys.*, 3, 1769-1778, 10.5194/acp-3-  
14 1769-2003, 2003.

15 Erbland, J., Vicars, W. C., Savarino, J., Morin, S., Frey, M. M., Frosini, D., Vince, E.,  
16 and Martins, J. M. F.: Air-snow transfer of nitrate on the East Antarctic Plateau - Part  
17 1: Isotopic evidence for a photolytically driven dynamic equilibrium, *Atmos. Chem.*  
18 *Phys.*, 13, 6403-6419, 10.5194/acp-13-6403-2013, 2013.

19 Fibiger, D. L., Hastings, M. G., Dibb, J. E., and Huey, L. G.: The preservation of  
20 atmospheric nitrate in snow at Summit, Greenland, *Geophys. Res. Lett.*, 40, 3484-  
21 3489, 10.1002/grl.50659, 2013.

22 Finkel, R. C., and Langway, C. C.: Changes in Precipitation chemistry at Dye 3,  
23 Greenland, *J. Geophys. Res.*, 91, 9849-9855, 10.1029/JD091iD09p09849, 1986.

24 Fischer, H., Wagenbach, D., and Kipfstuhl, J.: Sulfate and nitrate firn concentrations  
25 on the Greenland ice sheet 2. Temporal anthropogenic deposition changes, *J.*  
26 *Geophys. Res.*, 103, 21935-21942, 10.1029/98JD01886, 1998.

27 Flowers, B. A., Angerhofer, M. E., Simpson, W. R., Nakayama, T., and Matsumi, Y.:  
28 Nitrate radical quantum yield from peroxyacetyl nitrate photolysis, *J. Phys. Chem. A*,  
29 109, 2552-2558, 10.1021/Jp045529n, 2005.

30 Ford, K. M., Campbell, B. M., Shepson, P. B., Bertman, S. B., Honrath, R. E.,  
31 Peterson, M., and Dibb, J. E.: Studies of Peroxyacetyl nitrate (PAN) and its  
32 interaction with the snowpack at Summit, Greenland, *J. Geophys. Res.*, 107, 4102,  
33 10.1029/2001jd000547, 2002.

34 France, J. L., King, M. D., Frey, M. M., Erbland, J., Picard, G., Preunkert, S.,  
35 MacArthur, A., and Savarino, J.: Snow optical properties at Dome C (Concordia),  
36 Antarctica; implications for snow emissions and snow chemistry of reactive nitrogen,  
37 *Atmos. Chem. Phys.*, 11, 9787-9801, 10.5194/acp-11-9787-2011, 2011.

38 Frey, M. M., Savarino, J., Morin, S., Erbland, J., and Martins, J. M. F.: Photolysis  
39 imprint in the nitrate stable isotope signal in snow and atmosphere of East Antarctica  
40 and implications for reactive nitrogen cycling, *Atmos. Chem. Phys.*, 9, 8681-8696,  
41 DOI 10.5194/acp-9-8681-2009, 2009.

- 1 Fusco, A. C., and Salby, M. L.: Interannual variations of total ozone and their  
2 relationship to variations of planetary wave activity, *J Climate*, 12, 1619-1629,  
3 10.1175/1520-0442, 1999.
- 4 Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W.,  
5 Cowling, E. B., and Cosby, B. J.: The nitrogen cascade, *Bioscience*, 53, 341-356,  
6 2003.
- 7 Geng, L., Alexander, B., Cole-Dai, J., Steig, E. J., Savarino, J., Sofen, E. D., and  
8 Schauer, A. J.: Nitrogen isotopes in ice core nitrate linked to anthropogenic  
9 atmospheric acidity change, *Proceedings of the National Academy of Sciences*, 111,  
10 5808-5812, 10.1073/pnas.1319441111, 2014.
- 11 Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and  
12 annual burned area using the fourth-generation global fire emissions database  
13 (GFED4), *Journal of Geophysical Research: Biogeosciences*, 118, 317-328,  
14 10.1002/jgrg.20042, 2013.
- 15 Grannas, A. M., Jones, A. E., Dibb, J., Ammann, M., Anastasio, C., Beine, H. J.,  
16 Bergin, M., Bottenheim, J., Boxe, C. S., Carver, G., Chen, G., Crawford, J. H.,  
17 Domine, F., Frey, M. M., Guzman, M. I., Heard, D. E., Helmig, D., Hoffmann, M. R.,  
18 Honrath, R. E., Huey, L. G., Hutterli, M., Jacobi, H. W., Klan, P., Lefer, B.,  
19 McConnell, J., Plane, J., Sander, R., Savarino, J., Shepson, P. B., Simpson, W. R.,  
20 Sodeau, J. R., von Glasow, R., Weller, R., Wolff, E. W., and Zhu, T.: An overview of  
21 snow photochemistry: evidence, mechanisms and impacts, *Atmos. Chem. Phys.*, 7,  
22 4329-4373, 10.5194/acp-7-4329-2007, 2007.
- 23 Hastings, M. G., Steig, E. J., and Sigman, D. M.: Seasonal variations in N and O  
24 isotopes of nitrate in snow at Summit, Greenland: Implications for the study of nitrate  
25 in snow and ice cores, *J. Geophys. Res.*, 109, D20306, 10.1029/2004jd004991, 2004.
- 26 Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and  
27 Pfister, L.: Stratosphere-Troposphere Exchange, *Rev. Geophys.*, 33, 403-439, Doi  
28 10.1029/95rg02097, 1995.
- 29 Honrath, R. E., and Jaffe, D. A.: The Seasonal Cycle of Nitrogen-Oxides in the Arctic  
30 Troposphere at Barrow, Alaska, *J. Geophys. Res.*, 97, 20615-20630, 1992.
- 31 Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell,  
32 B.: Evidence of NO<sub>x</sub> production within or upon ice particles in the Greenland  
33 snowpack, *Geophys. Res. Lett.*, 26, 695-698, 10.1029/1999GL900077, 1999.
- 34 Hutterli, M. A., McConnell, J. R., Stewart, R. W., Jacobi, H. W., and Bales, R. C.:  
35 Impact of temperature-driven cycling of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) between air and  
36 snow on the planetary boundary layer, *J. Geophys. Res.*, 106, 15395-15404,  
37 10.1029/2001JD900102, 2001.
- 38 Jacobi, H. W., Weller, R., Bluszczyk, T., and Schrems, O.: Latitudinal distribution of  
39 peroxyacetyl nitrate (PAN) over the Atlantic Ocean, *J. Geophys. Res.*, 104, 26901-  
40 26912, 10.1029/1999JD900462, 1999.
- 41 Jacobi, H. W., Bales, R. C., Honrath, R. E., Peterson, M. C., Dibb, J. E., Swanson, A.  
42 L., and Albert, M. R.: Reactive trace gases measured in the interstitial air of surface

1 snow at Summit, Greenland, *Atmos. Environ.*, 38, 1687-1697,  
2 10.1016/j.atmosenv.2004.01.004, 2004.

3 Jin, J. J., Semeniuk, K., Manney, G. L., Jonsson, A. I., Beagley, S. R., McConnell, J.  
4 C., Rinsland, C. P., Boone, C. D., Walker, K. A., and Bernath, P. F.: Denitrification in  
5 the Arctic winter 2004/2005: Observations from ACE-FTS, *Geophys. Res. Lett.*, 33,  
6 L19814, 10.1029/2006gl027687, 2006.

7 Johnson, D. G., Jucks, K. W., Traub, W. A., and Chance, K. V.: Isotopic composition  
8 of stratospheric ozone, *J. Geophys. Res.*, 105, 9025-9031, 10.1029/1999JD901167,  
9 2000.

10 Kaiser, J., Hastings, M. G., Houlton, B. Z., Rockmann, T., and Sigman, D. M.: Triple  
11 oxygen isotope analysis of nitrate using the denitrifier method and thermal  
12 decomposition of N<sub>2</sub>O, *Anal. Chem.*, 79, 599-607, 10.1021/Ac061022s, 2007.

13 Kleinbohl, A., Bremer, H., Kullmann, H., Kuttippurath, J., Browell, E. V., Canty, T.,  
14 Salawitch, R. J., Toon, G. C., and Notholt, J.: Denitrification in the Arctic mid-winter  
15 2004/2005 observed by airborne submillimeter radiometry, *Geophys. Res. Lett.*, 32,  
16 L19811, 10.1029/2005gl023408, 2005.

17 Kramer, L. J., Helmig, D., Burkhart, J. F., Stohl, A., Oltmans, S., and Honrath, R. E.:  
18 Seasonal variability of atmospheric nitrogen oxides and non-methane hydrocarbons at  
19 the GEOSummit station, Greenland, *Atmos. Chem. Phys. Discuss.*, 14, 13817-13867,  
20 10.5194/acpd-14-13817-2014, 2014.

21 Kunasek, S. A., Alexander, B., Steig, E. J., Hastings, M. G., Gleason, D. J., and  
22 Jarvis, J. C.: Measurements and modeling of  $\Delta^{17}\text{O}$  of nitrate in snowpits from  
23 Summit, Greenland, *J. Geophys. Res.*, 113, D24302, 10.1029/2008jd010103, 2008.

24 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D.,  
25 Liou, S. C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J.,  
26 Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N.,  
27 McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850-2000)  
28 gridded anthropogenic and biomass burning emissions of reactive gases and aerosols:  
29 methodology and application, *Atmos. Chem. Phys.*, 10, 7017-7039, 10.5194/acp-10-  
30 7017-2010, 2010.

31 Legrand, M. R., and Delmas, R. J.: Formation of HCl in the Antarctic atmosphere,  
32 *Journal of Geophysical Research: Atmospheres*, 93, 7153-7168,  
33 10.1029/JD093iD06p07153, 1988.

34 Levy, H., Moxim, W. J., Klonecki, A. A., and Kasibhatla, P. S.: Simulated  
35 tropospheric NO<sub>x</sub>: Its evaluation, global distribution and individual source  
36 contributions, *J. Geophys. Res.*, 104, 26279-26306, 10.1029/1999JD900442, 1999.

37 Liang, M. C., Irion, F. W., Weibel, J. D., Miller, C. E., Blake, G. A., and Yung, Y. L.:  
38 Isotopic composition of stratospheric ozone, *J. Geophys. Res.*, 111, D02302,  
39 10.1029/2005jd006342, 2006.

40 Logan, J. A.: Nitrogen oxides in the troposphere: Global and regional budgets, *J.*  
41 *Geophys. Res.*, 88, 10785-10807, 10.1029/JC088iC15p10785, 1983.

1 Lyons, J. R.: Transfer of mass-independent fractionation in ozone to other oxygen-  
2 containing radicals in the atmosphere, *Geophys. Res. Lett.*, 28, 3231-3234,  
3 10.1029/2000GL012791, 2001.

4 Mann, G. W., Davies, S., Carslaw, K. S., and Chipperfield, M. P.: Factors controlling  
5 Arctic denitrification in cold winters of the 1990s, *Atmos. Chem. Phys.*, 3, 403-416,  
6 10.5194/acp-3-403-2003, 2003.

7 Mauersberger, K., Lammerzahl, P., and Krankowsky, D.: Stratospheric ozone isotope  
8 enrichments-revisited, *Geophys. Res. Lett.*, 28, 3155-3158, 10.1029/2001GL013439,  
9 2001.

10 Mayewski, P. A., Lyons, W. B., Spencer, M. J., Twickler, M. S., Buck, C. F., and  
11 Whitlow, S.: An ice-core record of atmospheric response to anthropogenic sulphate  
12 and nitrate, *Nature*, 346, 554 - 556, 10.1038/346554a0, 1990.

13 McCabe, J. R., Thiemens, M. H., and Savarino, J.: A record of ozone variability in  
14 South Pole Antarctic snow: Role of nitrate oxygen isotopes, *J. Geophys. Res.*, 112,  
15 D12303, 10.1029/2006jd007822, 2007.

16 Meusinger, C., Berhanu, T. A., Erbland, J., Savarino, J., and Johnson, M. S.:  
17 Laboratory study of nitrate photolysis in Antarctic snow. I. Observed quantum yield,  
18 domain of photolysis, and secondary chemistry, *The Journal of Chemical Physics*,  
19 140, doi:<http://dx.doi.org/10.1063/1.4882898>, 2014.

20 Michalski, G., Scott, Z., Kabling, M., and Thiemens, M. H.: First measurements and  
21 modeling of  $\Delta^{17}\text{O}$  in atmospheric nitrate, *Geophys. Res. Lett.*, 30, 1870,  
22 10.1029/2003gl017015, 2003.

23 Morin, S., Savarino, J., Bekki, S., Gong, S., and Bottenheim, J. W.: Signature of  
24 Arctic surface ozone depletion events in the isotope anomaly ( $\Delta^{17}\text{O}$ ) of atmospheric  
25 nitrate, *Atmos. Chem. Phys.*, 7, 1451-1469, 10.5194/acp-7-1451-2007, 2007.

26 Morin, S., Savarino, J., Frey, M. M., Yan, N., Bekki, S., Bottenheim, J. W., and  
27 Martins, J. M. F.: Tracing the Origin and Fate of  $\text{NO}_x$  in the Arctic Atmosphere Using  
28 Stable Isotopes in Nitrate, *Science*, 322, 730-732, 10.1126/science.1161910, 2008.

29 Morin, S., Sander, R., and Savarino, J.: Simulation of the diurnal variations of the  
30 oxygen isotope anomaly ( $\Delta^{17}\text{O}$ ) of reactive atmospheric species, *Atmos. Chem.*  
31 *Phys.*, 11, 3653-3671, 10.5194/acp-11-3653-2011, 2011.

32 Moxim, W. J., Levy, H., and Kasibhatla, P. S.: Simulated global tropospheric PAN:  
33 Its transport and impact on  $\text{NO}_x$ , *J. Geophys. Res.*, 101, 12621-12638,  
34 10.1029/96JD00338, 1996.

35 Mulvaney, R., and Wolff, E. W.: Evidence for Winter Spring Denitrification of the  
36 Stratosphere in the Nitrate Record of Antarctic Firn Cores, *J. Geophys. Res.*, 98,  
37 5213-5220, 10.1029/92JD02966, 1993.

38 Osborn, T. J.: Winter 2009/2010 temperatures and a record-breaking North Atlantic  
39 Oscillation index, *Weather*, 66, 19-21, 10.1002/Wea.660, 2011.

40 Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and  
41 transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the  
42 United States: Implications for policy, *J. Geophys. Res.*, 109, 10.1029/2003jd004473,  
43 2004.

1 Pye, H. O. T., Liao, H., Wu, S., Mickley, L. J., Jacob, D. J., Henze, D. K., and  
2 Seinfeld, J. H.: Effect of changes in climate and emissions on future sulfate-nitrate-  
3 ammonium aerosol levels in the United States, *J. Geophys. Res.*, 114,  
4 10.1029/2008jd010701, 2009.

5 Randel, W. J., Wu, F., and Stolarski, R.: Changes in column ozone correlated with the  
6 stratospheric EP flux, *J Meteorol Soc Jpn*, 80, 849-862, 10.2151/Jmsj.80.849, 2002.

7 Röhlisberger, R., Hutterli, M. A., Wolff, E. W., Mulvaney, R., Fischer, H., Bigler,  
8 M., Goto-Azuma, K., Hansson, M. E., Ruth, U., Siggaard-Andersen, M. L., and  
9 Steffensen, J. P.: Nitrate in Greenland and Antarctic ice cores: a detailed description  
10 of post-depositional processes, *Ann. Glaciol.*, 35, 209-216,  
11 10.3189/172756402781817220, 2002.

12 Savarino, J., and Legrand, M.: High northern latitude forest fires and vegetation  
13 emissions over the last millennium inferred from the chemistry of a central Greenland  
14 ice core, *J. Geophys. Res.*, 103, 8267-8279, 10.1029/97JD03748, 1998.

15 Savarino, J., Kaiser, J., Morin, S., Sigman, D. M., and Thiemens, M. H.: Nitrogen and  
16 oxygen isotopic constraints on the origin of atmospheric nitrate in coastal Antarctica,  
17 *Atmos. Chem. Phys.*, 7, 1925-1945, 10.5194/acpd-6-8817-2006, 2007.

18 Shepherd, T. G.: Dynamics, stratospheric ozone, and climate change, *Atmos Ocean*,  
19 46, 117-138, 10.3137/Ao.460106, 2008.

20 Singh, H. B., Ohara, D., Herlth, D., Bradshaw, J. D., Sandholm, S. T., Gregory, G. L.,  
21 Sachse, G. W., Blake, D. R., Crutzen, P. J., and Kanakidou, M. A.: Atmospheric  
22 Measurements of Peroxyacetyl Nitrate and Other Organic Nitrates at High-Latitudes:  
23 Possible Sources and Sinks, *J. Geophys. Res.*, 97, 16511-16522, 1992.

24 Sjostedt, S. J., Huey, L. G., Tanner, D. J., Peischl, J., Chen, G., Dibb, J. E., Lefer, B.,  
25 Hutterli, M. A., Beyersdorf, A. J., Blake, N. J., Blake, D. R., Sueper, D., Ryerson, T.,  
26 Burkhardt, J., and Stohl, A.: Observations of hydroxyl and the sum of peroxy radicals  
27 at Summit, Greenland during summer 2003, *Atmos. Environ.*, 41, 5122-5137,  
28 10.1016/j.atmosenv.2006.06.065, 2007.

29 Stroud, C., Madronich, S., Atlas, E., Ridley, B., Flocke, F., Weinheimer, A., Talbot,  
30 B., Fried, A., Wert, B., Shetter, R., Lefer, B., Coffey, M., Heikes, B., and Blake, D.:  
31 Photochemistry in the arctic free troposphere: NO<sub>x</sub> budget and the role of odd  
32 nitrogen reservoir recycling, *Atmos. Environ.*, 37, 3351-3364, 10.1016/S1352-  
33 2310(03)00353-4, 2003.

34 Talukdar, R. K., Burkholder, J. B., Schmoltner, A. M., Roberts, J. M., Wilson, R. R.,  
35 and Ravishankara, A. R.: Investigation of the Loss Processes for Peroxyacetyl Nitrate  
36 in the Atmosphere - UV Photolysis and Reaction with OH, *J. Geophys. Res.*, 100,  
37 14163-14173, 10.1029/95JD00545, 1995.

38 Thomas, J. L., Dibb, J. E., Huey, L. G., Liao, J., Tanner, D., Lefer, B., von Glasow,  
39 R., and Stutz, J.: Modeling chemistry in and above snow at Summit, Greenland - Part  
40 2: Impact of snowpack chemistry on the oxidation capacity of the boundary layer,  
41 *Atmos. Chem. Phys.*, 12, 6537-6554, 10.5194/acp-12-6537-2012, 2012.

42 Waugh, D. W., and Randel, W. J.: Climatology of arctic and antarctic polar vortices  
43 using elliptical diagnostics, *J. Atmos. Sci.*, 56, 1594-1613, 1999.

1 Weber, M., Dikty, S., Burrows, J. P., Garny, H., Dameris, M., Kubin, A., Abalichin,  
2 J., and Langematz, U.: The Brewer-Dobson circulation and total ozone from seasonal  
3 to decadal time scales, *Atmos. Chem. Phys.*, 11, 11221-11235, 10.5194/acp-11-  
4 11221-2011, 2011.

5 Whitlow, S., Mayewski, P. A., and Dibb, J. E.: A Comparison of Major Chemical-  
6 Species Seasonal Concentration and Accumulation at the South-Pole and Summit,  
7 Greenland, *Atmos. Environ.*, 26, 2045-2054, 10.1016/0960-1686(92)90089-4, 1992.

8 WMO: Scientific assessment of ozone depletion: 2006, Geneva50, 572, 2007.

9 Yang, J., Honrath, R. E., Peterson, M. C., Dibb, J. E., Sumner, A. L., Shepson, P. B.,  
10 Frey, M., Jacobi, H. W., Swanson, A., and Blake, N.: Impacts of snowpack emissions  
11 on deduced levels of OH and peroxy radicals at Summit, Greenland, *Atmos. Environ.*,  
12 36, 2523-2534, 10.1016/S1352-2310(02)00128-0, 2002.

13 Yang, Q. Z., Mayewski, P. A., Whitlow, S., Twickler, M., Morrison, M., Talbot, R.,  
14 Dibb, J., and Linder, E.: Global Perspective of Nitrate Flux in Ice Cores, *J. Geophys.*  
15 *Res.*, 100, 5113-5121, 10.1029/94JD03115, 1995.

16 Zatkan, M. C., Grenfell, T. C., Alexander, B., Doherty, S. J., Thomas, J. L., and Yang,  
17 X.: The influence of snow grain size and impurities on the vertical profiles of actinic  
18 flux and associated NO<sub>x</sub> emissions on the Antarctic and Greenland ice sheets, *Atmos.*  
19 *Chem. Phys.*, 13, 3547-3567, 10.5194/acp-13-3547-2013, 2013.

20 Zhou, X. L., Beine, H. J., Honrath, R. E., Fuentes, J. D., Simpson, W., Shepson, P. B.,  
21 and Bottenheim, J. W.: Snowpack photochemical production of HONO: a major  
22 source of OH in the Arctic boundary layer in springtime, *Geophys. Res. Lett.*, 28,  
23 4087-4090, 10.1029/2001GL013531, 2001.

24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

**Table 1.** Uncertainties of replicate measurements of reference gases and standards using the bacterial denitrifier method (Kaiser et al., 2007) at the University of Washington IsoLab.

**Table 2.** Frequency (number of years per decade) of years with double nitrate peaks ~~the spring nitrate peak~~ in each decade from 1960 to 2006 in the shallow ice core.

1

2 **Figure 1.**  $\delta^{18}\text{O}(\text{NO}_3^-)$  (a),  $\delta^{15}\text{N}(\text{NO}_3^-)$  (b),  $\Delta^{17}\text{O}(\text{NO}_3^-)$  (c) and concentrations of  $\text{NO}_3^-$   
3 (d),  $\text{Na}^+$  (e),  $\text{SO}_4^{2-}$  (f) and  $\text{Mg}^{2+}$  (g) in the snow samples (data are 3-point running  
4 averages). SP-1 data were plotted in gray with plus signs; SP-2 and SB data were  
5 plotted in black with circle and star signs, respectively. The pink and green stars  
6 represent samples in SB used to calculate the winter and 2005 spring mean  $\Delta^{17}\text{O}(\text{NO}_3^-)$   
7 ), respectively; while the red and blue plus signs represent samples in SP-1 used to  
8 calculate the winter and 2005 spring mean  $\Delta^{17}\text{O}(\text{NO}_3^-)$ , respectively. The vertical  
9 dashed line indicates the approximate months of snowfall at depths of 0, 0.5, 1.0, 1.5  
10 and 2.0 m by Method B as described in the text.

11

12 **Figure 2.** (a) Nitrate concentration record from the shallow ice core; (b), (c), (d) show  
13 seasonal peaks of  $\text{NO}_3^-$ ,  $\text{Na}^+$  and  $\text{Ca}^{2+}$ , respectively in two sections of the ice core  
14 (data shown in b, c and d are 3-point running averages). Vertical dashed lines in (a)  
15 indicate the years of snowfall at relevant depths.

16

17 **Figure 3.** Summit springtime  $\text{O}_3$  column density (February and March averages) and  
18 poleward eddy heat flux in the lower stratosphere (40-80 °N, January and February  
19 averages) from 1979 to 2006. The solid blue line is the average of springtime  $\text{O}_3$   
20 column density from 1979 to 2006 ( $390 \pm 50$  ( $1\sigma$ ) DU), and the dashed blue line is the  
21 linear least-squared regression of the  $\text{O}_3$  column density from 1979 to 2006. The  
22 symbols marked as red and green indicate ~~the~~ years with double nitrate peaks, while



1 the green symbols indicate one of the double nitrate peaks is accompanied with a  
 2 sulfate peak, a spring nitrate peak observed in the shallow ice core.

3 **Table 1.** Uncertainties of replicate measurements of reference gases and standards  
 4 using the bacterial denitrifier method (Kaiser et al., 2007) at the University of  
 5 Washington IsoLab.

		$\delta^{15}\text{N}$ (‰)	$\delta^{17}\text{O}$ (‰)	$\delta^{18}\text{O}$ (‰)	$\Delta^{17}\text{O}$ (‰)
Zero Enrichment <sup>a</sup> (N = 30)		0.02	0.10	0.01	0.10
IAEA-NO-3 (vs. reference gas) <sup>b</sup>	Short term <sup>d</sup> (N = 6)	5.1 ± 1.0	28.9 ± 0.4	56.1 ± 0.6	-0.3 ± 0.1
USGS35 (vs. reference gas) <sup>b</sup>	Short term (N = 6)	1.4 ± 1.6	63.6 ± 0.2	85.2 ± 0.4	19.3 ± 0.1
IAEA-NO-3 (normalized) <sup>c</sup>	Long term <sup>e</sup> (N > 100)	4.7 ± 0.8	12.9 ± 0.8	25.3 ± 1.3	-0.6 ± 0.4
	Short term (N = 6)	4.7 ± 1.0	12.9 ± 0.6	25.2 ± 0.8	-0.6 ± 0.3
USGS35 (normalized) <sup>c</sup>	Long term (N > 100)	2.5 ± 0.9	51.2 ± 0.9	57.1 ± 1.4	21.6 ± 0.4
	Short term (N = 6)	1.9 ± 2.1	51.1 ± 0.3	56.8 ± 0.5	21.6 ± 0.1


- 6 a. Continuous flow measurements of reference gases (O<sub>2</sub> and N<sub>2</sub>) relative to  
 7 themselves; values are one standard deviations (1σ);  
 8 b. Refers to raw values not corrected for any isotopic effects during the analytical  
 9 procedure; reported values are means ± RMSD (root-mean-square deviation);  
 10 c. Refers to corrected values using the least squares linear regression curve between  
 11 the measured and accepted values of reference materials; reported values are means  
 12 ± 1σ (standard deviation);  
 13 d. Refers to the time period (two days) when samples in this study were measured;  
 14 e. Refers to the period of instrument running since October, 2009 to present.

15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24 **Table 2.** Frequency (number of years per decade) of years with double nitrate peaks  
 25 the spring nitrate peak in each decade from 1960 to 2006 in the shallow ice core.

Decade	1960s	1970s	1980s	1990s	2000-2006
--------	-------	-------	-------	-------	-----------

1

2



3

4

**Figure 1.**  $\delta^{18}\text{O}(\text{NO}_3^-)$  (a),  $\delta^{15}\text{N}(\text{NO}_3^-)$  (b),  $\Delta^{17}\text{O}(\text{NO}_3^-)$  (c) and concentrations of  $\text{NO}_3^-$

5

(d),  $\text{Na}^+$  (e),  $\text{SO}_4^{2-}$  (f) and  $\text{Mg}^{2+}$  (g) in the snow samples (data are 3-point running

6

averages). SP-1 data were plotted in gray with plus signs; SP-2 and SB data were

7

plotted in black with circle and star signs, respectively. The pink and green stars

8

represent samples in SB used to calculate the winter and 2005 spring mean  $\Delta^{17}\text{O}(\text{NO}_3^-)$


9

(), respectively; while the red and blue plus signs represent samples in SP-1 used to


10

calculate the winter and 2005 spring mean  $\Delta^{17}\text{O}(\text{NO}_3^-)$ , respectively. The vertical

- 1 dashed line indicates the approximate months of snowfall at depths of 0, 0.5, 1.0, 1.5
- 2 and 2.0 m by Method B as described in the text.



- 3
- 4 **Figure 2.** (a) Nitrate concentration record from the shallow ice core; (b), (c), (d) show
- 5 seasonal peaks of  $\text{NO}_3^-$ ,  $\text{Na}^+$  and  $\text{Ca}^{2+}$ , respectively in two sections of the ice core
- 6 (data shown in b, c and d are 3-point running averages). Vertical dashed lines in (a)
- 7 indicate the years of snowfall at relevant depths.



1

2 **Figure 3.** Summit springtime O<sub>3</sub> column density (February and March averages) and  
 3 poleward eddy heat flux in the lower stratosphere (40-80 °N, January and February  
 4 averages) from 1979 to 2006. The solid blue line is the average of springtime O<sub>3</sub>  
 5 column density from 1979 to 2006 (390 ± 50 (1σ) DU), and the dashed blue line is the  
 6 linear least-squared regression of the O<sub>3</sub> column density from 1979 to 2006. The  
 7 symbols marked as red and green indicate the years with double nitrate peaks, while  
 8 the green symbols indicate one of the double nitrate peaks is accompanied with a  
 9 sulfate peak, a spring nitrate peak observed in the shallow ice core.