Atmos. Chem. Phys. Discuss., 14, 9705–9728, 2014 www.atmos-chem-phys-discuss.net/14/9705/2014/ doi:10.5194/acpd-14-9705-2014 © Author(s) 2014. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Organic matter matters for ice nuclei of agricultural soil origin

Y. Tobo¹, P. J. DeMott¹, T. C. J. Hill^{1,2}, A. J. Prenni¹, N. G. Swoboda-Colberg³, G. D. Franc^{2,†}, and S. M. Kreidenweis¹

¹Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, USA ²Department of Plant Sciences, University of Wyoming, Laramie, WY 82071, USA ³Department of Geology and Geophysics, University of Wyoming, Laramie, WY 82071, USA [†]deceased

Received: 29 March 2014 - Accepted: 1 April 2014 - Published: 11 April 2014

Correspondence to: Y. Tobo (ytobo@atmos.colostate.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.





Abstract

Heterogeneous ice nucleation is a crucial process for forming ice-containing clouds and subsequent ice-induced precipitation. The importance for ice nucleation of airborne desert soil dusts composed predominantly of minerals is relatively well understood.
On the other hand, the potential influence of agricultural soil dusts on ice nucleation has been poorly recognized, despite recent estimates that they may account for up to ~ 25% of the global atmospheric dust load. We have conducted freezing experiments with various dusts, including agricultural soil dusts derived from the largest dust source region in North America. Here we show evidence for the significant role of soil organic matter (SOM) in particles acting as ice nuclei (IN) under mixed-phase cloud conditions. We find that the ice nucleating ability of the agricultural soil dusts is similar to that of desert soil dusts, but is reduced to almost the same level as that of clay minerals (e.g., kaolinite) after either H₂O₂ digestion or dry heating to 300°C. In addition, based on chemical composition analysis, we show that organic-rich particles are more impor-

tant than mineral particles for the ice nucleating ability of the agricultural soil dusts at temperatures warmer than about -36°C. Finally, we suggest that such organic-rich particles of agricultural origin (namely, SOM particles) may contribute significantly to the ubiquity of organic-rich IN in the global atmosphere.

1 Introduction

- It has been shown that certain aerosol particles acting as ice nuclei (IN), such as soil dusts, soot, volcanic ash and biological materials, are required to trigger ice nucleation at temperatures warmer than about -36 °C (Szyrmer and Zawadzki, 1997; Hoose and Möhler, 2012; Murray et al., 2012). In particular, laboratory and modeling studies suggest that soil dusts are the most important IN sources at temperatures between about 26 °C and 15 °C because of their ice nucleation properties and abundances in the
- -36°C and -15°C because of their ice nucleation properties and abundances in the global atmosphere (Hoose et al., 2010; Murray et al., 2012). In the estimates from





these studies, however, all soil dust IN are regarded as minerals from desert sources and thus the contribution of agricultural soil dusts to ice nucleation has not been taken into account. Although a very large uncertainty remains regarding estimates for the contribution of agricultural emissions to the global atmospheric dust load within the ⁵ range of 0 to 50% (Mahowald et al., 2004; Tegen et al., 2004; Forster et al., 2007; Ginoux et al., 2012), the Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment Report (Forster et al., 2007) and a more recent study (Ginoux et al., 2012) suggest 0–20 and 25% as reasonable estimates, respectively.

So far, some laboratory experiments with samples immersed in supercooled water
 have shown that agricultural soil dusts having higher contents of organic matter may serve as better sources of IN than clay minerals (Schnell and Vali, 1972; Conen et al., 2011; O'Sullivan et al., 2014). However, the chemical composition of individual particles capable of nucleating ice has remained uncertain. Also, the focus of these earlier studies has centered on the role of agricultural soil dusts as IN at temperatures warmer
 than about –15 °C (Schnell and Vali, 1972; Conen et al., 2011; O'Sullivan et al., 2014), where clay minerals are less effective as IN (Szyrmer and Zawadzki, 1997; Murray et al., 2012).

In this study, we examine heterogeneous ice nucleation by aerosolized agricultural soil dusts under conditions above water saturation at temperatures warmer than about

-36 °C (i.e., mixed-phase cloud conditions where ice crystals coexist with liquid cloud droplets) and their chemical composition. In particular, we focus on the relative importance of organics (i.e., soil organic matter (SOM)) as nuclei for heterogeneous ice nucleation.

2 Materials and methods

Agricultural soil dusts were prepared using surface soils (0 to 5 cm in depth) collected on 17 May 2011 from sugar beet (42.12878°N, 104.39516°W, 1270 ma.m.s.l.) and grass/alfalfa fallow (42.12266°N, 104.38585°W, 1270 ma.m.s.l.) fields at the



Sustainable Agricultural Research and Extension Center (SAREC) near Lingle (mean annual temperature: 9.3 °C), Wyoming, USA. The agricultural fields are located within the largest dust source region in North America (Ginoux et al., 2012). Soil samples were air dried on an aluminum tray in clean conditions and then divided into particles

- smaller than 45 μm by dry sieving. China loess soils (CJ-1) (Nishikawa et al., 2000) were collected in an arid area in Gansu Province, China, and prepared for analysis by the National Institute for Environmental Studies in Japan. We also used the Clay Minerals Society kaolinite (KGa-1b) (Chipera and Bish, 2001; Murray et al., 2011), for comparison. In addition to the untreated samples, we prepared samples treated with
- ¹⁰ H_2O_2 and ones heated to 300 °C. The former samples were prepared by boiling gently in a 30 % H_2O_2 solution until all organic matter was removed, followed by rinsing with deionized water and drying. The latter samples were prepared by exposing to dry heat at 300 °C for about 2 h in a muffle furnace.
- The overview of the setup for dust aerosol generation and sampling is illustrated ¹⁵ in Fig. 1. Dry dust particles were generated using a self-built flask dust generator (nitrogen flow for dust generation: ~2Lmin⁻¹) and then passed through a cyclone (cut-point diameter: $3.5 \,\mu\text{m}$ at $2 \,\text{Lmin}^{-1}$) and 210 Po neutralizers. It has been reported that while the number-size distributions of airborne soil dust particles vary depending on the source area conditions (e.g., surface wind speed, soil characteristics) and
- ²⁰ long-range transport regimes, the mode diameters during their long-range transport typically range from ~ 2 μm down to submicrometer (Formenti et al., 2011; Kok, 2011). Here, we prepared particles with a mobility diameter of 600 nm selected in a differential mobility analyzer (DMA; Model 3081, TSI Inc.; sheath flow: 4.5 Lmin⁻¹, sample flow: 1 Lmin⁻¹). A condensation particle counter (CPC; Model 3010, TSI Inc.) was used to measure number exponentrations of the total exceed particles. A Colorade State Unit.
- ²⁵ measure number concentrations of the total aerosol particles. A Colorado State University continuous flow diffusion chamber (CFDC; Rogers et al., 2001; sheath flow: $8.5 Lmin^{-1}$, sample flow: $1.5 Lmin^{-1}$) was used to measure number concentrations of IN active under conditions above water saturation ($105.0 \pm 0.5\%$ relative humidity with respect to liquid water) at temperatures warmer than about -36 °C. Under





such conditions, heterogeneous ice nucleation processes including deposition nucleation (ice formation on the surfaces of insoluble nuclei from ice-supersaturated water vapor) and condensation/immersion freezing (ice formation during or following the condensation growth of aqueous droplets containing insoluble nuclei) are expected to be

- activated (Sullivan et al., 2010a, b; Tobo et al., 2012, 2013). These data were collected every second and then averaged for 150–180 s. The total dust aerosol particles and IN active at given temperatures were collected on a butvar film supported by Ni grids (EM Japan Co., Ltd.) using impactors. Then, the size, morphology and elemental composition of individual particles were analyzed manually after Au-coating (coating thickness:
- 2–3 nm) using a Quanta FEG MK2 scanning electron microscope (SEM; FEI Company) combined with an energy dispersive X-ray analyzer (EDX; Model 51-XMX0005, Oxford Instruments America Inc.).

3 Results and discussion

3.1 Freezing experiments

- ¹⁵ In Fig. 2a, we show the number fraction of particles capable of nucleating ice as a function of temperature. The results show that the ice nucleation properties of agricultural soil dusts obtained from sugar beet and grass/alfalfa fallow fields in Wyoming are similar to each other. After H₂O₂ treatment, both agricultural soil dusts experienced a significant reduction in their ice nucleating ability at temperatures warmer than about -36 °C,
- ²⁰ suggesting that they contain specific ice nucleation active constituents that can be removed by H₂O₂ treatment (most likely, organic matter). To evaluate the possibility that the ice nucleation properties of some mineral components might be affected by H₂O₂ treatment, we conducted freezing experiments with China loess soil dust (desert loess; a proxy for Asian natural soil dusts, Nishikawa et al., 2000) and kaolinite (Chipera and Bish, 2001; Murray et al., 2011) in the same manner. The results indicate that the





impact of H_2O_2 treatment on their ice nucleating ability is relatively small. In addition, we confirmed that dry heating to 300 °C has a similar impact to H_2O_2 digestion (Fig. 3).

In Fig. 2b, we compare the experimental data for agricultural soil dusts collected in Wyoming with the parameterizations for various dusts based on the number of ice

⁵ nucleation active sites per unit surface area (i.e., ice nucleation active site density: n_s). To estimate their surface area, we assume that all particles are spherical. The results show that the n_s values for the original Wyoming agricultural soil dusts are relatively similar to those for desert (Niemand et al., 2012) and agricultural (O'Sullivan et al., 2014) soil dusts from various locations in the world, at least in the temperature range examined. After H₂O₂ digestion, however, the n_s values for the Wyoming agricultural soil dusts are reduced to an almost comparable level to those for kaolinite and other clay minerals (e.g., montmorillonite, Atkinson et al., 2013). Around -36 °C (near to the limit for homogeneous freezing temperature of water in droplets, Koop et al., 2000), all samples presented here show relatively similar n_s values.

15 3.2 Identification of particle types

To identify particle types in the agricultural soil dusts used here, we examined the elemental composition of individual particles using SEM/EDX analysis. We found that the majority of the analyzed particles can be classified as either "organics" or "minerals" (Fig. 4). The major elements of organics are C, N and S (examples of the other elements: O, F, Na, Cl, K). It should be noted that we classify carbonaceous particles lacking N and S (e.g., soot-like particles as shown in Fig. 4e) as "others" and not organics. The major element of minerals is Si (examples of the other elements: C, O, F, Na, Mg, Al, Cl, K, Ca, Mn, Fe, Ti). Particles containing the major elements of both organics and minerals (i.e., C, N, Si and S) are categorized as "organo-mineral mixtures".

In Fig. 5a, we summarize the results of SEM/EDX analysis for the total dust aerosol particles and IN active at temperatures of -36 to -24 °C in the untreated 600 nm monodisperse sugar beet soil dust particles. The results show that mineral particles account for more than half of the total aerosol population and organic-rich particles for



about 40 %. However, the number fraction of mineral particles in the IN population decreases dramatically with increasing freezing temperatures. Correspondingly, the fraction of organic particles increases, accounting for nearly 90% of the IN population at around -24 °C. The fraction of organo-mineral mixture particles or other particles (mainly, soot-like and/or Na-rich particles) is only 2–7% in both the total aerosol and IN populations.

5

10

Furthermore, we estimated the fractions of organic and inorganic particles in the agricultural soil dust IN, based on the results of freezing experiments with the soil dusts before and after H_2O_2 treatment. Here, we assume that organic and inorganic particles account for 40 and 60% of the total agricultural dust aerosol particles, respectively (this assumption is based on the results of the SEM/EDX analysis) and that all organic compounds can be digested and removed after H_2O_2 treatment. The calculation method is detailed in Appendix A1. The temperature-dependent changes in the fractions of organic and inorganic IN estimated from this calculation (Fig. 5b) are approximately consistent with the results from the SEM/EDX analysis (Fig. 5a), indicating

¹⁵ proximately consistent with the results from the SEM/EDX analysis (Fig. 5a), indicating that the reduction of the ice nucleating ability of the agricultural soil dusts after H_2O_2 treatment (Fig. 2) can be explained mainly by the removal of organic matter.

Based on these results, we propose organic-rich particles (namely, SOM particles) as the most important component of agricultural soil dusts for ice nucleation in the

- temperature regime examined. We note that while phosphorus is known to be one of the major biological markers (Pósfai et al., 2003; Pratt et al., 2009; Creamean et al., 2013; Cziczo et al., 2013), no P-containing particles were found in all of the analyzed particles. This suggests that there was no measurable contribution of microorganisms (e.g., fungal spores, bacteria) to the numbers of the agricultural soil dusts or IN exam-
- ined here, although the possibility of the presence of some P-free microorganisms or plants (or their fragments) cannot be ruled out. Also, humic-like substances are known as one of the major components of SOM. However, immersion freezing experiments with certain standard humic-like substances (e.g., humic acid, fulvic acid) have indicated that while they can act as IN under mixed-phase cloud conditions (Fornea et al.,





2009; Wang and Knopf, 2011; Knopf and Alpert, 2013; Rigg et al., 2013; O'Sullivan et al., 2014), they are much less effective as IN than fertile soil dusts (O'Sullivan et al., 2014). Thus, the major sources of SOM particles having very high ice nucleating ability are not yet known.

- It is noteworthy that agricultural soil dusts collected in Wyoming have similar ice nucleating abilities to those collected in England (Fig. 2b). In this regard, O'Sullivan et al. (2014) suggested that mineral components are more important than biogenic components for the ice nucleating ability of the England soil dusts at temperatures colder than about -15°C. Their suggestion was based on the results from freezing experiments with the soil dusts before and after wet heating to 90°C. It is expected
- that wet heating to $90-100^{\circ}$ C deactivates only certain organic matter (e.g., proteins or proteinaceous compounds, Christner et al., 2008). In fact, it has been reported that wet heating to $90-100^{\circ}$ C is less effective than H₂O₂ digestion in reducing the ice nucleating ability of fertile soil dusts (Conen et al., 2011; O'Sullivan et al., 2014). On the other
- ¹⁵ hand, we applied treatments designed to remove and/or deactivate almost all organic matter (i.e., H_2O_2 digestion or dry heating to 300°C). Our results indicate that the presence of organic compounds (i.e., SOM particles) has a significant influence on the ice nucleating ability of agricultural soil dusts throughout the entire temperature range down to about -36°C.
- ²⁰ Our results also show that both agricultural and desert soil dusts have similar ice nucleating abilities and are more efficient IN than clay minerals over the wide temperature range examined (Fig. 2b). Since treatments to remove and/or deactivate organic matter have a small impact on the ice nucleating ability of China loess soil dust, the key ice nucleation active sites contained in desert soil dusts are presumed to be re-
- ²⁵ lated to inorganic compounds. In this regard, a recent study reported that the feldspar (in particular, K-feldspar) component can explain higher n_s values for desert soil dusts than clay minerals (Atkinson et al., 2013). Since we have not evaluated the feldspar content in dust samples used here, further investigations are required to verify this





hypothesis. Nevertheless, our results clearly demonstrate that the key IN type is quite different between agricultural and desert soil dusts.

3.3 Atmospheric implications

In order to estimate the contribution of agricultural soil dusts to the IN population in the atmosphere, we combined a global simulation of atmospheric concentrations of 5 different aerosol particle types at the 600 hPa pressure altitude (Hoose et al., 2010; Murray et al., 2012) with experimentally derived n_s values. The calculation method for potential IN number concentrations is detailed in Appendix A2. The n_s values used in the calculation are based on the parameterizations for desert soil dusts (Niemand et al., 2012) and for agricultural soil dusts derived from this study. In Fig. 6a, we show 10 the estimates of potential number concentrations of soil dust IN, based on the classical view that all dusts are of natural origin and that desert soil dusts are representative of natural dusts (Murray et al., 2012). In addition, we provide the estimates of potential number concentrations of soil dust IN by assuming that agricultural soil dusts account for either 5% (Tegen et al., 2004) or 25% (Ginoux et al., 2012) of the total soil dust 15 emissions (Fig. 6b). In these cases, agricultural soil dusts represent a relatively small but non-negligible contribution to the IN population, as compared with natural soil dusts. The influence of biological IN on ice clouds is also a controversial topic (Pratt et al., 2009; Creamean et al., 2013; Cziczo et al., 2013). In this regard, the results in Fig. 6b

- ²⁰ suggest that the contribution of agricultural soil dusts to the IN population may be more significant than that of microorganisms, as exemplified by certain well-known fungal spores (lannone et al., 2011; Murray et al., 2012), at least at temperatures below about –18°C and on a global scale. Considering that SOM particles play the dominant role in the ice nucleating ability of agricultural soil dusts at temperatures warmer than about
- -36°C (Fig. 5), the results in Fig. 6b suggest the possibility that SOM particles of agricultural origin may contribute strongly to the global atmospheric IN population.

The results presented here offer a possible explanation for the presence of organicrich particles found in residues within ice clouds and ice-phase precipitation. For





example, field studies have indicated that organic/sulfate/nitrate particles account for about 5-25 % of the residues involved in heterogeneous ice nucleation in ice-containing clouds, whereas mineral particles always predominate (DeMott et al., 2003; Richardson et al., 2007; Pratt et al., 2009; Creamean et al., 2013; Cziczo et al., 2013). Also,

- s some field studies have inferred the possible role of urban anthropogenic organic particles as IN at temperatures as warm as -20°C (Knopf et al., 2010; Wang et al., 2012b), although the composition of individual IN was not examined. Laboratory experiments have indicated that certain sulfates (e.g., ammonium sulfate) and oxidized organics exist as anhydrous salts or glassy solids at relatively cold temperatures and act as ef-
- fective IN (Abbatt et al., 2006; Murray et al., 2010; Wang et al., 2012a). On the other 10 hand, the experimental data do not readily explain heterogeneous ice nucleation by these materials at temperatures warmer than about -30°C. Here, we speculate that SOM particles of agricultural origin can be regarded as a possible source for the organic/nitrate/sulfate particles found in residues within ice clouds. This idea may also
- be supported by recent work showing that most organic matter contained in hailstones 15 originated from soils (Šanti-Temkiv et al., 2013), leading to speculation that SOM particles might participate in ice nucleation as well as be scavenged by ice-phase precipitation. Further field, laboratory and modeling studies will therefore be necessary to validate the hypothesis that SOM particles of agricultural origin are indeed an important
- source of nuclei for atmospheric ice nucleation.

Conclusions 4

25

In this study, we highlight the role of agricultural soil dusts as IN under mixed-phase cloud conditions. Our results indicate that the ice nucleating ability of agricultural soil dusts is comparable to that of desert soil dusts, but is reduced to almost the same level as clay minerals after treatments to remove and/or deactivate almost all organic matter (i.e., H₂O₂ digestion or dry heating to 300 °C). Based on chemical composition analysis of individual IN, we demonstrate that the presence of SOM particles, rather





than mineral particles, is largely responsible for the ice nucleating ability of the agricultural soil dusts at temperatures warmer than about -36 °C. In addition, we suggest the possibility that SOM particles of agricultural origin may be regarded as a possible significant source influencing the ubiquity of organic-rich IN and also residues found within ice clouds and ice-phase precipitation.

Appendix A

A1 Estimation of the composition of ice nuclei of agricultural soil origin

Here, we explain how to estimate the fraction of organic and inorganic IN of agricultural soil origin as shown in Fig. 5b, based on the n_s parameterizations obtained from freezing experiments. The number concentration of IN active at a given temperature *T*, $N_{\rm IN}(T)$, can be described by (Murray et al., 2012):

$$N_{\rm IN}(T) = N_{\rm total} \left(1 - \exp(-n_{\rm s}(T) s)\right) \tag{A1}$$

where N_{total} is the number concentration of total particles, $n_{s}(T)$ is the ice nucleation active site density and *s* is the surface area of a single particle. Similarly, the number concentrations of IN of agricultural soil origin, $N_{\text{IN} [agri. soildust]}(T)$, can be expressed as:

 $N_{\text{IN} [\text{agri. soil dust}]}(T) = N_{\text{total} [\text{agri. soil dust}]}(1 - \exp(-n_{\text{s} [\text{agri. soil dust}]}(T) s))$ (A2)

where $N_{\text{total [agri. soildust]}}$ is the number concentration of agricultural soil dust particles, $n_{s [agri. soildust]}(T)$ is the n_s value for untreated agricultural soil dust particles presented in Fig. A1a. When calculating *s*, all soil dust particles are assumed to be spherical particles having a diameter of 600 nm. Also, if only inorganic particles exist after the removal of organic matter by H_2O_2 treatment, the number concentration of inorganic IN of agricultural soil origin, $N_{\text{IN [inorganic]}}(T)$, can be expressed as:

²⁵
$$N_{\text{IN [inorganic]}}(T) = N_{\text{total [inorganic]}} (1 - \exp(-n_{\text{s [inorganic]}}(T) s))$$



(A3)

where $N_{\text{total [inorganic]}}$ is the number concentration of inorganic particles of agricultural soil origin, $n_{s [inorganic]}$ is the n_s value for H_2O_2 -treated agricultural soil dust particles presented in Fig. A1a. To obtain $N_{\text{total [inorganic]}}$ in Eq. (A3), we assume that inorganic (mostly, mineral) particles account for 60% of the total agricultural soil dust particles (i.e., organic: inorganic ratio = 4:6; see the top-left pie chart in Fig. 5b), with the choiceof the ratio guided by in the elemental composition analysis of individual agricultural soildust particles (see the top-left pie chart in Fig. 5a):

 $N_{\text{total [inorganic]}} = 0.6 N_{\text{total [agri. soil dust]}}$

By combining Eqs. (A2), (A3) and (A4), the fractions of both inorganic and organic particles in the agricultural soil dust IN, $f_{\text{IN [inorganic]}}(T)$ and $f_{\text{IN [organic]}}(T)$, respectively, can be estimated:

$$f_{\text{IN [inorganic]}}(T) = \frac{N_{\text{IN [inorganic]}}(T)}{N_{\text{IN [agri. soildust]}}(T)}$$

15

$$f_{\text{IN [organic]}}(T) = 1 - f_{\text{IN [inorganic]}}(T)$$
(A6)

A2 Estimation of the number concentrations of ice nuclei of agricultural soil origin

Here, we describe a possible method to estimate the global mean number concentrations of IN of agricultural soil origin under mixed-phase cloud conditions as shown in Fig. 6. According to modeling estimates, the zonal annual mean number concentrations of soil dusts (1 µm in diameter) at the 600 hPa pressure altitude range from 0.1 to 50 cm⁻³ (Hoose et al., 2010; Murray et al., 2012). If soil dusts in the global atmosphere can be regarded as consisting of only desert and agricultural soil dusts, then the zonal annual mean concentration of soil dusts, $N_{\text{total [soil dust]}}$ (= 0.1 to 50 cm⁻³), may be expressed as:

$$N_{\text{total [soil dust]}} = N_{\text{total [natural soil dust]}} + N_{\text{total [agri. soil dust]}}$$

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

(A7)

(A4)

(A5)

where $N_{\text{total [natural soil dust]}}$ is the number concentration of natural soil dust particles. Recently, Murray et al. (2012) estimated the zonal annual mean number concentration of IN of natural soil origin, $N_{\text{IN [natural soil dust]}}(T)$, using the formula:

$${}_{5} N_{\text{IN [natural soil dust]}}(T) = N_{\text{total [natural soil dust]}} (1 - \exp(-n_{\text{s [desert soil dust]}}(T) s))$$
(A8)

where $n_{s \text{[desert soil dust]}}(T)$ is the n_s value for desert soil dusts (Niemand et al., 2012). In this calculation, Murray et al. (2012) assumed that all dusts are of natural origin (i.e., $N_{\text{total [natural soil dust]}} : N_{\text{total [agri. soil dust]}} = 100 : 0$) and that natural soil dusts can be regarded as desert soil dusts. The results are shown in Fig. 6a. On the other hand, in Fig. 6b, we provide two estimates of the global mean number concentrations of both natural and agricultural soil dust IN by assuming that agricultural soil dusts account for 5% ($N_{\text{total [natural soil dust]}} : N_{\text{total [natural soil dust]}} = 75 : 25$) and by combining Eqs. (A2), (A7) and (A8).

- Acknowledgements. We thank M. Nishikawa for providing China loess soils; R. Baumgartner and J. Meeks for access to the fields at SAREC; K. Hara, A. Matsuki and Y. Iwasaka for their support for the preparation of SEM/EDX analysis; R. C. Sullivan for his assistance for the setup of dust generation systems. This research was funded in part by the National Science Foundation (NSF) under grants ATM-0841542, ATM-0841602 and AGS-1036028. Y. Tobo acknowledges the Japan Society for the Promotion of Science (JSPS) Postdoctoral Fellowships for
- 20 edges the Japan Society for the Promotion of Science (JSPS) Postdoctoral Fellowships for Research Abroad.

References

- Abbatt, J. P. D., Benz, S., Cziczo, D. J., Kanji, Z., Lohmann, U., and Möhler, O.: Solid ammonium sulfate aerosols as ice nuclei: a pathway for cirrus cloud formation, Science, 313, 1770–1773, 2006.
- 25
- Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., Dobbie, S., O'Sullivan, D., and Malkin, T. L.: The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds, Nature, 498, 355–358, 2013.





- Chipera, S. J. and Bish, D. L.: Baseline studies of the clay minerals society source clays: powder X-ray diffraction analyses, Clay. Clay Miner., 49, 398–409, 2001.
- Christner, B. C., Morris, C. E., Foreman, C. M., Cai, R., and Sands, D. C.: Ubiquity of biological ice nucleators in snowfall, Science, 319, 1214–1214, 2008.
- ⁵ Conen, F., Morris, C. E., Leifeld, J., Yakutin, M. V., and Alewell, C.: Biological residues define the ice nucleation properties of soil dust, Atmos. Chem. Phys., 11, 9643–9648, doi:10.5194/acp-11-9643-2011, 2011.
 - Creamean, J. M., Suski, K. J., Rosenfeld, D., Cazorla, A., DeMott, P. J., Sullivan, R. C., White, A. B., Ralph, F. M., Minnis, P., Comstock, J. M., Tomlinson, J. M., and Prather, K. A.:
- ¹⁰ Dust and biological aerosols from the Sahara and Asia influence precipitation in the western US, Science, 339, 1572–1578, 2013.
 - Cziczo, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M., Zondlo, M. A., Smith, J. B., Twohy, C. H., and Murphy, D. M.: Clarifying the dominant sources and mechanisms of cirrus cloud formation, Science, 340, 1320–1324, 2013.
- ¹⁵ DeMott, P. J., Cziczo, D. J., Prenni, A. J., Murphy, D. M., Kreidenweis, S. M., Thomson, D. S., Borys, R., and Rogers, D. C.: Measurements of the concentration and composition of nuclei for cirrus formation, P. Natl. Acad. Sci. USA, 100, 14655–14660, 2003.
 - Formenti, P., Schütz, L., Balkanski, Y., Desboeufs, K., Ebert, M., Kandler, K., Petzold, A., Scheuvens, D., Weinbruch, S., and Zhang, D.: Recent progress in understanding physical
- and chemical properties of African and Asian mineral dust, Atmos. Chem. Phys., 11, 8231– 8256, doi:10.5194/acp-11-8231-2011, 2011.
 - Fornea, A. P., Brooks, S. D., Dooley, J. B., and Saha, A.: Heterogeneous freezing of ice on atmospheric aerosols containing ash, soot, and soil, J. Geophys. Res., 114, D13201, doi:10.1029/2009JD011958, 2009.
- Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in atmospheric constituents and in radiative forcing, in: Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the International Panel on Climate Change, edited by: Solomon, S., Qin, D.,
- Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge Univ. Press, Cambridge, UK, 129–234, 2007.





- Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C., and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, Rev. Geophys., 50, RG3005, doi:10.1029/2012RG000388, 2012.
 Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of
- results from laboratory experiments, Atmos. Chem. Phys., 12, 9817–9854, doi:10.5194/acp-12-9817-2012, 2012.
 - Hoose, C., Kristjánsson, J. E., Chen, J.-P., and Hazra, A.: A classical-theory-based parameterization of heterogeneous ice nucleation by mineral dust, soot, and biological particles in a global climate model, J. Atmos. Sci., 67, 2483–2503, 2010.
- ¹⁰ Knopf, D. A. and Alpert, P. A.: A water activity based model of heterogeneous ice nucleation kinetics for freezing of water and aqueous solution droplets, Faraday Discuss., 165, 513–534, 2013.
 - Knopf, D. A., Wang, B., Laskin, A., Moffet, R. C., and Gilles, M. K.: Heterogeneous nucleation of ice on anthropogenic organic particles collected in Mexico City, Geophys. Res. Lett., 37, L11202, doi:10.1020/2010CL.042262.2010
- L11803, doi:10.1029/2010GL043362, 2010.
 - Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate the size of the global dust cycle, P. Natl. Acad. Sci. USA, 108, 1016–1021, 2011.
 - Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, Nature, 406, 611–614, 2000.
 - Mahowald, N. M., Rivera, G. D. R., and Luo, C.: Comment on "Relative importance of climate and land use in determining present and future global soil dust emission" by I. Tegen et al., Geophys. Res. Lett., 31, L24105, doi:10.1029/2004GL021272, 2004.
 - Murray, B. J., Wilson, T. W., Dobbie, S., Cui, Z., Al-Jumur, S. M. R. K., Möhler, O., Schnaiter, M.,
- ²⁵ Wagner, R., Benz, S., Niemand, M., Saathoff, H., Ebert, V., Wagner, S., and Kärcher, B.: Heterogeneous nucleation of ice particles on glassy aerosols under cirrus conditions, Nat. Geosci., 3, 233–237, 2010.
 - Murray, B. J., Broadley, S. L., Wilson, T. W., Atkinson, J. D., and Wills, R. H.: Heterogeneous freezing of water droplets containing kaolinite particles, Atmos. Chem. Phys., 11, 4191–4207, doi:10.5104/acp.11.4101.2011.
- ³⁰ doi:10.5194/acp-11-4191-2011, 2011.

20

Murray, B. J., O'Sullivan, D., Atkinson, J. D., and Web, M. E.: Ice nucleation by particles immersed in supercooled cloud droplets, Chem. Soc. Rev., 41, 6519–6554, 2012.





- Niemand, M., Möhler, O., Vogel, B., Vogel, H., Hoose, C., Connolly, P., Klein, H., Bingemer, H., DeMott, P., Skrotzki, J., and Leisner, T.: A particle-surface-area-based parameterization of immersion freezing on desert dust particles, J. Atmos. Sci., 69, 3077–3092, 2012.
- Nishikawa, M., Hao, Q., and Morita, M.: Preparation and evaluation of certified reference materials for Asian mineral dust, Global Environ. Res., 4, 103–113, 2000.
- rials for Asian mineral dust, Global Environ. Res., 4, 103–113, 2000.
 O'Sullivan, D., Murray, B. J., Malkin, T. L., Whale, T. F., Umo, N. S., Atkinson, J. D., Price, H. C., Baustian, K. J., Browse, J., and Webb, M. E.: Ice nucleation by fertile soil dusts: relative importance of mineral and biogenic components, Atmos. Chem. Phys., 14, 1853–1867, doi:10.5194/acp-14-1853-2014, 2014.
- ¹⁰ Pósfai, M., Li, J., Anderson, J. R., and Buseck, P. R.: Aerosol bacteria over the South Ocean during ACE-1, Atmos. Res., 66, 231–240, 2003.
 - Pratt, K. A., DeMott, P. J., French, J. R., Wang, Z., Westphal, D. L., Heymsfield, A. J., Twohy, C. H., Prenni, A. J., and Prather, K. A.: In situ detection of biological particles in cloud ice-crystals, Nat. Geosci., 2, 398–401, 2009.
- Richardson, M. S., DeMott, P. J., Kreidenweis, S. M., Cziczo, D. J., Dunlea, E. J., Jimenez, J. L., Thomson, D. S., Ashbaugh, L. L., Borys, R. D., Westphal, D. L., Casuccio, G. S., and Lersch, T. L.: Measurements of heterogeneous ice nuclei in the western United State in springtime and their relation to aerosol characteristics, J. Geophys. Res., 112, D02209, doi:10.1029/2006JD007500, 2007.
- Rigg, Y. J., Alpert, P. A., and Knopf, D. A.: Immersion freezing of water and aqueous ammonium sulfate droplets initiated by humic-like substances as a function of water activity, Atmos. Chem. Phys., 13, 6603–6622, doi:10.5194/acp-13-6603-2013, 2013.

25

30

- Rogers, D. C., DeMott, P. J., Kreidenweis, S. M., and Chen, Y.: A continuous-flow diffusion chamber for airborn measurements of ice nuclei, J. Atmos. Ocean. Tech., 18, 725–741, 2001.
- Šantl-Temkiv, T., Finster, K., Dittmar, T., Hansen, B. M., Thyrhaug, R., Nielsen, N. W., and Karlson, U. G.: Hailstones: a window into the microbial and chemical inventory of a storm cloud, PLoS ONE, 8, e53550, doi:10.1371/journal.pone.0053550, 2013.
- Schnell, R. C. and Vali, G.: Atmospheric ice nuclei from decomposing vegetation, Nature, 236, 163–165, 1972.
- Sullivan, R. C., Miñambres, L., DeMott, P. J., Prenni, A. J., Carrico, C. M., Levin, E. J. T., and Kreidenweis, S. M.: Chemical processing does not always impair heteroge-





9721

neous ice nucleation of mineral dust particles, Geophys. Res. Lett., 37, L24805, doi:10.1029/2010GL045540, 2010a.

- Sullivan, R. C., Petters, M. D., DeMott, P. J., Kreidenweis, S. M., Wex, H., Niedermeier, D., Hartmann, S., Clauss, T., Stratmann, F., Reitz, P., Schneider, J., and Sierau, B.: Irreversible loss
- of ice nucleation active sites in mineral dust particles caused by sulphuric acid condensation, 5 Atmos. Chem. Phys., 10, 11471–11487, doi:10.5194/acp-10-11471-2010, 2010b.
 - Szyrmer, W. and Zawadzki, I.: Biogenic and anthropogenic sources of ice-forming nuclei: a review, B. Am. Meteorol. Soc., 78, 209-228, 1997.
 - Tegen, I., Werner, M., Harrison, S. P., and Kohfeld, K. E.: Relative importance of climate and
- land use in determining present and future global soil dust emission, Geophys. Res. Lett., 10 31. L05105. doi:10.1029/2003GL019216. 2004.
 - Tobo, Y., DeMott, P. J., Raddatz, M., Niedermeier, D., Hartmann, S., Kreidenweis, S. M., Stratmann, F., and Wex, H.: Impacts of chemical reactivity on ice nucleation of kaolinite particles: a case study of levoglucosan and sulfuric acid. Geophys. Res. Lett., 39, L19803. doi:10.1029/2012GL053007.2012.
 - Tobo, Y., Prenni, A. J., DeMott, P. J., Huffman, A. J., McCluskey, C. S., Tian, G., Pöhlker, C., Pöschl, U., and Kreidenweis, S. M.: Biological aerosol particles as a key determinant of ice nuclei populations in a forest ecosystem, J. Geophys. Res.-Atmos., 118, 10100–10110, 2013.

15

- Wang, B. and Knopf, D. A.: Heterogeneous ice nucleation on particles composed of humic-like 20 substances impacted by O₃, J. Geophys. Res., 116, D03205, doi:10.1029/2010JD014964, 2011.
 - Wang, B., Lambe, A. T., Massoli, P., Onasch, T. B., Davidovits, P., Worsnop, D. R., and Knopf, D. A.: The deposition ice nucleation and immersion freezing potential of amorphous
- secondary organic aerosol: pathways for ice and mixed-phase cloud formation, J. Geophys. 25 Res., 117, D16209, doi:10.1029/2012JD018063, 2012a.
 - Wang, B., Laskin, A., Roedel, T., Gilles, M. K., Moffet, R. C., Tivanski, A. V., and Knopf, D. A.: Heterogeneous ice nucleation and water uptake by field-collected atmospheric particles below 273 K, J. Geophys. Res., 117, D00V19, doi:10.1029/2012JD017446, 2012b.



Discussion

Paper

Discussion Paper

Discussion Paper





Fig. 1. Schematics of freezing experiment systems. Dry dust particles were generated from the flask dust generator. The DMA selected particles with a mobility diameter of 600 nm for analysis by the CPC and CFDC. Impactors were used for direct sampling of the total particles and ice-nucleating particles.



Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper





Fig. 2. Ice nucleating ability for various dusts under mixed-phase cloud conditions. **(a)** Fraction of 600 nm monodisperse particles capable of nucleating ice. The data for untreated and H_2O_2 -treated samples are shown in each figure. **(b)** Ice nucleation active site densities for various dusts. The n_s parameterizations for various dusts are compared with the data for agricultural soil dusts (before and after H_2O_2 treatment) collected in Wyoming. The n_s parameterizations for China loess soil dust and kaolinite are based on the data presented in Fig. A1b. The n_s parameterizations for fertile soil dusts collected in England (O'Sullivan et al., 2014) and desert soil dusts (Niemand et al., 2012) are also shown. Error bars represent standard deviations.







Fig. 3. Same as Fig. 2a, but for the data for untreated and heated samples.







Fig. 4. Representative SEM/EDX images of sugar beet soil dust particles. **(a)** Organic particle (crystalline-shape type). **(b)** Organic particle (amorphous-shape type). **(c)** Organo-mineral mixture particle. **(d)** Mineral particle. **(e)** Soot-like particle. Scale bars, 300 nm. Red X-ray spectra show the elemental composition of each particle, and grey X-ray spectra represent the background signal caused by a butvar film supported by Ni mesh grids and Au-coating.







Fig. 5. Relative contributions of different particle types to agricultural soil dusts. **(a)** Compositions of the total dust aerosol particles (600 nm monodisperse sugar beet soil dust particles) and IN active at -36, -30 and -24° C determined by SEM/EDX analysis (n = 95, 58, 52 and 68, respectively). **(b)** Compositions of the IN active at -36, -30 and -24° C estimated from the freezing experiments with untreated and H_2O_2 -treated agricultural soil dusts. In this calculation, the percentages of organic and inorganic particles in the total dust aerosol particles are set to 40 and 60 %, respectively, and then the n_s parameterizations presented in Fig. A1a are used to estimate the number fractions of organic and inorganic IN.







Fig. 6. Estimates of global mean number concentrations of IN active under mixed-phase cloud conditions. **(a)** IN number concentrations for soil dusts, assuming that all dusts are of natural origin (i.e., desert origin). **(b)** IN number concentrations for agricultural and natural soil dusts, assuming that 5 and 25% of all dusts are of agricultural origin. Calculations are performed using the n_s parameterizations for desert soil dusts (Niemand et al., 2012) and for untreated agricultural soil dusts (Fig. A1a) and the range of the zonal annual mean concentrations of dusts at 600 hPa (lower and upper limits) (Hoose et al., 2010; Murray et al., 2012). Also provided for comparison are IN number concentrations for fungal spores estimated using the n_s parameterization for *Cladosporium* spores (lannone et al., 2011; Murray et al., 2012) and the range of the zonal mean concentrations of fungal spores at 600 hPa (lower and upper limits) (Hoose et al., 2011; Murray et al., 2012) and the range of the zonal mean concentrations of fungal spores at 600 hPa (lower and upper limits) (Hoose et al., 2011; Murray et al., 2012) and the range of the zonal mean concentrations of fungal spores at 600 hPa (lower and upper limits) (Hoose et al., 2011; Murray et al., 2012) and the range of the zonal mean concentrations of fungal spores at 600 hPa (lower and upper limits) (Hoose et al., 2010; Murray et al., 2012).







Fig. A1. Fit to ice nucleation active site densities for various dusts. (a) Parameterizations of n_s for untreated agricultural soil dusts ($\ln(n_s) = -0.4736T + 0.3644$; validity range: $-36^{\circ}C < T < -18^{\circ}C$) and for H_2O_2 -treated agricultural soil dusts ($\ln(n_s) = -0.6773T - 7.8436$; validity range: $-36^{\circ}C < T < -22^{\circ}C$). (b) Parameterizations of n_s for untreated China loess soil dust ($\ln(n_s) = -0.5230T - 1.5767$; validity range: $-36^{\circ}C < T < -18^{\circ}C$) and for untreated kaolinite ($\ln(n_s) = -0.9803T - 17.7764$; validity range: $-36^{\circ}C < T < -26^{\circ}C$). Error bars represent standard deviations.



