

1 **How sensitive is the recovery of stratospheric ozone to changes in**
2 **concentrations of very short lived bromocarbons?**

3
4 X. Yang^{1,2,*}, N. L. Abraham^{1,2}, A. T. Archibald^{1,2}, P. Braesicke^{1,2,**}, J. Keeble², P. J.
5 Telford^{1,2}, N. J. Warwick^{1,2}, J. A. Pyle^{1,2}
6

7 ¹National Centre for Atmospheric Science (NCAS-Climate), University of Cambridge,
8 Cambridge, UK

9 ²Centre for Atmospheric Science, Department of Chemistry, University of Cambridge,
10 Cambridge, UK

11 *Now at British Antarctic Survey, Cambridge, UK

12 **Now at Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research,
13 Karlsruhe, Germany

14
15
16 **Abstract**
17

18 Naturally produced very short-lived substances (VSLS) account for almost a quarter of the
19 current stratospheric inorganic bromine, Br_y. Following VSLS oxidation, bromine radicals
20 (Br and BrO) can catalytically destroy ozone. The extent to which possible increases in
21 surface emissions or transport of these VSLS bromocarbons to the stratosphere could
22 counteract the effect of halogen reductions under the Montreal Protocol is an important
23 policy question. Here by using a chemistry-climate model, UM-UKCA, we investigate the
24 impact of a hypothetical doubling (an increase of 5 ppt Br_y) in VSLS bromocarbons on ozone
25 and how the resulting ozone changes depend on the background concentrations of chlorine
26 and bromine. Our model experiments indicate that for the 5 ppt increase in Br_y from VSLS,
27 the ozone decrease in the lowermost stratosphere of the southern hemisphere (SH) may reach
28 up to 10% in the annual mean; the ozone decrease in the northern hemisphere (NH) is smaller
29 (4-6%). The largest impact on the ozone column is found in the Antarctic spring. There is a
30 significantly larger ozone decrease following the doubling in VSLS burden under a high
31 stratospheric chlorine background than under a low chlorine background, indicating the
32 importance of the inter-halogen reactions. For example, the decline of the high latitude, lower
33 stratospheric ozone concentration as a function of Br_y is higher by about 30-40% when
34 stratospheric Cl_y is ~3 ppb (present day) compared with Cl_y of ~0.8 ppb (a pre-industrial or
35 projected future situation). Bromine will play an important role in the future ozone layer.
36 However, even if bromine levels from natural VSLS were to increase significantly later this
37 century, changes in the concentration of ozone will likely be dominated by the decrease of
38 anthropogenic chlorine. Our calculation suggests that for a 5 ppt increase in Br_y from VSLS,
39 the Antarctic ozone hole recover date could be delayed by approximately 6-8 years,
40 depending on Cl_y levels.
41
42

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92

1. Introduction

It is now very well established that stratospheric ozone loss since the 1970s has occurred as a consequence of emissions of halogenated species into the atmosphere. These gases, the CFCs and halons, are generally long lived in the troposphere but at sufficient altitudes in the stratosphere they can be oxidized to liberate chlorine (as suggested by Molina and Rowland, 1974) and bromine atoms (see Wofsy et al, 1975), which can subsequently play a role in catalytic ozone destruction. The ‘ozone hole’ in Antarctica was reported by Farman and his colleagues in 1985 (Farman et al., 1985) and intensive field, laboratory and modeling research was soon able to demonstrate that the observed decline in polar ozone was indeed a result of ozone destruction by halogen chemistry. First investigations (for details see the review by Solomon, 1999) focused on the role of chlorine but subsequent research (see, e.g., the observationally based study of Salawitch et al. (1998) and the modelling study of Chipperfield and Pyle, 1998) showed that coupled chlorine-bromine reactions made a substantial contribution to the polar loss, as first suggested by McElroy et al. (1986). Following on from the scientific understanding that anthropogenic halogens were responsible for ozone loss, the Montreal Protocol was enacted to regulate production and consumption of ozone-depleting substances. The Protocol has been a major success and atmospheric concentrations of the regulated chlorine- and bromine-gases are now in decline (WMO, 2011).

Chlorine in the stratosphere arises mainly from industrial compounds, principally the CFCs, with a contribution of about 0.6 ppb (~20% relative to current values) from long-lived, naturally occurring CH_3Cl (see, for example, Harper and Hamilton, 2003). In contrast the bromine budget of the stratosphere is less well constrained (Ko et al., 1998; Dorf et al, 2008). It is now thought that about 5 ppt (~20 to 25%) of stratospheric bromine may come from the oxidation of halocarbons with lifetimes of less than about 6 months (so-called Very Short Lived Substances, VSLS), while the majority comes from methyl bromide and the anthropogenic halons. Thorough reviews of these halogen gases are available in the recent UNEP/WMO ozone assessment (Montzka, Reimann et al., 2011), where the total stratospheric bromine concentration in 2008 is given as 22.5 ppt. In the preindustrial era, the VSLS contribution may have been as much as 50% of the total atmospheric bromine loading. VSLS bromocarbons are mainly produced by macro- and microalgae (Carpenter and Liss, 2000; Quack and Wallace 2003; Yokouchi et al., 2005). CHBr_3 and CH_2Br_2 are the dominant contributors to bromine VSLS species. Although their tropospheric lifetime is relatively short (a few weeks for CHBr_3 and several months for CH_2Br_2), they can be effectively lifted through deep convective systems and transported into the upper troposphere and/or lower stratosphere (UTLS) to make a significant contribution to the total ambient bromine (Sturges et al., 2000; Yang et al., 2005; Salawitch, 2006). Being short-lived, there are large uncertainties in using atmospheric concentration measurements to estimate their global fluxes and their net contribution to the stratospheric bromine (Warwick et al., 2006; Liang et al., 2010; Pyle et al., 2011; Ordóñez et al., 2012; Ziska et al., 2013; Hossaini et al., 2013). A large range of contributions to stratospheric inorganic bromine of 1-10 ppt can be found in the literature (Dorf et al., 2008; Salawich et al., 2010; Schofield et. al., 2011; Aschmann et al., 2011; Tegtmeier et al., 2012; Stachnik et al., 2012; Hossaini et al., 2012a). The UNEP/WMO Ozone Assessment of 2011 reports the value as 6(3-8) ppt (Montzka, Reimann et al., 2011).

93 Stratospheric inorganic chlorine has increased from pre-industrial levels of ~0.6 ppb, to a
94 peak of greater than 3 ppb in the 1990s and, under the influence of the Montreal Protocol is
95 decreasing and is expected to continue to decrease during this century (WMO, 2011). The
96 sum of the concentrations of anthropogenic bromine compounds is now also declining. Under
97 the influence of the Montreal Protocol, the stratospheric ozone layer should recover later this
98 century.

99
100 While anthropogenic halogen loading to the stratosphere should decrease in response to the
101 Montreal Protocol, we do not know how the concentration of the VSLs bromocarbons will
102 change in the future. Since they are mainly produced by biochemical processes, mostly in the
103 ocean, their emissions could change with climate, perhaps depending on sea surface
104 temperature, on ocean pH or on surface wind stress (Gschwend et al., 1985; Keppler et al.,
105 2000). We currently do not have a good enough understanding to allow confident predictions.
106 Changes in transport to the stratosphere could also occur due to climate change. For example,
107 Hossaini et al. (2012b) predict 0.3-1.0 ppt increases in the direct transport of bromine source
108 gases to the stratosphere between 2000 and 2100 under RCPs 4.5 and 8.5 while the earlier
109 study of Dessens et al. (2009) projects a 1-2 ppt increase in total bromine in the lower
110 stratosphere.

111
112 Bromine is about 100 times more efficient than chlorine as an ozone sink in the high latitude
113 lower stratosphere, with an annual average global value of around 65 (Sinnhuber et al., 2009).
114 So, any future increases in stratospheric bromine, perhaps driven by increased emissions in
115 VSLs bromocarbon or their transport to the stratosphere, could affect stratospheric ozone
116 recovery. Our primary aim in this paper is to explore the sensitivity of ozone recovery to
117 changes in the concentrations of bromine in the stratosphere, arising from very short lived
118 bromocarbons.

119
120 Gas phase and heterogeneous inter-halogen reactions involving both bromine and chlorine
121 contribute significantly to ozone loss so that the ozone loss due to bromine should be
122 dependent on the chlorine concentration. A second aim of this paper is investigate how ozone
123 might respond to changes in VSLs loading when future stratospheric chlorine levels have
124 declined.

125
126 To address these aims, we use the UM-UKCA chemistry-climate model. We perform a
127 number of idealized experiments under two different stratospheric chlorine concentrations,
128 corresponding very roughly to the present day and to the end of the 21st century, and three
129 different bromine levels. The range of bromine concentrations chosen allows us to explore
130 the sensitivity of the coupled chlorine-bromine system. Each experiment is run twice, first
131 with a VSLs contribution of 5 ppt and then with an additional increase of 5 ppt. These
132 experiments allow us to explore how ozone loss due to short lived bromocarbons could vary
133 with background chlorine and bromine loading. We note that 5 ppt would represent a large
134 change in the contribution of shortlived bromine to the stratosphere; we reiterate that these
135 are idealized experiments designed to explore the sensitivity of ozone recovery to
136 hypothetical changes in bromine.

137
138 Other changes could also affect stratospheric ozone recovery (and the contribution of
139 bromine changes to ozone recovery). Several previous studies have looked at the impact of
140 VSLs bromocarbons on ozone trends in the recent past (e.g. Salawitch et al., 2005; Feng et
141 al., 2007; Sinnhuber et al, 2009). They show that the effect of VSLs on ozone is particularly
142 important under enhanced aerosol loading. Transient changes due to volcanic eruptions

143 would certainly affect the trajectory of recovery while a sustained increase in stratospheric
144 aerosol would also lead to a general decrease in ozone and an increased role of bromine, and
145 chlorine, heterogeneous processing. However, in the calculations here aerosol loading is held
146 constant. Climate change will also affect the recovery of stratospheric ozone, as is well
147 known. Salawitch et al. (2005) show that, while the effect of bromine on ozone is dominated
148 by the ClO+BrO reaction in the lowermost stratosphere, the reaction of BrO+HO₂ becomes
149 increasingly important with increasing VLSL. Thus any change in HO₂ in the future could
150 potentially affect the estimation of the ozone impact from these changes in halogens. We do
151 not deal with this in detail here although we do mention in Section 3 possible changes in the
152 flux through the BrO+HO₂ reaction under climate change.

153
154 The model and the experimental set-up are described in more detail in Section 2. The model
155 results are presented in Section 3. Concluding discussion is found in Section 4.

156 157 **2. Model and Experiments**

158
159 UM-UKCA is a chemistry-climate model; its dynamical core is the Met Office Unified
160 Model (UM) version 7.3 running in the HadGEM3-A configuration (similar to Hewitt et al.,
161 2011). The detailed chemistry scheme for the troposphere and stratosphere, CheST, combines
162 the schemes described by Morgenstern et al. (2009) and O'Connor et al. (2014) and was
163 further updated with tropospheric bromine chemistry based on our work in the *p*TOMCAT
164 chemistry transport model (Yang et al., 2005; 2010). The stratospheric halogen scheme is
165 improved by introducing a number of heterogeneous reactions on stratospheric particles
166 (polar stratospheric clouds (PSCs) and sulphate aerosol) to account for inter-halogen
167 (chlorine and bromine) reactivation as in our recent study (Braesicke et al., 2013). The
168 model's horizontal resolution is 3.75 degree in longitude and 2.5 degree in latitude on an
169 Arakawa-C grid. A hybrid sigma-geometric height coordinate is used to resolve the vertical
170 range from the surface to ~84 km with 60 levels. All experiments are performed for year
171 2000 boundary conditions with prescribed monthly sea surface temperatures (SSTs) and sea
172 ice conditions. The radiative species concentrations for CO₂, N₂O and CH₄ are taken for year
173 2000; ozone is calculated interactively. Apart from the halocarbons (eg CFCs, CH₃Br),
174 chemical emissions are for the present day. The sulphate aerosol field is a monthly
175 climatology. PSCs and ice particles are calculated online based on water vapour fields and
176 temperatures. To get significant results, each of the experiments was performed as a 40-year
177 time-slice integration with the first 20 years as spin-up and the last 20 years for analysis.

178
179 For the VLSL bromocarbons in the model, we have introduced tracers for 5 additional species
180 (CHBr₃, CH₂Br₂, CH₂BrCl, CHBr₂Cl and CHBrCl₂) with emissions based on the original
181 work (scenario 5) of Warwick et al. (2006), except for emissions of CH₂Br₂ which were
182 updated to 57 Gg yr⁻¹, 50% of the original flux but now much more in accord with Liang et al
183 (2010) and Ordóñez et al (2012). Our updated fluxes provide ~5 ppt of inorganic bromine to
184 the stratosphere.

185
186 We have 6 base experiments, using 2 different stratospheric chlorine loadings and 3 different
187 stratospheric bromine loadings designed to cover a range of past, present and possible future
188 halogen loadings. We chose stratospheric chlorine levels of ~0.8 ppb and ~3 ppb achieved by
189 applying fixed concentrations in the model's lowest layer for the major chlorine species,
190 CFCs, CH₃Cl and CCl₄ (from the A1 Scenario of WMO 2011). 3 ppb is roughly a present day
191 stratospheric chlorine loading. The low chlorine level of 0.8 ppb represents both a
192 preindustrial chlorine value (when CH₃Cl at 0.6 ppb was the major chlorine-carrier as well as

193 a likely chlorine level towards the end of the century, on a Montreal Protocol trajectory
194 towards ‘recovery’, according to projected mitigation scenarios for 2100. For stratospheric
195 inorganic bromine, we chose three levels of ~10 ppt, ~15 ppt and ~23 ppt to represent
196 approximately a low (~pre-industrial era, when ice core data suggests a methyl bromide
197 concentration of about 5 ppt (Saltzman et al., 2008)), a medium and a high (~present day)
198 level respectively. These values all include ~5 ppt from VSLS (as described above) with the
199 remaining Br_y coming from long-lived halons and CH₃Br. These values are somewhat
200 arbitrary; our simple aim is to explore a range of Br_y values and, based on them, to
201 investigate the ozone response to perturbed VSLS bromocarbon emissions.

202

203 Our study is designed to address how changes in VSLS bromocarbons would impact
204 stratospheric ozone, against a background of changes in stratospheric chlorine and bromine
205 loading. So, each base experiment is then repeated with the VSLS bromocarbon contribution
206 doubled, to about 10 ppt. In the next section for each base experiment we look at the
207 difference in the ozone results between the 2xVSLS and 1xVSLS (a ~5 ppt increase in
208 stratospheric Br_y) integrations.

209

210 3. Result and Discussions

211

212 Figure 1 shows the annual zonal mean ozone differences between pairs of runs in which the
213 VSLS bromine contribution was increased by 5 ppt. The plots show the 2xVSLS - 1xVSLS
214 ozone differences in percentage. The left column shows ozone changes for a low
215 stratospheric Cl_y of 0.8 ppb, representative either of a pre-industrial atmosphere or after a
216 substantial reduction in halogen loading. The right column has Cl_y of ~3 ppb, roughly
217 representative of the present day. The panels from top to bottom have, respectively, a high
218 inorganic bromine background of ~23 ppt (Figure 1a, 1b), of ~15 ppt (figure 1c, 1d) and a
219 low bromine background of ~10 ppt (figure 1e, 1f).

220

221 For a doubled VSLS emission, with an extra ~5 ppt Br_y in the stratosphere, the ozone
222 differences in these runs are statistically significant (P<0.05) (marked with dots in figure 1) in
223 most of the atmosphere below 20 km. The less significant signals above 20 km are likely due
224 to the strong dynamical feedback rather than chemical response, as addressed in a related
225 paper by Braesicke et al. (2013). Longer runs would be required to ensure statistical
226 significance in the middle and upper stratosphere where the relative changes in ozone are
227 small, and we focus here on the lowermost stratosphere where VSLS has a more important
228 role.

229

230 Below 20 km in the Southern Hemisphere the largest annual mean decrease is about 10%,
231 while in the northern hemisphere high latitudes, the ozone decreases by 4-6%. Near the
232 tropical tropopause, an ozone decline of 2-4% is modelled. Since the ozone concentration
233 near the tropopause is low, these losses are small in absolute terms (<10 ppbv). In terms of
234 the seasonal response, a maximum local ozone reduction of ~20% is simulated in the SH high
235 latitude during spring to early summer. The seasonal variation of column ozone loss in the
236 SH high latitude is discussed below.

237

238 It is interesting to see that ozone decrease is significant in most of the troposphere, with a
239 small deficit of <2% in the tropics; 2-4% in high latitudes of the NH and 2-6% in the SH. In
240 absolute terms, the ozone decline, in most of the free troposphere is smaller than 2 ppb.
241 Further experiments in which we switched off the inter-halogen heterogeneous reactions
242 suggest that the tropical response is due to *in situ* tropospheric chemistry. The changes in

243 tropospheric ozone in high latitudes, in contrast, are largely due to transport of ozone-
244 depleted air from the lower stratosphere.

245

246 When Cl_y is ~ 3 ppb, the maximum SH polar ozone decrease of about 10% shown in figure 1
247 corresponds to ~ 100 ppb, compared with ~ 70 ppb for a Cl_y mixing ratio of 0.8 ppb. In the
248 NH, the ozone decrease is about half of the magnitude simulated in the SH.

249

250 Comparing the left and right panels in Figure 1 for the same bromine loadings, we see greater
251 ozone decrease when the chlorine background is higher. Cycles involving both chlorine and
252 bromine radicals will be more efficient at a given bromine concentration when chlorine levels
253 are higher. To demonstrate this clearly, Fig 2 plots the relationship between the ozone and
254 total inorganic Br_y concentrations at ~ 15 km for (a) $80^\circ S$ and (b) $80^\circ N$ from our experiments
255 with the different chlorine levels. We note that in all cases ozone decreases almost linearly
256 with increasing Br_y concentrations; straight line fits are given for each Cl_y value. The gradient
257 of the ozone decrease as a function of increasing Br_y is larger by about 30-40% when Cl_y is
258 ~ 3 ppb compared with Cl_y of ~ 0.8 ppb. As would be expected given the greater potential for
259 heterogeneous chemistry, there is more ozone loss in the SH than in the NH under the same
260 Cl_y level.

261

262 Comparing the red and blue lines in Figure 2a, we can see that, for a given Br_y level, the
263 ozone concentrations under a pre-industrial (or ozone-recovered) Cl_y of 0.8 ppb are ~ 500 ppb
264 higher than that at current Cl_y of ~ 3 ppb in the SH. According to the fit lines in Figure 2a, for
265 a Br_y increase of 20 ppt (a roughly 4-fold increase of VLS emissions), the ozone decreases
266 are 190 ppb (red line) and 280 ppb (blue line), respectively, considerably less than the
267 chlorine-driven changes. So, even if bromine levels from natural VLS were to increase very
268 significantly later this century, changes in the concentration of ozone will likely be dominated
269 by the recovery of anthropogenic chlorine.

270

271 We have noted above that longer runs would be required to reveal statistically significant
272 changes in the pairwise ozone differences in the middle and upper stratosphere shown in
273 Figure 1. Consequently, changes in column ozone between the different experiments can be
274 somewhat noisy. However, spatial and temporal averaging, as well as the grouping of
275 experiments together, can improve the statistical significance of the column ozone changes.
276 Focusing on the importance of a 5 ppt Br_y increase under the two different chlorine loadings
277 we can achieve a reasonable estimate of the difference in column ozone by averaging the
278 results from the three different Br_y backgrounds for each of the 0.8 ppb and 3.0 ppb Cl_y
279 cases, respectively.

280

281 Figure 3 shows the seasonal evolution of column ozone in high southern latitudes, plotting
282 the average difference when VLS bromine is increased by 5 pppt for each of the two
283 background chlorine loadings. Again, the relative change in ozone is higher under the higher
284 chlorine loading. Under higher chlorine (red line) we model much lower column ozone
285 globally and a larger ozone hole compared with the $Cl_y=0.8$ ppb case (black line). For 3 ppb
286 of chlorine a more pronounced seasonal cycle emerges with the ozone differences peaking in
287 Antarctic spring. In spring (October), the modelled total column ozone decrease is about 9
288 DU for the 0.8 ppb chlorine case and about 11 DU for the 3 ppb case. Column differences in
289 the Arctic are only marginal significant, at best, (see figure 4) so we do not discuss the
290 northern hemisphere seasonal evolution.

291

292 Figure 4 shows annually averaged column ozone differences as a function of latitude. The

293 largest reductions in the column occur in high latitudes, as expected, with a peak reduction of
294 above 8 DU in southern high latitudes under the high chlorine scenario (Figure 4a), which is
295 about 3.5% of the total column ozone (Figure 4b). The changes in figure 4 are smaller at low
296 latitudes (~2 DU or ~1%) and appear to be relatively insensitive to the chlorine background.
297 Note the reduced area of statistical significance in the northern hemisphere (as indicated by
298 the horizontal lines in Figure 4b, see caption for details), especially under low chlorine
299 loading, consistent with much greater dynamical variability there.

300

301 All our integrations have used year 2000 boundary conditions so the impact of climate
302 change on the chemistry here has not been considered. Separate integrations looking at ozone
303 recovery under a range of different greenhouse gas scenarios, but with constant boundary
304 conditions for chlorine and bromine source gases, (see Banerjee et al., 2014) do throw some
305 light on possible changes in some key bromine reaction fluxes. For example, focussing just
306 on changes in the stratosphere, the change in the flux through the reaction $\text{HO}_2 + \text{BrO}$ between
307 2000 and 2100 under RCP8.5 is less than 10% in the lower stratosphere of high southern
308 latitudes, where the calculated changes reported here are highest. Slightly larger changes are
309 found in the tropical very low stratosphere, where ozone concentrations are low, and just
310 above the tropopause in high northern latitudes where dynamically-driven variability in
311 ozone is high. We believe that while climate change will certainly affect the impact of VSLs
312 on ozone recovery it will not change the general results presented here.

313

314 **4. Conclusions and Discussions**

315

316 Naturally produced VSLs bromocarbons account for almost a quarter of the current
317 stratospheric inorganic bromine and about half of that in the pre-industrial era. Fundamental
318 emission processes for the VSLs are not well understood so that we are unsure how
319 emissions might have changed in the past or how they will respond to future climate change.
320 Model calculations do suggest (e.g. Dessens et al, 2009; Hossaini et al, 2013) that transport to
321 the stratosphere, where these compounds play an important role in regulating lower
322 stratospheric ozone, will increase in the future. To what extent possible increases in VSLs
323 surface emissions or transport to the stratosphere could counteract the effect of halogen
324 reductions under the Montreal Protocol is an important research question, which we have
325 addressed here using a simple experimental design. In particular, we have asked to what
326 extent the impact of a hypothetical increase in VSLs (we use 5 ppt here) could affect ozone
327 in the lower stratosphere and how that impact depends on the background concentrations of
328 chlorine and bromine.

329

330 Our model experiments indicate that for a ~5 ppt increase in inorganic bromine from VSLs,
331 the depletion of stratospheric ozone can reach up to 10% in the annual mean in the lowermost
332 stratosphere of the SH polar region and 4-6% in the NH. Decreases in the ozone column are
333 largest in high latitudes, are greater for higher chlorine levels and are dominated by the ozone
334 reductions in the low stratosphere. The southern hemisphere high latitudes see the largest
335 reductions, with an annual average south of 70°S of around 8 DU (Figure 4a); changes in
336 middle and low latitudes are small.

337

338 Ozone losses following an increase in VSLs burden are highest under a high stratospheric
339 chlorine background, indicating the importance of the inter-halogen reactions. Although
340 bromine plays an important role in destroying atmospheric ozone, chlorine concentrations
341 largely dominate the changes in stratospheric ozone concentrations. Even if bromine levels
342 from natural VSLs were to increase significantly later this century, the concentration of

343 ozone will be dominated by the recovery of anthropogenic chlorine.

344

345 It is possible to make a crude estimate of the possible impact on ozone recovery of a
346 hypothetical increase of VLSL bromine. In our calculations, the increase of Br_y by 5 ppt leads
347 to a reduction in modelled springtime (October) Antarctic ozone of about 9 DU for the 0.8
348 ppb chlorine case and ~11 DU for the 3 ppb chlorine case. A range of chemistry-climate
349 models reported in WMO (2011) and Eyring et al. (2010) give a linearly averaged Antarctic
350 spring time (October) ozone recovery rate of about 1.4 DU/yr between the years 2025 and
351 2075 (see Figure 3.11 and Table 4 of those papers, respectively). So, our modelled decrease
352 of 9 and 11 DUs, due to the additional 5 ppt of bromine under two different chlorine levels,
353 correspond to a delayed recovery of about 6 and 8 years, respectively. We note, in passing,
354 that although Antarctic ozone recovery is predicted consistently by models for the second
355 half of this century, there are large model-model differences.

356

357 In these model studies we have only considered the effect of variations in chlorine and
358 bromine concentrations. We recognize that other changes, including changes in climate and
359 aerosol loading, could also modulate the impact on ozone of changed VLSL emissions.
360 Further studies are required.

361

362 **Acknowledgements:** JAP and XY thank the EU for support through SHIVA (SHIVA-
363 226224-EP7-ENV-2008-1) and the ERC for support through the ACCI project (Project
364 number 267760). We thank Tara Banerjee for information on changes in the $\text{HO}_2 + \text{BrO}$ flux
365 with climate change. We thank NCAS-CMS for modelling support. Model integrations have
366 been performed using the UK National Supercomputing Service HECToR.

367

References:

- 368 Aschmann, J., Sinnhuber, B.-M., Chipperfield, M. P., and Hossaini, R.: Impact of deep convection
369 and dehydration on bromine loading in the upper troposphere and lower stratosphere, *Atmos. Chem.*
370 *Phys.*, 11, 2671–2687, doi:10.5194/acp-11-2671-2011, 2011.
- 371 Banerjee, A., A.T. Archibald, A.C. Maycock, P. Telford, N.L. Abraham, Xin Yang, P. Braesicke and
372 J.A. Pyle, Lightning NO_x, a key chemistry-climate interaction: impacts of future climate change and
373 consequences for tropospheric oxidising capacity, *Atmos. Chem. Phys.*, ACP-2104-129, In press,
374 2014.
- 375 Braesicke, P., Keeble, J., Yang, X., Stiller, G., Kellmann, S., Abraham, N. L., Archibald, A. T.,
376 Telford, P., and Pyle, J. A.: Consistent circulation differences in the Southern Hemisphere caused by
377 ozone changes: a chemistry-climate model and observational study, *Atmos. Chem. Phys. Discuss.*, 13,
378 8455-8487, doi:10.5194/acpd-13-8455-2013, 2013.
- 379 Carpenter, L. J. and Liss, P. S.: On temperate sources of bromoform and other reactive organic
380 bromine gases, *J. Geophys. Res.*, 105, D16, 20539-20547, doi: 10.1029/2000JD900242, 2000.
- 381 Chipperfield, M. P., and Pyle, J. A.: Model sensitivity studies of Arctic ozone depletion, *J. Geophys.*
382 *Res.*, 103(D21), 28389–28403, doi:10.1029/98JD01960, 1998.
- 383 Dessens, O., Zeng, G., Warwick, N., Pyle, J.: Short-lived bromine compounds in the lower
384 stratosphere; impact of climate change on ozone, *Atmos. Sci. Lett.* 10, 201–206, doi:10.1002/asl.236,
385 2009.
- 386 Dorf, M., Butz, A., Camy-Peyret, C., Chipperfield, M. P., Kritten, L., and Pfeilsticker, K.: Bromine in
387 the tropical troposphere and stratosphere as derived from balloon-borne BrO observations, *Atmos.*
388 *Chem. Phys.*, 8, 7265–7271, doi:10.5194/acp-8-7265-2008, 2008.
- 389 Eyring, V., Cionni, I., Bodeker, G. E., Charlton-Perez, A. J., Kinnison, D. E., Scinocca, J. F.,
390 Waugh, D. W., Akiyoshi, H., Bekki, S., Chipperfield, M. P., Dameris, M., Dhomse, S., Frith, S. M.,
391 Garny, H., Gettelman, A., Kubin, A., Langematz, U., Mancini, E., Marchand, M., Nakamura, T.,
392 Oman, L. D., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Shepherd, T. G., Shibata, K.,
393 Tian, W., Braesicke, P., Hardiman, S. C., Lamarque, J. F., Morgenstern, O., Pyle, J. A., Smale, D.,
394 and Yamashita, Y.: Multi-model assessment of stratospheric ozone return dates and ozone recovery in
395 CCMVal-2 models, *Atmos. Chem. Phys.*, 10, 9451-9472, doi:10.5194/acp-10-9451-2010, 2010.
- 396 Farman, J. C., Gardiner, B. G., Shanklin, J. D.: Large losses of total ozone in Antarctica reveal
397 seasonal ClO_x/NO_x interaction, *Nature* 315, 207-210, doi:10.1038/315207a0, 1985.
- 398 Feng, W. Chipperfield, M.P., Dorf, M., Pfeilsticker, K and Ricaud, P.: Mid-latitude ozone changes:
399 Studies with a 3-D CTM forced by ERA-40 analyses, *Atmos. Chem. Phys.*, 7, 2357-2369, 2007.
- 400 Gschwend, P. M., Macfarlane, J. K., and Newman, K. A.: Volatile Halogenated Organic-Compounds
401 Released To Seawater From Temperate Marine Macroalgae, *Science*, 227, 1033-1035, 1985.
- 402 Harper, D. B. and Hamilton, J. T. G.: The global cycles of the naturally-occurring monohalomethanes,
403 in: *The Handbook of Environmental Chemistry Vol. 3/P, Natural production of organohalogen*
404 *compounds*, edited by: Gribble, G. W., 17–41, 2003.
- 405 Hewitt, H. T., Copsey, D., Culverwell, I. D., Harris, C. M., Hill, R. S. R., Keen, A. B.,
406 McLaren, A. J., and Hunke, E. C.: Design and implementation of the infrastructure of HadGEM3: the
407 next-generation Met Office climate modelling system, *Geosci. Model Dev.*, 4, 223-253,
408 doi:10.5194/gmd-4-223-2011, 2011.
- 409 Hossaini, R., Chipperfield, M. P., Feng, W., Breider, T. J., Atlas, E., Montzka, S. A., Miller, B. R.,
410 Moore, F., and Elkins, J.: The contribution of natural and anthropogenic very short-lived species to
411 stratospheric bromine, *Atmos. Chem. Phys.*, 12, 371-380, doi:10.5194/acp-12-371-2012, 2012a.
- 412 Hossaini, R., Chipperfield, M. P., Dhomse, S., Ordonez, C., Saiz-Lopez, A., Abraham, N. L.,
413 Archibald, A. T., Braesicke, P., Telford, P. J., Warwick, N. J., Yang, X., Pyle, J. A.: Modelling future
414 changes to the stratospheric source gas injection of biogenic bromocarbons, *Geophys. Res. Lett.*, 39,
415 doi:10.1029/2012GL053401, 2012b.
- 416 Hossaini, R., Mantle, H., Chipperfield, M. P., Montzka, S. A., Hamer, P., Ziska, F., Quack, B.,
417 Krüger, K., Tegtmeier, S., Atlas, E., Sala, S., Engel, A., Bönisch, H., Keber, T., Oram, D., Mills, G.,
418 Ordóñez, C., Saiz-Lopez, A., Warwick, N., Liang, Q., Feng, W., Moore, F., Miller, B. R.,
419 Marécal, V., Richards, N. A. D., Dorf, M., and Pfeilsticker, K.: Evaluating global emission

inventories of biogenic bromocarbons, *Atmos. Chem. Phys. Discuss.*, 13, 12485-12539, doi:10.5194/acpd-13-12485-2013, 2013.

Keppler, F., Eiden, R., Niedan, V., Pracht J. and Schöler H.F.: Halocarbons produced by natural oxidation processes during degradation of organic matter, *Nature*, 403, 298–301, 2000.

Ko, M. K. W., Sze, N.-D., Scott, C. J. and Weisenstein D.K.: On the relation between stratospheric chlorine/bromine loading and short-lived tropospheric source gases, *J. Geophys. Res.*, 102, 25,507–25,517, 1997.

Liang, Q., Stolarski, R. S., Kawa, S. R., Nielsen, J. E., Douglass, A. R., Rodriguez, J. M., Blake, D. R., Atlas, E. L., and Ott, L. E.: Finding the missing stratospheric Br_y: a global modeling study of CHBr₃ and CH₂Br₂, *Atmos. Chem. Phys.*, 10, 2269–2286, doi:10.5194/acp-10-2269-2010, 2010.

McElroy, M. B., Salawitch R.J., Wofsy S.C., Logan, J.A.: Reductions of Antarctic ozone due to synergistic interactions of chlorine and bromine, *Nature*, 321, 759–762, 1986.

Molina, M. J. and Rowland, F. S.: Stratospheric Sink for Chlorofluoromethanes: Chlorine Atom-Catalysed Destruction of Ozone. *Nature*, 249, 810–812, doi:10.1038/249810a0, 1974.

Montzka, S. A. and Reimann, S. et al: Ozone-depleting substances (ODSs) and related chemicals, in: *Scientific Assessment of Ozone Depletion: 2010*, Global Ozone Research and Monitoring Project, Report No. 52, Chapt. 1, edited by: World Meteorological Organization, Geneva, Switzerland, 2011.

Morgenstern, O., Braesicke, P., O'Connor, F. M., Bushell, A. C., Johnson, C. E., Osprey, S. M., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model – Part 1: The stratosphere, *Geosci. Model Dev.*, 2, 43–57, doi:10.5194/gmd-2-43-2009, 2009.

O'Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., Dalvi, M., Folberth, G. A., Sanderson, M. G., Telford, P. J., Young, P. J., Zeng, G., Collins, W. J., Pyle, J. A.: Evaluation of the new UKCA climate-composition model - Part 2: The Troposphere. *Geoscientific Model Development Discussions*, 6 (1), 1743--1857, doi:10.5194/gmdd-6-1743-2013, 2013.

Ordóñez, C., Lamarque, J.-F., Tilmes, S., Kinnison, D. E., Atlas, E. L., Blake, D. R., Sousa Santos, G., Brasseur, G., and Saiz-Lopez, A.: Bromine and iodine chemistry in a global chemistry-climate model: description and evaluation of very short-lived oceanic sources, *Atmos. Chem. Phys.*, 12, 1423–1447, doi:10.5194/acp-12-1423-2012, 2012.

Pyle J. A., Warwick, N., Yang, X., Young, P. J., and Zeng, G.: Climate/chemistry feedbacks and biogenic emissions, *Phil. Trans. R. Soc. A*, 365, 1727-1740, 2007.

Pyle, J.A., Ashfold, M. J., Harris, N. R. P., Robinson, A. D., Warwick, N. J., Carver, G. D., Gostlow, B., O'Brien, L. M., Manning, A. J., Phang, S. M., Yong, S. E., Leong, K. P., Ung, E. H., and Ong, S.: Bromoform in the tropical boundary layer of the Maritime Continent during OP3, *Atmos. Chem. Phys.*, 11, 529-542, doi:10.5194/acp-11-529-2011, 2011.

Quack B, Wallace, D. W. R.: Air–sea flux of bromoform: controls, rates and implications. *Global Biogeochem. Cycles*, 17(1), 1023, doi:10.1029/2002GB001890, 2003.

Salawitch, R. J.: Ozone depletion: a greenhouse warming connection, *Nature*, 392, 551–552, 1998.

Salawitch, R.J., Weisenstein, D.K., Kovalenko, L.J., Sioris, C.E., Wennberg, P.O., Chance, K., Ko, M.K.W. and McLinden, C.A.: Sensitivity of ozone to bromine in the lower stratosphere. *Geophysical Research Letters*, 32: doi: 10.1029/2004GL021504, 2005.

Salawitch, R. J.: Atmospheric chemistry - Biogenic bromine, *Nature*, 439, 275-277, 2006.

Salawitch, R. J., Canty, T., Kurosu, T., Chance, K., Liang, Q., da Silva, A., Pawson, S., Nielsen, J. E., Rodriguez, J. M., Bhartia, P. K., Liu, X., Huey, L. G., Liao, J., Stickel, R. E., Tanner, D. J., Dibb, J. E., Simpson, W. R., Donohoue, D., Weinheimer, A., Flocke, F., Knapp, D., Montzka, D., Neuman, J. A., Nowak, J. B., Ryerson, T. B., Oltmans, S., Blake, D. R., Atlas, E. L., Kinnison, D. E., Tilmes, S., Pan, L. L., Hendrick, F., Van Roozendaal, M., Kreher, K., Johnston, P. V., Gao, R. S., Johnson, B., Bui, T. P., Chen, G., Pierce, R. B., Crawford, J. H., and Jacob, D. J.: A new interpretation of total column BrO during Arctic spring, *Geophys. Res. Lett.*, 37, L21805, doi: 10.1029/2010gl043798, 2010.

Saltzman, E.S., M. Aydin, C. Tatum, and M.B. Williams, 2,000-year record of atmospheric methyl bromide from a South Pole ice core, *J. Geophys. Res.*, 113, D05304, doi: 10.1029/2007JD008919, 2008.

Schofield, R., Fueglistaler, S., Wohltmann, I., and Rex, M.: Sensitivity of stratospheric Br_y to uncertainties in very short lived substance emissions and atmospheric transport, *Atmos. Chem. Phys.*, 11, 1379–1392, doi:10.5194/acp-11-1379-2011, 2011.

475 Sinnhuber, B. M., Sheode, N., Sinnhuber, M., Chipperfield, M. P., and Feng, W.: The contribution of
476 anthropogenic bromine emissions to past stratospheric ozone trends: A modelling study, *Atmos.*
477 *Chem. Phys.*, 9(8), 2863–2871, 2009.

478 Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37,
479 275-316, doi:10.10291999RG900008, 1999.

480 Stachnik, R. A., Millán, L., Jarnot, R., Monroe, R., McLinden, C., Kühl, S., Puķite, J., Shiotani, M.,
481 Suzuki, M., Kasai, Y., Goutail, F., Pommereau, J. P., Dorf, M., and Pfeilsticker, K.: Stratospheric BrO
482 abundance measured by a balloon-borne submillimeterwave radiometer, *Atmos. Chem. Phys.*, 13,
483 3307-3319, doi:10.5194/acp-13-3307-2013, 2013.

484 Sturges, W., Oram, D., Carpenter, L., Penkett, S., and Engel, A.: Bromoform as a source of
485 stratospheric bromine, *Geophys. Res. Lett.*, 27(14), 2081–2084, doi:10.1029/2000GL011444, 2000.

486 Tegtmeier, S., Krüger, K., Quack, B., Pisso, I., Stohl, A., and Yang, X.: Emission and transport of
487 bromocarbons: from the West Pacific ocean into the stratosphere, *Atmos. Chem. Phys.*, 12, 10633-
488 10648, doi:10.5194/acp-12-10633-2012, 2012.

489 Warwick, N. J., Pyle, J. A., Carver, G. D., Yang, X., Savage, N. H., O'Connor, f. M., and Cox, R. A.:
490 Global modeling of biogenic bromocarbons, *J. Geophys. Res.*, 111, D24305,
491 doi:10.1029/2006JD007264, 2006.

492 WMO (World Meteorological Organization), *Scientific Assessment of Ozone Depletion: 2010*, Global
493 Ozone Research and Monitoring Project-Report No. 52, 516 pp., Geneva, Switzerland, 2011.

494 Wofsy, S., McElroy, M., and Yung, Y.: Chemistry of atmospheric bromine, *Geophys. Res. Lett.*, 2(6),
495 215–218, doi:10.1029/GL002i006p00215, 1975.


496 Yang, X., Cox, R. A., Warwick, N. J., Pyle, J. A., Carver, G. D., O'Connor, F. M., and Savage, N. H.:
497 Tropospheric bromine chemistry and its impacts on ozone: A model study. *J. Geophys. Res.*, 110,
498 D23311, doi:10.1029/2005JD006244, 2005.

499 Yang, X., Pyle, J. A., Cox, R. A., Theys, N., Van Roozendaal, M.: Snow-sourced bromine and its
500 implications for polar tropospheric ozone, *Atmos. Chem. Phys.*, 10, 7763-7773, doi:10.5194/acp-10-
501 7763-2010, 2010.


502 Yokouchi, Y., Hasebe, F., Fujiwara, M., Takashima, H., Shiotani, M., Nishi, N., Kanaya, Y.,
503 Hashimoto, S., Fraser, P., Toom-Sauntry, D., Mukai, H., and Nojiri, Y.: Correlations and emission
504 ratios among bromoform, dibromochloromethane, and dibromomethane in the atmosphere, *J.*
505 *Geophys. Res.-Atmos.*, 110, D23309, doi:10.1029/2005JD006303, 2005.

506 Ziska, F., Quack, B., Abrahamsson, K., Archer, S. D., Atlas, E., Bell, T., Butler, J. H., Carpenter, L.
507 J., Jones, C. E., Harris, N. R. P., Hepach, H., Heumann, K. G., Hughes, C., Kuss, J., Krüger, K., Liss,
508 P., Moore, R. M., Orlikowska, A., Raimund, S., Reeves, C. E., Reifenhäuser, W., Robinson, A. D.,
509 Schall, C., Tanhua, T., Tegtmeier, S., Turner, S., Wang, L., Wallace, D., Williams, J., Yamamoto, H.,
510 Yvon-Lewis, S., and Yokouchi, Y.: Global sea-to-air flux climatology for bromoform,
511 dibromomethane and methyl iodide, *Atmos. Chem. Phys. Discuss.*, 13, 5601–5648, doi:10.5194/acpd-
512 13-5601-2013, 2013.


Change in zonal Mean Ozone(%): 2X-1XVLSL Annual





515
516 Figure 1 Relative ozone changes (in %) arising from doubling VLSL emissions under 0.8 ppb
517 Cl_y (left column) and 3 ppb Cl_y (right column). The base case Br_y concentrations (with
518 1xVLSL, representing 5ppt of the total Br_y) increase from the bottom to the top row as
519 indicated in the panels. The dotted regions represent statistically significant signals on a
520 confidence level above 95% as calculated by Student's t-test.
521



522
 523 Figure 2 Relationship between stratospheric ozone concentrations and Br_y at ~15 km and (a)
 524 80°S and (b) 80°N. Straight lines have been fitted for each Cl_y concentration (red:
 525 Cl_y=0.8ppb; blue: Cl_y=3 ppb) and the quality of the fit is indicated by the correlation
 526 coefficient (r²).
 527
 528
 529
 530
 531
 532



533
 534 Figure 3 Relative change of column ozone (in %) due to a doubling of VLS Br_y averaged
 535 over 60°S-90°S. Each line represents one particular Cl_y level (black-open circles: Cl_y=0.8
 536 ppb; red-closed circles: Cl_y= 3 ppb). Note that we average the ozone difference over the three
 537 different Br_y concentrations for each Cl_y case. The individual results for different background
 538 Br_y are shown in Figures 1 and 2.



539
 540 Figure 4 Latitude distributions of annual mean total ozone changes due to a doubling of
 541 VLSLs. The top panel (a) shows the absolute changes in Dobson Units and the bottom panel
 542 (b) shows the relative changes in %. Each line represents one particular Cl_y level (black-open
 543 circles: $Cl_y=0.8$ ppb; red-closed circles: $Cl_y=3$ ppb). Note that we average the ozone
 544 difference over the three different Br_y concentrations for each Cl_y case. Corresponding
 545 horizontal lines indicate where the differences are significant in each of the three pairs of
 546 integrations above a confidence level of 99% as calculated by Student's t-test.
 547
 548