

[Interactive  
Comment](#)

# ***Interactive comment on “Constraining terrestrial ecosystem CO<sub>2</sub> fluxes by integrating models of biogeochemistry and atmospheric transport and data of surface carbon fluxes and atmospheric CO<sub>2</sub> concentrations” by Q. Zhu et al.***

**Q. Zhu et al.**

qzhu@lbl.gov

Received and published: 11 January 2015

1. Why was AIRS chosen, rather than a NIR sensor (GOSAT, SCIAMACHY) with greater sensitivity to the lower troposphere?

Response: We chose AIRS CO<sub>2</sub> data over GOSAT or SCIAMACHY due to the consideration of (1) spatial coverage and (2) data accuracy. SCIAMACHY CO<sub>2</sub> products have the smallest spatial coverage, roughly 8-20% of the global surface. The measurements over ocean are filtered out due to low surface reflectance. GOSAT CO<sub>2</sub>

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



products roughly have 32-44% spatial coverage, because that the areas with aerosol optical depth less than 0.5 are eliminated. In comparison with these two products, AIRS CO<sub>2</sub> products have much higher spatial coverage (about 95%). We used satellite CO<sub>2</sub> retrievals aiming to compensate the low spatial coverage of GLOBALVIEW-CO<sub>2</sub> network. Therefore, AIRS CO<sub>2</sub> is the most appropriate product for our purpose. Validated against a global network of ground-based high-resolution CO<sub>2</sub> measurements (TCCON network), SCIAMACHY data are systematically lower; GOSAT CO<sub>2</sub> retrievals have large random errors due to instrumental noise; the mean difference between AIRS and TCCON is small. Collectively, compared with surface CO<sub>2</sub> measurements, the relative accuracy of GOSAT, SCIAMACHY and AIRS CO<sub>2</sub> are 0.38%, 0.87%, and 0.25%, respectively (Miao et al., 2013).

2. What measurement and model/representativity uncertainties were assumed for the GLOBALVIEW and AIRS measurements used to constrain the fluxes?

Response: GLOBALVIEW-CO<sub>2</sub> observation errors are from data product (GLOBALVIEW-CO<sub>2</sub> 2013). The errors are roughly 0.5 ppm including the instrumental error and errors from the GLOBALVIEW data fitting procedure. The representation error (inability of transport model to represent the observed site location) is not considered. A previous study implied that the representation error is about 0.3 ppm (Baker et al., 2006). AIRS CO<sub>2</sub> errors are from AIRS CO<sub>2</sub> level-2 dataset version 5 (Susskind et al., 2011). A two (two adjacent FOVs) by two (two adjacent scan lines) array of AIRS CO<sub>2</sub> retrieval is used to determine the final retrieval of CO<sub>2</sub> concentration. The error represents the spatial coherence over the 2 by 2 array. We only used the level 2 “standard product”, in which the errors are less than 2 ppm. The CO<sub>2</sub> retrievals with errors larger than 2 ppm are placed in level 2 “support product”, which was not used. However, the representation error is not considered in the AIRS CO<sub>2</sub> level 2 products.

3. Are the prior and posterior fluxes only at monthly time scales? If so, why, and what effect might this have on the evaluation in concentration space? What time step was

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

the model running at?

Response: The model used to estimate the prior surface CO<sub>2</sub> fluxes has a monthly time step. And our prior and posterior fluxes are also at monthly time scale. The low temporal resolution of surface CO<sub>2</sub> flux is one of the potential limitations of our study. It limited our ability to capture the fine-scale (e.g. daily, diurnal) variation of CO<sub>2</sub> concentration. Therefore, we were only able to compare and validate our posterior CO<sub>2</sub> concentrations against monthly averaged CO<sub>2</sub> concentration data (both GLOBALVIEW-CO<sub>2</sub> and CONTRAIL).

4. Why only 2003? This makes it difficult to compare to other flux estimates, in many cases, as the inter-annual variability can be quite substantial. Also, was there a spin-up or spin-down time with the atmospheric inversion? Or is this not an issue with the specific 4D-Var method used here? I would strongly recommend expanding the analysis to a longer time period, even just a few years, in order to provide a better basis for analysis and comparison.

Response: We limited our study period in 2003 mainly because we aimed to illustrate the methodology and effectiveness of combining transport inversion and ecosystem model parameter inversion in estimating global NEP. The research question is whether or not an improved NEP prior (based on TEM model data assimilation) would benefit the estimate of posterior NEP. And whether or not the sequential data assimilation approach is a good way to carry out CO<sub>2</sub> inversion study? Therefore, it doesn't really matter which year we are conducting for our inversion and analysis. We agree with the reviewer's concern that limiting our study period in 2003 made us difficult to compare our inversion results with others. In this revision, we have applied the CO<sub>2</sub> inversion to a longer period to analyze the long-term trend and inter-annual variability of global NEP. However we decide not to show the results in this study, because it conflicts to our future publication plan regarding to the long-term trend and interannual variability of global NEP. Therefore, we only provided some results of our posterior NEP for 2004 and 2005 in this response letter, but not in the manuscript. Figure 1 shows the posterior

C11094

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



NEP over eleven TRANSCOM 3 land regions for 2004 and 2005. The annual total terrestrial ecosystem carbon sinks are  $-2.6 \text{ Pg C yr}^{-1}$  and  $-2.7 \text{ Pg C yr}^{-1}$ . In both 2004 and 2005, North American temperate is the largest carbon sink followed. North American Boreal, Europe, Eurasian Boreal, and South American Tropical regions are secondary largest carbon sinks. It is interesting to see that the South American tropical region is a much stronger carbon sink in 2005 compared with that in 2003 and 2004. It coincides with the Amazonia green-up during 2005 (Saleska et al., 2007). It implied that the atmospheric CO<sub>2</sub> concentration signals were able to reflect the enhanced land carbon sink for this region.

5. The approach holds a great deal of promise, but this attempt is rather heavily weighted towards the bottom-up side of things. The analysis does little to inform me of the added value of using the surface stations and satellite measurements. Perhaps it would be worthwhile to compare inversions using only the surface network with those using AIRS as well, to see what value the mid-tropospheric information brings to surface processes.

Response: In this revision, we conducted an additional CO<sub>2</sub> inversion using only GLOBALVIEW-CO<sub>2</sub> dataset. Comparing the differences of NEP among (1) prior; (2) posterior NEP using only GLOBALVIEW-CO<sub>2</sub>; and (3) posterior NEP using both GLOBALVIEW-CO<sub>2</sub> and AIRS CO<sub>2</sub>, we are able to show the added values of assimilating GLOBALVIEW-CO<sub>2</sub> and AIRS CO<sub>2</sub> datasets (Figure 2). We find that regional NEP is more effectively constrained over North America and Europe, where in situ measurements are abundant. GLOBALVIEW-CO<sub>2</sub> data generally provide more added values to posterior NEP, compared with AIRS CO<sub>2</sub> data. AIRS CO<sub>2</sub> data are most beneficial for European regional NEP, they provide almost equal added value to prior NEP, compared with GLOBALVIEW-CO<sub>2</sub> data. We added a paragraph and Figure 2 to the manuscript in this revision.

7. Here you're adding the land-use change flux to surface fluxes constrained by atmospheric measurements to argue that there's good agreement. But in both cases

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

(Peylin et al., 2013 and the current study) the atmospheric measurements have "seen" the land-use change signal, and this should be reflected (however erroneously) in the optimized fluxes, provided the prior uncertainty isn't too tight. Or am I missing something?

Response: We agree that the observed atmospheric CO<sub>2</sub> signals contain information about all the land surface CO<sub>2</sub> activities including the land-use induced CO<sub>2</sub> emissions. The definitions of natural land CO<sub>2</sub> flux are different in our study and Peylin 2013. Our inverted natural land fluxes only contain Net Ecosystem Production (NEP). The land-use induced CO<sub>2</sub> emission was considered as a part of fire emission, which had been subtracted before we inverted the NEP. Peylin et al. (2013)'s study defined that the natural land fluxes are the summation of NEP and land-use related fluxes (usually fires emission). Therefore, we need to add the land-use induced CO<sub>2</sub> emission back into our posterior NEP, in order to be comparable with the posterior natural land fluxes reported by Peylin et al. (2013).

## REFERENCE

Baker, D. F., Law, R. M., Gurney, K. R., Rayner, P., Peylin, P., Denning, A. S., ... & Zhu, Z. (2006). TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO<sub>2</sub> fluxes, 1988–2003. *Global Biogeochemical Cycles*, 20(1). Miao, R., Lu, N., Yao, L., Zhu, Y., Wang, J., & Sun, J. (2013). Multi-Year Comparison of Carbon Dioxide from Satellite Data with Ground-Based FTS Measurements (2003–2011). *Remote Sensing*, 5(7), 3431-3456. Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., Niwa, Y., Patra, P. K., Peters, W., Rayner, P. J., Rödenbeck, C., van der Laan-Luijkx, I. T., and Zhang, X.: Global atmospheric carbon budget: results from an ensemble of atmospheric CO<sub>2</sub> inversions, *Biogeosciences*, 10, 6699-6720, doi:10.5194/bg-10-6699-2013, 2013. Saleska, S. R., Didan, K., Huete, A. R., & Da Rocha, H. R. (2007). Amazon forests green-up during 2005 drought. *Science*, 318(5850), 612-612.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper




Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



**Fig. 1.** 2004 and 2005 posterior NEP after CO<sub>2</sub> inversions over 11 TRANSCOM3 defined land regions.


[Interactive  
Comment](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



**Fig. 2.** Bars are prior. Stars (\*), circles (o) represent the posterior mean of NEP updated by CO<sub>2</sub> inversion after assimilating only GLOBALVIEW-CO<sub>2</sub> dataset and GLOBALVIEW-CO<sub>2</sub> + AIRS CO<sub>2</sub> datasets, respectively.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)