

Interactive comment on “The oxidation capacity of the boreal forest: first simulated reactivities of O₃ and NO₃” by D. Mogensen

Anonymous Referee #2

Received and published: 17 February 2015

This manuscript by Mogensen et al. describes first simulation to evaluate oxidation strength by O₃ and NO₃ in a boreal forest using 1D atmospheric chemistry transport model (SOSAA; model to Simulate Organic vapours, Sulphuric Acid and Aerosols). The authors concluded that OH is the most reactive and followed by NO₃ and O₃, while O₃ is the fastest remover for pollutants when considering ambient levels of each oxidant. I do not agree with this though, as specifically commented below. They also found some missing OH reactivity due to unknown BVOCs and their oxidation products.

Overall, I truly appreciate that the authors' effort to study unexplored area and draw the attention about importance of NO₃ and O₃ reactivities in the forest environment to the atmospheric chemistry community. However, this manuscript is still poorly written and needs to be improved in many parts of the sections, particularly in the discussion

C12243

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



of the reactivities of oxidants by organics. Otherwise, this cannot be published in ACP. My comments and suggestions are below.

General Comments:

The title seems that this MS intended to focus on O₃ and NO₃ reactivities. Throughout the MS, however, description and discussion of OH and NO₃ reactivities are much more allocated than O₃ reactivity. In addition, reactivities of each oxidant by reacting with organics are poorly explained despite of their importance in the forest area. One big concern is that the authors did not use BVOC measurement data or did not justify BVOC input data when comparing the modeled value by MEGAN to any other available data at the site or to the similar vegetation types. This may cause substantial error in reactivity calculation. As the referee #1 pointed out, I agree that discrepancy of modeled vs observed OH reactivity needs to be thoroughly discussed. As well as, during daytime NO+O₃ → NO₂ +O₂ reaction is null cycle since O₃ returns back by NO₂ photolysis, so it is hard to say that O₃ is reacted with NO. To improve this, I suggest focusing more on organic compounds than inorganic compounds. Moreover, the lifetime of oxidant is equivalent to the inverse of reactivity, thus lifetime of O₃ is estimated ~13 hours based on the result in the abstract. This is considerably slow and not plausible in the forest environment if trees emit BVOCs. Further, when considering NO+O₃ reaction as null cycle, the O₃ lifetime is going to be even much slower. This is also required to discuss in the MS.

Specific comments:

Abstract: Overall, this part should be rewritten after revision of the MS.

- P.30948, L23: 'removing pollutants fastest', this sounds like inappropriate since O₃ also reacts with BVOC as mentioned previously in the MS, as well as O₃ by itself is generally not the fastest oxidant to remove either organic or inorganic compounds.

-P.30949 L18-19: What do you mean by 'the measurement is tricky'? Be specific here

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



and need good references.

-P.30950 L12-14: How did you come with '5% of the global emission of BVOCs'? Did you only account for the boreal forests or together with the temperate forests? Clarify this since you mentioned both in the previous sentence.

-P.30951 Section 2: Briefly describe the general information for each site such as vegetation type, climate, wind pattern, anthropogenic influences, and etc. . . Section 3.1-3.3: Even though table 1 shows some information for measured gases, it does not give any information for diurnal profile. So, I suggest describing general diurnal patterns for each gas at the site or generating some figures similar as figure 7 for measured gases.

-P.30955 L6-8: Explain what method you used to calculate uncertainty or add a reference.

-Section 4.4: What was the measurement uncertainties for NO₃ and N₂O₅?

-Section 5: How was treated the deposition in the model (e.g. surface removal and stomatal uptake)? In addition, how does the SOSAA deal with partitioning between gas and particle?

-Section 5.2: This section is poorly written and need to be more specific because BVOCs are presumably very important source in reactivity calculation in the forest environment. So, please describe a little more in detail how you estimated LAI or add a reference, and explain a bit more about vegetation composition such as dominant tree species and understory plants. In order to calculate reactivities of oxidants by reacting with BVOCs, mixing ratios of each BVOC at different heights are necessary, so describe how the BVOC mixing ratios were considered in detail rather than just stating use of MEGAN. Also, it needs to be discussed whether or not the BVOC input data used in the model were reliable when comparing them to the measurement data from the same site (or other similar sites, if not applicable).

-P.30960 L13: 'MCM v3.2' is mentioned first here, so give the full name here than in

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the section 5.4.

-P.30960 L22-23: Justify or add a reference why you used a constant mixing ratio of H₂ and CH₄.

-P.30969 L25-26: Discuss the reason why the time window of 7 pm-8pm was chosen to validate the turbulence. Also discuss what result was come out if you do this for daytime or at nighttime. Because 7pm should be nearly neutral condition of the air and it apparently cannot be representative for turbulence scheme.

-P.30972 L24-26: This sentence is not true, particularly for daytime measurement which apparently shows higher at ROH in-canopy than above-canopy in figure 7.(b). Moreover, modeled ROH result shows consistent lower value at above-canopy in figure 7.(c). Improve this appropriately.

-P.30973 L3-10: It is not clear whether or not the same size of data set was used to compare 'Measured' vs 'Modelled'. In figure 7. (a), it indicates maximum 10-12 days of measurement data were used for one month campaign period. Is it the same for the model? Please clarify this. In case they used different set, justify how those can be compared each other.

-P.30973 L22-23: Give specific number of isoprene concentration range rather than stating just 'very low'.

-P.30979 L6-8: As explained in general comment, with given number of O₃ reactivity, the lifetime of O₃ can be calculated by inverse number of reactivity, thus O₃ lifetime is estimated by ~11 hours during daytime and ~5.8 days at night. This is pretty slow, particularly at night. Was there decreasing trend in nighttime ozone concentration? If so, discuss what major processes of ozone removal take place at the site. Also, provide diurnal trend of O₃ and NO during the campaign. I think the model only accounts for very limited number of chemical reactions with O₃, so I believe there should be missing parts of O₃ reactivity for unknown/unaccounted compounds and reactions. At

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the end of this section, even though possibility of underestimation is mentioned, authors should thoroughly discuss about how much of reactivities they expect to underestimate. Otherwise, this whole manuscript will be less meaningful in terms of O₃ reactivities.

-P.30981 L1: As mentioned above, O₃ molecule by itself is not the fastest oxidant, but O₃ removes reactants much more than other oxidants do because it is the most abundant oxidant at the site. State this very clearly.

-P.30982 L4-5: Explain in detail how OH and NO₃ are reformed and why not for O₃?

-P.30982 L5-8: Again, rephrase 'how fast can the oxidant remove pollutants from the atmosphere',. 'pollutants' is not appropriate here since BVOCs are not the pollutants.

-P.30982 L11-13: With this sentence, it seems authors think opposite way to the rate of reaction between OH and O₃. The statement 'though it is only capable of doing it at a slow rate.' is generally wrong except for a few very reactive compounds with O₃ (e.g. beta-caryophyllene). For example, when comparing reaction rate for isoprene and most monoterpenes, the reaction with OH is much faster than O₃.

-P.30982 L16-19: I do not agree with this statement, because O₃ can react with the second or even higher generation products from primary reactants. Methacrolein and methyl-vinyl-ketone are good example since those are the second products from isoprene oxidation and those can react with O₃, though the reaction rates are relatively much slow with either isoprene or MVK&MACR compared to reaction with OH. The reason why O₃ has much less number of reactions than OH does is due to the reaction mechanism by O₃ for a variety of compounds has not broadly studied yet, particularly for high order generation products from organic compounds.

-P.30983 L26: Again, the word 'fast' is improper. Please consider it to replace with a relevant word or rephrase it.

-Section 8: This part needs to be rewritten after revision.

Table 1: I suggest replacing this table to diurnal profile plots for each compound.

C12247

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Table 3: List more specified compounds for monoterpenes and sesquiterpenes used to the model and show each reactivity rather than using total MTs and SQTs.

Figure 6: It is not easy to compare. I suggest making this figure to 1:1 scatter plots with regression curves. (e.g. in a figure, Meas vs TUV with black dots and Meas vs MODEL with red triangles.)

Figure 8: I suggest generating similar area plots that show contribution of each organic compound used in the model in addition to this plot. With this figure alone, it seems inorganic compounds are more important in the reactivities of oxidants.

Interactive comment on Atmos. Chem. Phys. Discuss., 14, 30947, 2014.

ACPD

14, C12243–C12248,
2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C12248

