

Interactive
Comment

Interactive comment on “Interactions of bromine, chlorine, and iodine photochemistry during ozone depletions in Barrow, Alaska” by C. R. Thompson et al.

A. Bogdan

anatoli.bogdan@uibk.ac.at

Received and published: 25 February 2015

It is believed that PSCs are formed by freezing aqueous aerosol drops which contain acids up to 30 wt %. Since ice is highly intolerant to impurities, after freezing mixed-phase particles are formed: an ice core enveloped with a freeze-concentrated solution (FCS). Since the rate of heterogeneous reactions depends on the surface phase state of PSC particles, there will be difference whether solid pure ice or mixed-phase PSC particles are considered in model(s). If the current models cannot take into account this important physical process (phase separation during freezing), it would be a good thing to mention it in introduction of the paper. Also it would be a good thing to give

C12390

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



some reasoning why it is not considered in the model used by the authors. About mixed-phase PSC particles the authors can find in a paper Bogdan, A., Molina, M. J., Tenhu, H., Mayer, E. & Loerting, T. 2010. "Formation of mixed-phase particles during the freezing of polar stratospheric ice clouds" Nature Chem., 2, 197-201.

Interactive comment on Atmos. Chem. Phys. Discuss., 14, 28685, 2014.

ACPD

14, C12390–C12391,
2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C12391

