

Interactive
Comment

Interactive comment on “Hygroscopic properties and mixing state of aerosol measured at the high altitude site Puy de Dôme (1465 m a.s.l.), France” by H. Holmgren et al.

H. Holmgren et al.

helene.holmgren@gmail.com

Received and published: 20 June 2014

We thank reviewer 2 for his numerous comments and remarks which greatly helped to improve the manuscript. We have done our best to follow the suggestions made, and our answers are reported below.

General comments: The authors present humidified tandem differential mobility analyzer (HTDMA) measurements and hygroscopic growth factors (GF) for aerosol sampled at the French high-altitude research station, Puy de Dome. Measurements are reported for September of 2008, December–May 2009, September–February 2010, and

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



continuously from January 2011 to December 2012. The stated goal of this paper is to explore how aerosol hygroscopic growth factors vary year-to-year, seasonally, diurnally, and with air mass type. Given the incomplete dataset for 2008–2012 it is hard to draw definite conclusions regarding whether the observed differences in GF are due to seasonally-varying aerosol characteristics or due to an analysis approach biased by inclusion or lack of data for a given season in a given year. A revised manuscript should limit the scope of the dataset to ensure an unbiased analysis. In addition, a number of speculative conclusions are drawn from the HTDMA data regarding aerosol composition and emissions source attribution that are currently not supported (some instances noted under specific comments below); if these observations are based on additional measurement data, they should be discussed more fully or otherwise removed. Finally, this manuscript would be greatly improved by including all of the measured HTDMA sizes (given the non-linear nature of the observed hygroscopicities) and by a more extensive and sophisticated discussion of aerosol mixing state that incorporates the GF “spread” of each mode.

Reply: Regarding the non-continuous aspect of the data set; previous to the publication of the paper in the present form, we actually did an analysis of the seasonal variation of hygroscopic aerosol properties over the time period ranging from January 2011 to December 2012, excluding the years 2008, 2009 and 2010, which were less complete. The average seasonal variation obtained using this restricted time period were very similar to the one obtained with the whole data set (and also undistinguishable from the seasonal variation obtained over the years 2011–2012), so we decided to keep all data in the submitted version. However, we agree that we can not draw robust conclusions on a year-to-year variability when using 2008, 2009 and 2010 that do not contain sufficient data. Consequently, we have modified the figures and text of the paper excluding the 2008–2010 period regarding the year-to-year variability. Because the year to year variability is shown to be small, we will keep them in the analysis when discussing the air mass type dependency, and all analysis that are split by season.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Regarding the number of studied particle diameters; we will follow the reviewer recommendation of showing the behaviour of all diameters in the Su et al. (2010) manner in the first general figures showing yearly and seasonal averages. However, for the mixing state quantifications and air mass analysis, the paper would be too heavy if we showed the behaviour of all diameters, and we wish to continue showing the behaviour of particles whose diameters are representative of the nucleation, Aitken and accumulation modes usually observed at the puy de Dôme station (Venzac et al. 2009). We believe studying intermediate diameter would make the study too heavy, without bringing new information. Nonetheless, figures 2 and 4 have been re-plotted, using the approach proposed by the reviewer (Su et al. 2010), see also comment and reply 3.

Finally, the GF spread information is already included in our analysis, by the fact that when the GF spread is higher than expected ($\sigma>0.10$) from the DMA transfer alone, two different hygroscopic modes are searched to explain this spread. This is similar to the analysis by Sjögren et al. (2008), who suggested the presence of two modes when the spread in the GF-PDF was greater than 0.15. The aspect of this type of analysis, with a strict mode fitting procedure, enables a more accurate idea of the aerosol mixing state. We have stressed this point in the analysis section of our paper.

Specific comments:

Comment 1: HTMDA measurements were made at PdD at six sizes: 25, 35, 50, 75, 110, and 165 nm, but only measurements for three sizes (25, 50 and 165 nm) are reported. Given the similar GF-derived kappas reported for 25 and 50 nm particles in Table 7 and the much higher kappa reported for 165 nm, there is clearly a non-linear size-dependence to kappa. Thus, it is essential that the authors add the other 3 measured diameters to the manuscript in order to provide greater size resolution in understanding how the aerosol composition is varying.

Reply 1: See the answer to this in the general comment above. We know include two additional figures of kappa PDF as a function of particle size for all particles measured.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



A small section is added to the text:

"The size dependant hygroscopicity solely due to the aerosol composition can be addressed by plotting K-PDF as a function of the aerosol diameters, calculated for 25, 35, 50, 75, 110, and 165 nm (Figure 2b) using the hygroscopicity distribution concept as described by Su et al., 2010. Both during 2011 and 2012, the aerosol hygroscopicity is increasing with increasing size independently of the Kelvin effect from 50 nm to larger sizes. There is also a tendency of the nucleation mode particles to be slightly more hygroscopic than the Aitken mode particles. Years 2011 and 2012 are very similar on average, with more dispersion in the hygroscopicity of the accumulation mode particles during 2012."

Comment 2: In section 3.1 and Figures 2-7 it does not seem appropriate to include a single month as the annual average for 2008 and half a year for the annual average in 2009 and 2010. This only leaves two years for assessing "year-to-year" annual average variability, which obviously cannot lead to very robust conclusions. It would make sense to me to focus on reporting average aerosol properties over the continuous 2011-2012 time period rather than trying to identify an inter-annual trend from this dataset. Similarly, the "unbalanced seasonal sampling" identified on Pg. 6767, Lines 17-18 is a major flaw in this analysis that it must be corrected, perhaps by again focusing on the only two continuous years 2011-2012 or by bringing in only the winter data from 2009-2012. As it currently stands, definitive conclusions cannot be drawn from regarding GF month-to-month variability.

Reply 2: The year-to-year analysis now only includes the continuous sampling years 2011-2012.

Comment 3: Given the large amount of size-resolved GF data being compared in the tables and in Figures 2 and 4, I recommend that the authors report the data using the hygroscopicity distribution concept as described by Su et al., *Atmos. Chem. Phys.*, 2010 (doi:10.5194/acp-10-7489-2010). Cumulative H (kappa, Dd) vs. Kappa curves

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

facilitate the same comparisons but would also more easily convey the contribution of non-hygroscopic aerosols as well as the means and spreads of the hygroscopic modes. Similarly, since there are six different HTDMA sizes, interpolated color maps similar to those in Figure 9 of Su et al. would be an especially nice way to display this dataset.

Reply 3: Figures of aerosol hygroscopicity (GF and kappa) as a function of size, similar to those presented in Su et al., 2010, now complete figures 2 and 4 (see reply 1).

Comment 4: I'd like to see a much more extensive discussion of aerosol mixing state beyond the current method, which defines an external mixture as when two or more modes are present. For example, the spread of each mode also contains information about mixing state, and should be more fully discussed. In the end how representative is an average GF reported in these tables (and those in other studies) of the individual aerosol particles typically present at this site?

Reply 4: See answer to the general comment. This information contained in the spread of each mode is taken into account by our method. We discuss the frequency with which more than one mode is observed (determined also from the GF spread), but also quantify the number fraction in each mode.

Comment 5: Please add kappa axes to Figures 1-7, 10-11 that correspond to the GF axes so that the reader is able to easily see the compositional differences across aerosol sizes without having to account for the (admittedly small) Kelvin size dependence of the GF.

Reply 5: Kappa axes have been added to the Figures.

Comment 6: Pg. 6762, Line 25-27: This statement is untrue and should be stricken. See for example, Sjögren et al., 2008, which is cited in the previous paragraph, who made measurements at Jungfraujoch during multiple, 1-month, campaigns spanning between different seasons in 2000-2005. Kammermann et al., 2010 (also cited in the previous paragraph) made measurements at Jungfraujoch for a continuous 13-month

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



period.

Reply 6: The sentence has been changed to ‘This is, to the authors’ knowledge, the first study that investigates hygroscopic properties measured at a high altitude site almost continuously over more than two years’.

Comment 7: Pg. 6763, Lines 5-6: I don’t understand the meaning of the definitions “background site” and “high altitude site” in the context of the citations and the HTDMA measurements. A description of what characteristics go into these labels would be preferable.

Reply 7: Asmi et al. 2011 define a high altitude site as a site at more than 1000 m a.s.l, and place Puy de Dôme in that category. “Background site” should be “rural background site”, which is defined as site with a distance 10-50 km from large pollution sources by Putaud et al. 2004. This has been corrected in the text. The reference Venzac et al. 2009 is removed.

Comment 8: Pg. 6764, Lines 15-19: I don’t understand how the authors are calibrating their DMAs using ammonium sulphate. I would think that this is not really a calibration, but rather a consistency check of the instrument operation using a known pure-component salt. This should be removed or, if I’m mistaken and a calibration was performed, more details are needed.

Reply 8: We are actually evaluating the accuracy of the RH sensors in the HTDMA. Since we know at a given RH how much ammonium sulphate should grow, we adjust the real RH attributed to the measurement method using this in the Gysel software (Gysel et al. 2009). This has been clarified in the text.

Comment 9: Pg. 6769, Lines 16-22: This result is surprising since I might expect shallow boundary layers during winter to prevent PBL air from reaching the high altitude site. Also, please provide support for the claim that “the nearly hydrophobic mode observed in winter and autumn originates from the presence of combustion aerosols

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



emitted from heating devices".

Reply 9: During the winter time the boundary layer height can raise over the height of the puy de Dome station bringing within it large quantities of anthropogenic aerosols that have been concentrated in the shallow boundary layer. These events are isolated and are short-lived, but are responsible for the hydrophobic mode observed during the winter. This is now better explained in the text, and we bring supporting information on the presence of higher concentrations of organics and nitrates are measured during the winter months at the puy de Dome site (Freney et al., 2011) and also a higher fraction of biomass burning aerosol during the winter at the site (Crippa et al., 2013). We can't provide support for the statement that "the nearly hydrophobic mode observed in winter and autumn originates from the presence of combustion aerosols emitted from heating devices", so this sentence has been removed.

Comment 10: Pg. 6770, Lines 19-22: Is there any evidence to support the statement: "in autumn and winter, the high degree of external mixing may also be explained by more combustion aerosol being injected high in the atmosphere due to the very strong convection in the warm outflow".

Reply 10: We have only visual evidences of many wood fires during autumn and the beginning of winter. We will remove this statement.

Comment 11: Pg 6772, Lines 1-18: Is this discussion regarding seasonal differences in BC or in regional biomass burning supported by any measurements or observations from the site. It is mentioned on Pg. 6763, Line 10 that particulate BC, NOx, and CO2 are measured at the PdD during the study period; are these combustion tracer measurements consistent with the discussion in this section over the HTDMA measurement time period? What is the observational evidence for the statement: "In winter and in PBL conditions, the contribution from biomass burning is greater than in summer, bringing with it a high fraction of primary organics and nitrates"?

Reply 11: Higher concentrations of organics and nitrates are measured during the

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

winter months at the puy de Dome site (Freney et al., 2011) and also a higher fraction of biomass burning aerosol have been identified during the winter at the site (Crippa et al., 2013).

Comment 12: Pg. 6772, Line 25 and discussion throughout: How is the height of the PBL or RL being assessed so as to apportion specific aerosol types as deriving from the PBL or the FT? Presumably, this is coming from the vertical component of the HYSPLIT back trajectories or possible sonde-based temperature profiles?

Reply 12: PBL heights as a function of the time of the day as a function of the season have been calculated using model outputs (Venzac et al. 2009), and retrieved from LIDAR measurements (Freney et al. 2011; Rose et al. 2013) BLH retrievals were cross-checked using meteorological parameters (Boulon et al. 2011) and modelling outputs (Venzac et al. 2009). This is now mentioned in the method/site description section of the text.

Comment 13: Pg. 6773, Line 5-8: It should be easy to detect periods influenced by NPF by examining the variation in the overall particle number concentrations referenced on Pg. 6773, Line 10. It would be very interesting to know how frequently NPF events were observed during this study period in explaining the HTDMA results.

Reply 13: The frequency of NPF events has already been mentioned in the text, pg 6773, lines 11-15, that: “In winter, the day time increase in the number fraction of less hygroscopic particles (NF1) is smaller than during the other seasons. This is consistent with observations by Rose et al. (2013), who report that NPF events are less frequent in winter (NPF event frequency 17%) than in spring, summer and autumn (NPF event frequency 26%, 27% and 24%, respectively).”

Comment 14 Pg. 6774, Line 2-4: I don't understand the connection between lower wind speeds and a higher PBL.

Reply 14: We only point to the fact that within the PBL, wind speeds are lower than in

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

the FT. Hence, when the site lays within the PBL, air masses are expected to be more local than when it lays in the FT.

Comment 15: Pg. 6774, Lines 15-21: I'm surprised that the HTDMA measurements at 25-265 nm would be substantially influenced by dust, which tends to be present at larger particle sizes (>~0.5-5 micron diameters).

Reply 15: Sjögren et al. (2008) report changes in the GF-PDF mainly for particle size 250 nm, and Van Dingenen et al. (2005) observed a strong less hygroscopic mode during dust events for particle sizes 100 and 200 nm. We only cite the results. This has been corrected in the text.

Comment 16: Pg. 6775, Lines 6-8: What is the basis for the statement that anthropogenically-influenced marine aerosol is more "aged" than those originating from the continental PBL?

Reply 16: This sentence was unclear and has been removed.

Comment 17: Pg. 6775, Lines 21-25: I don't understand the formation mechanisms for "freshly formed anthropogenic aerosol" and "aged anthropogenic aerosol". What type of particles are these – organics and sulphate? "Aging" of fresh combustion particles is mentioned in this sentence as well.

Reply 17: By "freshly formed anthropogenic aerosol", we mean fresh anthropogenic emissions (BC and primary organics), and by "aged anthropogenic particles" we mean secondary organic and inorganic compounds. This has been clarified in the text.

Comment 18: Pg. 6776, Lines 10-12: The meaning of the sentence "Continental aerosol displays the same trend as smaller particles, with higher hygroscopicity in the cold season" is unclear.

Reply 18: The sentence has been clarified.

Comment 19: Section 3.4: I don't understand the reason for using the Zhou et al.,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

2001 parameterization since it is not physically based and because it obscures the inherent size-dependent measurement variability by reducing the data to two arbitrary coefficients. I recommend cutting this parameterization and only including the kappa values as is currently done while also adding their standard deviation. In addition, the hygroscopicity distribution concept of Su et al., 2010 seems perfect for describing this dataset, while preserving the spread of the GF distribution and the frequency of observation.

Reply 19: We believe that using the Zhou et al. (2001) parameterization is straightforward for users who are not necessarily familiar with the kappa theory. Users can directly grow their aerosols in a realistic way, according to air mass type and season. We believe that this parameterisation is useful (and harmless in the paper) and would like to keep it.

Comment 20: Pg. 6777, Lines 13-18: The statement, “The parameterization and the kappa values are issued from a long term data set, and are therefore reliable for future use in studies in which the hygroscopic properties should be taken into account, such as for calculations of condensational sink or for calculation of size distributions at ambient humidities for calculations of optical properties. The parameterizations and kappa values are representative of western European aerosol in remote sites”, is unfounded and should be removed.

Reply 20: The pdD site has been shown to be representative of western European air masses over a large scale (Henne et al. 2010) and we do not believe that this statement is unfounded.

Comment 21: Pg. 6778, Lines 10-11: What is the basis for this statement regarding ageing of fresh continental combustion aerosols? Is this based on elevated BC measurements?

Reply 21: We are clearly saying that we are speculating. We will remove this sentence.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Comment 22: Figure 1: How were the error bars for the MDF distribution calculated? Showing the actual data points should be a clearer way to show what the raw distribution looks like.

Reply 22: Error bars indicate the estimated counting efficiency of the measurements, and are described in detail by Gysel et al. (2009). The MDF in the figure is the measurement distribution function for one HTDMA scan, and actually shows the raw distribution (corrected to 90% RH), i.e. the actual data points, for that scan.

Comment 23: Figures 2 and 4: Given the overlapping uncertainty envelopes, it is hard to distinguish which colour corresponds to which curve (especially at 165 nm). For example, red and magenta are nearly indistinguishable, as are purple and blue. Recasting these figures using the hygroscopicity distribution concept might be a clearer representation; otherwise, consider spinning each curve out into a separate subfigure.

Reply 23: These figures have been improved, and are also now complemented with kappa values as a function of diameter, similar to those of Su et al. 2010.

Comment 24: The stacked bar graphs in Figure 5-7, 10-11 are very interesting and a nice contribution to better quantitative understanding of the aerosol mixing state.

Reply 24: Thank you.

Minor comments:

Comment 25: Pg. 6760, Line 4: Research was conducted under this period but the data set is not continuous. This should be noted here as it is on Pg. 6767, Lines 17-21.

Reply 25: This has been clarified in the text.

Comment 26: Pg. 6760, Lines 17-20: I don't think speculative conclusions are appropriate for an abstract.

Reply 26: Yes, the speculative phrasing has been removed.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Comment 27: Pg. 6761, Line 12: Update to the latest IPCC report.

Reply 27: Reference has been updated.

Comment 28: Pg. 6762, Lines 12-13: Boundary layer aerosols often feed convectively-driven tropospheric clouds.

Reply 28: Yes, this is now mentioned in the text.

Comment 29: Pg. 6762, Line 25: Again, please note that the data set is not continuous.

Reply 29: This has been clarified in the text.

Comment 30: Pg. 6763, Line 18: Remove citation to Hervo et al., 2014, which is in preparation. It is grossly misleading and inappropriate to cite a non-published paper with a publication year.

Reply 30: The citation has been removed.

Comment 31: Pg. 6765, Line 1: Was the RH constant at 90% or was it varied. Why is recalculation necessary?

Reply 31: The measured RH value varies over the course of the measurements. In general this variation is small, $\pm 3\%$, however in some cases the variation can be larger. The data analysis procedure filters out any data that was measured with a greater range of RH than 87 to 93% and corrects all the others to 90% so that the measured GF are comparable.

Comment 32: Pg. 6765, Lines 22-26: Is this referring to the GF and sigma for each mode or for the entire PDF?

Reply 32: This is referring to the entire PDF. A large sigma suggests that the PDF should be divided into different modes.

Comment 33: Pg. 6766, Line 5: Reference Figure 8 here.

Reply 33 Thanks, reference to Figure 8 is added.

C3906

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Comment 34: Pg. 6766, Line 5: What is the “oceanic modified” air mass type referring to? How is the air mass modified?

Reply 34: Oceanic modified refers to air masses originating from the ocean, but have travelled over the continent and are thus influenced by continental sources. This has been clarified in the text.

Comment 35: Pg. 6770, Line 5: What is “modified sea salt”?

Reply 35: Modified sea salt refers to sea salt that has been modified by other species in the air, for example condensation of secondary continental species. This has been clarified in the text.

Comment 36: Pg. 6772, Lines 23-24: Stating that the less hygroscopic fraction increases during the day at the expense of the hygroscopic fraction does not tell the reader whether there are more less hygroscopic particles present or just fewer hygroscopic particles present during the day.

Reply 36: The changes are, of course, for one population relative to the other. We have now stressed this in the text.

Comment 37: Pg. 6773, Line 19: What height was used to initialize the back trajectory?

Reply 37: The HYSPLIT air mass backward trajectories were calculated for the arrival pressure 850 hPa at the height of the puy de Dôme. This has been added to section 2.3.2 of the text.

Comment 38: Pg. 6775, Line 16: Does CNN refer to “cloud condensation nuclei (CCN)”?

Reply 38: Yes, thanks. This is corrected.

Comment 39: Pg. 6778, Line 27: I don’t think “realistic” is the correct word here, as ambient RH conditions can vary from dry air masses to moist air masses.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

Reply 39: Thanks, we have changed the word to “useful”.

Comment 40: Pg. 6780, Line 29: Strike the Hervo et al., in preparation reference since it is not yet accepted for publication.

Reply 40: The reference has been removed.

Comment 41: Table 1 could be moved to the supplementary material as it is not directly relevant to the discussion.

Reply 41: Yes, the table has been moved to the supplementary material.

Comment 42: In Tables 2-4 and Figures 1-7, 10-11, please explicitly note that the GF is GF(90%) in either the caption or the Table heading/Figure axis labels.

Reply 42: Thanks, this has been added to the captions.

References

Asmi, A., Wiedensohler, A., Laj, P., Fjaeraa, A.-M., Sellegrí, K., Birmili, W., Weingartner, E., Baltensperger, U., Zdimal, V., Zikova, N., Putaud, J.-P., Marinoni, A., Tunved, P., Hansson, H.-C., Fiebig, M., Kivekäs, N., Lihavainen, H., Asmi, E., Ulevicius, V., Aalto, P. P., Swietlicki, E., Kristensson, A., Mihalopoulos, N., Kalivitis, N., Kalapov, I., Kiss, G., de Leeuw, G., Henzing, B., Harrison, R. M., Beddows, D., O'Dowd, C., Jennings, S. G., Flentje, H., Weinhold, K., Meinhardt, F., Ries, L., and Kulmala, M.: Number size distributions and seasonality of submicron particles in Europe 2008– 2009, *Atmos. Chem. Phys.*, 11, 5505–5538, doi:10.5194/acp-11-5505-2011, 2011.

Crippa, M., Canonaco, F., Lanz, V. A., Äijälä, M., Allan, J. D., Carbone, S., Capes, G., Dall'Osto, M., Day, D. A., DeCarlo, P. F., Di Marci, C. F., Ehn, M., Eriksson, A., Freney, E., Hildenbrandt Ruiz, L., Hillamo, R., Jimenez, J.-L., Junnunen, H., Kiendler-Scharr, A., Kortelainen, A.-M., Kulmala, M., Mensah, A. A., Mohr, C., Nemitz, E., O'Dowd, C., Ovadnevaite, J., Pandis, S. N., Petäjä, T., Poulin, L., Saarikoski, S., Sellegrí, K., Swietlicki, E., Tiitta, P., Worsnop, D. R., Baltensperger, U., and Prévôt, A. S. H.: Organic

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



aerosol components derived from 25 AMS datasets across Europe using a newly developed ME-2 based source apportionment strategy, *Atmos. Chem. Phys. Discuss.*, 13, 23325–23371, doi: 10.5194/acpd-13-23325-2013, 2013.

Freney, E. J., Sellegrí, K., Canonaco, F., Boulon, J., Hervo, M., Weigel, R., Pichon, J. M., Colomb, A., Prévôt, A. S. H., and Laj, P.: Seasonal variations in aerosol particle composition at the puy-de-Dôme research station in France, *Atmos. Chem. Phys.*, 11, 13047–13059, doi:10.5194/acp-11-13047-2011, 2011.

Gysel, M., McFiggans, G. B., and Coe, H.: Inversion of tandem differential mobility analyser (TDMA) measurements, *J. Aerosol Sci.*, 40, 134–151, 2009.

Kammermann, L., Gysel, M., Weingartner, E., and Baltensperger, U.: 13-month climatology of the aerosol hygroscopicity at the free tropospheric site Jungfraujoch (3580 m a.s.l.), *Atmos. Chem. Phys.*, 10, 10717–10732, doi:10.5194/acp-10-10717-2010, 2010.

Putaud, J.-P., Raes, F., Van Dingenen, R., Brüggeman, E., Facchini, M. C., Decesari, S., Fuzzi, S., Gehrig, R., Hüglin, C., Laj, P., Lorbeer, G., Maenhaut, W., Mihalopoulos, N., Müller, K., Querol, X., Rodriguez, S., Schneider, J., Spindler, G., Ten Brink, H., Tørseth, K., and Wiedensohler, A.: A European Aerosol Phenomenology-2: chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2579–2595, 2004.

Rose, C., Boulon, J., Hervo, M., Holmgren, H., Asmi, E., Ramonet, M., Laj, P., and Sellegrí, K.: Long-term observations of positive cluster ion concentration, sources and sinks at the high altitude site of the Puy de Dôme, *Atmos. Chem. Phys. Discuss.*, 13, 14927–14975, doi:10.5194/acpd-13-14927-2013, 2013.

Sjögren, S., Gysel, M., Weingartner, E., Alfarra, M. R., Duplissy, J., Cozic, J., Crosier, J., Coe, H., and Baltensperger, U.: Hygroscopicity of the submicrometer aerosol at the highalpine site Jungfraujoch, 3580 m a.s.l., Switzerland, *Atmos. Chem. Phys.*, 8,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Su, H., Rose, D., Cheng, Y. F., Gunthe, S. S., Massling, A., Stock, M., Wiedensohler, A., Andreae, M. O., and Pöschl, U.: Hygroscopic distribution concept for measurement data analysis and modelling of aerosol particle mixing state with regard to hygroscopic growth and CCN activation, *Atmos. Chem. Phys.*, 10, 7489–7503, doi: 10.5194/acp-10-7489-2010, 2010. Van Dingenen, R., Putaud, J.-P., Martins-Dos Santos, S., and Raes, F.: Physical aerosol properties and their relation to air mass origin at Monte Cimone (Italy) during the first MINATROC campaign, *Atmos. Chem. Phys.*, 5, 2203–2226, doi:10.5194/acp-5-2203-2005, 2005.

Venzac, H., Sellegrí, K., Villani, P., Picard, D., and Laj, P.: Seasonal variation of aerosol size distributions in the free troposphere and residual layer at the puy de Dôme station, France, *Atmos. Chem. Phys.*, 9, 1465–1478, doi:10.5194/acp-9-1465-2009, 2009.

Zhou, J., Swietlicki, E., Berg, O. H., Aalto, P. P., Hameri, K., Nilsson, E. D., and Leck, C.: Hygroscopic properties of aerosol particles over the central Arctic Ocean during summer, *J. Geophys. Res.*, 106, 32111–32123, doi:10.1029/2000JD900426, 2001

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 14, 6759, 2014.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

