

Supplement of

**The effects of nitrate on the heterogeneous uptake of sulfur dioxide
on hematite**

L. D. Kong et al.

Correspondence to: L. D. Kong (ldkong@fudan.edu.cn)

1 XRD pattern of the synthesized sample




Fig. S1. XRD pattern of the synthesized sample.

The synthesized sample presented some diffraction peaks in the wide angle X-ray diffraction region. These characteristic peaks that emerged can be indexed according to the hematite crystalline phase, and no other crystalline phase was found in the peaks, indicating that the synthesized sample was a fairly pure α -Fe₂O₃ sample.

2 Schematic diagrams of DRIFTS and White Cell-FTIR apparatus




Fig. S2. Schematic diagram of DRIFTS apparatus




Fig. S3. Schematic diagram of White Cell-FTIR apparatus

3 The variations of the fitting peak areas with reaction time




Fig. S4. The variations of peak areas at 1261, 1222 and 1156 cm^{-1} with reaction time

Upon exposure of SO_2 , the surface-adsorbed H_2O was gradually consumed and the surface acidity of pure hematite sample was gradually enhanced. The transformation of the adsorption mode of the surface-formed sulfate would occur with the increase of the surface acidity, and then the peak at 1219 cm^{-1} appeared (Yamaguchi et al., 1986; Persson and Lovgren, 1996; Faguy and Marinković, 1996; Hug, 1997). The slight decrease in intensity of the peak at 1219 cm^{-1} after it reaches the maximum may be due to partial dissolution of hematite along with consumption of surface acidic species, which would lead to the formation of Fe^{3+} ions and some other surface species on the water-containing surface (Chun and Quon, 1973; Shi et al., 2011).

In addition, it should be pointed that peak fitting can produce bias, and hence the fitting band at 1222 cm^{-1} corresponds to the band at 1219 cm^{-1} in Fig. 1.

4 Peak heights of the main peaks observed on FN-24




Fig. S5. Peak heights of the main peaks in different infrared spectral regions.

5 Chromatogram and EI mass spectrum of N₂O




Fig. S6. Chromatogram (0.1-0.8 min) of the gas-phase sample collected from the heterogeneous reaction of SO₂ on FN-24 in a closed glass bottle in the dark (a) and corresponding EI mass spectrum of N₂O (b).

6 Assignments of the broad peak in the region between 1100 and 900 cm⁻¹ observed on pure hematite

A broad peak containing several weak peaks, especially at the beginning of the reaction, is also observed between 1100 and 900 cm⁻¹. Several peaks in this region can be assigned to the stretching motion of adsorbed sulfite and/or bisulfite based on previous studies on SO₂ adsorption on different metal oxide surfaces (Zhang et al., 2006; Wu et al., 2011; Nanayakkara, et al., 2012). These peaks can be overlapped by the symmetric stretching of sulfate in this region (Nanayakkara et al., 2012; Zhang et al., 2006), and it is difficult to distinguish these absorption peaks in this region because of the increase of surface sulfate species and the decrease of surface sulfite species during the reaction. These results indicate that surface S(IV) species such as adsorbed or surface-coordinated H₂SO₃, HSO₃⁻ and SO₃²⁻ species were converted to sulfate in the reaction process and some S(IV) species have not been completely oxidized to sulfate on the pure hematite surface after the experiment.

References

- Chun, K. C. and Quon, J. E.: Capacity of ferric oxide particles to oxidize sulfur dioxide in air, *Environ. Sci. Technol.*, 7, 532-538, 1973.
- Faguy, P. W. and Marinković, N. S.: An in situ infrared study on the effect of pH on anion adsorption at Pt (111) electrodes from acid sulfate solutions, *Langmuir*, 12, 243-247, 1996.

- Hug, S. J.: In situ Fourier transform infrared measurements of sulfate adsorption on hematite in aqueous solutions, *J. Colloid Interface Sci.*, 188, 415-422, 1997.
- Nanayakkara, C. E., Pettibone, J., and Grassian, V. H.: Sulfur dioxide adsorption and photooxidation on isotopically-labeled titanium dioxide nanoparticle surfaces: Roles of surface hydroxyl groups and adsorbed water in the formation and stability of adsorbed sulfite and sulfate, *Phys. Chem. Chem. Phys.*, 14, 6957-6966, 2012.
- Persson, P. and Lovgren, L.: Potentiometric and spectroscopic studies of sulfate complexation at the goethite-water interface, *Geochim. Cosmochim. Ac.*, 60, 2789-2799, 1996.
- Shi, Z., Bonneville, S., Krom, M. D., Carslaw, K. S., Jickells, T. D., Baker, A. R., and Benning, L. G.: Iron dissolution kinetics of mineral dust at low pH during simulated atmospheric processing, *Atmos. Chem. Phys.*, 11, 995-1007, 2011.
- Wu, L. Y., Tong, S. R., Wang, W. G., and Ge, M. F.: Effects of temperature on the heterogeneous oxidation of sulfur dioxide by ozone on calcium carbonate, *Atmos. Chem. Phys.*, 11, 6593–6605, 2011.
- Yamaguchi, T., Jin, T., and Tanabe, K.: Structure of acid sites on sulfur-promoted iron oxide, *J. Phys. Chem.*, 90, 3148-3152, 1986.
- Zhang, X., Zhang, G., Chen, J., Wang, Y., Wang, X., An, Z., and Zhang, P.: Heterogeneous reactions of sulfur dioxide on typical mineral particles, *J. Phys. Chem. B*, 110, 12588-12596, 2006.