



Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz^{1,*}, L. J. Carpenter², R.-J. Huang^{1,3,4}, M. Kundel¹, J. Bosle¹, and T. Hoffmann¹

¹Institute of Inorganic and Analytical Chemistry, Johannes Gutenberg-University of Mainz, 55128 Mainz, Germany

²Department of Chemistry, University of York, Heslington, York YO10 5DD, UK

³School of Physics & Centre for Climate and Air Pollution Studies, Ryan Institute, National University of Ireland Galway, University Road, Galway, Ireland

⁴Laboratory of Atmospheric Chemistry, Paul Scherrer Institute (PSI), 5232 Villigen, Switzerland

*now at: the Max Planck Institute for Chemistry, 55128 Mainz, Germany

Received: 15 April 2014 – Accepted: 12 May 2014 – Published: 5 June 2014

Correspondence to: T. Hoffmann (t.hoffmann@uni-mainz.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In this study we present the results of an incubation study of different phytoplankton samples in F/2 aqueous media treated with elevated ozone levels. Halocarbon measurements show that the samples tested released bromoform and different iodocarbons including iodomethane, iodochloromethane and diiodomethane. Iodide and iodate levels in the liquid phase were representative of concentrations of surface water in a natural environment. Measurement of volatile iodine (I_2) emissions from two diatom samples (*Mediopyxis helysia* and *Porosira glacialis*) and the background sample (F/2-medium from locally seawater), showed that the quantity of I_2 evolved depends on the ozone concentration in the air. This behaviour was assumed to be caused by the oxidation reaction mechanism of iodine with ozone. The I_2 emission flux agrees with model calculations at different iodide concentrations. The I_2 emission of a natural plankton concentrate sample was, however, very low compared to other samples and showed no dependence on ozone. The reason for this was shown to be the low iodide concentration in the algae suspension, which seems to be the limiting factor in the oxidative formation of I_2 .

1 Introduction

Iodine chemistry plays an essential role in the marine boundary layer (MBL) due to its effect on the destruction of tropospheric ozone, perturbation of the HO_x/NO_x cycle and the formation of new particles and cloud condensation nuclei, thereby leading to changes in the global radiative forcing (Hoffmann et al., 2001; von Glasow and Crutzen, 2003; O'Dowd and Hoffmann, 2005; Bloss et al., 2005; Huang et al., 2010a, b). This essential role of iodine and of other activated halogens is shown in field measurements in the marine boundary layer (MBL), laboratory chamber experiments or incubation experiments of different algae and in atmospheric models (Carpenter, 2003; Küpper et al., 2008; Kundel et al., 2012; McFiggans et al., 2000). The biogeochemical cycle

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of iodine is controlled by large iodine exchanges from the oceans to the atmosphere, driven by marine biotic and abiotic production (Schall et al., 1997). Volatilized species are photolabile iodocarbons like CH_2I_2 , CH_3I , $\text{C}_2\text{H}_5\text{I}$, CH_2ICl , CH_2IBr and molecular iodine (I_2). Marine species like macroalgae and microalgae play a dominant role in the emission of these compounds (Carpenter et al., 1999; Huang et al., 2013; Saiz-Lopez and Plane, 2004).

Since molecular iodine and iodocarbons are photochemically instable (lifetimes between about some tens of seconds for I_2 and a few days for CH_3I) they are photolysed under UV-visible light to form $\text{I}\cdot$ atoms which are then instantly oxidised by ozone to form the iodine monoxide radical IO (g) (Hoffmann et al., 2001; Saiz-Lopez et al., 2006). Further oxidation reactions of IO in the gas phase then can form low volatile iodine oxides (I_xO_y) which may nucleate under certain conditions and form new particles.

Recently it was proposed that the ozone loss over the tropical Atlantic Ocean was higher than calculated from global atmospheric models, and that this additional ozone destruction is induced by halogens such as bromine and iodine (Read et al., 2008). Biogenic emissions, such as the already studied iodocarbon emissions by phytoplankton species, e.g. coccolithophorids, diatoms and chlorophytes, (Colomb et al., 2008) are too low to explain the differences in model calculations and observations (Mahajan et al., 2010), therefore additional sources of the reactive iodine species are discussed, one of them being the surface reaction of ozone with seawater.

Garland and Curtis first discovered that the emission of molecular iodine from the surface of artificial and natural seawater is proportional to the ozone concentration at the air/water interface (Garland and Curtis, 1981). Sakamoto and co-workers examined the reaction mechanism of the iodide oxidation by ozone at the air/water interface, resulting in the formation of the intermediates IOOO^- and HOI and the emission products IO and I_2 (Sakamoto et al., 2009). Further laboratory experiments show that different organics affect the reaction of iodide with ozone, e.g. fulvic acid enhances the I_2 formation, but not the formation of IO (Hayase et al., 2010, 2012).

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Since the formation of I_2 and IO from the air/water interface is dependent on the iodide concentration in seawater, the reaction path found by Garland and Curtis may explain elevated iodine emissions in areas of higher phytoplankton activity (Garland and Curtis, 1981). The ability of different phytoplankton, e.g diatoms, to reduce iodate, which is ubiquitous in the open ocean, to iodide was shown for natural and elevated iodate concentrations (Wong et al., 2002; Chance et al., 2007) and for the different growth states (Bluhm et al., 2010) of the phytoplankton cultures. A correlation of iodine species in the particle phase and average chlorophyll exposure of air masses along back trajectories was found by Lai et al. (2011), indicating the link between phytoplankton activity and emission of atmospheric iodine.

Since the formation of I_2 and IO is correlated to the iodide concentration (Sakamoto et al., 2009) and the iodide concentration of surface waters is correlated to phytoplankton (Bluhm et al., 2010), this study investigates links between phytoplankton, iodide concentrations in seawater and I_2 emission, utilising laboratory experiments of the reaction of the seawater surface with ozone.

2 Materials and methods

2.1 Experimental set-up



Two diatom cultures (*M. helysia*, *Porosyra glacialis*) were kept in F/2 seawater medium (Guillard, 1975). Both cultures were kept for at least 4 weeks prior the experiment and were then diluted in a 2 : 1 ratio in F/2 medium and homogenised by stirring. In addition to the diatom cultures, a plankton concentrate was collected from the North Sea (55°01.562' N; 8°27.113' E) on 24 May 2012 using a 80 μ m and 200 μ m Apstein plankton net and diluted using F/2 medium, the sample was dominated by colonies of the haptophyte *Phaeocystis sp.* and only a low amount of diatoms was present in the sample, as determined by a microscopic analyses.

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For each experiment, 1.5 L of the sample was introduced into a glass chamber tube (10 L), shown in Fig. 1, and three magnetic stirrers were switched on immediately. A continuous flow of synthetic air (3.4 L min^{-1}) was channelled over the stirred algae suspension in the first experiment with no ozone and in the second experiment with elevated ozone levels of 100 ppb. The ozone was generated using an UV radiation source and the resulting ozone levels were measured using an ozone analyzer (Dasibi Environmental Corp. Model 1008-RS, Glendale, USA). To measure the emission of I_2 and halocarbons, α -cyclodextrin-coated denuders (Huang and Hoffmann, 2009; Huang et al., 2010c) and adsorption tubes (Kundel et al., 2012) were mounted at the other end of the tube chamber together with the ozone monitor. The chamber outflow was sampled using two membrane pumps, one with 0.50 L min^{-1} for the denuders and the other using 0.15 L min^{-1} for the adsorption tubes. To assure an overpressure over the sampling time a U-shaped tube filled with ultra-pure water was mounted in the centre exit of the glass chamber to measure the overpressure hydrostatically. The whole set-up was wrapped with aluminium foil to prevent photolysis of I_2 and halocarbon compounds.

To monitor the emissions of I_2 and halocarbons from the liquid samples, an evaporation standard was added to the microalgae suspension in order to highlight any problems related to air sampling. This standard was 1,3-dibromopropane diluted in ultrapure water ($500 \mu\text{L}$ of $0.94 \mu\text{g L}^{-1}$ which was then diluted with the sample to 1.5 L). The standard was chosen given the results from a first set of experiments with *M. helysia* and *Coscinodiscus wailesii* which show no detectable traces of this compound. We decided not to add any iodine containing compounds to prevent interferences with the I_2 emission.

2.2 Halocarbon measurements

Air samples of 6.75 L sampling volume were preconcentrated at a flow rate of 150 mL min^{-1} on thermal desorption tubes filled with 100 mg Tenax TA 60/80 and 150 mg CarbotrapTM 20/40 both provided by Supelco (Bellefonte, PA, USA). The samples were analysed using a self-made thermal desorption device mounted on a gas

After adding 3 mL sodium acetate the sample was extracted with 100 μ L cyclohexane and 100 μ L 2,4,6-tribromoanilin (internal standard: IS) in cyclohexane (250 ppb).

Iodide and iodate were derivatized from seawater to form the same product as described for I_2 . Iodide was oxidized to form I_2 by using iodosobenzoate and iodate was reduced first to iodide and then oxidized to form I_2 . 10 mL aliquots of seawater were analysed for iodide and for total iodine, iodate was calculated by difference. The method for iodide derivatization was slightly changed from the one described by Mishra et al. (2000). The use of sodium hydrogen sulfite as an agent to reduce iodate to iodide is described by Schwehr and Santschi (2003).

$$I_{\text{seawater}} = I^- + IO_3^-$$

To measure iodide, 10 mL seawater were mixed with 1 mL ethylenediaminetetraacetic acid solution (0.5 %), 500 μ L phosphate buffer, 500 μ L N,N-dimethylaniline, 500 μ L iodosobenzoate and shaken. After adding 3 mL sodium acetate the sample was extracted with 100 μ L cyclohexane and 100 μ L 2,4,6-tribromoanilin (IS) in cyclohexane (250 ppb).

To measure iodate an aliquot of 10 mL seawater was mixed with 1 mL ethylenediaminetetraacetic acid solution (0.5 %), 1 mL hydrochloric acid (3.7 %) and 500 μ L sodium hydrogen sulfite solution (283.9 $\mu\text{mol L}^{-1}$) to reduce the iodate. Afterwards 500 μ L sodium acetate, 4 mL phosphate buffer, 500 μ L N,N-dimethylaniline, 500 μ L iodosobenzoate were added. After shaking the sample was again extracted with 100 μ L cyclohexane and 100 μ L 2,4,6-tribromoanilin (IS) in cyclohexane (250 ppb).

1 μ L of the cyclohexane extract was injected to the GC-MS System (6850 GC & 5973 MS, Agilent Technologies, Waldbronn, Germany) at a constant flow of 1 mL min^{-1} of helium (99.999 %), the chromatographic separation was performed using a capillary column FS Supreme 5 MS with length of 30 m, inner diameter of 0.25 mm and film thickness of 0.25 μm (CS Chromatographie Servieve, Langenwehe, Germany) with a temperature program starting at 50 $^\circ\text{C}$ (for 3 min), then heating up at 30 $^\circ\text{C min}^{-1}$ to 220 $^\circ\text{C}$ (for 3 min). The mass spectrometer measured in electron ionisation mode at 70 eV, the

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion standard was added to the different samples to recognize differences in emission rates of the organic compounds from the aqueous phase. The standard was added in a 10 to 100-fold excess compared to natural concentrations of bromocarbons in Atlantic seawater (Carpenter et al., 2000) to reduce the effect of natural 1,3-dibromopropane which may alter the mixing ratio of the evaporation standard measured. In the chosen concentration a natural abundance would change the result only by 1–10 % compared to the spike solution. The results of the measurements of 1,3-dibromopropane showed very constant values, as can be seen from the low standard deviation between the different samples and replicates. This result indicates a stable and reliable experimental setup in terms of evaporation of volatile compounds from the water surface and of the mixing of the bulk water.

The measured emission rates of the natural halocarbons show that the brominated compound, CHBr_3 , is elevated compared to the iodocarbons emission rates. This result actually fits to observations of the natural abundance of halocarbons in seawater as described in earlier studies (Roy et al., 2011). The emission rate of CHBr_3 is higher for the two diatom cultures (*M. helysia* and *P. glacialis*) than for the plankton samples containing *Phaeocystis sp.* and the background F/2 medium. Again, this result matches field and laboratory data showing a link between elevated CHBr_3 concentrations in seawater and the simultaneous occurrence of diatoms (Colomb et al., 2008; Quack et al., 2007; Moore et al., 1996).

The organoiodine compounds have two different emission patterns regarding the relative abundance of the different iodine containing volatiles: the first one is reflected by the F/2 media background and the plankton concentrate and the other by the two diatom cultures. The abundance of iodocarbons in **experiment** using the F/2 media background was dominated by CH_3I , followed by CH_2ICl and CH_2I_2 . Whereas for the diatom cultures, CH_2I_2 was the dominant iodocarbon emitted and CH_3I and CH_2ICl both showing lower emission rates. Additionally, the iodocarbon emission rates in the diatom samples were elevated compared to the F/2 background, showing that diatoms are capable of producing iodocarbons (Moore et al., 1996). To compare the natural

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



plankton concentrate with the cultivated diatom cultures one must keep in mind that chl- α concentrations are biomass tracers reflecting the abundance of phytoplankton. The results for the chl- α measurement, given in Table 1, clearly show that the natural plankton concentrate contains less biomass than the cultured diatoms. Therefore, we conclude that the lower iodocarbon emissions of the plankton concentrate compared to the diatom cultures is partly due to lower biomass density.

The emission flux summed for the three iodocarbons in the four samples F/2-medium, plankton concentrate, *P. glacialis* and *M. helysia* was in the range of 0.034–0.163 ng min⁻¹ m⁻², 0.025–0.098 ng min⁻¹ m⁻², 0.106–0.264 ng min⁻¹ m⁻² and 0.153–0.288 ng min⁻¹ m⁻², respectively. We are not aware of incubation studies investigating the flux of iodocarbons from micro algae suspensions to directly compare these results. To establish a connection to other experimental observations the results listed above are compared to incubation studies of marine aggregates producing iodocarbons and calculated emission fluxes for open sea water. Hughes et al. (2008) measured the iodocarbon production of different marine aggregates to be within 6 to 66 ng min⁻¹ L⁻¹. The production rate is difficult to compare to the presented results, since the flux in our study is based on the production by the microalgae species and evaporation from the surface, whereas Hughes et al. (2008) measured the production in the aqueous phase. Jones et al. (2009) calculated fluxes from seawater and gas phase concentrations. The sampling site was surrounded by fields of macro algae in open sea water at Roscoff, France. The flux of iodocarbons was estimated to 11.281 ng min⁻¹ m⁻², actually two orders of magnitude higher than the flux obtained in the present study. This may be explained by the higher iodocarbon emissions from macro algae exposed to intertidal dryness (Carpenter et al., 1999), compared to micro algae suspensions.

3.2 Iodide and iodate

The concentration of iodide and iodate is also shown in Table 1, again the concentrations were not dependent on the ozone concentration and therefore results are summarized for the six replicates of each sample. The iodate concentrations in the

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



background F/2 medium and in the three plankton samples were in the same range, with mean concentrations between 438 and 448 nmol L⁻¹. These iodate concentrations are in the range measured for the open ocean of 400 to 500 nmol L⁻¹ iodate in most oceanic regions (Bluhm et al., 2011). The ubiquitous abundance of iodate suggests that its concentration is not a limiting factor.

The iodide concentrations in the two diatom cultures, *P. glacialis* and *M. helysia*, are slightly elevated with mean values of 12.70 nmol L⁻¹ and 16.84 nmol L⁻¹, respectively, compared to the F/2 medium iodide concentration of 10.35 nmol L⁻¹ and the plankton concentrate iodide concentration of 6.47 nmol L⁻¹. This enhanced iodide concentration indicates the reduction of iodate by the two diatom cultures, which was also found by Bluhm et al. (2010) and Wong et al. (2002) for different phytoplankton species. The iodide concentrations in all samples are comparable with oceanic surface water concentrations, for example around 10–30 nmol L⁻¹ in the Weddel Sea surface water (Bluhm et al., 2011).

The low iodide concentration of the plankton concentrate sample compared to the F/2 background sample is surprising, but may be assigned to an overall low level of different nutrients, like phosphate and silicate, in the Wadden Sea of Sylt at springtime (Weisse et al., 1986), although the level of iodate was consistent.

3.3 Ozone measurements

The results of the ozone measurement for the samples: F/2 medium, *P. glacialis*, *M. helysia* and the plankton concentrate were normalized against a background measurement obtained using ultra-pure water in the chamber. This was performed in order to account for losses of ozone through wall reactions, losses on the water surface, and losses due to droplet formation from stirring. The ozone consumption was calculated using a Continuous Stirred-Tank Reactor (CSTR) approach with 668 ng min⁻¹ ozone (100 ppb) introduced into the chamber (total volume: 10 L, flow: 3.4 L min⁻¹ and residence time: 2.94 min). The difference between the introduced ozone flow and measured ozone flow is considered as consumed ozone, due to the oxidation of iodide

and other ozone depleting reactions in the samples. To calculate the consumed ozone, the flow rate was summarized over 45 min of the experiment. Ozone consumption was clearly observed for all samples. The F/2 background sample showed the weakest ozone consumption of 58 nmol, followed by the sample of *P. glacialis* with 186 nmol and the plankton concentrate with 253 nmol. The highest ozone consumption was shown by *M. helysia* with 335 nmol.

3.4 I₂ emissions

The I₂ emission rate was calculated by dividing the amount of I₂ by the sampling time and the suspension surface area. The results for the four samples are shown in Fig. 2. The F/2 medium and the two diatom samples, *M. helysia* and *P. glacialis* show significant higher emission rates when the ozone level is elevated (100 ppb O₃) compared to conditions where no ozone is present (0 ppb O₃). The difference between the high and low ozone conditions is small for the F/2 medium, increases for the *P. glacialis* sample and is highest for the *M. helysia* sample. The plankton concentrate does not show a significant dependence of the I₂ emission rate on the ozone level. The ozone-dependent increase in the I₂ emission rate of the other samples indicates that iodide, which is present at the air/water interface, is oxidised by ozone to form I₂, which is consistent with the results from artificial and natural seawater (Garland and Curtis, 1981; Sakamoto et al., 2009).

Figure 3 shows the change in I₂ emission rate ($[I_2 \text{ at } 100 \text{ ppb ozone}] - [I_2 \text{ at } 0 \text{ ppb ozone}]$) of the different samples as a function of the iodide concentration measured in the bulk water. A linear correlation fits the data well with a Pearson coefficient of $R^2 = 0.998$. This behaviour indicates a direct proportional relationship, which was also seen by Sakamoto et al. (2009) for small iodide concentrations (0–5 mmol L⁻¹). Carpenter et al. (2013) also observed that the I₂ emission is dependent on the aqueous iodide concentration. The proposed reaction sequence as shown in Reactions (R1)–(R5) explain the relationship between the iodide concentration in the aqueous phase

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and the I₂ emission (Sakamoto et al., 2009).



The plankton sample does not show an elevated I₂ emission at 100 ppb ozone compared to zero ozone. This observation indicates that in the plankton sample an additional I₂ loss process takes place. Reactions or partitioning of I₂ in an organic surface layer, which was discussed in Carpenter et al. (2013), would be one possibility to explain these results. In fact the specific microalgae found in the plankton concentrate, *Phaeocystis sp.*, is known to produce high amounts of organic matter (Eberlein et al., 1985). An alternative explanation is the low iodide concentration in the plankton concentrate. The iodide concentrations and ozone mixing ratios in this study represent more likely natural conditions compared to the study of Sakamoto et al. (iodide concentration between 0.01–50 mmol L⁻¹ and ozone mixing ratio from 2–298 ppm). However, the results presented here show that also at natural conditions a significant formation of I₂ by the ozone driven oxidation of iodide at the air/water interface takes place, until the iodide concentration gets too low.

Calculated emissions for the F/2 medium, *P. glacialis* and *M. helysia* were 8.32×10^5 , 1.47×10^6 and 2.40×10^6 molecules cm⁻² s⁻¹, respectively. Modelled emissions calculated using the kinetic model of the aqueous interfacial layer by Carpenter et al. (2013) for the iodide concentration measured were 1.16×10^6 , 1.67×10^6 and 2.91×10^6 molecules cm⁻² s⁻¹, respectively. The measured and modelled values agree well, showing that the model is able to predict emissions for natural iodide concentrations well.

Figure 4 shows the change in the I₂ emission rate plotted vs. the consumed ozone for the four different samples. This was done to see whether ozone depletion in the

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flow chamber is mainly driven by the iodide or if other factors are important. The graph shows that the ozone depletion correlates with the enhancement in the I_2 emission rate for the two diatom samples and for the F/2 medium. Therefore the formation ratio of I_2 in dependence on O_3 was calculated by $R(I_2) = n(I_2)/n(O_3)$, with $n(I_2) =$ amount of I_2 formed and $n(O_3) =$ amount of O_3 consumed during the experiment. $R(I_2)$ has a maximum value of 1, regarding to Reactions (R1)–(R5) if every molecule ozone which is consumed produces one molecule of I_2 . The formation ratio for the F/2 background sample was the highest with $R(I_2) = 0.14\text{‰}$, followed by the samples of *M. helysia* with $R(I_2) = 0.08\text{‰}$ and *P. glacialis* $R(I_2) = 0.07\text{‰}$. This means that a higher degree of biologic activity of the sample decreases the formation ratio. The decrease of I_2 emission in the surface reaction of ozone with iodide was also seen by Carpenter et al. when turning from iodide solutions to sea water, which contains more organic substances (Carpenter et al., 2013).

The plankton concentrate also depletes ozone, although there is no enhancement in I_2 emission. Therefore another mechanism in ozone depletion obviously takes place, possibly induced by other ozone reactive substances formed or excreted from *Phaeocystis sp.* Another explanation is a reduced release of I_2 and a higher release of HOI, which was not measured in this study. Indeed Carpenter and coworkers observed HOI as the main iodine compound released in their experiments, followed by I_2 (Carpenter et al., 2013).

4 Conclusions

Different phytoplankton suspensions were treated with high and low ozone levels. Halocarbons including bromoform, iodomethane, iodochloromethane and diiodomethane, were released from the suspensions independent of the ozone level. The use of an evaporation standard in the aqueous phase indicated that the emission rates of all gaseous organics were quite stable. The iodide and iodate concentration in the liquid phase also showed no dependence on the ozone level in the gas phase. The emis-

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sion flux of the iodocarbons was lower compared to the calculated flux at a natural site in Roscoff, France, an observation which emphasizes the higher emission of iodocarbons from macroalgae compared to microalgae. The iodide and iodate concentration of the investigated samples was comparable to surface water in the open ocean. The emission of I_2 showed a dependency on the ozone level in the air as well as from the iodide concentration in the sample suspension, as has been found previously (Carpenter et al., 2013 and other refs). For the two diatom samples *M. helysia* and *P. glacialis* and the F/2 medium background sample, a correlation was found for the I_2 emission and the ozone consumption during the experiment. The I_2 emissions from the plankton concentrate, taken in the Wadden Sea of Sylt, was lower than the other samples and showed no dependence on the ozone levels. An explanation could be the lower iodide concentration in the plankton sample, since iodide is the limiting factor for the oxidative reaction. Another explanation may be the preferred formation and emission of HOI when organic compounds are present in the liquid phase. The experiments showed that different algae (*M. helysia* and *P. glacialis*) suspensions are capable to emit I_2 by the reaction of ozone with dissolved iodide at the air/water interface. However, it remains unclear whether without the simultaneous measurement of HOI the iodine emission from aquatic systems can be fully understood.

Acknowledgements. The authors gratefully acknowledge the financial support of the German Research Foundation (DFG: Deutsche Forschungsgemeinschaft) within the Research Training Group 826: “Trace Analysis of Elemental Species: Development of Methods and Applications.”

References

Bloss, W. J., Lee, J. D., Johnson, G. P., Sommariva, R., Heard, D. E., Saiz-Lopez, A., Plane, J. M., McFiggans, G., Coe, H., Flynn, M., Williams, P., Rickard, A. R., and Fleming, Z. L.: Impact of halogen monoxide chemistry upon boundary layer OH and HO₂ concentrations at a coastal site, *Geophys. Res. Lett.*, 32, L06814, doi:10.1029/2004GL022084, 2005.

**Emission of iodine
containing volatiles
by selected
microalgal species**

U. R. Thorenz et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bluhm, K., Croot, P., Wuttig, K., and Lochte, K.: Transformation of iodate to iodide in marine phytoplankton driven by cell senescence, *Aquat. Biol.*, 11, 1–15, 2010.
- Bluhm, K., Croot, P. L., and Huhn, O. R. G.: Distribution of iodide and iodate in the Atlantic sector of the southern ocean during austral summer, *Deep-Sea Res. Pt. II*, 58, 2733–2748, 2011.
- Carpenter, L. J.: Iodine in the marine boundary layer, *Chem. Rev.*, 103, 4953–4962, 2003.
- Carpenter, L. J., Sturges, W. T., Penkett, S. A., Liss, P. S., Alicke, B., Hebestreit, K., and Platt, U.: Short-lived alkyl iodides and bromides at Mace Head, Ireland: links to biogenic sources and halogen oxide production, *J. Geophys. Res.-Atmos.*, 104, 1679–1689, 1999.
- Carpenter, L. J., Malin, G., Liss, P. S., and Kupper, F. C.: Novel biogenic iodine-containing trihalomethanes and other short-lived halocarbons in the coastal East Atlantic, *Global Biogeochem. Cy.*, 14, 1191–1204, 2000.
- Carpenter, L. J., MacDonald, S. M., Shaw, M. D., Kumar, R., Saunders, R. W., Parthipan, R., Wilson, J., and Plane, J. M. C.: Atmospheric iodine levels influenced by sea surface emissions of inorganic iodine, *Nat. Geosci.*, 6, 108–111, 2013.
- Chance, R., Malin, G., Jickells, T., and Baker, A. R.: Reduction of iodate to iodide by cold water diatom cultures, *Mar. Chem.*, 105, 169–180, 2007.
- Colomb, A., Yassaa, N., Williams, J., Peeken, I., and Lochte, K.: Screening volatile organic compounds (VOCs) emissions from five marine phytoplankton species by head space gas chromatography/mass spectrometry (HS-GC/MS), *J. Environ. Monit.*, 10, 325–330, 2008.
- Eberlein, K., Leal, M. T., Hammer, K. D., Hickel, W.: Dissolved organic substances during a *Phaeocystis pouchetii* bloom in the German Bight (North Sea), *Mar. Biol.*, 89, 311–316, 1985.
- Edlar, L.: *Phytoplankton and Chlorophyll*. Baltic Marine Biologists, vol 5, Univ. of Stockholm, Stockholm, 1979.
- Garland, J. A. and Curtis, H.: Emission of iodine from the sea-surface in the presence of ozone, *J. Geophys. Res.-Oc. Atm.*, 86, 3183–3186, 1981.
- Hayase, S., Yabushita, A., Kawasaki, M., Enami, S., Hoffmann, M. R., and Colussi, A. J.: Heterogeneous reaction of gaseous ozone with aqueous iodide in the presence of aqueous organic species, *J. Phys. Chem. A*, 114, 6016–6021, 2010.
- Hayase, S., Yabushita, A., and Kawasaki, M.: Iodine emission in the presence of humic substances at the water's surface, *J. Phys. Chem.*, 116, 5779–5783, 2012.

**Emission of iodine
containing volatiles
by selected
microalgae species**

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hoffmann, T., O'Dowd, C. D., and Seinfeld, J. H.: Iodine oxide homogeneous nucleation: an explanation for coastal new particle production, *Geophys. Res. Lett.*, 28, 1949–1952, 2001.
- Huang, R. J. and Hoffmann, T.: Development of a coupled diffusion denuder system combined with gas chromatography/mass spectrometry for the separation and quantification of molecular iodine and the activated iodine compounds iodine monochloride and hypoiodous acid in the marine atmosphere, *Anal. Chem.*, 81, 1777–1778, 2009.
- Huang, R.-J., Seitz, K., Buxmann, J., Pöhler, D., Hornsby, K. E., Carpenter, L. J., Platt, U., and Hoffmann, T.: In situ measurements of molecular iodine in the marine boundary layer: the link to macroalgae and the implications for O₃, IO, OIO and NO_x, *Atmos. Chem. Phys.*, 10, 4823–4833, doi:10.5194/acp-10-4823-2010, 2010a.
- Huang, R. J., Seitz, K., Neary, T., O'Dowd, C. D., Platt, U., and Hoffmann, T.: Observations of high concentrations of I₂ and IO in coastal air supporting iodine-oxide driven coastal new particle formation, *Geophys. Res. Lett.*, 37, L03803, doi:10.1029/2009GL041467, 2010b.
- Huang, R. J., Hou, X. L., and Hoffmann, T.: Extensive evaluation of a diffusion denuder for the quantification of atmospheric stable and radioactive molecular iodine, *Environ. Sci. Technol.*, 44, 5061–5066, 2010c.
- Huang, R.-J., Thorenz, U. R., Kundel, M., Venables, D. S., Ceburnis, D., Ho, K. F., Chen, J., Vogel, A. L., Küpper, F. C., Smyth, P. P. A., Nitschke, U., Stengel, D. B., Berresheim, H., O'Dowd, C. D., and Hoffmann, T.: The seaweeds *Fucus vesiculosus* and *Ascophyllum nodosum* are significant contributors to coastal iodine emissions, *Atmos. Chem. Phys.*, 13, 5255–5264, doi:10.5194/acp-13-5255-2013, 2013.
- Hughes, C., Malin, G., Turley, C., Keely, B., Nightingale, P., and Liss, P.: The production of volatile iodocarbons by biogenic marine aggregates. *Limnol. Oceanogr.*, 53, 867–872, 2008.
- Jeffrey, S. W. and Humphrey, G. F.: New spectrophotometric equations for determining chlorophylls A, B, C1 and C2 in higher plants, algae and natural phytoplankton, *Biochem. Physiol. Pfl.*, 167, 191–194, 1975.
- Jones, C. E., Hornsby, K. E., Dunk, R. M., Leigh, R. J., and Carpenter, L. J.: Coastal measurements of short-lived reactive iodocarbons and bromocarbons at Roscoff, Brittany during the RHaMBLe campaign, *Atmos. Chem. Phys.*, 9, 8757–8769, doi:10.5194/acp-9-8757-2009, 2009.
- Kundel, M., Thorenz, U. R., Petersen, J. H., Huang, R., Bings, N. H., and Hoffmann, T.: Application of mass spectrometric techniques for the trace analysis of short-lived iodine-containing volatiles emitted by seaweed, *Anal. Bioanal. Chem.*, 402, 3345–3357, 2012.

**Emission of iodine
containing volatiles
by selected
microalgal species**

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Küpper, F. C., Carpenter, L. J., McFiggans, G. B., Palmer, C. J., Waite, T. J., Boneberg, E.,
Woitsch, S., Weiller, M., Abela, R., Grolimund, D., Potin, P., Butler, A., Luther, G. W., Kro-
neck, P. M. H., Meyer-Klaucke, W., and Feiters, M. C.: Iodide accumulation provides help
with an inorganic antioxidant impacting atmospheric chemistry, *P. Natl. Acad. Sci. USA*, 105,
6954–6958, 2008.

Lai, S. C., Williams, J., Arnold, S. R., Atlas, E. L., Gebhardt, S., and Hoffmann, T.: Iodine con-
taining species in the remote marine boundary layer: a link to oceanic phytoplankton, *Geo-
phys Res. Lett.*, 38, L20801, doi:10.1029/2011GL049035, 2011.

Mahajan, A. S., Plane, J. M. C., Oetjen, H., Mendes, L., Saunders, R. W., Saiz-Lopez, A.,
Jones, C. E., Carpenter, L. J., and McFiggans, G. B.: Measurement and modelling of tropo-
spheric reactive halogen species over the tropical Atlantic Ocean, *Atmos. Chem. Phys.*, 10,
4611–4624, doi:10.5194/acp-10-4611-2010, 2010.

McFiggans, G., Plane, J. M., Allan, B. J., Carpenter, L. J., Coe, H., and O'Dowd, C.: A model-
ing study of iodine chemistry in the marine boundary layer, *J. Geophys. Res.-Atmos.*, 105,
14371–14385, 2000.

Mishra, S., Singh, V., Jain, A., and Verma, K. K.: Determination of iodide by derivatization to
4-iodo-N,N-dimethylaniline and gas chromatography-mass spectrometry, *Analyst*, 125, 459–
464, 2000.

Moore, R. M., Webb, M., Tokarczyk, R., and Wever, R.: Bromoperoxidase and iodoperoxidase
enzymes and production of halogenated methanes in marine diatom cultures. *J. Geophys.
Res.-Oceans*, 101, 20899–20908, 1996.

O'Dowd, C. D. and Hoffmann, T.: Coastal new particle formation: a review of the current state-
of-the-art, *Environ. Chem.*, 2, 245–255, 2005.

Quack, B., Peeken, I., Petrick, G., and Nachtigall, K.: Oceanic distribution and sources of bro-
moform and dibromomethane in the Mauritanian upwelling, *J. Geophys. Res.-Oceans*, 112,
C10006, doi:10.1029/2006JC003803, 2007.

von Glasow, R. and Crutzen, P. J.: *Tropospheric Halogen Chemistry*, edited by: Turekian, K. K.
and Holland, H. D., Volume 4, *Treatise on Geochemistry*, 21–64, 2003.

Read, K. A., Mahajan, A. S., Carpenter, L. J., Evans, M. J., Faria, B. V. E., Heard, D. E., Hop-
kins, J. R., Lee, J. D., Moller, S. J., Lewis, A. C., Mendes, L., McQuaid, J. B., Oetjen, H.,
Saiz-Lopez, A., Pilling, M. J., and Plane, J. M. C.: Extensive halogen-mediated ozone de-
struction over the tropical Atlantic Ocean, *Nature*, 453, 1232–1235, 2008.

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Roy, R.: Short-term variability in halocarbons in relation to phytoplankton pigments in coastal waters of the central eastern Arabian Sea, *Estuar Coast. Shelf. S.*, 88, 311–321, 2010.
- Roy, R., Pratihary, A., Narvenkar, G., Mochemadkar, S., Gauns, M., and Naqvi, S. W. A.: The relationship between volatile halocarbons and phytoplankton pigments during a *Trichodesmium* bloom in the coastal eastern Arabian Sea, *Estuar Coast. Shelf. S.*, 95, 110–118, 2011.
- 5 Saiz-Lopez, A. and Plane, J. M.: Novel iodine chemistry in the marine boundary layer, *Geophys. Res. Lett.*, 31, L04112, doi:10.1029/2003GL019215, 2004.
- Saiz-Lopez, A., Plane, J. M. C., McFiggans, G., Williams, P. I., Ball, S. M., Bitter, M., Jones, R. L., Hongwei, C., and Hoffmann, T.: Modelling molecular iodine emissions in a coastal marine environment: the link to new particle formation, *Atmos. Chem. Phys.*, 6, 883–895, doi:10.5194/acp-6-883-2006, 2006.
- 10 Sakamoto, Y., Yabushita, A., Kawasaki, M., and Enami, S.: Direct emission of I₂ molecule and IO radical from the heterogeneous reactions of gaseous ozone with aqueous potassium iodide solution, *J. Phys. Chem. A*, 113, 7707–7713, 2009.
- 15 Schall, C., Heumann, K. G., and Kirst, G. O.: Biogenic volatile organoiodine and organobromine hydrocarbons in the Atlantic Ocean from 42° N to 72° S, *Fresen. J. Anal. Chem.*, 359, 298–305, 1997.
- Schwehr, K. A. and Santschi, P. H.: Sensitive determination of iodine species, including organoiodine for freshwater and seawater samples using high performance liquid chromatography and spectrophotometric detection, *Anal. Chim. Acta*, 482, 59–71, 2003.
- 20 Thorenz, U. R., Kundel, M., Müller, L., and Hoffmann, T.: Generation of standard gas mixtures of halogenated, aliphatic, and aromatic compounds and prediction of the individual output rates based on molecular formula and boiling point, *Anal. Bioanal. Chem.*, 404, 2177–2183, 2012.
- 25 Weisse, T., Grimm, N., Hickel, W., and Martens, P.: Dynamics of *Phaeocystis pouchetii* blooms in the Wadden Sea of Sylt (German Bight, North Sea), *Estuar. Coast. Shelf. S.*, 23, 171–182, 1986.
- Wong, G. T., Piumsomboon, A. U., and Dunstan, W. M.: The transformation of iodate to iodide in marine phytoplankton cultures, *Mar. Ecol.-Prog. Ser.*, 237, 27–39, 2002.

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Table 1. Halocarbon emission rates, concentrations of chlorophyll α , iodide and iodate in the four different sample suspensions.

| Sample | | F/2 medium background Range (Mean) | <i>P. glacialis</i> Range (Mean) | <i>M. helysia</i> Range (Mean) | plankton concentrate Range (Mean) |
|--|--------------------------------------|--|--|--------------------------------------|---|
| CH ₃ I | ng min ⁻¹ m ⁻² | 0.024–0.102 (0.050) | 0.030–0.098 (0.064) | 0.046–0.116 (0.075) | 0.011–0.053 (0.027) |
| CH ₂ ICl | ng min ⁻¹ m ⁻² | 0.004–0.039 (0.019) | 0.003–0.039 (0.028) | 0.05–0.038 (0.031) | 0.003–0.021 (0.013) |
| CH ₂ I ₂ | ng min ⁻¹ m ⁻² | 0.006–0.022 (0.020) | 0.073–0.117 (0.096) | 0.057–0.134 (0.098) | 0.011–0.024 (0.019) |
| CHBr ₃ | ng min ⁻¹ m ⁻² | 0.445–0.479 (0.457) | 0.503–0.549 (0.528) | 0.442–0.548 (0.529) | 0.442–0.590 (0.461) |
| chl α | $\mu\text{g L}^{-1}$ | n.d. | 257.27 | 926.59 | 2.53 |
| Iodide | nmol L ⁻¹ | 6.60–15.69 (10.35) | 7.32–19.71 (12.70) | 9.90–21.94 (16.84) | 3.52–9.45 (6.47) |
| Iodate | nmol L ⁻¹ | 402–538 (428) | 408–478 (448) | 397–499 (446) | 424–478 (442) |
| 1,3-C ₃ H ₆ Br ₂ [*] | ng min ⁻¹ m ⁻² | 1.568 ± 0.009 | 1.570 ± 0.12 | 1.569 ± 0.200 | 1.552 ± 0.014 |

* Evaporation standard given as mean ± standard deviation. Chl α was measured for each sample once halocarbons, iodide, iodate mean values and ranges are calculated from 6 replicates.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.




Figure 1. Experimental setup of the chamber with the phytoplankton suspension.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Emission of iodine containing volatiles by selected microalgal species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 2. Iodine emission rates normalized for the surface area of the different samples at 0 ppb and 100 ppb ozone. The error bars represent the standard deviation of the three replicates of each experiment.

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.




Figure 3. Correlation of the change in the I₂ emission in dependency on the iodide concentration in the phytoplankton suspension.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Emission of iodine containing volatiles by selected microalgae species

U. R. Thorenz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Function of the change in the I_2 emission rate in dependency of the amount of consumed ozone.