

Interactive comment on “Comparison of GEOS-5 AGCM planetary boundary layer depths computed with various definitions” by E. L. McGrath-Spangler and A. Molod

Anonymous Referee #1

Received and published: 20 March 2014

General comments:

This manuscript compared 7 methods to diagnose PBL depth in the GEOS-5 model. Such investigation is important for applications that use the PBL depth as input. However there are some deficiencies in its current form (see comments below). Thus I recommend major revision before it can be accepted for publication.

Major comments:

1. All the PBL depth diagnose methods are actually based on vertical profiles of variables such as potential temperature, wind, TKE etc. Vertical profiles of these variables

C613

describe/illustrate the boundary layer structure more clearly. So the comparison of those PBL depth diagnosed by different methods (e.g., Fig. 6) should be discussed with the aid of vertical profiles of those relevant variables. Good example are Fig. 3 of (Nielsen-Gammon et al., 2008), Fig. 1 of (Seibert et al., 2000)

2. Table 1 lists the PBL depth diagnose method for the model simulation. What is the method used to diagnose PBL depth based on radiosonde observations (particularly at night)? A separate section is needed in method section to describe the observations.

3. I would expect the nighttime PBL depth diagnosed by method 6 ($Ricrit=0$) is 0 in most places since there is always temperature inversion near the surface. I am a little surprised to see significant nighttime PBL depth from this method.

4. Some of the text is repetitive from the figure captions, e.g., “The horizontal dashed lines indicate the PBL depth found using the total K_h (Method 1, Fig. 7a) and bulk Richardson number (Method 4, Fig. 7b).”, which are unnecessary in the text.

Specific comments:

1. LN21, page 6593, The sentence of “estimates the PBL depth as the model level below that which K_h falls below” does not make sense.

2. “A spatial map of the JJA skin temperature (Fig. 3b) shows the same pattern as the PBL depth”. This is not true for the Arabian Peninsula. Any reason why?

3. “In general, both local Richardson number methods (Methods 5 and 6) estimate PBL depths that are lower than the other methods throughout the diurnal cycle.” Please analyze vertical profiles of relevant variables such as potential temperature, wind, TKE, local Ri (pick up one time in the day and one at night)

4. “This has implications for estimating the shallow nocturnal boundary layer that has been shown to be relevant for constituent transport”. This is not true for some air quality models that do not use the variable of PBL depth to compute transport of constituents. For these models, constituent mixing does not depend on the diagnosed PBL depth.

C614

5. Many studies investigated/compared PBL depth diagnose method (e.g., (Helmis et al., 2012; Hu et al., 2010; Nielsen-Gammon et al., 2008)). They need to be better summarized in introduction.

References:

Helmis, C.G., Sgouros, G., Tombrou, M., Schafer, K., Munkel, C., Bossioli, E., Dandou, A., 2012. A Comparative Study and Evaluation of Mixing-Height Estimation Based on Sodar-RASS, Ceilometer Data and Numerical Model Simulations. *Bound-Lay Meteorol* 145, 507-526.

Hu, X.M., Nielsen-Gammon, J.W., Zhang, F.Q., 2010. Evaluation of Three Planetary Boundary Layer Schemes in the WRF Model. *J Appl Meteorol Clim* 49, 1831-1844.

Nielsen-Gammon, J.W., Powell, C.L., Mahoney, M.J., Angevine, W.M., Senff, C., White, A., Berkowitz, C., Doran, C., Knupp, K., 2008. Multisensor estimation of mixing heights over a coastal city. *J Appl Meteorol Clim* 47, 27-43.

Seibert, P., Beyrich, F., Gryning, S.E., Joffre, S., Rasmussen, A., Tercier, P., 2000. Review and intercomparison of operational methods for the determination of the mixing height. *Atmos Environ* 34, 1001-1027.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 14, 6589, 2014.