

Interactive comment on “Enhancement of aerosols in UTLS over the Tibetan Plateau induced by deep convection during the Asian summer monsoon” by Q. S. He et al.

P. Seifert

seifert@tropos.de

Received and published: 4 February 2014

Dear Authors, dear Editor,

I would like to point you to the likely explanation of the observations described in the underlying manuscript.

On 13 June 2011 the Nabro volcano, located in northeastern Africa, erupted. Approximately 1.3 Tg of SO₂ were released into the upper troposphere and lower stratosphere during the first days after the eruption.

Within the following months the plume spread over the whole northern hemisphere.

C66

Its development was documented from space (Clarisso et al., 2013) and from ground (Sawamura et al., 2013; Uchino et al., 2013, Mateshvili et al., 2013).

The time of occurrence, vertical distribution, and optical properties presented in the underlying manuscript fit perfect to the presence of layers stemming from the Nabro volcano and volcanic aerosol in general. Also the lidar ratio of 30 sr agrees well to volcanic aerosol (Wandinger et al., 1995) whereas small, freshly nucleated aerosol particle should have small sizes (and be non-absorbing) and should therefore feature rather high lidar ratios (Müller et al., 2007).

I strongly suggest to the Editor to propose a withdrawal of the manuscript because the presented interpretation of the measurements may lead to misled follow-up studies. Of course, the paper can be renamed and re-submitted to provide information about the Nabro aerosol above the Tibetan Plateau. Due to the high altitude of the lidar station, the data quality obtained from the stratospheric layers is apparently quite good. If the authors still claim that the observed aerosol layers stem from new-aerosol-particle formation, they should provide a thorough model that can explain the formation of stratospheric aerosol by deep convection.

Yours sincerely, Patric Seifert.

References:

Clarisso, L., Coheur, P.-F., Theys, N., Hurtmans, D., and Clerbaux, C.: The 2011 Nabro eruption, a SO₂ plume height analysis using IASI measurements, *Atmos. Chem. Phys. Discuss.*, 13, 31161–31196, doi:10.5194/acpd-13-31161-2013, 2013.

Sawamura, P., J. P. Vernier, J. E. Barnes, et al.: Stratospheric AOD after the 2011 eruption of Nabro volcano measured by lidars over the Northern Hemisphere Environmental Research Letters, Vol. 7, No. 3. (06 August 2012), 034013, doi:10.1088/1748-9326/7/3/034013, 2013.

Mateshvili, N., Fussen, D., Mateshvili, G., Mateshvili, I., Vanhellemont, F., Kyrölä, E.,

Tukiainen, S., Kujanpää, J., Bingen, C., Robert, C., Tétard, C., and Dekemper, E.: Nabro volcano aerosol in the stratosphere over Georgia, South Caucasus from ground-based spectrometry of twilight sky brightness, *Atmos. Meas. Tech.*, 6, 2563-2576, doi:10.5194/amt-6-2563-2013, 2013.

Müller, D., A. Ansmann, I. Mattis, M. Tesche, U. Wandinger, D. Althausen, and G. Pisani (2007), Aerosol-type-dependent lidar ratios observed with Raman lidar, *J. Geophys. Res.*, 112, D16202, doi:10.1029/2006JD008292.

Uchino, O., Sakai, T., Nagai, T., Nakamae, K., Morino, I., Arai, K., Okumura, H., Takubo, S., Kawasaki, T., Mano, Y., Matsunaga, T., and Yokota, T.: On recent(2008–2012) stratospheric aerosols observed by lidar over Japan, *Atmos. Chem. Phys.*, 12, 11975-11984, doi:10.5194/acp-12-11975-2012, 2012.

Ulla Wandinger, Albert Ansmann, Jens Reichardt, and Terry Deshler, "Determination of stratospheric aerosol microphysical properties from independent extinction and backscattering measurements with a Raman lidar," *Appl. Opt.* 34, 8315-8329 (1995)

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 14, 3169, 2014.