

Interactive
Comment

Interactive comment on “Development towards a global operational aerosol consensus: basic climatological characteristics of the International Cooperative for Aerosol Prediction Multi-Model Ensemble (ICAP-MME)” by W. R. Sessions et al.

W. R. Sessions et al.

reidj@nrlmry.navy.mil

Received and published: 17 September 2014

Response to Reviewer 1 Sessions et al.

We thank the reviewer very much for the thorough review. We have done our best to update the paper. Certainly all points of clarification have been made. Some major conceptual changes suggested were not possible however because of the nature of the multi-model ensemble. This work is based on operational models for operational customers, so we are constrained by what operational centers provide. We have done

C7169

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



our best to further clarify.

1. <size resolved model request> ICAP participants and the multi model components are all operationally based. Thus, we need to take them as they come-which is all mass based. Some do have finer mass bins (like fine and coarse mode dust), but here is no evidence in AOT space that these are any better from a bulk score point of view. We choose to not go down this path at this time, as each center has its own reasons for its configuration. Further, with 4 core models and 7 dust models, there is simply not enough information in the time series to explore size resolving issues. There are so many other issues that dominate-like source functions, data assimilation protocols, and scavenging. AEROCOM has been comparing such simulations for years, and there is more to do. The point that we are making is that the multi-model ensemble is better than any single model. That is, you can get forecast improvements (e.g., in means and biases) using a multi-model ensemble vice using a single model.

2.The results in Fig.10 indicate that in general the model creates large biases and RMSE for the cases with $AOT > 0.6$. Have you ever attempted to increase the spatial resolution to see if it makes any differences? No, as we are attempting to leverage currently available products to produce a better forecast, and those available products are limited by each centers resources. One degree is the current largest common denominator for all models, which is why we chose it. There has been discussion to increase the ICAP MME to half a degree resolution in the future. When all models reach that benchmark, we could redo the experiment. If we could hazard a guess for high AOT events (which we do in the final section), the biggest problem is nonlinearity in the meteorological source functions or boundary layer processes.

3.It would be interesting to show some results on the heavy air pollution events. Compared to the individual models, how does the ensemble model perform in capturing such Originally we chose not to do pollution specifically but it is not clean cut and deserves a separate paper. But as we show Cape Verde (which is the most clean cut), the reviewer is right that we should show a site where things fail. The two worst sites for

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



verification are Kanpur and Beijing. Kanpur has a much more contiguous data record, so we added a final plot and discussion to that section to show where models all go wrong .

Line 8: what is bulk error statistics? Please give some explanations. The statistics used are explained in text. Bulk here just refers to them being taken over numerous observations.

The advantage of using multi-model ensemble relative to the individual model is NOT shown in abstract. Model bias in specific regimes, e.g. biomass burning, have been previously found by many aerosol models. It would be more interesting to highlight what the ICAP-MME model can do better than individual models.

What is highlighted here is that those specific regime bias within the individual models are lessened through utilization of the ensemble as a consensus model. Indeed, the conclusion of the abstract clearly states the benefit of using an ensemble “However, there is an overall AOT low bias among models, particularly for high AOT events. Biomass burning regions have the most diversity in seasonal average AOT. The southern oceans, though low in AOT, nevertheless also have high diversity. In regard to root mean square error, as expected the ICAP-MME placed first over all models worldwide, and was typically first or second in ranking against all models at individual sites. “

This is the first time in the literature that operational models have been compared. While these models often have roots in climate systems, they fundamentally are run differently, and serve different purposes (e.g., use of data assimilation, concentration on events versus seasonal bias). For RMSE, the most common metric for operations (rather than bias for climate models), the ensemble product is clearly better than any single model. We expected this result because of similar findings in other geophysical parameters, but for the first time we have shown it is true for any atmospheric composition variable. Regarding bias, despite the use of data assimilation, there is still a low bias. We thought that interesting.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Introduction: Line 24: ‘as an artifact’ what does this mean? It is an artifact of the radiance observations via SEVERI in Merchant et al 2006, as opposed to the model output from Evans.

Line 6-9: ‘while single-model probabilistic ensemble forecasting is clearly enhancing model solutions (particularly in data sparse regions), multi-model ensembles are an ever increasing tool for forecasters’. As you stated that the former can clearly enhance the model solution, why the latter is becoming an increasing tool? This varies by center, but the first order answer is that single-model ensembles are computationally resource intensive, while ICAP relies upon readily available products that are already budgeted.. Moreover, multi model ensembles have a degree of model independence which improves accuracy. Single model ensembles are useful for some applications (e.g., probabilities), but they still have their own built in biases and lack independence. The strength in different groups developing their own models is that they come up with algorithms that are more independent of each other than members of a single model ensemble.

2.1 input models: I would suggest separating this sub-section into two parts, with one for aerosol models and other for dust-only model, to make the description more clearly. This has since been moved to an appendix.

Does the same emission inventory is employed in all the aerosol models? It is not clear. Please include the information on the emissions, especially anthropogenic emissions. The models are largely independent in dust and smoke, but there is some cross usage on pollution. We have added this to the appendix text.

p.14940, line 5: why not simply examine the seasonal mean like JJA, DJF, MAM, and SON? This is a one year time series, so we are short on data for statistical significance. Even so, the character of DJF+MAM and JJA+SON was not significantly distinguished to divide further. This makes sense for when one looks at aerosol meteorology, and how many aerosol events are in transitional periods, two super seasons based nom-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

inally on solstice months (D-M and Jun-N) makes a lot more sense. We have been meaning to write a paper about that-probably will soon!

p.14943, line 15: 'MODS', should be 'MODIS'? fixed!

sec. 2.3 line 27: 'Instances of cirrus contamination (Chew et al., 2011) were evident in the level 1.5 (and to a lesser extent) level 2 products'. This sentence seems incomplete. Do you want to say 'Instances . . . level 1.5 (. . .) COMPARED TO level 2'? No, it is present in both. The 'and' in the parentheses should be outside of them to state, "... in the level 1.5 and (to a lesser extent) level 2." They have been removed in favor of commas.

Bottom line in p. 14951: when you removed the top five percent of coarse observations in NA and EA. Is that possible to remove the data in clear sky but with the heavy dust event? The wording of "in clear sky but with heavy dust event" is unclear to me. If you are asking if it is possible that actual dust events were removed when our clearing assumes that the observations are spurious, then the answer is yes-but we think it unlikely. We did do a hand analysis to make sure this was not the case.

Fig. 2: AOD in Winter/Spring is higher than in Summer/Fall over North Africa, which is contrary to what we expect, any explanations? It depends on the site. Cape Verde it is slightly higher in summer in AERONET (also figure). What also may be throwing things off conceptually are:

1) Our two season scheme includes the low dust months of Oct-November 2) Statistics when there was QAed AERONET data. 3) Dust events for the second half of 2012 and into 2013 were potentially suppressed by anomalously wet conditions that year (http://www.cpc.ncep.noaa.gov/products/fews/AFR_CLIM/arch/arc2_wam_2012_anom.jpg) while there was a strong May dust season.

Please keep in mind, this is a verification and ensemble paper over a limited period, not a climatology paper.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



As for other sites like Ilorin, wintertime frontal activity brings dust to the region. In the summer, winds are more barotropic and easterly and thus the dust does not get that far south. Besides, the ITCZ is parked over it at that time of year. Just check out the AERONET website http://aeronet.gsfc.nasa.gov/cgi-bin/type_one_station_opera_v2_new Ilorin is more dusty and smoky in Dec-April. This said, we always worry about cirrus contamination in this part of the world. We hope the next version of AERONET will be able to detect it.

Table 1 and 2: It would be good to have a map plot showing the locations of these sites. Site map added as Figure 1b.

Fig. 5 and top line on p.14957: bias in Baegnyeong improves with time, why? Does that imply there is problematic in the analysis? Analysis issues are potentially the causes. (p14957, L3-5)

Table 4 and line 13 on p.14957: what metric shown in Table 4? biases or RMSE? We made a typo in the header, not the caption. These are RMSE values.

Fig. 6 and line 14 on p.14957: small dots are each model's value. What value? AOT? These are model AOT values, with marker face indicated which model.

Line 3-5 on p. 'On average, the RMSE's of the 1 day forecasts of ICAP-MME run approximately 50% of the climatological mean. Dust AOT forecasting is superior to overall fine and coarse mode 5 AOT, running approximately 1/3rd of climatological AOT.' How do you come up to this conclusion? Please show more details. We have modified the language a bit for clarity. "Based on the slope of RMSE against mean AOT value for each site in Figure 7, the RMSE's of the 1 day forecasts of ICAP-MME run approximately 50% of the climatological mean AOT value. Dust AOT forecasting is superior to overall fine and coarse mode AOT, running approximately 1/3rd of climatological AOT. Again, this is part reflects the importance of the dust species by centers. Further, the AERONET Cape Verde site (in which RMSE is particularly skillful) is a common benchmark site for Saharan dust-hence models are typically tuned for the region."

Line 13 on p.14958: please give the definition of fractional gross error. Fixed.

Line 20 on p.14958: 'although the dominance of the ICAP-MME generally increases in time, particularly for dust'. What do you mean? We changed the verbage a bit for clarification: "Like bias, forecasting skill for all models and the ensemble mean degrades in time. Although the relative performance of the ICAP-MME mean relative to the member models increases in time, particularly for dust."

Fig.9. how is the rank and histogram computed here? Did you put all the results from the individual models and the ensemble together for this calculation? Please add more details on how you obtain the results in order to help the readers to understand. We added a bit more to the explanation. It is the result of taking all the member models at a valid time for a particular observation, then ranking that observation as if it were another member.

Line 20 on p.14966: if 'Experience has shown however that equal weighting in a consensus style appears to provide the most robust results overall, and this is backed up on both practical and theoretical grounds (DeSole et al., 2013)', Then why 'we intend to convert the ICAP-MME to a super ensemble where models are weighted by their scores (e.g., Krishnamurti et al., 1999; Casanova and Ahrens, 2009)'. It is confusing here. Also, can you give some explanations on why equal weighting is most robust? You have cut off the important first half of the "we intend" sentence. The statement is that we receive questions about future intent, including whether or not to produce a weighted super-ensemble, which we answer by stating it would be less robust. The short answer is that while there have been many expectations from super ensembles, in reality they rarely stand the test of time as they require prior knowledge of error characteristics. This is difficult to obtain as a) models are always improving, b) even "bad" models can result in an improved consensus, and c) one training data set often is not applicable to another season. But at some point we need to address the issue more directly. For now, we clarified this statement.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C7176

