



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Concentrations and solubility of trace elements in fine particles at a mountain site, southern China: regional sources and cloud processing

T. Li¹, Y. Wang¹, W. J. Li², J. M. Chen^{1,2}, T. Wang³, and W. X. Wang²

¹School of Environmental Science and Engineering, Shandong University, Jinan, Shandong 250100, China

²Environment Research Institute, Shandong University, Jinan, Shandong 250100, China

³Department of Civil and Environmental Engineering, the Hong Kong Polytechnic University, Hong Kong, China

Received: 10 February 2015 – Accepted: 14 April 2015 – Published: 5 May 2015

Correspondence to: Y. Wang (wy@sdu.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The concentrations and solubility of twelve trace elements in PM_{2.5} at Mt. Lushan, southern China, were investigated during the summer of 2011 and the spring of 2012. The average PM_{2.5} mass was $55.2 \pm 20.1 \mu\text{g m}^{-3}$ during the observation period. Temporal variations of all trace elements including total and water-soluble fractions with several dust storm spikes for total fraction Al and Fe were observed. The enrichment factor (EF) values were one order of magnitude higher for the water-soluble fractions vs. the total fractions of trace elements. Four major emission sources were classified by principal component analysis (PCA), namely nonferrous metal mining and smelting (for Cr, As, Ba and parts of Zn), coal combustion (for Pb, Zn, Se, Cu and Mn), crustal materials (for Al and Fe) and municipal solid waste incineration (for Cd and Mo). Trajectory cluster analysis and the potential source contribution function (PSCF) consistently identified the Yangtze River Delta (YRD), the Pearl River Delta (PRD) and parts of Hunan and Jiangxi as the major source regions and pathways for anthropogenic elements, while northern China was identified for crustal elements. In contrast, the local Jiangxi area has become the most significant contributor to the solubility of most trace elements, apart from the YRD with severe air pollution. In addition, the solubility alteration of trace elements in cloud events was investigated and transmission electron microscopy (TEM) analysis indicated that the irreversible alteration of particle morphology by cloud processing was highly responsible for the enhancement of element solubility. Our work implies an important role of regional anthropogenic pollution and cloud processing in the evolution of trace element solubility during transport.

1 Introduction

Epidemiologic studies have associated long-term exposure to highly inhalable ambient fine particulate matter ($\leq 2.5 \mu\text{m}$, PM_{2.5}) with many adverse health outcomes such as cardiovascular and respiratory morbidity and mortality, whether in urban or rural

ACPD

15, 13001–13042, 2015

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



areas (Hoek et al., 2013; Cao et al., 2012; Weichenthal et al., 2014). Sustained exposure to high concentrations of PM air pollution has been suggested to contribute to a decrease in life expectancy of approximately 3.0 y in China (Chen et al., 2013). The long atmospheric lifetimes of days or even weeks give fine particles opportunities to be subject to long-range transboundary or intercontinental transport in the air and to deposition toward remote areas, carrying abundant anthropogenic pollutants and affecting ecosystems (Mahowald, 2011). Fine particles are also responsible for regional and global climate change and the hydrologic cycle interference through radiative forcing directly by reflecting sunlight and indirectly by changing cloud properties and precipitation acting as cloud condensation nuclei (Kaufman et al., 2002).

Although trace elements constitute only a small proportion of fine particle mass, their negative impacts on human health and ecosystems have attracted considerable attention because of their toxicity and bioaccumulation by inhalation and deposition. Toxicological and epidemiologic studies often suggest trace metals, particularly water-soluble metals which are more easily bioavailable, as the critical components harmful to the cardiopulmonary system through oxidative chemistry (Cakmak et al., 2014; Costa and Dreher, 1997). Soluble trace elements, especially the transition metals, are likely to be the primary drivers of the generation of reactive oxygen species, inducing cellular inflammation (Charrier et al., 2014; Cheung et al., 2012; Shafer et al., 2010). Besides, trace metal ions also play an important catalysis role in secondary aqueous-phase (Harris et al., 2013) and particle-phase (Clements et al., 2013) sulphate formation, heterogeneous production and elimination of gas-phase hydrogen peroxide (Guo et al., 2014a) as well as multiphase cloud chemistry (Deguillaume et al., 2004). Importantly, most anthropogenic trace elements in ambient air are considered to be mainly concentrated on fine particles, strengthening their toxicity. Therefore, understanding both the concentrations and water solubility of fine particle trace elements is essential for revealing their bioavailability in ambient environments and reactivity in atmospheric chemistry.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diverse characteristics and sources of atmospheric trace elements have been investigated in various areas. For example, regional background, urban and industrial sites in Spain were found to have clearly different concentration levels and source origins of trace elements in PM (Querol et al., 2007); technogenic metalliferous fine aerosols observed in urban Montseny demonstrated the spatiotemporal variability of trace metal pollution (Moreno et al., 2011); traffic, static combustion and crustal were suggested as the main sources for Fe, Zn, Pb and Cu in PM_{2.5} in an urban background area in Edinburgh (Heal et al., 2005); meanwhile, stationary industrial emissions from coal combustion were confirmed as the contributor to substantial atmospheric lead pollution of PM_{2.5} in Shanghai (Chen et al., 2008). The United States has already established a spatially gridded national emissions inventory of PM_{2.5} trace elements in speciation profiles with the dominant sources of crustal, biomass burning, coal combustion and industrial processing (Reff et al., 2009). In Asia, atmospheric metallic element pollution by human activities has been surveyed, with the highest metal concentrations occurring in China (Fang et al., 2010). Coal combustion, nonferrous metal smelting and iron and steel manufacturing are found to be the major disproportionate sources of atmospheric hazardous trace elements in China according to the latest anthropogenic emissions inventory, which highlighted an increasing trend in air pollution by toxic heavy metals (Cheng et al., 2015).

Recently, many field observations have found that even remote mountains and oceans have been polluted by anthropogenic trace elements in fine particles, significantly confirming the contributions of long-range transport (Deng et al., 2011; Schwab, 2004; Fomba et al., 2013). A few studies on dust in Asia (Takahashi et al., 2011; Hsu et al., 2013) indicate that the water solubility of trace elements could be modified during long-range transport, probably affected by emission sources, anthropogenic pollution, acid processing and photochemical aging, etc. (Hsu et al., 2010). However, little information about fine particle trace elements has been reported with regard to the identification of source regions or pathways, and the main mechanism of trace element dissolution by atmospheric processing is still unclear.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



This paper provides a dataset of concentration and solubility levels of trace elements in fine particles at Mt. Lushan, a high, rural mountain within an acid precipitation area in southern China. We focused on identifying the potential source region distributions for individual elements, aiming to evaluate the contributions of local, regional and long-range transport. Regional contributions to element solubility were also preliminarily investigated. One condensation-evaporation cloud cycle can increase aerosol solubility, and atmospheric aerosols may experience up to ten cloud cycles before removal (Desboeufs et al., 2001; Spokes et al., 1994); hence, we intensively discussed the influence of actual cloud processing on the alteration of fine particle trace element solubility, considering the rather frequent cloud events at Mt. Lushan. The size, mixing state and chemical composition of individual metal particles in cloud residues were detected using transmission electron microscopy (TEM) to investigate the effects of microphysical properties on the evolution of element solubility.

2 Methodology

2.1 Site description

Mt. Lushan is one of the National Geoparks of China, with an area of 302 km², and is situated in northern Jiangxi Province, southern China, adjacent to the Yangtze River and Poyang Lake, which is the largest freshwater lake in China, with an area greater than 4000 km² (Fig. 1). Mt. Lushan is significantly influenced by a subtropical monsoon and mountain climate, with frequent cloud events occurring from spring to autumn. The sampling site is located in Guniubei (115°59' E, 29°35' N, 1165 m a.s.l.), the top of Guling town where most residents work in tourism or related services and little pollution is produced. There are intensive coal-fired power plants located at the eastern coastal area, and abundant mineral resources and many large non-ferrous industries exist in Jiangxi and in neighbouring provinces. Therefore, Mt. Lushan is a favourable platform

for observing the long-range transport of regional air pollution and for evaluating the influence of atmospheric particles on the local ecological environment.

2.2 Sample collection

Daily PM_{2.5} samples were collected on quartz fibre filters (MK 360, 90 mm, Munktell, Switzerland) during the summer of 2011 (8 August–23 September) and the spring of 2012 (18 March–20 May). The filters were heated at 500 °C for 3 h prior to sampling, subsequently balanced at 25 °C and 50 ± 5 % relative humidity for at least 24 h and weighed three times before and after collection. The sampling procedures were conducted during non-rainy and non-cloudy periods by employing a PM_{2.5} impactor (TH-150A, Tianhong Co., China) with a medium flow rate of 100 L min⁻¹. Seventy-six valid ambient samples and 6 blank filters were obtained and stored in the dark and at -20 °C for laboratory analysis.

Individual cloud residues in two selected cloud events (on 11 September 2011 and 18 April 2012) were trapped onto copper TEM grids coated with carbon film using a single-stage cascade impactor with a 0.5 mm-diameter jet nozzle under a flow rate of 0.5 L min⁻¹. The grids were sealed in a dry plastic tube and stored in desiccators. More detailed information can be found in (Li et al., 2013).

2.3 Extraction procedure

2.3.1 Total fraction (acid)

To obtain the total concentration of trace elements in fine particles with the least loss of semi-volatile elements, one quarter of the samples and blank filters were digested with an acid mixture (5 mL 65 % HNO₃ and 2 mL 30 % H₂O₂, guaranteed reagent grade, Kemiou Co., China) in closed Teflon vessels using a microwave digestion system according to a controlled gradient temperature procedure: reaching 120 °C in 5 min and remaining there for 5 min, then reaching 160 °C in 5 min and remaining there for 2 min,

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and finally reaching 185 °C in 5 min and remaining there for 15 min. After cooling to room temperature, the solutions were filtered and subsequently diluted to 25 mL with high-purity deionised water ($\geq 18 \text{ M}\Omega\text{CM}$) in acid-cleaned brown glass bottles.

2.3.2 Water-soluble fraction (water)

5 Half of the samples and blank filters were extracted with 50 mL high-purity deionised water ($\geq 18 \text{ M}\Omega\text{CM}$) using ultra-sonication for 1 h followed by filtration. Solutions of approximately 25 mL were preserved in acid-cleaned brown glass bottles with 1 % v/v high pure hydrochloric acid to lessen adsorption of the elements by the glass walls. All solutions were stored at 4 °C until instrument analysis.

10 2.4 Chemical and physical analysis

The concentrations of the total and water-soluble fractions of twelve trace elements (including Al, Cr, Mn, Fe, Cu, Zn, As, Se, Mo, Cd, Ba and Pb) were measured by Inductively Coupled Plasma Mass Spectrometry (ICP-MS, Agilent 7500a) based on the EPA 200.8 method using internal standard substances of Li, Sc, Ge, Y, In, Tb and Bi to eliminate matrix interference. National standard materials (soil, GSS-4, China) were digested for calculating element recoveries by the acid extraction method. All recoveries were found to be a little lower than 100 % (Table S1), ranging from 93.0% for Cd to 72.7 % for Se, primarily ascribed to volatilisation losses and acid mixture that is not strong enough to release all mineral constituents from the silica matrix into the solution. Reagent and filter blanks were also prepared and analysed for background element content. The detection limits of the ICP-MS analysis for the total fraction elements determined were calculated as three times the SD (3δ) of the six blank values (Table S1 in the Supplement).

Individual metal particles in cloud residues on the TEM grids were analysed with
25 a JEM-2100 TEM operating at 200 kV accompanied by a semi-quantitative determina-
tion of elemental composition by an energy-dispersive X-ray spectrometer (EDS) that

25

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



W_{ij} to reduce the effect of small values of n_{ij} and to better reflect the uncertainty in values for these cells (Polissar et al., 2001). The weighting function reduced the PSCF values when the total number of the endpoints in a particular cell (n_{ij}) was less than approximately three times the average value (n_{Ave}) of the endpoints per each cell, defined as

$$W_{ij} = \begin{cases} 1.00 & n_{ij} > 3n_{Ave} \\ 0.70 & 1.5n_{Ave} < n_{ij} \leq 3n_{Ave} \\ 0.42 & n_{Ave} < n_{ij} \leq 1.5n_{Ave} \\ 0.17 & 0 < n_{ij} \leq n_{Ave} \end{cases} \quad (2)$$

3 Results and discussion

3.1 Elemental composition

3.1.1 General characterisation

The ambient atmospheric concentrations of PM_{2.5} and trace elements including total and water-soluble fractions at Mt. Lushan are summarised in Table 1, and the comparison with typical mountains and megacities in China are listed in Table S1. The average PM_{2.5} concentration at Mt. Lushan in spring ($54.7 \mu\text{g m}^{-3}$) was approximately that of summer ($55.9 \mu\text{g m}^{-3}$), more than twice the WHO air quality guideline for daily PM_{2.5} of $25 \mu\text{g m}^{-3}$, but still lower than many urban sites at ground level such as Beijing ($118.5 \mu\text{g m}^{-3}$) and Guangzhou ($81.7 \mu\text{g m}^{-3}$) (Yang et al., 2011). It was clearly contrary to Mt. Tai (Deng et al., 2011), where PM_{2.5} exhibited a concentration 2–3 times higher in summer ($123.1 \mu\text{g m}^{-3}$) than in spring ($46.6 \mu\text{g m}^{-3}$), suggesting a less-pronounced seasonal variation of PM_{2.5} at Mt. Lushan.

The total fraction concentrations of twelve trace elements in spring were comparable with those in summer at Mt. Lushan, with the exception of higher Al in spring.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Overall, elements Al and Fe contributed the highest concentration, and the other elements decreased from 258.3 ng m^{-3} for Zn to 2.0 ng m^{-3} for Mo. The dominant elements (Al, Fe and Zn) exhibited significantly lower concentrations at Mt. Lushan than Mt. Tai (1200 ng m^{-3} for Al, 810 ng m^{-3} for Fe and 400 ng m^{-3} for Zn) (Deng et al., 2011) and slightly higher than Mt. Gongga (295.8 ng m^{-3} for Al, 224.0 ng m^{-3} for Fe and 154.6 ng m^{-3} for Zn) (Yang et al., 2009a), probably due to the geomorphology and climatology biases. As shown in Table S1, all elements (except Ba) have lower concentration levels in comparison with Mt. Dinghu, a national nature reserve influenced by atmospheric pollution (Yang et al., 2009b) and with typical megacities in China such as Beijing, Guangzhou (Yang et al., 2011) and Shanghai (Chen et al., 2008). For water-soluble fractions, Zn transformed into the primary elements, followed by Al and Fe, with higher concentrations in summer than spring; other trace elements (from Pb to Mo) were still comparable between summer and spring.

3.1.2 Temporal variations

Figure 2 displays the temporal variations of trace element concentrations in $\text{PM}_{2.5}$ at Mt. Lushan for total and water-soluble fractions during the sampling campaigns. Most total fraction elements exhibited more intense variances in spring than in summer, with unique temporal patterns such as the highly variable pattern observed for aluminium and the relatively steady pattern of arsonium. Similar temporal patterns were observed for the water-soluble and total fractions for most elements, except for Al and Fe, which had drastic variations with many spike episodes for total fractions but lower constant concentration for water-soluble fractions. The dependence of temporal variations of water-soluble elements on their total fractions were also examined (Fig. S1). The concentrations of water-soluble Zn, Cu, Mn, As, Se and Cd were found to be rather consistent with their total fraction according to their high correlations ($R^2 = 0.50\text{--}0.78$), whereas there was only a weak association between the total concentrations and water-soluble fractions of Al, Fe and Cr.

Several spikes in element concentration were observed in specific periods. The first spike (Spike I) for all elements from 11–13 August 2011 was ascribed to the short sampling time due to the interruption by an unexpected thundershower. However, the significantly elevated total concentration of Al, Fe and Mn during 23–25 March 2012 (Spike II) and 25 and 26 April 2012 (Spike III) with no distinct elevation in the other elements were observed (Fig. 2), likely contributed by the dust storms originating from the Gobi desert and the Taklimakan desert in spring according to the MODIS image, the OMI Aerosol Index and air mass backward trajectory analysis (Fig. S2). These features were similar with the Asian Dust observed at Huaniao Island in the East China Sea, where concentrations of Al, Fe, Mn and Ba 3–4 times higher were observed with no significant difference for Zn, Cu, As and Cd in the air compared with non-dust days (Guo et al., 2014b). In addition, the dust storms contributed little to the water-soluble fractions of all trace elements.

3.2 Source identification

3.2.1 Correlations between individual elements

The Pearson correlation coefficients r between each element for both total fraction and water-soluble fraction were examined (Table S3). Moderate significant associations of Fe with elements Al, Mn and Cr are found to exist both in total and water-soluble fractions with comparable coefficients between 0.40–0.68. Correlations for other trace elements in water-soluble fraction are stronger than in total fraction, suggesting the higher possibilities of common sources for some water-soluble elements. For example, the correlation levels of Zn–Cu, Zn–Se, Pb–Zn and Pb–Se are dramatically high for water-soluble fractions ($0.72 < r < 0.82$) and moderate for total fractions ($0.25 < r < 0.52$), which could be attributed to their common anthropogenic sources of coal-fired power plants and industrial processes based on the suggestion of the spherical morphology and elemental composition of individual metal particles via the TEM examination (Li et al., 2014).

3.2.2 Anthropogenic pollution

The enrichment factor (EF) for individual trace elements was applied to identify their general crustal and anthropogenic sources and evaluate the degree of pollution using the following formula:

$$EF_i = (X_i/X_R)_{\text{aerosol}} / (X'_i/X'_R)_{\text{crust}} \quad (3)$$

where EF_i is the enrichment factor of element i ; X_i and X_R are the concentrations of element i and reference element R in aerosol, respectively; X'_i and X'_R are the background contents of elements in Chinese soils (Wei et al., 1991); and Al is selected as the reference element for calculation. Element sources are classified into three groups with the following standard: $EF < 10$ is considered to be a crustal origin without enrichment; $10 < EF < 100$ comes from mixed origins (crustal and anthropogenic sources); $EF > 100$ indicates air pollution from an anthropogenic origin. Figure 3 describes the EF values for both the total and water-soluble fractions of each element in a decreasing order. Elements with $EF > 100$, including Se, Cd, Zn, Pb, As, Mo and Cu, were found to be highly enriched, ranging from hundreds to tens of thousands in both fractions, indicating the severe anthropogenic pollution of atmospheric trace elements. Cr, Ba and Mn were likely to be from mixed origins because the majority of their EF values fell within 10–100. Fe was predominantly from a crustal origin based on its $EF < 10$. It is worth noting that the water-soluble fractions of most elements (except Cr and Fe) have higher EF values by approximately one order of magnitude than the total fractions, especially for elements with extremely high EF values, implying the inclination of enormous pollution toward the water-soluble fraction of anthropogenic elements.

3.2.3 Emission source classification

The principal component analysis (PCA) was performed in detail to classify the main sources of trace elements in $PM_{2.5}$ at Mt. Lushan. The PCA results for trace elements in total fraction are shown in Fig. 4 with four major components identified.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The first principal component was considered primarily from nonferrous metal mining and metallurgical smelting, accounting for 34.4 % of the total variance with high loadings for Cr, As, and Ba and considerable loadings for Zn and Mo. Although there were diverse EF values which suggested anthropogenic origins mixed with crustal materials ranging from dozens for Cr and Ba to thousands for As, Zn and Mo, these total fraction elements still presented considerably moderate correlations. According to the provincial output statistics of nonferrous metals in 2010 (Cheng et al., 2015) and the Overall Planning of Mining Resources of 2008–2015 for Jiangxi and adjacent Hunan Province, there are abundant nonferrous minerals (such as barite, realgar and sphalerite, etc.) and many large-scale metallurgical industries within and near Jiangxi Province. In addition, high concentrations of Pb and Zn in cloud drop residues at Mt. Lushan were suggested deriving from fine particles contributed by nonferrous smelting in Jiangxi Province (Li et al., 2013).

The second principal component that contributed to high loadings for Pb, Cu, Se, Zn, and Mn was identified as coal combustion. Coal combustion is representatively considered to be the source of most atmospheric trace elements in China. All species in this component except Mn exhibited high EF values, indicating their anthropogenic origins. The semi-volatile Se is usually used as a tracer for determining atmospheric sources and even the pathways of coal combustion pollutants, because it is primarily produced in high temperature combustion and can undergo long-range transport with fine particles after the rapid gas-to-particle conversion (Husain et al., 2004; Wen and Carignan, 2007). Coal combustion is also becoming the major contributor to atmospheric Pb with the nationwide phase-out of leaded gasoline in China (Xu et al., 2012). The high correlations between Se and the other species in Component 2 clearly indicated the likely sources of coal combustion.

The third component in particular explained most of the Al and Fe which are the typical signatures of crustal materials or soils, along with relatively high correlations and low EF values.

Municipal solid waste incineration was regarded as the primary sources of fine particle Cd and Mo at Mt. Lushan, explaining 8.4 % of the variance. Both Cd and Mo had low concentrations but high EF values, with a strong association in total fraction ($r = 0.71$). Jakob et al. (1995) found that Cd in fly ash of municipal solid waste incinerators could evaporate into the atmosphere completely during heat treatment. The rapid increase in atmospheric Cd emissions from municipal solid waste incineration in recent years and the unbalanced large emissions in east-central and southeast China (Tian et al., 2012) were responsible for fine particle Cd at Mt. Lushan.

3.3 Potential source region distributions

Three-day air mass backward trajectories were calculated to estimate the potential source regions and pathways of air pollutants, making full use of the geographical advantages of the high-altitude observation site and the long-range transport capacity of fine particles. The air masses associated with PM_{2.5} samples were classified into five clusters (Fig. 5a). The air masses of the Southwest (SW) cluster originated from the South China Sea and passed through the Pearl River Delta (PRD), Hunan and Jiangxi Province with the highest occurrence frequency of 28.9 %. Cluster East (E) mainly came from the East China Sea and partly from the Circum-Bohai Sea Region, traversing the Yangtze River Delta (YRD) and Anhui Province at a frequency of 27.6 %. Air masses of cluster Local (L) were found to be advancing toward or revolving around Mt. Lushan slowly from ground-level within areas covering distances of approximately 200 km away from the observation site, contributing 22.4 % of the occurrence frequency. Clusters Northwest (NW) and North (N) both went through the China Loess Plateau and accounted for 13.2 and 7.9 % of the occurrence frequency, respectively. The primary difference was that the former mainly originated from Taklimakan and the Gobi Desert, whereas the latter was from remote Siberia and Mongolia with a much higher wind speed.

Figure 5b shows the averaged PM_{2.5} concentration of different clusters. Cluster E exhibited the highest PM_{2.5} concentration of $67.0 \mu\text{g m}^{-3}$, followed by cluster L

($59.6 \mu\text{g m}^{-3}$) and the similar clusters NW and N, whereas cluster SW had the lowest concentration ($45.3 \mu\text{g m}^{-3}$). The order of $\text{PM}_{2.5}$ concentration seemed to be consistent with the PSCF result (Fig. 5c) which integrated $\text{PM}_{2.5}$ concentration with air mass occurrence frequency. As shown in Fig. 5c, the YRD and southern Anhui Province were the most likely source regions of $\text{PM}_{2.5}$ at Mt. Lushan. These economically developed regions exhibit massive energy consumption and industrial production, becoming important anthropogenic source regions in eastern China. The local area of Jiangxi, neighbouring southern Anhui and eastern Hunan were also significant $\text{PM}_{2.5}$ source regions, where many nonferrous metal smelting industries are widely distributed. However, the contributions to $\text{PM}_{2.5}$ at Mt. Lushan by long-range transport of air masses from northern China (Inner Mongolia and the Loess Plateau) were much less than from eastern China and local regions, suggesting nearly negligible impacts of mineral dusts or soil transport on fine particle mass concentration at Mt. Lushan compared with regional anthropogenic pollution. In addition, the $\text{PM}_{2.5}$ concentration in five clusters or different source regions were highly dependent on the inorganic ions, especially the secondary ions (Li et al., 2014). Interestingly, the PRD contributed much less concentrations to fine particles in spite of its high occurrence frequency, probably due to particle scavenging by frequent rainfall during the Asian summer monsoon.

Figure 6a–l illustrates the distinctive potential source region distributions for individual elements in fine particles at Mt. Lushan. Crustal elements Al and Fe exhibited similar source region distributions (Fig. 6a and b). The highest PSCF values in northern China including the Loess Plateau, Henan and eastern Hubei Province indicated that these areas were very important source regions and pathways for fine particle Al and Fe, with vast contributions of long-range transport of mineral dusts or soils. The PRD and eastern Hunan Province were likely to be other important source regions for Al and Fe probably due to the influence of mining activities, whereas the YRD in eastern China and the local area of Jiangxi Province made much lower contributions.

Fine particle As (Fig. 6c) had an almost parallel pattern of source region distribution with Fe, except the discordance that the PRD, eastern Guangxi, Hunan and

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jiangxi Province were identified as the most important source regions and pathways, attributable to the emissions of the nonferrous metals mining and smelting industries. Northern China also tended to be an important potential source region, suggesting the possible contribution of natural sources for As. Similar to As, the PRD, Hunan and Jiangxi Province, as well as northern China were also potential source regions for fine particle Mn (Fig. 6d). However, the YRD area was characterised by extremely high PSCF values for Mn, indicating an important anthropogenic source region.

Fine particle Pb, Zn, Cu and Se (Fig. 6e–h), which were identified primarily from coal combustion by the PCA analysis, had similar source region distributions with some discrepancies, generally coinciding with the geographical distribution of coal-fired power plants in China which were aggregated within coastal areas such as the YRD, the PRD and Shandong Province (Fig. S3a). The YRD area including Jiangsu, Zhejiang and southern Anhui Province was an especially pronounced source region for Pb, Zn, Cu and Se at Mt. Lushan. The significantly high PSCF values in the YRD region were attributable to the large coal consumption by intensive power plants in Jiangsu, Zhejiang and even Shandong Province and to the differences in trace element contents in raw coal such as high Se contents in Anhui Province, both of which made great contributions to emissions of atmospheric trace elements (Tian et al., 2014). Meanwhile, the PRD, Hunan and the local area of Jiangxi Province also played a remarkable role in source region distributions of Zn, Cu and Se at Mt. Lushan, for which coal-fired power plants and nonferrous metals smelting industries in these areas were primarily responsible. Unexpectedly, northern China rather than the PRD area was also identified as a potential source region of Pb, possibly attributed to the partial contributions of long-range transport of mineral dusts for fine particle Pb at Mt. Lushan.

As shown in Fig. 6i and j, the YRD, the PRD, eastern Hunan and western Jiangxi were identified as the common likely source regions and pathways for fine particle Cd and Mo at Mt. Lushan. The identified source regions corresponded very well with the uneven regional distribution of municipal solid waste (MSW) incineration plants in China (Fig. S3b), with 183.5, 175.1 and 146.6 kg of atmospheric Cd emitted by MSW

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



incineration from Jiangsu, Zhejiang and Guangdong, respectively, the top three largest emitting provinces in 2010 (Tian et al., 2012). Figure 6k shows an apparently more homogeneous source region distribution for Ba, with the PRD and areas surrounding Mt. Lushan as potential source regions which have a high natural background content of barite. On the contrary, few potential source regions were identified for Cr (Fig. 6l) besides the PRD, parts of Hunan Province and a small region in central Zhejiang with low PSCF values.

It should be noted that the potential source region distributions for trace elements at Mt. Lushan were very different with $PM_{2.5}$. The YRD, the PRD and Hunan Province were likely to be the major source regions for most anthropogenic elements (except Cu within eastern Jiangxi), and northern China was likely for crustal elements (Al and Fe). However, the contribution probabilities of areas near Mt. Lushan for most elements were obviously lower than other source regions. That revealed that long-range transport and regional anthropogenic sources, rather than local sources, made much more significant contributions to fine particle trace elements at Mt. Lushan, which could be ascribed to the different types and quantities of emissions as well as the protection policies for Mt. Lushan and its adjacent Poyang Lake.

In addition, the order of mass concentration of individual trace elements for five air mass clusters ending at Mt. Lushan (Fig. S4) was in accordance with the distributions of likely source regions identified by PSCF values, further suggesting greater contributions by regional sources and long-range transport to fine particle trace elements at Mt. Lushan. For example, clusters N and NW contributed the highest concentration for crustal Al and Fe and anthropogenic Zn and As, followed by the highest cluster SW, E and the lowest cluster L; exceptionally, cluster L made the highest contribution to the combustion element Cu.

3.4 Trace element solubility

3.4.1 Comparison of individual element solubility

In this study, element solubility is defined as the percentage of water-soluble concentrations of an element divided by its total concentration. Figure S5 compares the average water solubility of individual elements in PM_{2.5} at Mt. Lushan with other sites in decreasing order. The most soluble elements were As, Mn, Cu, Zn and Se with a solubility reaching up to 60–70 %; followed by Cd, Pb, Ba and Mo with moderate solubility, approximately 40–50 %; whereas the least soluble elements were Fe, Al and Cr with a solubility less than 30 %. Compared with TSP over the East China Sea (Hsu et al., 2010), some elements (Mn, Cu, Ba, Mo, Fe and Al) were significantly highly soluble in PM_{2.5} at Mt. Lushan, whereas other species (As, Zn, Se and Cd) showed the opposite with the exception of comparable Pb and Cr. Similarly, compared with the PM_{2.5} at Edinburgh, UK (Heal et al., 2005), the elements Mn, Cu and Fe exhibited higher water solubility, whereas the elements As, Zn and Pb were comparable, and the elements Cd and Cr were lower at Mt. Lushan. A large gap in anthropogenic elements (As, Mn, Cu and Pb) between Mt. Lushan and Nanjing (Hu et al., 2012) was also observed. Experimental conditions such as digestion methods or analysis instruments may result in the solubility discrepancies at different sites, but they are far from the determinant.

3.4.2 Effect of emission sources on element solubility

Element solubility is associated with four major emission sources identified by EF value and PCA analysis and displayed in Fig. 7. Crustal elements with mean EF < 10 had low average solubility (< 30 %) besides the highly dissolved Mn that was possibly impacted by a combustion source; the solubility of elements related to mining and metallurgy varied between < 10–70 % with a 10–1000 range in EF value; coal combustion elements characterised by high EF values over 100 exhibited a high solubility of approximately 45–75 %, contributable to the combustion process during which oxidisable elements in

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on the effects on climate (Jickells et al., 2005). As shown in Fig. 8a, northern China, identified as the major source region for crustal Fe, made little contribution to its solubility. The significantly low solubility of fine particle Fe in northern China, much lower than the PSCF criteria value, was consistent with the very low soluble Fe (solubility less than 2%) from desert dusts or soils (Baker et al., 2006; Jickells et al., 2005). The experiment performed by (Mackie et al., 2006) suggests that the amount of readily released Fe does not increase after the uplift and abrasion of soils in areas with little atmospheric pollution; thus, the low solubility of the Fe from northern China could be attributed to the initial absence of dissolved Fe in dusts or soils derived from the Gobi desert or Loess Plateau. On the contrary, the relatively much higher PSCF values for Fe in the local area of Jiangxi, southern Anhui and the YRD indicated greater contributions of anthropogenic pollution in these regions to the solubility of fine particle Fe at Mt. Lushan, because the high solubility of Fe in the YRD and Anhui Province was just in accordance with the large amounts of anthropogenic emissions of SO₂ and NO_x in these regions (Zhang et al., 2009), which could significantly promote the dissolution of aerosol Fe acting as acid precursors (Schulz et al., 2012; Solmon et al., 2009). A significant correlation (0.629) of iron solubility and sulphate concentration in our fine particle samples was further verified, similar to the correspondence of high iron solubility to the sulphur content in ambient fine particles observed by (Oakes et al., 2012). Moreover, the outstanding enhancement of fine particle Fe solubility in local regions close to Mt. Lushan was principally recognised to be an effect of cloud processing, an important atmospheric process controlling Fe dissolution in aerosols (discussed in Sect. 3.4.4). The regional distributions for the solubility of crustal Al was similar with Fe, except for the lower PSCF values near Mt. Lushan (Fig. 8b).

Anhui, Jiangsu and Jiangxi Province were identified as the most important contributors to the high solubility of fine particle As at Mt. Lushan (Fig. 8c), ascribed to the large amount of atmospheric As emissions from anthropogenic sources (approximately 200 tons in 2010) in these provinces (Cheng et al., 2015); in contrast, little contributions were made to As solubility by the PRD and Hunan Province where the highest As con-

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



centrations were provided. For the elements from coal combustion (including Mn, Pb, Zn, Cu and Se, Fig. 8d–h) and MSW incineration (including Cd and Mo, Fig. 8i and j), the highest contributions of element solubility were uniformly focused within the local area of Jiangxi Province, the YRD and parts of Fujian Province, generally coinciding with their significantly high concentrations emitted from anthropogenic sources in eastern China. The solubility pattern for Ba (Fig. 8k) was rather similar to its concentration distribution, whereas the pattern of high solubility for Cr in Anhui Province (Fig. 8l) was very different to the concentration pattern.

It can be concluded that the solubility of fine particle trace elements from local and regional source regions were much higher than that by long-range transport from south-western China and northern China. The high contributions for particle element solubility in the YRD region should be attributed to the serious air pollution. However, on account of the relatively less air pollution released from local sources, the frequent cloud processing was considered as the significant contributor to the high solubility of trace elements in local regions near Mt. Lushan.

Additionally, the statistics on the solubility of individual fine particle elements at Mt. Lushan for five air mass categories are shown in Fig. S7. In general, the fine particle elements derived from cluster L exhibited the highest solubility whether for crustal or anthropogenic elements, closely followed by cluster E; elements from cluster SW showed moderate solubility, ascribed to the wash-out of soluble constituents by frequent rainfall in summer monsoon; elements from cluster NW and N were the least soluble, primarily impacted by crustal materials from northern China. It was demonstrated that the statistical results for most elements fairly agreed with the solubility distributions identified by PSCF analysis.

3.4.4 Evolution of element solubility during cloud processing

Cloud processing might be of great importance in affecting aerosol dissolution at Mt. Lushan where frequent cloud events occurred. To determine the evolution of element solubility during cloud processing with less disturbance of natural factors and

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



human activities, cloud events were selected with the following conditions: (1) non-raining cloud to avoid scavenging of particles by precipitation, (2) short duration and time interval of cloud processing and particle sampling to minimise the interference of external aerosols, (3) identical air mass trajectory of the cloud and particles with low wind speed to ensure consistent origins of the air masses. Table S4 shows the alteration of individual element solubility in fine particles before and after the two selected cloud events on 11 September 2011 and on 18 April 2012.

It was evident that after cloud processing on 11 September 2011, most fine particle elements exhibited a drastic increase in solubility. For instance, post-cloud particle Al became much more soluble compared with the pre-cloud particle, with solubility changing from 1.7 to 28.7 %. Other elements, such as Cu, Zn, As, Se, Cd and Ba, also showed an approximately one- to two-fold increase in solubility. The cloud processing on 18 April 2012 also brought about a significantly large number of solubility increments for the crustal elements Al and Fe, but it resulted in less increments for anthropogenic elements such as Cu, As, Se and Pb due to their already higher solubility in pre-cloud particles which might have undergone intensive cloud cycles during the prolonged residence time in the air. Generally, the evidently higher solubility for the post-cloud fine particle elements after the two cloud events, overpassing their overall average solubility, corroborated the remarkable dissolution efficiencies of aerosol elements by cloud processing at Mt. Lushan.

Interestingly, the increase in aerosol element solubility after both cloud events was coupled with the elevation of aerosol sulphate concentration as shown in Table S4. For example, post-cloud sulphate particles (4 h) exhibited a sharp elevation of $10 \mu\text{g m}^{-3}$ after the cloud processing (3 h) on 11 September 2011. Based on the very short intervals between cloud events and post-cloud particle sampling, the increased sulphate particles were most likely contributed by cloud water sulphate, which could be generated by the rapid atmospheric aqueous-phase oxidation of S(IV) with a rate greater than $100 \% \text{ h}^{-1}$ (Seinfeld and Pandis, 2012), much quicker than the gas-phase oxidation of SO_2 with a rate of approximately $1 \% \text{ h}^{-1}$ (Newman, 1981). In addition, the higher sul-

phate concentrations in cloud water and post-cloud particles on 11 September 2011 (41.70 mg L^{-1} and $24.72 \mu\text{g m}^{-3}$, respectively) from cluster E compared with those on 18 April 2012 (26.78 mg L^{-1} and $17.34 \mu\text{g m}^{-3}$, respectively) from cluster L were likely to be associated with the larger amount of anthropogenic SO_2 gases from the YRD in eastern China because acidic substances, such as sulphate formed by their precursor gases (e.g., SO_2), are prone to exist in fine particles and lead to acidification of aerosols (Ren et al., 2011); and the pH of cloud droplets modified by these dissolved acidic substances in particles could in turn influence the dissolution of aerosol trace elements (Deguillaume et al., 2005). As a consequence, the enhancement of trace element solubility during cloud processing were probably dependent on the corresponding heterogeneous formation of sulphate.

To further understand the effect of cloud processing on aerosol elements dissolution, individual cloud droplet residues were collected during the two cloud events, and the micro-morphology and composition of metal particles were examined by TEM-EDS analysis. Representative TEM images and the EDS spectra of major metal particles are shown in Fig. 9. Plenty of Pb-rich, Fe-rich and fly ash particles were observed to be the dominating metal particles in the two cloud events, consistent with the major types of metal-associated particles (Pb-rich, fly ash, Fe-rich and Zn-rich) in cloud droplets at Mt. Lushan (Li et al., 2013). Almost all of metal particles were found to be internally mixed within S-rich cloud residues with the encapsulation of cloud water and presented a nearly spherical shape or aggregation, suggesting the likely sources to be high-temperature coal combustion by coal-fired power plants and industries, during which industrial gases containing abundant SO_2 and metals were released, followed by the generation of metal-sulphate particles (Gieré et al., 2006).

Figure 9 also demonstrates the small diameters of the metal particles in cloud residues much smaller than $1 \mu\text{m}$, especially for the aggregation of spherical Fe nanoparticles embedded in aged S-rich residues (Fig. 9c). The unique, large specific surface of nano-sized Fe particles is able to effect a higher absorption intensity and more complicated surface chemistry during cloud processing, generating greater

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



quantities of surface complexes with acidic anions such as sulphate, and improving the dissolution rates of iron particles (Rubasinghege et al., 2010). Shi et al. (2009) suggested that the variations in pH during cloud processing would induce the formation of amorphous Fe nanoparticles and an increase in iron solubility (reactivity) in Saharan
dusts. More importantly, the solubilisation of aerosol elements caused by the change in particle morphology during cloud processing is irreversible (Deguillaume et al., 2005). Finally, along with the dissipating of cloud water in droplets, the Fe nanoparticles and sulphate residues remain in post-cloud particles, resulting in an attractive enhancement of Fe solubility. Therefore, we hypothesise that irreversible changes or alterations of particle morphology such as specific surface area, which may result from the acidification and heterogeneous reactions during cloud processing, should be the critical factor for the significant enhancement of solubility not only for Fe, but also for many other trace elements in cloud-processed fine particles.

4 Conclusions

The characteristics of fine particle trace elements at the summit of a mountain in southern China were investigated. The PM_{2.5} concentration ($55.2 \pm 20.1 \mu\text{g m}^{-3}$) was twice the WHO guideline but much lower than that in urban sites, with little seasonal variation. The total fractions of Al and Fe and the water-soluble fraction of Zn were the dominant elements and similar temporal variations between the total and water-soluble fractions for all species along with several spikes contributed by dust storms for Al and Fe were observed. The enrichment factor analysis implied the pollution trace elements to be more easily concentrated in water-soluble fractions with higher correlations. Nonferrous metal mining and smelting, coal combustion, crustal materials and municipal solid waste incineration were further classified as the major emission sources, which might initially determine the trace element solubility with interferences such as total concentrations. The YRD including Jiangsu, Zhejiang and parts of Anhui Province were indicated as the most important source regions of PM_{2.5} by trajectory cluster analysis and

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



PSCF results, and this result was discrepant with the source distributions for individual trace elements. The YRD as well as the PRD, eastern Hunan and western Jiangxi were identified as the primary source regions and pathways for combustion-related elements (Mn, Pb, Zn, Cu, Se, Cd and Mo) emitted by regional coal-fired power plants and municipal solid waste incineration, whereas the long-range transport of mineral dusts from northern China was the major contributor to crustal Al and Fe. Interestingly, local sources contributed relatively less to most trace elements. In contrast, the YRD region and the local area of Jiangxi Province were very likely to make the greatest contributions to element solubility rather than long-range transport from northern or southern China. A significant enhancement in trace element solubility and corresponding increase in aerosol sulphate were observed during two selected cloud events. TEM-EDS analysis of cloud residues suggested that the irreversible alteration of particle morphology by heterogeneous reactions or acidification with sulphate during cloud processing could crucially increase element solubility.

This study highlighted the contributions of regional anthropogenic pollution and local cloud processing to trace element solubility of PM_{2.5} at Mt. Lushan. We further expect that fine particles could undergo several cloud-processing cycles during long-range transport with massive clouds covering the surface of the earth, promoting trace element dissolution and increasing health and ecological risks.

The Supplement related to this article is available online at doi:10.5194/acpd-15-13001-2015-supplement.

Acknowledgements. This research was funded by the National Natural Science Foundation of China (21177073 and 41075092). We acknowledge the NOAA Air Resources Laboratory (ARL) for provision of the HYSPLIT trajectory model. We are grateful for all the staff of Lushan Mountain Meteorological Station for offering the observation platform and their assistance during the field campaigns. Thanks also due to Taixing Yue at Environmental Monitoring Central Station of Shandong Province for the help in the ICP-MS measurements.

References

- Baker, A. R., Jickells, T. D., Witt, M., and Linge, K. L.: Trends in the solubility of iron, aluminium, manganese and phosphorus in aerosol collected over the Atlantic Ocean, *Mar. Chem.*, 98, 43–58, doi:10.1016/j.marchem.2005.06.004, 2006.
- 5 Cakmak, S., Dales, R., Kauri, L. M., Mahmud, M., Van Ryswyk, K., Vanos, J., Liu, L., Kumarathasan, P., Thomson, E., Vincent, R., and Weichenthal, S.: Metal composition of fine particulate air pollution and acute changes in cardiorespiratory physiology, *Environ. Pollut.*, 189, 208–214, doi:10.1016/j.envpol.2014.03.004, 2014.
- Cao, J., Xu, H., Xu, Q., Chen, B., and Kan, H.: Fine particulate matter constituents and cardiopulmonary mortality in a heavily polluted Chinese city, *Environ. Health Persp.*, 120, 373–378, doi:10.1289/ehp.1103671, 2012.
- Charrier, J. G., McFall, A. S., Richards-Henderson, N. K., and Anastasio, C.: Hydrogen peroxide formation in a surrogate lung fluid by transition metals and quinones present in particulate matter, *Environ. Sci. Technol.*, 48, 7010–7017, doi:10.1021/es501011w, 2014.
- 15 Chen, J., Tan, M., Li, Y., Zheng, J., Zhang, Y., Shan, Z., Zhang, G., and Li, Y.: Characteristics of trace elements and lead isotope ratios in PM_{2.5} from four sites in Shanghai, *J. Hazard. Mater.*, 156, 36–43, doi:10.1016/j.jhazmat.2007.11.122, 2008.
- Chen, Y., Ebenstein, A., Greenstone, M., and Li, H.: Evidence on the impact of sustained exposure to air pollution on life expectancy from China's Huai River policy, *P. Natl. Acad. Sci. USA*, 110, 12936–12941, 2013.
- 20 Cheng, K., Wang, Y., Tian, H., Gao, X., Zhang, Y., Wu, X., Zhu, C., and Gao, J.: Atmospheric emission characteristics and control policies of five precedent-controlled toxic heavy metals from anthropogenic sources in China, *Environ. Sci. Technol.*, 49, 1206–1214, doi:10.1021/es5037332, 2015.
- 25 Cheung, K., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Diurnal trends in oxidative potential of coarse particulate matter in the Los Angeles Basin and their relation to sources and chemical composition, *Environ. Sci. Technol.*, 46, 3779–3787, doi:10.1021/es204211v, 2012.
- Clements, A. L., Buzcu-Guven, B., Fraser, M. P., Kulkarni, P., and Chellam, S.: Role of particulate metals in heterogenous secondary sulfate formation, *Atmos. Environ.*, 75, 233–240, doi:10.1016/j.atmosenv.2013.04.038, 2013.
- 30

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Costa, D. L. and Dreher, K. L.: Bioavailable transition metals in particulate matter mediate cardiopulmonary injury in healthy and compromised animal models, *Environ. Health Persp.*, 105, 1053–1060, 1997.

Deguillaume, L., Leriche, M., Monod, A., and Chaumerliac, N.: The role of transition metal ions on HO_x radicals in clouds: a numerical evaluation of its impact on multiphase chemistry, *Atmos. Chem. Phys.*, 4, 95–110, doi:10.5194/acp-4-95-2004, 2004.

Deguillaume, L., Leriche, M., Desboeufs, K., Mailhot, G., George, C., and Chaumerliac, N.: Transition metals in atmospheric liquid phases: sources, reactivity, and sensitive parameters, *Chem. Rev.*, 105, 3388–3431, doi:10.1021/cr040649c, 2005.

Deng, C., Zhuang, G., Huang, K., Li, J., Zhang, R., Wang, Q., Liu, T., Sun, Y., Guo, Z., Fu, J. S., and Wang, Z.: Chemical characterization of aerosols at the summit of Mountain Tai in Central East China, *Atmos. Chem. Phys.*, 11, 7319–7332, doi:10.5194/acp-11-7319-2011, 2011.

Desboeufs, K. V., Losno, R., and Colin, J. L.: Factors influencing aerosol solubility during cloud processes, *Atmos. Environ.*, 35, 3529–3537, doi:10.1016/S1352-2310(00)00472-6, 2001.

Draxler, R. R. and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY Website, available at: <http://www.arl.noaa.gov/HYSPLIT.php> (last access: 1 February 2015), NOAA Air Resources Laboratory, College Park, MD, 2014.

Fang, G. C., Huang, Y. L., and Huang, J. H.: Study of atmospheric metallic elements pollution in Asia during 2000–2007, *J. Hazard. Mater.*, 180, 115–121, doi:10.1016/j.jhazmat.2010.03.120, 2010.

Fomba, K. W., Müller, K., van Pinxteren, D., and Herrmann, H.: Aerosol size-resolved trace metal composition in remote northern tropical Atlantic marine environment: case study Cape Verde islands, *Atmos. Chem. Phys.*, 13, 4801–4814, doi:10.5194/acp-13-4801-2013, 2013.

Gieré, R., Blackford, M., and Smith, K.: TEM study of PM_{2.5} emitted from coal and tire combustion in a thermal power station, *Environ. Sci. Technol.*, 40, 6235–6240, 2006.

Guo, J., Tilgner, A., Yeung, C., Wang, Z., Louie, P. K., Luk, C. W., Xu, Z., Yuan, C., Gao, Y., Poon, S., Herrmann, H., Lee, S., Lam, K. S., and Wang, T.: Atmospheric peroxides in a polluted subtropical environment: seasonal variation, sources and sinks, and importance of heterogeneous processes, *Environ. Sci. Technol.*, 48, 1443–1450, doi:10.1021/es403229x, 2014a.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Guo, L., Chen, Y., Wang, F., Meng, X., Xu, Z., and Zhuang, G.: Effects of Asian dust on the atmospheric input of trace elements to the East China Sea, *Mar. Chem.*, 163, 19–27, doi:10.1016/j.marchem.2014.04.003, 2014b.
- Harris, E., Sinha, B., van Pinxteren, D., Tilgner, A., Fomba, K. W., Schneider, J., Roth, A., Gnauk, T., Fahlbusch, B., Mertes, S., Lee, T., Collett, J., Foley, S., Borrmann, S., Hoppe, P., and Herrmann, H.: Enhanced role of transition metal ion catalysis during in-cloud oxidation of SO₂, *Science*, 340, 727–730, doi:10.1126/science.1230911, 2013.
- Heal, M. R., Hibbs, L. R., Agius, R. M., and Beverland, I. J.: Total and water-soluble trace metal content of urban background PM₁₀, PM_{2.5} and black smoke in Edinburgh, UK, *Atmos. Environ.*, 39, 1417–1430, doi:10.1016/j.atmosenv.2004.11.026, 2005.
- Hoek, G., Krishnan, R. M., Beelen, R., Peters, A., Ostro, B., Brunekreef, B., and Kaufman, J. D.: Long-term air pollution exposure and cardio-respiratory mortality: a review, *Environ. Health Glob.*, 12, 43, doi:10.1186/1476-069X-12-43, 2013.
- Hopke, P., Barrie, L., Li, S. M., Cheng, M. D., Li, C., and Xie, Y.: Possible sources and preferred pathways for biogenic and non-sea-salt sulfur for the high Arctic, *J. Geophys. Res.-Atmos.*, 100, 16595–16603, 1995.
- Hsu, S.-C., Lin, F.-J., and Jeng, W.-L.: Seawater solubility of natural and anthropogenic metals within ambient aerosols collected from Taiwan coastal sites, *Atmos. Environ.*, 39, 3989–4001, doi:10.1016/j.atmosenv.2005.03.033, 2005.
- Hsu, S.-C., Wong, G. T. F., Gong, G.-C., Shiah, F.-K., Huang, Y.-T., Kao, S.-J., Tsai, F., Candice Lung, S.-C., Lin, F.-J., Lin, I. I., Hung, C.-C., and Tseng, C.-M.: Sources, solubility, and dry deposition of aerosol trace elements over the East China Sea, *Mar. Chem.*, 120, 116–127, doi:10.1016/j.marchem.2008.10.003, 2010.
- Hsu, S.-C., Lin, F.-J., Liu, T.-H., Lin, S.-H., Kao, S.-J., Tseng, C.-M., and Huang, C.-H.: Short time dissolution kinetics of refractory elements Fe, Al, and Ti in Asian outflow-impacted marine aerosols and implications, *Atmos. Environ.*, 79, 93–100, doi:10.1016/j.atmosenv.2013.06.037, 2013.
- Hu, X., Zhang, Y., Ding, Z., Wang, T., Lian, H., Sun, Y., and Wu, J.: Bioaccessibility and health risk of arsenic and heavy metals (Cd, Co, Cr, Cu, Ni, Pb, Zn and Mn) in TSP and PM_{2.5} in Nanjing, China, *Atmos. Environ.*, 57, 146–152, doi:10.1016/j.atmosenv.2012.04.056, 2012.
- Husain, L., Ghauri, B., Yang, K., Khan, A. R., and Rattigan, O. V.: Application of the SO₄²⁻/Se tracer technique to study SO₂ oxidation in cloud and fog on a time scale of minutes, *Chemosphere*, 54, 177–183, doi:10.1016/s0045-6535(03)00531-9, 2004.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jakob, A., Stucki, S., and Kuhn, P.: Evaporation of heavy metals during the heat treatment of municipal solid waste incinerator fly ash, *Environ. Sci. Technol.*, 29, 2429–2436, doi:10.1021/es00009a040, 1995.

Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, *Science*, 308, 67–71, doi:10.1126/science.1105959, 2005.

Kaufman, Y. J., Tanré, D., and Boucher, O.: A satellite view of aerosols in the climate system, *Nature*, 419, 215–223, 2002.

Li, W., Wang, Y., Collett Jr., J. L., Chen, J., Zhang, X., Wang, Z., and Wang, W.: Microscopic evaluation of trace metals in cloud droplets in an acid precipitation region, *Environ. Sci. Technol.*, 47, 4172–4180, doi:10.1021/es304779t, 2013.

Li, W., Chi, J., Shi, Z., Wang, X., Chen, B., Wang, Y., Li, T., Chen, J., Zhang, D., Wang, Z., Shi, C., Liu, L., and Wang, W.: Composition and hygroscopicity of aerosol particles at Mt. Lu in south China: implications for acid precipitation, *Atmos. Environ.*, 94, 626–636, doi:10.1016/j.atmosenv.2014.06.003, 2014.

Mackie, D. S., Peat, J. M., McTainsh, G. H., Boyd, P. W., and Hunter, K. A.: Soil abrasion and eolian dust production: implications for iron partitioning and solubility, *Geochem. Geophys.*, 7, 1–11, doi:10.1029/2006gc001404, 2006.

Mahowald, N.: Aerosol indirect effect on biogeochemical cycles and climate, *Science*, 334, 794–796, doi:10.1126/science.1207374, 2011.

Moreno, T., Querol, X., Alastuey, A., Reche, C., Cusack, M., Amato, F., Pandolfi, M., Pey, J., Richard, A., Prévôt, A. S. H., Furger, M., and Gibbons, W.: Variations in time and space of trace metal aerosol concentrations in urban areas and their surroundings, *Atmos. Chem. Phys.*, 11, 9415–9430, doi:10.5194/acp-11-9415-2011, 2011.

Newman, L.: Atmospheric oxidation of sulfur dioxide: a review as viewed from power plant and smelter plume studies, *Atmos. Environ.*, 15, 2231–2239, doi:10.1016/0004-6981(81)90255-9, 1981.

Oakes, M., Ingall, E. D., Lai, B., Shafer, M. M., Hays, M. D., Liu, Z. G., Russell, A. G., and Weber, R. J.: Iron solubility related to particle sulfur content in source emission and ambient fine particles, *Environ. Sci. Technol.*, 46, 6637–6644, doi:10.1021/es300701c, 2012.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Polissar, A. V., Hopke, P. K., and Harris, J. M.: Source regions for atmospheric aerosol measured at Barrow, Alaska, *Environ. Sci. Technol.*, 35, 4214–4226, 2001.
- Querol, X., Viana, M., Alastuey, A., Amato, F., Moreno, T., Castillo, S., Pey, J., de la Rosa, J., Sánchez de la Campa, A., Artíñano, B., Salvador, P., García Dos Santos, S., Fernández-Patier, R., Moreno-Grau, S., Negral, L., Minguillón, M. C., Monfort, E., Gil, J. I., Inza, A., Ortega, L. A., Santamaría, J. M., and Zabalza, J.: Source origin of trace elements in PM from regional background, urban and industrial sites of Spain, *Atmos. Environ.*, 41, 7219–7231, doi:10.1016/j.atmosenv.2007.05.022, 2007.
- Quispe, D., Pérez-López, R., Silva, L. F. O., and Nieto, J. M.: Changes in mobility of hazardous elements during coal combustion in Santa Catarina power plant (Brazil), *Fuel*, 94, 495–503, doi:10.1016/j.fuel.2011.09.034, 2012.
- Reff, A., Bhave, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D., and Houyoux, M.: Emissions inventory of PM_{2.5} trace elements across the United States, *Environ. Sci. Technol.*, 43, 5790–5796, doi:10.1021/es802930x, 2009.
- Ren, L., Wang, W., Wang, Q., Yang, X., and Tang, D.: Comparison and trend study on acidity and acidic buffering capacity of particulate matter in China, *Atmos. Environ.*, 45, 7503–7519, doi:10.1016/j.atmosenv.2010.08.055, 2011.
- Rubasinghege, G., Lentz, R. W., Scherer, M. M., and Grassian, V. H.: Simulated atmospheric processing of iron oxyhydroxide minerals at low pH: roles of particle size and acid anion in iron dissolution, *P. Natl. Acad. Sci. USA*, 107, 6628–6633, doi:10.1073/pnas.0910809107, 2010.
- Schulz, M., Prospero, J. M., Baker, A. R., Dentener, F., Ickes, L., Liss, P. S., Mahowald, N. M., Nickovic, S., Garcia-Pando, C. P., Rodriguez, S., Sarin, M., Tegen, I., and Duce, R. A.: Atmospheric transport and deposition of mineral dust to the ocean: implications for research needs, *Environ. Sci. Technol.*, 46, 10390–10404, doi:10.1021/es300073u, 2012.
- Schwab, J. J.: Aerosol chemical composition in New York state from integrated filter samples: urban/rural and seasonal contrasts, *J. Geophys. Res.*, 109, D16S05, doi:10.1029/2003jd004078, 2004.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: from Air Pollution to Climate Change*, John Wiley & Sons, New Jersey, 2012.
- Shafer, M. M., Perkins, D. A., Antkiewicz, D. S., Stone, E. A., Quraishi, T. A., and Schauer, J. J.: Reactive oxygen species activity and chemical speciation of size-fractionated atmospheric

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



particulate matter from Lahore, Pakistan: an important role for transition metals, *J. Environ. Monitor.*, 12, 704–715, 2010.

Shi, Z., Krom, M. D., Bonneville, S., Baker, A. R., Jickells, T. D., and Benning, L. G.: Formation of iron nanoparticles and increase in iron reactivity in mineral dust during simulated cloud processing, *Environ. Sci. Technol.*, 43, 6592–6596, 2009.

Solmon, F., Chuang, P. Y., Meskhidze, N., and Chen, Y.: Acidic processing of mineral dust iron by anthropogenic compounds over the north Pacific Ocean, *J. Geophys. Res.*, 114, D02305, doi:10.1029/2008jd010417, 2009.

Spokes, L. J., Jickells, T. D., and Lim, B.: Solubilisation of aerosol trace metals by cloud processing: a laboratory study, *Geochim. Cosmochim. Ac.*, 58, 3281–3287, doi:10.1016/0016-7037(94)90056-6, 1994.

Takahashi, Y., Higashi, M., Furukawa, T., and Mitsunobu, S.: Change of iron species and iron solubility in Asian dust during the long-range transport from western China to Japan, *Atmos. Chem. Phys.*, 11, 11237–11252, doi:10.5194/acp-11-11237-2011, 2011.

Tian, H., Gao, J., Lu, L., Zhao, D., Cheng, K., and Qiu, P.: Temporal trends and spatial variation characteristics of hazardous air pollutant emission inventory from municipal solid waste incineration in China, *Environ. Sci. Technol.*, 46, 10364–10371, doi:10.1021/es302343s, 2012.

Tian, H., Liu, K., Zhou, J., Lu, L., Hao, J., Qiu, P., Gao, J., Zhu, C., Wang, K., and Hua, S.: Atmospheric emission inventory of hazardous trace elements from China's coal-fired power plants-temporal trends and spatial variation characteristics, *Environ. Sci. Technol.*, 48, 3575–3582, doi:10.1021/es404730j, 2014.

Wang, Y., Zhang, X., and Draxler, R. R.: TrajStat: GIS-based software that uses various trajectory statistical analysis methods to identify potential sources from long-term air pollution measurement data, *Environ. Modell. Softw.*, 24, 938–939, 2009.

Wei, F., Chen, J., Wu, Y., and Zheng, C.: Study of the background contents of 61 elements of soils in China, *Environmental Science (in Chinese)*, 12, 12–19, 1991.

Weichenthal, S., Villeneuve, P. J., Burnett, R. T., van Donkelaar, A., Martin, R. V., Jones, R. R., DellaValle, C. T., Sandler, D. P., Ward, M. H., and Hoppin, J. A.: Long-term exposure to fine particulate matter: association with nonaccidental and cardiovascular mortality in the agricultural health study cohort, *Environ. Health Persp.*, 122, 609–615, doi:10.1289/ehp.1307277, 2014.

Wen, H. and Carignan, J.: Review on atmospheric selenium: emissions, speciation and fate, *Atmos. Environ.*, 41, 7151–7165, doi:10.1016/j.atmosenv.2007.07.035, 2007.

- Werner, M. L., Nico, P. S., Marcus, M. A., and Anastasio, C.: Use of micro-XANES to speciate chromium in airborne fine particles in the Sacramento Valley, Environ. Sci. Technol., 41, 4919–4924, doi:10.1021/es070430q, 2007.
- Xu, H. M., Cao, J. J., Ho, K. F., Ding, H., Han, Y. M., Wang, G. H., Chow, J. C., Watson, J. G., Khol, S. D., Qiang, J., and Li, W. T.: Lead concentrations in fine particulate matter after the phasing out of leaded gasoline in Xi'an, China, Atmos. Environ., 46, 217–224, doi:10.1016/j.atmosenv.2011.09.078, 2012.
- Yang, F., Tan, J., Zhao, Q., Du, Z., He, K., Ma, Y., Duan, F., Chen, G., and Zhao, Q.: Characteristics of PM_{2.5} speciation in representative megacities and across China, Atmos. Chem. Phys., 11, 5207–5219, doi:10.5194/acp-11-5207-2011, 2011.
- Yang, Y., Wang, Y., Wen, T., Li, W., Zhao, Y., and Li, L.: Elemental composition of PM_{2.5} and PM₁₀ at Mount Gongga in China during 2006, Atmos. Res., 93, 801–810, doi:10.1016/j.atmosres.2009.03.014, 2009a.
- Yang, Y., Wang, Y., Wen, T., Zhao, Y., and Li, J.: Element characteristics and sources of PM_{2.5} at Mount Dinghu in 2006, Environmental Science (in Chinese), 30, 988–992, 2009b.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, Atmos. Chem. Phys., 9, 5131–5153, doi:10.5194/acp-9-5131-2009, 2009.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Table 1. Concentrations (mean \pm SD) of PM_{2.5} ($\mu\text{g m}^{-3}$) and trace elements (ng m^{-3}) including total and water soluble fraction at Mt. Lushan.

	Overall		Summer, 2011		Spring, 2012	
	Total	Water soluble	Total	Water soluble	Total	Water soluble
PM _{2.5}	55.2 \pm 20.1		55.9 \pm 21.8		54.7 \pm 18.9	
Al	449.1 \pm 441.1	73.3 \pm 93.5	369.1 \pm 464.4	108.0 \pm 121.7	515.1 \pm 415.2	48.9 \pm 57.3
Fe	331.1 \pm 236.2	71.5 \pm 70.5	330.4 \pm 250.9	101.9 \pm 70.5	331.6 \pm 227.3	15.7 \pm 12.9
Zn	258.3 \pm 162.8	172.9 \pm 105.8	274.3 \pm 129.5	211.9 \pm 104.4	245.3 \pm 186.1	142.3 \pm 97.5
Pb	68.2 \pm 49.3	29.4 \pm 23.8	65.4 \pm 47.3	37.8 \pm 27.5	70.5 \pm 51.4	22.7 \pm 18.3
Ba	63.8 \pm 54.2	23.4 \pm 17.0	66.0 \pm 71.1	19.3 \pm 15.0	61.9 \pm 33.6	26.7 \pm 18.0
Mn	22.2 \pm 12.2	15.5 \pm 8.1	24.7 \pm 12.5	19.0 \pm 8.8	20.2 \pm 11.6	12.8 \pm 6.3
As	21.5 \pm 19.6	14.8 \pm 11.9	22.3 \pm 16.3	12.4 \pm 6.1	20.9 \pm 22.2	17.0 \pm 15.1
Cr	13.7 \pm 17.2	0.7 \pm 0.5	18.2 \pm 19.5	0.8 \pm 0.6	9.6 \pm 13.8	0.3 \pm 0.1
Cu	12.4 \pm 9.6	9.9 \pm 8.9	13.2 \pm 6.4	10.7 \pm 6.9	11.7 \pm 11.6	9.3 \pm 10.2
Se	7.0 \pm 3.3	5.0 \pm 2.9	7.6 \pm 2.9	5.2 \pm 2.9	6.5 \pm 3.5	4.8 \pm 2.9
Cd	2.5 \pm 1.8	1.3 \pm 0.9	2.8 \pm 1.9	1.3 \pm 0.6	2.2 \pm 1.6	1.2 \pm 1.1
Mo	2.0 \pm 2.0	0.6 \pm 0.4	2.6 \pm 2.2	0.7 \pm 0.5	1.5 \pm 1.7	0.5 \pm 0.3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.




Figure 1. The location of Mt. Lushan and adjacent Yangtze River and Poyang Lake.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 2. Temporal variations of individual trace element concentrations in $PM_{2.5}$ at Mt. Lushan. Boxes indicate the concentration spikes.

Concentrations and solubility of trace elements in fine particles

T. Li et al.




Figure 3. Enrichment factor (EF) values of fine particle trace elements at Mt. Lushan in total and water soluble fraction.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Loadings of (a) Component 1, (b) Component 2, (c) Component 3 and (d) Component 4 for trace elements in total fraction in $PM_{2.5}$.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 5. (a) Three-days mean trajectories arriving at Mt. Lushan, (b) averaged $PM_{2.5}$ concentration of five clusters, and (c) the likely source regions of $PM_{2.5}$ identified using PSCF plots during observation period.

Concentrations and solubility of trace elements in fine particles

T. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 6. Likely source regions of individual elements in $PM_{2.5}$ at Mt. Lushan, identified using PSCF plots. Note the criteria are 75th percentile for Al and mean concentration for others.

Concentrations and solubility of trace elements in fine particles

T. Li et al.




Figure 7. Scatter plot of aerosol element solubility vs. logarithmic EF value, computed with geometric mean in 99 % confidence interval. Colored shade areas represent different sources classified by EF value and PCA analysis.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Concentrations and solubility of trace elements in fine particles

T. Li et al.




Figure 8. Solubility distributions in likely source regions for individual elements in $PM_{2.5}$ at Mt. Lushan, identified by PSCF with the mean solubility values as criteria.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Concentrations and solubility of trace elements in fine particles

T. Li et al.




Figure 9. Typical TEM images and corresponding EDS spectra of metal particles embedded in individual S-rich cloud residues collected during cloud events. **(a)** Pb-S. **(b)** Fly ash-S. **(c)** Fe-S. The dotted red circles indicate the examined area of EDS. Cu* signals result from copper TEM grid.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)