

Interactive comment on “Modelled and observed changes in aerosols and surface solar radiation over Europe between 1960 and 2009” by S. T. Turnock et al.

S. T. Turnock et al.

Correspondence to: S. T. Turnock

(s.t.turnock12@leeds.ac.uk)

We would like to thank both of the anonymous reviewers for their helpful and constructive comments. Below we have responded to each comment in turn and made alterations to the manuscript where appropriate (shown enclosed in “*speech marks and italic font*”). The referee comments are shown first in **grey shading** and the author’s response is shown below in normal font.

Response to Anonymous Referee #1

General Comments:

1. The study focuses mostly on mass properties of atmospheric aerosol, but tells little about aerosol size distribution representation in the model. Aerosol size distributions have large effects on optical properties and aerosol-cloud interactions. In order to make the study stronger, I highly recommend including a figure or table comparing present day aerosol size distribution properties over the different European locations (or a few representative locations) as presented by Asmi et al., (2012). Asmi et al., (2012) presents a comprehensive analysis of present day aerosol size distributions in several locations across Europe and the data is freely available. The model results could help explain differences between modeled and observed AOD and shed some light on changes in CCN over Europe in the recent past. For example a simple comparison of the model ability to capture particle concentrations in the ranges 30-50nm, >50nm and >100nm, would complement the existing comparison. If the model represents well present day size distributions, changes over time in those parameters would be of special value to the community.

We have evaluated simulated aerosol number concentrations at sites across Europe using present day observational data from Asmi et al. (2011). The evaluation has used observational data for particles larger than 30 nm (N30), 50 nm (N50) and 100 nm (N100) dry diameter. An additional figure (see below for new Figure 11) shows the comparison between modelled and observed annual mean particle number concentrations at N30, N50 and N100 dry diameters across sites over Europe for the period 2008-2009. This figure shows that the model underpredicts particle number concentrations across all sizes over Europe by factor of 1.5-2.0. The model substantially underestimates the low particle number concentrations across all size ranges at the high latitude arctic sites of Zeppelin and Pallas. The model also underestimates larger size particles more than smaller sized ones, as indicated by the larger NMBF shown in the figure. These results are in accordance with the findings of the multi model comparison of global aerosol microphysical models to size distribution observational data undertaken by Mann et al. (2014). An underestimation of the aerosol size distribution across Europe could possibly account for some of the underestimation in simulated AOD and may also contribute to some of the discrepancies in the simulated surface solar radiation. However, factors other than aerosols might contribute to discrepancies between the modelled and observed AOD and surface solar radiation (e.g. humidity, clouds), which would require further work to investigate. We have modified the original manuscript to include a section and a new figure on the evaluation of the modelled present day aerosol size distributions. In addition, we have also included an amended Figure 4 showing the simulated change in annual mean particle number concentration across the different size fractions for Europe. This shows a large reduction (>50%) in aerosol number concentration over Europe across all sizes from 1970 to 2009. Taking into account the underestimation in present day aerosol number concentrations the simulated changes in aerosol number could be even larger, with potentially large impacts on climate.




Figure 11. Annual mean observed vs modelled aerosol number concentrations for 2008-2009 over the size fractions a) N30, b) N50 and c) N100

The following sections of the original manuscript have been modified to include details of the evaluation of simulated aerosol size distributions.

An additional sentence on Page 13458, line 6 has been modified.

“Here we compare the HadGEM3-UKCA coupled chemistry-climate model for the period 1960 to 2009 against extensive ground based observations of sulfate aerosol mass (1978–2009), total suspended particle matter (SPM, 1978–1998), PM₁₀ (1997–2009), aerosol optical depth (AOD, 2000–2009), aerosol size distributions (2008-2009) and surface solar radiation (SSR, 1960–2009) over Europe.”

An additional sentence on Page 13458, line 10 has been modified.

“The model underestimates observed sulfate aerosol mass (normalised mean bias factor (NMBF) = -0.4), SPM (NMBF = -0.9), PM₁₀ (NMBF = -0.2), aerosol number concentrations (N30 NMBF = -0.85), 50 nm (N50 NMBF = -0.65) and 100 nm (N100 NMBF = -0.96) and aerosol optical depth (AOD, NMBF = -0.01) but slightly overpredicts SSR (NMBF = 0.02).”

The sentence on Page 13463, line 2 has been modified.

“We evaluate the ability of the model to consistently capture observed changes in bulk in situ aerosol properties (PM, aerosol size distributions and chemical components) as well as radiative properties (AOD, SSR) over Europe.”

The sentence on Page 13467, line 24 has been modified.

“Ground-based measurements of aerosols used in this study are listed in Table 1 and include aerosol mass concentrations (sulfate and total mass) from the EMEP network (<http://www.emep.int>), AOD from AERONET (<http://aeronet.gsfc.nasa.gov/>), aerosol size distributions from EUSAAR (European

Supersites for Atmospheric Aerosol Research) and GUAN (German Ultrafine Aerosol Network) (Asmi et al., 2011) and SSR from the GEBA database (<http://www.geba.ethz.ch/history/index>).”

A new section has been added before section 2.2.2 on page 13468 to briefly mention the aerosol size distribution data.

“Section 2.2.2 Aerosol size distribution

Aerosol size distribution data over Europe from EUSAAR and GUAN ground based monitoring sites have previously been collected and processed by Asmi et al., (2011) over the period 2008-2009. Here we use the same 17 low altitude sites as in the multi model comparison by Mann et al., (2014) to compare modelled and observed aerosol number concentrations at three different size fractions (>30 nm (N30), >50 nm (N50) and >100 nm (N100) dry diameter) over the period 2008-2009.”

Figure 4a has been revised to include an additional panel showing the simulated change in aerosol number concentrations for the total, N30, N50 and N100 size fractions over the period of 1960 to 2009. The paragraph starting on Page 13471, line 9 has been modified to the following.

“Figure 4b shows the change in simulated European surface aerosol number concentrations over the total, N30, N50 and N100 size fractions between 1960 and 2009. Aerosol number concentrations across all size fractions increase from 1960 to a peak in 1970 before declining to concentrations that are 50% lower in 2009.

Simulated European AOD (Fig. 4c) increases from 1960 to a peak in 1973 before decreasing till 2009 to an AOD that is lower than that simulated 1960. Figure 4d shows that simulated European SSR decreases from 1960 to 1980 (“dimming”), then increases until 2009 (“brightening”).”

A new section has been inserted on Page 13476, line 1 before Section 3.2.3 Aerosol optical depth. Within this new section a new Figure 11 has also been included to show the model and observed aerosol number size distribution comparison (as shown above). The new section is as follows:

“Section 3.2.3 Aerosol size distribution

Figure 11a-c compares simulated and observed annual mean surface aerosol number concentrations averaged over the period 2008-2009 for particles of greater than 30 nm, 50 nm and 100nm in dry diameter. The model underestimates the observed aerosol number concentrations as all size fractions, with a slightly larger model bias in larger sized particles (N100 NMBF = -0.96, N50 NMBF = -0.65 and N30 NMBF = -0.85). The model substantially underestimates aerosol number concentrations over all size fractions at the high latitude Arctic monitoring locations of Pallas and Zeppelin. In addition, simulated aerosol number concentrations are underestimated to a larger extent at the more polluted, central European monitoring sites of Ispra, Preila, Bösels and K-Pusztas.

The model is able to reproduce the observed aerosol number concentrations within a factor of 2 at the majority of European monitoring locations, which is in agreement with the recent intercomparison and evaluation of global aerosol microphysical models (Mann et al., 2014). In addition, Mann et al., (2014) also found that all of the evaluated models underestimated aerosol number concentrations in the Arctic.

This suggests that N50 concentrations (a proxy for CCN concentrations) are slightly underpredicted by the model in the present day but have declined across Europe (Fig 4b) by 50% since the 1970s, with a large potential impact on the European radiative balance. Although over the more recent period of 2001-2010 no discernible trend was found in aerosol number concentration by Asmi et al., (2013) across three central European monitoring sites.”

A new sentence has been inserted on Page 13479, line 6.

“The model also underestimates aerosol number concentrations across the N30 (NMBF = -0.85), N50 (NMBF = -0.65) and N100 (NMBF = -0.96) size fractions in 2008-2009. Underestimation of surface sulfate, PM_{10} and aerosol number concentration is therefore not manifested in the simulation of AOD or SSR.”

The sentence on Page 13481, line 21 has been amended to:

“We evaluated the model against European observations of sulfate aerosol mass, total suspended particulate matter (SPM), PM₁₀ mass concentrations, aerosol number concentrations, aerosol optical depth (AOD) and surface solar radiation (SSR).”

A new sentence has been inserted on Page 13481, on line 27.

“In addition, the model underpredicts present day aerosol number concentrations over the size fractions of N30 (NMBF= -0.85), N50 (NMBF= -0.65) and N100 (NMBF= -0.96).”

Table 1 has also been modified to include the additional details as follows.

Table 1. Details of the ground based observations used in this study

Data Source	Measurements	Period	Total Number of Sites Available
EMEP	Total sulfate aerosol mass concentrations	1978-2009	97
EMEP	Total suspended particle mass concentrations	1978-2005	42
EMEP	PM ₁₀ Mass Concentrations	1996-2009	52
EUSAAR and GUAN	Aerosol number concentrations greater than 30 nm (N30), 50 nm (N50) and 100 nm (N100) in dry diameter	2008-2009	24
AERONET	Aerosol optical depth (AOD)	1994-2009	21
GEBA	Surface Solar Radiation (SSR)	1928-2009	50

2. Show, if possible, an additional plot similar to figure 12 of the seasonal SSR changes with time in addition to the annual values. It would be interesting to learn if the model captures the SSR seasonal cycle better than it does AOD and sulfate aerosol mass seasonal cycles, and the possible reasons why it does or does not.

We have calculated the seasonal mean statistics for SSR (see table below) and have amended a comment to the manuscript based on these results. The results show that as well as a small model bias in annual mean SSR (shown by small values of NMBF), the seasonal cycle in SSR is quite well reproduced by the model across all time periods (NMBF < 0.1). However, the model was previously shown to have larger seasonal biases in sulfate aerosol mass (DJF NMBF = -2.19 and JJA NMBF = 0.694), PM₁₀ (DJF NMBF = -0.12 and JJA NMBF = -0.339) and AOD (DJF NMBF = 0.258 and JJA NMBF = -0.167). This suggests that like for annual means, seasonal model biases in aerosol properties are not manifested in the model simulation of the seasonal cycle of SSR. The seasonal cycle in SSR is probably more strongly determined by variations in factors other than aerosols e.g. clouds.

We have included the following with the manuscript to briefly mention the seasonal cycle in SSR. A new sentence on page 13477, line 8.

“An evaluation of seasonal mean modelled and observed SSR also showed small differences in the NMBF over each season for all time periods, although RMSE is larger and the spatial correlation is smaller in summer.”

The sentence on page 13479, line 7 has been modified to:

“Underestimation of annual and seasonal mean surface sulfate and PM₁₀ is therefore not manifested in the simulation of the annual mean and seasonal cycle of AOD or SSR.”

Time Period	Season	Type	Absolute Change in SSR ($W m^{-2}$)	S.D. of mean value	% change in SSR	Calculated linear trend ($W m^{-2} yr^{-1}$)	-2 * S.E. trend	+ 2 * S.E. trend	r^2	NMBF	RMSE
1960-74	Annual	Observed	-0.79	2.40	-1	-0.011	-0.309	0.286	0.838	0.002	7.72
		Model + ARE	-1.67	1.89	-1	-0.088	-0.317	0.141			
		Model - ARE	-0.20	2.12	-0.2	0.069	-0.237	0.376	0.748	0.045	11.04
	DJF	Observed	-3.94	2.50	-10.6	-0.37	-0.601	-0.137	0.906	0.042	6.20
		Model + ARE	-2.60	2.51	-6.8	-0.20	-0.494	0.085			
		Model - ARE	-2.46	2.65	-6.4	-0.19	-0.500	0.124	0.847	0.108	7.45
	MAM	Observed	-0.73	4.76	-0.5	0.06	-0.527	0.650	0.552	-0.023	14.68
		Model + ARE	-1.53	4.06	-1.0	-0.03	-0.536	0.470			
		Model - ARE	-0.59	4.36	-0.4	0.06	-0.479	0.600	0.511	0.046	16.00
	JJA	Observed	2.40	5.94	1.2	0.39	-0.315	1.094	0.450	0.025	18.03
		Model + ARE	-3.32	4.02	-1.6	-0.13	-0.628	0.359			
		Model - ARE	0.48	4.73	0.2	0.26	-0.314	0.824	0.539	0.116	24.00
	SON	Observed	-0.61	3.50	-0.8	-0.09	-0.522	0.339	0.906	-0.023	7.93
		Model + ARE	1.50	2.59	2.1	0.11	-0.201	0.429			
		Model - ARE	1.77	2.97	2.4	0.15	-0.210	0.509	0.888	0.056	7.94
1975-89	Annual	Observed	-2.24	3.42	-2	-0.259	-0.658	0.140	0.803	0.029	8.43
		Model + ARE	3.12	3.06	3	0.304	-0.036	0.643			
		Model - ARE	1.89	3.50	1.6	0.187	-0.191	0.564	0.757	0.063	11.84
	DJF	Observed	-0.39	2.18	-1.1	-0.13	-0.393	0.127	0.905	0.099	6.92
		Model + ARE	1.06	1.87	2.8	0.03	-0.205	0.256			
		Model - ARE	0.79	1.96	2.1	0.00	-0.241	0.245	0.861	0.077	8.50
	MAM	Observed	-3.19	7.42	-2.1	-0.32	-1.221	0.584	0.485	0.000	14.00
		Model + ARE	2.87	5.95	1.9	0.30	-0.424	1.015			
		Model - ARE	1.87	6.06	1.2	0.22	-0.526	0.958	0.540	0.036	18.63
	JJA	Observed	-7.08	10.60	-3.5	-0.69	-1.947	0.569	0.450	0.046	20.03
		Model + ARE	5.06	9.70	2.5	0.56	-0.602	1.722			
		Model - ARE	2.01	10.73	-0.1	0.28	-1.041	1.603	0.637	0.102	30.40
	SON	Observed	1.71	3.31	9.5	0.10	-0.302	0.511	0.879	0.016	7.87
		Model + ARE	3.79	3.01	5.2	0.35	0.026	0.666			
		Model - ARE	2.90	3.02	2.6	0.25	-0.102	0.596	0.889	0.038	9.76
1990-09	Annual	Observed	5.78	3.47	5	0.369	0.153	0.584	0.896	0.043	8.00
		Model + ARE	3.98	3.04	3	0.316	0.125	0.507			
		Model - ARE	0.27	2.92	0.2	0.09	-0.185	0.366	0.828	0.053	10.23
	DJF	Observed	1.19	1.66	3.5	0.09	-0.037	0.214	0.930	0.078	5.91
		Model + ARE	0.27	2.01	0.7	0.02	-0.135	0.184			
		Model - ARE	-0.82	2.07	-2.2	-0.05	-0.211	0.116	0.890	0.030	7.31
	MAM	Observed	9.29	6.79	6.1	0.64	0.196	1.092	0.672	0.009	12.22
		Model + ARE	4.01	4.00	2.6	0.29	0.006	0.581			
		Model - ARE	0.25	4.29	0.2	0.06	-0.281	0.400	0.633	0.023	16.03
	JJA	Observed	6.07	7.70	3.0	0.36	-0.230	0.949	0.613	0.075	21.59
		Model + ARE	5.90	6.85	2.7	0.56	0.083	1.038			
		Model - ARE	-0.29	7.18	-0.1	0.18	-0.384	0.746	0.671	0.101	29.40
	SON	Observed	6.58	4.58	9.5	0.38	0.066	0.701	0.920	0.034	6.74
		Model + ARE	3.81	4.01	5.2	0.29	0.001	0.579			
		Model - ARE	1.92	4.16	2.6	0.17	-0.152	0.492	0.880	0.031	9.24

- Please mention in the text the mean global radiative forcing induced by changes in European aerosol emissions between periods with largest and smallest aerosol loadings. This value makes a direct comparison to GHG global forcing direct, and tells how important the regional aerosol perturbation is.

In our simulations, global emissions were changed. Without running further simulations it is not possible to obtain the global mean radiative forcing changes solely induced by European aerosol precursor emissions. We would need to run additional simulations over 50 years whereby the emissions from Europe are fixed to be able to isolate their effect on the globe, which was considered outside the scope of work for this paper. However, changes in global aerosol radiative forcing (RF) due to changes in global emissions can be calculated from our existing simulations. We have added a new Table to the manuscript (Table 6, see below) where we compare changes in global mean RF with European mean RF that we already reported. This table highlights the importance of regional changes in aerosol radiative forcing when compared to global changes and also the magnitude of changes compared to greenhouse gas forcing between 1960 and 2010. In addition, if only a global view is taken then important regional changes with opposite signs can be masked.

We have also added the following comment to the manuscript that relates to the Table below.

The sentence on Page 13480, line 1 has been changed to the following.

“Figure 15 shows the changes in simulated European mean top of atmosphere (TOA) outgoing shortwave radiation, relative to a 1980 to 2000 mean, under all-sky (a) and clear-sky conditions (b) with the numbers presented in Table 6.”

An additional sentence has been included on page 13481, line 9.

“Table 6 shows that the change in European mean aerosol RF is much larger than the change in global mean aerosol RF of $+0.4 \text{ W m}^{-2}$. Shindell et al. (2013) reported a multi-model evaluation of aerosol RF over the period 1850 to 2100, with 6 of the 9 models reporting a positive increase in global mean aerosol RF between 1980 and 2000, qualitatively similar to the results presented here. At the global scale, our simulated change in all-sky aerosol RF between 1970 and 2009 is ~40% of the magnitude of change in global CO_2 radiative forcing over the same period. At the European scale, the simulated change in all-sky aerosol RF is more than three times the change in global mean CO_2 RF. This indicates the large regional impact that decreasing aerosol concentrations have had on the radiative balance and climate over Europe compared to other forcing agents.”

Table 6. Global and European shortwave top of atmosphere all-sky and clear-sky aerosol radiative forcing, relative to a 1980 to 2000 mean. Values for 1972 are included as this is when the minimum aerosol radiative effect occurs over Europe. For comparison the global carbon dioxide radiative forcing (relative to 1750) from the IPCC fifth assessment report is shown in the last column (Myhre et al., 2013)

Year	All-Sky Aerosol RF (W m^{-2})		Clear-sky Aerosol RF (W m^{-2})		Estimates of CO_2 radiative forcing (Myhre et al., 2013)
	Global	Europe	Global	Europe	Global
1960	+ 0.9	+ 0.8	+ 0.4	+ 0.2	+ 0.7
1970	- 0.1	- 1.0	+ 0.01	- 0.1	+ 0.9
1972	- 0.1	- 1.4	+ 0.2	- 1.6	N/A
1980	- 0.4	- 0.6	+ 0.01	- 0.5	+ 1.1
1990	+ 0.01	+ 0.1	+ 0.04	- 0.1	+ 1.3
2000	+ 0.4	+ 1.7	- 0.1	+ 1.1	+ 1.5
2009	+ 0.3	+ 2.1	+ 0.1	+ 1.9	+ 1.8

Specific Comments:

Page 3(13459), line 26. How much have NO_x , CO and BC emissions decreased between 1980 and 2010?

We have changed line 26 on Page 3(13459) to include the percentage change in NO_x , CO and BC emissions between 1980 and 2010 and it reads as follow:

“Anthropogenic emissions of oxides of nitrogen (NO_x), carbon monoxide (CO) and black carbon (BC) have also decreased over Europe between 1980 and 2010 by 30%, 58% and 55% respectively (Granier et al., 2011).”

Page 13(13469), line 19. How was the interpolation made? Please add a short sentence explaining how this was done.

We have changed line 19 on Page 13(13469) to mention that the values from the four nearest model squares were used to linearly interpolate to the actual measurement location.

“Model values were linearly interpolated to each measurement site using the relative contribution from the four closest surrounding model grid squares.”

Response to Anonymous Referee #2

General Comments:

1. Discussion of the model evaluation is good, but could be presented more clearly. Comparison of modeled and observed 1978-2009 averages is a challenging task. Potentially significant details are lost in such a long-term average contains significant emission decreases. For example, according Figure 5, summer sulfate is underestimated early in the record is overestimated (quite significantly) before coming into closer agreement near the end of the record. Such details are likely lost in Figure 6, which very clearly depicts spatial biases, but how have those biases changed throughout the time period? This is important when assessing trends and impacts on SSR.

We use a combination of Figure 5 and 6 to show both spatial and temporal variability and trends in simulated and observed sulfate concentrations. Figure 5 shows the spatial distribution of annual mean sulfate concentrations in 1980, 1990, 2000 and 2009 as well temporal trends in annual mean, summertime and wintertime concentrations. Figure 6 shows summertime and wintertime NMBF at the European scale and also by region, with the range in NMBF reflecting variability in model bias over time. These show that wintertime sulfate is under estimated across all regions and for all of the period 1978 to 2009. Summertime sulfate is overestimated early in the record before coming into closer agreement towards the end of record, as indicated by the referee. Figure 6 shows that summertime concentrations are mostly underestimated in Northern and Southern Europe, but overestimated in central and eastern Europe. Combining Figure 5 and 6 together indicates that central European sites could be leading to the model overestimating summertime sulfate concentrations early in the record. The simulated trend in sulfate compared to that observed is dominated by discrepancies at these locations within different regions of Europe and over different time periods. In order to examine the reasons behind the changes in biases across the period 1978-2009 would require a more detailed investigation of simulated and observed summertime sulfate concentrations across monitoring sites within each region, which is outside the scope of the current work for this paper.

We have amended the lines 1-5 on Page 13472, line 1 to the following:

“In summer, observed sulfate is underestimated by the model at the start of the period before coming into closer agreement towards the end of the period (Figs., 5e, 6). The model underestimates observed summertime sulfate across northern and southern Europe (NMBF between 0 and -1), and overestimates sulfate in central and eastern Europe (NMBF <1). The model consistently underestimates wintertime sulfate (NMBF of -1 to -6) across all the European regions, with the largest discrepancy occurring in northern Europe.”

2. The importance of seasonality needs to be further discussed, including in SSR. Differing errors in the sulfate simulation exist summer and winter. This is important when considering differing insolation and hygroscopic growth that impact AOD and forcing.

See response to REVIEWER 1, general comment number 2 on this point.

3. The potential role of nitrate needs to be further highlighted. Nitrate likely plays an important role, particularly in winter, and might be able to resolve some of the issues mentioned in the manuscript, but also may exacerbate other problems in simulating AOD/SSR.

Nitrate is an important aerosol component particularly in northern Europe over winter and in more recent years as the concentration of SO₂ and sulfate aerosol has reduced. Future versions of HadGEM3-UKCA will include nitrate aerosol but this version currently does not and therefore we are unable to include its effects. However, due to its seasonal and temporal variability nitrate is only anticipated to make a difference to more recent comparisons over northern and western Europe in wintertime and could aid in the reduction of model observational bias here.

In Section 3.2.2 on page 13475, lines 4-12 we state that the contribution from nitrate aerosol has been shown to be about 1-3 μg m⁻³ over Europe and could therefore account for some of the discrepancy in the comparison of the simulated aerosol mass to that observed (Fagerli and Aas, 2008; Bellouin et al., 2011; Pozzer et al., 2012). It is anticipated to have a much smaller contribution

in 1980s due to the preference of ammonium sulfate aerosol formation over ammonium nitrate. Therefore the inclusion of nitrate within the model would not aid in reducing the model bias in aerosol mass over this period. Additionally the inclusion of nitrate aerosol could further exacerbate the model evaluation with AOD in winter and also potentially SSR (although as discussed above other factors e.g. clouds are likely to be dominant). This would likely occur in more recent years and over more northern and western areas of Europe where nitrate is likely to be a dominant component whereas, over southern Europe the model evaluation is less likely to be affected as sulfate remains the dominant aerosol component (Aan de burgh et al., 2011).

We have added additional statements on the seasonal, spatial and temporal importance of nitrate at the following points.

The following sentence has been inserted on Page 13475, line 6.

“This could be particularly important in reducing the model bias in winter and over northern and north west Europe where nitrate concentrations are anticipated to be largest.”

The following sentence has been inserted on Page 13475, line 12.

“Inclusion of nitrate aerosol would reduce the model bias in PM_{10} concentrations over the more recent years but have a smaller fractional impact in the 1980s-1990s due to the dominance of sulfate concentrations.”

4. In general, the figures in this manuscript are well done. However, many are tightly cramped into the ACPD structure. Please ensure that the figures are of adequate size to best display their fine detail.

We will endeavour to ensure that the figures are of adequate size and clarity during the type setting process for ACP.

Specific Comments:

Page 13464, Lines 12-13 – The meteorology is as close as the model can simulate, but not necessarily a match. This should be clarified.

We have changed the sentence on page 13464, lines 12-13 to read as follows:

“By nudging to re-analysis data, we ensure that the model produces a realistic representation of the meteorological conditions under which aerosol observations were taken.”

Page 13456, Lines 6-7 – This is potentially an important omission, particularly in northwestern continental Europe. Later in the manuscript the implications of this omission should be detailed.

See response to general comment 3.

Page 13466, Lines 1-10 – This is the standard procedure, but in this application it does not allow for segregation of the direct and first indirect effect.

As the reviewer has pointed out the double-call radiation configuration for all-sky and clear-sky effects does not allow for a direct separation of the direct and 1st indirect aerosol radiative effects. However, the radiative effects calculated under clear-skies are considered to represent the contribution from the direct aerosol radiative effect and thus an indication of the 1st indirect effect can be obtained by subtracting the radiative effects calculated under all-skies from those under clear-skies. From this calculation the change in cloud albedo effect over the period 1973-2009 has been stated on page 13480, line 20-24. Therefore we have included the following text within the methods section on page 13466, line 10:

“An estimate of the cloud albedo effect (first aerosol indirect) can be obtained by subtracting the radiative effect obtained under all-sky conditions from those under clear skies.”

Page 13466, Line 14 – “Time-averaged sporadic” is an oxymoron. Can this be rephrased? Shouldn't sporadic eruptions be allowed to occur at the appropriate place/time in the model to best reconstruct their impact?

The inventory of SO₂ emissions for explosively erupting volcanoes from the inventories by Andres and Kasgnoc (1998) and Halmer et al. (2002) does provide emissions from volcanoes at the appropriate location and date of occurrence. The sentence on page 13466, line 14 has been reworded to the following to improve clarity:

“Emissions from tropospheric volcanoes (both continuous and explosively erupting) are included from Andres and Kasgnoc (1998) and Halmer et al. (2002) using the AEROCOM recommendations.”

Page 13466, Lines 20-21 – Are the biogenic emissions not dependent on local meteorological conditions? Is this an acceptable approximation when looking at monthly mean values?

The inventory of Guenther et al. (1995) is used to provide monthly mean monoterpene emission fluxes here. This inventory has been derived based on a global model that simulates the flux of monoterpenes from natural sources, taking into factors such as light and temperature dependence. Therefore the fluxes used from this inventory are generated with local meteorological conditions taken into consideration.

Page 13467, Lines 13-15 – It is unclear what is meant here. Are ammonia emissions entirely omitted or just fixed to a constant value? This is an important detail.

Ammonia emissions are included within the model from the MACCity emissions inventory. Ammonia is used within the chemistry scheme and is available as an advected tracer for the aerosol scheme. However, due to the absence of ammonium nitrate within the aerosol scheme ammonia is not utilised by the aerosol scheme within HadGEM3-UKCA and it is assumed that sulfate aerosol is of the form ammonium sulfate.

For clarity we have amended page 13467 lines 13-15 to the following.

“The emissions of NH₃ across Europe (not used as part of the aerosol scheme in this study due to the absence of ammonium nitrate) increased continuously over the period 1960–2009, driven largely by the agriculture sector.”

Page 13469, Line 19 – How was the model data interpolated to each measurement site? A description of the method, even if just by name, or a reference is necessary since many model-observation studies simply compare the box average to measurements.

See response to reviewer 1, specific comment 2 for details on this point.

Page 13471, Line 15 – Are the values plotted in Figure 5 5-year averages or a single year?

The values in the contour plot on Figure 5 are single year annual mean concentrations for 1980, 1990, 2000 and 2009. We have slightly amended line 15 on page 13471 to reflect this as follows:

“Figure 5a–d compares simulated single year annual mean sulfate concentrations over Europe for 1980, 1990, 2000 and 2009 against observations.”

Page 13471, Line 24 – “slightly” should be omitted.

Deleted as requested.

Page 13472, Line 13-16 – Is wet deposition data available? Can it be added to the suite of observations used to validate the model?

Sulfate wet deposition data is available over Europe from EMEP but has not been processed and used as part of the current study.

Page 13476 – Is the discussion SSR clear-sky or all-sky? Is data available to evaluate COD?

The discussion is based on all-sky simulated and observed surface solar radiation data. Line 23, page 13476 has been amended below to reflect this:

“Figure 13 shows simulated and observed annual mean all-sky SSR anomalies across Europe between 1960 and 2009, relative to a 1980–2000 mean.”

The model evaluation of cloud optical depth was considered outside the scope of this paper and its feasibility has not been investigated further.

Page 134678, Lines 3-12 – Can this be used as a critique of the emission inventory?

The uncertainties in emissions inventory will certainly contribute to the errors in the modelled representation of SSR over the period 1960-1974. However, there are other additional factors e.g. clouds, humidity, observational errors contributing to the SSR over this period. Therefore it would be difficult to say with any confidence (and additional observations) that the uncertainties in the emissions inventories are causing the errors in the SSR.

Page 13481, Line 1-2 – The percentage change is not what matters, but the absolute magnitude.

The sentence on page 13481, line 1-2 has been amended to include the absolute change in sulfate concentrations from the two studies. This shows that there is still a larger decrease in sulfate concentrations over Europe than the USA, which reflects the difference in the changes in aerosol radiative effects between the two regions.

“The smaller changes in aerosol RF reported by Leibensperger et al. (2012) are possibly related to the smaller reductions in sulfate aerosol mass concentrations observed over the USA of $\sim 0.8 \mu\text{g S m}^{-3}$ (40 %), when compared to that observed over Europe of $1.2 \mu\text{g S m}^{-3}$ (70 %).”

Page 13501, Figure 3 caption – It is unclear what is meant by “within each particular network.” I think that phrase can be removed.

Agreed and has been removed as requested.

Page 13503, Figure 5 – The legend is not clear, particularly the observation (solid) and model (dotted) lines beneath the annual, DJF, and JJA lines. The S.D. estimates could be shown as colored rectangles rather than lines.

The legends on Figure 5 and Figure 8 have been amended to reflect the above comments.

References

Aan de Brugh, J. M. J., Schaap, M., Vignati, E., Dentener, F., Kahnert, M., Sofiev, M., Huijnen, V. and Krol, M. C.: The European aerosol budget in 2006, *Atmos. Chem. Phys.*, 11(3), 1117–1139, doi:10.5194/acp-11-1117-2011, 2011.

Andres, R. J. and Kasgnoc, A. D.: A time-averaged inventory of subaerial volcanic sulfur emissions, *J. Geophys. Res.*, 103(D19), 25251, doi:10.1029/98JD02091, 1998.

Asmi, A., Wiedensohler, A., Laj, P., Fjaeraa, A.-M., Sellegri, K., Birmili, W., Weingartner, E., Baltensperger, U., Zdimal, V., Zikova, N., Putaud, J.-P., Marinoni, A., Tunved, P., Hansson, H.-C., Fiebig, M., Kivekäs, N., Lihavainen, H., Asmi, E., Ulevicius, V., Aalto, P. P., Swietlicki, E., Kristensson, A., Mihalopoulos, N., Kalivitis, N., Kalapov, I., Kiss, G., de Leeuw, G., Henzing, B., Harrison, R. M., Beddows, D., O’Dowd, C., Jennings, S. G., Flentje, H., Weinhold, K., Meinhardt, F., Ries, L. and Kulmala, M.: Number size distributions and seasonality of

submicron particles in Europe 2008–2009, *Atmos. Chem. Phys.*, 11(11), 5505–5538, doi:10.5194/acp-11-5505-2011, 2011.

Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J. and Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, *J. Geophys. Res.*, 116(D20), D20206, doi:10.1029/2011JD016074, 2011.

Fagerli, H. and Aas, W.: Trends of nitrogen in air and precipitation: Model results and observations at EMEP sites in Europe, 1980–2003, *Environ. Pollut.*, 154(3), 448–461 [online] Available from: <http://www.sciencedirect.com/science/article/pii/S0269749108000523> (Accessed 9 December 2013), 2008.

Halmer, M. M., Schmincke, H.-U. and Graf, H.-F.: The annual volcanic gas input into the atmosphere, in particular into the stratosphere: a global data set for the past 100 years, *J. Volcanol. Geotherm. Res.*, 115(3-4), 511–528, doi:10.1016/S0377-0273(01)00318-3, 2002.

Mann, G. W., Carslaw, K. S., Reddington, C. L., Pringle, K. J., Schulz, M., Asmi, A., Spracklen, D. V., Ridley, D. A., Woodhouse, M. T., Lee, L. A., Zhang, K., Ghan, S. J., Easter, R. C., Liu, X., Stier, P., Lee, Y. H., Adams, P. J., Tost, H., Lelieveld, J., Bauer, S. E., Tsigaridis, K., van Noije, T. P. C., Strunk, A., Vignati, E., Bellouin, N., Dalvi, M., Johnson, C. E., Bergman, T., Kokkola, H., von Salzen, K., Yu, F., Luo, G., Petzold, A., Heintzenberg, J., Clarke, A., Ogren, J. A., Gras, J., Baltensperger, U., Kaminski, U., Jennings, S. G., O'Dowd, C. D., Harrison, R. M., Beddows, D. C. S., Kulmala, M., Viisanen, Y., Ulevicius, V., Mihalopoulos, N., Zdimal, V., Fiebig, M., Hansson, H.-C., Swietlicki, E. and Henzing, J. S.: Intercomparison and evaluation of global aerosol microphysical properties among AeroCom models of a range of complexity, *Atmos. Chem. Phys.*, 14(9), 4679–4713, doi:10.5194/acp-14-4679-2014, 2014.

Myhre, G., Shindell, D., Breon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T. and Zhang, H.: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.

Pozzer, A., de Meij, A., Pringle, K. J., Tost, H., Doering, U. M., van Aardenne, J. and Lelieveld, J.: Distributions and regional budgets of aerosols and their precursors simulated with the EMAC chemistry-climate model, *Atmos. Chem. Phys.*, 12(2), 961–987, doi:10.5194/acp-12-961-2012, 2012.

Shindell, D. T., Lamarque, J.-F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P. J., Lee, Y. H., Rotstayn, D., Mahowald, N., Milly, G., Faluvegi, G., Collins, W. J., Conley, A. J., Dalsoren, S., Easter, R., Ghan, S., Horowitz, L., Liu, X., Myhre, G., Nagashima, T., Naik, V., Rumbold, S. T., Skeie, R., Sudo, K., Szopa, S., Takemura, T., Voulgarakis, A., Yoon, J.-H. and Lo, F.: Radiative forcing in the ACCMIP historical and future climate simulations, *Atmos. Chem. Phys.*, 13, 2939–2974, doi:10.5194/acp-13-2939-2013, 2013.

Modelled and Observed Changes in Aerosols and Surface Solar Radiation over Europe between 1960 and 2009

**S. T. Turnock¹, D. V. Spracklen¹, K. S. Carslaw¹, G. W. Mann^{1,2}, M. T. Woodhouse^{1,*},
P. M. Forster¹, J. Haywood³, C. E. Johnson³, M. Dalvi³, N. Bellouin⁴, and
A. Sanchez-Lorenzo⁵**

¹Institute of Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, UK

²National Centre for Atmospheric Science, University of Leeds, UK

³Met Office, Fitzroy Road, Exeter, Devon, UK

⁴Department of Meteorology, University of Reading, Reading, UK

⁵Pyrenean Institute of Ecology, Spanish National Research Council, Zaragoza, Spain

*Now at CSIRO Ocean and Atmosphere, Aspendale, Victoria, Australia

Correspondence to: Steven Turnock
(s.t.turnock12@leeds.ac.uk)

Abstract

Substantial changes in anthropogenic aerosols and precursor gas emissions have occurred over recent decades due to the implementation of air pollution control legislation and economic growth. The response of atmospheric aerosols to these changes and the impact on climate are poorly constrained, particularly in studies using detailed aerosol chemistry climate models. Here we compare the HadGEM3-UKCA coupled chemistry-climate model for the period 1960 to 2009 against extensive ground based observations of sulfate aerosol mass (1978-2009), total suspended particle matter (SPM, 1978-1998), PM₁₀ (1997-2009), aerosol optical depth (AOD, 2000-2009), [aerosol size distributions \(2008-2009\)](#) and surface solar radiation (SSR, 1960-2009) over Europe. The model underestimates observed sulfate aerosol mass (normalised mean bias factor (NMBF) = -0.4), SPM (NMBF = -0.9), PM₁₀ (NMBF = -0.2) ~~and~~, [aerosol number concentrations \(N30 NMBF = -0.85, N50 NMBF = -0.65 and N100 NMBF = -0.96\)](#) and aerosol optical depth (AOD, NMBF = -0.01) but slightly overpredicts SSR (NMBF = 0.02). Trends in aerosol over the observational period are well simulated by the model, with observed (simulated) changes in sulfate of -68% (-78%), SPM of -42% (-20%), PM₁₀ of -9% (-8%) and AOD of -11% (-14%). Discrepancies in the magnitude of simulated aerosol mass do not affect the ability of the model to reproduce the observed SSR trends. The positive change in observed European SSR (5%) during 1990-2009 ('brightening') is better reproduced by the model when aerosol radiative effects (ARE) are included (3%), compared to simulations where ARE are excluded (0.2%). The simulated top-of-the-atmosphere aerosol radiative forcing over Europe under all-sky conditions increased by ~~3~~ [3.1](#) W m⁻² during the period 1970-2009 in response to changes in anthropogenic emissions and aerosol concentrations.

1 Introduction

Aerosols can cause acid deposition, degradation of atmospheric visibility, changes to the Earth's radiative balance, and are also a major source of air pollution, which affects human

health. Aerosols interact with climate by absorbing and reflecting incoming solar radiation and by modifying the microphysical properties of clouds. These effects have been defined in the latest Intergovernmental Panel on Climate Change (IPCC) report (Boucher et al., 2013) as aerosol-radiation interactions (ARI) and aerosol-cloud interactions (ACI). Aerosols (also referred to as Particulate Matter (PM)) are detrimental to air quality and human health, as particles below a certain size can penetrate into the lungs causing respiratory and cardiovascular problems (COMEAP, 2010). Strategies that attempt to mitigate climate change and poor air quality are inherently connected and have the potential to induce both benefits and penalties for either depending on the particular species targeted (Arneth et al., 2009; Ramanathan and Feng, 2009; Fiore et al., 2012).

Here we use a global coupled chemistry-climate model to improve our understanding of changes in aerosols over Europe from 1960 to 2009. The climate impact of aerosols over this period, in response to emission changes, was calculated as an aerosol radiative forcing. An assessment of the confidence in this effect was obtained from the ability of the model to reproduce observed long term changes in a number of aerosol properties including mass concentrations and aerosol optical depth.

Anthropogenic emissions of aerosol particles and their precursors have increased significantly since pre-industrial times. For example, global SO₂ emissions have increased by a factor of 60 from the pre-industrial to a peak in the 1970s (Lamarque et al., 2010; Granier et al., 2011; Smith et al., 2011). However, from the 1980s onwards regional reductions in anthropogenic emissions (mainly North America and Europe) have occurred due to air quality mitigation strategies, which has led to a decline in European SO₂ emissions of 73% between 1980 and 2004 (Vestreng et al., 2007; Hand et al., 2012). ~~European emissions of other anthropogenic species such as~~ Anthropogenic emissions of oxides of nitrogen (NO_x), carbon monoxide (CO) and black carbon (BC) have also decreased over Europe between 1980 and 2010 by 30%, 58% and 55% respectively (Granier et al., 2011). Conversely, anthropogenic emissions from East Asia have increased due to population growth and economic development, leading to an increase in SO₂ emissions of a factor of seven from the 1960s to the present day (Smith et al., 2011).

Changes in anthropogenic emissions and aerosol concentrations affect the Earth's climate (Arneth et al., 2009; Ramanathan and Feng, 2009; Fiore et al., 2012). The effect of past and future changes in emissions on aerosols and their associated climate impacts is uncertain (Penner et al., 2010; Chalmers et al., 2012). In addition, emission inventories of aerosols and their precursor species account for large uncertainty in models (de Meij et al., 2006). It is therefore important to understand and evaluate the changes to aerosol processes and properties that have occurred over recent decades where we have aerosol measurements.

Ground-based monitoring networks providing observations of aerosol concentrations and physio-chemical properties were established following air pollution control legislation. The longest continuous measurements of aerosols are available in North America and Europe from the 1970s to present day. In Europe, observations of aerosol mass concentrations (both sulfate and total) are available from the European Monitoring and Evaluation Programme (EMEP) network (Tørseth et al., 2012) and similar data for North America are available from the Integrated Monitoring of Protected Visual Environments (IMPROVE) network (<http://vista.cira.colostate.edu/improve/>). In addition, Aerosol Optical Depth (AOD) has been monitored in Europe over the last decade by the ground-based Aerosol Robotic Network (AERONET) (Holben et al., 1998). There are limitations in the spatial and temporal extent of the data from these networks, as well as the consistency of the instrumental techniques used and components measured throughout the monitoring period. However they do provide a useful source of multi decadal aerosol data with which to evaluate model predictions. Several studies have analysed long-term trends in observed aerosols. Tørseth et al. (2012) used observations from the EMEP network to show a decline in sulfate aerosol mass from the 1970s to present day and a decline in PM_{10} (mass of particles of diameter $< 10\mu\text{m}$) from 2000 to 2009. Asmi et al. (2013) and Barmpadimos et al. (2012) reported a reduction in the aerosol number and mass concentration over the last 10 to 15 years across Europe. However, Collaud Coen et al. (2013) found no significant changes in aerosol optical properties over Europe for a similar period. Similarly Harrison et al. (2008) analysed changes in aerosol mass concentrations over the UK and reported relatively stable concentrations over

2000 to 2010, even when emission reductions are anticipated to have occurred.

Evaluating the ability of chemistry-climate models to reproduce observed trends is necessary in order to reliably predict the climate effects of aerosols over this period. Many studies qualitatively match the direction of observed trends in aerosol but underestimate both absolute concentrations and the magnitude of observed trends (Berglen et al., 2007; Colette et al., 2011; Koch et al., 2011; Chin et al., 2014). Leibensperger et al. (2012) used the Chemical Transport Model (CTM) GEOS-Chem to evaluate aerosol trends over the USA at decadal time slices and found that sulfate but not BC was represented well by the model. A multi-model assessment of aerosol trends in Europe over the last decade showed models successfully simulate observed negative trends in PM₁₀ but fail to reproduce the positive trends observed at some locations and typically underestimate absolute concentrations (Colette et al., 2011). Observed reductions in sulfate over Europe have also been underestimated by other studies (Berglen et al., 2007; Koch et al., 2011). Over Europe, Skeie et al. (2011) reproduced the observed change in decadal average sulfate concentrations but overestimated nitrate aerosol concentrations.

Several studies (Lamarque et al., 2010; Shindell et al., 2013) have also assessed long term changes in AOD. Lamarque et al. (2010) evaluated simulations of present day AOD against AERONET observations and reported a relatively good reproduction of inter-annual variability, except in regions of high AOD. Shindell et al. (2013) assessed AOD from the models involved in the Atmospheric Chemistry and Climate Model Inter comparison Project (AC-CMIP) and showed that most captured the observed trends between 1980 to 2010 relatively well, although many under predicted in the present day, particularly over East Asia.

Changes in surface solar radiation (SSR) do not provide a direct measurement of aerosols but they can be used to infer their influence on the surface radiation balance. The Global Energy Balance Archive (GEBA) provides long term observations of SSR from the 1950s until present day over a large part of Europe (Sanchez-Lorenzo et al., 2013). These measurements can also be used as a general measure to validate radiation balance predictions from global climate models against the observed surface variations. Such observations have shown that the European 'dimming' period of the 1980s and the subsequent 'brightening'

period of the 1990s-2000s can be attributed partly to changes in clouds, concentrations of aerosols and aerosol-cloud effects (Wild, 2009). Model simulations have been evaluated against SSR observations and demonstrated issues in simulating the timing and magnitude of observed SSR trends. In an assessment of the models contributing to the 5th Coupled Model Intercomparison Project (CMIP5), Allen et al. (2013) showed that the dimming trend over Europe was underestimated in all models, potentially due to an under-represented aerosol direct effect. However, the CMIP5 models were able to reproduce the observed brightening trend. Folini and Wild (2011) performed transient simulations with ECHAM5-HAM over Europe (with interactive aerosols) and found that simulated reductions in SSR occurred too early in the model whereas, the increase in SSR was correctly timed. Using a regional climate model over Europe driven by reanalysis data, Chiacchio et al. (2015) overestimated SSR, simulated a premature onset of dimming and demonstrated that only simulations including aerosols were able to reproduce the observed brightening trend. In addition, Koch et al. (2011) simulated the correct inter-decadal variability in SSR but underestimated the magnitude of observed SSR.

These previous studies have either simplified their treatment of aerosols or the model was evaluated against a limited range of aerosol properties over a relatively coarse spatial and temporal scale. The importance of studies at regional spatial scales was also highlighted as changes in aerosols at this level can potentially be masked by compensatory changes observed on the global scale. Here we simulate monthly-mean aerosol concentrations from 1960 to 2009 using the HadGEM3-UKCA global chemistry climate model, which includes aerosol microphysics (aerosol number and mass size distributions). We evaluate the ability of the model to consistently capture observed changes in bulk in situ aerosol properties (PM, [aerosol size distributions](#) and chemical components) as well as radiative properties (AOD, SSR) over Europe. We also calculate the regional top of atmosphere radiative perturbations due to simulated changes in aerosols. This has enabled a detailed regional analysis and evaluation of the changing radiative impact of aerosols due to variations in emissions. Section 2 describes the HadGEM3-UKCA model, the simulations performed and the long term observations used. Section 3 discusses and evaluates the simulated changes to Euro-

pean aerosols and surface solar radiation. Section 4 presents aerosol radiative forcing over Europe. Conclusions are presented in Sect. 5.

2 Methods

2.1 Model Description and Simulations

2.1.1 General

We used the coupled chemistry-climate model HadGEM3-UKCA to study the interaction between chemistry, aerosols and the impacts on the radiation balance of the climate system. HadGEM3-UKCA is part of the third generation of the Met Office's Hadley Centre Global Environment Model (HadGEM) family, which incorporates an online treatment of chemistry and aerosols through the United Kingdom Chemistry and Aerosols (UKCA) programme. The Met Office Unified Model (UM) acts as the dynamical core and provides the components for atmospheric transport and tracer mixing. This is based on the dynamics implemented by Davies et al. (2005) and includes processes such as large-scale advection, convective uplift and boundary layer mixing. A description of the atmosphere-only version of HadGEM3 is provided in Hewitt et al. (2011). Morgenstern et al. (2009) and O'Connor et al. (2014) describe the incorporation and evaluation of UKCA into HadGEM.

HadGEM3-UKCA is used here in atmosphere-only mode. We output monthly 3-D aerosol and radiation fields for the years 1960-2009 at a resolution of $1.875^\circ \times 1.25^\circ$ (approximately 140 km at mid latitudes) with 63 vertical levels up to 40 km. The 3-D meteorological fields were nudged at 6-hourly intervals to the European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis (ERA-40) (Uppala et al., 2005) for the years 1960 to 2000 and ERA-Interim (Dee et al., 2011) for 2000 to 2009. Sea surface temperatures and sea ice fields were prescribed in accordance with those used in CMIP5 (Hurrell et al., 2008). Coupling between the land surface and atmosphere was simulated using the Met Office Surface Exchange Scheme (MOSES; Essery et al. (2002)). By nudging to re-analysis data,

we ensure that the ~~meteorological conditions in the model match those model produces a~~ realistic representation of the meteorological conditions under which aerosol observations were taken.

5 Tropospheric ozone- HO_X - NO_X -VOC chemistry was calculated using the mechanism described by O'Connor et al. (2014), which includes reactions of odd oxygen (O_X), nitrogen (NO_Y), hydrogen ($\text{HO}_X = \text{OH} + \text{HO}_2$) and CO, as well as methane and other short chain non-methane volatile organic compounds (VOCs). The scheme has been extended to include additional sulfur (Mann et al., 2010), monoterpene (Spracklen et al., 2006) and isoprene (Scott et al., 2014) chemistry.

10 The Fast-J photolysis scheme is implemented within UKCA to calculate online photolysis rates based on the distribution of clouds, ozone and aerosols (Wild et al., 2000). Dry and wet deposition of gas phase species is described in O'Connor et al. (2014).

HadGEM3-UKCA uses the modal aerosol scheme of the Global Model of Aerosol Processes (GLOMAP-mode) (Mann et al., 2010). GLOMAP-mode uses log-normal modes to represent the aerosol size distribution and simulates the evolution of the size-resolved number and mass of aerosol particles with different compositions. GLOMAP simulates the interaction of various aerosol processes including size-resolved primary emissions, cloud processing, new particle formation, hygroscopic growth, coagulation, condensation, deposition and scavenging. Log normal modes are used to represent aerosols in the nucleation (diameter (D) < 10 nm), Aitken (D 10 – 100 nm), accumulation (D 100 nm - 1 μm) and coarse (D > 1 μm) modes. In this study the model is set up to simulate sulfate, BC, organic carbon (OC) and sea salt aerosol in 5 different modes (4 soluble and 1 insoluble Aitken modes). Secondary organic aerosol (SOA) is formed from products of monoterpene oxidation, which generated at 13% yield and assumed to be involatile (Spracklen et al., 2006). There is no representation of ammonium nitrate in this version of the model. Mineral dust is simulated using a separate 6-bin scheme developed by Woodward (2001) and covers particle sizes from 0.03 to 30 μm in radius.

15
20
25

2.1.2 Aerosol Radiative Effects

The Edwards-Slingo radiation code (Edwards and Slingo, 1996) calculates changes in the Earth's radiation balance from chemical and aerosol species. ARI (aerosol direct effects) are calculated according to Bellouin et al. (2013) from waveband-averaged scattering and absorption coefficients obtained from modelled size distributions and a volume-weighted average of component refractive indices within each mode. GLOMAP-mode provides the aerosol fields for the calculation of ACI on-line within the model, in accordance with that described in Bellouin et al. (2013). Simulated cloud condensation nuclei (CCN) concentrations are used to calculate cloud droplet number (CDN) concentrations based on the empirical relationship derived by Jones et al. (2001). The cloud albedo effect is calculated using simulated CDN. The coupling of UKCA to the precipitation scheme potentially allows for the effect of aerosols on cloud lifetime (rapid adjustments from ACI) to be diagnosed. This can be done by calculating the influence they have on CDN concentrations, cloud effective radius and ultimately the auto-conversion of cloud drops to precipitation, as described in Jones et al. (2001). However, in this study, because we nudge to reanalysis fields, the changes in the radiative balance due to aerosols are not allowed to feedback on to the meteorology.

The radiation scheme was called twice each time step (every 30 minutes) in a double-call radiation configuration (Bellouin et al., 2013). The first call uses the modelled aerosol fields and the second call uses an aerosol-free atmosphere for the clear-sky radiative calculation and a prescribed default CDN field (based on an aerosol climatology) for the cloud radiation calculation. This approach therefore eliminates the model's radiatively-driven response to changes in aerosols, enabling the radiative forcing from ARI and ACI to be calculated with no rapid adjustments to clouds permitted from either effect. Here we calculate an aerosol radiative forcing (only from the direct and first indirect effects) as the difference between monthly values in each year of the simulation and the monthly mean aerosol radiative state in the period 1980-2000. [An estimate of the cloud albedo effect \(first aerosol indirect\) can](#)

be obtained by subtracting the radiative effect obtained under all-sky conditions from those under clear skies.

2.1.3 Aerosol Emissions

Sea salt emissions are calculated using the surface wind speed and the parameterisation of Gong (2003). Emissions from tropospheric volcanoes (both continuous and ~~time-averaged sporadic explosively erupting~~) are included from ~~Andres and Kasgnoc (1998) and Halmer et al. (2002)~~ Andres and Kasgnoc (1998) and Halmer et al. (2002) using the AEROCOM recommendations. Dimethyl sulfide (DMS) emissions are calculated from monthly sea-water concentration fields (Kettle and Andreae, 2000) and a wind speed dependent air-sea exchange parameterisation (Liss and Merlivat, 1986). Wildfires (biomass burning) emissions are from the MACCity inventory (Granier et al., 2011) for 1980 to 2009 and from the RETRO inventory from 1960 to 1980 (Schultz et al., 2008). Monoterpene emissions from vegetation are prescribed as monthly mean fields from Guenther et al. (1995). Dust emissions from the surface are calculated on-line using a 6-bin scheme (Woodward, 2001) and depend on the particle size distribution of the soil, surface soil type (represented by a bare soil fraction), soil moisture and wind speed. Dust is emitted when the wind speed exceeds a threshold value and if the soil moisture is below a certain level. Monthly mean anthropogenic emissions of CO, SO₂, NO_x, OC and BC from 1960-2009 are taken from the MACCity inventory (Granier et al., 2011). Emissions are provided at a 0.5° by 0.5° resolution and are based on those from ACCMIP and the IPCC Reference Concentration Pathway (RCP) scenario 8.5 for energy, transportation, industry, shipping, agriculture, residential and waste sectors.

Figure 1 shows the European and regional domains used throughout this study. Figure 2 shows the emissions of SO₂, OC, BC, NH₃ and NO_x across Europe from the MACCity inventory between 1960 and 2009. Anthropogenic emissions of SO₂ over the European domain have increased from 33 Tg/yr in 1960 to a peak of 46 Tg/yr in 1980 before decreasing at a relatively constant rate to 11 Tg/yr in 2009 (Fig. 2a). Between 1980 and 2009, European anthropogenic SO₂ emissions in this dataset declined by 70%, in agreement

with previous assessments (Vestreng et al., 2007; Tørseth et al., 2012). The emissions of NO_x have decreased by 20% and followed a similar temporal trend to SO_2 . A continuous decline in BC and OC emissions occurred from the 1960s to present day, due to reductions in the residential sector, partially offset by recent increases from the transportation sector.

5 The emissions of NH_3 across Europe (not used as part of the aerosol scheme in this study due to the absence of ammonium nitrate) increased continuously over the period 1960-2009, driven largely by the agriculture sector.

Figure 2b to f shows the annual emissions of each species from the MACCity inventory across the individual European regions. Emissions of OC (Fig. 2f) and SO_2 (Fig. 2b) have decreased across all the different European regions. Emissions of NO_x and BC (Fig. 2c and e) increase until the 1980s-1990s before declining across all regions, except in southern Europe. Anthropogenic emissions from northern and central Europe have declined from the 1980s onwards, whereas emissions from southern Europe have either increased or remained unchanged.

15 2.2 Observations

Ground-based measurements of aerosols used in this study are listed in Table 1 and include aerosol mass concentrations (sulfate and total mass) from the EMEP network (<http://www.emep.int>), AOD from AERONET (<http://aeronet.gsfc.nasa.gov/>) ~~and~~, aerosol size distributions from EUSAAR (European Supersites for Atmospheric Aerosol Research) and GUAN (German Ultrafine Aerosol Network) (Asmi et al., 2011) and SSR from the GEBA database (<http://www.geba.ethz.ch/history/index>). All data used in this study is presented as either monthly or annual averages. Figure 1 shows the location of all the measurement sites along with the 5 regions of Europe. Figure 3 shows the temporal evolution in the number of measurement sites used in this analysis for each observation. We note that

20

25 the number of locations reporting sulfate and SSR has declined since 2000.

2.2.1 EMEP Observations

The EMEP network has reported the concentrations of sulfate and total aerosol mass at locations across Europe from 1978 until present day (Tørseth et al., 2012). Measurements of total Suspended Particle Matter (SPM) are an early measure of particulate matter and available from 1978 to 2005, with most measurements from Germany, Switzerland and Spain. However, only measurements up to and including 1998 are used due to the reduced availability of data in the period 1999-2005 (Fig. 3). The SPM measurements cover all particle sizes and may be influenced by local sources of very large particles (diameter $> 10\mu\text{m}$). Measurements of PM_{10} are available from EMEP from 1996 until present day. Sulfate aerosol mass measurements are available from 1978 until present day.

We used sulfate aerosol mass, PM_{10} and total SPM from the sites that have been continuously operating for more than 5 years (Fig. 1a). The measurements were made using different measurement techniques and time frequencies (hourly and daily). The raw data were screened to remove any anomalous data points according to the flag in the original data records. Monthly and annual mean values were then calculated from the screened data for sites that had more than 75% of measurements in the averaging period.

2.2.2 Aerosol size distribution

Aerosol size distribution data over Europe from the EUSAAR and GUAN ground based monitoring sites have previously been collected and processed by Asmi et al. (2011) over the period 2008-2009. Here we use the same 17 low altitude sites as in the multi model comparison by Mann et al. (2014) to compare modelled and observed aerosol number concentrations at three different size fractions ($>30\text{ nm}$ (N30), $>50\text{ nm}$ (N50) and $>100\text{ nm}$ (N100) dry diameter) over the period 2008-2009.

2.2.3 AOD

The AERONET program is a ground-based network of sun photometers, currently with more than 200 sites providing aerosol optical, microphysical and radiative properties

(Holben et al., 1998). Observed AOD values are available over Europe from the mid-1990s but most sites only started operating within the last ten years (Fig. 3) and there are relatively few consistent long term datasets available before 2000. We used the Level 2.0 data product (cloud-screened and quality assured) from 20 sites that have been operating for longer than 5 years over Europe between 2000 and 2009. AOD measurements have the best record for wavelengths of 440, 500 and 675 nm. We used the 440 nm wavelength as it has the best spatial and temporal coverage.

2.2.4 Surface Solar Radiation

GEBA contains worldwide measurements of energy fluxes at the surface from more than 2000 sites, with the highest density over Europe. Monthly mean values of incident SSR (expressed as mean irradiance, in W m^{-2}) have been obtained from 56 sites across Europe starting before the 1970s, provided by Sanchez-Lorenzo et al. (2013), and including more than 20 sites that have data in the 1960s. The length of this observational record enabled the model evaluation to be extended prior to the availability of sulfate data. These measurements therefore enable an indirect evaluation of aerosol changes in the model across the entire time period of the simulations and also validation of how changes in aerosols can affect the Earth's radiation balance.

2.3 Model Evaluation Metrics

Comparisons were made using monthly and annual mean values at individual monitoring locations and also across Europe as a whole. Model values were [linearly](#) interpolated to each measurement site [using the relative contribution from the four closest surrounding model grid squares](#). The absolute and percentage change in the simulated and observed values of sulfate, SPM, PM_{10} , AOD and SSR were calculated as the difference between the mean of the initial 5 years and most recent 5 years of data.

The temporal trend in simulated and observed data was calculated by fitting an ordinary

least squares linear model to the data using the function below:

$$Y_i = a + bX_i (i = 1, \dots, n) \quad (1)$$

The standard error (SE) of the trend line was used to provide an assessment of the error. For each simulated and observed trend, +/- two SE in the gradient was applied to provide an uncertainty range. Firstly the standard deviation of the residuals (σ) was determined and then used to calculate the SE.

The simulated temporal trends were evaluated by comparing against observed trends; if the gradient of the simulated and observed trends are within +/- two SE of each other we considered them to be similar.

An assessment of model accuracy is provided here by calculating the normalised mean bias factor (NMBF) of the model when compared to the observations (Yu et al., 2006). This metric is symmetric (i.e. not biased towards under prediction or over prediction) and is not biased when a low number of observed values are used. It is defined as:

$$NMBF = S [\exp(|\ln(\bar{M}/\bar{O})|) - 1], \quad (2)$$

Where

$$S = (\bar{M} - \bar{O}) / |\bar{M} - \bar{O}|. \quad (3)$$

where M represents the model values and O represents the observed values. The sign of the NMBF indicates whether the model underestimates (negative) or overestimates (positive) the observed values. For a negative NMBF, the model values are a factor of $(1 - NMBF)$ below the observed values and for a positive NMBF the model values are a factor of $(1 + NMBF)$ above the observations (Yu et al., 2006). That is, $NMBF = -0.5$ means the model is a factor 1.5 low biased and $NMBF = 0.5$ means the model is a factor 1.5 high biased.

The goodness of fit between the model and observations is obtained by calculating the square of the linear Pearson correlation coefficient. A measure of the difference between model and observational values is provided by calculating the Root Mean Square Error (RMSE).

3 Results

3.1 Simulated European Aerosols and Surface Solar Radiation 1960 to 2009

Figure 4a shows the simulated European (land only) mass concentrations of the different aerosol components over the period 1960-2009. Simulated PM₁₀ declines from the early 1980s until 2009, coinciding with the reduction in anthropogenic emissions (Fig. 2). Continually decreasing mass concentrations of BC and OC are simulated between 1960 and 2009. Simulated sulfate mass concentrations increase from 1960 to a peak around 1980 and decrease thereafter.

Figure 4b shows ~~that simulated European SSR decreases~~ the change in simulated European surface aerosol number concentrations over the total N30, N50 and N100 size fractions between 1960 and 2009. Aerosol number concentrations across all size fractions increase from 1960 to 1980 ('dimming'), then increases until 2009 ('brightening'). a peak in 1970 before declining to concentrations that are 50% lower in 2009.

Simulated European AOD (Fig. 4bc) increases from 1960 to a peak in 1973 before decreasing till 2009 to an AOD that is lower than that simulated 1960. Figure 4d shows that simulated European SSR decreases from 1960 to 1980 ('dimming'), then increases until 2009 ('brightening').

3.2 Model Evaluation

3.2.1 Sulfate Aerosol Mass

Figures 5a-d compare simulated single year annual mean sulfate concentrations over Europe ~~in for~~ 1980, 1990, 2000 and 2009 against observations. Observed and simulated annual mean European sulfate aerosol concentrations declined by 68% and 78% respectively over the period 1978-2009 (Table 2). In 1980 (Fig. 5a) simulated and observed sulfate concentrations were largest in central Europe. In 2000 and 2009 (Fig. 5c-d) both simulated and observed sulfate concentrations in Central Europe have declined substantially, with the largest simulated concentrations over south east Europe.

The model underestimates European annual mean sulfate aerosol mass concentrations in the period 1978-2009 (NMBF = -0.384, Table 2), with summertime concentrations ~~slightly~~ overestimated (NMBF = 0.694) and wintertime concentrations underestimated (NMBF = -2.19). Figure 6 shows the range of model bias in seasonal sulfate across European regions for the period 1978-2009. In summer, ~~the model underestimates observed~~ sulfate is underestimated by the model at the start of the period before coming into closer agreement towards the end of the period (Figs. 5e,6). The model underestimates observed summertime sulfate across northern and southern Europe (NMBF between 0 and -1), and overestimates sulfate in central and eastern Europe (NMBF <1). The model consistently underestimates wintertime sulfate (NMBF of -1 to -6) across all the European regions, with the largest discrepancy occurring in northern Europe.

An underprediction of wintertime European sulfate concentrations has been previously reported and may be due to an underestimation of oxidants in the model (Berglen et al., 2007; Manktelow et al., 2007; Langmann et al., 2008). In wintertime over northern Europe, the region with largest model bias, low concentrations of H₂O₂ limit in-cloud aqueous phase oxidation of SO₂. Under these conditions oxidation by ozone is the dominant sulfate formation mechanism (Kreidenweis, 2003). We hypothesise that oxidation by ozone could be under-represented in the model, resulting in an underestimation of wintertime sulfate. This will be explored further in a future publication. Model underestimation could also be due to an artificially high wet deposition rate of sulfate, caused by an enhanced occurrence of drizzle precipitation within this version of the model (Walters et al., 2011).

Although the model underestimates absolute sulfate concentrations, the simulated trend over the period 1978-2009 ($-0.04 \pm 0.002 \mu\text{g S m}^{-3} \text{ yr}^{-1}$) is in good agreement with the observed trend ($-0.05 \pm 0.002 \mu\text{g S m}^{-3} \text{ yr}^{-1}$), at least on a European-wide annual mean basis (Fig. 5e). These trends can be considered similar as they are within two standard errors of each other. The largest decline in both simulated and observed sulfate concentrations occurred during 1980-2000, when average concentrations changed by $-0.05 \mu\text{g S m}^{-3} \text{ yr}^{-1}$. Between 1980 and 2000, average simulated and observed concentrations declined by 50-60%, corresponding with a 60-70% decrease in anthropogenic emissions of

SO₂ (Fig. 2). A smaller reduction in sulfate aerosol mass of 13-18% was simulated and observed in the period 2000-2009, when average concentrations changed by $-0.015 \mu\text{g S m}^{-3} \text{ yr}^{-1}$. Figure 7 compares the simulated and observed linear trends in sulfate concentrations at all measurement locations. The model reproduces the observed linear trends in the annual mean concentrations. However, linear trends in wintertime sulfate aerosol mass are underestimated (by a factor of 3) but overestimated in summertime (by a factor of 1.3). This corresponds with the calculated NMBFs for winter and summertime mass concentrations, which showed an under and over prediction respectively.

3.2.2 Total Aerosol Mass

Figures 8a and b compare simulated and observed annual mean total SPM concentrations over Europe in 1980 and 1990. The spatial changes in SPM over this period are less distinct than that for sulfate. Simulated and observed SPM decreases over central Europe and increases over eastern Europe between 1980 and 1990. In contrast to sulfate aerosol mass, the simulated 20% decrease in SPM mass concentrations in the period 1978-1998 is considerably lower than the 50% observed decrease.

The model underpredicts the observed European annual mean SPM mass concentrations, with a NMBF of -0.88 in the period 1978-1998 (Table 3). A consistent underprediction of observed SPM concentrations was modelled across all European regions in both summer and winter (Fig. 9a). The model substantially underpredicts SPM concentrations in wintertime for southern and eastern Europe (NMBF of -8 to -0.5). In summertime and across all European regions the model underpredicts observations by a smaller amount (NMBF of 0 to -2). The large underprediction of observed SPM mass concentrations could indicate that an additional emission source or process for generating supermicron aerosol mass is missing from the model, particular in the 1980s and early 1990s when the model bias is largest.

Figure 8c shows that the larger observed trend in European annual mean SPM mass over the period 1978-1998 of $-1.19 \pm 0.22 \mu\text{g m}^{-3} \text{ yr}^{-1}$ is substantially different to the simulated trend of $-0.26 \pm 0.12 \mu\text{g m}^{-3} \text{ yr}^{-1}$ (Table 3). Figure 10a shows that the observed trends

in SPM mass concentrations are underpredicted at all the measurement locations, with little seasonal differences. The calculated trends in simulated and observed SPM values are considered to be different as they are outside the range of +/- two standard errors of each trend line (Table 3).

5 Figures 8d and e compare simulated and observed annual mean PM₁₀ mass concentrations over Europe in 2000 and 2009. A slight reduction in PM₁₀ mass concentrations of 8-9% was both observed and simulated over this period (Table 3), with the largest reductions occurring over central and north-eastern continental Europe.

The model generally underestimates observed PM₁₀ mass concentrations (NMBF of 0 to -1) for the majority of European regions and across most of the evaluated years (Fig. 9b). An exception occurs across northern and north-western Europe in wintertime where the model overpredicts concentrations (NMBF of 0 to 2). This is potentially caused by an overestimation of sea salt aerosol (as also seen in studies by Mann et al. (2010, 2012)), so mostly affects coastal locations. Overall the model simulates European PM₁₀ mass concentrations between 1997 and 2009 within a factor of 2 and is much improved when compared to the simulation of total SPM between 1978 and 1998.

The temporal changes in PM₁₀ mass concentrations (Fig. 8f) highlight the smaller difference between simulated and observed PM₁₀ across Europe compared to SPM. The linear trends for observed ($-0.27 \pm 0.24 \mu\text{g m}^{-3} \text{ yr}^{-1}$) and simulated ($-0.14 \pm 0.16 \mu\text{g m}^{-3} \text{ yr}^{-1}$) PM₁₀ mass concentrations over the period 1997-2009 are similar (gradients within twice the standard error of each other) (Table 3). However, Fig. 10b shows that the magnitude of the observed downward trends is slightly underpredicted at the majority of measurement locations, with little difference between summer and winter.

The model underpredicts SPM mass concentrations by up to $20 \mu\text{g m}^{-3}$ in the 1980s and PM₁₀ by less than $5 \mu\text{g m}^{-3}$ in the 2000s. This larger underprediction in the 1980s could be due to errors in the measurement of SPM, as most of these observations are not well documented (Tørseth et al., 2012) and may have substantial uncertainty. We compared SPM and PM₁₀ observations during a period when both variables were observed at 6 monitoring sites in Spain, and found SPM was greater than PM₁₀ by 6-17 $\mu\text{g m}^{-3}$. Taking this

into account, along with the better model agreement for PM₁₀ mass, indicates that coarse sized particles ($D > 10 \mu\text{m}$) are under-represented by the model. Potential anthropogenic sources of coarse particles that are not represented in the model include road traffic dust and construction sources.

5 Underprediction of aerosol mass could be due to underestimation of aerosol sources, as well as missing aerosol sources from the model. The model does not include nitrate aerosol which could account for $1\text{-}3 \mu\text{g m}^{-3}$ of aerosol mass over Europe (Fagerli and Aas, 2008; Bellouin et al., 2011; Pozzer et al., 2012). This could be particularly important in reducing the model bias in winter and over northern and north west Europe where nitrate concentrations are anticipated to be largest.

10 The reductions in SO₂ emissions and increase in NH₃ emissions across Europe over the last 30 years (Fig. 2) could have important impacts on aerosol composition (Fagerli and Aas, 2008). In historical periods with high SO₂ emissions, sulfate aerosol will dominate and nitrate concentrations are likely to be small. In the recent past and future, declines in SO₂ and sulfate aerosol mass, coupled with an increase in NH₃ emissions may lead to increased nitrate aerosol concentrations. Inclusion of nitrate aerosol would reduce the model bias in PM₁₀ concentrations over more recent years but have a smaller fractional impact in the 1980s-1990s due to the dominance of sulfate concentrations.

15 The model also doesn't include primary biological aerosol sources, which may contribute $1\text{-}2 \mu\text{g m}^{-3}$ to PM₁₀ mass over Europe (Heald and Spracklen, 2009); the contribution to $D > 10 \mu\text{m}$ is not known.

20 Uncertainty in aerosol precursor emissions will also contribute to the model-observation discrepancy. In particular, domestic wood burning and wild fires could contribute up to 50% of OC locally over Europe and may be underestimated in emission datasets (Hodzic et al., 2007; Langmann et al., 2008; Manders et al., 2012). The calculation of SOA is also a large

25 uncertainty, particularly the proportions from anthropogenic and biogenic sources. Global aerosol models typically underpredict the amount of organic aerosols in the atmosphere (Tsigaridis et al., 2014), particularly from anthropogenic sources (Volkamer et al., 2006; Farina et al., 2010; Spracklen et al., 2011b). Anthropogenic sources (or anthropogenically modified biogenic sources) that are not accounted for here may contribute up to $3 \mu\text{g m}^{-3}$

of SOA over Europe (~~Spracklen et al., 2011a~~)([Spracklen et al., 2011b](#)).

Nevertheless, even using the upper estimates of some of these potential missing sources, there still appears to be a model underprediction of total aerosol mass particularly during the early period (1980-1990), suggesting that additional sources or processes are missing within the model or that removal processes are overestimated.

3.2.3 [Aerosol Size Distribution](#)

[Figure 11a-c compares simulated and observed annual mean surface aerosol number concentrations averaged over the period 2008-2009 for particles of greater than 30 nm, 50 nm and 100 nm dry diameter. The model underestimates the observed aerosol number concentrations as all size fractions, with a slightly larger model bias in larger sized particles \(N100 NMBF = -0.96, N50 NMBF = -0.65 and N30 NMBF = -0.85\). The model substantially underestimates aerosol number concentrations over all size fractions at the high latitude Arctic monitoring locations of Pallas and Zeppelin. In addition, simulated aerosol number concentrations are underestimated to a larger extent at the more polluted, central European monitoring sites of Ispra, Preila, Bösel and K-Puszta.](#)

[The model is able to reproduce the observed aerosol number concentrations within a factor of 2 at the majority of European monitoring locations, which is in agreement with the recent intercomparison and evaluation of global aerosol microphysical models \(Mann et al., 2014\). In addition, Mann et al. \(2014\) also found that all of the evaluated models underestimated aerosol number concentrations in the Arctic.](#)

[This suggests that N50 concentrations \(a proxy for CCN concentrations\) are slightly underpredicted by the model in the present day but have declined across Europe \(Fig. 4b\) by 50% since the 1970s, with a large potential impact on the European radiative balance. Although over the more recent period of 2001-2010 no discernible trend was found in aerosol number concentration by Asmi et al. \(2013\) across three central European monitoring sites.](#)

3.2.4 Aerosol Optical Depth

Figure 12a-b compares simulated and observed annual mean AOD at 440 nm in 2000 and 2009. The largest simulated and observed AOD occurs over eastern and south-eastern Europe. The model is relatively unbiased against annual mean AOD (NMBF= -0.013). The model captures the observed seasonal cycle in European AOD with highest AOD in the summer and lowest in the winter, but overestimates wintertime AOD (NMBF = 0.258) and underestimates summertime AOD (NMBF = -0.167) (Table 4). These seasonal biases are of opposite sign to those for sulfate and PM₁₀. However, we note that simulation of AOD requires information on aerosol optics, aerosol size distribution and atmospheric humidity meaning it is difficult to relate to comparisons of surface aerosol mass.

Observed and simulated AOD has declined over the period 2000 to 2009 (Fig. 12c). Table 4 shows that at the three monitoring locations with the longest data records (9-10 years), simulated and observed AOD has declined by a similar magnitude (11-14%) but AOD is underestimated by the model (NMBF = -0.296). The observed AOD trend at the three long term sites in the period 2000-2009 is $-0.007 \pm 0.004 \text{ yr}^{-1}$ and is similar to that modelled of $-0.006 \pm 0.0008 \text{ yr}^{-1}$ (Table 4). The trend in observed wintertime AOD ($-0.006 \pm 0.002 \text{ yr}^{-1}$) is well captured by the model ($-0.007 \pm 0.002 \text{ yr}^{-1}$). The larger observed summertime trend of $-0.014 \pm 0.003 \text{ yr}^{-1}$ is underestimated ($-0.005 \pm 0.002 \text{ yr}^{-1}$). The ability of the model to reproduce the decline in AOD is similar to that for sulfate and PM₁₀.

3.2.5 Surface Solar Radiation

Figure 13 shows simulated and observed annual mean [all-sky](#) SSR anomalies across Europe between 1960 and 2009, relative to a 1980-2000 mean. The long term mean was based on the period 1980-2000, considered to be the period with the most reliable observations. Evaluation of SSR has been split into three distinct time periods (1960-1974, 1975-1989 and 1990-2009) based on the changes in the observed all-sky SSR values (Table 5). The observed SSR anomaly is generally positive and relatively constant in the period 1960-1974. There is a decrease in observed SSR from 1975 until the late 1980s, after which

a strong increase in SSR is observed from 1990 to 2009.

Between 1990 and 2009 both the observed and simulated European SSR increases by 5.8 and 4.0 W m^{-2} respectively, with similar positive linear trends (0.37 to 0.32 $\text{W m}^{-2} \text{yr}^{-1}$ - Table 5). The highest spatial correlation ($r^2 = 0.90$) between modelled and observed SSR values occurs in the period 1990-2009, whereas the bias (NMBF = 0.04) and error (RMSE = 8.0) in SSR are similar to that in 1975-1989. [An evaluation of seasonal mean modelled and observed SSR also showed small differences in the NMBF over each season for all time periods, although RMSE is larger and the spatial correlation is smaller in summer.](#) The positive SSR trend ('brightening') observed between the late 1980s and 2009 is reproduced by the model, but simulated brightening begins several years earlier than observed. The positive trend in SSR anomaly across Europe from the mid-1980s to present day corresponds with the observed and simulated decrease in aerosol concentrations. Figure 13 also shows the modelled all-sky SSR anomalies without aerosol radiative effects (ARE). Without ARE the simulated trend in SSR is underestimated ($0.09 \pm 0.14 \text{ W m}^{-2} \text{yr}^{-1}$), suggesting that changes in aerosol concentrations are a dominant driver of SSR trends during this period. The simulated positive trend in SSR including ARE presented here is in agreement with other studies over this period (Wild, 2009; Allen et al., 2013; Sanchez-Lorenzo et al., 2013; Chiacchio et al., 2015).

In the period 1975-1989 both the modelled and observed SSR anomalies are generally negative, which coincides with the maximum anthropogenic emissions and atmospheric aerosol loading. Over this period the observed SSR decreases by an average of 2.2 W m^{-2} , whilst simulated SSR increases by 3.1 W m^{-2} (Table 5). A similar discrepancy in sign and magnitude is also apparent in the linear trends of the model ($0.30 \pm 0.17 \text{ W m}^{-2} \text{yr}^{-1}$) and observations ($-0.26 \pm 0.20 \text{ W m}^{-2} \text{yr}^{-1}$). This reflects the models inability to simulate the timing and magnitude of the observed dimming trend in SSR values between 1975 and 1989 (Fig. 13). Over this period the simulated and observed SSR values have a lower correlation ($r^2 = 0.80$) and larger error (RMSE = 8.43) than the 1990-2009 period, indicating a slightly poorer representation by the model. The model without ARE also shows similar disagreements (Table 5), potentially indicating that errors in the simulation of aerosol are

not causing simulated discrepancy in SSR during this period.

Over the period 1960-1974 simulated European annual mean SSR remained relatively constant (Fig. 13) and is of a similar magnitude (Table 5) to that observed. Over this period the observed SSR anomalies are positive (compared to a 1980-2000 mean), whilst the modelled SSR anomalies are negative. The small observed trend of $-0.01 \pm 0.15 \text{ W m}^{-2} \text{ yr}^{-1}$ over the period 1960-1974 is overestimated by the model ($-0.09 \pm 0.11 \text{ W m}^{-2} \text{ yr}^{-1}$). The stronger simulated negative trend in SSR between 1960 and 1974 indicates that the dimming observed in the period 1974-1989 occurs too early in the model. The model without ARE does not show a dimming trend over this period (but does have a large uncertainty, Table 5), which implies that the discrepancy in the SSR trend could be due to uncertainties in simulated aerosols.

Understanding the discrepancy in simulated SSR prior to 1980 is difficult because aerosol observations are not available. Possible causes of model discrepancy include errors in simulated aerosol, problems with the observations, or the ECMWF reanalysis product. With regards to observational uncertainties, there were fewer observations of SSR before 1970 (Fig. 3) and there is also a larger correction factor associated with data from the available sites (Sanchez-Lorenzo et al., 2013). This suggests that observational error may be larger in the early period. Prior to 2000, the model is forced by the ERA-40 reanalysis. ERA-40 was improved in the 1970s by the inclusion of additional measurements, most notably from satellites (Uppala et al., 2005). Larger errors in the reanalyses prior to 1980 (Uppala et al., 2005) could cause errors in the generation of clouds by the host GCM, which would affect simulated SSR.

3.2.6 Evaluation Summary

Figure 14 summarises the comparison between simulated and observed sulfate, SPM, PM₁₀, AOD and SSR across Europe, separately for their entire operational period and also for the period 2000-2009 (when PM₁₀ and AOD observations are available). The model underpredicts SPM, PM₁₀ and sulfate aerosol mass over both periods. The largest underprediction occurs for SPM (1978-1998, NMBF = -0.88), with smaller underpredictions for

sulfate (1978-2009, NMBF = -0.38) and PM₁₀ (1997-2009, NMBF = -0.22). Simulated European annual mean SSR has a smaller model bias (1960-2009, NMBF = 0.02). Over the period 2000-2009, the model has comparatively small biases in AOD (NMBF = -0.013) and SSR (NMBF = 0.036) but larger biases for sulfate (NMBF = -0.71) and PM₁₀ (NMBF = -0.22). ~~Underestimation of surface sulfate and~~ The model also underestimates aerosol number concentrations across the N30 (NMBF = -0.85), N50 (NMBF = -0.65) and N100 (NMBF = -0.96) size fractions in 2008-2009. Underestimation of annual and seasonal mean surface sulfate, PM₁₀ and aerosol number concentration is therefore not manifested in the simulation of the annual and seasonal cycle of AOD or SSR. Calculation of AOD requires information on the aerosol vertical profile, aerosol optics, aerosol size distribution and atmospheric humidity. Simulation of SSR strongly depends on model representation of clouds. A direct comparison of model performance in simulating surface aerosol mass with AOD or SSR is therefore complicated. Figure 15 shows the spatial correlation and variability (represented by the standard deviation in observed values normalised to the modelled values, SD_{obs}/SD_{mod}) in sulfate, SPM, PM₁₀, AOD and SSR. In general SSR and sulfate are better simulated in terms of spatial correlation and variability, with poorer model simulation of SPM, PM₁₀ and AOD.

The observed negative trends in sulfate, PM₁₀ and AOD ($-0.05 \mu\text{g S m}^{-3} \text{ yr}^{-1}$, $-0.27 \mu\text{g m}^{-3} \text{ yr}^{-1}$ and -0.007 yr^{-1}) are all well reproduced by the model ($-0.04 \mu\text{g S m}^{-3} \text{ yr}^{-1}$, $-0.14 \mu\text{g m}^{-3} \text{ yr}^{-1}$ and -0.006 yr^{-1}). Over the period 1990 to 2009, observed trends in SSR ($0.37 \text{ W m}^{-2} \text{ yr}^{-1}$) are also well simulated by the model when ARE are included ($0.32 \text{ W m}^{-2} \text{ yr}^{-1}$), but poorly simulated when ARE are excluded ($0.09 \text{ W m}^{-2} \text{ yr}^{-1}$). This confirms that being able to simulate the decline in aerosol concentrations over Europe is important for reproducing the observed brightening trend in SSR between 1990 and 2009. Prior to 1990, the model does not simulate trends in SSR as well, but few aerosol observations are available to determine the reason for model failure, which could be caused by issues with simulated aerosol, clouds or with the observations.

4 European Aerosol Radiative Forcing Trends

Figure 16 shows the changes in simulated European mean top of atmosphere (TOA) outgoing radiation, relative to a 1980 to 2000 mean, under all-sky (a) and clear-sky conditions (b) with the numbers presented in Table 6. Here we define this difference in TOA radiation as a radiative forcing (RF) between the current year and the long term mean state. European mean all-sky RF (Fig. 16a) decreases by $\gtrsim 2.0 \text{ W m}^{-2}$ (cooling trend) over the period 1960–1972, corresponding with the increase in simulated aerosol loading. From 1973 to 2009, European mean all-sky radiation increases by $\gtrsim 3.0 \text{ W m}^{-2}$ (warming trend), corresponding to the simulated reduction of aerosols. All-sky RF showed the largest increase of 6.0 W m^{-2} over central Europe between 1973 and 2009, which is consistent with this region having experienced the largest change in anthropogenic emissions (Fig. 2) and aerosol concentrations (Fig. 8). Other regions of Europe have a similar temporal change but with a smaller magnitude.

The simulated clear-sky aerosol TOA RF (Fig. 16b) is similar to that simulated under all-sky conditions. European mean clear-sky RF decreased by $\gtrsim 1.5 \text{ W m}^{-2}$ between 1960 and 1972 (cooling) and from 1973 to 2009 it increased by $\gtrsim 3.0 \text{ W m}^{-2}$ (warming). Marmier et al. (2007) reported a similar change of $+2.0 \text{ W m}^{-2}$ in the direct shortwave RF from sulfate aerosols over Europe between 1980 and 2000. This indicates the strong influence directly exerted by aerosols on the European radiative balance in response to changes in anthropogenic emissions. An estimate of the cloud albedo effect is obtained as the difference between the all-sky and clear-sky RF. Over the period 1973–2009 the cloud albedo effect is estimated to have increased by 0.44 W m^{-2} (warming), indicating that is a relatively small change when compared to that from the direct effect.

The changes in aerosol RF we simulate over Europe are slightly larger than those calculated for the USA by Leibensperger et al. (2012) of approximately $+1 \text{ W m}^{-2}$ for the direct effect and $+1 \text{ W m}^{-2}$ for the indirect (first and second) effects. The smaller changes in aerosol RF reported by Leibensperger et al. (2012) are possibly related to the smaller reductions in sulfate aerosol mass concentrations observed over the USA of $\sim 0.8 \mu\text{g S m}^{-3}$

(40%), when compared to that observed over Europe of $1.2 \mu\text{g S m}^{-3}$ (70%).

The calculated changes in all-sky TOA RF indicate the extent to which changes in anthropogenic emissions over the last 50 years have affected the European radiative balance. Reductions in anthropogenic aerosols have resulted in a positive response in the European radiative balance. We estimate that the magnitude of these emission reductions has caused European mean all-sky RF to increase by 3.0 – 3.1 W m^{-2} between the mid-1970s and 2009, mainly due to the direct aerosol effect (as shown by similar changes in the clear-sky RF). Table 6 shows that the change in European mean aerosol RF is much larger than the change in global mean aerosol RF of $+0.4 \text{ W m}^{-2}$. Shindell et al. (2013) reported a multi-model evaluation of aerosol RF over the period 1850 to 2100, with 6 of the 9 models reporting a positive increase in global mean aerosol RF between 1980 and 2000, qualitatively similar to the results presented here. At the global scale, our simulated change in all-sky aerosol RF between 1970 and 2009 is ~40% of the magnitude of change in global CO_2 radiative forcing over the same period. At the European scale, the simulated change in all-sky aerosol RF is more than three times the change in global mean CO_2 RF. This indicates the large regional impact that decreasing aerosol concentrations have had on the radiative balance and climate over Europe compared to other forcing agents.

The agreement between the model and observations in the changes in aerosols and in the brightening period of the surface radiation balance between the 1990 and 2009 provides confidence in the magnitude and temporal change of the simulated TOA RF over this period when most of the change occurs (2.0 W m^{-2}). Future work needs to explore the potential climate implications from these changes to the radiative balance. It will be important to understand the role of European air quality legislation in observed emission reductions as this may have important implications when considering the impact of future air quality mitigation measures on climate.

5 Conclusions

We used the HadGEM3-UKCA coupled chemistry climate model to simulate changes in aerosols between 1960 and 2009, a period over which anthropogenic sources of aerosol have changed substantially. We evaluated the model against European observations of sulfate aerosol mass, total suspended particulate matter (SPM), PM₁₀ mass concentrations, aerosol number concentrations, aerosol optical depth (AOD) and surface solar radiation (SSR). We also calculated the impact of changes in atmospheric aerosols on European aerosol radiative forcing.

The model underpredicts sulfate aerosol mass concentrations (NMBF = -0.4), SPM (1978-1998, NMBF = -0.9) and PM₁₀ (1997-2009, NMBF = -0.2). In addition, the model underpredicts present day aerosol number concentrations over the size fractions of N30 (NMBF= -0.85), N50 (NMBF= -0.65) and N100 (NMBF= -0.96). Underestimation of aerosol number and mass could be due to uncertainties in the observations (Tørseth et al., 2012), an overestimation of deposition processes or underestimated sources of PM including nitrate, anthropogenic SOA, domestic biomass combustion, dust and primary biological aerosol particles. The larger underestimation of particles with diameter >10 μm suggests that the sources of such particles may be more uncertain and are not well treated by the model. The model particularly underestimates sulfate in winter and over northern Europe potentially due to an under-representation of the in-cloud oxidation of sulfur species to sulfate via reaction with ozone or an enhanced wet deposition rate, caused by artificially high drizzle precipitation. Bias in simulated AOD (2000-2009, NMBF = -0.01) are smaller than for surface aerosol concentrations. Calculation of AOD requires information on the aerosol vertical profile, aerosol optics, aerosol size distribution and atmospheric humidity and complicates any direct comparison between surface aerosol mass and AOD.

Observed trends in surface aerosol mass and AOD were generally well represented by the model. Sulfate aerosol mass declines by ~~68-78% in both~~ 68% in the observations and by 78% in the model between 1978 and 2009, consistent with the decrease in SO₂ emissions over Europe. The observed European annual mean SPM decreased by 42% between 1978

and 1998, compared to a simulated decrease of 20%. Between 1997 and 2009 ~~an 8-9% decrease in~~ PM_{10} mass concentrations ~~was both observed and modelled~~ decreased by 9% in the observations and by 8% in the model. Between 2000 and 2009 ~~a decrease in AOD of 11-14% was observed and modelled~~ AOD decreased by 11% in the observations and by 14% in the model at observation sites with more than 9 years of data.

The all-sky European SSR was shown to increase between 1990 and 2009 in both the model (4.0 W m^{-2}) and observations (5.8 W m^{-2}) ('brightening'). In the model simulation where aerosol radiative effects were excluded European all-sky SSR increased by only 0.3 W m^{-2} . This comparison suggests that observed brightening post-1990 is driven by changes in aerosol that are well captured by the model. Accounting for changes to aerosols is therefore important in being able to reproduce the European brightening trend from the 1990 to 2009.

Prior to 1990, there are discrepancies between observed and simulated all-sky SSR anomalies. Specifically, the model is unable to reproduce the magnitude and timing of the observed reduction in SSR values ('dimming'). ~~Lack~~ These errors in SSR coincide with the largest model bias in observed SPM concentrations between 1978 and 1980 (8c) but the lack of extensive aerosol observations prior to 1980, prevents isolation of the cause of this model ~~discrepancy~~ observational discrepancy in SSR. Possible reasons include errors in simulated aerosols, errors associated with the meteorological reanalysis fields, and issues with the measurement data (less SSR observations were available before 1970).

From the peak in aerosol loading in the early 1970s European all-sky aerosol TOA radiative forcing has increased by ~~3.0-3.1~~ 3.1 W m^{-2} , due to a reduction in the aerosol cooling effect. Changes in European mean RF are mainly due to changes in the direct aerosol effect, as shown by ~~a similar magnitude of change in the~~ the similarity between clear-sky RF) and all-sky RF. The largest RF is over central Europe ($+6.0 \text{ W m}^{-2}$), which has seen the largest change in anthropogenic emissions and aerosol concentrations. The change in European mean aerosol RF is much larger than that globally and also three times as large as the change in global mean CO_2 RF over the period 1970-2009. Our evaluation showed that the model is able to reproduce the observed changes in SSR over the period 1990-2009,

during which two-thirds of the simulated RF occurred (2.0 W m^{-2}). Our evaluation therefore provides confidence in the simulated changes of TOA RF. The reductions in anthropogenic aerosol emissions over this period have resulted in a positive response in the radiative balance over Europe ~~due a reduction in the strength of the aerosol cooling effect, which may have contributed to increasing temperatures across Europe~~ (Philipona et al., 2009). The ~~magnitude of these changes are similar~~ RF simulated in this work are similar in magnitude to those reported by Marmer et al. (2007) over Europe and by Leibensperger et al. (2012) over the USA.

The change in anthropogenic aerosol emissions over the period 1970-2009, in part due to measures to improve air quality, has led to a considerable reduction in the concentrations of aerosols over Europe. This decrease in aerosols has reduced the aerosol radiative cooling effect over Europe. Attempts to improve European air quality over the last 30 to 40 years has potentially had non-negligible impacts on European climate (Arneth et al., 2009; Philipona et al., 2009; Ramanathan and Feng, 2009; Fiore et al., 2012) and should be the subject of future study. These air quality - climate interactions should be considered when designing any future measures to improve air quality and mitigate climate change.

Acknowledgements. ~~We thank the~~ Steven Turnock would like to acknowledge the funding for his PhD studentship from the Natural Environment Research Council (NERC) and Met Office ~~for providing the funding for this PhD studentship.~~ For making their data available to be used in this study we would like to acknowledge the EMEP, GEBA and AERONET measurement networks along with any data managers involved in data collection. We would also like to acknowledge Ari Asmi for providing the aerosol size distribution data from the EUSAAR and GUAN networks and Carly Reddington pre-processing this dataset for use in the model evaluation. Anthropogenic and biomass-burning emissions from the MACCity dataset were retrieved from the ECCAD emissions server. This work was also made possible by participation in the EU Framework 7 PEGASOS project (no: 265148). We acknowledge use of the MONSooN system, a collaborative facility supplied under the Joint Weather and Climate Research Programme, a strategic partnership between the Met Office and the Natural Environment Research Council. Matthew Woodhouse would like to thank the Royal Society for support via the International Exchange Scheme.

References

- Allen, R. J., Norris, J. R., and Wild, M.: Evaluation of multidecadal variability in CMIP5 surface solar radiation and inferred underestimation of aerosol direct effects over Europe, China, Japan, and India, *Journal of Geophysical Research: Atmospheres*, 118, 6311–6336, doi:10.1002/jgrd.50426, <http://doi.wiley.com/10.1002/jgrd.50426>, 2013.
- Andres, R. J. and Kasgnoc, A. D.: A time-averaged inventory of subaerial volcanic sulfur emissions, *Journal of Geophysical Research*, 103, 25 251, doi:10.1029/98JD02091, <http://doi.wiley.com/10.1029/98JD02091>, 1998.
- Arnth, A., Unger, N., Kulmala, M., and Andreae, M. O.: Clean the Air, Heat the Planet?, *Science*, 326, 672–673, doi:10.1126/science.1181568, <http://www.sciencemag.org/cgi/doi/10.1126/science.1181568>, 2009.
- Asmi, A., Wiedensohler, A., Laj, P., Fjaeraa, A.-M., Sellegri, K., Birmili, W., Weingartner, E., Baltensperger, U., Zdimal, V., Zikova, N., Putaud, J.-P., Marinoni, A., Tunved, P., Hansson, H.-C., Fiebig, M., Kivekäs, N., Lihavainen, H., Asmi, E., Ulevicius, V., Aalto, P. P., Swietlicki, E., Kristensson, A., Mihalopoulos, N., Kalivitis, N., Kalapov, I., Kiss, G., de Leeuw, G., Henzing, B., Harrison, R. M., Beddows, D., O'Dowd, C., Jennings, S. G., Flentje, H., Weinhold, K., Meinhardt, F., Ries, L., and Kulmala, M.: Number size distributions and seasonality of submicron particles in Europe 2008–2009, *Atmospheric Chemistry and Physics*, 11, 5505–5538, doi:10.5194/acp-11-5505-2011, <http://www.atmos-chem-phys.net/11/5505/2011/acp-11-5505-2011.html>, 2011.
- Asmi, a., Collaud Coen, M., Ogren, J. a., Andrews, E., Sheridan, P., Jefferson, a., Weingartner, E., Baltensperger, U., Bukowiecki, N., Lihavainen, H., Kivekäs, N., Asmi, E., Aalto, P. P., Kulmala, M., Wiedensohler, a., Birmili, W., Hamed, a., O'Dowd, C., G Jennings, S., Weller, R., Flentje, H., Fjaeraa, a. M., Fiebig, M., Myhre, C. L., Hallar, a. G., Swietlicki, E., Kristensson, a., and Laj, P.: Aerosol decadal trends – Part 2: In-situ aerosol particle number concentrations at GAW and AC-TRIS stations, *Atmospheric Chemistry and Physics*, 13, 895–916, doi:10.5194/acp-13-895-2013, <http://www.atmos-chem-phys.net/13/895/2013/>, 2013.
- Barnpadimos, I., Keller, J., Oderbolz, D., Hueglin, C., and Prévôt, A.: One decade of parallel fine (PM_{2.5}) and coarse (PM₁₀–PM_{2.5}) particulate matter measurements in Europe: trends and variability, *Atmospheric Chemistry and Physics*, 12, 3189–3203, <http://www.atmos-chem-phys.net/12/3189/2012/acp-12-3189-2012.pdf>, 2012.

- Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, *Journal of Geophysical Research*, 116, D20 206, doi:10.1029/2011JD016074, <http://doi.wiley.com/10.1029/2011JD016074>, 2011.
- 5 Bellouin, N., Mann, G. W., Woodhouse, M. T., Johnson, C., Carslaw, K. S., and Dalvi, M.: Impact of the modal aerosol scheme GLOMAP-mode on aerosol forcing in the Hadley Centre Global Environmental Model, *Atmospheric Chemistry and Physics*, 13, 3027–3044, doi:10.5194/acp-13-3027-2013, <http://www.atmos-chem-phys.net/13/3027/2013/acp-13-3027-2013.html>, 2013.
- 10 Berglen, T. F., Myhre, G., Isaksen, I. S., Vestreng, V., and Smith, S. J.: Sulphate trends in Europe: are we able to model the recent observed decrease?, *Tellus B*, 59, 773–786, doi:10.3402/tellusb.v59i4.17056, <http://www.tellusb.net/index.php/tellusb/article/view/17056>, 2007.
- Boucher, O., Randall, P., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2013.
- 15 Chalmers, N., Highwood, E. J., Hawkins, E., Sutton, R., and Wilcox, L. J.: Aerosol contribution to the rapid warming of near-term climate under RCP 2.6, *Geophysical Research Letters*, 39, 2–7, doi:10.1029/2012GL052848, <http://www.agu.org/pubs/crossref/2012/2012GL052848.shtml>, 2012.
- 20 Chiacchio, M., Solmon, F., Giorgi, F., Stackhouse, P., and Wild, M.: Evaluation of the radiation budget with a regional climate model over Europe and inspection of dimming and brightening, *Journal of Geophysical Research: Atmospheres*, 120, 1951–1971, doi:10.1002/2014JD022497, <http://doi.wiley.com/10.1002/2014JD022497>, 2015.
- 25 Chin, M., Diehl, T., Tan, Q., Prospero, J. M., Kahn, R. A., Remer, L. A., Yu, H., Sayer, A. M., Bian, H., Geogdzhayev, I. V., Holben, B. N., Howell, S. G., Huebert, B. J., Hsu, N. C., Kim, D., Kucsera, T. L., Levy, R. C., Mishchenko, M. I., Pan, X., Quinn, P. K., Schuster, G. L., Streets, D. G., Strode, S. A., Torres, O., and Zhao, X.-P.: Multi-decadal aerosol variations from 1980 to 2009: a perspective from observations and a global model, *Atmospheric Chemistry and Physics*, 14, 3657–3690, doi:10.5194/acp-14-3657-2014, <http://www.atmos-chem-phys.net/14/3657/2014/acp-14-3657-2014.html>, 2014.
- 30

- Colette, A., Granier, C., Hodnebrog, O., Jakobs, H., Maurizi, A., Nyiri, A., Bessagnet, B., D'Angiola, A., D'Isidoro, M., Gauss, M., Meleux, F., Memmesheimer, M., Mieville, A., Rouïl, L., Russo, F., Solberg, S., Stordal, F., and Tampieri, F.: Air quality trends in Europe over the past decade: a first multi-model assessment, *Atmospheric Chemistry and Physics*, 11, 11 657–11 678, doi:10.5194/acp-11-11657-2011, <http://www.atmos-chem-phys.net/11/11657/2011/acp-11-11657-2011.html>, 2011.
- Collaud Coen, M., Andrews, E., Asmi, a., Baltensperger, U., Bukowiecki, N., Day, D., Fiebig, M., Fjaeraa, a. M., Flentje, H., Hyvärinen, a., Jefferson, a., Jennings, S. G., Kouvarakis, G., Lihavainen, H., Lund Myhre, C., Malm, W. C., Mihapopoulos, N., Molenaar, J. V., O'Dowd, C., Ogren, J. a., Schichtel, B. a., Sheridan, P., Virkkula, a., Weingartner, E., Weller, R., and Laj, P.: Aerosol decadal trends – Part 1: In-situ optical measurements at GAW and IMPROVE stations, *Atmospheric Chemistry and Physics*, 13, 869–894, doi:10.5194/acp-13-869-2013, <http://www.atmos-chem-phys.net/13/869/2013/>, 2013.
- COMEAP: The Mortality Effects of Long-term Exposure to Particulate Air Pollution in the United Kingdom, Tech. rep., Health Protection Agency for the Committee on the Medical Effects of Air Pollutants, 2010.
- Davies, T., Cullen, M. J. P., Malcolm, A. J., Mawson, M. H., Staniforth, A., White, A. A., and Wood, N.: A new dynamical core for the Met Office's global and regional modelling of the atmosphere, *Quarterly Journal of the Royal Meteorological Society*, 131, 1759–1782, doi:10.1256/qj.04.101, <http://doi.wiley.com/10.1256/qj.04.101>, 2005.
- de Meij, a., Krol, M., Dentener, F., Vignati, E., Cuvelier, C., and Thunis, P.: The sensitivity of aerosol in Europe to two different emission inventories and temporal distribution of emissions, *Atmospheric Chemistry and Physics Discussions*, 6, 3265–3319, doi:10.5194/acpd-6-3265-2006, <http://www.atmos-chem-phys-discuss.net/6/3265/2006/>, 2006.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, doi:10.1002/qj.828, <http://doi.wiley.com/10.1002/qj.828>, 2011.

- Edwards, J. M. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model, *Quarterly Journal of the Royal Meteorological Society*, 122, 689–719, doi:10.1002/qj.49712253107, <http://doi.wiley.com/10.1002/qj.49712253107>, 1996.
- Essery, R. L. H., Best, M. J., Betts, R., and Cox, P. M.: Explicit Representation of Subgrid Heterogeneity in a GCM Land Surface Scheme, *Journal of Hydrometeorology*, 4, 530–543, 2002.
- Fagerli, H. and Aas, W.: Trends of nitrogen in air and precipitation: Model results and observations at EMEP sites in Europe, 1980–2003, *Environmental Pollution*, 154, 448–461, <http://www.sciencedirect.com/science/article/pii/S0269749108000523>, 2008.
- Farina, S. C., Adams, P. J., and Pandis, S. N.: Modeling global secondary organic aerosol formation and processing with the volatility basis set: Implications for anthropogenic secondary organic aerosol, *Journal of Geophysical Research*, 115, D09202, doi:10.1029/2009JD013046, <http://doi.wiley.com/10.1029/2009JD013046>, 2010.
- Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S., Eyring, V., Folberth, G. a., Ginoux, P., Horowitz, L. W., Josse, B., Lamarque, J.-F., MacKenzie, I. a., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S. T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T., and Zeng, G.: Global air quality and climate., *Chemical Society reviews*, 41, 6663–83, doi:10.1039/c2cs35095e, <http://www.ncbi.nlm.nih.gov/pubmed/22868337>, 2012.
- Folini, D. and Wild, M.: Aerosol emissions and dimming/brightening in Europe: Sensitivity studies with ECHAM5-HAM, *Journal of Geophysical Research*, 116, D21104, doi:10.1029/2011JD016227, <http://www.agu.org/pubs/crossref/2011/2011JD016227.shtml>, 2011.
- Gong, S. L.: A parameterization of sea-salt aerosol source function for sub- and supermicron particles, *Global Biogeochemical Cycles*, 17, 1097, doi:10.1029/2003GB002079, <http://doi.wiley.com/10.1029/2003GB002079>, 2003.
- Granier, C., Bessagnet, B., Bond, T. C., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Matsui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., Aardenne, J., Werf, G. R., and Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163–190, doi:10.1007/s10584-011-0154-1, <http://www.springerlink.com/index/10.1007/s10584-011-0154-1>, 2011.
- Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., Mckay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and

- Zimmerman, P.: A global model of natural volatile organic compound emissions, *Journal of Geophysical Research*, 100, 8873, doi:10.1029/94JD02950, <http://doi.wiley.com/10.1029/94JD02950>, 1995.
- 5 Halmer, M., Schmincke, H.-U., and Graf, H.-F.: The annual volcanic gas input into the atmosphere, in particular into the stratosphere: a global data set for the past 100 years, *Journal of Volcanology and Geothermal Research*, 115, 511–528, doi:10.1016/S0377-0273(01)00318-3, <http://www.sciencedirect.com/science/article/pii/S0377027301003183>, 2002.
- 10 Hand, J. L., Schichtel, B. a., Malm, W. C., and Pitchford, M. L.: Particulate sulfate ion concentration and SO₂ emission trends in the United States from the early 1990s through 2010, *Atmospheric Chemistry and Physics*, 12, 10353–10365, doi:10.5194/acp-12-10353-2012, <http://www.atmos-chem-phys.net/12/10353/2012/>, 2012.
- Harrison, R. M., Stedman, J., and Derwent, D.: New Directions: Why are PM10 concentrations in Europe not falling?, *Atmospheric Environment*, 42, 603–606, doi:10.1016/j.atmosenv.2007.11.023, <http://linkinghub.elsevier.com/retrieve/pii/S1352231007010795>, 2008.
- 15 Heald, C. L. and Spracklen, D. V.: Atmospheric budget of primary biological aerosol particles from fungal spores, *Geophysical Research Letters*, 36, 1–5, doi:10.1029/2009GL037493, 2009.
- Hewitt, H. T., Copley, D., Culverwell, I. D., Harris, C. M., Hill, R. S. R., Keen, a. B., McLaren, a. J., and Hunke, E. C.: Design and implementation of the infrastructure of HadGEM3: the next-generation Met Office climate modelling system, *Geoscientific Model Development*, 4, 223–253, doi:10.5194/gmd-4-223-2011, <http://www.geosci-model-dev.net/4/223/2011/>, 2011.
- 20 Hodzic, A., Madronich, S., Bohn, B., Massie, S., Menut, L., and Wiedinmyer, C.: Wildfire particulate matter in Europe during summer 2003: meso-scale modeling of smoke emissions, transport and radiative effects, *Atmospheric Chemistry and Physics*, 7, 4043–4064, doi:10.5194/acp-7-4043-2007, <http://www.atmos-chem-phys.net/7/4043/2007/acp-7-4043-2007.html>, 2007.
- 25 Holben, B., Eck, T., Slutsker, I., Tanré, D., Buis, J., Setzer, A., Vermote, E., Reagan, J., Kaufman, Y., Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sensing of Environment*, 66, 1–16, doi:10.1016/S0034-4257(98)00031-5, <http://www.sciencedirect.com/science/article/pii/S0034425798000315>, 1998.
- 30 Hurrell, J. W., Hack, J. J., Shea, D., Caron, J. M., and Rosinski, J.: A New Sea Surface Temperature and Sea Ice Boundary Dataset for the Community Atmosphere Model, *Journal of Climate*, 21, 5145–5153, doi:10.1175/2008JCLI2292.1,

<http://journals.ametsoc.org/doi/abs/10.1175/2008JCLI2292.1#.U6G54GGXQUc.mendeley>, 2008.

Jones, A., Roberts, D. L., Woodage, M. J., and Johnson, C. E.: Indirect sulphate aerosol forcing in a climate model with an interactive sulphur cycle, *Journal of Geophysical Research*, 106, 20 293, doi:10.1029/2000JD000089, <http://doi.wiley.com/10.1029/2000JD000089>, 2001.

Kettle, A. J. and Andreae, M. O.: Flux of dimethylsulfide from the oceans: A comparison of updated data sets and flux models, *Journal of Geophysical Research*, 105, 26 793, doi:10.1029/2000JD900252, <http://doi.wiley.com/10.1029/2000JD900252>, 2000.

Koch, D., Bauer, S. E., Del Genio, A., Faluvegi, G., McConnell, J. R., Menon, S., Miller, R. L., Rind, D., Ruedy, R., Schmidt, G. a., and Shindell, D. T.: Coupled Aerosol-Chemistry–Climate Twentieth-Century Transient Model Investigation: Trends in Short-Lived Species and Climate Responses, *Journal of Climate*, 24, 2693–2714, doi:10.1175/2011JCLI3582.1, <http://journals.ametsoc.org/doi/abs/10.1175/2011JCLI3582.1>, 2011.

Kreidenweis, S. M.: Modification of aerosol mass and size distribution due to aqueous-phase SO₂ oxidation in clouds: Comparisons of several models, *Journal of Geophysical Research*, 108, 4213, doi:10.1029/2002JD002697, <http://doi.wiley.com/10.1029/2002JD002697>, 2003.

Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, a., Klimont, Z., Lee, D., Liousse, C., Mieville, a., Owen, B., Schultz, M. G., Shindell, D. T., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, <http://www.atmos-chem-phys.net/10/7017/2010/>, 2010.

Langmann, B., Varghese, S., Marmer, E., Vignati, E., Wilson, J., Stier, P., and O'Dowd, C.: Aerosol distribution over Europe: a model evaluation study with detailed aerosol microphysics, *Atmospheric Chemistry and Physics*, 8, 1591–1607, <http://www.atmos-chem-phys.net/8/1591/2008/>, 2008.

Leibensperger, E. M., Mickley, L. J., Jacob, D. J., Chen, W.-T., Seinfeld, J. H., Nenes, a., Adams, P. J., Streets, D. G., Kumar, N., and Rind, D.: Climatic effects of 1950–2050 changes in US anthropogenic aerosols – Part 1: Aerosol trends and radiative forcing, *Atmospheric Chemistry and Physics*, 12, 3333–3348, doi:10.5194/acp-12-3333-2012, <http://www.atmos-chem-phys.net/12/3333/2012/>, 2012.

Liss, P. and Merlivat, L.: Air-Sea Gas Exchange Rates: Introduction and Synthesis, in: *The Role of Air-Sea Exchange in Geochemical Cycling*, pp. 113–127, D. Reidel, 1986.

- Manders, a. M. M., van Meijgaard, E., Mues, a. C., Kranenburg, R., van Ulft, L. H., and Schaap, M.: The impact of differences in large-scale circulation output from climate models on the regional modeling of ozone and PM, *Atmospheric Chemistry and Physics*, 12, 9441–9458, doi:10.5194/acp-12-9441-2012, <http://www.atmos-chem-phys.net/12/9441/2012/>, 2012.
- 5 Manktelow, P. T., Mann, G. W., Carslaw, K. S., Spracklen, D. V., and Chipperfield, M. P.: Regional and global trends in sulfate aerosol since the 1980s, *Geophysical Research Letters*, 34, L14 803, doi:10.1029/2006GL028668, <http://doi.wiley.com/10.1029/2006GL028668>, 2007.
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. a., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, *Geoscientific Model Development*, 3, 519–551, doi:10.5194/gmd-3-519-2010, <http://www.geosci-model-dev.net/3/519/2010/>, 2010.
- 10 Mann, G. W., Carslaw, K. S., Ridley, D. a., Spracklen, D. V., Pringle, K. J., Merikanto, J., Korhonen, H., Schwarz, J. P., Lee, L. a., Manktelow, P. T., Woodhouse, M. T., Schmidt, a., Breider, T. J., Emmer-son, K. M., Reddington, C. L., Chipperfield, M. P., and Pickering, S. J.: Intercomparison of modal and sectional aerosol microphysics representations within the same 3-D global chemical transport model, *Atmospheric Chemistry and Physics*, 12, 4449–4476, doi:10.5194/acp-12-4449-2012, <http://www.atmos-chem-phys.net/12/4449/2012/>, 2012.
- 15 Mann, G. W., Carslaw, K. S., Reddington, C. L., Pringle, K. J., Schulz, M., Asmi, A., Spracklen, D. V., Ridley, D. A., Woodhouse, M. T., Lee, L. A., Zhang, K., Ghan, S. J., Easter, R. C., Liu, X., Stier, P., Lee, Y. H., Adams, P. J., Tost, H., Lelieveld, J., Bauer, S. E., Tsigaridis, K., van Noije, T. P. C., Strunk, A., Vignati, E., Bellouin, N., Dalvi, M., Johnson, C. E., Bergman, T., Kokkola, H., von Salzen, K., Yu, F., Luo, G., Petzold, A., Heintzenberg, J., Clarke, A., Ogren, J. A., Gras, J., Baltensperger, U., Kaminski, U., Jennings, S. G., O'Dowd, C. D., Harrison, R. M., Beddows, D. C. S., Kulmala, M., Viisanen, Y., Ulevicius, V., Mihalopoulos, N., Zdimal, V., Fiebig, M., Hansson, H.-C., Swietlicki, E., and Henzing, J. S.: Intercomparison and evaluation of global aerosol microphysical properties among AeroCom models of a range of complexity, *Atmospheric Chemistry and Physics*, 14, 4679–4713, doi:10.5194/acp-14-4679-2014, <http://www.atmos-chem-phys.net/14/4679/2014/acp-14-4679-2014.html>, 2014.
- 20 25 30 Marmer, E., Langmann, B., Fagerli, H., and Vestreng, V.: Direct shortwave radiative forcing of sulfate aerosol over Europe from 1900 to 2000, *Journal of Geophysical Research*, 112, D23S17, doi:10.1029/2006JD008037, <http://doi.wiley.com/10.1029/2006JD008037>, 2007.

- Morgenstern, O., Braesicke, P., O'Connor, F. M., Bushell, A. C., Johnson, C. E., Osprey, S. M., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model – Part 1 : The stratosphere, *Geoscientific Model Development*, 2, 43–57, www.geosci-model-dev.net/2/43/2009, 2009.
- Myhre, G., Shindell, D., Breon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- O'Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., Dalvi, M., Folberth, G. A., Sanderson, M. G., Telford, P. J., Voulgarakis, A., Young, P. J., Zeng, G., Collins, W. J., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model – Part 2: The Troposphere, *Geoscientific Model Development*, 7, 41–91, doi:10.5194/gmd-7-41-2014, <http://www.geosci-model-dev.net/7/41/2014/gmd-7-41-2014.html>, 2014.
- Penner, J. E., Prather, M. J., Isaksen, I. S. a., Fuglestedt, J. S., Klimont, Z., and Stevenson, D. S.: Short-lived uncertainty?, *Nature Geoscience*, 3, 587–588, doi:10.1038/ngeo932, <http://www.nature.com/doi/finder/10.1038/ngeo932>, 2010.
- Philipona, R., Behrens, K., and Ruckstuhl, C.: How declining aerosols and rising greenhouse gases forced rapid warming in Europe since the 1980s, *Geophysical Research Letters*, 36, L02806, doi:10.1029/2008GL036350, <http://onlinelibrary.wiley.com/doi/10.1029/2008GL036350/full>, 2009.
- Pozzer, A., de Meij, A., Pringle, K. J., Tost, H., Doering, U. M., van Aardenne, J., and Lelieveld, J.: Distributions and regional budgets of aerosols and their precursors simulated with the EMAC chemistry-climate model, *Atmospheric Chemistry and Physics*, 12, 961–987, doi:10.5194/acp-12-961-2012, <http://www.atmos-chem-phys.net/12/961/2012/acp-12-961-2012.html>, 2012.
- Ramanathan, V. and Feng, Y.: Air pollution, greenhouse gases and climate change: Global and regional perspectives, *Atmospheric Environment*, 43, 37–50, doi:10.1016/j.atmosenv.2008.09.063, <http://www.sciencedirect.com/science/article/pii/S1352231008008583>, 2009.
- Sanchez-Lorenzo, A., Wild, M., and Trentmann, J.: Validation and stability assessment of the monthly mean CM SAF surface solar radiation dataset over Europe against a homogenized surface dataset (1983–2005), *Remote Sensing of Environment*, 134, 355–366, doi:10.1016/j.rse.2013.03.012, <http://www.sciencedirect.com/science/article/pii/S0034425713000916>, 2013.

- Schultz, M. G., Heil, A., Hoelzemann, J. J., Spessa, A., Thonicke, K., Goldammer, J. G., Held, A. C., Pereira, J. M. C., and van het Bolscher, M.: Global wildland fire emissions from 1960 to 2000, *Global Biogeochemical Cycles*, 22, n/a–n/a, doi:10.1029/2007GB003031, <http://doi.wiley.com/10.1029/2007GB003031>, 2008.
- 5 Scott, C. E., Rap, a., Spracklen, D. V., Forster, P. M., Carslaw, K. S., Mann, G. W., Pringle, K. J., Kivekäs, N., Kulmala, M., Lihavainen, H., and Tunved, P.: The direct and indirect radiative effects of biogenic secondary organic aerosol, *Atmospheric Chemistry and Physics*, 14, 447–470, doi:10.5194/acp-14-447-2014, 2014.
- 10 Shindell, D. T., Lamarque, J.-F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P., Lee, Y. H., Rotstajn, D., Mahowald, N., Milly, G., Faluvegi, G., Collins, W., Conley, A., Dalsoren, S., Easter, R., Ghan, S., Horowitz, L., Liu, X., Myhre, G., Nagashima, T., Naik, V., Rumbold, S., Skeie, R., Sudo, K., Szopa, S., Takemura, T., Voulgarakis, A., Yoon, J.-H., and Lo, F.: Radiative forcing in the ACCMIP historical and future climate simulations, *Atmospheric Chemistry and Physics*, 13, 2939–2974, doi:10.5194/acp-13-2939-2013, 2013.
- 15 Skeie, R. B., Berntsen, T. K., Myhre, G., Tanaka, K., Kvalevåg, M. M., and Hoyle, C. R.: Anthropogenic radiative forcing time series from pre-industrial times until 2010, *Atmospheric Chemistry and Physics*, 11, 11827–11857, doi:10.5194/acp-11-11827-2011, <http://www.atmos-chem-phys.net/11/11827/2011/>, 2011.
- 20 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, a., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmospheric Chemistry and Physics*, 11, 1101–1116, doi:10.5194/acp-11-1101-2011, <http://www.atmos-chem-phys.net/11/1101/2011/>, 2011.
- Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Mann, G. W., and Sihto, S.-L.: The contribution of boundary layer nucleation events to total particle concentrations on regional and global scales, *Atmospheric Chemistry and Physics*, 6, 5631–5648, doi:10.5194/acp-6-5631-2006, <http://www.atmos-chem-phys.net/6/5631/2006/acp-6-5631-2006.html>, 2006.
- 25 Spracklen, D. V., Carslaw, K. S., Pöschl, U., Rap, a., and Forster, P. M.: Global cloud condensation nuclei influenced by carbonaceous combustion aerosol, *Atmospheric Chemistry and Physics*, 11, 9067–9087, doi:10.5194/acp-11-9067-2011, <http://www.atmos-chem-phys.net/11/9067/2011/>, 2011a.
- 30 Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, a., and Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol bud-

- get, *Atmospheric Chemistry and Physics*, 11, 12 109–12 136, doi:10.5194/acp-11-12109-2011, <http://www.atmos-chem-phys.net/11/12109/2011/>, 2011b.
- 5 Tørseth, K., Aas, W., Breivik, K., Fjæ raa, a. M., Fiebig, M., Hjellbrekke, a. G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, *Atmospheric Chemistry and Physics*, 12, 5447–5481, doi:10.5194/acp-12-5447-2012, <http://www.atmos-chem-phys.net/12/5447/2012/>, 2012.
- 10 Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E., Bellouin, N., Benedetti, A., Bergman, T., Berntsen, T. K., Beukes, J. P., Bian, H., Carslaw, K. S., Chin, M., Curci, G., Diehl, T., Easter, R. C., Ghan, S. J., Gong, S. L., Hodzic, A., Hoyle, C. R., Iversen, T., Jathar, S., Jimenez, J. L., Kaiser, J. W., Kirkevå g, A., Koch, D., Kokkola, H., Lee, Y. H., Lin, G., Liu, X., Luo, G., Ma, X., Mann, G. W., Mihalopoulos, N., Morcrette, J.-J., Müller, J.-F., Myhre, G., Myriokefalitakis, S., Ng, N. L., O'Donnell, D., Penner, J. E., Pozzoli, L., Pringle, K. J., Russell, L. M., Schulz, M., Sciare, J., Seland, O., Shindell, D. T., Sillman, S., Skeie, R. B., Spracklen, D., Stavrou, T., Steenrod, S. D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., van Noije, T., van Zyl, P. G., von Salzen, K., Yu, F., Wang, Z., Zaveri, R. A., Zhang, H., Zhang, K., Zhang, Q., and Zhang, X.: The AeroCom evaluation and intercomparison of organic aerosol in global models, *Atmospheric Chemistry and Physics*, 14, 10 845–10 895, doi:10.5194/acp-14-10845-2014, <http://www.atmos-chem-phys.net/14/10845/2014/acp-14-10845-2014.html>, 2014.
- 20 Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Berg, L. V. D., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012, doi:10.1256/qj.04.176, <http://doi.wiley.com/10.1256/qj.04.176>, 2005.
- 25 Vestreng, V., Myhre, G., Fagerli, H., Reis, S., and Tarrasón, L.: Twenty-five years of continuous sulphur dioxide emission reduction in Europe, *Atmospheric Chemistry and Physics*, 7, 3663–3681, doi:10.5194/acp-7-3663-2007, <http://www.atmos-chem-phys.net/7/3663/2007/acp-7-3663-2007.html>, 2007.
- 30

Volkamer, R., Jimenez, J. L., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L. T., Worsnop, D. R., and Molina, M. J.: Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophysical Research Letters*, 33, L17811, doi:10.1029/2006GL026899, <http://doi.wiley.com/10.1029/2006GL026899>, 2006.

5 Walters, D. N., Best, M. J., Bushell, A. C., Copsey, D., Edwards, J. M., Falloon, P. D., Harris, C. M., Lock, A. P., Manners, J. C., Morcrette, C. J., Roberts, M. J., Stratton, R. A., Webster, S., Wilkinson, J. M., Willett, M. R., Boutle, I. A., Earnshaw, P. D., Hill, P. G., MacLachlan, C., Martin, G. M., Moufouma-Okia, W., Palmer, M. D., Petch, J. C., Rooney, G. G., Scaife, A. A., and Williams, K. D.: The Met Office Unified Model Global Atmosphere 3.0/3.1 and JULES Global Land 3.0/3.1 configurations, *Geoscientific Model Development*, 4, 919–941, doi:10.5194/gmd-4-919-2011, <http://www.geosci-model-dev.net/4/919/2011/gmd-4-919-2011.html>, 2011.

10 Wild, M.: Global dimming and brightening: A review, *Journal of Geophysical Research*, 114, D00D16, doi:10.1029/2008JD011470, <http://www.agu.org/pubs/crossref/2009/2008JD011470.shtml>, 2009.

15 Wild, O., Zhu, X., and Prather, M. J.: Fast-J: Accurate Simulation of In- and Below-Cloud Photolysis in Tropospheric Chemical Models, *Journal of Atmospheric Chemistry*, 37, 245–282, doi:10.1023/A:1006415919030, <http://0-link.springer.com.wam.leeds.ac.uk/article/10.1023/A:1006415919030>, 2000.

20 Woodward, S.: Modeling the atmospheric life cycle and radiative impact of mineral dust in the Hadley Centre climate model, *Journal of Geophysical Research*, 106, 18 155, doi:10.1029/2000JD900795, http://apps.webofknowledge.com/full_record.do?product=UA&search_mode=GeneralSearch&qid=918, 2001.

25 Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. a., Anderson, T. L., Balkanski, Y., Belouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T., and Zhou, M.: A review of measurement-based assessments of the aerosol direct radiative effect and forcing, *Atmospheric Chemistry and Physics*, 6, 613–666, doi:10.5194/acp-6-613-2006, <http://www.atmos-chem-phys.net/6/613/2006/>, 2006.




Figure 1. Location of measurements used in this study for a) sulfate aerosol mass (*), total aerosol mass (◇) and sites that have measured both (×), b) AOD (□) and surface solar radiation (+). Regional European definitions are North West Europe (green), Northern Europe (Red), Central Europe (Blue), Eastern Europe (Orange) and Southern Europe (Purple).




Figure 2. Annual European emissions (Gg yr^{-1}) of aerosols and aerosol precursors from the MACC-inventory over the period 1960-2009. a) Total European emissions of sulfur dioxide (SO_2 , orange), organic carbon (OC, light green), black carbon (BC, black), ammonia (NH_3 , light purple) and oxides of nitrogen (NO_x , blue). b) to f) shows European regional emissions for b) sulfur dioxide SO_2 , c) oxides of nitrogen NO_x , d) ammonia NH_3 , e) black carbon BC and f) organic carbon OC. Regions are as defined in Fig. 1: northern (red), north west (green), central (blue), eastern (orange), southern (purple).




Figure 3. Temporal variation in the number of locations measuring surface solar radiation (SSR), sulfate aerosol mass (SO₄), total suspended particle mass (SPM), PM₁₀ mass and aerosol optical depth (AOD ~~within each particular network~~).




Figure 4. Simulated European annual mean a) PM_{10} and composition resolved surface mass concentration, b) surface solar radiation aerosol number concentration for particles with dry diameter >3 nm (total), >30 nm (N30), >50 nm (N50) and >100 nm (N100), c) aerosol optical depth (AOD) at 440 nm and d) surface solar radiation (SSR) between 1960 and 2009. Mean values are calculated over the European land domain shown in Fig. 1




Figure 5. Annual mean sulfate aerosol mass concentrations ($\mu\text{g S m}^{-3}$) from the lowest model level with observations overplotted in circles for a) 1980, b) 1990, c) 2000 and d) 2009. e) **shows a time Time** series of annual and seasonal mean observed (solid lines) and modelled (dashed lines) sulfate concentrations averaged across all measurement locations for each particular year. Shaded areas show ± 1 standard deviation of the modelled (light grey) and observed (dark grey) annual mean values, with the hatching identifying areas of overlap.




Figure 6. European and sub-regional Normalised Mean Bias Factors for summertime (red) and wintertime (blue) sulfate aerosol mass concentrations across all the years that data was available. The solid line shows the median value, the boxes show the 25th and 75th percentile values with the error bars showing the maximum and minimum values and the circles representing outliers (values > 1.5 * inter quartile range).




Figure 7. Annual (black), summertime (red) and wintertime (blue) trends in modelled and observed sulfate aerosol mass concentrations ($\mu\text{g S m}^{-3} \text{ yr}^{-1}$) at each individual monitoring location over their operational period between 1978 and 2009.




Figure 8. Annual mean suspended particulate matter (SPM) mass concentrations ($\mu\text{g m}^{-3}$) from the lowest model level with observations overplotted in circles for a) 1980 and b) 1990. d) and e) are the same as a) and b) but for PM₁₀ mass concentrations in 2000 and 2009. A time series of annual and seasonal mean observed (solid lines) and modelled (dashed lines) SPM-c) SPM and PM₁₀-f) PM₁₀ concentrations averaged across all measurement locations for each particular year. Shaded areas show ± 1 standard deviation of the modelled (light grey) and observed (dark grey) annual mean values, with the hatching identifying areas of overlap.




Figure 9. European and sub-regional Normalised Mean Bias Factors for summertime (red) and wintertime (blue) [SPM](#)-a) [suspended particulate matter \(SPM\)](#) and [PM₁₀](#)-b) [PM₁₀](#) mass concentrations across all the years that data was available. The solid line shows the median value, the boxes show the 25th and 75th percentile values with the error bars showing the maximum and minimum values and the circles representing outliers (values > 1.5 * inter quartile range).




Figure 10. Annual (black), summertime (red) and wintertime (blue) calculated linear trends in modelled and observed **SPM**-a) suspended particulate matter (SPM) and **PM₁₀**-b) PM₁₀ mass concentrations ($\mu\text{g S m}^{-3} \text{ yr}^{-1}$) at each individual monitoring location over their operational period between 1978 to 2002 for SPM and 1996 to 2009 for PM₁₀.




Figure 11. Annual mean observed vs simulated aerosol number concentrations for a) N30 (particles greater than 30 nm diameter), b) N50 (>50 nm) and c) N100 (>100 nm) for monitoring sites across Europe.




Figure 12. Annual mean ~~Aerosol Optical Depth~~ aerosol optical depth (AOD) at 440 nm with observations from the AERONET sites overplotted for a) 2000 and b) 2009. c) ~~A time~~ Time series of monthly and seasonal mean observed (solid lines) and modelled (dashed lines) AOD at 440 nm averaged across all measurement locations. d) European Normalised Mean Bias Factors (NMBF) of AOD for annual (orange), summertime (red) and wintertime (blue) across all the years and sites that data was available. The annual NMBF for the 3 sites with the longest continuous record (> 9 years) is shown in yellow.




Figure 13. Observed (black line) and simulated (red line) European annual mean all-sky surface solar radiation (SSR) anomalies (Wm^{-2}) relative to a 1980 to 2000 average. Simulated all-sky SSR not including aerosol radiative effects (ARE) is also shown as the additional (orange coloured line). Values are calculated as the average across all measurement locations within Europe (see Fig. 1). The standard deviation of the annual mean for each year is shown by the shaded areas of dark grey for the observations and light grey for the model with the hatching representing where the areas overlap.




Figure 14. European Normalised Mean Bias Factors for sulfate aerosol mass, suspended particulate matter (SPM) mass, PM_{10} mass, aerosol optical depth (AOD) and surface solar radiation (SSR) across all the years that data was available (see Tables 2 to 5) and ~~in~~ separately for the period 2000-2009. The solid line shows the median value, the boxes show the 25th and 75th percentile values with the error bars showing the maximum and minimum values and the circles representing outliers (values $> 1 * \text{inter quartile range}$).




Figure 15. Taylor diagram comparing modelled-simulated and observed European sulfate mass concentrations, suspended particulate matter (SPM), PM₁₀, aerosol optical depth (AOD) and SSR surface solar radiation (for the three different time periods SSR)for-. Comparisons are made across all the years that data is observations are available and in-. We additionally show results for the period 2000-2009, where sulfate, PM₁₀, AOD and SSR are consistently available. For AOD, we separately plot the 3 locations with the longest observational record. For SSR we also across three different time periods (1960-1974, 1975-1989, 1990-2009). Correlation coefficients are plotted against the observed standard deviations observed for each component that have been normalised relative to the calculated model-simulated standard deviation values.




Figure 16. European and sub-regional top of atmosphere aerosol radiative forcing (Wm^{-2}), relative to a 1980 to 2000 average, under **all-sky** (a) **all-sky** and **clear-sky** (b) **clear-sky** conditions. European regions are as defined in Fig. 1 -

Table 1. Details of the ground based observations used in this study.

Data Source	Measurements	Period	Total Number of Sites Available
EMEP	Total Sulfate Aerosol Mass Concentrations	1978-2009	97
EMEP	Total Suspended Particle Mass Concentrations	1978-2005	42
EMEP	PM ₁₀ Mass Concentrations	1996-2009	52
EUSAAR and GUAN	Aerosol number concentrations greater than 30 nm (N30), 50 nm (N50) and 100 nm (N100) in dry diameter	2008-2009	24
AERONET	Aerosol Optical Depth (AOD)	1994-2009	21
GEBA	Surface Solar Radiation (SSR)	1928-2009	50

Table 2. Statistical summary of modelled and observed annual and seasonal (DJF and JJA) mean sulfate at all long term (>20 years data) measurement sites (34 sites) between 1978 to 2009. Absolute and relative (%) changes in concentrations are calculated as the difference between the mean of the initial 5 years of data minus the mean of the last 5 years of data. Trends that are above or below the value of twice of the standard error of the trend (95% confidence) are highlighted in bold.

Component	Type	Absolute Change in Concentration ($\mu\text{g S m}^{-3}$)	Standard Deviation of Mean Value	% Change in Concentration	Calculated Linear Trend ($\mu\text{g S m}^{-3} \text{ yr}^{-1}$)	-2 * S.E. of trend	+ 2 * S.E. of trend	r^2	NMBF	RMSE
Sulfate Annual	Observed	-1.23	0.466	-68%	-0.048	-0.053	-0.043	0.434	-0.384	0.334
	Model	-1.15	0.424	-78%	-0.044	-0.048	-0.040			
Sulfate DJF	Observed	-1.67	0.65	-73%	-0.064	-0.073	-0.055	0.385	-2.19	0.978
	Model	-0.53	0.20	-75%	-0.020	-0.023	-0.017			
Sulfate JJA	Observed	-1.39	0.57	-70%	-0.056	-0.064	-0.048	0.434	0.694	1.17
	Model	-3.35	1.34	-86%	-0.138	-0.152	-0.124			

Table 3. Statistical summary of modelled and observed annual and seasonal (DJF and JJA) mean SPM concentrations at all long term (>20 years data) measurement sites (34 sites) between 1978 to 1998. A similar comparison is presented for PM₁₀ mass concentrations at all long term (>10 years data) measurement sites (16 sites) between 1997 to 2009. Absolute and relative (%) changes in concentrations are calculated as the difference between the mean of the initial 5 years of data minus the mean of the last 5 years of data. Trends that are above or below the value of twice of the standard error of the trend (95% confidence) are highlighted in bold.

Component	Type	Absolute Change in Concentration ($\mu\text{g m}^{-3}$)	Standard Deviation of Mean Value	% Change in Concentration	Calculated Linear Trend ($\mu\text{g m}^{-3} \text{ yr}^{-1}$)	-2 * S.E. of trend	+ 2 * S.E. of trend	r ²	NMBF	RMSE
SPM Annual	Observed	-18.32	7.96	-42%	-1.19	-1.41	-0.97	0.104	-0.88	18.04
	Model	-4.11	2.36	-20%	-0.26	-0.39	-0.14			
SPM DJF	Observed	-24.50	10.81	-51%	-1.52	-1.91	-1.13	0.044	-0.852	19.39
	Model	-3.95	3.82	-19%	-0.27	-0.52	-0.02			
SPM JJA	Observed	-14.80	7.74	-35%	-1.04	-1.36	-0.72	0.354	-0.796	16.41
	Model	-5.06	2.94	-23%	-0.38	-0.51	-0.25			
PM ₁₀ Annual	Observed	-1.50	1.84	-9%	-0.27	-0.50	-0.03	0.037	-0.221	3.35
	Model	-1.08	1.16	-8%	-0.14	-0.30	0.02			
PM ₁₀ DJF	Observed	-1.53	2.35	-9%	-0.29	-0.61	0.03	0.023	-0.12	4.01
	Model	-2.40	3.04	-15%	-0.29	-0.73	0.14			
PM ₁₀ JJA	Observed	0.19	1.87	1%	-0.02	-0.31	0.27	0.295	-0.339	4.66
	Model	3.12	2.75	30%	0.32	-0.06	0.70			

Table 4. Statistical summary of modelled and observed annual and seasonal (DJF and JJA) mean ~~Aerosol Optical Depth~~ aerosol optical depth (AOD) at all long term (>7 years data) measurement sites (13 sites) between 2000 to 2009. A similar comparison is presented at the 3 measurement locations that contain annual mean observations throughout every year between 2000 and 2009 (AOD 3 Sites). Absolute and relative (%) changes in concentrations are calculated as the difference between the mean of the initial 5 years of data minus the mean of the last 5 years of data. Trends that are above or below the value of twice of the standard error of the trend (95% confidence) are highlighted in bold.

Component	Type	Absolute Change in AOD	Standard Deviation of Mean Value	% Change in AOD	Calculated Linear Trend (AOD yr ⁻¹)	-2 * S.E. of trend	+ 2 * S.E. of trend	r ²	NMBF	RMSE
AOD	Observed	-0.03	0.022	-12%	-0.0061	-0.009	-0.003	0.031	-0.013	0.026
	Model	-0.006	0.014	-2%	0.00005	-0.003	0.003			
AOD 3 Sites	Observed	-0.03	0.037	-11%	-0.007	-0.014	-0.001	0.011	-0.296	0.066
	Model	-0.03	0.019	-14%	-0.006	-0.007	-0.004			
AOD DJF	Observed	-0.024	0.026	-13%	-0.0063	-0.010	-0.002	0.024	0.258	0.050
	Model	-0.040	0.027	-16%	-0.0074	-0.011	-0.004			
AOD JJA	Observed	-0.072	0.047	-23%	-0.014	-0.019	-0.008	0.039	-0.167	0.053
	Model	-0.030	0.021	-12%	-0.005	-0.008	-0.002			

Table 5. Statistical summary of ~~modelled~~ simulated and observed annual mean surface solar radiation (SSR) for three different time periods ~~of 1960 to 1974~~ (1960-1974, ~~1975 to 1989~~ 1975-1989 and ~~1990 to 2009~~ 1990-2009) at the 20 measurement sites across Europe, which have a continuous 50 year data record. Absolute and relative (%) changes in ~~anomaly values~~ SSR are calculated as the difference between the mean of the initial 5 years of data minus the mean of the last 5 years of data for the time period considered. Trends that are above or below the value of twice of the standard error of the trend (95% confidence) are highlighted in bold.

Component	Type	Time Period	Absolute Change in SSR (W m ⁻²)	Standard Deviation of Mean Value	% Change in SSR	Calculated Linear Trend (W yr ⁻¹)	-2 * S.E. of trend	+ 2 * S.E. of trend	r ²	NMBF	RMSE
All Europe	Observed		-0.79	2.40	-1	-0.011	-0.309	0.286			
	Model +	1960-1974	-1.67	1.89	-1	-0.088	-0.317	0.141	0.838	0.002	7.72
	ARE										
	Model -		-0.20	2.12	-0.2	0.069	-0.237	0.376	0.748	0.045	11.04
	ARE										
	Observed		-2.24	3.42	-2	-0.259	-0.658	0.140	0.803	0.029	8.43
	Model +	1975-1989	3.12	3.06	3	0.304	-0.036	0.643	0.803	0.029	8.43
	ARE										
	Model -		1.89	3.50	1.6	0.187	-0.191	0.564	0.757	0.063	11.84
	ARE										
	Observed		5.78	3.47	5	0.369	0.153	0.584	0.896		8.00
	Model +	1990-2009	3.98	3.04	3	0.316	0.125	0.507	0.896	0.043	8.00
ARE											
Model -		0.27	2.92	0.2	0.09	-0.185	0.366	0.828	0.053	10.23	
ARE											

Table 6. Global and European shortwave top of atmosphere all-sky and clear-sky aerosol radiative forcing (RF), relative to a 1980 to 2000 mean. Values for 1972 are included as this is when the simulated minimum aerosol RF occurs over Europe. For comparison we include global carbon dioxide RF (relative to 1750) from the IPCC fifth assessment report (Myhre et al., 2013).

Year	All-sky Aerosol RF (W m^{-2})		Clear-sky Aerosol RF (W m^{-2})		Estimates of CO_2 radiative forcing (Myhre et al., 2013)
	Global	Europe	Global	Europe	
1960	+0.9	+0.8	+0.4	+0.2	+0.7
1970	-0.1	-1.0	+0.01	-0.1	+0.9
1972	-0.1	-1.4	+0.2	-1.6	N/A
1980	-0.4	-0.6	+0.01	-0.5	+1.1
1990	+0.01	+0.1	+0.04	-0.1	+1.3
2000	+0.4	+1.7	-0.1	+1.1	+1.5
2009	+0.3	+2.1	+0.1	+1.9	+1.8