

Does GOSAT capture the true seasonal cycle of XCO₂?

Hannakaisa Lindqvist^{1,2}, Christopher W. O'Dell^{1,4}, Sourish Basu^{3,4},
Hartmut Boesch⁵, Frédéric Chevallier⁶, Nicholas Deutscher^{7,8}, Liang Feng⁹,
Brendan Fisher¹⁰, Frank Hase¹¹, Makoto Inoue^{12,13}, Rigel Kivi¹⁴, Isamu Morino¹²,
Paul I. Palmer⁹, Robert Parker⁵, Matthias Schneider¹¹, Ralf Sussmann¹⁵, and
Yukio Yoshida¹²

¹Department of Atmospheric Science, Colorado State University, CO, USA

²Finnish Meteorological Institute, Helsinki, Finland

³NOAA Earth System Research Laboratory, Boulder, CO, USA

⁴Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

⁵National Centre for Earth Observation, Department of Physics and Astronomy, University of Leicester, Leicester, UK

⁶Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, Gif-sur-Yvette, France

⁷School of Chemistry, University of Wollongong, Wollongong, Australia

⁸Institute of Environmental Physics, University of Bremen, Bremen, Germany

⁹National Centre for Earth Observation, School of GeoSciences, University of Edinburgh, Edinburgh, UK

¹⁰Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

¹¹Karlsruhe Institute of Technology, IMK-ASF, Karlsruhe, Germany

¹²National Institute for Environmental Studies (NIES), Tsukuba, Japan

¹³Department of Biological Environment, Akita Prefectural University, Akita, Japan

¹⁴Finnish Meteorological Institute, Arctic Research Centre, Sodankylä, Finland

¹⁵Karlsruhe Institute of Technology, IMK-IFU, Garmisch-Partenkirchen, Germany

Correspondence to: Hannakaisa Lindqvist (hannakaisa.lindqvist@fmi.fi)

Abstract.

The seasonal cycle accounts for a dominant mode of total column CO₂ (XCO₂) annual variability and is connected to CO₂ uptake and release; it thus represents an important quantity to test the accuracy of the measurements from space. We quantitatively evaluate the XCO₂ seasonal cycle of the
5 Greenhouse Gases Observing Satellite (GOSAT) observations from the Atmospheric Carbon Observations from Space (ACOS) retrieval system, and compare average regional seasonal cycle features to those directly measured by the Total Carbon Column Observing Network (TCCON). We analyze the mean seasonal cycle amplitude, dates of maximum and minimum XCO₂, as well as the regional growth rates in XCO₂ through the fitted trend over several years. We find that GOSAT generally
10 captures the seasonal cycle amplitude within 1.0 ppm accuracy compared to TCCON, except in Europe, where the difference exceeds 1.0 ppm at two sites, and the amplitude captured by GOSAT is generally shallower compared to TCCON. This bias over Europe is not as large for the other GOSAT retrieval algorithms (NIES v02.21, RemoTeC v2.35, UoL v5.1, and NIES PPDF-S v.02.11), although they have significant biases at other sites. The ACOS bias correction was found to partially explain

15 the shallow amplitude over Europe. The impact of the co-location method and aerosol changes in the
ACOS algorithm were also tested, and found to be few tenths-of-a-ppm and mostly non-systematic.
We find generally good agreement in the date of minimum XCO₂ between ACOS and TCCON, but
ACOS generally infers a date of maximum XCO₂ 2-3 weeks later than TCCON. We further analyze
the latitudinal dependence of the seasonal cycle amplitude throughout the Northern Hemisphere,
20 and compare the dependence to that predicted by current optimized models that assimilate in-situ
measurements of CO₂. In the zonal averages, models are consistent with the GOSAT amplitude to
within 1.4 ppm, depending on the model and latitude. We also show that the seasonal cycle of XCO₂
depends on longitude especially at the mid-latitudes: the amplitude of GOSAT XCO₂ doubles from
West U.S. to East Asia at 45 – 50°N, which is only partially shown by the models. In general, we
25 find that model-to-model differences can be larger than GOSAT-to-model differences. These results
suggest that GOSAT retrievals of the XCO₂ seasonal cycle may be sufficiently accurate to evaluate
land surface models in regions with significant discrepancies between the models.

1 Introduction

Satellites provide unprecedented spatial coverage of the variability of atmospheric carbon dioxide
30 (CO₂) through retrievals of column mean dry mole fractions of CO₂ (XCO₂). XCO₂ shows tem-
poral variability on different timescales: diurnal, synoptic, seasonal, inter-annual, and long-term
(Olsen and Randerson, 2004; Keppel-Aleks et al., 2011). Variability is determined by the collec-
tive impact of CO₂ fluxes resulting from fossil fuel emissions, biosphere-atmosphere exchange, and
ocean-atmosphere exchange, and the imprint of these on regional XCO₂ can be strongly influenced
35 by atmospheric dynamics, in addition to the regional origin of the fluxes. While the secular trend
and multi-year interhemispheric CO₂ gradient are driven by the global build-up of CO₂ from fossil
fuel combustion mainly in the Northern Hemisphere, the seasonal variability is mainly controlled
by variations in the terrestrial biospheric fluxes (Palmer et al., 2008; Keppel-Aleks et al., 2011). The
ocean-atmosphere and fossil fuel CO₂ fluxes are, although seasonally varying, only minor contribu-
40 tors to the XCO₂ seasonal variability in the Northern Hemisphere. Therefore, the seasonal cycle of
XCO₂ bears the signature of large-scale biospheric flux patterns, especially their north-south distri-
bution.

Regional biospheric CO₂ fluxes are a critical output of land surface models that describe the
biosphere-atmosphere carbon exchange in larger modeling systems, such as coupled climate - carbon
45 cycle models (Pitman, 2003). Inverse model systems use these land surface models in conjunction
with atmospheric transport models, and optimize their CO₂ flux estimates by assimilating CO₂ mea-
surements, but especially in regions where the in-situ measurement network has sparse coverage, the
inverse models can strongly disagree about the seasonality and magnitude of the fluxes (Lindqvist et
al., 2015, in prep.). Recently, this disagreement has been found to lead to large regional discrepan-

50 cies of several ppm in the seasonal cycle amplitudes of modeled XCO₂ (Keppel-Aleks et al., 2012;
Peng et al., 2015; Lindqvist et al., 2015, in prep.). This finding suggests that regional XCO₂ seasonal
cycles may be indicative of local fluxes, and hence that satellite-measured XCO₂ may be useful in
evaluating model fidelity without resorting to full carbon flux inversions. It is also another reminder
that there may be much to be gained by assimilating space-based XCO₂ retrievals which vastly ex-
55 pand the current in-situ measurement network; a lesson shown previously by a number of studies
(e.g., Rayner and O'Brien, 2001; Chevallier et al., 2007; Takagi et al., 2011; Maksuytov et al., 2013;
Takagi et al., 2014). In particular, the strength of the seasonal cycle drawdown is fundamentally
connected to the magnitude of the carbon sink during the growing season. By studying the GOSAT
XCO₂ seasonal cycle and its inter-annual variability, Wunch et al. (2013) showed that the variability
60 in the drawdown correlates with surface temperature in the boreal regions, and Guerlet et al. (2013b)
found a reduced carbon uptake during the 2010 Northern Hemisphere summer.

The Greenhouse Gases Observing Satellite (GOSAT; Yokota et al., 2009) and the Orbiting Car-
bon Observatory -2 (OCO-2; Crisp et al., 2004) are indeed designed to make near-global XCO₂
retrievals that will constrain the inverse model systems enough to provide a picture of the global
65 carbon cycle with respect to regional sources and sinks. However, a crucial question still lingers:
are the satellite observations accurate enough to reliably capture the seasonal variability of XCO₂?
The question is fair because satellite-retrieved XCO₂ is subject to biases in the retrieval system (e.g.,
Wunch et al., 2011b), and also sampling biases due to the seasonally-dependent amount of solar
radiation (e.g., Liu et al., 2014). Both of these may have an impact on the measured seasonal cycle.
70 For the Atmospheric CO₂ Observations from Space (ACOS) retrieval system (O'Dell et al., 2012;
Crisp et al., 2012), known biases in GOSAT retrievals are corrected using a global bias correction
(Wunch et al., 2011b) but some parameters of the bias correction vary seasonally, for example sur-
face albedo. Potential remaining biases, their seasonality, and impact on the seasonal cycles of XCO₂
are best identified through evaluation of the GOSAT seasonal cycle against the best available inde-
75 pendent data — those from the Total Carbon Column Observing Network, TCCON (Wunch et al.,
2011a). There have been several studies that compare GOSAT retrievals against the TCCON, some
of them introducing novel methods for comparisons (Wunch et al., 2011b; Nguyen et al., 2014),
some concentrating on quantifying biases in a specific retrieval algorithm (Butz et al., 2011; Cogan
et al., 2012; Yoshida et al., 2013), and some focusing more on the intercomparison of different re-
80 trieval algorithms (Buchwitz et al., 2013; Oshchepkov et al., 2013a; Reuter et al., 2013; Dils et al.,
2014). Overall, the collective message from the validation studies is that the agreement of GOSAT
and TCCON has improved (i.e., the satellite biases have decreased) substantially from the earliest
validation efforts (Morino et al., 2011), owing to major improvements and updates in the retrieval
algorithms and the development of more sophisticated comparison methods. However, less attention
85 has been paid to the evaluation of the seasonal cycle. Reuter et al. (2013, p. 1776) touched on this
by showing averages of the seasonal cycle amplitude differences between all GOSAT retrievals and

TCCON (and also a model, CarbonTracker CT2011_oi). More recently, Kulawik et al. (2015) studied the seasonality of GOSAT-TCCON biases (using the ACOS B3.4 retrieval algorithm for GOSAT data) and found notable station-to-station variability in the biases, but also persisting seasonal biases in latitudinally averaged results. These seasonal biases were reflected in the seasonal cycle amplitudes.

In this paper, we continue the evaluation of the GOSAT seasonal cycle from Kulawik et al. (2015). Five years of GOSAT observations and the updated TCCON GGG2014 retrievals lengthen the co-located time series sufficiently to evaluate the seasonal cycles regionally at 12 TCCON sites in the Northern Hemisphere. We extend the seasonal cycle analysis to four other retrieval algorithms to identify potential biases characteristic to the ACOS retrievals. Although the emphasis of the study is on these TCCON comparisons, we also compare the GOSAT seasonal cycle against models that assimilate in-situ data; because of their connection to measurements, models may be a reasonable representation of the truth in areas with high assimilated data density, such as North America or Western Europe. This seasonal cycle evaluation study lays important ground work to the analysis of OCO-2 observations that also use the ACOS retrieval system and are, therefore, likely to be affected by any seasonal biases present in the GOSAT/ACOS retrievals that are due to the ACOS system or ACOS a priori inputs.

2 GOSAT

The Greenhouse Gases Observing Satellite (GOSAT), developed by Japan Aerospace Exploration Agency (JAXA), was launched in January 2009 to make near-global greenhouse gas measurements from a polar orbit (Yokota et al., 2009). GOSAT measures reflected solar near-infrared radiation with a Fourier transform spectrometer (TANSO-FTS; Kuze et al., 2009). The diameter of a GOSAT sounding footprint is approximately 10 km, and the soundings repeat in a three-day cycle. We used GOSAT data taken in two primary modes: glint over oceans, and nadir view over land. Nadir data over land has two gain states: high gain (H) for most of the data, and medium (M) over bright surfaces, such as deserts.

Several retrieval algorithms have been developed for retrieving the column-averaged CO₂ from the GOSAT near-infrared measurements; these algorithms have been recently reviewed and compared by Oshchepkov et al. (2013a) and Reuter et al. (2013). In this paper, we concentrate on the evaluation of the Atmospheric CO₂ Observations from Space build 3.5 (ACOS B3.5) retrieval algorithm (Crisp et al., 2012). The ACOS retrieval algorithm is described in detail by O'Dell et al. (2012). The most significant subsequent updates and improvements to the operational algorithm include updated spectroscopy for the 1.6 μm and 2.1 μm CO₂ absorption bands, moving from static to dynamic vertical pressure levels, an improved prior profile of CO₂, and a complete change in the treatment of aerosol and cloud scattering. Instead of a globally constant aerosol model that was incorporated in

ACOS B3.4 and earlier versions, B3.5 uses Modern-Era Retrospective Analysis for Research and Applications (MERRA) reanalysis data of five aerosol types (mineral dust, sea salt, black carbon, sulphates, and organic carbon) to determine two most common types at a given GOSAT sounding location, and applies their respective optical properties in the retrieval.

3 Validation data

3.1 TCCON

The Total Carbon Column Observing Network (TCCON) is currently composed of 21 operating Fourier transform spectrometers that make ground-based measurements of atmospheric XCO₂ and other gases (Wunch et al., 2011a). Their validated and calibrated higher precision and accuracy compared to satellite observations, coupled with the fact that they measure the same quantity in essentially the same way as the satellites, though looking directly at the sun rather than sunlight reflected off the Earth, so are not affected by surface albedo, make them an ideal, independent validation source for GOSAT (Wunch et al., 2010; Messerschmidt et al., 2011). Though the seasonal cycle of TCCON has itself never been explicitly validated by comparison with aircraft, we implicitly assume that our inferred TCCON seasonal cycles for XCO₂ can be taken as truth, similar to the assumption in several previous studies (Messerschmidt et al., 2011; Keppel-Aleks et al., 2012; Wunch et al., 2013), though in principle sub-ppm seasonal biases could remain. For instance, the TCCON retrieval performs a post-hoc airmass bias correction (Wunch et al., 2011a), errors in which could lead to small but nontrivial differences in the TCCON seasonal cycle. However, it is beyond the scope of this work to validate the accuracy of the TCCON seasonal cycle.

In this study, we used data from all Northern Hemisphere TCCON sites that had 1) at least two years of coincidental measurements with GOSAT; and 2) enough co-located data (see Sect. 4.1) to evaluate a seasonal cycle; i.e., both ACOS and TCCON observations available at the proximity of the site through most seasons. The first criterion eliminated the Four Corners and Caltech/Pasadena sites, while the second eliminated the northernmost sites of Ny Ålesund and Eureka which have very little co-located data due to the high latitude. We did not include the Southern Hemisphere sites because the seasonal changes in XCO₂ at those sites are minor, making the definition of an average seasonal cycle more ambiguous and sensitive to inter-annual variability. We therefore chose to focus on the Northern Hemisphere, which has both a larger seasonal cycle amplitude, and a larger quantity of TCCON stations against which to compare. The sites that were used in this study are shown in Fig. 1. For these sites, we analyzed all co-located data between April 23, 2009, and December 31, 2013. We used the newest available GGG2014 TCCON retrievals for each site: Bialystok (Messerschmidt et al., 2012; Deutscher et al., 2014), Bremen (Notholt et al., 2014), Garmisch (Sussmann and Rettinger, 2014), Izaña (Blumenstock et al., 2014), JPL (Wennberg et al., 2014a), Karlsruhe (Hase et al., 2014), Lamont (Wennberg et al., 2014c), Orleans (Warneke et al., 2014), Park Falls (Washenwelder et al.,

2006; Wennberg et al., 2014b), Saga (Kawakami et al., 2014), Sodankylä (Kivi et al., 2014), and Tsukuba (Ohyama et al., 2009; Morino et al., 2014). TCCON data were obtained from the TCCON Data Archive website at <http://tccon.ornl.gov/>.

160 **3.2 Model CO₂ data**

Because evaluation against TCCON is limited to 12 sites in the Northern Hemisphere, another validation source is necessary for obtaining a more thorough view of the accuracy of the GOSAT seasonal cycle. Therefore, we also analyzed XCO₂ from three models that assimilate in-situ CO₂ measurements to optimize their fluxes. The models were CarbonTracker (CT2013B; Peters et al., 2007, with updates documented at <http://carbontracker.noaa.gov/>), MACC 13.1 (Chevallier et al., 2010, documentation and data available at <http://www.copernicus-atmosphere.eu/catalogue>), and the University of Edinburgh model (UoE; Feng et al., 2009, 2011, <http://www.palmergroup.org>). Relevant model properties are listed in Table 1. The models were resampled at GOSAT/ACOS observations in latitude, longitude and time, and integrated over all atmospheric layers to form the column-averaged CO₂. The ACOS averaging kernel correction was first considered for CT2013B, but as it had only a very minor effect on the total column (generally < 0.1 ppm difference in monthly averages), it was subsequently neglected for all models. However, seasonal effects of the averaging kernel correction are briefly assessed in Sect. 5.3. All model results were available from the beginning of GOSAT data (April 23, 2009) but have different end dates: UoE and CT2013B run until the end of December 2012, and MACC 13.1 is available until the end of December 2013.

4 Methods

In this section, we describe the co-location of ground-based and satellite remote sensing measurements, filtering and bias correction for GOSAT/ACOS, and the averaging kernel correction, and define the average seasonal cycle. We demonstrate these steps with an example TCCON site at Park Falls, Wisconsin, U.S.

4.1 Co-locating GOSAT and TCCON

ACOS retrievals of GOSAT soundings are estimates of total column XCO₂. Therefore, the issue of co-locating GOSAT soundings with TCCON soundings boils down to the question of whether we expect both sounding locations to have the same atmospheric XCO₂. Any co-location technique is an assumption about the geographical region over which we expect XCO₂ to be the same as a TCCON retrieval, within some tolerance. For example, a geometrical co-location criterion, where we consider all GOSAT soundings within some fixed distance of a TCCON station, assumes that in the real atmosphere the variation of XCO₂ over that distance is smaller than said tolerance. Similarly, co-locating using the 700 hPa potential temperature (Wunch et al., 2011b) assumes that air with the

190 same transport history – in so far as it is reflected in the 700 hPa potential temperature – will have the same XCO₂ (within said tolerance). However, neither of these co-location techniques account for the fact that ultimately atmospheric XCO₂ is a convolution of surface fluxes and transport. Therefore, in our paper we have applied the NOAA/Basu co-location technique (Guerlet et al., 2013a) which uses a modelled atmospheric XCO₂ field to delineate the region around a TCCON station over
195 which we expect XCO₂ to be constant within some tolerance (0.5 ppm). Since the model is run with realistic surface fluxes and atmospheric transport, we expect this co-location technique to account for XCO₂ variations due to both. To set upper spatiotemporal limits for the co-located soundings, the GOSAT soundings were required to be within $\pm 22.5^\circ$ in longitude and $\pm 7.5^\circ$ in latitude from the TCCON site, and acquired on the same day, within 2 hours of each other. The TCCON soundings
200 were interpolated to local noon to exclude any effects from the diurnal cycle of XCO₂. In practice, the NOAA/Basu co-location technique has several advantages: high co-location data volume, good accuracy, and good sampling of parameter space, such as surface albedo. It should also be noted that the performance of this technique does not depend on the absolute accuracy of simulated XCO₂; all that is required is for the spatial gradient of three day average XCO₂ over a few thousand kilometers
205 to be correct to within some tolerance.

The NOAA/Basu co-location technique is visually demonstrated for the Park Falls TCCON site in Fig. 2a. All GOSAT soundings over almost five years of co-located observations at Park Falls are mapped in Fig. 2b, which shows that the exact locations of the co-located GOSAT soundings are to a minor extent dependent on the season.

210 4.2 Data processing

We used GOSAT/ACOS B3.5 level 2 data, which has been pre-filtered and cloud-screened (O’Dell et al., 2012; Taylor et al., 2012). All available ACOS soundings (land H and M gain, ocean glint) were used at each site, but for the northern mid-latitude sites, most, if not all, data were land gain H soundings (see Table 3). After the co-location, the ACOS soundings were filtered using a post-
215 processing filter that removed bad data, such as data from poor spectral fits or containing larger amounts of aerosols, from the soundings. In total, filtering removed 47% of the H gain over land, 45% of M gain over land, and 40% of glint soundings that had been co-located with the TCCON sites considered in this study. An example of the effect of post-processing filtering is shown in Fig. 3, in the upper panels.

220 We also corrected for the known retrieval biases via a multi-parameter linear regression similar to Wunch et al. (2011b) but optimized for B3.5. The optimization is done with respect to all TCCON data and an average of eight inversion-based models. Model results are used for bias correction only when the models agree with each other to within 1 ppm of the total XCO₂ for a given sounding. The bias correction algorithm performed a correction to the retrieved XCO₂ based on different parame-

225 ters. Bias correction is optimized globally, not regionally, but separately for land (nadir, gains H and M) and ocean (glint) soundings.

When comparing two different remote-sensing measurements, the results are not comparable before the difference due to the retrieval averaging kernels has been considered (Rodgers and Connor, 2003). Since the averaging kernels of TCCON and ACOS are quite similar, it was sufficient to follow 230 the correction introduced by Wunch et al. (2011b), and further implemented in Nguyen et al. (2014). The effects of the averaging kernel correction for TCCON and bias correction for GOSAT/ACOS soundings are presented in Fig. 3, in the lower left panel. For model results, the averaging kernel corrections were not applied.

Finally, we calculated daily averages of co-located GOSAT/ACOS and TCCON retrievals. This 235 way, days with multiple soundings are not more dominant in the seasonal cycle fit than the days with fewer soundings. Time series of daily averages are shown in Fig. 3, in the lower right panel.

4.3 Seasonal cycle

In what follows, we parameterize the seasonal cycle of XCO_2 as a skewed sine wave with an upward trend, and find that it is generally a good model for the time series of XCO_2 in the Northern 240 Hemisphere. We fitted an average seasonal cycle to the daily XCO_2 averages using the following six-parameter function

$$f(t) = a_0 + a_1 t + a_2 \sin(\omega[t - a_3] + \cos^{-1}[a_4 \cos(\omega[t - a_5])]), \quad (1)$$

where t is the time in days and $\omega = 2\pi/T$, where T is 365 days. The first two terms with the parameters a_0 and a_1 (denoting the average growth rate) fit for a linear trend, and the third term, a 245 sine wave with a time-dependent phase, fits for the seasonal cycle parameters $a_2 - a_5$. As an example, we give the parameters for both TCCON and ACOS fits at Park Falls in Table 2. In particular, $2|a_2|$ denotes the peak-to-peak amplitude of the sine wave and is, from here forwards, used to define the seasonal cycle amplitude. The nonlinear least squares fit was solved using a standard gradient-expansion algorithm. For Park Falls, the seasonal cycle fits for TCCON and ACOS are shown in 250 Fig. 3, lower right panel, and the resulting seasonal cycle amplitude is 8.4 ± 0.1 ppm for TCCON, and 8.6 ± 0.2 ppm for ACOS. The errors of the fitted parameters are driven by the standard deviations σ of each daily XCO_2 , initially requiring $\sigma_{ACOS} \geq 1.5$ ppm and $\sigma_{TCCON} \geq 0.3$ ppm. Because the true errors in daily-averaged XCO_2 are not well known, we scaled the σ of each daily-averaged XCO_2 by multiplying them with the minimized quantity χ to yield $\chi^2 = 1$ from the least squares 255 fit. For TCCON data fits, the original χ^2 values varied between $2 < \chi^2 < 10$, while for ACOS, the values were typically $\chi^2 < 1$, which implies that the initial errors σ_{TCCON} may have been underestimated and σ_{ACOS} overestimated. The fitting errors are purely statistical, and do not take into account systematic errors in the data. A more traditional Fourier series fit with an annual and semi-annual cycle (Wunch et al., 2013) was also tried, and the fitted seasonal cycle amplitudes were virtually

260 identical (well within the fitting errors), but because some strange behavior during unobserved times
of year could result, we opted for the fit in Eq. (1). To ensure that the amplitude and phase of the sea-
sonal cycle were not determined largely by the fit function, we assessed the fit-minus-data residuals
for both TCCON and ACOS, and could not identify any systematic signatures in the residuals.

We recognize that there could be inter-annual variability in some or all of the fitted parameters,
265 and that our results can be affected by that variability; especially we can expect sites with shorter co-
located time series to be more sensitive. However, we do not fit for inter-annual variability because
we are interested in identifying potential systematic errors in the average seasonal cycle captured by
GOSAT and, in particular, the ACOS retrieval system. For the purposes of evaluating the average
seasonal cycle of XCO₂, it is important to compare observations from the same time interval, which
270 we take into account by co-locating the observations from TCCON and GOSAT.

5 Results and discussion

5.1 Evaluation against TCCON

Seasonal cycles for co-located TCCON and GOSAT/ACOS B3.5 XCO₂ soundings were studied at
12 TCCON sites in the Northern Hemisphere. Detrended average seasonal cycles for both retrievals
275 at each site are shown in Fig. 4. Detrending removed a linear trend, i.e. XCO₂ average growth rate,
that varied between 1.88–2.39 ppm/year for ACOS and 2.03–2.58 ppm/year for TCCON retrievals,
depending on the site. We estimated the sensitivity of the average seasonal cycle parameters of
Eq. (1) to the fitted trend from the error covariance matrix associated to the best-fit parameters. The
error in the trend was generally weakly negatively correlated with the error in the seasonal cycle
280 amplitude, for both TCCON and ACOS. The phase-related parameters $a_3 - a_5$ were not correlated
with the trend. Therefore, the error from removing the trend should statistically have little effect on
the parameters of the average seasonal cycle. Descriptive fit parameters together with the associated
errors are collected in Table 3. Instead of showing the fitted values for the three parameters $a_3 - a_5$
of the phase term in Eq. (1), the average dates of annual maximum and minimum XCO₂ are listed.

285 The global average growth rate in CO₂ is accurately captured by long-term ground-based mea-
surements of CO₂ concentration, such as the Mauna Loa record (Keeling et al., 1976). Global annual
trends for the years 2009–2013 varied between 1.66 ppm/year and 2.53 ppm/year (Ed Dlugokencky
and Pieter Tans, NOAA/ESRL, www.esrl.noaa.gov/gmd/ccgg/trends/, 30.3.2015). The accuracy of
the TCCON-inferred regional XCO₂ growth rates is not precisely known, though agreement of 0.1–
290 0.2 ppm/year in the global growth rate has been obtained via assimilation of TCCON data in an in-
verse modeling framework (Chevallier et al., 2011). According to Table 3, GOSAT shows a slightly
lower XCO₂ growth rate than TCCON at many validation sites, of order 0.2 ppm/year (around 10%).
Only at JPL, the trend fitted for GOSAT is modestly larger than that of TCCON. There are several
explanations for this. Firstly, GOSAT showing a generally lower trend than TCCON is not surpris-

295 ing but rather a sign of a potentially inaccurate correction for radiometric degradation that is caused
by minor contamination of the instrument over time (Kuze et al., 2014). Secondly, time series of a
little over 2 years of co-located data (like those of Saga, JPL, and Tsukuba) are arguably too short to
distinguish a trend from inter-annual variability. However, the trend captured by GOSAT may be of
minor significance compared to its measurements of the seasonal cycle: errors in capturing the trend
300 may result in errors of the order of a few tenths-of-a-ppm while errors in capturing the seasonal cycle
may have a more significant impact, though this will depend on the detailed set-up of each inverse
modeling system.

The phase of the seasonal cycle is relatively well captured by GOSAT/ACOS. The timing of
the (detrended) maximum concentration varies from March 8 to May 16 for TCCON, and from
305 March 27 to May 21 for GOSAT. The satellite observes the maximum later than the TCCON at the
European sites, but obtains good agreement elsewhere. At the European sites, the difference extends
up to 2–3 weeks, and is likely connected with the biased amplitude inferred by ACOS discussed
below. While the maximum occurs within two spring months depending on location, the minimum is
more seasonally restricted, varying from August 15 to September 27 for TCCON, and from August
310 14 to September 25 for GOSAT. During the minimum, the Northern Hemisphere receives solar
light abundantly and is not snow-covered, so the number of co-located soundings is larger and the
minimum is well captured by the satellite, within 6 days from TCCON, except for Tsukuba and
Bremen. These values are generally in good agreement (within a few days) with Wunch et al. (2013,
p. 9451), except for the TCCON seasonal cycle maximum date at the European sites Bialystok and
315 Bremen. However, regarding the difference in the dates of the maximum, Kulawik et al. (2015)
found a much smaller phase difference in Europe by calculating cross-correlation of the data points
to determine the phase shift. Because our results were based on the fitted seasonal cycles instead of
the actual data, we evaluated the statistical errors of the dates of the maximum and minimum XCO_2
with a Monte Carlo approach, using the error covariance matrices associated with the fitted function
320 parameters. The deviations from the fit maximum and minimum followed a normal distribution with
an average σ of 3.5 days for the TCCON maximum date, and 6.1 days for ACOS maximum date,
reflecting a notable uncertainty in the fitted phase and thus explaining at least partially the difference
between our results and those of Kulawik et al. (2015). The corresponding average σ for the date of
the minimum were 2.2 days (TCCON) and 3.6 days (ACOS).

325 The seasonal cycle amplitudes are presented in Fig. 5a, in addition to Table 3. The amplitude is
captured within the error bars of the regression at four sites: Izaña, Lamont, Saga, and Park Falls.
The largest absolute differences are 1.6 ppm at Tsukuba and 1.1 ppm at Bremen and Orleans, which
are also the largest relative differences (28%, 14% and 15%). Within 1.0 ppm difference, the ampli-
tude is captured at all other sites. It should be noted that Tsukuba only has data for two years and
330 therefore substantial uncertainty in both the trend and amplitude, whereas the Bremen and Orleans
sites have sufficient data for evaluating an average seasonal cycle. A closer inspection of Figs. 4 and

5a reveals that the amplitude seen by GOSAT/ACOS is systematically shallower than TCCON at all five TCCON sites in continental Europe. This bias appears to be regionally very concentrated, because at the Northern European site Sodankylä, GOSAT captures the seasonal cycle reasonably well (within 0.8 ppm), considering the site suffers from data (and sunlight) deficiency in winter. Kulawik et al. (2015) noted the low bias as well, although they grouped all TCCON sites within latitudes 46 – 53°N together and found that, at this latitude range, the seasonal cycle of ACOS was biased low by 0.7 ± 0.7 ppm.

We explored several possible explanations for the low-biased seasonal cycle amplitude over continental Europe. First, we repeated the analysis using GOSAT/ACOS B3.4 retrievals (instead of B3.5), which have two constant aerosol types in the retrieval, different filtering, and bias correction. This did not have a systematic effect: the seasonal cycle amplitude of GOSAT increased at Bremen (+0.3 ppm) and Orleans (+0.5 ppm), and decreased at Bialystok (–0.2 ppm), Garmisch (–0.2 ppm), and Karlsruhe (–0.4 ppm).

Next, we introduced variations to the co-location method to quantify its impact to the seasonal cycle amplitude. Our default co-location technique was the NOAA/Basu method with 0.50 ppm CO₂ gradient, maximum latitude difference 7.5°, and longitude 22.5°. We experimented with four modifications to it: 1) latitude 5.0°, longitude 15°, 2) latitude 2.5°, longitude 7.5°, 3) 0.25 ppm CO₂ gradient, and 4) 1.0 ppm CO₂ gradient. The latter increased the number of co-located points while the three former reduced it by making the co-location requirement stricter. We found that a smaller longitude-latitude box and a tighter CO₂ gradient led to a better match-up in terms of the seasonal cycle amplitude at Bialystok (difference only 0.1 ppm), but not in other European sites where the difference either did not change or increased. The ACOS seasonal cycle amplitude at Garmisch site turned out to be highly dependent on the co-location details, varying from 5.0 ppm to 5.9 ppm in these tests. The TCCON amplitudes changed typically only 0.1 ppm, but the fitting errors increased as the number of co-located soundings decreased. We also found that the co-location box dimensions had an impact on the seasonal cycle at JPL, which is located in the Los Angeles basin where large CO₂ gradients could be expected. With the default technique, the amplitude for ACOS was 0.5 ppm shallower than TCCON (10% difference), but when decreasing the box size, the difference was reduced to 0.1 ppm (2%).

In our last experiment, we tested the impact of the ACOS B3.5 bias correction for H gain over land; as Table 3 shows, all co-located soundings at the continental European sites were land gain H. We found that the bias correction increased the seasonal cycle amplitude at Park Falls by 1.4 ppm, mostly due to a correction for dust aerosol optical depth and surface albedo in the 2.1 μm band, but the bias correction had only a 0.1 ppm total impact on the amplitude at the European sites. It turned out that two of the bias correction parameters (related to the retrieved surface pressure and vertical CO₂ gradient) made the seasonal cycle over Europe consistently shallower by 0.3 – 0.4 ppm, depending on the site (see Fig. 5b). However, these parameters did not affect the seasonal cycle amplitude at

Park Falls or Lamont, which are the two main sites used when optimizing the ACOS bias correction.

370 An interesting finding is that removing these two terms from the bias correction made the ACOS seasonal cycle amplitude (Fig. 5b) and trend (not shown) agree better with TCCON at 10 of the 12 sites, even though it made the scatter worse in single-sounding statistics. This implies that the bias correction might be improved by designing it based on aggregated soundings in addition to single observation statistics.

375 **5.2 Evaluation against other retrieval algorithms**

To further study the discrepancies of GOSAT and TCCON, we repeated the seasonal cycle analysis for four other retrieval algorithms, taking into account their individual bias corrections: RemoTeC v2.35 (Butz et al., 2011; Guerlet et al., 2013a), University of Leicester (UoL) v5.1 (Cogan et al., 2012), NIES PPDF-S v.02.11 (Oshchepkov et al., 2013b), and NIES v02.21 (Yoshida et al., 2013),
380 which is the operational GOSAT retrieval algorithm with the bias correction applied. The seasonal cycle amplitude, the trend, and the days of maximum and minimum (detrended) XCO₂ are presented in Fig. 6 together with their daily averages RMS error with respect to the TCCON fit. RemoTeC had a shorter time series than the other retrievals, and was therefore not included in the Saga, JPL, and Tsukuba results. UoL data did not include glint soundings, which may cause some differences at
385 coastal or island sites. Also, only ACOS and NIES retrievals included a sufficient amount of co-located soundings for successfully fitting a seasonal cycle at Sodankylä.

Overall, the five algorithms performed qualitatively similarly but show notable scatter at most validation sites and in most of the fitted parameters. Also, no algorithm clearly outperforms another. The only systematic difference is that all algorithms except NIES generally capture a smaller mean
390 growth rate than TCCON, whereas NIES retrieves a higher trend. This may be due to different corrections for radiometric degradation in the different algorithms, but could also result from other factors, such as bias correction. For example, NIES v02.21 and NIES PPDF-S v.02.11 have different growth rates despite the use of similar corrections for radiometric degradation. The TCCON seasonal cycle amplitude is captured by GOSAT at almost every site but by a different retrieval: as shown in
395 Sect. 5.1, ACOS has a very good agreement with TCCON at the North American sites as well as Izaña and Saga but, in continental Europe, NIES and NIES PPDF-S perform generally the best. ACOS, RemoTeC, and UoL all show a low-biased amplitude in continental Europe, and NIES, UoL, and NIES PPDF-S are biased high elsewhere. If considering only those sites with longer time series, the scatter between the algorithms is around 1 ppm.

400 The maximum and minimum days of the seasonal cycle reflect the drawdown season and are dependent on latitude and climate region. Both TCCON and GOSAT capture an earlier start of drawdown at the continental European sites compared to the other sites, the latest start being at the southernmost site, Izaña. The ACOS and NIES PPDF-S algorithms appear to be generally best in phase with TCCON regarding the date of maximum XCO₂. At the continental European sites,

405 GOSAT and TCCON fits for the maximum day differ by several weeks, TCCON being systematically earlier. The minimum is better captured by all retrievals, with the spread varying from a few days to about 20 days; the performance of the individual algorithms is very site-specific.

Since none of the retrieval algorithms clearly outperformed the others at every TCCON site, we repeated the analysis for the ensemble median algorithm EMMA (Reuter et al., 2013), which combines all individual retrievals into one data set of median XCO₂ values. Even though EMMA had the 410 smallest RMS error at four TCCON sites overall, it did not perform systematically better or worse than the individual retrieval algorithms in capturing the seasonal cycle of XCO₂.

5.3 Evaluation against models

The seasonal cycle amplitude of GOSAT/ACOS B3.5 was also compared to the inverse model systems 415 MACC 13.1, CT2013B, and UoE in the Northern Hemisphere. As described in Sect. 3.2, these models have been optimized against assimilated flask and in-situ CO₂ measurements, though not exactly same data sets nor using the exact same weighting. For the comparison, latitudes from 0° to 70° were divided into 5° latitude bins (see Fig. 1 for the map), and the GOSAT/ACOS soundings within one latitude bin were collected into a single time series. The seasonal cycle was fitted 420 on the daily averages of GOSAT/ACOS XCO₂ and the resampled models. The resulting seasonal cycle amplitudes are shown in Fig. 7. The amplitude increases significantly from the tropics towards high latitudes for both GOSAT and the models. Although the results are qualitatively similar, the models can show close to 2 ppm differences within latitude bands. ACOS is in excellent agreement to MACC from 0°N to 50°N, whereas CT2013B and UoE have a shallower seasonal cycle from 425 the tropics up to 35°N. Differences in the model seasonal cycle can be caused by a number of error sources, including their prior, transport, and inversion. Tropical and subtropical latitudes include large regions where the data constraint is weaker; therefore, the land surface prior (and its particular implementation) may impact the inversion results more than at those regions where the measurement network is dense. Both UoE and CT2013B use a variant of CASA as their biospheric 430 flux model, as presented in Table 1 (in fact, CT2013B uses a unique combination of two flavors of CASA (Andy Jacobson, personal communication, April 17, 2015)). Even though different versions of CASA can differ in their seasonal cycle magnitude, our results may imply that the seasonal cycle of CASA fluxes is too shallow in some tropical regions or biomes. We first did the comparison using earlier versions of CarbonTracker (CT2011 and CT2013), and found that CarbonTracker and UoE 435 results were nearly identical in these regions (see CT2013 and UoE in Figs. 7 and 8), which was surprising because the two models were different in every aspect (transport, in-situ data selection, inversion) except for their prior biospheric fluxes. However, a significant correction to the transport model's vertical mixing was introduced in CT2013B. This led to an increase of about 0.5 ppm in the CarbonTracker's seasonal cycle amplitude at all latitudes.

440 At 50 – 60°N in Fig. 7, ACOS agrees better with UoE and CT2013B. From 60° to 70°, ACOS
has a higher seasonal cycle amplitude than most models. A similar result was also obtained by Be-
likov et al. (2014) using GOSAT/NIES v02.00 retrievals, NIES transport model, and LMDZ model.
However, at high boreal latitudes, the satellite observations are associated with larger errors that are
not reflected in the purely statistical fitting errors. ACOS results at these latitudes should therefore
445 be interpreted with caution.

We tested how the ACOS bias correction and model averaging kernel correction affected the
latitudinally averaged seasonal cycle amplitudes. The ACOS bias correction decreased the amplitude
about 0.5 ppm at latitudes 10 – 40°N, but increased the amplitude at 40 – 70°N. The maximum
increase was 1.0 ppm at latitudes 50 – 60°N, implying that before the bias correction, ACOS was in
450 better agreement with MACC at these latitudes, but that after the bias correction, ACOS agreed better
with UoE and CT2013B. Even though validation against models is part of the ACOS bias correction,
the TCCON sites are likely to dominate the bias correction at mid-latitudes. We studied the potential
seasonal impact of the averaging kernel correction for CT2013B. We found that the averaging kernel
correction systematically decreased the model seasonal cycle amplitude in the Northern Hemisphere
455 by 0.15 ppm on average. Overall, these changes are minor and do not affect our general conclusions
about the model comparisons.

The latitudinal dependence of the CO₂ seasonal cycle amplitude has been previously shown in e.g.
"the flying carpet" plot presented by Conway et al. (1994, Fig. 4), but we would like to emphasize
that the amplitude can also depend on longitude. Especially in the mid-latitudes, its increase from
460 west to east is notable; this is demonstrated in Fig. 8 for latitude band 45 – 50°N, where the seasonal
cycle amplitude of GOSAT/ACOS is 6.4 ppm over the longitudes 180W–120W, and is doubled at
120E–180E. The increased seasonal cycle is likely due to the large seasonal sink of the boreal forests,
accrued in the total column as the observation point is moved eastward. These GOSAT observations
considered were taken over land, so in practice, this means that the seasonal cycle amplitude is
465 dampened from the Eastern Asia over the North Pacific Ocean to the North-West United States. In
the lower troposphere, this dampening above 30°N latitude was shown by Nakazawa et al. (1992)
who analyzed a three-year time series (1984–1986) of CO₂ measurements onboard container ships.
The model results in Fig. 8 show a similar pattern of amplitude enhancement towards east, albeit
the seasonal cycle amplitude of MACC is 2–3 ppm shallower compared to those of the other models
470 and ACOS in the Eastern Asia. Despite this large discrepancy in the east where the data volume
is small (see Fig. 8, right vertical axis), the zonally-averaged seasonal cycle amplitudes of MACC
and ACOS agree within 0.1 ppm at the same latitude band (45 – 50°N). The CT2013B amplitudes
are consistently higher than ACOS at all longitudes in Fig. 8, but they agree within 0.1 ppm in the
Eastern Asia. Of the three models, UoE is most consistent with ACOS, agreeing about the seasonal
475 cycle amplitude to within 1 ppm at these specific regions. The northern and mid-latitudinal regions

of Asia are again regions where the in-situ measurement coverage is very limited, which explains the large spread between the individual model results.

6 Conclusions

The seasonal cycle of XCO₂ is profoundly connected to the biospheric fluxes that determine the global terrestrial net CO₂ sink. Satellite measurements of XCO₂ by the Greenhouse Gases Observing Satellite (GOSAT) and the Orbiting Carbon Observatory (OCO-2) expand the current in-situ measurement network tremendously and therefore have the potential to improve flux inversions. However, the satellite-measured seasonal cycle of XCO₂ can be affected by different retrieval biases, such as biases related to seasonally-varying parameters (e.g., surface albedo) and a sampling bias due to the seasonal variation in solar radiation. Mischaracterization of the seasonal cycle could lead to errors in the inverse model systems that assimilate satellite CO₂ data. Motivated by this, we evaluated the seasonal cycle of GOSAT observations using ACOS B3.5 retrievals from years 2009–2013.

Three independent approaches were used for the evaluation of the XCO₂ seasonal cycle: comparisons against the Total Carbon Column Observing Network (TCCON), other GOSAT retrievals (UoL v5.1, NIES v02.21, NIES PPDF-S v.02.11, and RemoTeC v2.35), and comparisons to optimized inversion models that assimilate in-situ measurements of CO₂. We found that ACOS captures the seasonal cycle amplitude of TCCON with an accuracy of better than 1.0 ppm at most of the 12 TCCON sites in the Northern Hemisphere considered in this study. As we also inferred the mean annual growth rate at each TCCON site in order to remove it, we found agreement of generally better than 0.2 ppm/year in this quantity, with the ACOS-inferred growth rate most often being lower than TCCON. Over continental Europe, the seasonal cycle amplitude as measured by ACOS was biased low at all five sites, the largest difference being 1.1 ppm at Bremen and Orleans. We also found that ACOS generally captured the seasonal cycle phase within a few days, except over Europe where the differences were 2–3 weeks, with ACOS measuring the date of maximum XCO₂ later than TCCON. Several other algorithms also had minor low biases in their seasonal cycle amplitudes over Europe. We explored the cause of the low bias for ACOS, and found that the bias correction parameters related to the retrieved surface pressure and vertical CO₂ gradient were partially responsible, explaining 16 – 48% of the difference. This suggests that the bias correction might benefit from considering aggregated soundings in addition to deviations at single-sounding level. Also, the selection of the co-located soundings was found to affect the seasonal cycle amplitude at few sites. Especially at JPL, which is in the Los Angeles basin, the agreement with TCCON improved notably when the co-location criteria were made sufficiently tight to not include soundings taken too far from the basin.

510 Model comparisons at latitudes 0–70°N revealed that qualitatively the models and satellite obser-
vations agreed well, but also that the model-to-model differences were (at most latitude bands stud-
ied) larger than model-to-ACOS differences. From the tropics up to 50°N, the zonally-averaged sea-
sonal cycle amplitude of ACOS was in very good agreement with MACC 13.1, while between 50 –
60°N, ACOS agreed better with the University of Edinburgh model and CarbonTracker CT2013B.
515 Both of the latter models had seasonal cycle amplitudes shallower than ACOS or MACC at tropical
and subtropical latitudes, where the models lack direct constraints from measurements over land and
are thus more affected by their prior fluxes (or by extra-tropical or ocean measurements through
long-range transport). Therefore, the shallower seasonal cycle amplitude might be connected to their
prior land surface models that are different variants of CASA. However, to verify this, one should
520 investigate also the impact of transport, data assimilation, and inversion system differences. We also
found that the longitudinal changes in the seasonal cycle amplitude at mid-latitudes can be notable.
In particular, we showed that at 45–50°N latitudes, the amplitude of the GOSAT XCO₂ seasonal cy-
cle doubles from the North-West U.S. to Eastern Asia. The model results showed a gradient as well,
although it was 1–3 ppm shallower, depending on the model. We also noticed that the averaging
525 kernel correction can systematically decrease the seasonal cycle amplitude by up to 0.2 ppm.

Based on our study, the GOSAT/ACOS seasonal cycle error is of the order of 1.0 ppm near
TCCON stations and likely to be of this size in other parts of the world, though may be influenced
by the a priori accuracy of jointly retrieved parameters, such as those related to aerosols. As model-
to-model differences in the XCO₂ seasonal cycle amplitude can be several ppm at regions poorly
530 sampled by in-situ measurements, GOSAT observations could potentially be used directly (without
elaborate inversions) to evaluate model differences at these regions. This idea is explored in more
detail in a work under preparation (Lindqvist et al., 2015, in prep.).

Acknowledgements. HL and CO wish to acknowledge support from the NASA Jet Propulsion Laboratory and
the OCO-2 project, via JPL subcontract number 1439002. HL also acknowledges funding from the Academy
535 of Finland, via project number 285421. HB and RP acknowledge the National Centre for Earth Observation
NCEO and the ESA GHG-CCI project. FC acknowledges the European Commission (grant agreement No.
630080, MACC III). We wish to thank the referees (Susan Kulawik and an anonymous referee) and Andy
Jacobson for their comments and constructive feedback to the manuscript. We also gratefully acknowledge all
the data providers. TCCON data were obtained from the TCCON Data Archive, operated by the California
540 Institute of Technology from the website at <http://tcon.ornl.gov/>. TCCON work at Garmisch has been funded
by the ESA GHG-cci project via subcontract with the University of Bremen and by the EC within the INGOS
project. CarbonTracker CT2013B results were provided by NOAA ESRL, Boulder, Colorado, USA from the
website at <http://carbontracker.noaa.gov>. Finally, we thank our colleagues for the use of their GOSAT retrievals:
André Butz and Otto Hasekamp for RemoTeC v2.35, and Andrey Bril for NIES PPDF-S v.02.11.

545 References

- Belikov, D. A., Bril, A., Maksuytov, S., Oshchepkov, S., Saeki, T., Takagi, H., Yoshida, Y., Ganshin, A., Zhuravlev, R., Aoki, S., and Yokota, T.: Column-averaged CO₂ concentrations in the subarctic from GOSAT retrievals and NIES transport model simulations, *Polar Science*, 8, 129–145, 2014.
- Blumenstock, T., Hase, F., Schneider, M., García, O. E., and Sepúlveda, E.: TCCON data from Izaña, Tenerife, Spain, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. 550 <http://dx.doi.org/10.14291/tcon.ggg2014.izana01.R0/1149295>, 2014.
- Notholt, J., Petri, C., Warneke, T., Deutscher, N., Buschmann, M., Weinzierl, C., Macatangay, R., Grupe, P.: TCCON data from Bremen, Germany, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. 555 <http://dx.doi.org/10.14291/tcon.ggg2014.bremen01.R0/1149275>, 2014.
- Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante R., Bergamaschi, P., Blumenstock, T., Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J. P., Butz, A., Chedin, A., Chevallier, F., Crevoisier, C. D., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., Kaminski, T., Laeng, A., Lichtenberg, G., De Maziere, M., Noel, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G. P., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D. W. T., Kuze, A., O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., and Yoshida, Y.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, *Remote Sensing of Environment*, 560 doi:10.1016/j.rse.2013.04.024, pp. 19, (in press), 2013.
- Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophys. Res. Lett.*, 38, L14812, doi:10.1029/2011GL047888, 2011.
- 570 Chevallier, F., Breon, F.-M., and Rayner, P. J.: Contribution of the Orbiting Carbon Observatory to the estimation of CO₂ sources and sinks: Theoretical study in a variational data assimilation framework, *J. Geophys. Res.*, 112, D09307, doi:10.1029/2006JD007375, 2007.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröhlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morgui, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.: CO₂ surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *J. Geophys. Res.*, 115, D21307, doi:10.1029/2010JD013887, 2010.
- 575 Chevallier, F., Deutscher, N. M., Conway, T. J., Ciais, P., Ciattaglia, L., Dohe, S., Fröhlich, M., Gomez-Pelaez, A. J., Griffith, D., Hase, F., Haszpra, L., Krummer, P., Kyrö, E., Labuschagne, C., Langenfelds, R., Machida, T., Maignan, F., Matsueda, H., Morino, I., Notholt, J., Ramonet, M., Sawa, Y., Schmidt, M., Sherlock, V., Steele, P., Strong, K., Sussmann, R., Wennberg, P., Wofsy, S., Worthy, D., Wunch, D., and Zimnoch, M.: Global CO₂ fluxes inferred from surface air-sample measurements and from TCCON retrievals of the CO₂ total column, *Geophys. Res. Lett.*, 38, L24810, doi:10.1029/2011GL049899, 2011.

- 585 Cogan, A. J., Boesch, H., Parker, R. J., Feng, L., Palmer, P. I., Blavier, J.-F. L., Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., Warneke, T., and Wunch, D.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): Comparison with ground-based TCCON observations and GEOS-Chem model calculations, *J. Geophys. Res.*, 117, D21301, doi:10.1029/2012JD018087, 2012.
- Conway, T. J., Tans, P. P., Waterman, L. S., Thoning, K. W., Kitzis, D. R., Masarie, K. A., and Zhang, N.:
590 Evidence for interannual variability of the carbon cycle from the National Oceanic and Atmospheric Administration/Climate Monitoring and Diagnostics Laboratory Global Air Sampling Network, *J. Geophys. Res. Atmos.*, 99, D11, 22831–22855, doi:10.1029/94JD01951, 1994.
- Crisp, D., Atlas, R. M., Breon, F.-M., Brown, L. R., Burrows, J. P., Ciais, P., Connor, B. J., Doney, S. C., Fung, I. Y., Jacob, D. J., Miller, C. E., O'Brien, D., Pawson, S., Randerson, J. T., Rayner, P., Salawitch, R. J., Sander,
595 S. P., Sen, B., Stephens, G. L., Tans, P. P., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Yung, Y. L., Kuang, Z., Chudasama, B., Sprague, G., Weiss, B., Pollock, R., Kenyon, D., and Schroll, S.: The orbiting carbon observatory (OCO) mission, *Advances in Space Research*, 34(4), 700–709, 2004.
- Crisp, D., Fisher, B., O'Dell, C., Frankenberg, C., Basilio, R., Bösch, H., Brown, L. R., Castano, R., Connor, B., Deutscher, N. M., Eldering, A., Griffith, D., Gunson, M., Kuze, A., Mandrake, L., McDuffie, J.,
600 Messerschmidt, J., Miller, C. E., Morino, I., Natraj, V., Notholt, J., O'Brien, D. M., Oyafuso, F., Polonsky, I., Robinson, J., Salawitch, R., Sherlock, V., Smyth, M., Suto, H., Taylor, T. E., Thompson, D. R., Wennberg, P. O., Wunch, D., and Yung, Y. L.: The ACOS CO₂ retrieval algorithm – Part II: Global XCO₂ data characterization, *Atmos. Meas. Tech.*, 5 (4), 687–707, doi:10.5194/amt-5-687-2012, 2012.
- Deutscher, N., Notholt, J., Messerschmidt, J., Weinzierl, C., Warneke, T., Petri, C., Grupe, P., and Katrynski,
605 K.: TCCON data from Bialystok, Poland, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.bialystok01.R0/1149277>, 2014.
- Dils, B., Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Parker, R., Guerlet, S., Aben, I., Blumenstock, T., Burrows, J. P., Butz, A., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., De Maziere, M., Notholt, J., Sussmann, R., Warneke, T., Griffith, D., Sherlock, V., and Wunch, D.:
610 The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, *Atmos. Meas. Tech.*, 7, 1723–1744, 2014.
- Feng, L., Palmer, P. I., Boesch, H., and Dance, S.: Estimating surface CO₂ fluxes from space-borne CO₂
615 dry air mole fraction observations using an ensemble Kalman Filter, *Atmos. Chem. Phys.*, 9, 2619–2633, doi:10.5194/acp-9-2619-2009, 2009.
- Feng, L., Palmer, P. I., Yang, Y., Yantosca, R. M., Kawa, S. R., Paris, J.-D., Matsueda, H., and Machida, T.: Evaluating a 3-D transport model of atmospheric CO₂ using ground-based, aircraft, and space-borne data, *Atmos. Chem. Phys.*, 11, 2789–2803, doi:10.5194/acp-11-2789-2011, 2011.
- 620 Guerlet, S., Basu, S., Butz, A., Krol, M., Hahne, P., Houweling, S., Hasekamp, O. P., and Aben, I.: Reduced carbon uptake during the 2010 Northern Hemisphere summer from GOSAT, *Geophys. Res. Lett.*, 40(10), 2378–2383, doi:10.1002/grl.50402, 2013b.
- Guerlet, S., Butz, A., Schepers, D., Basu, S., Hasekamp, O. P., Kuze, A., Yokota, T., Blavier, J.-F., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kyrö, E., Morino, I., Sherlock, V., Sussmann, R., Galli, A., and Aben,

- 625 I.: Impact of aerosol and thin cirrus on retrieving and validating XCO₂ from GOSAT shortwave infrared measurements, *J. Geophys. Res. Atmos.*, 118, 4887–4905, doi:10.1002/jgrd.50332, 2013a.
- Hase, F., Blumenstock, T., Dohe, S., Groß, J., and Kiel, M.: TCCON data from Karlsruhe, Germany, Release GGG2014R1, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A.
- 630 <http://dx.doi.org/10.14291/tcon.ggg2014.karlsruhe01.R1/1182416>, 2014.
- Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Taura, C., Fukamachi, T., Sakashita, M.: TCCON data from Saga, Japan, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A.
- <http://dx.doi.org/10.14291/tcon.ggg2014.saga01.R0/1149283>, 2014.
- 635 Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdahl, C. A., Guenther, P. R., Waterman, L. S., and Chin, J. F.: Atmospheric carbon dioxide variations at Mauna Loa observatory, Hawaii, *Tellus*, 28(6), 538–551, 1976.
- Keppel-Aleks, G., Wennberg, P. O., and Schneider, T.: Sources of variations in the total column carbon dioxide, *Atmos. Chem. Phys.*, 11, 3581–3593, doi:10.5194/acp-11-3581-2011, 2011.
- Keppel-Aleks, G., Wennberg, P. O., Washenfelder, R. A., Wunch, D., Schneider, T., Toon, G. C., Andres, R. J.,
- 640 Blavier, J.-F., Connor, B., Davis, K. J., Desai, A. R., Messerschmidt, J., Notholt, J., Roehl, C. M., Sherlock, V., Stephens, B. B., Vay, S. A., and Wofsy, S. C.: The imprint of surface fluxes and transport on variations in total column carbon dioxide, *Biogeosciences*, 9(3), 875–891, doi:10.5194/bg-9-875-2012, 2012.
- Kivi, R., Heikkinen, P., and Kyrö, E.: TCCON data from Sodankylä, Finland, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory,
- 645 Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.sodankyla01.R0/1149280>, 2014.
- Kulawik, S. S., Wunch, D., O'Dell, C., Frankenberg, C., Reuter, M., Oda, T., Chevallier, F., Sherlock, V., Buchwitz, M., Osterman, G., Miller, C., Wennberg, P., Griffith, D. W. T., Morino, I., Dubey, M., Deutscher, N. M., Notholt, J., Hase, F., Warneke, T., Sussmann, R., Robinson, J., Strong, K., Schneider, M., and Wolf, J.: Consistent evaluation of GOSAT, SCIAMACHY, CarbonTracker, and MACC through comparisons to
- 650 TCCON, *Atmos. Meas. Tech. Discuss.*, 8, 6217–6277, doi:10.5194/amtd-8-6217-2015, 2015.
- Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the greenhouse gases observing satellite for greenhouse gases monitoring, *Appl. Opt.*, 48, 6716–6733, 2009.
- Kuze, A., Taylor, T. E., Kataoka, F., Bruegge, C. J., Crisp, D., Harada, M., Helmlinger, M., Inoue, M.,
- 655 Kawakami, S., Kikuchi, N., Mitomi, Y., Murooka, J., Naitoh, M., O'Brien, D. M., O'Dell, C. W., Ohyama, H., Pollock, H., Schwandner, F. M., Shiomi, K., Suto, H., Takeda, T., Tanaka, T., Urabe, T., Yokota, T., and Yoshida, Y.: Long-term vicarious calibration of GOSAT short-wave sensors: Techniques for error reduction and new estimates of radiometric degradation factors, *Geoscience and Remote Sensing, IEEE Transactions on*, 52(7), 3991–4004, 2014.
- 660 Lindqvist, H., O'Dell, C., Schuh, A., Baker, D., Cheeseman, M., Baker, I., Haynes, K., and Denning, S.: The potential of GOSAT XCO₂ to evaluate land surface models, In preparation, 2015.
- Liu, J., Bowman, K. W., Lee, M., Henze, D. K., Bousseres, N., Brix, H., Collatz, G. J., Menemenlis, D., Ott, L., Pawson, S., Jones, D., and Nassar, R.: Carbon monitoring system flux estimation and attribution: impact

- of ACOS-GOSAT XCO₂ sampling on the inference of terrestrial biospheric sources and sinks, *Tellus B*, 66, doi:<http://dx.doi.org/10.3402/tellusb.v66.22486>, 2014.
- 665 Maksuytov, S., Takagi, H., Valsala, V. K., Saito, M., Oda, T., Saeki, T., Belikov, D. A., Saito, R., Ito, A., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., and Yokota, T.: Regional CO₂ flux estimates for 2009–2010 based on GOSAT and ground-based CO₂ observations, *Atmos. Chem. Phys.*, 13, 9351–9373, doi:10.5194/acp-13-9351-2013, 2013.
- 670 Messerschmidt, J., Chen, H., Deutscher, N. M., Gerbig, C., Grupe, P., Katrynski, K., Koch, F.-T., Lavrič, J. V., Notholt, J., Rodenbeck, C., Ruhe, W., Warneke, T., and Weinzierl, C.: Automated ground-based remote sensing measurements of greenhouse gases at the Bialystok site in comparison with collocated in situ measurements and model data, *Atmos. Chem. Phys.*, 12(15), 6741–6755, doi:10.5194/acp-12-6741-2012, 2012.
- Messerschmidt, J., Geibel, M. C., Blumenstock, T., Chen, H., Deutscher, N. M., Engel, A., Feist, D. G., Gerbig, 675 C., Gisi, M., Hase, F., Katrynski, K., Kolle, O., Lavrič, J. V., Notholt, J., Palm, M., Ramonet, M., Rettinger, M., Schmidt, M., Sussmann, R., Toon, G. C., Truong, F., Warneke, T., Wennberg, P. O., Wunch, D., and Xueref-Remy, I.: Calibration of TCCON column-averaged CO₂: the first aircraft campaign over European TCCON sites, *Atmos. Chem. Phys.*, 11, 10765–10777, doi:10.5194/acp-11-10765-2011, 2011.
- Morino, I., Matsuzaki, T., Ikegami, H., and Shishime, A.: TCCON data from Tsukuba, Ibaraki, 680 Japan, 125HR, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.tsukuba02.R0/1149301>, 2014.
- Morino, I., Uchino, O., Inoue, M., Yoshida, Y., Yokota, T., Wennberg, P. O., Toon, G. C., Wunch, D., Roehl, C. M., Notholt, J., Warneke, T., Messerschmidt, J., Griffith, D. W. T., Deutscher, N. M., Sherlock, V., Connor, B., 685 Robinson, J., Sussmann, R., and Rettinger, M.: Preliminary validation of column-averaged volume mixing ratios of carbon dioxide and methane retrieved from GOSAT short-wavelength infrared spectra, *Atmos. Meas. Tech.*, 4, 1061–1076, doi:10.5194/amt-4-1061-2011, 2011.
- Nakazawa, T., Murayama, S., Miyashita, K., Aoki, S., and Tanaka, M.: Longitudinally different variations of lower tropospheric carbon dioxide concentrations over the North Pacific Ocean, *Tellus*, 44B, 161–172, 1992.
- 690 Nguyen, H., Osterman, G., Wunch, D., O'Dell, C., Mandrake, L., Wennberg, P., Fisher, B., and Castano, R.: A method for collocating satellite XCO₂ data to ground-based data and its application to ACOS-GOSAT and TCCON, *Atmos. Meas. Tech.*, 7, 2631–2644, doi:10.5194/amt-7-2631-2014, 2014.
- O'Dell, C. W., Connor, B., Bösch, H., O'Brien, D., Frankenberg, C., Castano, R., Christi, M., Eldering, D., Fisher, B., Gunson, M., McDuffie, J., Miller, C. E., Natraj, V., Oyafuso, F., Polonsky, I., Smyth, M., Taylor, T., 695 Toon, G. C., Wennberg, P. O., and Wunch, D.: The ACOS CO₂ retrieval algorithm – Part 1: Description and validation against synthetic observations, *Atmos. Meas. Tech.*, 5(1), 99–121, doi:10.5194/amt-5-99-2012, 2012.
- Ohyama, H., Morino, I., Nagahama, T., Machida, T., Suto, H., Oguma, H., Sawa, Y., Matsueda, H., Sugimoto, N., Nakane, H., and Nakagawa, K.: Column-averaged volume mixing ratio of CO₂ measured with ground-based Fourier transform spectrometer at Tsukuba, *J. Geophys. Res.*, 114, D18303, doi:10.1029/2008JD011465, 2009.
- 700 Olsen, S. C. and Randerson, J. T.: Differences between surface and column atmospheric CO₂ and implications for carbon cycle research, *J. Geophys. Res.*, 109, D02301, doi:10.1029/2003JD003968, 2004.

- Oshchepkov, S., Bril, A., Yokota, T., Wennberg, P. O., Deutscher, N. M., Wunch, D., Toon, G. C., Yoshida, Y., O'Dell, C. W., Crisp, D., Miller, C. E., Frankenberg, C., Butz, A., Aben, I., Guerlet, S., Hasekamp, O., Boesch, H., Cogan, A., Parker, R., Griffith, D., Macatangay, R., Notholt, J., Sussmann, R., Rettinger, M., Sherlock, V., Robinson, J., Kyrö, E., Heikkinen, P., Feist, D. G., Morino, I., Kadyrov, N., Belikov, D., Maksyutov, S., Matsunaga, T., Uchino, O., and Watanabe, H.: Effects of atmospheric light scattering on spectroscopic observations of greenhouse gases from space. Part 2: Algorithm intercomparison in the GOSAT data processing for CO₂ retrievals over TCCON sites, *J. Geophys. Res. Atmos.*, 118(3), 1493–1512, doi:10.1002/jgrd.50146, 2013a.
- Oshchepkov, S., Bril, A., Yokota, T., Yoshida, Y., Blumenstock, T., Deutscher, N. M., Dohe, S., Macatangay, R., Morino, I., Notholt, J., Rettinger, M., Petri, C., Schneider, M., Sussman, R., Uchino, O., Velasco, V., Wunch, D., and Belikov, D.: Simultaneous retrieval of atmospheric CO₂ and light path modification from space-based spectroscopic observations of greenhouse gases: Methodology and application to GOSAT measurements over TCCON sites, *Appl. Opt.*, 52, 1339–1350, 2013b.
- Palmer, P. I., Barkley, M. P., and Monks, P. S.: Interpreting the variability of space-borne CO₂ column-averaged volume mixing ratios over North America using a chemistry transport model, *Atmos. Chem. Phys.*, 8, 5855–5868, 2008.
- Peng, S., Ciais, P., Chevallier, F., Peylin, P., Cadule, P., Sitch, S., Piao, S., Ahlström, A., Huntingford, C., Levy, P., Li, X., Liu, Y., Lomas, M., Poulter, B., Viovy, N., Wang, T., Wang, X., Zaehle, S., Zeng, N., Zhao, F., and Zhao, H.: Benchmarking the seasonal cycle of CO₂ fluxes simulated by terrestrial ecosystem models, *Global Biogeochem. Cycles*, 29, 46–64, doi:10.1002/2014GB004931, 2015.
- Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L. M. P., Petron, G., Hirsch, A. I., Worthy, D. E. J., van der Werf, G. R., Randerson, J. T., Wennberg, P. O., Krol, M. C., and Tans, P. P.: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker, *PNAS*, November 27, 2007, 104(48), 18925–18930, 2007.
- Pitman, A. J.: The evolution of, and revolution in, land surface schemes designed for climate models, *Int. J. Climatol.*, 23, 479–510, doi: 10.1002/joc.893, 2003.
- Rayner, P. J. and O'Brien, D. M.: The utility of remotely sensed CO₂ concentration data in surface source inversions, *Geophys. Res. Lett.*, 28(1), 175–178, 2001.
- Reuter, M., Bösch, H., Bovensmann, H., Bril, A., Buchwitz, M., Butz, A., Burrows, J. P., O'Dell, C. W., Guerlet, S., Hasekamp, O., Heymann, J., Kikuchi, N., Oshchepkov, S., Parker, R., Pfeifer, S., Schneising, O., Yokota, T., and Yoshida, Y.: A joint effort to deliver satellite retrieved atmospheric CO₂ concentrations for surface flux inversions: the ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771–1780, doi:10.5194/acp-13-1771-2013, 2013.
- Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *J. Geophys. Res.*, 108, D3, 4116, doi:10.1029/2002JD002299, 2003.
- Sussmann, R. and Rettinger, M.: TCCON data from Garmisch, Germany, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.garmisch01.R0/1149299>, 2014.
- Takagi, H., Houweling, S., Andres, R. J., Belikov, D., Bril, A., Bösch, H., Butz, A., Guerlet, S., Hasekamp, O., Maksyutov, S., Morino, I., Oda, T., O'Dell, C. W., Oshchepkov, S., Parker, R., Saito, M., Uchino, O., Yokota,

T., Yoshida, Y., and Valsala, V.: Influence of differences in current GOSAT XCO₂ retrievals on surface flux estimation, *Geophys. Res. Lett.*, 41, 2598–2605, doi:10.1002/2013GL059174, 2014.

745 Takagi, H., Sacki, T., Oda, T., Saito, M., Valsala, V., Belikov, D., Saito, R., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., Yokota, T., and Maksuytov, S.: On the benefit of GOSAT observations to the estimation of regional CO₂ fluxes, *SOLA*, 7, 161–164, doi:10.2151/sola.2011-041, 2011.

Taylor, T. E., O'Dell, C. W., O'Brien, D. M., Kikuchi, N., Yokota, T., Nakajima, T. Y., Ishida, H., Crisp, D., and
750 Nakajima, T.: Comparison of cloud-screening methods applied to GOSAT near-infrared spectra, *Geoscience and Remote Sensing, IEEE Transactions on*, 50(1), 295–309, doi:10.1109/TGRS.2011.2160270, 2012.

Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N., Petri, C., Grupe, P., Vuillemin, C., Truong, F., Schmidt, M., Ramonet, M., and Parmentier, E.: TCCON data from Orleans, France, Release GGG2014R0. TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A.
755 <http://dx.doi.org/10.14291/tcon.ggg2014.orleans01.R0/1149276>, 2014.

Washenfelder, R. A., Toon, G. C., Blavier, J.-F. L., Yang, Z., Allen, N. T., Wennberg, P. O., Vay, S. A., Matross, D. M., and Daube, B. C.: Carbon dioxide column abundances at the Wisconsin Tall Tower site, *J. Geophys. Res.*, 111(D22), 1–11, doi:10.1029/2006JD007154, 2006.

760 Wennberg, P. O., Roehl, C., Blavier, J.-F., Wunch, D., Landeros, J., and Allen, N.: TCCON data from Jet Propulsion Laboratory, Pasadena, California, USA, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.jpl02.R0/1149297>, 2014a.

Wennberg, P. O., Roehl, C., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R., Keppel-Aleks, G., Allen, N., Ayers, J.: TCCON data from Park Falls, Wisconsin, USA, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.parkfalls01.R0/1149161>, 2014b.

765 Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., Allen, N., Dowell, P., Teske, K., Martin, C., and Martin, J.: TCCON data from Lamont, Oklahoma, USA, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.lamont01.R0/1149159>, 2014c.

770 Wunch, D., Toon, G. C., Blavier, J. F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The total carbon column observing network, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369(1943), 2087–2112, doi:10.1098/rsta.2010.0240, 2011a.

775 Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jimenez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, *Atmos. Meas. Tech.*, 3, 1351–1362, doi:10.5194/amt-3-1351-2010, 2010.

- Wunch, D., Wennberg, P. O., Messerschmidt, J., Parazoo, N. C., Toon, G. C., Deutscher, N. M., Keppel-Aleks, G., Roehl, C. M., Randerson, J. T., Warneke, T., and Notholt, J.: The covariation of Northern Hemisphere summertime CO₂ with surface temperature in boreal regions, *Atmos. Chem. Phys.*, 13, 9447–9459, doi:10.5194/acp-13-9447-2013, 2013.
- 785 Wunch, D., Wennberg, P. O., Toon, G. C., Connor, B. J., Fisher, B., Osterman, G. B., Frankenberg, C., Mandrake, L., O'Dell, C., Ahonen, P., Biraud, S. C., Castano, R., Cressie, N., Crisp, D., Deutscher, N. M., Eldering, A., Fisher, M. L., Griffith, D. W. T., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyrö, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C. E., Morino, I., Notholt, J., Oyafuso, F. A., Rettinger, M., Robinson, J., Roehl, C. M., Salawitch, R. J., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D. R., Uchino, O., Warneke, T., and Wofsy, S. C.: A method for evaluating bias in global measurements of CO₂ total columns from space, *Atmos. Chem. Phys.*, 11, 12317–12337, doi:10.5194/acp-11-12317-2011, 2011b.
- 790 Yokota, T., Yoshida, Y., Eguchi, N., Ota, Y., Tanaka, T., Watanabe, H., and Maksuytov, S.: Global concentrations of CO₂ and CH₄ retrieved from GOSAT: first preliminary results, *Sola*, 5, 160–163, 2009.
- Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmos. Meas. Tech.*, 6 (6), 1533–1547, 2013.
- 800




Figure 1. Twelve Northern Hemisphere TCCON sites used for GOSAT validation in this study.




Figure 2. (a) An example of the GOSAT-TCCON co-locations using the NOAA/Basu technique (Guerlet et al., 2013a) at Park Falls TCCON station (Wisconsin, USA). All GOSAT/ACOS soundings from 8-11 Aug 2009 are shown with filled circles. The dynamical criterion based on the modelled XCO_2 fields and a 0.5 ppm tolerance from the value at the TCCON location limits the number of co-located satellite soundings (green circles). The soundings marked with yellow symbols did not pass the co-location criteria. (b) All co-located GOSAT/ACOS soundings from Apr 2009 to Dec 2013 at the Park Falls TCCON, coloured according to the month of observation.

Table 1. Models used in the evaluation of the GOSAT seasonal cycle.

Model	Biosphere	Transport	Resolution of the model run (lon x lat x time x layers)
CT2013B	CASA/GFED2 and CASA/GFED3.1	TM5 / ERA-interim, ECMWF	$3^\circ \times 2^\circ \times 3 \text{ h} \times 25$
UoE	CASA/GFED	GEOS-Chem / GEOS5	$5^\circ \times 4^\circ \times 3 \text{ h} \times 47$
MACC 13.1	ORCHIDEE	LMDZ / ECMWF	$3.75^\circ \times 1.9^\circ \times 3 \text{ h} \times 39$




Figure 3. An example of data processing and the seasonal cycle fitting procedure at Park Falls. The upper left panel shows time series of the retrieved XCO₂ for all co-located TCCON (black) and GOSAT/ACOS (pink) soundings. The upper right figure shows only those ACOS L2 soundings that pass the post-processing filters. The lower left figure has bias correction applied for ACOS data and averaging kernel correction considered for TCCON soundings. The lower right panel shows the daily averages of XCO₂ and the respective seasonal cycle fits.

Table 2. Parameters defining the fitted seasonal cycle functions of co-located TCCON and ACOS soundings at Park Falls.

Retrieval	a_0 (ppm)	a_1 (ppm/day)	a_2 (ppm)	a_3 (days)	a_4	a_5 (days)
TCCON	384.5	0.006050	-4.224	-111.4	0.6803	-307.9
ACOS	384.8	0.005904	-4.311	-112.2	0.7585	-268.5




Figure 4. Detrended, best-fit seasonal cycles for GOSAT/ACOS (pink) and TCCON (black) at 12 validation sites in the Northern Hemisphere. The sites are organized according to their latitude (Sodankylä highest, Izaña lowest). The dashed lines depict the times of year with zero or little co-located soundings. On the vertical axis, one tick interval corresponds to 1.0 ppm XCO₂.




Figure 5. Seasonal cycle amplitude for ACOS (vertical axis) and TCCON (horizontal axis) for all the 12 NH sites used in the validation. The dashed black line corresponds to the one-to-one line, and the gray lines denote ± 1.0 ppm. Panel (a) shows the standard bias-corrected ACOS B3.5, and Panel (b) shows ACOS B3.5 with a modified bias correction (see Sect. 5.1 for details).




Figure 6. Comparison of the GOSAT and TCCON XCO₂ time series using the following parameters: root-mean-square (RMS) error (upper left panel), average trend (middle left panel), seasonal cycle amplitude (middle right panel), and the days of maximum and minimum XCO₂ (bottom row). Five retrieval algorithms were included to describe GOSAT observations. TCCON values were based on ACOS B3.5 co-located soundings. The 12 Northern Hemisphere validation sites are shown on the horizontal axis, their latitude increasing from left to right.




Figure 7. Latitudinal dependence of the seasonal cycle amplitude for bias-corrected ACOS B3.5 soundings and for three models resampled at the satellite soundings. For CarbonTracker, we show both CT2013 and CT2013B results, their difference being a major correction in the TM5 transport model. The left vertical axis shows the seasonal cycle amplitude in ppm, while the right vertical axis indicates the number of soundings that fall within each 5° latitude band.




Figure 8. Longitudinal dependence of the seasonal cycle amplitude within the latitude band $45 - 50^\circ\text{N}$. The left vertical axis shows the seasonal cycle amplitude in ppm, while the right vertical axis indicates the number of soundings that fall within each 60° longitude bin. This latitude zone is highlighted in the world map where also the locations of the continents can be seen.

Table 3. Parameters describing the XCO₂ seasonal cycle for TCCON and bias-corrected GOSAT/ACOS B3.5. The fraction of gain H soundings over land is also shown. The validation sites are sorted according to their latitude.

Site	Time series (month/year)	Retrieval	Growth rate (ppm/year)	Amplitude (ppm)	Date of max. XCO ₂	Date of min. XCO ₂	Fraction of land gain H
Izaña	5/2009–10/2013	TCCON	2.41 ± 0.02	5.3 ± 0.1	May 16	Sep 19	12.2%
		GOSAT	2.22 ± 0.04	5.3 ± 0.2	May 18	Sep 17	
Saga	8/2011–10/2013	TCCON	2.39 ± 0.09	6.7 ± 0.2	May 7	Sep 13	77.7%
		GOSAT	1.92 ± 0.26	6.7 ± 0.4	Apr 28	Sep 14	
JPL	5/2011–6/2013	TCCON	2.34 ± 0.07	5.1 ± 0.2	May 2	Sep 27	87.2%
		GOSAT	2.39 ± 0.11	4.6 ± 0.3	May 21	Sep 25	
Tsukuba	8/2011–12/2013	TCCON	2.58 ± 0.10	5.7 ± 0.2	Apr 23	Sep 10	91.9%
		GOSAT	2.20 ± 0.22	7.3 ± 0.5	Apr 23	Aug 26	
Lamont	4/2009–12/2013	TCCON	2.33 ± 0.02	5.3 ± 0.1	May 4	Sep 20	96.5%
		GOSAT	2.14 ± 0.03	5.2 ± 0.1	May 6	Sep 15	
Park Falls	4/2009–12/2013	TCCON	2.21 ± 0.03	8.4 ± 0.1	Apr 22	Aug 15	100%
		GOSAT	2.16 ± 0.04	8.6 ± 0.2	Apr 27	Aug 14	
Garmisch	5/2009–10/2013	TCCON	2.03 ± 0.04	6.6 ± 0.1	Mar 25	Aug 27	100%
		GOSAT	1.90 ± 0.07	5.7 ± 0.2	Apr 17	Aug 24	
Orleans	8/2009–11/2013	TCCON	2.29 ± 0.04	7.3 ± 0.1	Mar 30	Aug 28	100%
		GOSAT	2.04 ± 0.07	6.2 ± 0.3	Apr 13	Aug 22	
Karlsruhe	4/2010–11/2013	TCCON	2.25 ± 0.06	7.3 ± 0.2	Mar 21	Aug 24	100%
		GOSAT	2.05 ± 0.09	6.5 ± 0.2	Mar 27	Aug 27	
Bremen	4/2009–4/2013	TCCON	2.21 ± 0.06	7.7 ± 0.3	Mar 8	Sep 3	100%
		GOSAT	1.88 ± 0.09	6.6 ± 0.3	Apr 10	Aug 24	
Bialystok	4/2009–10/2013	TCCON	2.18 ± 0.03	8.1 ± 0.1	Mar 16	Aug 18	100%
		GOSAT	1.99 ± 0.06	7.5 ± 0.2	Apr 5	Aug 17	
Sodankylä	5/2009–10/2013	TCCON	2.15 ± 0.04	8.7 ± 0.3	Apr 16	Aug 15	100%
		GOSAT	2.05 ± 0.09	9.5 ± 0.5	Apr 24	Aug 17	