

Response to the reviews

In this response, we first list all referee comments (shown with *blue italic text*), answer them and indicate the changes made to the manuscript. Then we list other changes made (mainly updates). A document tracking the changes made to the manuscript published in ACPD has been generated automatically using 'latexdiff', and can be found in the end of this response.

Review #1

The NOAA/Basu coincidence criteria seem to lead to huge footprints! Figure 2 shows the coincident observations for comparison with the Park Falls, WI TCCON site. This footprint would overlap the footprint of the TCCON site at Lamont, OK (since the latitude limit is 7.5 deg and these sites are separated by only 10 deg lat). These two sites have a seasonal that differ by several ppm, and there is a subset of ACOS data that is assumed to represent both sites. This is possibly justifiable, since transport in the free troposphere connects the two sites dynamically, but I think some justification of the criteria used, beyond a citation, is required.

Regarding the possible overlap of "footprints" of Park Falls and Lamont TCCON sites, the referee is correct that that is entirely possible. Such an overlap, however, is entirely physical; all that means is that there is a region in between the two where the XCO₂ is within 0.5 ppm of the XCO₂ at the two sites (which still allows for up to a ppm difference between the two sites). Since our colocation criterion is dynamic, such overlaps are more likely in the northern hemisphere winter, when the north-south gradient of XCO₂ over the conterminous United States is small (see examples in Figs. A and B in this response letter). During the northern hemisphere summer, when the north-south gradient is much larger and regions of constant XCO₂ are elongated zonally, such an overlap is less likely (see examples in Figs. C and D), reflecting the reality that XCO₂ at Lamont and Park Falls are different by more than a ppm. Thus, an itinerant overlap between the "footprints" of Park Falls and Lamont TCCON sites is completely consistent with the XCO₂ seasonal cycles at the two sites differing by several ppm.




Figure A. An example of co-located GOSAT soundings in the proximity of Park Falls TCCON station (star symbol) in winter. The co-locations accepted by the NOAA/Basu technique are denoted with green symbols.




Figure B. As in Fig. A but for Lamont TCCON station.




Figure C. As in Fig. A but for Park Falls station in summer.




Figure D. As in Fig. A but for Lamont station in summer.

As a reaction to the referee's comment, we have added a descriptive example figure to the paper (Fig. 2a), and clarified and expanded the text in Sect. 4.1 followingly:

“ACOS retrievals of GOSAT soundings are estimates of total column XCO_2 . Therefore, the issue of co-locating GOSAT soundings with TCCON soundings boils down to the question of whether we expect both sounding locations to have the same atmospheric XCO_2 . Any co-location technique is an assumption about the geographical region over which we expect XCO_2 to be the same as a TCCON retrieval, within some tolerance. For example, a geometrical co-location criterion, where we consider all GOSAT soundings within some fixed distance of a TCCON station, assumes that in the real atmosphere the variation of XCO_2 over that distance is smaller than said tolerance. Similarly, co-locating using the 700 hPa potential temperature (Wunch et al., 2011b) assumes that air with the same transport history – in so far as it's reflected in the 700 hPa potential temperature

– will have the same XCO₂ (within said tolerance). However, neither of these co-location techniques account for the fact that ultimately atmospheric XCO₂ is a convolution of surface fluxes and transport. Therefore, in our paper we have applied the NOAA/Basu co-location technique (Guerlet et al., 2013a) which uses a modelled atmospheric XCO₂ field to delineate the region around a TCCON station over which we expect XCO₂ to be constant within some tolerance (0.5 ppm). Since the model is run with realistic surface fluxes and atmospheric transport, we expect this co-location technique to account for XCO₂ variations due to both. To set upper spatiotemporal limits for the co-located soundings, the GOSAT soundings were required to be within $\pm 22.5^\circ$ in longitude and $\pm 7.5^\circ$ in latitude from the TCCON site, and acquired on the same day, within 2 hours of each other. The TCCON soundings were interpolated to local noon to exclude any effects from the diurnal cycle of XCO₂. In practice, the NOAA/Basu co-location technique has several advantages: high co-location data volume, good accuracy, and good sampling of parameter space, such as surface albedo. It should also be noted that the performance of this technique does not depend on the absolute accuracy of simulated XCO₂; all that is required is for the spatial gradient of three day average XCO₂ over a few thousand kilometers to be correct to within some tolerance.

The NOAA/Basu co-location technique is visually demonstrated for the Park Falls TCCON site in Fig. 2a. All GOSAT soundings over almost five years of co-located observations at Park Falls are mapped in Fig. 2b, which shows that the exact locations of the co-located GOSAT soundings are to a minor extent dependent on the season.”

The coincidence criteria are not plotted for any European TCCON sites, which are located quite close together. It seems that the autocorrelation between the GOSAT soundings used to compute the seasonal cycle for Bialystock, Bremen, Karlsruhe, Orleans, and Garmisch would be quite high. How does this affect the significance of the bias found by the authors with respect to the seasonal cycle?

To address this, we checked the number of GOSAT soundings that are shared co-locations for all five sites in continental Europe. This number turned out to be surprisingly low: only 18 GOSAT soundings were shared by all sites (this is about 1% of all co-located soundings per site). Next, we paired up the sites to test what is the maximum number of shared soundings between two sites. The results varied between 6% and 44%. The maximum of shared soundings was between Garmisch (44%) and Karlsruhe (39%); all other combinations shared less than a third of their soundings with another site. Considering that the fractions of shared soundings were less than half for every site, the shallow seasonal cycle issue seems to exist separately for all these sites.

When attempting to explain the discrepancy between TCCON and GOSAT, the authors use TCCON retrievals from both GGG2012 and GGG2014, and note that “The difference comes likely from the extended time series and the additional measurements present in the GGG2014 version”. Please clarify what is meant here. Are an additional 1-2 years worth of data going into the second data set? Are more TCCON data used to calculate its mean annual cycle since GGG2014 is better able to

account for e.g., solar zenith angle bias so fewer observations required to be excluded? It seems that for this comparison, the authors should take the average only of the same unique soundings.

We have updated all TCCON data used in the paper to GGG2014 versions that have now become available for all stations. To avoid confusion, we removed all references to GGG2012 from the paper.

Also on p 16476, the authors write that “a shallow-biased GOSAT seasonal cycle over Europe contradicts.. a stronger carbon sink over Europe”. This is not necessarily true, since the sink only depends on the imbalance between assimilation and respiration and does not require a strong seasonal difference between these two quantities. I don’t think there is necessarily a contradiction to resolve, but I definitely don’t understand the explanatory sentence that follows referencing Reuter.

Due to the current controversy around and about the European carbon sink, and the fact that this paper does not aim to contribute directly to that discussion, we decided to remove the mention of the European carbon sink entirely.

p 16479, If there are biases unique to individual retrieval algorithms that degrade the agreement with TCCON, is it really a good idea to brush these biases under the rug and attempt to use a composite dataset that contains a regionally inconsistent set of biases (EMMA)? I don’t see the logical link that supports the use of the EMMA.

During the revision, we tested the performance of EMMA and have modified this part in the text followingly: “Since none of the retrieval algorithms clearly outperformed the others at every TCCON site, we repeated the analysis for the ensemble median algorithm EMMA (Reuter et al., 2013), which combines all individual retrievals into one data set of median XCO₂ values. Even though EMMA had the smallest RMS error at four TCCON sites overall, it did not perform systematically better or worse than the individual retrieval algorithms in capturing the seasonal cycle of XCO₂.”

p 16483, The authors discuss the role of prior fluxes as driving differences in the GOSAT-to-model comparison, but should also include discussion of the differences in the transport models at the heart of each inversion.

The role of transport models (and the need of further studies on the matter) is already mentioned in several parts in the text (in particular in Sect. 5.3), and because this paper is focused on the validation of the seasonality of satellite-retrieved XCO₂, we do not wish to expand our speculation about the models further without additional results.

Figures: Figure 4: This is the closest to raw data that we see for most of the sites, so I would like it if this figure were bigger so the reader could interpret it better. Maybe two columns, but the same height so that the tick marks could be better spaced and labeled?

We updated the figure following these suggestions. In the updated version, the

seasonal cycles and especially the differences between TCCON and ACOS stand out more, which is good. We also added in dashed lines for the unobserved times of year.

Figure 8 and discussion in the text of longitude-dependence of seasonal cycle: It would be nice if there were a figure in the paper that showed the position of the continents and some indicator of the dominant tropospheric transport pattern to allow the reader to better interpret. For example, is the relatively higher amplitude between 120-60W due to the fact that those measurements are made over a continent? Or is it due to the fact that the jet is northwesterly at this location?

We added a small map of the Northern Hemisphere to the top of the figure, and highlighted the region in question. From this map, the locations of the continents with respect to the latitude-longitude regions can be conveniently seen. We also added this sentence to the text: “The increased seasonal cycle is likely due to the large seasonal sink of the boreal forests, accrued in the total column as the observation point is moved eastward.”

wording: Abstract: “The seasonal cycle ... represents an important variable to accurately measure from space”, but we can’t measure the seasonal cycle since the long-term upward trend necessitates pre-treatment of the measurements. Therefore, the wording might be better as “represents an important quantity to test the accuracy of measurements from space” or something like that.

Modified as suggested.

“GOSAT agrees with the models”. Why should the data have to agree with models? Perhaps better worded as “models are consistent with the GOSAT amplitude to within 1.4 ppm”

Corrected as suggested.

p 16465 “This finding not only suggests that regional XCO₂ can be indicative of local fluxes”. I find this wording confusing.

We modified the sentence as follows: “This finding suggests that regional XCO₂ seasonal cycles may be indicative of local fluxes, and hence that satellite-measured XCO₂ may be useful in evaluating model fidelity without resorting to full carbon flux inversions. It is also another reminder that there may be much to be gained by assimilating space-based XCO₂ retrievals which vastly expand the current in-situ measurement network; a lesson shown previously by a number of studies (e.g., Rayner and O’Brien, 2001; Chevallier et al., 2007; Takagi et al., 2011, 2014; Maksuytov et al., 2013).”

p 16467 “GOSAT measures scattered solar”. I think “reflected” sunlight is more accurate term.

Corrected as suggested.

Review #2

General comments:

The amplitude of the GOSAT fit should be viewed with caution above 60N where the gaps in the seasonal cycle could cause significant fit errors. When comparing to models, the same data gaps should be applied to both the models and the GOSAT and TCCON data.

The models have been resampled at exact GOSAT soundings in latitude, longitude and time, and therefore take into account the gaps in the satellite data.

The amplitude and phase of the fit may be partially prescribed by the fit function that is used, e.g. the fit of data far from the peak could affect the peak location and amplitude, so it is important to assess the fit minus data residuals for signal. The seasonal cycle peak and minimum might be more accurately calculated with a local smoothing function rather than a prescribed globally fit function. For this paper, plots and assessment of fit minus data residual signals, especially near the peak and minimum, and discussion of the above should be included if there are residual signals.

Based on this comment, we made the plots of the residuals for each TCCON site, and found that there was no systematic signal left in the residuals. We identified few non-systematic, small-scale features at a few sites (for example at Bialystok) but these were something that would be extremely difficult to fit out anyway. We added the following sentence to Sect. 4.3 to briefly summarize these studies: "To ensure that the amplitude and phase of the seasonal cycle were not determined largely by the fit function, we assessed the fit-minus-data residuals for both TCCON and ACOS, and could not identify any systematic signatures in the residuals."

"As model-to-model differences in XCO₂ can be several ppm at regions poorly sampled by in-situ measurements, GOSAT observations that measure seasonal cycle amplitude to within 1.0 ppm, based on this study, could potentially be used directly (without elaborate inversions) to evaluate model differences at these regions."

The statement that GOSAT observations that measure seasonal cycle amplitude to within 1.0 ppm globally should be qualified. The satellite retrievals depend on a priori knowledge of the interferent species, like aerosols, temperature, and water, which will be better constrained in Europe and North America where most TCCON stations are. These errors may be larger in other parts of the world. The statement should be modified to something like "whereas the ACOS-GOSAT seasonal cycle error is on the order of 1.0 ppm near TCCON stations and likely to be of this size in other parts of the world, though may be influenced by the a priori accuracy of jointly retrieved parameters."

This should be updated in the text and conclusions.

Based on this reasonable comment, we modified the text in the conclusions as follows: "Based on our study, the GOSAT/ACOS seasonal cycle error is of the

order of 1.0 ppm near TCCON stations and likely to be of this size in other parts of the world, though may be influenced by the a priori accuracy of jointly retrieved parameters, such as those related to aerosols. As model-to-model differences in the XCO₂ seasonal cycle amplitude can be several ppm at regions poorly sampled by in-situ measurements, GOSAT observations could potentially be used directly (without elaborate inversions) to evaluate model differences at these regions. This idea is explored in more detail in a work under preparation (Lindqvist et al., 2015)."

Specific Comments:

Page 4 line 100: "likely to be affected by any seasonal biases present in the GOSAT/ACOS retrievals that are due to the ACOS system itself." change to "likely to be affected by any seasonal biases present in the GOSAT/ACOS retrievals that are due to the ACOS system or ACOS a priori inputs."

Corrected as suggested.

Page 5, line 130 "Their validated and calibrated higher precision and accuracy compared to satellite observations, coupled with the fact that they measure the same quantity in essentially the same way as the satellites" change to "coupled with the fact that they measure the same quantity in essentially the same way as the satellites, though looking directly at the sun rather than sunlight reflected off the earth, so are not affected by surface albedo, "

Corrected as suggested.

Page 5, line 145. The southern hemisphere amplitude is small, however it is has large flux uncertainties and less in situ data, so that satellites could add significant guidance to models. I would not discount it but rather state why your analysis is not appropriate for it or that you choose to focus on the northern hemisphere.

We added the following sentence to the manuscript Sect. 3.1: "We therefore chose to focus on the Northern Hemisphere, which has both a larger seasonal cycle amplitude, and a larger quantity of TCCON stations against which to compare."

Page 5, line 190. It doesn't seem like TCCON should be hyphenated at a line break.g. TC-CON.

Corrected throughout the LaTeX document.

Page 7, line 219, "Finally, we calculated daily averages of both GOSAT/ACOS and TCCON retrievals." What is the local time of TCCON that is averaged? Is it the time averaged for TCCON around the time of the GOSAT observations? Please state.

We modified the sentence to clarify this: "Finally, we calculated daily averages of co-located GOSAT/ACOS and TCCON retrievals."

Page 8, line 235. The "daily error" for GOSAT/ACOS and TCCON are of interest, so state what they are.

This confusing term has been replaced with “ σ of each daily-averaged XCO₂”.

Page 8, line 235. The fit chosen may also not be the correct seasonal fit, so it is important to note whether the TCCON error (in particular since TCCON errors are smaller) is randomly distributed about the fit. This can be shown with a difference plot, e.g. with green dots around the dashed lines in figure 4, or in a separate figure, in particular for a case where there are larger differences in the maximum location.

According to our additional studies made during the revision, the fit - TCCON residuals are small and randomly distributed, with no systematic signal.

Page 8, Equation 1. $\cos^{-1}()$ has a domain issue in that $\cos^{-1}(x)$ will range from 0 to π , rather than $-\pi$ to π . I can't quite wrap my mind around what $\sin(\cos^{-1}(\cos(wt)))$ does. Could you give the fit values for a_0 - a_5 for at least one example, e.g. Park Falls. I assume that the $\cos^{-1}()$ term is to give a time-dependent phase. Is this a standard equation for fitting a seasonal cycle? Is there a reference for this fit? It doesn't matter if there is a reference if it does a good job; the quality of the fit should be assessed by looking at residuals of fit-data (see general comments).

Unfortunately we do not have a special reference for the fit; the function is one of many ways of creating a so-called skewed sine wave, and to our knowledge has not been used in a seasonal cycle context before. This domain issue pointed out by the referee is true for Eq. (1) and in practice means that, for certain parameter combinations, the fitted function has unphysical discontinuities and regions where it does not exist. However, it turned out that such parameter combinations (even though they were allowed by the nonlinear fitting procedure) never resulted in the lowest chi square values, and were excluded on that basis. As an example, we added the fit parameter values to Table 2 for the TCCON and ACOS fits at Park Falls.

Page 9, Line 286, "The satellite observes the maximum later than the TCCON at the European sites, but obtains good agreement elsewhere. At the European sites, the difference extends up to 2–3 weeks, and is likely connected with the biased amplitude inferred by ACOS discussed below." Fitting can create phase differences if the fitting function does not match the data shape (see general comments). Can a plot be shown of the GOSAT/ACOS and TCCON data for a station where there is a phase difference between TCCON and GOSAT so that the reader can see that the data supports the fit shape? Kulawik et al., 2015 used cross-correlation to determine phase shift and found a much smaller phase difference in Europe, which seems in disagreement of your findings.

This is an interesting observation and definitely worth a comment in the text. We decided not to add another figure, however, but instead explain how we derived error statistics for the fitted maximum and minimum, because these statistics reflect the statistical uncertainty in the fit. We expanded the text as follows: “However, regarding the difference in the dates of the maximum, Kulawik et al. (2015) found a much smaller phase difference in Europe by using cross-correlation to determine the phase shift. Because our results were based on the fitted seasonal cycles instead of the actual data, we evaluated the statistical

errors of the dates of the maximum and minimum XCO₂ with a Monte Carlo approach, using the error covariance matrices associated with the fitted function parameters. The deviations from the fit maximum and minimum followed a normal distribution with an average σ of 3.5 days for the TCCON maximum date, and 6.1 days for ACOS maximum date, reflecting a notable uncertainty in the fitted phase and thus explaining at least partially the difference between our results and those of Kulawik et al. (2015). The corresponding average σ for the date of the minimum were 2.2 days (TCCON) and 3.6 days (ACOS)."

Page 12, line 390, " These results can be interpreted to support the ensemble median algorithm EMMA introduced by Reuter et al. (2013), which combines all individual retrievals into one data set that globally has the best agreement with TCCON." It would be useful to add EMMA to Figure 6.

We repeated the analysis for EMMA and it turned out that EMMA was neither the best nor the worst when compared to TCCON by the measures that we use in Fig. 6. However, we would prefer not to replace Fig. 6 with a version where EMMA is included because the figure is already quite busy with symbols and because EMMA did not outperform the other algorithms in any of the panels. We updated the sentence in the text accordingly: "Since none of the retrieval algorithms clearly outperformed the others at every TCCON site, we repeated the analysis for the ensemble median algorithm EMMA (Reuter et al., 2013), which combines all individual retrievals into one data set of median XCO₂ values. Even though EMMA had the smallest RMS error at four TCCON sites overall, it did not perform systematically better or worse than the individual retrieval algorithms in capturing the seasonal cycle of XCO₂."

Page 12, line 405, " The seasonal cycle was fitted on the daily averages of GOSAT/ACOS XCO2 and the resampled models." The models were presumably sampled in the daytime? It is important to match the approximate GOSAT overpass time. Also, see general comments, gaps in the GOSAT data can result in differences from a complete seasonal cycle.

We agree that matching the GOSAT overpass time is important, and therefore (as explained in Sect. 3.2) all modeled XCO₂ data were resampled at exact GOSAT observations in latitude, longitude and time, so the model seasonal cycles include the same gaps as the satellite data does.

Page 13, line 427, "From 60_ to 70_, ACOS has a higher seasonal cycle amplitude than most models." North of 60N the gaps in GOSAT seasonal data are such that the peak fit of the seasonal cycle is likely outside of the seasonal span of GOSAT data, see general comments. To compare to model fits, both models and data should have the same data gaps.

As already mentioned in the general comments, the models and the satellite data both have the same data gaps because we resample the model values at the GOSAT soundings. The winter gap north of 60 degrees latitude is indeed wide, but it appears that we observe the maximum or at least the time very close to it (at least in most years) because the XCO₂ values increase during the first (roughly) 10-20 days before they start to decrease.

Page 13, line 440, that the averaging kernel correction results in a modest systematic effect on the seasonal cycle amplitude is an important finding which should be mentioned in the conclusions. A seasonally dependent 0.2 ppm error could have a significant impact on flux estimates.

We added this finding in the Conclusions as follows: “We also noticed that the averaging kernel correction can systematically decrease the seasonal cycle amplitude by up to 0.2 ppm.”

Page 15, line 508. Accuracy of GOSAT/ACOS results has dependence on prior information of the interferents and some caution is a warranted regarding the accuracy far from TCCON sites.

As pointed out in our replies to the General comments, we have modified this sentence to take into account the fair and valid concerns of the referee. It is true that without further validation studies we simply cannot know how accurate the GOSAT soundings are far from the TCCON sites, although we do not expect the accuracy to deteriorate notably, because the TCCON sites used in validation already cover a variety of different atmospheric and geographic conditions.

Figure 2. The tan background makes the colors hard to see.

The figure colors have been changed (currently this is Figure 2b). We also added in the US state borders and the provincial borders for Canada.

Figure 5 label: Refer to Panel (a) and Panel (b) rather than Panel a and Panel b.

Corrected as suggested.

Other changes

- We updated the Bremen TCCON data from GGG2012 retrieval version to GGG2014 retrieval. This led to an improved match of GOSAT/ACOS and TCCON.
- We also updated the Karlsruhe TCCON data from retrieval GGG2014 r0 to GGG2014 r1. This had only minor effects on the comparisons but led to an improved match of GOSAT/ACOS and TCCON.
- The above-mentioned data updates led to updated versions of Figs. 4, 5 and 6, and Table 3.
- We found an error in the definition of Eq. (1) parameter ω and corrected that.
- We updated the journal reference of Kulawik et al. (2015).
- We updated the Acknowledgements to include one more funding agency and both referees.

Manuscript prepared for Atmos. Chem. Phys.
with version 2014/07/29 7.12 Copernicus papers of the L^AT_EX class copernicus.cls.
Date: 2 October 2015

Does GOSAT capture the true seasonal cycle of XCO₂?

Hannakaisa Lindqvist^{1,2}, Christopher W. O'Dell¹, Sourish Basu^{3,4},
Hartmut Boesch⁵, Frédéric Chevallier⁶, Nicholas Deutscher^{7,8}, Liang Feng⁹,
Brendan Fisher¹⁰, Frank Hase¹¹, Makoto Inoue^{12,13}, Rigel Kivi¹⁴, Isamu Morino¹²,
Paul I. Palmer⁹, Robert Parker⁵, Matthias Schneider¹¹, Ralf Sussmann¹⁵, and
Yukio Yoshida¹²

¹Department of Atmospheric Science, Colorado State University, CO, USA

²Finnish Meteorological Institute, Helsinki, Finland

³NOAA Earth System Research Laboratory, Boulder, CO, USA

⁴Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

⁵National Centre for Earth Observation, Department of Physics and Astronomy, University of Leicester, Leicester, UK

⁶Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, Gif-sur-Yvette, France

⁷School of Chemistry, University of Wollongong, Wollongong, Australia

⁸Institute of Environmental Physics, University of Bremen, Bremen, Germany

⁹National Centre for Earth Observation, School of GeoSciences, University of Edinburgh, Edinburgh, UK

¹⁰Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

¹¹Karlsruhe Institute of Technology, IMK-ASF, Karlsruhe, Germany

¹²National Institute for Environmental Studies (NIES), Tsukuba, Japan

¹³Department of Biological Environment, Akita Prefectural University, Akita, Japan

¹⁴Finnish Meteorological Institute, Arctic Research Centre, Sodankylä, Finland

¹⁵Karlsruhe Institute of Technology, IMK-IFU, Garmisch-Partenkirchen, Germany

Correspondence to: Hannakaisa Lindqvist (hannakaisa.lindqvist@fmi.fi)

Abstract.

The seasonal cycle accounts for a dominant mode of total column CO₂ (XCO₂) annual variability and is connected to CO₂ uptake and release; it thus represents an important ~~variable to accurately measure~~ quantity to test the accuracy of the measurements from space. We quantitatively evaluate the XCO₂ seasonal cycle of the Greenhouse Gases Observing Satellite (GOSAT) observations from the Atmospheric Carbon Observations from Space (ACOS) retrieval system, and compare average regional seasonal cycle features to those directly measured by the Total Carbon Column Observing Network (TCCON). We analyze the mean seasonal cycle amplitude, dates of maximum and minimum XCO₂, as well as the regional growth rates in XCO₂ through the fitted trend over several years. We find that GOSAT generally captures the seasonal cycle amplitude within 1.0 ppm accuracy compared to TCCON, except in Europe, where the difference exceeds 1.0 ppm at two sites, and the amplitude captured by GOSAT is generally shallower compared to TCCON. This bias over Europe is not as large for the other GOSAT retrieval algorithms (NIES v02.21, RemoTeC v2.35, UoL v5.1, and NIES PPDF-S v.02.11), although they have significant biases at other sites. The ACOS bias cor-

15 rection was found to partially explain the shallow amplitude over Europe. The impact of the ~~TCCON~~
~~retrieval version~~, co-location method ~~7~~, and aerosol changes in the ACOS algorithm were also tested,
and found to be few tenths-of-a-ppm and mostly non-systematic. We find generally good agreement
in the date of minimum XCO₂ between ACOS and TCCON, but ACOS generally infers a date of
maximum XCO₂ 2-3 weeks later than TCCON. We further analyze the latitudinal dependence of
20 the seasonal cycle amplitude throughout the Northern Hemisphere, and compare the dependence
to that predicted by current optimized models that assimilate in-situ measurements of CO₂. In the
zonal averages, ~~GOSAT agrees with the models~~ models are consistent with the GOSAT amplitude to
within 1.4 ppm, depending on the model and latitude. We also show that the seasonal cycle of XCO₂
depends on longitude especially at the mid-latitudes: the amplitude of GOSAT XCO₂ doubles from
25 West U.S. to East Asia at 45 – 50°N, which is only partially shown by the models. In general, we
find that model-to-model differences can be larger than GOSAT-to-model differences. These results
suggest that GOSAT retrievals of the XCO₂ seasonal cycle may be sufficiently accurate to evaluate
land surface models in regions with significant discrepancies between the models.

1 Introduction

30 Satellites provide unprecedented spatial coverage of the variability of atmospheric carbon dioxide
(CO₂) through retrievals of column mean dry mole fractions of CO₂ (XCO₂). XCO₂ shows tem-
poral variability on different timescales: diurnal, synoptic, seasonal, inter-annual, and long-term
(Olsen and Randerson, 2004; Keppel-Aleks et al., 2011). Variability is determined by the collec-
tive impact of CO₂ fluxes resulting from fossil fuel emissions, biosphere-atmosphere exchange, and
35 ocean-atmosphere exchange, and the imprint of these on regional XCO₂ can be strongly influenced
by atmospheric dynamics, in addition to the regional origin of the fluxes. While the secular trend
and multi-year interhemispheric CO₂ gradient are driven by the global build-up of CO₂ from fossil
fuel combustion mainly in the Northern Hemisphere, the seasonal variability is mainly controlled
by variations in the terrestrial biospheric fluxes (Palmer et al., 2008; Keppel-Aleks et al., 2011). The
40 ocean-atmosphere and fossil fuel CO₂ fluxes are, although seasonally varying, only minor contribu-
tors to the XCO₂ seasonal variability in the Northern Hemisphere. Therefore, the seasonal cycle of
XCO₂ bears the signature of large-scale biospheric flux patterns, especially their north-south distri-
bution.

Regional biospheric CO₂ fluxes are a critical ~~part~~ output of land surface models that describe the
45 biosphere-atmosphere carbon exchange in larger modeling systems, such as ~~carbon cycle and climate~~
coupled climate - carbon cycle models (Pitman, 2003). Inverse model systems use these land surface
models in conjunction with atmospheric transport models, and optimize their CO₂ flux estimates by
assimilating CO₂ measurements, but especially in regions where the in-situ measurement network
has sparse coverage, the inverse models can strongly disagree about the seasonality and magnitude

50 of the fluxes (Lindqvist et al., 2015, in prep.). Recently, this disagreement has been found to lead to large regional discrepancies of several ppm in the seasonal cycle amplitudes of modeled XCO₂ (Keppel-Aleks et al., 2012; Peng et al., 2015; Lindqvist et al., 2015, in prep.). This finding ~~not only~~ suggests that regional XCO₂ ~~can seasonal cycles may~~ be indicative of local fluxes ~~and, and hence~~ that satellite-measured XCO₂ may be useful in ~~constraining the models even without inversions, but~~ ~~also is evaluating model fidelity without resorting to full carbon flux inversions.~~ It is also another reminder that there ~~is potentially may be~~ much to be gained by assimilating space-based XCO₂ retrievals ~~that which~~ vastly expand the current in-situ measurement network; a lesson shown previously by a number of studies (e.g., Rayner and O'Brien, 2001; Chevallier et al., 2007; Takagi et al., 2011; Maksuytov et al., 2013; Takagi et al., 2014). In particular, the strength of the seasonal cycle drawdown is fundamentally connected to the magnitude of the carbon sink during the growing season. By studying the GOSAT XCO₂ seasonal cycle and its inter-annual variability, Wunch et al. (2013) showed that the variability in the drawdown correlates with surface temperature in the boreal regions, and Guerlet et al. (2013b) found a reduced carbon uptake during the 2010 Northern Hemisphere summer.

65 The Greenhouse Gases Observing Satellite (GOSAT; Yokota et al., 2009) and the Orbiting Carbon Observatory -2 (OCO-2; Crisp et al., 2004) are indeed designed to make near-global XCO₂ retrievals that will constrain the inverse model systems enough to provide a picture of the global carbon cycle with respect to regional sources and sinks. However, a crucial question still lingers: are the satellite observations accurate enough to reliably capture the seasonal variability of XCO₂?
70 The question is fair because satellite-retrieved XCO₂ is subject to biases in the retrieval system (e.g., Wunch et al., 2011b), and also sampling biases due to the seasonally-dependent amount of solar radiation (e.g., Liu et al., 2014). Both of these may have an impact on the measured seasonal cycle. For the Atmospheric CO₂ Observations from Space (ACOS) retrieval system (O'Dell et al., 2012; Crisp et al., 2012), known biases in GOSAT retrievals are corrected using a global bias correction
75 (Wunch et al., 2011b) but some parameters of the bias correction vary seasonally, for example surface albedo. Potential remaining biases, their seasonality, and impact on the seasonal cycles of XCO₂ are best identified through evaluation of the GOSAT seasonal cycle against the best available independent data — those from the Total Carbon Column Observing Network, TCCON (Wunch et al., 2011a). There have been several studies that compare GOSAT retrievals against the TCCON, some
80 of them introducing novel methods for comparisons (Wunch et al., 2011b; Nguyen et al., 2014), some concentrating on quantifying biases in a specific retrieval algorithm (Butz et al., 2011; Cogan et al., 2012; Yoshida et al., 2013), and some focusing more on the intercomparison of different retrieval algorithms (Buchwitz et al., 2013; Oshchepkov et al., 2013a; Reuter et al., 2013; Dils et al., 2014). Overall, the collective message from the validation studies is that the agreement of GOSAT
85 and TCCON has improved (i.e., the satellite biases have decreased) substantially from the earliest validation efforts (Morino et al., 2011), owing to major improvements and updates in the retrieval

algorithms and the development of more sophisticated comparison methods. However, less attention has been paid to the evaluation of the seasonal cycle. Reuter et al. (2013, p. 1776) touched on this by showing averages of the seasonal cycle amplitude differences between all GOSAT retrievals and TCCON (and also a model, CarbonTracker CT2011_oi). More recently, Kulawik et al. (2015) studied the seasonality of GOSAT-TCCON biases (using the ACOS B3.4 retrieval algorithm for GOSAT data) and found notable station-to-station variability in the biases, but also persisting seasonal biases in latitudinally averaged results. These seasonal biases were reflected in the seasonal cycle amplitudes.

In this paper, we continue the evaluation of the GOSAT seasonal cycle from Kulawik et al. (2015). Five years of GOSAT observations and the updated TCCON GGG2014 retrievals lengthen the co-located time series sufficiently to evaluate the seasonal cycles regionally at 12 TCCON sites in the Northern Hemisphere. We extend the seasonal cycle analysis to four other retrieval algorithms to identify potential biases characteristic to the ACOS retrievals. Although the emphasis of the study is on these TCCON comparisons, we also compare the GOSAT seasonal cycle against models that assimilate in-situ data; because of their connection to measurements, models may be a reasonable representation of the truth in areas with high assimilated data density, such as North America or Western Europe. This seasonal cycle evaluation study lays important ground work to the analysis of OCO-2 observations that also use the ACOS retrieval system and are, therefore, likely to be affected by any seasonal biases present in the GOSAT/ACOS retrievals that are due to the ACOS system ~~itself~~ for ACOS a priori inputs.

2 GOSAT

The Greenhouse Gases Observing Satellite (GOSAT), developed by Japan Aerospace Exploration Agency (JAXA), was launched in January 2009 to make near-global greenhouse gas measurements from a polar orbit (Yokota et al., 2009). GOSAT measures ~~scattered-reflect~~ ed solar near-infrared radiation with a Fourier transform spectrometer (TANSO-FTS; Kuze et al., 2009). The diameter of a GOSAT sounding footprint is approximately 10 km, and the soundings repeat in a three-day cycle. We used GOSAT data taken in two primary modes: glint over oceans, and nadir view over land. Nadir data over land has two gain states: high gain (H) for most of the data, and medium (M) over bright surfaces, such as deserts.

Several retrieval algorithms have been developed for retrieving the column-averaged CO₂ from the GOSAT near-infrared measurements; these algorithms have been recently reviewed and compared by Oshchepkov et al. (2013a) and Reuter et al. (2013). In this paper, we concentrate on the evaluation of the Atmospheric CO₂ Observations from Space build 3.5 (ACOS B3.5) retrieval algorithm (Crisp et al., 2012). The ACOS retrieval algorithm is described in detail by O'Dell et al. (2012). The most significant subsequent updates and improvements to the operational algorithm include updated

spectroscopy for the 1.6 μm and 2.1 μm CO_2 absorption bands, moving from static to dynamic vertical pressure levels, an improved prior profile of CO_2 , and a complete change in the treatment of aerosol and cloud scattering. Instead of a globally constant aerosol model that was incorporated in
125 ACOS B3.4 and earlier versions, B3.5 uses Modern-Era Retrospective Analysis for Research and Applications (MERRA) reanalysis data of five aerosol types (mineral dust, sea salt, black carbon, sulphates, and organic carbon) to determine two most common types at a given GOSAT sounding location, and applies their respective optical properties in the retrieval.

3 Validation data

130 3.1 TCCON

The Total Carbon Column Observing Network (TCCON) is currently composed of 21 operating Fourier transform spectrometers that make ground-based measurements of atmospheric XCO_2 and other gases (Wunch et al., 2011a). Their validated and calibrated higher precision and accuracy compared to satellite observations, coupled with the fact that they measure the same quantity in
135 essentially the same way as the satellites, [though looking directly at the sun rather than sunlight reflected off the Earth, so are not affected by surface albedo](#), make them an ideal, independent validation source for GOSAT (Wunch et al., 2010; Messerschmidt et al., 2011). Though the seasonal cycle of TCCON has itself never been explicitly validated by comparison with aircraft, we implicitly assume that our inferred TCCON seasonal cycles for XCO_2 can be taken as truth, similar to
140 the assumption in several previous studies (Messerschmidt et al., 2011; Keppel-Aleks et al., 2012; Wunch et al., 2013), though in principle sub-ppm seasonal biases could remain. For instance, the TCCON retrieval performs a post-hoc airmass bias correction (Wunch et al., 2011a), errors in which could lead to small but nontrivial differences in the TCCON seasonal cycle. However, it is beyond the scope of this work to validate the accuracy of the TCCON seasonal cycle.

145 In this study, we used data from all Northern Hemisphere TCCON sites that had 1) at least two years of coincidental measurements with GOSAT; and 2) enough co-located data (see Sect. 4.1) to evaluate a seasonal cycle; i.e., both ACOS and TCCON observations available at the proximity of the site through most seasons. The first criterion eliminated the Four Corners and Caltech/Pasadena sites, while the second eliminated the northernmost sites of Ny Ålesund and Eureka which have
150 very little co-located data due to the high latitude. We did not include the Southern Hemisphere sites because the seasonal changes in XCO_2 at those sites are minor, making the definition of an average seasonal cycle more ambiguous and sensitive to inter-annual variability. [We therefore chose to focus on the Northern Hemisphere, which has both a larger seasonal cycle amplitude, and a larger quantity of TCCON stations against which to compare](#). The sites that were used in this study are shown in
155 Fig. 1. For these sites, we analyzed all co-located data between April 23, 2009, and December 31, 2013. We used the newest available [GGG2014](#) TCCON retrievals for [every site. GGG2014 retrievals](#)

were available for the following sites in our study each site: Bialystok (Messerschmidt et al., 2012; Deutscher et al., 2014), Bremen (Notholt et al., 2014), Garmisch (Sussmann and Rettinger, 2014), Izaña (Blumenstock et al., 2014), JPL (Wennberg et al., 2014a), Karlsruhe (Hase et al., 2014), Lamont (Wennberg et al., 2014c), Orleans (Warneke et al., 2014), Park Falls (Washenwelder et al., 2006; Wennberg et al., 2014b), Saga (Kawakami et al., 2014), Sodankylä (Kivi et al., 2014), and Tsukuba (Ohyama et al., 2009; Morino et al., 2014), whereas for Bremen, we used the GGG2012 retrievals. TCCON data were obtained from the TCCON Data Archive website at <http://tcon.ornl.gov/>.

3.2 Model CO₂ data

Because evaluation against TCCON is limited to 12 sites in the Northern Hemisphere, another validation source is necessary for obtaining a more thorough view of the accuracy of the GOSAT seasonal cycle. Therefore, we also analyzed XCO₂ from three models that assimilate in-situ CO₂ measurements to optimize their fluxes. The models were CarbonTracker (CT2013B; Peters et al., 2007, with updates documented at <http://carbontracker.noaa.gov>), MACC 13.1 (Chevallier et al., 2010, documentation and data available at <http://www.copernicus-atmosphere.eu/catalogue>), and the University of Edinburgh model (UoE; Feng et al., 2009, 2011, <http://www.palmergroup.org>). Relevant model properties are listed in Table 1. The models were resampled at GOSAT/ACOS observations in latitude, longitude and time, and integrated over all atmospheric layers to form the column-averaged CO₂. The ACOS averaging kernel correction was first considered for CT2013B, but as it had only a very minor effect on the total column (generally < 0.1 ppm difference in monthly averages), it was subsequently neglected for all models. However, seasonal effects of the averaging kernel correction are briefly assessed in Sect. 5.3. All model results were available from the beginning of GOSAT data (April 23, 2009) but have different end dates: UoE and CT2013B run until the end of December 2012, and MACC 13.1 is available until the end of December 2013.

4 Methods

In this section, we describe the co-location of ground-based and satellite remote sensing measurements, filtering and bias correction for GOSAT/ACOS, and the averaging kernel correction, and define the average seasonal cycle. We demonstrate these steps with an example TCCON site at Park Falls, Wisconsin, U.S.

4.1 Co-locating GOSAT and TCCON

~~GOSAT/ACOS B3.5 ACOS retrievals of GOSAT soundings are estimates of total column XCO₂ observations were first co-located with the TCCON soundings, which were interpolated to local noon to exclude any effects from the diurnal cycle of. Therefore, the issue of co-locating GOSAT soundings with TCCON soundings boils down to the question of whether we expect both sounding locations to have~~

190 ~~the same atmospheric XCO₂. The Any~~ co-location ~~can be done in several ways that were described~~
~~and compared by Nguyen et al. (2014). In this study, we used~~ technique is an assumption about
the geographical region over which we expect XCO₂ to be the same as a TCCON retrieval, within
some tolerance. For example, a geometrical co-location criterion, where we consider all GOSAT
195 ~~the variation of XCO₂ over that distance is smaller than said tolerance. Similarly, co-locating using~~
~~the 700 hPa potential temperature (Wunch et al., 2011b) assumes that air with the same transport~~
~~history – in so far as it is reflected in the 700 hPa potential temperature – will have the same XCO₂~~
~~(within said tolerance). However, neither of these co-location techniques account for the fact that~~
~~ultimately atmospheric XCO₂ is a convolution of surface fluxes and transport. Therefore, in our~~
200 ~~paper we have applied~~ the NOAA/Basu co-location technique ~~that considers atmospheric transport~~
~~of CO~~(Guerlet et al., 2013a) which uses a modelled atmospheric XCO₂ ~~in addition to spatiotemporal~~
~~proximity of the TCCON and GOSAT observations (Guerlet et al., 2013a). This field to delineate the~~
~~region around a TCCON station over which we expect XCO₂ to be constant within some tolerance~~
~~(0.5 ppm). Since the model is run with realistic surface fluxes and atmospheric transport, we expect~~
205 ~~this co-location technique to account for XCO₂ variations due to both. To set upper spatiotemporal~~
~~limits for the co-located soundings, the GOSAT soundings were required to be within ±22.5° in~~
~~longitude and ±7.5° in latitude from the TCCON site, and acquired on the same day, within 2 hours~~
~~of each other. The TCCON soundings were interpolated to local noon to exclude any effects from~~
~~the diurnal cycle of XCO₂. In practice, the NOAA/Basu co-location~~ technique has several advan-
210 tages: high co-location data volume, good accuracy, and good sampling of parameter space, such as
surface albedo. ~~It should also be noted that the performance of this technique does not depend on~~
~~the absolute accuracy of simulated XCO₂; all that is required is for the spatial gradient of three day~~
~~average XCO₂ over a few thousand kilometers to be correct to within some tolerance.~~

The NOAA/Basu co-location ~~technique works as follows. Temporally, any~~ technique is visually
215 ~~demonstrated for the Park Falls TCCON site in Fig. 2a. All GOSAT soundings over almost five~~
~~years of co-located observations need to be acquired on the same day, within 2 hours of each~~
~~other. The spatial region of matching TCCON and GOSAT changes dynamically based on how~~
~~the inversion-derived estimates of local CO₂ surface fluxes are transported with the TM5 transport~~
~~model: the region around a TCCON site over which modeled XCO₂ does not differ by more than~~
220 ~~0.5 ppm from its value at the TCCON site sets the boundaries for co-location (as an upper spatial~~
~~limit, GOSAT soundings need to be within ±22.5° in longitude and ±7.5° in latitude from the~~
~~TCCON site). At Park Falls, all co-located GOSAT soundings~~ observations at Park Falls are mapped
in Fig. 2b, which shows that the exact locations of the co-located GOSAT soundings are to a minor
extent dependent on the season.

225 4.2 Data processing

We used GOSAT/ACOS B3.5 level 2 data, which has been pre-filtered and cloud-screened (O’Dell et al., 2012; Taylor et al., 2012). All available ACOS soundings (land H and M gain, ocean glint) were used at each site, but for the northern mid-latitude sites, most, if not all, data were land gain H soundings (see Table 3). After the co-location, the ACOS soundings were filtered using a post-
230 processing filter that removed bad data, such as data from poor spectral fits or containing larger amounts of aerosols, from the soundings. In total, filtering removed 47% of the H gain over land, 45% of M gain over land, and 40% of glint soundings that had been co-located with the TCCON sites considered in this study. An example of the effect of post-processing filtering is shown in Fig. 3, in the upper panels.

235 We also corrected for the known retrieval biases via a multi-parameter linear regression similar to Wunch et al. (2011b) but optimized for B3.5. The optimization is done with respect to all TCCON data and an average of eight inversion-based models. Model results are used for bias correction only when the models agree with each other to within 1 ppm of the total XCO₂ for a given sounding. The bias correction algorithm performed a correction to the retrieved XCO₂ based on different parameters. Bias correction is optimized globally, not regionally, but separately for land (nadir, gains H and
240 M) and ocean (glint) soundings.

When comparing two different remote-sensing measurements, the results are not comparable before the difference due to the retrieval averaging kernels has been considered (Rodgers and Connor, 2003). Since the averaging kernels of TCCON and ACOS are quite similar, it was sufficient to follow
245 the correction introduced by Wunch et al. (2011b), and further implemented in Nguyen et al. (2014). The effects of the averaging kernel correction for TCCON and bias correction for GOSAT/ACOS soundings are presented in Fig. 3, in the lower left panel. For model results, the averaging kernel corrections were not applied.

Finally, we calculated daily averages of ~~both~~ co-located GOSAT/ACOS and TCCON retrievals.
250 This way, days with multiple soundings are not more dominant in the seasonal cycle fit than the days with fewer soundings. Time series of daily averages are shown in Fig. 3, in the lower right panel.

4.3 Seasonal cycle

In what follows, we parameterize the seasonal cycle of XCO₂ as a skewed sine wave with an upward trend, and find that it is generally a good model for the time series of XCO₂ in the Northern
255 Hemisphere. We fitted an average seasonal cycle to the daily XCO₂ averages using the following six-parameter function

$$f(t) = a_0 + a_1 t + a_2 \sin(\omega[t - a_3] + \cos^{-1}[a_4 \cos(\omega[t - a_5])]), \quad (1)$$

where t is the time in days and ~~ω is the annual period of~~ $\omega = 2\pi/T$, where T is 365 days. The first two terms with the parameters a_0 and a_1 (denoting the average growth rate) fit for a linear trend,

260 and the third term, a sine wave with a time-dependent phase, fits for the seasonal cycle parameters $a_2 - a_5$. As an example, we give the parameters for both TCCON and ACOS fits at Park Falls in Table 2. In particular, $2|a_2|$ denotes the peak-to-peak amplitude of the sine wave and is, from here forwards, used to define the seasonal cycle amplitude. The nonlinear least squares fit was solved using a standard gradient-expansion algorithm. For Park Falls, the seasonal cycle fits for TCCON and ACOS are shown in Fig. 3, lower right panel, and the resulting seasonal cycle amplitude is 8.4 ± 0.1 ppm for TCCON, and 8.6 ± 0.2 ppm for ACOS. The errors of the fitted parameters are driven by the standard deviations σ of each daily XCO₂, initially requiring $\sigma_{\text{ACOS}} \geq 1.5$ ppm and $\sigma_{\text{TCCON}} \geq 0.3$ ppm. Because the true errors in daily-averaged XCO₂ are not well known, we scaled the ~~daily errors σ of each daily-averaged XCO₂~~ by multiplying them with the minimized quantity χ to yield $\chi^2 = 1$ from the least squares fit. For TCCON data fits, the original χ^2 values varied between $2 < \chi^2 < 10$, while for ACOS, the values were typically $\chi^2 < 1$, which implies that the initial errors σ_{TCCON} may have been underestimated and σ_{ACOS} overestimated. The fitting errors are purely statistical, and do not take into account systematic errors in the data. A more traditional Fourier series fit with an annual and semi-annual cycle (Wunch et al., 2013) was also tried, and the fitted seasonal cycle amplitudes were virtually identical (well within the fitting errors), but because some strange behavior during unobserved times of year could result, we opted for the fit in Eq. (1). To ensure that the amplitude and phase of the seasonal cycle were not determined largely by the fit function, we assessed the fit-minus-data residuals for both TCCON and ACOS, and could not identify any systematic signatures in the residuals.

280 We recognize that there could be inter-annual variability in some or all of the fitted parameters, and that our results can be affected by that variability; especially we can expect sites with shorter co-located time series to be more sensitive. However, we do not fit for inter-annual variability because we are interested in identifying potential systematic errors in the average seasonal cycle captured by GOSAT and, in particular, the ACOS retrieval system. For the purposes of evaluating the average seasonal cycle of XCO₂, it is important to compare observations from the same time interval, which we take into account by co-locating the observations from TCCON and GOSAT.

5 Results and discussion

5.1 Evaluation against TCCON

290 Seasonal cycles for co-located TCCON and GOSAT/ACOS B3.5 XCO₂ soundings were studied at 12 TCCON sites in the Northern Hemisphere. Detrended average seasonal cycles for both retrievals at each site are shown in Fig. 4. Detrending removed a linear trend, i.e. XCO₂ average growth rate, that varied between ~~1.90 — 2.39~~ 1.88 — 2.39 ppm/year for ACOS and ~~2.02 — 2.58~~ 2.03 — 2.58 ppm/year for TCCON retrievals, depending on the site. We estimated the sensitivity of the average seasonal cycle parameters of Eq. (1) to the fitted trend from the error covariance matrix associated to the

295 best-fit parameters. The error in the trend was generally weakly negatively correlated with the error
in the seasonal cycle amplitude, for both TCCON and ACOS. The phase-related parameters $a_3 - a_5$
were not correlated with the trend. Therefore, the error from removing the trend should statistically
have little effect on the parameters of the average seasonal cycle. Descriptive fit parameters together
with the associated errors are collected in Table 3. Instead of showing the fitted values for the three
300 parameters $a_3 - a_5$ of the phase term in Eq. (1), the average dates of annual maximum and minimum
XCO₂ are listed.

The global average growth rate in CO₂ is accurately captured by long-term ground-based mea-
surements of CO₂ concentration, such as the Mauna Loa record (Keeling et al., 1976). Global annual
trends for the years 2009–2013 varied between 1.66 ppm/year and 2.53 ppm/year (Ed Dlugokencky
305 and Pieter Tans, NOAA/ESRL, www.esrl.noaa.gov/gmd/ccgg/trends/, 30.3.2015). The accuracy of
the TCCON-inferred regional XCO₂ growth rates is not precisely known, though agreement of 0.1–
0.2 ppm/year in the global growth rate has been obtained via assimilation of TCCON data in an in-
verse modeling framework (Chevallier et al., 2011). According to Table 3, GOSAT shows a slightly
lower XCO₂ growth rate than TCCON at many validation sites, of order 0.2 ppm/year (around 10%).
310 Only at JPL, the trend fitted for GOSAT is modestly larger than that of TCCON. There are several
explanations for this. Firstly, GOSAT showing a generally lower trend than TCCON is not surpris-
ing but rather a sign of a potentially inaccurate correction for radiometric degradation that is caused
by minor contamination of the instrument over time (Kuze et al., 2014). Secondly, time series of a
little over 2 years of co-located data (like those of Saga, JPL, and Tsukuba) are arguably too short to
315 distinguish a trend from inter-annual variability. However, the trend captured by GOSAT may be of
minor significance compared to its measurements of the seasonal cycle: errors in capturing the trend
may result in errors of the order of a few tenths-of-a-ppm while errors in capturing the seasonal cycle
may have a more significant impact, though this will depend on the detailed set-up of each inverse
modeling system.

320 The phase of the seasonal cycle is relatively well captured by GOSAT/ACOS. The timing of the
(detrended) maximum concentration varies from March 16–8 to May 16 for TCCON, and from April
1–March 27 to May 21 for GOSAT. The satellite observes the maximum later than the TCCON at the
European sites, but obtains good agreement elsewhere. At the European sites, the difference extends
up to 2–3 weeks, and is likely connected with the biased amplitude inferred by ACOS discussed be-
325 low. While the maximum occurs within two spring months depending on location, the minimum is
more seasonally restricted, varying from August 15 to September 27 for TCCON, and from August
14 to September 25 for GOSAT. During the minimum, the Northern Hemisphere receives solar light
abundantly and is not snow-covered, so the number of co-located soundings is larger and the mini-
mum is well captured by the satellite, within 6 days from TCCON, except for Tsukuba and Bremen.
330 These values are generally in good agreement (within a few days) with Wunch et al. (2013, p. 9451),
except for the TCCON seasonal cycle maximum date at the European sites Bialystok and Bremen.

~~We~~ However, regarding the difference in the dates of the maximum, Kulawik et al. (2015) found a much smaller phase difference in Europe by calculating cross-correlation of the data points to determine the phase shift. Because our results were based on the fitted seasonal cycles instead of the actual data, we evaluated the statistical errors of the dates of the maximum and minimum XCO₂ with a Monte Carlo approach, using the error covariance matrices associated with the fitted function parameters. On average, the TCCON maximum date had an error of 3.5 days, while the error for ACOS maximum date was 6.1 days for the TCCON maximum date, and 6.1 days for ACOS maximum date, reflecting a notable uncertainty in the fitted phase and thus explaining at least partially the difference between our results and those of Kulawik et al. (2015). The corresponding average statistical errors σ for the date of the minimum were 2.2 days (TCCON) and 3.6 days (ACOS).

The seasonal cycle amplitudes are presented in Fig. 5a, in addition to Table 3. The amplitude is captured within the error bars of the regression at four sites: Izaña, Lamont, Saga, and Park Falls. The largest absolute differences are 1.6 ppm at Tsukuba and 1.1 ppm at Bremen and Orleans, which are also the largest relative differences (28% and 18, 14% and 15%). Within 1.0 ppm difference, the amplitude is captured at most sites, excluding Orleans, Bremen, and Tsukuba all other sites. It should be noted that the latter Tsukuba only has data for two years and therefore substantial uncertainty in both the trend and amplitude, whereas the former two Bremen and Orleans sites have sufficient data for evaluating an average seasonal cycle. A closer inspection of Figs. 4 and 5a reveals that the amplitude seen by GOSAT/ACOS is systematically shallower than TCCON at all five TCCON sites in continental Europe. This bias appears to be regionally very concentrated, because at the Northern European site Sodankylä, GOSAT captures the seasonal cycle reasonably well (within 0.8 ppm), considering the site suffers from data (and sunlight) deficiency in winter. Kulawik et al. (2015) noted the low bias as well, although they grouped all TCCON sites within latitudes 46 – 53°N together and found that, at this latitude range, the seasonal cycle of ACOS was biased low by 0.7 ± 0.7 ppm. Intuitively, a shallow-biased GOSAT seasonal cycle over Europe contradicts with the message from several recent flux inversion studies (??), where the inversions using GOSAT XCO₂ observations inferred a stronger carbon sink over Europe compared to the inversions that assimilated in-situ measurements only, and compared with bottom-up inventories. However, according to the results by ?, these two results are not in a conflict. They showed in their regional flux inversion experiment that the sink enhancement is due to a North-West to South-East gradient in XCO₂ over Europe, and that most of the additional uptake takes place in Eastern Europe.

We explored several possible explanations for the low-biased seasonal cycle amplitude over continental Europe. First, we repeated the analysis using GOSAT/ACOS B3.4 retrievals (instead of B3.5), which have two constant aerosol types in the retrieval, different filtering, and bias correction. This did not have a systematic effect: the seasonal cycle amplitude of GOSAT increased at

Bremen (+0.3 ppm) and Orleans (+0.5 ppm), and decreased at Bialystok (−0.2 ppm), Garmisch
370 (−0.2 ppm), and Karlsruhe (−0.4 ppm).

~~We also studied the differences between TCCON-GGG2012 and GGG2014 retrievals, and found
that in the latter, the seasonal cycle amplitudes in Europe were shallower by up to 0.4 ppm (Orleans).
The difference comes likely from the extended time series and the additional measurements present
in the GGG2014 version. It is therefore possible that some of the discrepancy between GOSAT and
375 Bremen TCCON is due to the use of the GGG2012 retrieval.~~

Next, we introduced variations to the co-location method to quantify its impact to the seasonal
cycle amplitude. Our default co-location technique was the NOAA/Basu method with 0.50 ppm
CO₂ gradient, maximum latitude difference 7.5°, and longitude 22.5°. We experimented with four
modifications to it: 1) latitude 5.0°, longitude 15°, 2) latitude 2.5°, longitude 7.5°, 3) 0.25 ppm CO₂
380 gradient, and 4) 1.0 ppm CO₂ gradient. The latter increased the number of co-located points while
the three former reduced it by making the co-location requirement stricter. We found that a smaller
longitude-latitude box and a tighter CO₂ gradient led to a better match-up in terms of the seasonal
cycle amplitude at Bialystok (difference only 0.1 ppm), but not in other European sites where the
difference either did not change or increased. The ACOS seasonal cycle amplitude at Garmisch site
385 turned out to be highly dependent on the co-location details, varying from 5.0 ppm to 5.9 ppm in
these tests. The TCCON amplitudes changed typically only 0.1 ppm, but the fitting errors increased
as the number of co-located soundings decreased. We also found that the co-location box dimensions
had an impact on the seasonal cycle at JPL, which is located in the Los Angeles basin where large
CO₂ gradients could be expected. With the default technique, the amplitude for ACOS was 0.5 ppm
390 shallower than TCCON (10% difference), but when decreasing the box size, the difference was
reduced to 0.1 ppm (2%).

In our last experiment, we tested the impact of the ACOS B3.5 bias correction for H gain over land;
as Table 3 shows, all co-located soundings at the continental European sites were land gain H. We
found that the bias correction increased the seasonal cycle amplitude at Park Falls by 1.4 ppm, mostly
395 due to a correction for dust aerosol optical depth and surface albedo in the 2.1 μm band, but the bias
correction had only a 0.1 ppm total impact on the amplitude at the European sites. It turned out
that two of the bias correction parameters (related to the retrieved surface pressure and vertical CO₂
gradient) made the seasonal cycle over Europe consistently shallower by 0.3 – 0.4 ppm, depending
on the site (see Fig. 5b). However, these parameters did not affect the seasonal cycle amplitude at
400 Park Falls or Lamont, which are the two main sites used when optimizing the ACOS bias correction.
An interesting finding is that removing these two terms from the bias correction made the ACOS
seasonal cycle amplitude (Fig. 5b) and trend (not shown) agree better with TCCON at 10 of the 12
sites, even though it made the scatter worse in single-sounding statistics. This implies that the bias
correction might be improved by designing it based on aggregated soundings in addition to single
405 observation statistics.

5.2 Evaluation against other retrieval algorithms

To further study the discrepancies of GOSAT and TCCON, we repeated the seasonal cycle analysis for four other retrieval algorithms, taking into account their individual bias corrections: RemoTeC v2.35 (Butz et al., 2011; Guerlet et al., 2013a), University of Leicester (UoL) v5.1 (Cogan et al., 410 2012), NIES PPDF-S v.02.11 (Oshchepkov et al., 2013b), and NIES v02.21 (Yoshida et al., 2013), which is the operational GOSAT retrieval algorithm with the bias correction applied. The seasonal cycle amplitude, the trend, and the days of maximum and minimum (detrended) XCO₂ are presented in Fig. 6 together with their daily averages RMS error with respect to the TCCON fit. RemoTeC had a shorter time series than the other retrievals, and was therefore not included in the Saga, JPL, and 415 Tsukuba results. UoL data did not include glint soundings, which may cause some differences at coastal or island sites. Also, only ACOS and NIES retrievals included a sufficient amount of co-located soundings for successfully fitting a seasonal cycle at Sodankylä.

Overall, the five algorithms performed qualitatively similarly but show notable scatter at most validation sites and in most of the fitted parameters. Also, no algorithm clearly outperforms another. The only systematic difference is that all algorithms except NIES generally capture a smaller 420 mean growth rate than TCCON, whereas NIES retrieves a higher trend. This may be due to different corrections for radiometric degradation in the different algorithms, but could also result from other factors, such as bias correction. For example, NIES v02.21 and NIES PPDF-S v.02.11 have different growth rates despite the use of similar corrections for radiometric degradation. The TCCON 425 seasonal cycle amplitude is captured by GOSAT at almost every site but by a different retrieval: as shown in Sect. 5.1, ACOS has a very good agreement with TCCON at the North American sites as well as Izaña and Saga but, in continental Europe, NIES and NIES PPDF-S perform generally the best. ACOS, RemoTeC, and UoL all show a low-biased amplitude in continental Europe, and NIES, UoL, and NIES PPDF-S are biased high elsewhere. If considering only those sites with longer 430 time series, the scatter between the algorithms is around 1 ppm. ~~These results can be interpreted to support the ensemble median algorithm EMMA introduced by Reuter et al. (2013), which combines all individual retrievals into one data set that globally has the best agreement with TCCON.~~

The maximum and minimum days of the seasonal cycle reflect the drawdown season and are dependent on latitude and climate region. Both TCCON and GOSAT capture an earlier start of 435 drawdown at the continental European sites compared to the other sites, the latest start being at the southernmost site, Izaña. The ACOS and NIES PPDF-S algorithms appear to be generally best in phase with TCCON regarding the date of maximum XCO₂. At the continental European sites, GOSAT and TCCON fits for the maximum day differ by several weeks, TCCON being systematically earlier. The minimum is better captured by all retrievals, with the spread varying from a few 440 days to about 20 days; the performance of the individual algorithms is very site-specific.

Since none of the retrieval algorithms clearly outperformed the others at every TCCON site, we repeated the analysis for the ensemble median algorithm EMMA (Reuter et al., 2013), which

combines all individual retrievals into one data set of median XCO₂ values. Even though EMMA had the smallest RMS error at four TCCON sites overall, it did not perform systematically better or worse than the individual retrieval algorithms in capturing the seasonal cycle of XCO₂.

445

5.3 Evaluation against models

The seasonal cycle amplitude of GOSAT/ACOS B3.5 was also compared to the inverse model systems MACC 13.1, CT2013B, and UoE in the Northern Hemisphere. As described in Sect. 3.2, these models have been optimized against assimilated flask and in-situ CO₂ measurements, though not exactly same data sets nor using the exact same weighting. For the comparison, latitudes from 0° to 70° were divided into 5° latitude bins (see Fig. 1 for the map), and the GOSAT/ACOS soundings within one latitude bin were collected into a single time series. The seasonal cycle was fitted on the daily averages of GOSAT/ACOS XCO₂ and the resampled models. The resulting seasonal cycle amplitudes are shown in Fig. 7. The amplitude increases significantly from the tropics towards high latitudes for both GOSAT and the models. Although the results are qualitatively similar, the models can show close to 2 ppm differences within latitude bands. ACOS is in excellent agreement to MACC from 0°N to 50°N, whereas CT2013B and UoE have a shallower seasonal cycle from the tropics up to 35°N. Differences in the model seasonal cycle can be caused by a number of error sources, including their prior, transport, and inversion. Tropical and subtropical latitudes include large regions where the data constraint is weaker; therefore, the land surface prior (and its particular implementation) may impact the inversion results more than at those regions where the measurement network is dense. Both UoE and CT2013B use a variant of CASA as their prior biospheric flux model, as presented in Table 1 (in fact, CT2013B uses a unique combination of two flavors of CASA (Andy Jacobson, personal communication, April 17, 2015)). Even though different versions of CASA can differ in their seasonal cycle magnitude, our results may imply that the seasonal cycle of CASA fluxes is too shallow in some tropical regions or biomes. We first did the comparison using earlier versions of CarbonTracker (CT2011 and CT2013), and found that CarbonTracker and UoE results were nearly identical in these regions (see CT2013 and UoE in Figs. 7 and 8), which was surprising because the two models were different in every aspect (transport, in-situ data selection, inversion) except for their prior biospheric fluxes. However, a significant correction to the transport model's vertical mixing was introduced in CT2013B. This led to an increase of about 0.5 ppm in the CarbonTracker's seasonal cycle amplitude at all latitudes.

465

470

At 50 – 60°N in Fig. 7, ACOS agrees better with UoE and CT2013B. From 60° to 70°, ACOS has a higher seasonal cycle amplitude than most models. A similar result was also obtained by Belikov et al. (2014) using GOSAT/NIES v02.00 retrievals, NIES transport model, and LMDZ model. However, at high boreal latitudes, the satellite observations are associated with larger errors that are not reflected in the purely statistical fitting errors. ACOS results at these latitudes should therefore be interpreted with caution.

475

We tested how the ACOS bias correction and model averaging kernel correction affected the latitudinally averaged seasonal cycle amplitudes. The ACOS bias correction decreased the amplitude about 0.5 ppm at latitudes 10 – 40°N, but increased the amplitude at 40 – 70°N. The maximum increase was 1.0 ppm at latitudes 50 – 60°N, implying that before the bias correction, ACOS was in better agreement with MACC at these latitudes, but that after the bias correction, ACOS agreed better with UoE and CT2013B. Even though validation against models is part of the ACOS bias correction, the TCCON sites are likely to dominate the bias correction at mid-latitudes. We studied the potential seasonal impact of the averaging kernel correction for CT2013B. We found that the averaging kernel correction systematically decreased the model seasonal cycle amplitude in the Northern Hemisphere by 0.15 ppm on average. Overall, these changes are minor and do not affect our general conclusions about the model comparisons.

The latitudinal dependence of the CO₂ seasonal cycle amplitude has been previously shown in e.g. "the flying carpet" plot presented by Conway et al. (1994, Fig. 4), but we would like to emphasize that the amplitude can also depend on longitude. Especially in the mid-latitudes, its increase from west to east is notable; this is demonstrated in Fig. 8 for latitude band 45 – 50°N, where the seasonal cycle amplitude of GOSAT/ACOS is 6.4 ppm over the longitudes 180W–120W, and is doubled at 120E–180E. The increased seasonal cycle is likely due to the large seasonal sink of the boreal forests, accrued in the total column as the observation point is moved eastward. These GOSAT observations considered were taken over land, so in practice, this means that the seasonal cycle amplitude is dampened from the Eastern Asia over the North Pacific Ocean to the North-West United States. In the lower troposphere, this dampening above 30°N latitude was shown by Nakazawa et al. (1992) who analyzed a three-year time series (1984–1986) of CO₂ measurements onboard container ships. The model results in Fig. 8 show a similar pattern of amplitude enhancement towards east, albeit the seasonal cycle amplitude of MACC is 2–3 ppm shallower compared to those of the other models and ACOS in the Eastern Asia. Despite this large discrepancy in the east where the data volume is small (see Fig. 8, right vertical axis), the zonally-averaged seasonal cycle amplitudes of MACC and ACOS agree within 0.1 ppm at the same latitude band (45 – 50°N). The CT2013B amplitudes are consistently higher than ACOS at all longitudes in Fig. 8, but they agree within 0.1 ppm in the Eastern Asia. Of the three models, UoE is most consistent with ACOS, agreeing about the seasonal cycle amplitude to within 1 ppm at these specific regions. The northern and mid-latitudinal regions of Asia are again regions where the in-situ measurement coverage is very limited, which explains the large spread between the individual model results.

6 Conclusions

The seasonal cycle of XCO₂ is profoundly connected to the biospheric fluxes that determine the global terrestrial net CO₂ sink. Satellite measurements of XCO₂ by the Greenhouse Gases Observ-

ing Satellite (GOSAT) and the Orbiting Carbon Observatory (OCO-2) expand the current in-situ
515 measurement network tremendously and therefore have the potential to improve flux inversions.
However, the satellite-measured seasonal cycle of XCO₂ can be affected by different retrieval bi-
ases, such as biases related to seasonally-varying parameters (e.g., surface albedo) and a sampling
bias due to the seasonal variation in solar radiation. Mischaracterization of the seasonal cycle could
lead to errors in the inverse model systems that assimilate satellite CO₂ data. Motivated by this,
520 we evaluated the seasonal cycle of GOSAT observations using ACOS B3.5 retrievals from years
2009–2013.

Three independent approaches were used for the evaluation of the XCO₂ seasonal cycle: com-
parisons against the Total Carbon Column Observing Network (TCCON), other GOSAT retrievals
(UoL v5.1, NIES v02.21, NIES PPDF-S v.02.11, and RemoTeC v2.35), and comparisons to opti-
525 mized inversion models that assimilate in-situ measurements of CO₂. We found that ACOS captures
the seasonal cycle amplitude of TCCON with an accuracy of better than 1.0 ppm at most of the 12
TCCON sites in the Northern Hemisphere considered in this study. As we also inferred the mean
annual growth rate at each TCCON site in order to remove it, we found agreement of generally better
than 0.2 ppm/year in this quantity, with the ACOS-inferred growth rate most often being lower than
530 TCCON. Over continental Europe, the seasonal cycle amplitude as measured by ACOS was biased
low at all five sites, the largest difference being ~~18% at Bremen~~ 1.1 ppm at Bremen and Orleans.
We also found that ACOS generally captured the seasonal cycle phase within a few days, except
over Europe where the differences were 2–3 weeks, with ACOS measuring the date of maximum
XCO₂ later than TCCON. Several other algorithms also had minor low biases in their seasonal cy-
535 cle amplitudes over Europe. We explored the cause of the low bias for ACOS, and found that the
bias correction parameters related to the retrieved surface pressure and vertical CO₂ gradient were
partially responsible, explaining 16 – 48% of the difference. This suggests that the bias correction
might benefit from considering aggregated soundings in addition to deviations at single-sounding
level. Also, the selection of the co-located soundings was found to affect the seasonal cycle ampli-
540 tude at few sites. Especially at JPL, which is in the Los Angeles basin, the agreement with TCCON
improved notably when the co-location criteria were made sufficiently tight to not include soundings
taken too far from the basin.

Model comparisons at latitudes 0 – 70°N revealed that qualitatively the models and satellite obser-
vations agreed well, but also that the model-to-model differences were (at most latitude bands stud-
545 ied) larger than model-to-ACOS differences. From the tropics up to 50°N, the zonally-averaged sea-
sonal cycle amplitude of ACOS was in very good agreement with MACC 13.1, while between 50 –
60°N, ACOS agreed better with the University of Edinburgh model and CarbonTracker CT2013B.
Both of the latter models had seasonal cycle amplitudes shallower than ACOS or MACC at tropical
and subtropical latitudes, where the models lack direct constraints from measurements over land and
550 are thus more affected by their prior fluxes (or by extra-tropical or ocean measurements through

long-range transport). Therefore, the shallower seasonal cycle amplitude might be connected to their prior land surface models that are different variants of CASA. However, to verify this, one should investigate also the impact of transport, data assimilation, and inversion system differences. We also found that the longitudinal changes in the seasonal cycle amplitude at mid-latitudes can be notable.

555 In particular, we showed that at 45–50°N latitudes, the amplitude of the GOSAT XCO₂ seasonal cycle doubles from the North-West U.S. to Eastern Asia. The model results showed a gradient as well, although it was 1–3 ppm shallower, depending on the model. We also noticed that the averaging kernel correction can systematically decrease the seasonal cycle amplitude by up to 0.2 ppm.

560 Based on our study, the GOSAT/ACOS seasonal cycle error is of the order of 1.0 ppm near TCCON stations and likely to be of this size in other parts of the world, though may be influenced by the a priori accuracy of jointly retrieved parameters, such as those related to aerosols. As model-to-model differences in the XCO₂ seasonal cycle amplitude can be several ppm at regions poorly sampled by in-situ measurements, GOSAT observations ~~that measure seasonal cycle amplitude to within 1.0 ppm, based on this study,~~ could potentially be used directly (without elaborate inversions) 565 to evaluate model differences at these regions. This idea is explored in more detail in a work under preparation (Lindqvist et al., 2015, in prep.).

Acknowledgements. HL and CO wish to acknowledge support from the NASA Jet Propulsion Laboratory and the OCO-2 project, via JPL subcontract number 1439002. HL also acknowledges funding from the Academy of Finland, via project number 285421. HB and RP acknowledge the National Centre for Earth Observation 570 NCEO and the ESA GHG-CCI project. FC acknowledges the European Commission (grant agreement No. 630080, MACC III). We wish to thank the referees (Susan Kulawik and an anonymous referee) and Andy Jacobson for ~~his~~their comments and constructive feedback to the manuscript. We also gratefully acknowledge all the data providers. TCCON data were obtained from the TCCON Data Archive, operated by the California Institute of Technology from the website at <http://tcon.ornl.gov/>. TCCON work at Garmisch has been funded 575 by the ESA GHG-cci project via subcontract with the University of Bremen and by the EC within the INGOS project. CarbonTracker CT2013B results were provided by NOAA ESRL, Boulder, Colorado, USA from the website at <http://carbontracker.noaa.gov>. Finally, we thank our colleagues for the use of their GOSAT retrievals: André Butz and Otto Hasekamp for RemoTeC v2.35, and Andrey Bril for NIES PPDF-S v.02.11.

References

- 580 ~~Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., Krummel, P., Steele, P.,
Langenfelds, R., Torn, M., Biraud, S., Stephens, B., Andrews, A., and Worthy, D.: Global CO₂
fluxes estimated from GOSAT retrievals of total column CO₂, *Atmos. Chem. Phys.*, 13, 8695–8717,
doi:10.5194/acp-13-8695-2013, 2013.~~ Belikov, D. A., Bril, A., Maksuytov, S., Oshchepkov, S., Saeki,
T., Takagi, H., Yoshida, Y., Ganshin, A., Zhuravlev, R., Aoki, S., and Yokota, T.: Column-averaged
585 CO₂ concentrations in the subarctic from GOSAT retrievals and NIES transport model simulations,
Polar Science, 8, 129–145, 2014. Blumenstock, T., Hase, F., Schneider, M., García, O. E., and
Sepúlveda, E.: TCCON data from Izaña, Tenerife, Spain, Release GGG2014R0, TCCON data archive,
hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak
Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.izana01.R0/1149295>, 2014. [Notholt,
590 J., Petri, C., Warneke, T., Deutscher, N., Buschmann, M., Weinzierl, C., Macatangay, R., Grupe, P.:
TCCON data from Bremen, Germany, Release GGG2014R0, TCCON data archive, hosted by the
Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee,
U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.bremen01.R0/1149275>, 2014.](http://dx.doi.org/10.14291/tccon.ggg2014.bremen01.R0/1149275) Buchwitz, M., Reuter, M.,
Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante R., Bergamaschi, P., Blumenstock, T.,
595 Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J. P., Butz, A., Chedin, A., Chevallier, F., Crevoisier,
C. D., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., Kaminski, T., Laeng,
A., Lichtenberg, G., De Maziere, M., Noel, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M.,
Sussmann, R., Stiller, G. P., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D. W. T., Kuze, A.,
O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., and Yoshida,
600 Y.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparison and quality assessment of
near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, *Remote Sensing of Environment*,
doi:10.1016/j.rse.2013.04.024, pp. 19, (in press), 2013. Butz, A., Guerlet, S., Hasekamp, O., Schepers,
D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G.,
Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and
605 Warneke, T.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophys. Res. Lett.*, 38, L14812,
doi:10.1029/2011GL047888, 2011. Chevallier, F., Breon, F.-M., and Rayner, P. J.: Contribution of the Orbit-
ing Carbon Observatory to the estimation of CO₂ sources and sinks: Theoretical study in a variational data
assimilation framework, *J. Geophys. Res.*, 112, D09307, doi:10.1029/2006JD007375, 2007. Chevallier, F.,
Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröh-
610 lich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M.,
Machida, T., Maignan, F., Matsueda, H., Morgui, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M.,
Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.: CO₂
surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *J.*
Geophys. Res., 115, D21307, doi:10.1029/2010JD013887, 2010. Chevallier, F., Deutscher, N. M., Conway,
615 T. J., Ciais, P., Ciattaglia, L., Dohe, S., Fröhlich, M., Gomez-Pelaez, A. J., Griffith, D., Hase, F., Haszpra, L.,
Krummer, P., Kyrö, E., Labuschagne, C., Langenfelds, R., Machida, T., Maignan, F., Matsueda, H., Morino,
I., Notholt, J., Ramonet, M., Sawa, Y., Schmidt, M., Sherlock, V., Steele, P., Strong, K., Sussmann, R.,
Wennberg, P., Wofsy, S., Worthy, D., Wunch, D., and Zimnoch, M.: Global CO₂ fluxes inferred from surface

air-sample measurements and from TCCON retrievals of the CO₂ total column, *Geophys. Res. Lett.*, 38, L24810, doi:10.1029/2011GL049899, 2011. [Chevallier, F., Palmer, P. I., Feng, L., Boesch, H., O'Dell, C. W., and Bousquet, P.: Toward robust and consistent regional CO₂ flux estimates from in-situ and spaceborne measurements of atmospheric CO₂, *Geophys. Res. Lett.*, 41, 1065–1070, doi:10.1002/2013GL058772, 2014.](#)

Cogan, A. J., Boesch, H., Parker, R. J., Feng, L., Palmer, P. I., Blavier, J.-F. L., Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., Warneke, T., and Wunch, D.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): Comparison with ground-based TCCON observations and GEOS-Chem model calculations, *J. Geophys. Res.*, 117, D21301, doi:10.1029/2012JD018087, 2012.

Conway, T. J., Tans, P. P., Waterman, L. S., Thoning, K. W., Kitzis, D. R., Masarie, K. A., and Zhang, N.: Evidence for interannual variability of the carbon cycle from the National Oceanic and Atmospheric Administration/Climate Monitoring and Diagnostics Laboratory Global Air Sampling Network, *J. Geophys. Res. Atmos.*, 99, D11, 22831–22855, doi:10.1029/94JD01951, 1994. Crisp, D., Atlas, R. M., Breon, F.-M., Brown, L. R., Burrows, J. P., Ciais, P., Connor, B. J., Doney, S. C., Fung, I. Y., Jacob, D. J., Miller, C. E., O'Brien, D., Pawson, S., Randerson, J. T., Rayner, P., Salawitch, R. J., Sander, S. P., Sen, B., Stephens, G. L., Tans, P. P., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Yung, Y. L., Kuang, Z., Chudasama, B., Sprague, G., Weiss, B., Pollock, R., Kenyon, D., and Schroll, S.: The orbiting carbon observatory (OCO) mission, *Advances in Space Research*, 34(4), 700–709, 2004. Crisp, D., Fisher, B., O'Dell, C., Frankenberg, C., Basilio, R., Bösch, H., Brown, L. R., Castano, R., Connor, B., Deutscher, N. M., Eldering, A., Griffith, D., Gunson, M., Kuze, A., Mandrake, L., McDuffie, J., Messerschmidt, J., Miller, C. E., Morino, I., Natraj, V., Notholt, J., O'Brien, D. M., Oyafuso, F., Polonsky, I., Robinson, J., Salawitch, R., Sherlock, V., Smyth, M., Suto, H., Taylor, T. E., Thompson, D. R., Wennberg, P. O., Wunch, D., and Yung, Y. L.: The ACOS CO₂ retrieval algorithm – Part II: Global XCO₂ data characterization, *Atmos. Meas. Tech.*, 5 (4), 687–707, doi:10.5194/amt-5-687-2012, 2012. Deutscher, N., Notholt, J., Messerschmidt, J., Weinzierl, C., Warneke, T., Petri, C., Grupe, P., and Katrynski, K.: TCCON data from Bialystok, Poland, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.bialystok01.R0/1149277>, 2014. Dils, B., Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Parker, R., Guerlet, S., Aben, I., Blumenstock, T., Burrows, J. P., Butz, A., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., De Maziere, M., Notholt, J., Sussmann, R., Warneke, T., Griffith, D., Sherlock, V., and Wunch, D.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, *Atmos. Meas. Tech.*, 7, 1723–1744, 2014. Feng, L., Palmer, P. I., Boesch, H., and Dance, S.: Estimating surface CO₂ fluxes from space-borne CO₂ dry air mole fraction observations using an ensemble Kalman Filter, *Atmos. Chem. Phys.*, 9, 2619–2633, doi:10.5194/acp-9-2619-2009, 2009. Feng, L., Palmer, P. I., Yang, Y., Yantosca, R. M., Kawa, S. R., Paris, J.-D., Matsueda, H., and Machida, T.: Evaluating a 3-D transport model of atmospheric CO₂ using ground-based, aircraft, and space-borne data, *Atmos. Chem. Phys.*, 11, 2789–2803, doi:10.5194/acp-11-2789-2011, 2011. Guerlet, S., Basu, S., Butz, A., Krol, M., Hahne, P., Houweling, S., Hasekamp, O. P., and Aben, I.: Reduced carbon uptake during the 2010 Northern Hemisphere summer from GOSAT, *Geophys. Res. Lett.*, 40(10), 2378–2383, doi:10.1002/grl.50402, 2013b. Guerlet, S., Butz, A., Schepers, D., Basu, S., Hasekamp, O. P., Kuze, A., Yokota, T., Blavier, J.-F., Deutscher, N. M., Griffith, D. W.

T., Hase, F., Kyrö, E., Morino, I., Sherlock, V., Sussmann, R., Galli, A., and Aben, I.: Impact of aerosol and thin cirrus on retrieving and validating XCO₂ from GOSAT shortwave infrared measurements, *J. Geophys. Res. Atmos.*, 118, 4887–4905, doi:10.1002/jgrd.50332, 2013a. Hase, F., Blumenstock, T., Dohe, S., Groß, J., and Kiel, M.: TCCON data from Karlsruhe, Germany, Release ~~GGG2014R0~~[GGG2014R1](https://doi.org/10.14291/tccon.ggg2014.karlsruhe01.R0R1/H1492701182416), TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.karlsruhe01.R0R1/H1492701182416>, 2014. Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Taura, C., Fukamachi, T., Sakashita, M.: TCCON data from Saga, Japan, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.saga01.R0/1149283>, 2014. Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdahl, C. A., Guenther, P. R., Waterman, L. S., and Chin, J. F.: Atmospheric carbon dioxide variations at Mauna Loa observatory, Hawaii, *Tellus*, 28(6), 538–551, 1976. Keppel-Aleks, G., Wennberg, P. O., and Schneider, T.: Sources of variations in the total column carbon dioxide, *Atmos. Chem. Phys.*, 11, 3581–3593, doi:10.5194/acp-11-3581-2011, 2011. Keppel-Aleks, G., Wennberg, P. O., Washenfelder, R. A., Wunch, D., Schneider, T., Toon, G. C., Andres, R. J., Blavier, J.-F., Connor, B., Davis, K. J., Desai, A. R., Messerschmidt, J., Notholt, J., Roehl, C. M., Sherlock, V., Stephens, B. B., Vay, S. A., and Wofsy, S. C.: The imprint of surface fluxes and transport on variations in total column carbon dioxide, *Biogeosciences*, 9(3), 875–891, doi:10.5194/bg-9-875-2012, 2012. Kivi, R., Heikkinen, P., and Kyrö, E.: TCCON data from Sodankylä, Finland, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.sodankyla01.R0/1149280>, 2014. Kulawik, S. ~~S.~~, Wunch, D., O'Dell, C. ~~W.~~, Frankenberg, C., ~~Jacobson, A. R.~~, Reuter, M., Oda, T., Chevallier, F., Sherlock, V., Buchwitz, M., Osterman, G., ~~and~~ Miller, C., [Wennberg, P., Griffith, D. W. T., Morino, I., Dubey, M., Deutscher, N. M., Notholt, J., Hase, F., Warneke, T., Sussmann, R., Robinson, J., Strong, K., Schneider, M., and Wolf, J.](https://doi.org/10.5194/amtd-8-6217-2015): Consistent evaluation of GOSAT, SCIAMACHY, CarbonTracker, and MACC through comparisons to TCCON, *Atmos. Chem. Phys. Meas. Tech. Discuss.*, ~~submitted in Feb.~~ [8, 6217–6277, doi:10.5194/amtd-8-6217-2015](https://doi.org/10.5194/amtd-8-6217-2015), 2015. Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the greenhouse gases observing satellite for greenhouse gases monitoring, *Appl. Opt.*, 48, 6716–6733, 2009. Kuze, A., Taylor, T. E., Kataoka, F., Bruegge, C. J., Crisp, D., Harada, M., Helmlinger, M., Inoue, M., Kawakami, S., Kikuchi, N., Mitomi, Y., Murooka, J., Naitoh, M., O'Brien, D. M., O'Dell, C. W., Ohyama, H., Pollock, H., Schwandner, F. M., Shiomi, K., Suto, H., Takeda, T., Tanaka, T., Urabe, T., Yokota, T., and Yoshida, Y.: Long-term vicarious calibration of GOSAT short-wave sensors: Techniques for error reduction and new estimates of radiometric degradation factors, *Geoscience and Remote Sensing, IEEE Transactions on*, 52(7), 3991–4004, 2014. Lindqvist, H., O'Dell, C., Schuh, A., Baker, D., Cheeseman, M., Baker, I., Haynes, K., and Denning, S.: The potential of GOSAT XCO₂ to evaluate land surface models, In preparation, 2015. Liu, J., Bowman, K. W., Lee, M., Henze, D. K., Bousserez, N., Brix, H., Collatz, G. J., Menemenlis, D., Ott, L., Pawson, S., Jones, D., and Nassar, R.: Carbon monitoring system flux estimation and attribution: impact of ACOS-GOSAT XCO₂ sampling on the inference of terrestrial biospheric sources and sinks, *Tellus B*, 66, doi:<http://dx.doi.org/10.3402/tellusb.v66.22486>, 2014. Maksuytov, S., Takagi, H., Valsala, V. K., Saito, M.,

Oda, T., Saeki, T., Belikov, D. A., Saito, R., Ito, A., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., and
700 Yokota, T.: Regional CO₂ flux estimates for 2009–2010 based on GOSAT and ground-based CO₂ observa-
tions, *Atmos. Chem. Phys.*, 13, 9351–9373, doi:10.5194/acp-13-9351-2013, 2013. Messerschmidt, J., Chen,
H., Deutscher, N. M., Gerbig, C., Grube, P., Katrynski, K., Koch, F.-T., Lavrič, J. V., Notholt, J., Roden-
beck, C., Ruhe, W., Warneke, T., and Weinzierl, C.: Automated ground-based remote sensing measurements
of greenhouse gases at the Bialystok site in comparison with collocated in situ measurements and model
705 data, *Atmos. Chem. Phys.*, 12(15), 6741–6755, doi:10.5194/acp-12-6741-2012, 2012. Messerschmidt, J.,
Geibel, M. C., Blumenstock, T., Chen, H., Deutscher, N. M., Engel, A., Feist, D. G., Gerbig, C., Gisi, M.,
Hase, F., Katrynski, K., Kolle, O., Lavrič, J. V., Notholt, J., Palm, M., Ramonet, M., Rettinger, M., Schmidt,
M., Sussmann, R., Toon, G. C., Truong, F., Warneke, T., Wennberg, P. O., Wunch, D., and Xueref-Remy,
I.: Calibration of TCCON column-averaged CO₂: the first aircraft campaign over European TCCON sites,
710 *Atmos. Chem. Phys.*, 11, 10765–10777, doi:10.5194/acp-11-10765-2011, 2011. Morino, I., Matsuzaki, T.,
Ikegami, H., and Shishime, A.: TCCON data from Tsukuba, Ibaraki, Japan, 125HR, Release GGG2014R0,
TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Lab-
oratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.tsukuba02.R0/1149301>,
2014. Morino, I., Uchino, O., Inoue, M., Yoshida, Y., Yokota, T., Wennberg, P. O., Toon, G. C., Wunch,
715 D., Roehl, C. M., Notholt, J., Warneke, T., Messerschmidt, J., Griffith, D. W. T., Deutscher, N. M., Sherlock,
V., Connor, B., Robinson, J., Sussmann, R., and Rettinger, M.: Preliminary validation of column-averaged
volume mixing ratios of carbon dioxide and methane retrieved from GOSAT short-wavelength infrared spec-
tra, *Atmos. Meas. Tech.*, 4, 1061–1076, doi:10.5194/amt-4-1061-2011, 2011. Nakazawa, T., Murayama, S.,
Miyashita, K., Aoki, S., and Tanaka, M.: Longitudinally different variations of lower tropospheric carbon
720 dioxide concentrations over the North Pacific Ocean, *Tellus*, 44B, 161–172, 1992. Nguyen, H., Osterman,
G., Wunch, D., O'Dell, C., Mandrake, L., Wennberg, P., Fisher, B., and Castano, R.: A method for colo-
cating satellite XCO₂ data to ground-based data and its application to ACOS-GOSAT and TCCON, *Atmos.*
Meas. Tech., 7, 2631–2644, doi:10.5194/amt-7-2631-2014, 2014. O'Dell, C. W., Connor, B., Bösch, H.,
O'Brien, D., Frankenberg, C., Castano, R., Christi, M., Eldering, D., Fisher, B., Gunson, M., McDuffie, J.,
725 Miller, C. E., Natraj, V., Oyafuso, F., Polonsky, I., Smyth, M., Taylor, T., Toon, G. C., Wennberg, P. O.,
and Wunch, D.: The ACOS CO₂ retrieval algorithm – Part 1: Description and validation against synthetic
observations, *Atmos. Meas. Tech.*, 5(1), 99–121, doi:10.5194/amt-5-99-2012, 2012. Ohshima, H., Morino,
I., Nagahama, T., Machida, T., Suto, H., Oguma, H., Sawa, Y., Matsueda, H., Sugimoto, N., Nakane, H., and
Nakagawa, K.: Column-averaged volume mixing ratio of CO₂ measured with ground-based Fourier trans-
730 form spectrometer at Tsukuba, *J. Geophys. Res.*, 114, D18303, doi:10.1029/2008JD011465, 2009. Olsen,
S. C. and Randerson, J. T.: Differences between surface and column atmospheric CO₂ and implications for
carbon cycle research, *J. Geophys. Res.*, 109, D02301, doi:10.1029/2003JD003968, 2004. Oshchepkov, S.,
Bril, A., Yokota, T., Wennberg, P. O., Deutscher, N. M., Wunch, D., Toon, G. C., Yoshida, Y., O'Dell, C.
W., Crisp, D., Miller, C. E., Frankenberg, C., Butz, A., Aben, I., Guerlet, S., Hasekamp, O., Boesch, H.,
735 Cogan, A., Parker, R., Griffith, D., Macatangay, R., Notholt, J., Sussmann, R., Rettinger, M., Sherlock, V.,
Robinson, J., Kyrö, E., Heikkinen, P., Feist, D. G., Morino, I., Kadyrov, N., Belikov, D., Maksuytov, S.,
Matsunaga, T., Uchino, O., and Watanabe, H.: Effects of atmospheric light scattering on spectroscopic ob-
servations of greenhouse gases from space. Part 2: Algorithm intercomparison in the GOSAT data processing

for CO₂ retrievals over TCCON sites, *J. Geophys. Res. Atmos.*, 118(3), 1493–1512, doi:10.1002/jgrd.50146,

740 2013a. Oshchepkov, S., Bril, A., Yokota, T., Yoshida, Y., Blumenstock, T., Deutscher, N. M., Dohe, S., Macatangay, R., Morino, I., Notholt, J., Rettinger, M., Petri, C., Schneider, M., Sussman, R., Uchino, O., Velasco, V., Wunch, D., and Belikov, D.: Simultaneous retrieval of atmospheric CO₂ and light path modification from space-based spectroscopic observations of greenhouse gases: Methodology and application to GOSAT measurements over TCCON sites, *Appl. Opt.*, 52, 1339–1350, 2013b. Palmer, P. I., Barkley,

745 M. P., and Monks, P. S.: Interpreting the variability of space-borne CO₂ column-averaged volume mixing ratios over North America using a chemistry transport model, *Atmos. Chem. Phys.*, 8, 5855–5868, 2008. Peng, S., Ciais, P., Chevallier, F., Peylin, P., Cadule, P., Sitch, S., Piao, S., Ahlström, A., Huntingford, C., Levy, P., Li, X., Liu, Y., Lomas, M., Poulter, B., Viovy, N., Wang, T., Wang, X., Zaehle, S., Zeng, N., Zhao, F., and Zhao, H.: Benchmarking the seasonal cycle of CO₂ fluxes simulated by terrestrial ecosystem models, *Global Biogeochem. Cycles*, 29, 46–64, doi:10.1002/2014GB004931, 2015. Peters, W., Jacobson, A.

750 R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L. M. P., Petron, G., Hirsch, A. I., Worthy, D. E. J., van der Werf, G. R., Randerson, J. T., Wennberg, P. O., Krol, M. C., and Tans, P. P.: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker, *PNAS*, November 27, 2007, 104(48), 18925–18930, 2007. Pitman, A. J.: The evolution of, and revolution in, land surface schemes designed for climate models, *Int. J. Climatol.*, 23, 479–510, doi: 10.1002/joc.893, 2003. Rayner, P. J. and O’Brien, D. M.: The utility of remotely sensed CO₂ concentration data in surface source inversions, *Geophys. Res. Lett.*, 28(1), 175–178, 2001. ~~Reuter, M., Buchwitz, M., Hilker, M., Heymann, J., Schneising, O., Pillai, D., Bovensmann, H., Burrows, J. P., Bösch, H., Parker, R., Butz, A., Hasekamp, O., O’Dell, C. W., Yoshida, Y., Gerbig, C., Nehrkorn, T., Deutscher, N. M., Warneke, T., Notholt, J., Hase,~~

760 ~~F., Kivi, R., Sussmann, R., Machida, T., Matsueda, H., and Sawa, Y.: Satellite-inferred European carbon sink larger than expected, *Atmos. Chem. Phys.*, 14, 13739–13753, doi:10.5194/acp-14-13739-2014, 2014.~~ Reuter, M., Bösch, H., Bovensmann, H., Bril, A., Buchwitz, M., Butz, A., Burrows, J. P., O’Dell, C. W., Guerlet, S., Hasekamp, O., Heymann, J., Kikuchi, N., Oshchepkov, S., Parker, R., Pfeifer, S., Schneising, O., Yokota, T., and Yoshida, Y.: A joint effort to deliver satellite retrieved atmospheric CO₂ concentrations for surface flux inversions: the ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771–1780, doi:10.5194/acp-13-1771-2013, 2013. Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *J. Geophys. Res.*, 108, D3, 4116, doi:10.1029/2002JD002299, 2003. Sussmann, R. and Rettinger, M.: TCCON data from Garmisch, Germany, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee,

770 U.S.A. <http://dx.doi.org/10.14291/tccon.ggg2014.garmisch01.R0/1149299>, 2014. Takagi, H., Houweling, S., Andres, R. J., Belikov, D., Bril, A., Bösch, H., Butz, A., Guerlet, S., Hasekamp, O., Maksyutov, S., Morino, I., Oda, T., O’Dell, C. W., Oshchepkov, S., Parker, R., Saito, M., Uchino, O., Yokota, T., Yoshida, Y., and Valsala, V.: Influence of differences in current GOSAT XCO₂ retrievals on surface flux estimation, *Geophys. Res. Lett.*, 41, 2598–2605, doi:10.1002/2013GL059174, 2014. Takagi, H., Sacki, T., Oda, T., Saito, M., Valsala, V., Belikov, D., Saito, R., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., Yokota, T., and Maksyutov, S.: On the benefit of GOSAT observations to the estimation of regional CO₂ fluxes, *SOLA*, 7, 161–164, doi:10.2151/sola.2011-041, 2011. Taylor, T. E., O’Dell, C. W., O’Brien, D. M., Kikuchi, N., Yokota, T., Nakajima, T. Y., Ishida, H., Crisp, D., and Nakajima, T.: Comparison of cloud-screening meth-

ods applied to GOSAT near-infrared spectra, *Geoscience and Remote Sensing, IEEE Transactions on*, 50(1), 295–309, doi:10.1109/TGRS.2011.2160270, 2012. Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N., Petri, C., Grupe, P., Vuillemin, C., Truong, F., Schmidt, M., Ramonet, M., and Parmentier, E.: TCCON data from Orleans, France, Release GGG2014R0. TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.orleans01.R0/1149276>, 2014. Washenfelder, R. A., Toon, G. C., Blavier, J.-F. L., Yang, Z., Allen, N. T., Wennberg, P. O., Vay, S. A., Matross, D. M., and Daube, B. C.: Carbon dioxide column abundances at the Wisconsin Tall Tower site, *J. Geophys. Res.*, 111(D22), 1–11, doi:10.1029/2006JD007154, 2006. Wennberg, P. O., Roehl, C., Blavier, J.-F., Wunch, D., Landeros, J., and Allen, N.: TCCON data from Jet Propulsion Laboratory, Pasadena, California, USA, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.jpl02.R0/1149297>, 2014a. Wennberg, P. O., Roehl, C., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R., Keppel-Aleks, G., Allen, N., Ayers, J.: TCCON data from Park Falls, Wisconsin, USA, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.parkfalls01.R0/1149161>, 2014b. Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., Allen, N., Dowell, P., Teske, K., Martin, C., and Martin, J.: TCCON data from Lamont, Oklahoma, USA, Release GGG2014R0, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.lamont01.R0/1149159>, 2014c. Wunch, D., Toon, G. C., Blavier, J. F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The total carbon column observing network, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369(1943), 2087–2112, doi:10.1098/rsta.2010.0240, 2011a. Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jimenez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, *Atmos. Meas. Tech.*, 3, 1351–1362, doi:10.5194/amt-3-1351-2010, 2010. Wunch, D., Wennberg, P. O., Messerschmidt, J., Parazoo, N. C., Toon, G. C., Deutscher, N. M., Keppel-Aleks, G., Roehl, C. M., Randerson, J. T., Warneke, T., and Notholt, J.: The covariation of Northern Hemisphere summertime CO₂ with surface temperature in boreal regions, *Atmos. Chem. Phys.*, 13, 9447–9459, doi:10.5194/acp-13-9447-2013, 2013. Wunch, D., Wennberg, P. O., Toon, G. C., Connor, B. J., Fisher, B., Osterman, G. B., Frankenberg, C., Mandrake, L., O’Dell, C., Ahonen, P., Biraud, S. C., Castano, R., Cressie, N., Crisp, D., Deutscher, N. M., Eldering, A., Fisher, M. L., Griffith, D. W. T., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyrö, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C. E., Morino, I., Notholt, J., Oyafuso, F. A., Rettinger, M., Robinson, J., Roehl, C. M., Salawitch, R. J., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D. R., Uchino, O., Warneke, T., and Wofsy, S. C.: A method for evaluating bias in global measurements of CO₂ total columns from space, *Atmos.*

Chem. Phys., 11, 12317–12337, doi:10.5194/acp-11-12317-2011, 2011b. Yokota, T., Yoshida, Y., Eguchi,
820 N., Ota, Y., Tanaka, T., Watanabe, H., and Maksuytov, S.: Global concentrations of CO₂ and CH₄ retrieved
from GOSAT: first preliminary results, *Sola*, 5, 160–163, 2009. Yoshida, Y., Kikuchi, N., Morino, I., Uchino,
O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P.
O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B.,
825 Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe,
S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their
validation using TCCON data, *Atmos. Meas. Tech.*, 6 (6), 1533–1547, 2013.




Figure 1. Twelve Northern Hemisphere TCCON sites used for GOSAT validation in this study.




Figure 2. All co-located GOSAT/ACOS B3.5 soundings around (a) An example of the TCCON GOSAT-TCCON co-locations using the NOAA/Basu technique (Guerlet et al., 2013a) at Park Falls TCCON station (Wisconsin, USA). The co-location technique was NOAA All GOSAT/Basu ACOS soundings from 8-11 Aug 2009 are shown with 0.50 ppm CO filled circles. The dynamical criterion based on the modelled XCO₂ gradient, latitude limit of 7.5°, fields and longitude limit a 0.5 ppm tolerance from the value at the TCCON location limits the number of 22.5°; see Guerlet et al. (2013a) for details of co-located satellite soundings (green circles). The soundings marked with yellow symbols did not pass the method co-location criteria. (b) All co-located GOSAT/ACOS soundings from Apr 2009 to Dec 2013 at the Park Falls TCCON, coloured according to the month of observation.




Figure 3. An example of data processing and the seasonal cycle fitting procedure at Park Falls. The upper left panel shows time series of the retrieved XCO₂ for all co-located TCCON (black) and GOSAT/ACOS (pink) soundings. The upper right figure shows only those ACOS L2 soundings that pass the post-processing filters. The lower left figure has bias correction applied for ACOS data and averaging kernel correction considered for TCCON soundings. The lower right panel shows the daily averages of XCO₂ and the respective seasonal cycle fits.

Table 1. Models used in the evaluation of the GOSAT seasonal cycle.

Model	Biosphere	Transport	Resolution of the model run (lon x lat x time x layers)
CT2013B	CASA/GFED2 and CASA/GFED3.1	TM5 / ERA-interim, ECMWF	3° x 2° x 3 h x 25
UoE	CASA/GFED	GEOS-Chem / GEOS5	5° x 4° x 3 h x 47
MACC 13.1	ORCHIDEE	LMZD / ECMWF	3.75° x 1.9° x 3 h x 39

Table 2. [Parameters defining the fitted seasonal cycle functions of co-located TCCON and ACOS soundings at Park Falls.](#)

Retrieval	a_0 (ppm)	a_1 (ppm/day)	a_2 (ppm)	a_3 (days)	a_4	a_5 (days)
TCCON	384.5	0.006050	-4.224	-111.4	0.6803	-307.9
ACOS	384.8	0.005904	-4.311	-112.2	0.7585	-268.5




Figure 4. Detrended, best-fit seasonal cycles for GOSAT/ACOS (pink) and TCCON (black) at 12 validation sites in the Northern Hemisphere. The sites are organized according to their latitude (Sodankylä highest, Izaña lowest, Sodankylä highest). The dashed lines depict the times of year with zero or little co-located soundings. On the vertical axis, one tick interval corresponds to 1.0 ppm XCO₂.




Figure 5. Seasonal cycle amplitude for ACOS (vertical axis) and TCCON (horizontal axis) for all the 12 NH sites used in the validation. The dashed black line corresponds to the one-to-one line, and the gray lines denote ± 1.0 ppm. Panel (a) shows the standard bias-corrected ACOS B3.5, and Panel (b) shows ACOS B3.5 with a modified bias correction (see Sect. 5.1 for details).




Figure 6. Comparison of the GOSAT and TCCON XCO₂ time series using the following parameters: root-mean-square (RMS) error (upper left panel), average trend (middle left panel), seasonal cycle amplitude (middle right panel), and the days of maximum and minimum XCO₂ (bottom row). Five retrieval algorithms were included to describe GOSAT observations. TCCON values were based on ACOS B3.5 co-located soundings. The 12 Northern Hemisphere validation sites are shown on the horizontal axis, their latitude increasing from left to right.




Figure 7. Latitudinal dependence of the seasonal cycle amplitude for bias-corrected ACOS B3.5 soundings and for three models resampled at the satellite soundings. For CarbonTracker, we show both CT2013 and CT2013B results, their difference being a major correction in the TM5 transport model. The left vertical axis shows the seasonal cycle amplitude in ppm, while the right vertical axis indicates the number of soundings that fall within each 5° latitude band.




Figure 8. Longitudinal dependence of the seasonal cycle amplitude within the latitude band $45 - 50^\circ\text{N}$. The left vertical axis shows the seasonal cycle amplitude in ppm, while the right vertical axis indicates the number of soundings that fall within each 60° longitude bin. [This latitude zone is highlighted in the world map where also the locations of the continents can be seen.](#)

Table 3. Parameters describing the XCO₂ seasonal cycle for TCCON and bias-corrected GOSAT/ACOS B3.5. The fraction of gain H soundings over land is also shown. The validation sites are sorted according to their latitude.

Site	Time series (month/year)	Retrieval	Growth rate (ppm/year)	Amplitude (ppm)	Date of max. XCO ₂	Date of min. XCO ₂	Fraction land
Izaña	5/2009–10/2013	TCCON	2.41 ± 0.02	5.3 ± 0.1	May 16	Sep 19	
		GOSAT	2.22 ± 0.04	5.3 ± 0.2	May 18	Sep 17	12
Saga	8/2011–10/2013	TCCON	2.39 ± 0.09	6.7 ± 0.2	May 7	Sep 13	
		GOSAT	1.92 ± 0.26	6.7 ± 0.4	Apr 28	Sep 14	77
JPL	5/2011–6/2013	TCCON	2.34 ± 0.07	5.1 ± 0.2	May 2	Sep 27	
		GOSAT	2.39 ± 0.11	4.6 ± 0.3	May 21	Sep 25	87
Tsukuba	8/2011–12/2013	TCCON	2.58 ± 0.10	5.7 ± 0.2	Apr 23	Sep 10	
		GOSAT	2.20 ± 0.22	7.3 ± 0.5	Apr 23	Aug 26	91
Lamont	4/2009–12/2013	TCCON	2.33 ± 0.02	5.3 ± 0.1	May 4	Sep 20	
		GOSAT	2.14 ± 0.03	5.2 ± 0.1	May 6	Sep 15	96
Park Falls	4/2009–12/2013	TCCON	2.21 ± 0.03	8.4 ± 0.1	Apr 22	Aug 15	
		GOSAT	2.16 ± 0.04	8.6 ± 0.2	Apr 27	Aug 14	10
Garmisch	5/2009–10/2013	TCCON	2.03 ± 0.04	6.6 ± 0.1	Mar 25	Aug 27	
		GOSAT	1.90 ± 0.07	5.7 ± 0.2	Apr 17	Aug 24	10
Orleans	8/2009–11/2013	TCCON	2.29 ± 0.04	7.3 ± 0.1	Mar 30	Aug 28	
		GOSAT	2.04 ± 0.07	6.2 ± 0.3	Apr 13	Aug 22	10
Karlsruhe	4/ 2010–10 <u>2010–11</u> /2013	TCCON	2.21 ± 0.06 <u>2.25 ± 0.06</u>	7.4 ± 0.1 <u>7.3 ± 0.2</u>	Mar 19 <u>21</u>	Aug 23 <u>24</u>	
		GOSAT	2.01 ± 0.08 <u>2.05 ± 0.09</u>	6.4 ± 0.2 <u>6.5 ± 0.2</u>	Apr 1 <u>Mar 27</u>	Aug 27	10
Bremen	4/2009–4/2013	TCCON	2.02 ± 0.09 <u>2.21 ± 0.06</u>	7.9 ± 0.3 <u>7.7 ± 0.3</u>	Mar 20 <u>8</u>	Sep 5 <u>3</u>	
		GOSAT	1.91 ± 0.14 <u>1.88 ± 0.09</u>	6.5 ± 0.4 <u>6.6 ± 0.3</u>	Apr 8 <u>10</u>	Aug 22 <u>24</u>	10
Bialystok	4/2009–10/2013	TCCON	2.18 ± 0.03	8.1 ± 0.1	Mar 16	Aug 18	
		GOSAT	1.99 ± 0.06	7.5 ± 0.2	Apr 5	Aug 17	10
Sodankylä	5/2009–10/2013	TCCON	2.15 ± 0.04	8.7 ± 0.3	Apr 16	Aug 15	
		GOSAT	2.05 ± 0.09	9.5 ± 0.5	Apr 24	Aug 17	10