

# Influence of the ambient humidity on the concentration of natural deposition ice nuclei

María Laura López and Eldo E. Ávila

FaMAF, Universidad Nacional de Córdoba, IFEG-CONICET, Córdoba, Argentina.

*Correspondence to:* E. E. Ávila  
(avila@famaf.unc.edu.ar)

## Abstract.

This study reports measurements of deposition ice nuclei (IN) concentration at ground level during the period July-December 2014 in Córdoba, Argentina. The measurements were carried out at temperature of  $-25^{\circ}\text{C}$  and a 15 % supersaturation over ice. They were performed on days with  
5 different thermodynamic conditions, including rainy days. The effect of the relative humidity at ground level ( $\text{RH}_{amb}$ ) on the IN concentration was analyzed. The number of IN activated varied from  $1\text{ L}^{-1}$  at  $\text{RH}_{amb}$  of 25% to  $30\text{ L}^{-1}$  at  $\text{RH}_{amb}$  of 90%. In general, a linear trend between the IN concentration and the  $\text{RH}_{amb}$  was found, suggesting that this variable must be related to the effectiveness of the aerosols acting as IN. From the backward trajectories analysis, it was found that the  
10 link between IN concentration and  $\text{RH}_{amb}$  is independent of the origin of the air masses. The role of biological-IN and nucleation occurring in pores and cavities were discussed as a possible mechanism to explain the increase on the IN concentration during high ambient relative humidity events. This work provides valuable measurements of IN concentration from the Southern Hemisphere where the data of deposition ice nuclei are sparse.

## 15 1 Introduction

Natural ice nuclei (IN) are considered important in atmospheric processes since they induce freezing in clouds, thus initiating an efficient mechanism for cloud particles to reach a precipitating size. Consequently, many studies have been undertaken to determine their absolute concentrations, their origin and their chemical composition (DeMott et al. (2011); Hoose and Möhler (2012)).

20 Homogeneous freezing of supercooled droplets occurs at temperatures near  $-40^{\circ}\text{C}$ , but at temperatures warmer than approximately  $-36^{\circ}\text{C}$  heterogeneous nucleation occurs (e.g. Cantrell and

Heymtsfield (2005)), which involves four different modes: condensation freezing, immersion freezing, contact freezing and deposition nucleation (Vali, 1985). Cloud condensation nuclei (CCN) can serve as IN at temperatures below 0°C during the condensation freezing mode. During the immersion-freezing mode, a particle suspended in the droplet initiates freezing. In the contact-freezing mode, ice is formed by the collision of cloud droplets with interstitial aerosols within the cloud. When the environment is supersaturated with respect to ice and subsaturated with respect to liquid water, ice deposits on the IN directly from the vapor phase. This last mode is known as deposition nucleation. Measurements performed in the regime below water saturation are important in the understanding of the deposition nucleation (DeMott et al., 2011).

Temperature and supersaturation ratio are the two main parameters determining the deposition nucleation. Several reports have provided measurements of IN concentration for different temperature and supersaturation conditions. Studies linking IN concentration to temperature have shown an exponential trend in the dependence of IN on temperature (e.g. Fletcher (1962); Hussain and Saunders (1984); Meyers et al. (1992); López and Ávila (2013)). This same correlation was found when studying IN concentration regarding supersaturation (e.g. Rogers (1982); Al-Naimi and Saunders (1985); Hussain and Saunders (1984); Cooper (1980); López and Ávila (2013)).

Although temperature and supersaturation are recognized as the two main parameters in the activation of the IN, a significant divergence among the results of IN quantification is found in the literature when IN measurements are carried out at around the same temperature and supersaturation conditions (DeMott et al. (2011); Hoose and Möhler (2012)). In general, the results show that the ice-nuclei content of the atmosphere varies considerably from day to day and from place to place. For instance, important anomalies of atmospheric ice nuclei were observed under thunderclouds (Isono and Tanaka (1966); Isaac and Douglas (1973)). Increments at ground level ice nucleus concentrations by a factor of 10 or higher have been measured during precipitation from convective storm precipitation. These anomalies were associated with downdrafts in and under the thunderclouds. Prenni et al. (2013) also studied the impact of precipitation at a forested site on the concentration and composition of IN. They showed that at ground level IN-concentrations increase during rain events.

Thus, our knowledge of ice formation through the nucleation modes is still scarce and limited. As a consequence, the ice nucleation processes are currently very poorly represented in global climate models (e.g. Hoose et al. (2010)). Furthermore, the study of ice cloud formation over a wide range of atmospherically relevant temperatures and humidity values is challenging (e.g. DeMott et al. (2011); Murray et al. (2012)). Particularly, studies on anomalies in atmospheric ice nucleus concentration are important, since they will give us clues for solving problems on the origin and the nature of atmospheric ice nuclei. The major goal of this work is the study of the factors that regulate the effectiveness of aerosols on deposition ice nucleation. Regarding the possible effect of precipitation on the IN-concentration, this study reports measurements of IN-concentration at ground level during

the period July-December 2014 in Córdoba, Argentina. The experiments were performed on days  
60 with different thermodynamic conditions, including precipitation days, and the effect of the relative  
humidity at ground level on the IN concentration was analyzed. IN-concentrations in deposition  
nucleation mode were determined at a temperature of approximately  $-25^{\circ}\text{C}$  and supersaturation of  
approximately 15 % over ice. The ambient relative humidity at ground level was measured during  
all the experiments, in order to link this variable to the IN concentration.

## 65 2 Experimental

Measurements were carried out in Córdoba, a mediterranean city of Argentina, located in the center  
of the country, in a semiarid region (latitude  $-31.4^{\circ}\text{C}$ ; longitude  $-64.18^{\circ}\text{C}$ ; 470 m.a.s.l.). Córdoba  
is the second largest city in the country, with approximately 1.3 million inhabitants. The climate  
is sub-humid and the prevailing wind direction is NE. The annual average precipitation is close to  
70 700 mm. However, the area is affected by severe and persistent dry periods that occur cyclically.  
Snowfalls are infrequent; thus, precipitations consist mostly of rain. Rainfall is highly seasonal,  
concentrated mainly in summer (December-February) having accumulated values higher than 400  
mm, while the dry winter season (June-August) has a cumulative precipitation depth close to 20 mm.

In a previous work we have quantified the IN-concentration (López and Ávila, 2013) in depo-  
75 sition nucleation mode at ground level in Córdoba, for temperatures between  $-15^{\circ}\text{C}$  and  $-30^{\circ}\text{C}$ ,  
and ice supersaturation ranging between 2 and 20 %. In the present work the experiments were  
carried out following the same procedure as described by López and Ávila (2013) and a total of  
63 measurements of IN concentration were carried out during 45 days, since 28 July 2014 until 10  
December 2014 (winter and spring seasons). This procedure involves a cloud chamber of 46 L in  
80 volume placed inside a cold room. Controlled volumes of humid air were injected from the exterior  
(at ambient temperature) in order to increase the relative humidity in the cloud chamber. Since the  
same volume of air is removed while this air is injected, the number of aerosols and the pressure  
inside the chamber are kept around constant. The increasing of the relative humidity over ice inside  
the cloud chamber ( $\text{RH}_i$ ) is explained according the adiabatic isobaric mixing of two masses of air  
85 with different relative humidity and temperature (Curry and Webster, 1999). The supersaturation is  
achieved starting with a  $\text{RH}_i$  slightly lower than 100% in the cloud chamber and introducing con-  
trolled air injections from outside. Since the increase of  $\text{RH}_i$  for each air injection is known, it is  
possible to estimate the increase produced by a known number of injections, even once the vapor is  
supersaturated over ice. Once the supersaturation is reached, the IN are activated and then grown at  
90 the expense of vapor until ice crystals fall down onto a dish placed on the floor of the cold chamber.  
This dish contains a supercooled aqueous solution of cane sugar. Once in the sugar solution, the  
ice crystals grow and reach a size large enough to be counted with the naked eye, as can be seen in  
Figure 1. For a more detailed description of the procedure, see López and Ávila (2013).

In all the measurements carried out in the present work the conditions of temperature (T) and supersaturation over ice ( $S_i$ ) were settled at around  $-25^\circ\text{C}$  and 15%, respectively. These values were chosen because, under these conditions, the number of activated IN are significant and can be easily determined (López and Ávila, 2013). Furthermore, these T and  $S_i$  values are experimentally quick to reach, which facilitate carrying out the experiments. In order to determine  $\text{RH}_i$  and T in the cloud chamber the Testo 435-4 multifunction measuring instrument was used. This instrument measures relative humidity in the temperature range  $[-50, 150]^\circ\text{C}$ . When  $\text{RH}_i$  is measured over ice at temperatures below  $0^\circ\text{C}$ , the resolution is 0.1%. The experimental uncertainty associated with the determination of T and  $S_i$  was  $\pm 2^\circ\text{C}$  and  $\pm 2\%$ , respectively. These errors arise from variations in the experimental conditions involved in each measurement, such as difference in the temperature between the beginning and the end of the air injections, variability in the  $\text{RH}_i$  increase between the injections, number of injections until reaching the particular  $S_i$ , and accuracy of the humidity sensor.

Thus, considering that T and  $S_i$  were the same for all the experiments (apart from the inherent errors), the IN concentration was studied as a function of the ambient relative humidity at ground level ( $\text{RH}_{amb}$ ); this is the relative humidity of the air mass injected from the exterior.  $\text{RH}_{amb}$  was measured during all the experiments by using the instrument EE31 Series Model D (E+E Elektronik), which had a remote sensing probe for relative humidity measurements in the temperature range  $[-40, 180]^\circ\text{C}$ . The uncertainty in the determination of  $\text{RH}_{amb}$  was  $\pm 1\%$ .

### 3 Results and discussions

The experiments were all performed for a supersaturated atmosphere with respect to ice and subsaturated with respect to liquid water; thus, only the deposition IN were quantified. The IN concentration was determined at a temperature of approximately  $-25^\circ\text{C}$  and supersaturation of approximately 15% over ice. Figure 2 shows the temporal variations of the IN concentration values during the period under study. The grey points in the graph correspond to measurements performed during a rain event or immediately after rain had stopped. The error associated in the estimation of IN concentration did not exceed 10%. There are some time periods without experimental data mainly because the cold chamber was unable to achieve the expected temperature. The period of study is too short to perform a reliable statistical test showing a seasonal behavior of IN concentration; nevertheless, the results do not seem to indicate that there is any particular trend during this period.

In order to study the effect of the  $\text{RH}_{amb}$  in the IN concentration, Figure 3 displays the IN concentration as a function of the  $\text{RH}_{amb}$ . Again, the grey points indicate measurements performed on rainy days. IN concentration values can be seen to increase with  $\text{RH}_{amb}$ . The best linear fitting of the data yield values of 0.39 and  $-9.2$  for the slope and ordinate, respectively, being the square of the correlation coefficient ( $R^2$ ) equal to 0.71. The linear trend between the IN concentration and the  $\text{RH}_{amb}$  is remarkable and the dispersion of the experimental points is likely related to the natural

variability in the IN concentration and to the inherent errors associated with the determination of T,  
130  $S_i$ ,  $RH_{amb}$ , and IN-number, which are present in each experiment.

The number of IN activated varies from  $\sim 1 \text{ L}^{-1}$  at  $RH_{amb} \sim 25\%$  to  $\sim 30 \text{ L}^{-1}$  at  $RH_{amb} \sim 90\%$ . This variation is consistent with the results reported in López and Ávila (2013) on the same site, who found that IN concentration values varied between 2 and  $35 \text{ L}^{-1}$  at temperature of  $\sim -25^\circ\text{C}$  and  $\sim 15\%$  of supersaturation over ice. Thus, the variability in the IN concentration in the previous work  
135 could be related to the variability in  $RH_{amb}$  at the time the measurements were carried out (this variable was not recorded in such experiments). Current results are also in agreement with those reported by Prenni et al. (2013), who showed that, at ground level, the IN concentration rises by a factor of up to 40 during rain. Actually, here we observe that not only rainy days are required for increasing IN concentrations but also high relative humidity is needed, as shown in Figure 3 during  
140 rainfall events with low humidity.

A variety of insoluble particles such as volcanic ash, mineral dust, soot, metallic particles, or primary biological particles have been suggested as IN. However, the requirements needed for a surface to be an efficient IN are not completely understood (Pruppacher and Klett (1997); DeMott (2010); Hoose et al. (2010)). Prenni et al. (2013), Hader et al. (2014) and Bigg et al. (2015) presented evi-  
145 dence that the rising of IN concentration during rain events can be related to the increase of biological particle concentration. It has been shown that biological particles can be released during precipitation through mechanical mechanisms. For instance, winds produced by storms can release biological particles from leaf surfaces and soil. The release of these biological particles include biological IN (bacteria, fungal spores, and pollen) which can impact the IN population. This assumption was  
150 supported by different studies reporting both: the release of biological particles during rain events and the role of these particles as sources of IN (Huffman et al. (2012); Huffman et al. (2013)). Furthermore, Wright et al. (2014) suggested that high relative humidity may also trigger the release of IN at ground level. They provided observational evidence that a fraction of the biological particles released during high RH can serve as immersion mode IN while droplets cool during ascent.

155 It is important to remark that unlike the present study, where the deposition nucleation mode was examined, the previous studies examined condensation and immersion freezing modes. We have no evidence, so far, that the increase of IN concentration in the present work can be analyzed in terms of biological particles. The role of the biological particles acting as deposition ice nuclei need to be studied in more detail.

160 Mineral dust aerosols seem to play a relevant role in deposition nucleation. They have active sites which are able to start ice nucleation. These active sites are linked to surface defects, like cracks, capillaries, pores or cavities which allow water vapor to condense in them under vapor pressures below saturation (Fukuta (1966); Fletcher (1969)). Water vapor condenses in a capillary of a wettable particle under a pressure below water saturation ( $p$ ), according to Kelvin equation

$$165 \quad RT r \ln \left( \frac{p}{p_b} \right) = -2\gamma \nu_l \quad (1)$$

$R$  is the gas constant,  $T$  the absolute temperature,  $r$  the radius of the curved water surface,  $p_b$  the water vapor pressure over a flat water surface,  $\gamma$  the surface tension of water and  $\nu_l$  the molar volume of liquid water. Recently, Marcolli (2014) hypothesized that ice nucleation below water saturation occurs predominantly in water confined between closely spaced surfaces, such as cavities and pores that may form in aerosol particles. In this sense, it is plausible to expect that as the  $RH_{amb}$  increases, more water molecules from the surrounding environment can be adsorbed in cavities and pores. As a result, the condensation and the subsequent freezing can occur in the cavity when the relative humidity is lower than 100% outside. Considering this fact, a significant enhancement in ice nucleation efficiency should be expected at higher ambient relative humidity, which is consistent with the results reported here. Even though further work is needed to test and corroborate this hypothesis, the results presented in this work may support the idea that the deposition nucleation can be a pore condensation and freezing which occur in the defects on the surface of some aerosol particles as proposed by Marcolli (2014).

In order to relate the origin of the air masses to the behavior of IN measurements regarding RH, backward trajectories were calculated for all days when IN concentration measurements were carried out. Back trajectories were calculated by the NOAA HYSPLIT model (Hybrid Single-Particle Lagrangian Integrated Trajectory model) (Draxler and Rolph, 2003). Air mass back trajectories were calculated at 1000 m above mean sea level (AMSL), 48 h before the arrival time to the sampling site. NOAA trajectories were calculated every day of IN concentration measurements. The air mass back trajectories were subdivided into four groups, regarding the quadrants from the Compass Rose. As an example, Figure 4 shows of air mass back trajectories representative of the four studied sectors, which will be referred as North-West quadrant (example: Figure 4a), North-East quadrant (example: Figure 4b), South-West quadrant (example: Figure 4c) and South-East quadrant (example: Figure 4d). The lower sample number was obtained coming from NW quadrant (7.5%), while the highest numbers of IN measurements were obtained coming from NE quadrant (42.5%). Trajectories coming from SE and SW quadrants were 22.5% and 27.5%, respectively. Figure 5 shows the IN concentration measurements grouped following the previous classification of the origin of air masses. Observe that the number of IN measurements from NW quadrant is too scarce to make an analysis. However, the analysis from the three remaining quadrants show the increase of IN concentration with the increase in  $RH_{amb}$ , in the same way that it was observed in Figure 3. Results show that, from the studied period, variations in  $RH_{amb}$  are not explained because of the origin of the air masses. This is to say that, the relation between IN concentration and the ambient relative humidity is independent from the origin of the air masses. This fact discards the possibility that the increase in IN concentration is due to changes in the sources of the aerosols acting as ice nuclei. Thus, HYSPLIT results seem to be consistent with the previous assumption regarding to the importance of the ambient relative humidity on the ability of aerosols to act as IN.

#### 4 Summary and concluding remarks

Studies of heterogeneous ice nucleation with natural aerosols under a broad range of thermodynamic conditions are important, because they can provide clues for solving problems regarding the origin and the nature of atmospheric IN. This study reports measurements of deposition IN concentration at ground level during the period July-December 2014 in Córdoba, Argentina. The experiments were carried out at temperature  $(-25 \pm 2)^\circ\text{C}$  and supersaturation  $(15 \pm 2)\%$  over ice. The effect of the relative humidity at ground level on the IN concentration was analyzed and a linear trend between the IN concentration and the  $\text{RH}_{amb}$  was found, suggesting that this variable must be related to the ability of the aerosols acting as IN. This assumption seems to be corroborated by the analysis of the backward trajectories of the air masses, which shows that the relation between IN concentration and the ambient relative humidity is independent from the origin of the air masses. Therefore, the variability observed on IN concentration cannot be explained regarding the changes in the sources of the aerosols acting as ice nuclei.

The role of biological-IN and nucleation occurring in pores and cavities (as suggested by Marcolli (2014)) was discussed as a possible mechanism to explain the increase on the IN concentration during high ambient relative humidity events. More ice nucleation studies should be carried out to address the differentiation between these mechanisms.

This work contributes to the study of the factors affecting the IN-concentration. Laboratory studies of heterogeneous ice nucleation with natural aerosols are important for the possibility to transfer the results into parameterizations for numerical models of clouds and climate. Furthermore, this work provides valuable measurements of IN concentration from the Southern Hemisphere where the data of deposition ice nuclei are sparse.


*Acknowledgements.* We thank Secretaría de Ciencia y Tecnología de la Universidad Nacional de Córdoba (UNC), Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), and Agencia Nacional de Promoción Científica (FONCYT) for the support. We thank José Barcelona for the technical assistance.

## References

- Al-Naimi, R. and Saunders, C.P.R.: Measurements of natural deposition and condensation-freezing ice nuclei with a continuous flow chamber, *Atmos. Environ.*, 19, 1871-1882, 1985.
- 230 Bigg, E.K., Soubeyrand, S. and Morris, C.E.: Persistent after-effects of heavy rain on concentrations of ice nuclei and rainfall suggest a biological cause, *Atmos. Chem. Phys.*, 15, 2313-2326, 2015.
- Cantrell, W. and Heymsfield, A.: Production of ice in tropospheric clouds: A review, *Bull. Amer. Meteor. Soc.*, 86, 795-807, 2005.
- Cooper, W.A.: A method of detecting contact ice nuclei using filter samples. 8th Intl. Conf. on Cloud Physics, 235 Clermont-Ferrand, France, 665-668, 1980.
- Curry, J.A., and Webster, P.J.: *Thermodynamics of Atmospheres and Oceans*, Academic Press, 1999.
- DeMott, P.J.: New Directions: Need for defining the numbers and sources of biological aerosols acting as ice nuclei, *Atmospheric Environment* 44, 1944-1945, 2010.
- DeMott, P.J., Möhler, O., Stetzer, O., Vali, G., Levin, Z., Petters, M.D., Murakami, M., Leisner, T., Bundke, 240 U., Klein, H., Kanji, Z.A., Cotton, R., Jones, H., Benz, S., Brinkmann, M., Rzesanke, D., Saathoff, H., Nicolet, M., Saito, A., Nillius, B., Bingemer, H., Abbatt, J., Ardon, K., Ganor, E., Georgakopoulos, D.G. and Saunders, C.: Resurgence in ice nuclei measurement research, *Bull. Amer. Meteor. Soc.*, 92, 1623-1635, 2011.
- Draxler, R.R., and Rolph, G.D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) 245 Model Access Via NOAA ARL READY Website. NOAA Air Resources Laboratory, Silver Spring, MD <http://www.arl.noaa.gov/ready/hysplit4.html>, 2003.
- Fletcher, N.H.: *The Physics of Rain Clouds*, Cambridge Univ. Press., Cambridge, UK, 1962.
- Fletcher, N. H.: Active sites and ice crystal nucleation, *J. Atmos. Sci.*, 26, 12661271, 1969.
- Fukuta, N.: Activation of atmospheric particles as ice nuclei in cold and dry air, *J. Atmos. Sci.*, 23, 741-750, 250 1966.
- Hader, J.D., Wright, T.P. and Petters, M.D.: Contribution of pollen to atmospheric ice nuclei concentrations, *Atmos. Chem. Phys.*, 14, 5433-5449, 2014.
- Hoose, C., Kristjansson, J.E. and Burrows, S.M.: How important is biological ice nucleation in clouds on a global scale?, *Environ. Res. Lett.*, 5(2), 7, 2010.
- 255 Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys. Discuss.*, 12, 9817-9854, 2012.
- Huffman, J.A., Sinha, B., Garland, R.M., Snee-Pollmann, A., Gunthe, S.S., Artaxo, P., Martin, S.T., Andreae, M.O. and Pöschl, U.: Size distributions and temporal variations of biological aerosol particles in the Amazon rainforest characterized by microscopy and real-time UV-APS fluorescence techniques during AMAZE-08, 260 *Atmos. Chem. Phys.*, 12, 11997-12019, 2012.
- Huffman, J.A., Prenni, A.J., DeMott, P.J., Pöhlker, C., Mason, R.H., Robinson, N.H., Fröhlich-Nowoisky, J., Tobo, Y., Després, V.R., Garcia, E., Gochis, D.G., Harris, E., Müller-Germann, I., Ruzene, C., Schmer, B., Sinha, B., Day, D.A., Andreae, M.O., Jimenez, J.L., Gallagher, M., Kreidenweis, S.M., Bertram, A.K. and Pöschl, U.: High concentrations of biological aerosol particles and ice nuclei during and after rain, *Atmos. 265 Chem. Phys.*, 13, 6151-6164, 2013.
- Hussain, K. and Saunders, C.P.R.: Ice nucleus measurement with a continuous flow chamber, *Q. J. R. Meteorol.*



- Soc., 110, 75-84, 1984.
- Isaac, G.A. and Douglas, R.H.: Ice nucleus concentrations at  $-20^{\circ}\text{C}$  during convective storms, *J. Appl. Meteorol.*, 12, 1183-1190, 1973.
- 270 Isono, K. and Tanaka, T.: Sudden increase of ice nucleus concentration associated with a thunderstorm, *J. Meteorol. Soc. Jpn.*, 44, 255-259, 1966.
- López, M.L. and Ávila, E.E.: Measurements of natural deposition ice nuclei in Córdoba, Argentina, *Atmos. Chem. Phys.*, 13, 3111-3119, 2013.
- Marcolli, C.: Deposition nucleation viewed as homogeneous or immersion freezing in pores and cavities, *Atmos. Chem. Phys.*, 14, 2071-2104, 2014.
- 275 Meyers, M.P., DeMott, P.J. and Cotton, W.R.: New primary ice-nucleation parameterizations in an explicit cloud model, *J. Appl. Meteorol.*, 31, 708-721, 1992.
- Möhler, O., DeMott, P.J., Vali, G. and Levin, Z.: Microbiology and atmospheric processes: the role of biological particles in cloud physics, *Biogeosciences*, 4, 1059-1071, 2007.
- 280 Möhler, O., Georgakopoulos, D.G., Morris, C.E., Benz, S., Ebert, V., Hunsmann, S., Saathoff, H., Schnaiter, M. and Wagner, R.: Heterogeneous ice nucleation activity of bacteria: new laboratory experiments at simulated cloud conditions, *Biogeosciences*, 5, 1425-1435, 2008.
- Murray, B.J., O'Sullivan, D., Atkinson, J.D. and Webb, M.E.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519-6554, 2012.
- 285 Prenni, A.J., Tobo, Y., Garcia, E., DeMott, P.J., Huffman, J.A., McCluskey, C.S., Kreidenweis, S.M., Prenni, J.E., Pöhlker, C. and Pöschl, U.: The impact of rain on ice nuclei populations at a forested site in Colorado, *Geophys. Res. Lett.*, 40, 2013.
- Pruppacher, H.R. and Klett, J.D.: *Microphysics of Clouds and Precipitation*, Atmospheric and Oceanographic Sciences Library, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- 290 Rogers, D.C.: Field and laboratory studies of ice nucleation in winter orographic clouds, Ph.D. dissertation, Dept. of Atmospheric Science, Univ. of Wyoming, Laramie, 161 pp, 1982.
- Vali, G.: Nucleation Terminology, *J. Aerosol S.*, 16 (6), 575-576, 1985.
- Wright, T.P., Hader, J.D., McMeeking, G.R. and Petters, M.D.: High relative humidity as a trigger for widespread release of ice nuclei, *Aerosol Science and Technology*, 48:iv, 2014.



**Fig. 1.** Images of crystals once they grown on the sugar solution.




Fig. 2. Data of IN concentration measured throughout the period studied.





Fig. 3. Data of IN concentration at different ambient relative humidity.



**Fig. 4.** Location of Argentina and air mass pathways of different quadrants: North-West, North-East, South-West and South-East.




Fig. 5. IN concentration at different ambient relative humidity for the different origins of the air masses.