

Responses to Reviews

Manuscript acpd- 15, C89-91 2015

Authors response to Interactive comment on “Effects of global change during the 21st century on the nitrogen cycle” by D. Fowler et al.

The authors would like to thank the reviewers for the time spent reviewing the manuscript and the numerous helpful comments. Please find below in black the reviewer comments and in BLUE the authors responses.

Anonymous Referee #1

This manuscript is a substantial review of many aspects of the atmospheric N cycle and its coupling to the land surface and biosphere now and through the next century. Overall it is somewhat useful—but in its current form it is trying to be too encyclopedic and as a result fails to present a useful overview and/or to serve as a guide to the primary literature. This is even more of a problem because in the sections I am most familiar with, the review is quite incomplete. There are many authors of this manuscript, but it doesn't appear to have benefitted from sufficient internal review among the authors. It would be better if the authors divided the review into two or three companion papers where a manageable number of experts could make sure the review is complete and comprehensive enough to represent the current state of affairs in our science and where all the authors would feel sufficiently invested that they would read and comment on every aspect of the manuscript.

Response. The reviewer is correct in identifying the main aim of the paper in the opening sentence, and the authors accept that there are areas in which the review is incomplete. The missing areas are addressed in the next comment and reply. Further, we accept that the paper was insufficiently reviewed by the authors, prior to submission. With the substantial changes to the submitted ACPD paper, into the current revised form, this general point has been addressed.

A single paper or three shorter papers: We acknowledge the simplicity of dividing the review into three companion papers, but there are two important reasons for retaining the manuscript as a single paper. First, the main objective is to identify the climate sensitive components of the global nitrogen cycle and estimate the likely changes this century. This requires a single paper. Second, the paper is one of a set of reviews for which the companions are the papers by Fuzzi et al on particulate matter and Monks et al () on ozone. Dividing the papers on nitrogen would not meet the objectives of the wider reviews in this special issue. The paper is not intended to be encyclopaedic, the subject area is very large and a full review would be several times the scale of this manuscript. In revising the introduction we have made it clear that the focus of the paper is on those parts of the nitrogen cycle that are sensitive to changes in climate or land use and for which there is new, published information or on which the authors have been able to commission additional modelling.

Reviewer: in the sections I am most familiar with, the review is quite incomplete.

In focussing the review on processes and fluxes known to be sensitive to the changes this century, some important fluxes were omitted, notably the oxidized organic fluxes of PAN and related compounds and HONO. Additional text has been included in the revised manuscript on PAN section 6.5 pp37 and for HONO 7.00 p39, with additional references to publications on these compounds.

Reviewer: In addition to this larger structural issue, the manuscript is simply long and could in places be dramatically trimmed without loss. If it stays as one document, it is in many places repetitive—clearly written by committee—and would benefit from a single author/editor coalescing key ideas that frame the discussion once and only once.

Response: The paper has been edited to remove unnecessary phrases throughout, as indicated in the track-changes version of the manuscript. In addition to the removal of substantial quantity of text the old fig 8 has been removed as this is not really necessary. The additional text required to satisfy other reviewer comments has compensated for these deletions, so that the manuscript is similar in length to that submitted to ACPD. However, it is a more complete document. While it is long, relative to the subject and the available literature it is short, we have focussed on the main fluxes, their likely responses to changes in climate. Fully a third (30 pp) of the paper is the reference section, and this is a subset of those reviewed. We accept that there was repetition, especially of the scale of uncertainty, which was unhelpful and irritating to read, and has been removed from this revision.

Reviewer: Finally, this is a subject that has been at least partially reviewed and had special issues devoted to it quite recently. It would help the reader to have those items pointed out and to have some clarification about the intended utility of this review compared to other recent reviews in the early part of the introduction.

Response: The Introduction has been expanded (pages 4 to 6) to make clear the focus of the paper and how it differs from other recent reviews and special issues.

Some examples and specific comments:

Reviewer pg 1755 line 15: Sentences like this one appear throughout the manuscript. As a topic sentence for a paragraph it is useless and it undermines the utility of whatever follows. I recommend careful review of all references in the text to complexity, uncertainty and a challenging future and that the authors insure they are making a specific statement that a reader can interpret with each one. In most cases it would be better to write instead that current understanding provides upper and lower bounds on values for some property and then to give a central value with some discussion of why that value is more likely than the extremes. (see also pg 1757 lines 2-4 and lines 6-8 and page 1791 lines 4-6)

Response: These statements have been deleted or where considered necessary have been followed by estimates of the magnitude of the uncertainty.

93 Section 2.2.3. needs to be edited. There are almost more sentences summarizing or
 94 directing the reader to another point in the text than sentences with content.
 95
 96 Response: Point accepted, the text has been edited to clarify.
 97
 98 Reviewer section 4.1.1 The role of bidirectional exchange in setting the spatial patterns of
 99 emission and deposition downwind from large sources should be more clearly discussed.
 100 section
 101 Response: Point accepted and text on the point has been added, (section 10.2 pp59).
 102 Quantifying the effect is a research or modelling task which has not been reported.
 103
 104 Reviewer: 4.3.1. equilibrium and photostationary state are similar but not identical. There
 105 are probably some more recent global analyses of the N budget than the references
 106 used here.
 107
 108 Response: Point taken, this sentence is not needed and has been deleted.
 109
 110 Reviewer: There are certainly many important papers documenting trends of NO emissions at
 111 a continental scale and documenting revised understanding of the processes
 112 that were at the heart of the global analyses in the references quoted.
 113
 114 Response: The point made in 4.3.1 is that the combination of satellite remote sensing data
 115 with bottom-up inventories has led to substantial improvements in estimates of the global
 116 source strength for NO_x and is up to date.
 117
 118 Reviewer: section 4.3.3. the general statements about climate in this section belong earlier in
 119 the document,
 120
 121 Response: The text has been revised and moved to the introduction (p6).
 122
 123 Reviewer: section 5. My understanding is that the PAN lifetime is set more by dynamical
 124 time constants of convective lofting and transport rates from the free troposphere to the
 125 PBL than temperature itself.
 126
 127 Response: The work by Doherty et al (2013) using a coupled chemistry climate model shows
 128 that PAN lifetime is reduced in a warmer climate, and that decomposition is the main cause
 129 of the change.
 130
 131 Reviewer: section 5.1. I think the recent paper by Romps et al. Science 2014 is a more useful
 132 basis for estimating climate effects than the references here.
 133
 134 Response: The recent paper by Romps et al is now included, and, this paper supports the
 135 conclusions of the authors that the temperature response is close to the lower end of the range
 136 reported. The text has been amended to make this point.
 137
 138
 139 Reviewer: section 6. most of this section focusses on reduced organic nitrogen. If that is the
 140 only subject of interest then the section should state that early on and make clear it is not
 141 a review of oxidized organic N. On the other hand, if oxidized organic N is important,
 142 the section is missing an organizing concept for discussing it and is completely lacking

references to recent papers and reviews on gas phase and aerosol oxidized organic nitrogen.

Response: The authors fully accept the point made here, the focus in the manuscript was on reduced organic nitrogen in this section, this has been clarified. We also accept that more reference to PAN, PPN and MPAN is appropriate, even though the magnitude of the fluxes makes their contribution to the global cycling of reactive nitrogen small. Additional text has been added to address this point.

Reviewer: section 7.1 It isn't clear why this topic is in the global change section and not appearing earlier in a section about the state of our understanding of the present day N cycle, Also, this section is missing most of the recent literature on field and lab studies of NO_x and PAN fluxes.

Response: Section 7.1 is on surface -atmosphere exchange processes, and naturally this includes atmospheric transport. We accept that some of this could be presented earlier, but keeping the surface-atmosphere material together as a section appears equally valid. Additional references to recent field and laboratory NO fluxes have been included in section 7.1.4. A section on PAN has been included in section 6.6 (p 37).

Reviewer: sections 7.2-10 What would make this review useful would be to somehow link key ideas in the earlier sections to some assessment of what is knowable about the future. Sections 7.2,8,9 and 10 don't do that. They would stand entirely on their own without any of the preceding sections, hence my suggestion to split this document into several papers.

Response: The point about whether this should be one or three papers has been considered, and rejected, as above. The overall effects of changes this century on global nitrogen cycling are brought together in section 10, and in summary form in figure 15.

Reviewer #2

This paper is a comprehensive review of the nitrogen cycle, its future changes and policy implications. It is challenging to address some many topics in a single paper, but the authors manage to incorporate substantial amount of information in the manuscript. I am in favor of its publication. However, there are some important issues that need to be addressed as listed below.

Reviewer: Structure. This paper is not well structured and balanced which undermines its importance. The introduction is rather short and doesn't give a clear overview of the whole nitrogen cycle.

190 Response : The Introduction has been revised and extended to make the approach and
 191 structure clear. (Section 1 pp4 to 6). The paper has been extensively edited to remove
 192 unnecessary text and duplication of general comments.

193 Reviewer: Then it immediate follows by sections describing individual processes, in which
 194 different structures have been adopted. For example, in the section of marine nitrogen
 195 fixation, its influencing factors have been intensive discussed (in several sub-sections)
 196 which is not the case in the terrestrial nitrogen fixation section. Overall, I found the
 197 figures are easier to understand than the texts. I agree with Referee #1 that substantial
 198 editorial work is needed to compile and synthesize work from individual contributors.

199

200 Response: The point is accepted, there was insufficient editing of the manuscript to bring
 201 the different styles together and make the review more readable. The differences have been
 202 reduced by editing into a more consistent style with fewer levels within the sections and
 203 editing of the text to ease the transitions between different contributors.

204

205 Reviewer: Policy implications.

206 This paper is a contribution to the special issue “Atmospheric composition change:
 207 science for policy”. If that’s the main purpose, this review can be to shortened to a large
 208 extent. Processes of less importance and larger uncertainties can be summarized into
 209 a single section without detailed discussions.

210

211 Response: The authors have some sympathy with the approach suggested. However, a
 212 review must present the evidence on which to draw conclusions. Both Reviewers detail
 213 missing areas of research on reactive nitrogen compounds and on balance the authors
 214 agree that the review needs to present the evidence and a reasonable cross section of the
 215 very different areas of study within the global nitrogen cycle. The policy section is short
 216 relative to other main sections, as much of this material is recently reviewed in the European
 217 Nitrogen Assessment (Sutton et al 2011) and Our Nutrient World (Sutton et al 2013a).

218

219 (3) Missing processes.

220 With regard to the exchange of molecular and reactive nitrogen between the atmosphere
 221 and terrestrial surfaces, the review should not miss to address recent findings
 222 indicating that cryptogamic covers on ground and plant surfaces may account for as
 223 much as half of the biological nitrogen fixation on land (Elbert et al. 2012; Porada et
 224 al., 2014) and that nitrous acid (HONO) can be reversibly deposited and emitted in
 225 large quantities comparable nitrogen oxide (NO) (Su et al. 2011; Kulmala and Petäjä
 226 2011; Oswald et al. 2013; Maljanen et al. 2013; Donaldson et al. 2014; VandenBoer
 227 et al. 2015).”

228

229 Response: The point is accepted and new text on these processes has been included in
 230 section 7 (p38) on HONO, and on nitrogen fixation in section 2.1.

231

232 Reviewer: Abstract: p1749 line 3, what is the meaning of “input of reactive nitrogen from
 233 human activities”, NOx emission or nitrogen fixation?

234

235 Response: This is clarified in the abstract with the inclusion of “including combustion related
 236 NOx, industrial and agricultural N fixation, “.

237

238 Reviewer: Introduction: p1750 line 13, I would suggest including an introduction to the
 239 nitrogen cycle and relevant processes in the beginning.

240

241 Response: Point accepted. New text added to the introduction.

242

243 Reviewer: p1750 line 14, it is better to first explain what’s reactive nitrogen for policy makers.

244 [Response: Definition added within the introduction.](#)

245 Reviewer p1751 line 8-16, this paragraph should appear before the last one “The negative
246 effects of human N : : :”

247 [Response: Paragraph moved as suggested.](#)

248 [Response: Paragraph moved as suggested.](#)

249 Reviewer: p1753 line 11, “Biological nitrogen fixation (BNF) is currently estimated to provide
250 a global annual input of approximately : : :”, a reference is missing.

251 [Response: Reference added.](#)

252 [Response: Reference added.](#)

253 Reviewer: 1756 line 20, Q10 should be defined before use.

254 [Response: Definition added.](#)

255 Reviewer: 1757 line 20, why not adopt the same structure for terrestrial N fixation

256 [Response: The two sections have been edited into a similar format.](#)

257 [Response: The two sections have been edited into a similar format.](#)

258 Reviewer: p1761 line 19, “: : : by effects which decrease N₂ fixation.”, such as?

259 [Response: Sentence deleted.](#)

260 [Response: Sentence deleted.](#)

261 Reviewer: p1761 line 21 “... providing a strong negative feedback to increases in
262 atmospheric CO₂”, please explain what’s the negative feedback

263 [Response: Phrase deleted](#)

264 Reviewer: p1764 line 16 “plus combustion related emissions : : :”, is combustion related
265 emissions considered as fixation or emissions?

266 [Response The NO_x fixed in combustion is fixation while the emission from combustion is the
267 total of the NO_x fixed plus the oxidation of fuel N less any emission control.](#)

268 [Response The NO_x fixed in combustion is fixation while the emission from combustion is the
269 total of the NO_x fixed plus the oxidation of fuel N less any emission control.](#)

270 Reviewer: p1771 line 8, “kg”, g should be a subscript

271 [Response: Agreed, and corrected.](#)

272 Reviewer: p1771 line 8, please use either parentheses or brackets for NH₃ and NH_{sw}, the
273 meaning of “sw” should be explained.

274 [Response: sw is surface water and has been added.](#)

275 Reviewer: p1771 line 13 use either “exp” (Eq. 2) or “e” throughout the manuscript.

276 [Response: done.](#)

277 Reviewer: p1773 line 8 “ the global NH₃ emission” should be “that the global ocean NH₃
278 emission”

279 [Response: Done.](#)

280 p1778 line 23, why discuss denitrification in the nitrification section?

281 [Response: This was to provide context for the following paragraph.](#)

282 [Response: This was to provide context for the following paragraph.](#)

283 [Response: This was to provide context for the following paragraph.](#)

284 [Response: This was to provide context for the following paragraph.](#)

285 [Response: This was to provide context for the following paragraph.](#)

286 [Response: This was to provide context for the following paragraph.](#)

287 [Response: This was to provide context for the following paragraph.](#)

288 [Response: This was to provide context for the following paragraph.](#)

289 [Response: This was to provide context for the following paragraph.](#)

290 [Response: This was to provide context for the following paragraph.](#)

291 [Response: This was to provide context for the following paragraph.](#)

292 [Response: This was to provide context for the following paragraph.](#)

293 [Response: This was to provide context for the following paragraph.](#)

294 [Response: This was to provide context for the following paragraph.](#)

295 [Response: This was to provide context for the following paragraph.](#)

296 [Response: This was to provide context for the following paragraph.](#)

297 [Response: This was to provide context for the following paragraph.](#)

298 Figure 16, some texts were half-covered

299 Response: The figure has been corrected and edited to improve clarity.

300

301

302

303

304

305

306

307

308

309

310

311 **Effects of global change during the 21st century on the nitrogen cycle**

312

313 David Fowler¹, -Claudia E. Steadman^{1,2}, David Stevenson², Mhairi Coyle¹, -Robert M. Rees³
314 , Ute M. Skiba¹, Mark A. Sutton¹, J. Neil Cape¹, Anthony J. Dore¹, Massimo Vieno^{1,2}, -David
315 Simpson⁴, Sönke Zaehle⁵, Benjamin D. Stocker⁶, -Matteo Rinaldi⁷, M. Cristina Facchini⁷,
316 Chris R. Flechard⁸, Eiko Nemitz¹, Marsailidh Twigg¹, Jan Willem Erisman⁹, [Klaus](#)
317 [Butterbach-Bahl](#)¹⁰ and James N. Galloway¹⁰

318 1 Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

319 2 School of GeoSciences, University of Edinburgh, Edinburgh, UK

320 3 Scotland's Rural College (SRUC), West Mains Road, Edinburgh, EH9 3JG, UK

321 4 Norwegian Meteorological Institute, EMEP-MSC-W, P.O. Box 43, 0313 Blindern, Norway

322 5 Biogeochemical Integration Department, Max Planck Institute for Biogeochemistry, Hans-
323 Knöll-Strausse 10, 07745 Jena, Germany

324 6 Imperial College, Department of Life Sciences, Silwood Park, Ascot, SL5 7PY, UK

325 7 ISAC-CNR, via P. Gobetti 101, 40129 Bologna, Italy

326 8 INRA, Agrocampus Ouest, UMR 1069 SAS, 35042 Rennes, France
 327 9 Louis Bolk Institute, Hoofdstraat 24, 3972 LA, Driebergen, The Netherlands
 328 10 [Institute for Meteorology and Climate Research \(IMK-IFU\), Garmisch-Partenkirchen,](#)
 329 [Germany](#)
 330 11 Department of Environmental Sciences, University of Virginia, Charlottesville, VA, 22904-
 331 4123, USA

332

333 Contents

334	1.	INTRODUCTION
335	2.	BIOLOGICAL NITROGEN FIXATION
336	2.1	Terrestrial nitrogen fixation
337		<i>2.1.1 Effects of climate change on terrestrial biological nitrogen fixation</i>
338		
339	2.2	Marine biological nitrogen fixation in the 21st century
340		2.2.1 Factors affecting nitrogen fixation
341		2.2.1.1 Light
342		2.2.1.2 Temperature
343		2.2.1.3 Oxygen
344		2.2.1.4 Salinity
345		2.2.1.5 Trace metals and phosphorus
346		2.2.1.6 Stratification
347		2.2.1.7 Carbon dioxide
348		2.2.2 Present-day and pre-industrial estimates
349		<i>2.2.3 Effects of global change on marine biological nitrogen fixation</i> <i>2.2.3 Future</i>
350		<i>impacts</i>
351	2.3	Global changes in natural BNF 2010 to 2100
352		
353	3.	ANTHROPOGENIC FIXATION OF NITROGEN IN THE 21ST CENTURY
354	4.	EFFECTS OF ENVIRONMENTAL CHANGES ON THE FATE OF N _r IN
355		TERRESTRIAL AND MARINE ECOSYSTEMS
356	4.1	Terrestrial ecosystems: Emissions of NH ₃ from terrestrial ecosystems
357		through the 21st century
358		4.1.1 Effects of changes in climate on terrestrial emissions of NH ₃
359	4.2	Ammonia exchange over the oceans in the 21st century
360		4.2.1 Factors affecting the flux of ammonia between the atmosphere and the
361		ocean
362		4.2.2 Flux estimates
363		4.2.3 Future impacts
364	4.3	Terrestrial emissions of Nitric Oxide and Nitrous Oxide
365		4.3.1 Global sources of NO and N ₂ O in the atmosphere

Commented [MKE1]: See page 15 - there is a new subheading here but level is not clear - should it be 2.3 or 2.2.4? Title is: Global changes in natural BNF 2010 to 2100. Check with David and then edit contents page here and level indication in text on page 15.

366	4.3.1.1 NO _x
367	4.3.1.2 N ₂ O
368	4.3.2 Soil processes responsible for NO and N ₂ O emissions
369	4.3.2.1 Denitrification
370	4.3.2.2 Nitrification
371	4.3.3 Effects of climate change on NO and N ₂ O emissions
372	4.3.3.1 Climate change effects on NO and N ₂ O emissions and the importance
373	of changes in regional hydrology
374	4.3.3.2 Land-use and land management effects on denitrification
375	4.3.3.3 Atmospheric composition change and denitrification
376	5. ATMOSPHERIC PROCESSING – CHEMISTRY
377	5.1 Lightning-climate interactions
378	6. ORGANIC NITROGEN
379	6.1 Atmospheric relevance
380	6.2 Chemical composition
381	6.3 Organic nitrogen sources
382	6.4 Effects of future climate change on ON
383	<u>6.5 Peroxyacetyl nitrate (PAN), peroxypropionyl nitrate (PPN) and</u>
384	<u>peroxymethacryloyl nitrate (MPAN)</u>
385	
386	7. <u>EFFECTS OF GLOBAL CHANGE ON ECOSYSTEM/ATMOSPHERE</u>
387	<u>EXCHANGE OF REACTIVE NITROGEN</u>
388	<u>IMPACTS OF GLOBAL CHANGE ON ECOSYSTEM/ATMOSPHERE REACTIVE</u>
389	<u>NITROGEN EXCHANGE PROCESSES</u>
390	7.1 Impacts on processes regulating surface N _r sink/source strength
391	7.1.1 Vertical atmospheric transport
392	7.1.2 Stomatal exchange
393	7.1.3 Non-stomatal plant surfaces
394	7.1.4 Soil surface exchange
395	7.1.5 Chemical interactions during the exchange process
396	7.2 Regional and global projections for nitrogen deposition
397	8. THE EFFECTS OF CLIMATE AND LAND USE CHANGES ON THE WET
398	REMOVAL OF NITROGEN COMPOUNDS FROM THE ATMOSPHERE
399	9. EFFECTS OF CLIMATE AND LAND USE CHANGES ON C-N RESPONSES
400	IN TERRESTRIAL ECOSYSTEMS
401	10. DISCUSSION AND POLICY IMPLICATIONS OF THE RESPONSES OF THE
402	NITROGEN CYCLE TO GLOBAL CHANGE
403	10.1 Emissions and cycling
404	10.2 Effects of changes in atmospheric composition on long range transport of N _r
405	10.3 Costs
406	10.4 Policies to reduce the impacts of N _r
407	REFERENCES
408	

Formatted: Indent: Left: 1.27 cm, First line: 0 cm

Abstract

The global nitrogen (N) cycle at the beginning of the 21st century has been shown to be strongly influenced by the inputs of reactive nitrogen (N_r) from human activities, including combustion related NO_x , industrial and agricultural N fixation, estimated to be 210 Tg N yr⁻¹ in 2010 which is approximately equal to the sum ~~of biological~~ of biological N fixation in unmanaged terrestrial and marine ecosystems. According to current projection trajectories, changes in climate and land use during the 21st century will increase both biological and anthropogenic fixation, bringing the total to approximately 600 Tg N yr⁻¹ by around 2100. The fraction contributed directly by human activities is unlikely to increase substantially if increases in nitrogen use efficiency in agriculture are achieved and control measures on combustion related emissions implemented.

Some N cycling processes emerge as particularly sensitive to climate change. One of the largest responses to climate in the processing of N_r is the emission to the atmosphere of NH_3 , which is estimated to increase from 65 Tg N yr⁻¹ in 2008 to 93 Tg N yr⁻¹ in 2100 assuming a change in global surface temperature of 5°C ~~even~~ in the absence of increased anthropogenic activity. With changes in emissions in response to increased demand for animal products the combined effect would be to increase NH_3 emissions to 132 Tg N yr⁻¹. Another major change is the effect of climate changes in aerosol composition and specifically the increased sublimation of NH_4NO_3 close to the ground to form HNO_3 and NH_3 in a warmer climate which deposit more rapidly to terrestrial surfaces than aerosols. ~~combined with changes in temperature~~. Inorganic aerosols over the polluted regions especially in Europe and North America were dominated by $(NH_4)_2SO_4$ in the 1970s to 1980s, and large reductions in emissions of SO_2 have removed most of the SO_4^{2-} from the atmosphere in these regions. Inorganic aerosols from anthropogenic emissions are now dominated by NH_4NO_3 , a volatile aerosol which contributes substantially to PM_{10} and human health effects globally as well as eutrophication and climate effects. The volatility of NH_4NO_3 and rapid dry deposition of the vapour phase dissociation products, HNO_3 and NH_3 , is estimated to be reducing the transport distances, deposition footprints and inter-country exchange of N_r in these regions.

There have been important policy initiatives on components of the global N cycle. These have ~~For the most part they have~~ been regional or country-based and have delivered substantial reductions of inputs of N_r to sensitive soils, waters and the atmosphere. To date there date there ~~have been no attempts to develop a global strategy to regulate human inputs to the nitrogen cycle~~. However, considering the magnitude of global N_r use, potential future increases, and the very large leakage of N_r in many forms to soils, waters and the atmosphere, international action is required. Current legislation will not deliver the scale of reductions globally there is a very long way to go before evidence ~~for recovery from the effects of N_r deposition on sensitive ecosystems, or a decline in N_2O emissions to the global atmosphere, are likely to be detected~~. Such changes would require substantial improvements in nitrogen use efficiency across the global economy combined with optimisation of transport and food consumption patterns. This would allow reductions in N_r use, inputs to the atmosphere and deposition to sensitive ecosystems. Such changes would offer substantial economic and environmental co-benefits which could help motivate the necessary actions.

451

452 **1 INTRODUCTION**

453 While nitrogen is abundant, comprising 80% of the atmosphere, its form, as molecular nitrogen
 454 N₂ is largely unavailable to biota. Specialized organisms are able to fix nitrogen and transform
 455 it into compounds available for synthesis amino acids and other metabolic products. Before
 456 human activities contributed to nitrogen fixation, the global nitrogen cycling in soils,
 457 vegetation the atmosphere and oceans relied entirely on microbial biological fixation, plus a
 458 small contribution from lightning. Human activities began to substantially contribute to the
 459 global nitrogen cycle at the beginning of the 20th century through combustion, which creates
 460 fixed nitrogen as NO_x, industrial NH₃ production (by the Haber-Bosch process) and by growing
 461 nitrogen fixing crops. The global nitrogen (N) cycle has been perturbed extensively by human
 462 activity over the last 100 years with approximately two thirds of the annual flux of reactive
 463 nitrogen (N_r, which includes all compounds of nitrogen following fixation of molecular
 464 nitrogen N₂) ~~entering~~ entering the atmosphere at the beginning of the 21st century being
 465 anthropogenic in origin (Galloway et al., 2004, Fowler et al., 2013). This has led to widespread
 466 negative consequences through directly contributing to radiative forcing of climate, reductions
 467 in biodiversity at regional scales in terrestrial ecosystems and in damage to human health
 468 through aerosols and ozone production (Erisman et al., 2013, Sutton et al., 2011). Human
 469 modification of the N cycle also has substantial benefits, through sustaining the food supply to
 470 a global human population of seven billion and stimulating global CO₂ sequestration by
 471 terrestrial and marine ecosystems (Zaehle, 2013; Sutton et al., 2013b).

472 The damage by N_r to ecosystems, human health and climate result from leakage of N
 473 compounds from its use in agriculture, industry and transport (Erisman et al., 2013). A
 474 particular feature of the N cycle is the combination of the large number of forms, both oxidised
 475 and reduced, in which N_r exists, with biological and chemical transformations allowing the
 476 same emitted molecule of N_r to take part in a series of effects, both negative and positive, before
 477 being transformed back to molecular nitrogen and returned to the atmospheric reservoir. This
 478 has been termed the nitrogen cascade (Galloway et al., 2003) and substantially complicates an
 479 assessment of the pathways and effects of N_r in the environment.

480

481 The negative effects of human N fixation, are substantial and have been estimated to be 70-320
 482 billion Euros annually for Europe (Sutton et al., 2011; Brink et al., 2011). A comprehensive
 483 global assessment of the costs of human use of fixed N has yet to be made. However, the scale
 484 of European use, at ~17 Tg N annually, represents only 8% of the total anthropogenic N_r fixed
 485 annually (210 Tg N yr⁻¹). As the local hot spots of N_r use in North America and especially in
 486 East and South Asia, show values of emission and deposition similar to or larger than in Europe,
 487 it is likely that the global costs of human use of N_r are therefore an order of magnitude greater
 488 than those for Europe. This would be consistent with a preliminary estimate of global damage
 489 costs associated with N pollution of 800 (200-2000) billion US dollars per year (Sutton et al.,
 490 2013b).

Formatted: Superscript

The damage by N_r to ecosystems, human health and climate result from leakage of N compounds from its use in agriculture, industry and transport (Erisman et al., 2013). A particular feature of the N cycle is the combination of the large number of forms, both oxidised and reduced, in which N_r exists, with biological and chemical transformations allowing the same emitted molecule of N_r to take part in a series of effects, both negative and positive, before being transformed back to molecular nitrogen and returned to the atmospheric reservoir. This has been termed the nitrogen cascade (Galloway et al., 2003) and substantially complicates an assessment of the pathways and effects of N_r in the environment.

Recent analyses of the global N cycle have focussed on the magnitude of current fluxes (Fowler et al., 2013), effects of human activity on the processes and effects on human health, climate and ecosystems, especially in the regional assessments in Europe (Sutton et al. 2011, 2013b) European Nitrogen Assessment (ENA) and in the United States of America US assessments (Sutton et al. 2011, 2013b; (Davidson et al., 2012; Fowler et al., 2013). The extensive conversions of N_r in the environment mediated by biological and chemical processes are sensitive to environmental conditions and thus are likely to respond to changes in climate over coming decades. Thus the current global N cycle is likely to change, regardless of future changes in human activities or human intervention to regulate losses to the environment.

The likely responses of the exchanges of N_r between and within the major global reservoirs in coming decades to changes in climate and land use have not been considered to date, and are the extensively and are the focus of this review. The paper focuses on the effects of global changes expected this century on the processes and fluxes within the global N cycle.

Recent assessments of state of scientific understanding include 14 papers published by The Royal Society on the global nitrogen cycle (Phil. Trans. R.Soc.B368 2013). These relatively short papers focus on components of the global nitrogen cycle in the atmosphere, terrestrial marine and Polar Regions, and include a global overview (Fowler et al 2013). The coverage is not encyclopaedic and the main focus is on terrestrial ecosystems and the atmosphere. The effects of climate changes in the 21st century are not treated in detail within these papers.

The potential impacts of consequences of changes in climate and land use on the global nitrogen cycle are considerable in both the range and magnitude of effects. The processes which regulate transfers between the atmosphere and terrestrial and marine reservoirs are generally sensitive to aspects of climate that are expected to change, including temperature, absolute humidity and precipitation (Sutton et al., 2013b). Many of the major transfers are mediated by biological processes, especially microbiological transformations, which are very sensitive to changes in climate, especially in temperature or humidity. The exchange fluxes of N_r compounds at the Earth's surface, including emission and deposition, are regulated by a combination of atmospheric transfer and surface reactions and biological regulation through stomatal exchange and soil microbiology. These processes therefore include physical, chemical and biological interactions combining to regulate the overall process. Most of the components of the pathway are sensitive to climate, and while the response of some components to specific changes in the environment may be predicted, the overall process relies on measurements to constrain the potential range of effects (Fowler et al., 2009; Monks et al., 2009).

Formatted: Superscript

532 Some of the effects appear straightforward, such as increases in emission fluxes of nitric oxide
533 (NO) from soils and ammonia (NH₃) from vegetation with temperature, but when the full range
534 of expected changes in climate and the number and phase in which the N_r compounds reside
535 are included, the responses become complex and harder to quantify. To consider the whole N
536 cycle and interactions with climate and land use change requires a coupled global climate and
537 ~~global~~ N cycle model, which to date has not been achieved. While parts of the biogeochemistry
538 have been incorporated in global climate models, especially those linked to ozone chemistry
539 and emissions of oxidised N (Stevenson et al., 2006), many of the interactions of reduced
540 nitrogen compounds have yet to be included (Sutton et al., 2013b). In the absence of ~~a~~the
541 global modelling needed to quantify the interactions there have been a number of model
542 investigations at regional scales. There have also been modelling studies of interactions
543 between the carbon (~~C~~) and N cycles which provide useful insight to biogeochemical
544 interactions (Zaehle et al., 2013).

545 This paper explores current knowledge of the sensitivity of biological nitrogen fixation,
546 emissions, atmospheric processing and removal of N_r compounds to changes in climate and
547 land use, defined here as:

548 (a) Climate change: This refers to the change of the primarily environmental drivers
549 temperature and rainfall (amount, frequency, seasonal distribution), both affecting
550 soil environmental conditions but also site and landscape hydrology, vegetation
551 cover and substrate supply. Land use is also influenced, since farmers will adapt
552 land use and land management as climate changes (Kicklighter et al., 2014).

553 (b) Land use change: This refers to changes in vegetation cover, land use and
554 management resulting in changes in substrate supply to the soil microbial
555 community, but also triggers changes in soil and catchment hydrology. Policy and
556 economic drivers also influence the uptake of measures aimed at promoting
557 mitigation in the agricultural sector (MacLeod et al., 2010).

558 (c) Atmospheric composition change: This is mainly due to rising CO₂ concentrations,
559 resulting in reductions in plant transpiration and increasing levels of soil moisture
560 (e.g. Long et al., 2004), but also to changes in regional O₃ concentrations - affecting
561 plant performance and, thus, e.g. plant litter production or transpiration – and / or
562 atmospheric deposition of reactive nitrogen (Sutton et al., 2011), which is not only
563 an additional N_r source for soil microbial processes but also drives forest C
564 sequestration and changes in soil C and N stocks (De Vries et al., 2014).

565
566 ~~from the available literature.~~ The focus is on responses of the flow of nitrogen through
567 terrestrial and marine ecosystems and the atmosphere to changes this century and includes new
568 modelling and analysis as well as the published literature. ~~Discussion of the likely~~
569 ~~consequences for the overall functioning of the global N cycle is provided to the extent that~~
570 ~~this is currently feasible.~~ Consequences for human health, ecosystems and food production of
571 these likely responses are briefly considered.

Formatted: Indent: Left: 1.27 cm, Hanging: 0.63 cm,
Numbered + Level: 1 + Numbering Style: a, b, c, ... + Start at:
1 + Alignment: Left + Aligned at: 1.27 cm + Indent at: 1.9
cm

The structure of the review follows the pathway from fixation of atmospheric nitrogen, by both biological and industrial processes to emission of gaseous N_r compounds ~~into the through~~ atmospheric ~~and processing and~~ removal by dry and wet deposition. Interactions between global nitrogen and carbon cycles are included as they represent key areas of development of Earth system models and are a focus of wider academic interest in the global nitrogen cycle.

Some components of the nitrogen cycle are well supported by recent literature and extensive measurements, as in the case of surface-atmosphere exchange processes, and oxidized nitrogen compounds in the atmosphere, while others are poorly supported by measurements and recent research. This variability in knowledge leads to different approaches in the sections of the paper, concentrating on those components which have been subject to recent publications, or new modelling, specifically developed for this paper.

The review concludes with a brief discussion of the policy implications of climate-nitrogen cycle interactions, as this an important driver of the research agenda and provides context, and has been the subject of several recent publications (Sutton et al., 2011 and Sutton et al., 2013a).

2 BIOLOGICAL NITROGEN FIXATION

Biological nitrogen fixation (BNF) is currently estimated to provide a global annual input of ~~over 273300~~ Tg $N\ yr^{-1}$ to the biosphere (Fowler et al 2013) making it the largest single global input of N_r , although there are significant uncertainties about the magnitude and spatial distribution of fluxes (Fig. 1). ~~If we assume that the global N cycle was in an approximate equilibrium prior to industrialisation, BNF would have been balanced by the reductive processes of denitrification returning molecular nitrogen (N_2) to the atmosphere, with estimates of around 260 Tg $N\ yr^{-1}$ arising from terrestrial and oceanic sources (Galloway et al., 2004).~~ The process of fixation is undertaken by a very limited range of highly specialised microorganisms that share an ability to use the nitrogenase enzyme to split the triple bond present in atmospheric N_2 and combine it with hydrogen to produce a source of N_r . Although the process is highly energy demanding, it is performed at ambient temperature and pressure unlike the industrial Haber-Bosch process that requires the reactants to be combined in the presence of an iron catalyst at between 300-500°C in a reaction vessel at 20 MPa. Two main groups of organisms are responsible; free-living bacteria and algae (which are widespread in fresh water, oceans and uncultivated soils and often form mutualistic associations with a range of plant species) and symbiotic bacteria (mostly belonging to the genus *Rhizobium*) which form symbiotic associations with the roots of plants (mostly belonging to the family *Leguminosae*).

2.1 Terrestrial nitrogen fixation

In terrestrial environments, a wide diversity of both symbiotic and free-living N fixers contribute to BNF in non agricultural soils, but ~~again~~ a lack of measurements results in large

uncertainties in reported values. A meta-analysis of published data compiled from a large number of individual measurements of N fixation carried out in diverse ecosystems reported an average annual global flux of 195 Tg N with a range of 100-290 (Cleveland et al., 1999), although this was later revised downwards to 128 Tg (Galloway et al., 2004). It is thought that tropical environments are particularly important in contributing to terrestrial BNF, although these areas are associated with the least frequent measurements. Recent measurements of BNF by methanotrophs in pristine peatland at high latitude by Vile et al. (2014) suggest appreciable fixation inputs in these environments which have not been included in global estimates to date. Using net carbon uptake methods, Porada et al. (2014) also suggest significant contributions to global nitrogen fixation from lichens and bryophytes.

Using an N balance approach in which the global N cycle is assumed to be in steady state, BNF can be estimated as the difference between inputs and outputs of N within a global context. This approach has suggested that preindustrial terrestrial BNF in natural ecosystems was only 44 Tg N yr⁻¹ (Vitousek et al., 2013), however, such a small low-value questions whether current rates of natural BNF reported by Cleveland and others from up-scaling may have been be overestimated. The recent estimate of BNF in natural terrestrial ecosystems of 58 Tg N annually by Vitousek et al. (2013) is substantially smaller than other recent syntheses of the literature, which are generally in excess of 100 Tg N annually, but subject to large uncertainty. The most recent measurements of BNF in peatlands, which, although representing 3% of the world's land surface, contain approximately 25% of the world's soil carbon, suggest an additional source in these regions in the range of 4.8 to 62.3 kg N ha⁻¹ annually and a mean value of 25.8 kg N ha⁻¹ annually (Vile et al., 2014). Net carbon uptake by lichens and bryophytes has also been used to estimate nitrogen requirement and indirectly nitrogen fixation by Porada et al. (2014), also suggesting a significant contribution to global N fixation by these plant communities. Given these new measurement-based values for extensive ecosystems, the value for global BNF in natural ecosystems seems unlikely to be smaller than 100 Tg N annually and the values proposed by Galloway et al. (2004) of 128 Tg N yr⁻¹ is used here for 2010.

Biological N fixation provides a large significant input of fixed N to agricultural systems. Prior to the development of synthetic fertilizers at the beginning of the 20th century, most of the N used to produce crops and livestock would have been derived from this source. The current input is estimated to be approximately 60 Tg N yr⁻¹, taken as the central value in the range 50-70 Tg yr⁻¹ from Herridge et al. (2008). This value is divided mainly between the grain legumes (peas and beans) and forage legumes (such as clover and alfalfa) contributing 21 and 19 Tg yr⁻¹ respectively (Herridge et al., 2008). Estimates of BNF by the grain legumes are generally considered to be more reliable than those from forage crops since comprehensive records of the former are maintained by FAO (FAO, 2012). Other minor inputs of N by BNF in agriculture include symbiotic N fixation from tropical savannas used for grazing (14 Tg) free living micro-organisms associated with rice paddies (5 Tg), and sugar cane (0.5 Tg).

The uncertainties associated with global estimates of BNF make predictions of future changes particularly challenging. During the 20th century, there has been a rapid growth in the cultivation of leguminous crops contributing to an increase in associated BNF (Galloway et al.,

Commented [FD2]: This is new...and merits a comment to referees

2004). Future growth of legume crops will be constrained by the land area available to agriculture, and increases in production are most likely to occur when legumes are grown in place of other species. Emissions of nitrous oxide (N₂O) resulting from the growth of legume crops is generally ~~small~~^{low} by comparison with other crops, and the IPCC guidelines on greenhouse gas reporting assumes that the N input resulting from legume production is not associated with any N₂O emissions (IPCC, 2006). For this reason, increases in legume cultivation have been promoted as an opportunity to reduce N₂O emissions from agricultural systems by reducing emission intensity of fixed N inputs to agricultural systems (Luscher et al., 2014). Legumes also continue to provide the main source of N input to low input agricultural systems and organic farming globally.

2.1.1 Effects of climate change on terrestrial ~~b~~Biological ~~n~~Nitrogen ~~f~~Fixation (BNF)

Formatted: Font: Bold, Italic

Biological N fixation associated with non-agricultural ecosystems is susceptible to changes in environmental conditions. A framework for understanding the environmental controls determining the rates of BNF in the biosphere has been proposed in which there is a coupling between N, C and Phosphorus (P) cycling (Houlton et al., 2008; Vitousek et al., 2002). Free-living and symbiotic organisms with the potential to fix N are at a selective advantage in environments with low P availability, however, the high energy costs of BNF require adequate supplies of available fixed C. ~~Thus Net Primary Production (NPP) can be used as a proxy for changes in BNF, that can be useful for modelling purposes (Vitousek et al., 2002).~~ The temperature sensitivity of the nitrogenase enzyme responsible for the fixation process has been clearly demonstrated in a global meta-analysis of fixation rates across dominant terrestrial biomes; the optimal temperature for fixation was found to be 25.2°C with a very sharp decline in rates of fixation below 5°C and above 40°C (Houlton et al., 2008). Projected global increases in temperature are therefore likely to be associated with increases in BNF, providing that sufficient water is available to maintain NPP. However, other environmental changes may counteract increases resulting from climate change. The process of BNF is often down regulated by the presence of fixed N. Agricultural experiments have consistently shown lower rates of fixation in the presence of high concentrations of soil mineral N and organic N inputs (Ledgard and Steele 1992). The biological responses to temperature are generally positive and Q₁₀ values (~~defined as the response factor for a 10 degree C temperature change-~~) are often in the range 1.5 to 3 ~~in the range of~~ for soil temperatures between 5°C and 25°C, outside which non-linearities are common. Taking a Q₁₀ of 2 and temperature increases by 2100 of 4°C, which appear probable (IPCC, 2013), the natural, terrestrial BNF in 2100 is likely to be 170 Tg N annually.




Figure 1: Summary of estimated global terrestrial contributions to biological N fixation in 2000. Values in Tg N yr⁻¹ (based on Table 1 and activity projections).

Table 1: Global terrestrial contributions to biological N fixation in 2000. Values in Tg N per year with the range of estimates in brackets.

Agricultural system or ecosystem	Organism	Annual N fixation(T g yr ⁻¹) and range	References
Grain legumes	Legume rhizobia	21 (10 – 21)	Herridge et al., 2008; Smil, 1999)
Forage legumes	Legume rhizobia	18.5 (12-25)	Herridge et al., 2008
Rice	Azolla	5 (4-6)	Herridge et al., 2008; Smil, 1999

Other croplands	Endophytic and free living bacteria	3.5	Herridge et al., 2008
Tropical savanas (used for agriculture)	Endophytic and free living bacteria	12 (5-42)	Cleveland et al., 1999; Herridge et al., 2008
Non agricultural ecosystems	Legume rhizobia and free living bacteria and algae	128 (44-290)	Cleveland et al., 1999; Galloway et al., 2004; Vitousek et al., 2013
Total		188 (77-387)	

2.2 Marine biological nitrogen fixation in the 21st century

Marine [biological](#) N fixation ~~is the biological conversion of dinitrogen gas (N₂) into NH₃, and~~ is performed by a diverse range of diazotrophs in plankton, microbial mat communities, sea grasses, coral reefs and sea sediments. Cyanobacteria of the genus *Trichodesmium* have been particularly well studied due to their prevalence and their formation of large blooms. Biological N fixation rates vary by species, and can be limited by temperature, light, oxygen, salinity, molybdenum, iron, and P. ~~As a result, there is a great deal of variability in both the species composition of diazotrophs present in the various ocean basins, as well as the rate of N fixation, which changes regionally and seasonally.~~

Estimates have been made for global N fixation in the oceans, both by extrapolating from biological measurements, and by modelling the biogeochemistry. ~~However, there is a great deal of uncertainty, due to the difficulties in accounting for the large regional and seasonal differences.~~ Recent reviews include Carpenter and Capone, 2008; Moore et al., 2013; and Voss et al., 2013.

Future changes to the ocean including increasing carbon dioxide (CO₂) concentrations, increasing stratification, and increasing temperatures, will likely result in an increase in marine nitrogen fixation. Nitrogen fixation leads to an increase in bioavailable N present in the form of ammonium and dissolved organic N (Mulholland et al., 2006). An increase in N₂ fixation would therefore lead to an increase in the amount of N_r available to enable further processes in the N cycle.

The objective of this section is to characterise the current state of knowledge ~~about on~~ marine BNF, ~~and the likely effects of changes in climate on marine BNF as well as the major areas of uncertainty regarding BNF trends in~~ the 21st century. ~~The various factors limiting marine N~~

726 fixation, the regional differences, and the predicted future impacts of changing conditions are
727 discussed in more detail below.

728 2.2.1 Factors affecting marine nitrogen fixation

729 2.2.1.1 ~~Light~~

730 ~~Light~~: Nitrogen fixers have strong preferences for specific light conditions. Depending upon
731 the species, either light or darkness is required. Many non-heterocystous cyanobacteria fix
732 nitrogen at night, however members of the genus *Trichodesmium* fix N only in the presence of
733 light (Capone et al., 1997). *Trichodesmium* are therefore present at the surface of the ocean,
734 and maximum fixation occurs at midday (Carpenter and Capone, 2008). Light sensitive
735 diazotrophs like *Trichodesmium* could be affected by decreasing solar irradiance due to the
736 presence of more clouds, resulting in a decrease in N₂ fixation.

737 2.2.1.2 Temperature

738 ~~Temperature~~: Enzyme activity generally increases with temperature, and this is true for
739 nitrogen-fixing enzymes (nitrogenases). Staal et al. (2003) found that on short time scales, three
740 strains of cyanobacteria exhibited a Q₁₀ ranging from 1.08 to 4.72. *Trichodesmium* exhibited a
741 Q₁₀ of 1.12 for N₂ fixation in darkness from 20-35°C, and a Q₁₀ of 2.06 from 15-20°C. In the
742 presence of light, *Trichodesmium* exhibited a Q₁₀ of 1.64 for 15-20°C, and 1.84 for 20-35°C.
743 Fu et al. (2014) exposed strains of *Trichodesmium* and *Crocospaera* to varying temperatures
744 in the laboratory and found maximum N fixation to occur between 24-28°C and 28-30°C,
745 respectively.

746 Increasing temperatures will likely cause the rate of N fixation to increase, both because
747 enzyme activity increases at higher temperatures, and because the increase in sea surface
748 temperatures will lead to an expansion of habitat suitable for diazotrophs (Hutchins et al.,
749 2009). Boyd and Doney (2002) predict that habitat expansion will lead to an increase in N
750 fixation of 27%.

751 Until recently, there was little evidence of marine diazotrophic activity in the cooler waters
752 present at high latitudes (>50 degrees) (Carpenter and Capone, 2008). A recent study found
753 substantial N fixation in the surface of the Canadian Arctic (Blais et al., 2012). These recent
754 discoveries suggest diazotrophs may be fixing N in areas previously thought to be too cold for
755 large levels of BNF.

756 2.2.1.3 Oxygen

757 ~~Oxygen~~: Most nitrogen-fixing enzymes are inactivated by oxygen. Diazotrophs
758 generally deal with this by performing N fixation either at night to avoid oxygen produced
759 during photosynthesis, or within thick walled cells called heterocysts which maintain a
760 localised anaerobic environment.

761 Nitrogen fixation has generally not been considered in oxygen minimum zone (OMZ) systems
762 (Carpenter and Capone, 2008). Due to the removal of N_r by denitrification and anaerobic

Formatted: Font: Italic

Formatted: Font: Italic

763 ammonium oxidation, OMZs have low concentrations of N_r relative to P (Canfield, 2006), and
764 the conditions in these sites may be suitable for N fixation. Modelling efforts have considered
765 N₂ fixation in OMZs (Canfield, 2006; Moore and Doney, 2007).

766 Expanding OMZs may increase areas conducive to denitrification and anaerobic ammonium
767 oxidation. If nitrogen-fixing bacteria exist in balance with denitrification (Deutsch et al., 2007),
768 then the increase in denitrification may lead to a corresponding increase in N₂ fixation. Oxygen
769 minimum zones may also lead to an increase in the release of trace metals (Noble et al., 2012)
770 and P from sediments, which could stimulate increased N₂ fixation.

771 2.2.1.4 Salinity

772 Salinity: Diazotrophs may be able to live in a variety of saline conditions. For example,
773 a *Trichodesmium* isolate was found to grow over a salinity range of 22-43 psu, but maximum
774 growth and nitrogenase activity occurred over a narrow range of 33-37 psu (Fu and Bell, 2003).
775 Changes in salinity are not expected to have a large effect on N fixation.

776 2.2.1.5 Trace metals and phosphorus

777 Trace metals and phosphorus: Nitrogenase requires both iron and molybdenum.
778 Nitrogen fixation is limited by iron in approximately 35-75% of the oceans, globally (Moore
779 et al., 2002; Berman-Frank et al., 2001). Molybdenum is generally not growth limiting (Paerl
780 et al., 1987; Paulsen et al., 1991) as it is readily present in seawater. However, sulphate may
781 inhibit the uptake of molybdenum, because sulphate is also present, and is stereochemically
782 similar to molybdate (Howarth and Cole, 1985; Marino et al., 2003).

783 Phosphorus is an essential nutrient necessary for organisms, however surface waters today are
784 thought to be more limited by N rather than P over much of the oceans (Moore et al., 2007).
785 Approximately 4% of the world oceans are limited by P (Moore et al., 2002).

786 Aeolian dust deposition leads to higher levels of iron reaching the subtropical North Atlantic
787 Ocean. Under present day conditions, P may therefore be more limiting for diazotrophs in the
788 North Atlantic, and iron may be more limiting in the North Pacific Ocean (Prospero and Lamb,
789 2003). Climate change may affect the transport of aeolian dust. If drier areas become drier,
790 and/or wind speed increases, the amount of dust transported from continents to the oceans may
791 increase, which would increase nitrogen fixation in areas limited by iron. However, if the areas
792 that receive the dust are limited by other nutrients, then the increase in dust transport would
793 have little effect.

794 2.2.1.6 Stratification

795 Stratification: A strengthening of ocean stratification may lead to a decrease in nutrient
796 upwelling, which would in turn lead to a shortage of N at the surface, which may cause an
797 expansion of nitrogen-limited subtropical gyres (Sarmiento et al., 2004) and possibly
798 encourage an increased rate of N fixation.

799 2.2.1.7 Carbon dioxide

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

~~Carbon dioxide:~~ Both model and laboratory studies of *Trichodesmium* isolates have shown an increase in N₂ fixation associated with increasing atmospheric CO₂ concentrations. Studies with *Trichodesmium* cultures have reported a range of measurements for the increase in N₂ fixation associated with increasing CO₂ concentrations from present day levels (375-380 ppm) to projected 2100 levels (~750-1000 ppm). Studies have reported an increase in rates of around 35-65% (Hutchins et al., 2007; Barcelos e Ramos et al., 2007; Kranz et al., 2009), and as high as 100-121% (Hutchins et al., 2007; Levitan et al., 2007). Barcelos e Ramos et al. (2007) predicted that N₂ fixation rates for *Trichodesmium* would increase by 50% from 60-85 Tg N yr⁻¹ in 2005 to 90-128 Tg N yr⁻¹ by year 2100 with projected increases in CO₂ concentrations under a business-as-usual emission scenario (scenario IS92a).

Hutchins et al. (2009) estimated that N₂ fixation by *Trichodesmium* alone will rise from present day levels of 60 Tg N yr⁻¹ (Mahaffey et al., 2005) to 80-100 Tg N yr⁻¹ by 2100, based on the response of a *Trichodesmium* isolate to increasing CO₂ levels. Hutchins et al. (2007) found that N₂ fixation rates for *Trichodesmium* levelled off at 1250 and 1500ppm, suggesting that N₂ fixation rates may stop increasing with increasing CO₂ levels by the year 2100.

Recent evidence indicates that unicellular cyanobacteria may fix at least as much N as *Trichodesmium* (Montoya et al., 2004). A laboratory study using the unicellular cyanobacterium *Crocospaera watsonii* found that elevating CO₂ levels from 380ppm to 750ppm increased N₂ fixation rates by 40% (Fu et al., 2008), when not limited by iron. Based on measurements of the increase in N₂ fixation rates associated with CO₂ increases for seven strains of *Trichodesmium* and *Crocospaera*, Hutchins et al. (2013) predict that over the next 100 years, N₂ fixation rates will increase by 4-23% for these seven strains. More evidence is needed to determine if other diazotrophs will be similarly affected by rising CO₂ concentrations.

~~Increasing CO₂ concentrations will likely also lead to an increase in N fixation, although this increase may be offset by effects which decrease N₂ fixation.~~ Barcelos e Ramos et al. (2007) predicted that N₂ fixation rates would increase by 50% by year 2100 with projected increases in CO₂ concentration_s, ~~providing a strong negative feedback to increases in atmospheric CO₂.~~

Anthropogenic N fertilization of the ocean leads to an increase in marine uptake of CO₂, however this may lead to an increase in N₂O emissions. Duce et al. (2008) applied Redfield stoichiometry to estimates of anthropogenic N_r deposition of 54 Tg N yr⁻¹ and anthropogenic CO₂ uptake by the ocean of $\sim 2.2 \pm 0.5$ Pg C yr⁻¹ (IPCC, 2007), and calculated that the ocean may take up an additional 10% of atmospheric anthropogenic CO₂ as a result of atmospheric deposition of N_r. However, up to two-thirds of the decrease in radiative forcing generated by this drawdown of CO₂ may be offset by an increase in radiative forcing associated with an increase in the emissions of N₂O (Duce et al, 2008). A decrease in pH due to ocean acidification from rising CO₂ levels may lead to a decrease in the bioavailability of iron (Shi et al., 2010), which may in turn lead to a decrease in N₂ fixation for diazotrophs in areas where iron is limiting.

839 Table 2 provides a summary of the factors influencing marine N fixation, and the expected
840 effects on marine BNF in the 21st century.

841 **Table 2.** Summary of future impacts of factors affecting marine nitrogen fixation.

Factor	Effect on N ₂ Fixation
CO ₂ increase (and decrease in pH)	+ 35% to 121% by 2100
Temperature increase leading to expansion of diazotroph habitat	+ 27%
Temperature increase leading to faster enzyme activity	+
Stratification leading to shortage of nutrients in surface waters	+
Decreasing solar irradiance due to increased cloud cover	-?
Dust containing iron	+ or -
Increase in oxygen minimum zones	+
Increase in nitrogen export from rivers	-
Increase in deposition of reactive nitrogen	-
Improved measurement methods	+
Phosphorus	Limiting nutrient

Formatted Table

842

843 2.2.2 Present-day and pre-industrial estimates

844 Estimates of global ocean N₂ fixation (shown in Fig. 2) range from 75 to 200 Tg N yr⁻¹
845 (Galloway et al., 2004; Carpenter and Capone, 2008; Moore et al., 2006; Deutsch et al., 2007;
846 Eugster and Gruber, 2012; Luo et al., 2012), with recent estimates at around 130-140 Tg N yr⁻¹
847 (Deutsch et al., 2007; Eugster and Gruber, 2012; Luo et al., 2012). Deutsch et al. (2007)
848 estimated global ocean N fixation to be 140 Tg N yr⁻¹, using observed nutrient concentrations
849 and an ocean circulation model. Eugster and Gruber (2012) used two methods to estimate the
850 preindustrial global nitrogen fixation rate in the oceans to be 131 Tg N yr⁻¹ (94, 175) and 134
851 Tg N yr⁻¹ (117, 150), by combining geochemical observations with a two-dimensional box
852 model. Deutsch et al. (2007) and Eugster and Gruber (2012) found that the rates of N₂ fixation
853 were higher in the Pacific Ocean than the Atlantic. Luo et al. (2012) compiled a global database
854 of diazotroph abundances and N₂ fixation rates, and estimated the global pelagic (open ocean)
855 N₂ fixation rate to be 140 ± 9.2 Tg N yr⁻¹ (arithmetic mean ± one standard error). One possible
856 limitation of this [approach data set](#) is that 99% of the data were collected within the range of
857 40°S to 55°N, and if substantial N₂ fixation is found to occur outside of this range, [this estimate](#)
858 [it](#) may be an underestimate. Luo et al. (2014) applied a multiple linear regression model to the

859 same database of field observations and found an estimate of N₂ fixation of 74 (51–110) Tg N
860 yr⁻¹ for the open ocean.

861 Luo et al. (2012) note that the most common method for field measurements of N₂ fixation has
862 recently been found to underestimate the rates for *Trichodesmium* by 62% (Großkopf et al.,
863 2012). ~~so future estimates may be higher.~~ Extrapolating from the differences found between
864 the ¹⁵N₂ tracer bubble-addition and dissolution methods, Großkopf et al. (2012) estimate that
865 the global marine N fixation rate measured using the new method would be 177 ± 8 Tg N yr⁻¹.

866 Although recent midpoint estimates appear to have coalesced at around 130–140 Tg N yr⁻¹,
867 there is still a great deal of uncertainty due to the large variance in measurements ~~(5 to 8 orders~~
868 ~~of magnitude)~~ (Luo et al., 2012), and recent measurements of nitrogen fixation rates in areas
869 not previously thought to have high levels of diazotrophy.

870 ~~2.2.3 Future Effects of global change on marine biological nitrogen fixation impacts~~

871 ~~Many factors will affect future rates of BNF.~~ The most ~~important~~~~dramatic~~ effects will likely be
872 due to temperature and increasing CO₂ concentrations. ~~We estimate that m~~Marine BNF will
873 increase from present day estimates of 140 (100–200) Tg N yr⁻¹ to 166 (120–240) Tg N yr⁻¹ due
874 to temperature effects alone. Present day BNF estimates were scaled up using the Q₁₀ of 1.64
875 for *Trichodesmium* (15–20°C) ~~found by~~ (Staal et al. ~~(2003)~~).

876 ~~The implications of the various factors affecting BNF for the future are described in more detail~~
877 ~~below.~~

878 ~~Diazotrophs have a competitive advantage over nondiazotrophs under conditions when there~~
879 ~~is a shortage of N_i relative to iron and P (Dutkiewicz et al., 2012). The projected increase in~~
880 ~~atmospheric deposition and river export of N_i to the ocean may therefore lead to a decrease in~~
881 ~~N fixation.~~

882 In addition to the factors discussed above, estimates of N fixation may increase in the future
883 even if the ~~actual~~~~true~~ rate remained constant. This is because the most common method for
884 taking field measurements of marine N fixation has recently been found to underestimate the
885 rate, so future estimates of N fixation may increase as the methods become more accurate
886 (Großkopf et al., 2012). In addition, recent evidence suggests that regions such as coastal,
887 aphotic, and arctic regions may exhibit more N₂ fixation than previously thought.

888 Taken together, the factors discussed above suggest that ~~marine~~ N fixation ~~will~~~~may~~ increase in
889 the future, which ~~may~~~~will likely~~ lead to an increase in ocean drawdown of CO₂. Several
890 feedbacks may offset this increase. Increasing rates of N₂ fixation may drive areas to P and iron
891 limitation, thereby limiting ultimate N₂ fixation rates. ~~The many complex factors influencing~~
892 ~~N fixation make it difficult to draw strong conclusions about future rates of marine N fixation~~
893 ~~and its influence on the C cycle.~~

894 ~~2.3.4 Global changes in natural BNF 2010 to 2100.~~

Formatted: Font: Bold

Formatted: Font: Not Italic


Formatted: Font: Bold

Formatted: Font: Bold

895 ~~It appears likely that overall-global BNF will increase during this century in marine and~~
 896 ~~terrestrial ecosystems, both managed and natural ecosystems.~~ The total terrestrial natural N
 897 fixation by the end of this century suggested from these arguments is ~~170-200~~ Tg N annually,
 898 approximately ~~40%-40%~~ larger than the value at the beginning of the 20th century. Marine BNF
 899 is projected to increase from ~~120 TgN yr⁻¹ to 166 TgN yr⁻¹ by 2100, an increase of 38% on the~~
 900 ~~2010 value, 273 to 400 Tg N yr⁻¹ by 2100, an increase of 45% on 2010 values.~~

Formatted: Subscript

901



902

903 **Figure 2:** Summary of pre-industrial (blue), present (green), and future (purple) estimates of marine biological
 904 nitrogen fixation (BNF). Estimate from Carpenter and Capone (2008) represents their summary of the range
 905 presented in the literature, and includes no midpoint. Luo et al., 2012 values are arithmetic mean \pm standard error,
 906 so range limits may not be directly comparable to other estimate range limits. Estimates for fixation by
 907 *Trichodesmium* alone by Barcelos e Ramos et al. (2007) (60-85 Tg N yr⁻¹ in 2005, 90-128 Tg N yr⁻¹ by year 2100)
 908 and Hutchins et al. (2009) (80-100 Tg N yr⁻¹ by 2100) are included in text but not presented in figure because the
 909 estimates in the figure are for total marine BNF.

910

911 3 ANTHROPOGENIC FIXATION OF NITROGEN IN THE 21ST CENTURY

912 ~~The Human~~ demand for fixed N through the 21st century will be driven by requirements for
 913 food and industrial N_r use. ~~There is also unintended nitrogen fixation resulting from plus~~
 914 combustion related ~~NO_x emissions, emissions~~ Set against these drivers for increased N_r
 915 fixation, ~~and the extent of~~ control measures to mitigate emissions ~~will regulate the net~~
 916 ~~anthropogenic net anthropogenic contribution to global nitrogen fixation, and their effects.~~ A
 917 range of authors have considered the available scenarios and possible development trajectories
 918 including most recently Winiwarter et al. (2013) and Bouwman et al. (2013). The scenarios
 919 and time scales used by these authors differ, with Bouwman et al. (2013) projecting trends ~~only~~
 920 to 2050, rather than the end of the century. However, given the uncertainty in projections of
 921 this kind, they provide a useful guide for the likely trends. The projections from Winiwarter et
 922 al. (2013) are based on story lines and methodologies similar to those of the RCP

Formatted: Subscript

(Representative Concentration Pathways) as used in the 2014 IPCC assessment of climate change through the 21st century. However, while based on the RCP scenarios, only the N_r from combustion is taken directly from the IPCC methods as N_r formation was not a focus of the scenario developments for the IPCC. Together with mineral fertilizer and industrial use of N_r, fixation is projected in the range -140 Tg N yr⁻¹ to 235 Tg N yr⁻¹ by 2100, depending on the RCP chosen and compares with their estimate of 170 Tg N yr⁻¹ in 2000. The year 2000 value is smaller than the estimate of 210 Tg N yr⁻¹ by Fowler et al. (2013), but within the uncertainties shown in each synthesis. The projections from Winiwarter et al. (2013) imply modest overall change in N_r production by human activity through the 21st century as a consequence of gradual increases in efficiency compensating for increases in demand for fertilizer and industrial N_r applications, combined with the considerable improvements expected for reductions in nitrogen oxide (NO_x) emissions from combustion resulting from expected emission controls. Indeed, there have been important reductions in emissions of combustion N_r, as NO_x to the atmosphere throughout Europe, North America and other highly developed economies. Typically these have reduced NO_x emissions by about 50% over the last 30 years in these regions. Similar controls are likely for combustion emissions in the rapidly developing economies of Asia in the decades ahead. However, for reduced N, the global trend has been a monotonic increase in N_r fixation for most countries in the world outside Europe, and the social trends in rapidly developing economies towards increased meat consumption seem likely to continue the trend. Given these historical trends and the unwillingness of governments throughout the world to regulate the supply of reduced N_r for agriculture and industry, the assumption that N_r production will remain constant through the 21st century seems implausible.

A substantial increase in nitrogen use efficiency (NUE) seems likely, as has been achieved in European agriculture over the last 30 years, but this is unlikely to prevent a continued increase in global agricultural nitrogen use. Given that human N_r production doubled between 1980 and 2010, a period in which global population increased by 2.5 billion, and medium-most projections estimates project show a similar population increase by the later years of during the 21st century, the demand for food and other nitrogen consuming activities (transport, heating and consumer goods) will most likely lead to a substantial increase more in industrial N fixation. Assuming NUE increases, it is possible that anthropogenic N fixation only grows only by 30% between 2010 and 2100. This simplistic assumption would lead to 2100 N_r production through the Haber-Bosch process of 160 Tg-N yr⁻¹ and total annual anthropogenic production N_r of 247 Tg N yr⁻¹.





Figure 3: Global biological nitrogen fixation in natural ecosystems and by human activity in 2010 (left) and 2100 (right).

The global changes in fixation discussed above are summarised in Fig. 3, which show large increases in the total N fixed from 446 Tg N yr^{-1} in 2010 to 586 Tg N yr^{-1} in 2100 accompanied by substantial increases in the uncertainties of the component fluxes.

The N_r fixed by BNF and human activity is then used by and transformed within ecosystems and products of the chemical and biological processing cascade through terrestrial and marine ecosystems and the atmosphere. It is important now to consider the effect of changes in the environment this century on the fate of the N_r .

4 EFFECTS OF ENVIRONMENTAL CHANGES ON THE FATE OF N_r IN TERRESTRIAL AND MARINE ECOSYSTEMS

4. EFFECTS OF ENVIRONMENTAL CHANGES ON THE FATE OF N_r IN TERRESTRIAL AND MARINE ECOSYSTEMS

Formatted: No bullets or numbering

The total fixation of N through natural (BNF), combustion and Haber Bosch processes is projected to increase during the remainder of the 21st century, possibly to approximately 600 Tg N, an increase of 50% over values at the beginning of the century, ~~but subject to large uncertainties~~ (Fig. 3). The subsequent fate of the N_r in terrestrial and marine ecosystems and the responses of the different pools of N_r to changes in climate, and especially temperature ~~and hydrology~~, are now considered for terrestrial and marine ecosystems.

The fixed N, whether by natural processes in soils and the oceans or by human activities is predominantly in the reduced form as ammonia (NH₃) or ammonium (NH₄⁺) initially. Once formed, N_r is readily transformed in the environment and it is important to describe the likely effects of changes in the environment on the fate of N_r, and quantify, where possible the probable impacts due to climate and land use change, ~~in this century~~. In short, which are the components of the N cycle that are most responsive to expected changes in climate and land use ~~in this century~~?

4.1 Terrestrial ecosystems: Emissions of NH₃ from terrestrial ecosystems through the 21st century

The global total emissions of NH₃ to the atmosphere at the beginning of the 21st century have been estimated by Sutton et al. (2013b), at 59.3 Tg N yr⁻¹ of which 33 Tg N yr⁻¹ is from livestock and crops. The N_r fixed industrially through NH₃ manufacture, mainly for fertilizers, is currently 120 Tg N yr⁻¹, thus the emissions to the atmosphere from livestock and crops represent roughly a quarter of the annual fertilizer production ~~annually~~, effectively fertilizing the atmosphere. This substantial quantity is ~~of course simply~~ the unintentional leakage of the N_r from farming systems, due to the volatility of NH₃. Also presented in this analysis of global emissions of NH₃ are values for emissions from all other major sources. Given the spatial and temporal variability in emission rates and the sensitivity to climate, and especially temperature, the range of different emission estimates is small among the seven different estimates (35 to 65 Tg yr⁻¹) summarised, which reflects the fact that these estimates are not fully independent. It is suggested by Sutton et al. (2013b) that overall uncertainty is around ±30%, pointing to a range of emissions ~~a uncertainty range~~ for 2008 of 46 Tg N yr⁻¹ to 85 Tg N yr⁻¹.

The forces which have governed overall industrial production of fixed N have largely been economic, responding to the demand for food and the response functions between crop productivity and fertilizer use (Jensen et al., 2011) in which the economic benefits of increased yields have driven global N fertilizer use demand. At the same time, an increase in global meat consumption per capita (Erisman et al., 2008) ~~will have~~ has magnified fertilizer requirements and NH₃ emissions (Westhoek et al., 2014). This includes both the NH₃ emissions from fertilizer in growing animal feeds and the ammonia emissions from livestock manures, in animal houses, manure storage, land application and from grazing animals, where the used of housed livestock substantially increases emissions compared with pasture-only systems.

Global projections for future N_r use have not generally included possible control measures to reduce emissions of NH₃ to the atmosphere, which would increase the NUE. ~~There are~~ exceptions, in the case of the Netherlands and in Denmark, where policies to reduce the leakage

of N_r to the environment led to substantial reductions in atmospheric emissions (EMEP, 2014). Although first NUE estimates have now been provided for each country in the world (Sutton et al., 2013a), it is a matter for ongoing and future analysis to show how these have evolved over time and to demonstrate the quantitative relationships between reduction between N emissions, including NH_3 , and improvement of NUE. In the global projections of N_r use through the 21st century provided by both Erismann et al. (2008, using the SRES approach) and Winiwarter et al. (2013, using the RCP approach), scenarios included the potential to improve crop NUE, while Sutton et al. (2013a), examined the N savings possible also as a result of improving NUE across the full agri-food chain.

While subject to uncertainty, the global demand for food is likely to increase by 40% by 2050 due to population growth and a changing diet (Godfray et al., 2010), especially in the rapidly developing regions, unless additional actions are taken. The largest uncertainties in estimating future emissions of NH_3 to the atmosphere are the consumption drivers (food amount, food choice), the amounts of fertilizer and manure N applied and the effect of climate on the fraction emitted (van Vuuren et al., 2011a; and Sutton et al., 2013b, respectively). Excluding the climatic interaction (which is addressed below), emissions resulting from demand for food and industrial uses, have been estimated by van Vuuren et al. (2011a) to increase from 60 Tg N yr⁻¹ in 2000 to between 70 and 80 Tg N yr⁻¹ by 2100.

4.1.1 Effects of changes in climate on terrestrial emissions of NH_3

The processes of exchange of NH_3 between terrestrial ecosystems and the atmosphere have been subject to detailed field studies and intercomparisons of methods (Sutton et al., 1995, 1998, 2009; Flechard et al. 1999, 2013) and are discussed further in this review. The most recent estimates of the influence of climate change on emissions of NH_3 are by Sutton et al (2013b).

The surface atmosphere exchange of NH_3 is generally described numerically using a resistance analogy in which the vertical flux (F_t), is given by the potential difference between the surface ($\chi(z_0)$) and a reference height in the atmosphere ($\chi(z)$) divided by the sum of resistances in the pathway from the source to the reference height and comprising $R_a(z)$ and R_b , the turbulent atmospheric and quasi-laminar boundary layer resistances respectively.

$$F_t = [\chi(z_0) - \chi(z)] / [R_a(z) + R_b] \quad \text{Equation (1)}$$

In most ecosystems, the concentration at the surface, ($\chi(z_0)$) is non-zero, due to presence of NH_4^+ in the apoplast of vegetation. In these conditions the value of $\chi(z_0)$ is proportional to a ratio $\chi = [NH_4^+]/[H^+]$ of the canopy/ground surface, where according to the thermodynamics:

$$\chi = 161500/T \exp^{(-10380/T)} [NH_4^+]/[H^+] \quad \text{Equation (2)}$$


Temperatures (T) are in Kelvin and the scheme is represented schematically in Fig. 4.

The bi-directional exchange between surface and atmosphere of NH_3 modifies the spatial patterns of NH_3 fluxes in the landscape, with reduced emission or even deposition downwind of large sources (Fowler et al., 1998). Quantifying changes in NH_3 emission this

1052 century requires knowledge of apoplast and leaf litter NH_4^+ and pH, scaled through the coming
 1053 decades over global vegetation. The data [required](#) to calculate net exchange fluxes in this way
 1054 are not available. However, Sutton et al. (2013b) argue that by examining model ecosystems
 1055 and their exchange of NH_3 a surrogate for the likely change may be seen in empirical data.
 1056 When it comes to global upscaling of NH_3 emissions, this also needs to bear in mind that the
 1057 wide range of different terrestrial NH_3 sources are likely to have differing temperature
 1058 responses, due to the role of different interacting factors.

1059 To illustrate these effects, a model ecosystem was used for which both a global modelling
 1060 framework and field measurements are uniquely available, namely NH_3 emission from seabird
 1061 colonies. In addition to the availability of measurements and modelling, they are also globally
 1062 important sources of NH_3 and are distributed geographically across a broad range of climates,
 1063 with minimal human intervention, so that the effects of climate differences can be assessed
 1064 without confounding management interactions (Blackall et al., 2007; Riddick et al., 2012;
 1065 Sutton et al., 2013b). This approach demonstrated a strong climate dependence in the ammonia
 1066 emissions, with the modelling approach (incorporating Eq. (2)), agreeing closely with the
 1067 measured datasets.

1068
 1069



1070

1071 **Figure 4:** A resistance analogue of NH_3 exchange including cuticular, stomatal and pathways to soil (Sutton et
 1072 al., 2013b). Two methods for cuticular exchange schemes are shown: 1, Steady-state uptake according to a
 1073 varying cuticular resistance (R_w); 2, Dynamic exchange with a reservoir of NH_4^+ using a varying capacitance (C_d)
 1074 and charge (Q_d). Within-canopy transfer (R_{ac}), cuticular adsorption/desorption (R_d) and stomatal exchange (R_s).
 1075 Also shown are the air concentration (χ_a), cuticular concentration (χ_d), stomatal compensation point (χ_s), litter/soil
 1076 surface concentration (χ_l) and the canopy compensation point (χ_c).

Combining all sources of NH₃ emission globally, studies provided the data to model likely responses of terrestrial NH₃ emissions to a 5 degree increase in global temperature and showed that emissions in 2008 of 65 Tg-NH₃-N (45-85), increased to 93 (64-125) Tg-NH₃-N in 2100 (Sutton et al., 2013b), based on anthropogenic activity levels for 2008. This may be compared with an estimated increase in NH₃ emissions based on increased anthropogenic activities (excluding the climatic response), and of no-change for natural sources, of 42% (33- 52%) increase by 2100. Combining the increases in anthropogenic activity expected up to 2100 according to the RCP8.5 (Lamarque et al., 2011), with the estimated effect of climate warming on emissions, gives an overall estimate of NH₃ emissions for 2100 of 132(89-179) Tg N yr⁻¹. As Sutton et al. (2013b, supplementary material) point out, this value is nearly a factor of three higher than that included in the currently mapped EDGAR database, which is a consequence of including: a) additional sources (including oceans, see further below), b) the effect of the climate change feedback and c) the anticipated increase in anthropogenic activities.

4.2 Ammonia exchange over the oceans in the 21st century

In marine ecosystems NH₃/NH₄⁺ is produced by phytoplankton and other organisms. Although the aqueous-phase partitioning between NH₃ and its protonated form NH₄⁺ is dominated by NH₄⁺, the majority of emissions are in the form of NH₃. Ammonium is quickly assimilated by phytoplankton, so NH₃ and NH₄⁺ are usually present in low concentrations in the surface ocean.

4.2.1 Factors affecting the flux of ammonia between the atmosphere and the ocean

The exchange of ammonia between the ocean and the atmosphere depends on several factors: the concentrations of ammonia in the surface layer of the ocean and in the boundary layer of the atmosphere, temperature, and wind speed (Johnson et al., 2008). The flux across the atmosphere-ocean interface can be described by (Liss and Slater, 1974):

$$F = k_g k_H \{ (NH_{3(g)}) - K_H [NH_{3(sw)}] \} \quad \text{Equation (3)}$$

where F is the flux between the atmosphere and the ocean (mol m⁻² s⁻¹), k_g is the gas-phase transfer velocity (m s⁻¹), the NH₃ concentrations are given in mol m⁻³, and K_H is the dimensionless Henry's Law coefficient for ammonia. NH_{3(sw)} refers to surface water NH₃ concentration.

The Henry's law constant for ammonia can be calculated as follows (McKee, 2001):

$$K_H = (17.93 - (T/273.15)) e^{(4092/T) - 9.70} \quad \text{Equation (4)}$$

where T is temperature in Kelvin.

The concentration of NH₃ present in seawater depends on the partitioning between NH₃ and NH₄⁺, which is affected by pH, salinity, and temperature. This dissociation can be described by the logarithmic acid dissociation constant, pK_a (Bell et al., 2007):

$$pK_a = 10.0423 - (0.0315536 T) + (0.003071 S) \quad \text{Equation (5)}$$

1112 where T is the temperature in °C, and S is salinity in g/kg. Chemical reactions and transport of
1113 NH₃ into the atmosphere (from terrestrial emissions) and the ocean (from biological activity,
1114 deposition and river export) also affect the levels of NH₃ present.

1115 4.2.2 Flux estimates


1116 The present-day direction of NH₃ flux is believed to be from the atmosphere to the oceans at
1117 high latitudes, where the oceans are colder, allowing more gases such as NH₃ to be dissolved.
1118 In contrast, the oceans are believed to be a source of NH₃ emissions at lower latitudes, where
1119 the oceans are warmer, promoting a greater partitioning to the gas phase (Eq. (4)) and Johnson
1120 et al., 2008). When considering the global oceans together, the net flux is believed to be a small
1121 emission from the oceans to the atmosphere (Bouwman et al., 1997; Dentener and Crutzen,
1122 1994; Galloway et al., 2004).

1123 Ocean-atmosphere NH₃ fluxes vary across regions and seasons, and observations are limited,
1124 ~~making it difficult to extrapolate regionally and temporally.~~ As a result, there are only a few
1125 quantitative estimates of global flux, all of which are highly uncertain. Dentener and Crutzen
1126 (1994) estimated the flux to be 7.0 Tg N yr⁻¹ from the oceans to the atmosphere. They did this
1127 by taking a distribution of dimethylsulphide (DMS) emissions modelled by Bates et al. (1987),
1128 and assuming equal molar emissions of NH₃ and DMS. Although both DMS and NH₃ are
1129 produced by phytoplankton, assuming an equal molar relationship is acknowledged by
1130 Dentener and Crutzen (1994) to be ~~rather~~ speculative. The relationship between NH₃ and DMS
1131 may have been tightly coupled under pre-industrial conditions, but this is unlikely to be true
1132 under the present strong anthropogenic influences on the N cycle (Johnson and Bell, 2008).
1133 ~~especially in coastal waters.~~

1134 An independent estimate of global ocean NH₃ emissions was provided by Bouwman et al.,
1135 (1997) who applied an ocean carbon cycle model to calculate an NH₃ flux of 8.2 Tg N yr⁻¹ for
1136 1990. However, this did not account for non-zero atmospheric NH₃ concentrations, and
1137 Bouwman et al. (1997) acknowledged that doing so might reduce the net sea-atmosphere
1138 emission flux by a factor of two. For comparison, Galloway et al. (2004) estimated
1139 preindustrial, present, and future marine NH₃ emissions using a compensation point approach
1140 to be 5.6 Tg N yr⁻¹. However, it is unlikely that the flux would remain constant over these time
1141 periods, given the human perturbations to the N cycle.

1142 Steadman et al. (in preparation), have improved ~~on~~ these estimates by implementing the bi-
1143 directional flux calculation method described by Johnson et al. (2008), following Eqs. (3-5).
1144 ~~This—accountings~~ for both regional and temporal patterns in ocean and atmospheric
1145 concentrations of NH₃ and temperature. The flux is calculated by dividing the ocean surface
1146 into 5 degree grid squares, and determining the gas transfer velocity and the Henry's law
1147 constant for NH₃ within each grid square, using temperature, pH, and wind speed. The resulting
1148 estimated NH₃ flux for 2005 is 5.7 Tg N yr⁻¹ from the ocean to the atmosphere. Atmospheric
1149 concentrations of NH₃ were obtained from STOCHEM (Lamarque et al., 2013; Derwent et al.,
1150 2003) model output. Surface ocean NH₃ and NH₄⁺ concentrations were obtained from the
1151 British Oceanographic Data Centre.

1152 These recent analyses suggest that the global NH_3 emission estimate of 8.2 Tg N yr^{-1} of
 1153 Bouwman et al. (1997), which was incorporated into the global emissions estimates of Sutton
 1154 et al. (2013b), summarised above, may be overestimated. If so, the estimates of Sutton et al.
 1155 (2013b) should be reduced by around 2.5 Tg N yr^{-1} for 2008, giving total emissions of around
 1156 $63 (44\text{--}82) \text{ Tg N yr}^{-1}$, again based on $\pm 30\%$ uncertainty.



1157
 1158 **Figure 5:** Summary of pre-industrial (blue), present (green), and future (purple, for 2100) estimates of marine
 1159 ammonia flux from ocean to atmosphere.

1160 1161 4.2.3 Future impacts

1162 The future ocean-atmosphere flux of NH_3 will be affected by increasing temperatures,
 1163 increasing terrestrial NH_3 emissions, and ocean acidification from elevated CO_2 levels which
 1164 lowers the pH of the water. The mechanism of both of these effects is an alteration of the
 1165 partitioning of ammonia and ammonium in the ocean, as illustrated by Eq. (5). The lower pH
 1166 results in a greater relative concentration of ammonium. The lower concentration of NH_3 will
 1167 therefore result in lower emissions. Climate change and ocean acidification will also have
 1168 indirect effects on ocean ecosystems, leading to changes in plankton populations and species
 1169 composition.

1170 The decreased marine emissions of NH_3 in the future, combined with increasing N deposition
 1171 and export from rivers, suggest that the future oceans may accumulate more reactive N, leading
 1172 to eutrophication and OMZs. Some of the additional N_r may result in an increase in
 1173 denitrification and associated N_2O emissions.

1174 The expected temperature and pH changes in the ocean associated with climate change and
 1175 ocean acidification will likely have a large effect on the ammonia flux. Based on the estimates
 1176 of Bouwman et al. (1997), Eq. (2) and a 5°C warming scenario, Sutton et al. (2013b,
 1177 supplementary material) estimated that ocean NH_3 emissions would increase to 15 Tg N yr^{-1} .

1178 However, as noted above, the baseline may have been an overestimate, while the interaction
1179 with rising CO₂ levels was not included.

1180 Preliminary model results suggest that after accounting for the increasing temperatures and
1181 terrestrial emissions associated with RCP8.5, and the expected ocean acidification (a decrease
1182 in mean surface ocean pH of 0.31, from 8.14 in 2000 to 7.83 in 2100 (IPCC, 2013)), the
1183 estimated future NH₃ flux for 2100 is 1.7 Tg N yr⁻¹. ~~However, the flux depends greatly on~~
1184 ~~temperature, ocean acidification, and terrestrial NH₃ emission estimates.~~ If temperature
1185 increases and increasing terrestrial ammonia emissions are accounted for, but ocean
1186 acidification neglected (the effect of pH is excluded), the estimated emission for 2100 would
1187 be 9.2 Tg N yr⁻¹. If atmospheric NH₃ concentrations and ocean pH were to remain at 2000
1188 levels, but temperatures increase as expected under RCP8.5, the estimated 2100 ammonia
1189 emission is 10.9 Tg N yr⁻¹. Comparison of the bars in Fig. 5 shows that in relative terms the
1190 effect of ocean acidification is the largest driver, providing more than a factor of three
1191 difference in the flux calculated by Eq. (3).

1192 ~~These varying estimates demonstrate the high level of uncertainty in the magnitude of future~~
1193 ~~atmosphere-ocean fluxes, and show that climate change and terrestrial NH₃ emissions may lead~~
1194 ~~to dramatic changes in global NH₃ processes. These estimates do not account for changes in~~
1195 ~~oceanic NH₃ concentrations, which will likely increase due to increased biological activity,~~
1196 ~~export of nutrients from rivers, and atmospheric deposition. These changes may be especially~~
1197 ~~dramatic in coastal areas, which are strongly influenced by anthropogenic activity and would~~
1198 ~~contribute to increased NH₃ emissions from the coastal zone. Databases of existing~~
1199 ~~observations and projections from marine biogeochemical models will help improve our~~
1200 ~~understanding of future fluxes.~~

1201 4.3 — Terrestrial emissions of nitric oxide and nitrous oxide

1202 4.3.1- Global sources of NO and N₂O in the atmosphere

1203 4.3.1.1 NO_x

1204 NO_x. Sources of atmospheric NO_x (NO+NO₂) are soils, natural fires, lightning, transport from
1205 the stratosphere and combustion of fossil fuels. The sinks are in both soil through microbial
1206 uptake and the atmosphere, through reactions with OH (Miyazaki et al., 2012; Logan et al.,
1207 1983). ~~In the troposphere the highly reactive, short-lived molecule NO is in equilibrium with~~
1208 ~~NO₂, through photochemical formation and destruction of ozone (O₃).~~ Global NO_x emissions
1209 have increased 3 to 6 fold since the industrial revolution due to increased fossil fuel and
1210 biomass burning (Prather and Ehhalt, 2001). Recent new estimates of global NO_x emissions
1211 based on a combination of a top down inventory based on satellite observations, and bottom-
1212 up inventory, ~~the a posteriori inventory,~~ have constrained the global emissions budget to 40 Tg
1213 N yr⁻¹ (Jaeglé et al., 2005; Martin et al., 2003). Fuel combustion (fossil and biofuel) were the
1214 largest source, contributing 58% to the total budget, followed by soils (22%), biomass burning
1215 (14%), lightning (8%), stratospheric/tropospheric exchange (0.2%) and aircraft (0.1%) (Jaeglé
1216 et al., 2005; Martin et al., 2003). Largest soil contributions were from the African and
1217 Australian continents (39% of total), whereas in the more industrialised ~~and wealthier~~ US and

Formatted: Font: Italic

1218 Europe soil emissions contributed only with 12 and 18% to total emissions [respectively](#) (Fig.
1219 6). ~~Discrepancies between the bottom-up inventory and the *a posteriori* inventory were largest~~
1220 ~~for soil emissions, by 68%.~~ The monthly satellite NO_x data, links peak soil derived NO_x
1221 emissions with the onset of the rainy seasons in North equatorial Africa, and N fertilization of
1222 agricultural land in the northern and mid latitudes. ~~These observations imply that the Yienger~~
1223 ~~and Levy (1995) emission factors together with the Wang et al. (1998) algorithm for canopy~~
1224 ~~exchange need to be revised upward substantially (Jaeglé et al., 2005).~~

1225 Hudman et al. (2012) improved the presentation of soil NO_x emissions in global models by
1226 replacing the simple emission factors (Yienger and Levy, 1995) with equations representing
1227 spatial and temporal patterns of soil moisture, temperature, pulsing, fertilizer, manure and
1228 atmospheric N deposition and biome. ~~This~~ BDSNP model (Berkeley-Dalhousie Soil NO_x
1229 Parameterization) was coupled to [a](#) global [chemistry](#)-transport model GEOS-Chem, which
1230 normally used the Yienger and Levy (1995) (YL95) scheme for soil emissions (Wang et al.,
1231 1998), but retained the YL95 canopy reduction component. The new model calculated larger
1232 emissions for the below canopy emissions (10.7 Tg N yr⁻¹) relative to the YL95 approach (7.4
1233 Tg N yr⁻¹). Total above canopy soil NO_x emissions were calculated at 9 Tg N yr⁻¹, in good
1234 agreement with the Jaeglé et al. (2005) study. The new model was validated using satellite
1235 nitrogen dioxide (NO₂) data provided by OMI (Ozone Monitoring Instrument, Hudman et al.,
1236 2010). Their model was able to reproduce the monsoon induced soil NO peak in North
1237 equatorial Africa and the interannual variability of soil NO_x fluxes over the Great Plains in the
1238 US.

1239




Figure 6: Spatial distribution of NO_x emissions for the year 2000 from the main sources: Fossil & biofuel combustion, biomass burning and soils and main region. Data are the *a posteriori* data (top down and bottom-up NO_x emission inventory from Jaeglé et al. (2005)). (Jaeglé's data were grouped into Asia = East Asia, Mid Asia, SE Asia/India, Japan in Jaeglé et al.; Africa = S and N equatorial Africa and S Africa).

4.3.1.2 N₂O

N₂O: Nitrous oxide, is a long-lived (114yr) greenhouse gas, contributing to 10% of the global radiative forcing (Denman et al., 2007), and in the stratosphere is now the main cause of stratospheric O₃ depletion (Ravishankara et al., 2009). Microbial denitrification and nitrification processes are responsible for 87% of the annual global N₂O budget (18.8 Tg N yr⁻¹; Syakila and Kroeze, 2011), with contributions from natural soils (35%), agriculture (27%) and oceans (25%). Non biological sources are responsible for the remaining 13% through fossil fuel combustion, biofuel and biomass burning and industrial processes. Atmospheric N₂O concentrations have been rising since the industrial revolution from 270 ppb to over 319 ppb. It has always been assumed that increased N fertilizer use is responsible for this rise. Recent measurements of isotopic N₂O composition (^{14/15}N) in the atmosphere are consistent with this assumption (Park et al., 2012), and N fertilized agricultural soils are responsible for almost 16% of global annual N₂O emissions. All agricultural activities are responsible for two-thirds of the total anthropogenic N₂O emissions (Davidson and Kanter, 2014), and more than one-third is associated with animal production.

Natural soil emissions are the largest single global source of N₂O, with largest emissions from the warm wet regions in the Amazon, South-East Asia and Africa. Unfortunately, these are

Formatted: Font: Italic

also the regions for which data coverage is poor [relative to Europe and North America](#). Using [Based-on](#) an artificial neural network approach and ~~the few~~ available field observations Zhuang et al. (2012) calculated that 30% of the total natural soil contribution was from tropical evergreen broadleaved forests followed by 17% for woody savannas. Their total estimate for global soil emissions was 3.4, ranging from 2.0 – 4.6 Tg N yr⁻¹ for the year 2000. This is lower than the range 6-7 Tg N yr⁻¹ used by Syakila and Kroeze (2011), but with the very large uncertainties these values are probably not significantly different. ~~from each other.~~

[4.3.2 Soil processes responsible for NO and N₂O emissions](#)

~~*[4.3.2 Soil processes responsible for NO and N₂O emissions](#)*~~

[4.3.2.1 Denitrification](#)

[Denitrification](#): — Denitrification is the major N loss pathway for N_r (Fig. 7). The ratio of the denitrification products N₂O and N₂ depends on localised environmental conditions in the soil. This microbial process, performed by archaea, bacteria or fungi, using oxidised nitrogen compounds such as nitrate or nitrite as an alternative electron acceptor in the absence of oxygen (Butterbach-Bahl et al., 2013), removes approximately 30-40% of N_r inputs to watersheds (Seitzinger et al., 2006).

Microbial denitrification is a highly spatially distributed process occurring in soils, sediments or water bodies if environmental conditions become unfavourable for aerobic degradation of organic matter. It mostly occurs at aerobic/anaerobic interfaces in soils, e.g. in riparian zones where lateral water flow from upstream regions provides a steady influx of nitrogen oxides as well as dissolved organic C to the waterlogged and oxygen (O₂) depleted soil zones in such topographic depressions. Thus, riparian areas are hotspots of denitrification (Pinay et al., 2007) as well as often hotspots of soil N₂O emissions (Jungkunst et al., 2008; Butterbach-Bahl and Dannenmann, 2011). Denitrification is a heterotrophic process in which nitrate (NO₃⁻) is used as a terminal electron acceptor during the oxidation of C substrates (Groffman et al., 2006). Thus, at least three pre-conditions need to be fulfilled: a) oxygen depletion, b) availability of nitrogen oxides and c) availability of easily degradable C substrates. ~~Given these pre-conditions and the presence of an active denitrifying microbial community, thereby noting that many microbes are not only capable to breakdown organic matter under aerobic conditions, but can activate the denitrification enzyme chain under anaerobiosis, and the removal of nitrates by denitrification is a highly effective process. In wastewater Ddenitrification is the key process in wastewater treatment plants to removing NO₃⁻, but it is also a major loss pathway of N fertilizers in agriculture provided to arable fields to increase crop production.~~ Loss rates of N₂O from fertilized cropland due to denitrification have been reported to be up to 240 kg N ha⁻¹ yr⁻¹ (Barton et al., 1999), thus, potentially even exceeding fertilizer application ~~ation~~ rates.

Denitrification is activated if soils become water-saturated or water-logged, e.g. due to heavy rainfall or irrigation. The sudden increase in soil moisture, blocking macro- and micropores with soil water, decreases O₂ diffusion into soil by approximately a factor of four. Since microbial metabolism as well as plant root respiration continues, the soil becomes anaerobic. Thus, besides being spatially distributed, with certain hotspots such as riparian areas in

Formatted: Font: Italic

Formatted: Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

the landscape, denitrification is also temporally highly discontinuous and is a so-called “hot-moment” phenomenon (Groffman et al., 2006, 2009).

4.3.2.2 Nitrification

Nitrification: –Biological and abiotic processes in soils are responsible for the production and consumption of NO and N₂O. Principal microbial processes leading to NO/N₂O production are nitrification and denitrification, and nitrifier denitrification (Fig. 7). There may be other theoretically feasible processes, which have not yet been identified in soils (Medinets et al., 2015; Schreiber et al., 2012). Chemodenitrification, the chemical decomposition of NO₂, is an important source of NO in acid soils and soils rich in humic acids (Stevenson et al., 1970; Zumft, 1997). Reduction of nitrite (NO₂⁻) to NH₄⁺ is also known to be a source of N₂O in some reduced environments (Fig. 7; Baggs, 2008)

Nitrification is the microbial oxidation of NH₄⁺ to nitrate (NO₃⁻), with hydroxylamine and NO₂⁻ as essential intermediates. This process occurs in all soils and aqueous systems and involves a wide range of microorganisms. ~~Whereas in~~ In the denitrification pathway NO and N₂O are obligate intermediates; ~~whereas in nitrification~~ they are by-products of nitrification and thought to operate when conditions are suboptimal for further oxidation to NO₃⁻ (Conrad et al., 1996; Baggs, 2008). Both processes take place in the same soil microsites, but even with modern technologies such as isotopic labelling, use of microelectrodes and molecular analysis, it is difficult to unravel the detailed biological pathways responsible for NO and N₂O production under different conditions (Schreiber et al., 2012). Generally, NO emissions are considered to be associated with nitrification conditions and N₂O emissions with denitrification conditions.

Formatted: Font: Italic




Figure 7: A schematic diagram of the microbial processes contributing to N_2O production (adapted from Baggs, 2008).

4.3.3 Effects of climate change on NO and N_2O emissions

How will NO and N_2O emission rates change as a result of combined changes in land use, climate and atmospheric composition? This is a rather complex question, illustrated by Fig. 8, in which the very different processes which combine to regulate these microbial transformations are overlain by larger scale processes and changes. With this complexity the answer will be different for different landscapes. To predict the potential for nitrification and denitrification and emissions of the pollutants NO and N_2O , effects of environmental change need to be established at local scales as they will depend on local effects of climate, land use and atmospheric composition changes:

(a) *Climate change:* This refers to the change of the primary environmental drivers temperature and rainfall (amount, frequency, seasonal distribution), both affecting soil environmental conditions but also site and landscape hydrology, vegetation cover and substrate supply. Finally, this will also affect land use, since it can be expected that farmers will adapt land use and land management as climate changes (Kicklighter et al., 2014).

~~(b) Land use change: This refers to changes in vegetation cover, land use and management resulting in changes in substrate supply to the soil microbial community, but also triggers changes in soil and catchment hydrology. Changes in land use can be driven by climate change (Kicklighter et al., 2014), but economic factors, such as e.g. the recent trend to bioenergy production (e.g. Leemans et al., 1996), might be of much greater importance on time scales of years to decades. Policy and economic drivers also influences the uptake of measures aimed at promoting mitigation in the agricultural sector (MacLeod et al., 2010).~~

~~(c) Atmospheric composition change: this mainly links to rising CO₂ concentrations, resulting in reductions in plant transpiration and increasing levels of soil moisture (e.g. Long et al., 2004), but also to changes in regional O₃ concentrations affecting plant performance and, thus, e.g. plant litter production or transpiration or atmospheric deposition of reactive nitrogen (Sutton et al., 2011), which is not only an additional Nr source for soil microbial processes but also drives forest C sequestration and changes in soil C and N stocks (De Vries et al., 2014), and by this affects soil environmental conditions.~~

~~4.3.3.1 Climate change effects on NO and N₂O emissions and the importance of changes in regional hydrology~~

~~Climate change: Without doubt mMicrobial activity will increase with temperature if the process itself is not limited by the availability and supply of the respective substrate(s), such as easily degradable C and oxidised N compounds. If global and regional temperatures continue to increase, there is a potential for denitrification and nitrification rates also to increase. The study by Luo et al. (2014), for example, shows that in grassland soils undergoing experimental warming of 2°C over a period of 10 years, key metabolic pathways related to C and N turnover accelerated. In the case of denitrification, this increase was 12%. However, if summer temperatures increase whilst summer rainfall decreases, denitrification rates would decrease substantially, since the most important environmental precondition for denitrification, anaerobic conditions are not provided. These conditions would however be favourable forte NO and N₂O emissions via nitrification.~~

An additional consequence of rising temperature will be increased rates of transpiration and evaporation (Long et al., 2004). In conclusion, it can be assumed, that changes in soil moisture as driven by changes in rainfall patterns and amounts and evapotranspiration fluxes will very likely dominate the overall response and overwhelm any direct temperature effects on denitrification, nitrification and NO and N₂O emissions. Moreover, expected changes in the hydrological cycle at regional to continental scales will affect not only the seasonality of soil moisture changes, catchment and watershed hydrology, and the size and temporal expansion and shrinking of riparian zones (Pinay et al., 2007). Thus, when considering climate change effects on NO and N₂O emissions one must include changes in rainfall (amount, frequency and seasonality), evapotranspiration and associated changes in surface and subsurface water flows and catchment/watershed hydrology in the focus (Butterbach-Bahl and Dannenmann, 2011).

Formatted: Font: Italic




Figure 8: A summary of the processes influencing the responses of NO and N₂O emissions to changes in climate and land use.

4.3.3.2 Land use and land management effects on denitrification.

Land use: Expected changes in climate are already triggering changes in land use and land management. The area of irrigated agricultural land is expanding quickly not only in semi-arid but also in humid temperate climates to adapt agriculture to predicted temporal water scarcity due to climate change (Trost et al., 2013). Reviewing the existing literature on irrigation effects on soil N₂O emissions Trost et al. (2013) found that in most cases irrigation increased N₂O emissions in a range from ± 50 to $\pm 150\%$, which is very likely caused by increased denitrification activity in such soils. Irrigation may increase (Liu et al., 2010) or reduce NO emissions (Abalos et al., 2013), depending on the wetness of the soil.

The large scale introduction of no-till agriculture, especially in Latin America (Abdalla et al., 2013), may affect N₂O and NO emissions. A study by Rosa et al. (2014) which addresses denitrification activity in no-till production systems in the Argentinian pampas, suggests that increased soil aggregate stability in no-till systems, and its effects on C sequestration, water infiltration, soil aeration and microbial habitat provision, is the most important factor for explaining changes in denitrification activity, rather than by changes of the microbial community (Attard et al., 2011). For a cereal field in Scotland, UK no-till increased N₂O emissions, but decreased NO emissions, whereas tillage had the opposite effect and increased NO but decreased N₂O emissions (Skiba et al., 2002).

Formatted: Font: Italic

4.3.3.3 Atmospheric composition change and denitrification

Atmospheric composition change—specifically the main component is the increasing concentration of atmospheric CO₂, and in some regions still increasing levels of tropospheric O₃, and atmospheric deposition of N_r to terrestrial ecosystems, influences ecosystem nutrient status and productivity. Increasing levels of atmospheric CO₂ increases water use efficiency of plant photosynthesis, resulting in increased soil moisture levels and hence increased N₂O emissions by denitrification or nitrification, (e.g. Kammann et al., 2008), but probably reduced NO emissions. Also rhizodeposition of easily degradable C compounds has been shown to increase (Singh et al., 2010) as a result of additional inputs of N_r to soil by atmospheric deposition, i.e. the other denitrification substrate besides labile organic C compounds, the overall effect of atmospheric composition change on denitrification should be to increase denitrification.

To predict quantitatively how climate change will influence terrestrial denitrification and nitrification rates and associated NO and N₂O emissions it is necessary to know both the quantities of N_r used by agriculture and the effects of climate on the soil processing. The balance of evidence suggests a net increase of NO and N₂O emissions due to the increases in N_r use to need to feed a growing population and increased demand for biofuels. For N₂O IPCC (2013) climate simulations, using a new set of scenarios (representative concentration scenarios RCP2.6, RCP4.5, RCP6 and RCP8.5), suggests an average increase of N₂O by 1.6 Tg N₂O-N (range -1.4 to 4.5 Tg N₂O-N) between 2010 and 2050. A similar increase in the remaining half of the 21st century would lead to an increase in emissions by 28% over the century to 3.2 Tg N yr⁻¹ in 2100. An increase in soil NO emissions during the 21st century of similar magnitude to those for N₂O seem likely, as emissions of both gases are primarily driven by agricultural and biofuel production. This would lead to soil emissions of NO in 2100 of 11.5 Tg N yr⁻¹.


It is clear that predicted changes in rainfall and regional hydrological cycles are more important than direct effects of temperature for large scale denitrification activity. Increases in precipitation at higher latitudes appear common to many climate model projections for the later decades of this century (IPCC, 2013), but the variability in magnitude and distribution precludes clear regional quantification. Likewise the drying of the Mediterranean basin is a common feature in some climate model simulations. Such a response would decrease N₂O emissions, but could increase NO emissions. However if soils are irrigated N₂O emissions may increase and NO emissions decrease. These expected changes are overlaid by changes in land use and land management, which are also partly triggered by climate change. Moreover, changes in atmospheric composition are indirectly feeding back on denitrification activity in soils too, e.g. by affecting plant performance and thus, nutrient and water flows. To better understand climate change effects on regional and global denitrification and nitrification activities multi-factorial climate (e.g. Mikkelsen et al., 2008) and land use/ land management change experiments are needed. Such studies have still only been run for relatively short term, which hampers the detection of interactive and nonlinear effects, or the identification of thresholds and tipping points (Luo et al., 2011). Study data from multifactorial experiments are needed to test and improve process descriptions in biogeochemical and landscape models.

Formatted: Font: Italic

which are finally the tools for assessing denitrification and nitrification rates including NO and N₂O emissions at landscape to continental and global levels. Ultimately only such models, linked to hydrological, climate and dynamic landuse models, might allow us to get a better idea of how NO and N₂O emissions may change under future environmental and land management conditions, even though modelling interactions at various temporal and spatial scales remains a major future challenge.

5 ATMOSPHERIC PROCESSING - CHEMISTRY

Higher temperatures increase the rates of almost all chemical conversions: the higher kinetic energies associated with warmer temperatures means reactions proceed faster. Temperature has particularly important effects on two equilibria involving reactive nitrogen (Cox and Roffey, 1977; Feick and Hainer, 1954):



Higher temperatures push both these equilibria towards the right, i.e. resulting in thermal decomposition of gaseous peroxyacetyl nitrate (PAN: CH₃COO₂NO₂) and ammonium nitrate (NH₄NO₃) aerosol particles, effectively reducing the atmospheric lifetimes of these two species. The impacts of 21st century climate change on global atmospheric composition, via reaction (Eq. (6)), have been investigated by Doherty et al. (2013). For a temperature increase of +3K (typical for 2100 relative to present-day), the PAN lifetime in the troposphere approximately halves (from 4 to 2.5 hours at mean surface temperatures of 290 K; and from 6 to 3 months at mid- to upper-tropospheric temperatures of 250 K). PAN is the main cAs PAN is a major component of tropospheric NO_y, so climate change may significantly reduce the size of the NO_y reservoir, reducing the long-range (or intercontinental) transport of NO_y (Doherty et al., 2013).

Liao et al. (2006) find that climate change effects (specifically the SRES A1B scenario from 2000 to 2050) leads to reduced concentrations of NH₄⁺ aerosols over East Asia, and attribute this to temperature increases acting via Eq. (7). Similar results were found over the US (Pye et al., 2009).

Changes in the stratospheric source of HNO₃ are also likely as a consequence of a changing climate. Much like the predicted increase in tropospheric O₃ from enhanced stratosphere-troposphere exchange (STE), driven by a more intense Brewer-Dobson Circulation, the stratospheric O₃ enters the troposphere with some NO_y as HNO₃. This is a small source currently estimated to be ~1 Tg N yr⁻¹, but STE is projected to increase by 50-100% over the 21st century, so this NO_y source may ~double. Stratospheric NO_y and O₃ may show different trends, so it may be more complicated than just knowing the STE air mass flux (most models just add NO_y with a fixed ratio to O₃).

5.1 Lightning-climate interactions

Lightning NO is an important natural source of tropospheric NO_x, especially the tropical upper troposphere (Schumann and Huntrieser, 2007). Nitric oxide (NO) is formed following the dissociation of molecular oxygen and N by the lightning discharge in air. Atmospheric composition is modified as described in the companion paper by Monks et al. (2014).

The effects of climate change on lightning and NO_x production have been investigated by Toumi et al (1996), and by Reeve and Toumi, (1999) suggesting increases in both lightning and NO_x production. The estimates of increased NO_x production in a warmer climate are rather variable and range from 4 to 60% per degree K (surface temperature change), Schumann and Huntrieser, 2007; and Williams, 2005).— More recent analyses by Romps et al. (2014) based on the eConvective #Available #Potential Eenergy (CAPE) and precipitation rate indicate values of 12± 5% °K⁻¹.

Taking a value towards the lower end of the range of reported temperature responses, of 10% K⁻¹ and a temperature change of 4°K by 2100, yields an increase in lightning NO_x production from 5 to 7 Tg N yr⁻¹.

6 ———ORGANIC NITROGEN

Gaseous organic nitrogen exists in the atmosphere in both oxidized (peroxy acetyl nitrate PAN, and related compounds) and reduced forms, including amines, amino acids and urea. Organic nitrogen also occurs in particulate matter and in rain and snow and from the methodology for chemical analysis it is not always clear in whether the organic nitrogen was in reduced or oxidized form. -The oxidized gaseous forms and PAN especially, are important as a reservoir of ozone precursors for photochemical oxidant formation (Singh and Hanst, 1981), while interest in the gaseous reduced forms and ON in aerosols and precipitation is primarily through their contribution to nitrogen deposition (Jickells et al., 2013).

~~Atmospheric organic nitrogen (ON) has received increasing attention in the last two decades, nevertheless its characterisation and understanding are far from complete and it represents by far the least understood component of the atmospheric N cycle.~~

Recently, Cape et al. (2011), Cornell et al. (2011) and Jickells et al. (2013) reviewed Organic nitrogen in the atmosphere. Organic nitrogen (ON) and it has been reviewed recently by Cape et al. (2011), Cornell et al. (2011) and Jickells et al. (2013). These reviews cover available -measurements, atmospheric cycling, and providing also a technical discussion on analytical methods, their comparability and statistical caveats for data treatment. Here we-In this section provide a brief and-a general description of atmospheric ON, its sources, relevant properties and relations to the N cycle, -with-a-and the likely effects of changes this century are described. Gaseous oxidized organic nitrogen compounds are treated separately, as the literature, and motivation are distinct.

~~viewpoint coherent to the aims of this review.~~

6.1 Atmospheric relevance

Organic nitrogen is a ubiquitous component of the atmosphere, mainly found in aerosols and precipitation, although present also in the gas phase (Cape et al., 2011; Cornell et al., 2011; Jickells et al., 2013). Atmospheric concentration data for the aerosol fraction in literature often refer to the water-soluble fraction of ON (WSON), more frequently investigated as it is considered to be more bioavailable (Seitzinger and Sanders, 1999) and climate relevant. Aerosol WSON atmospheric concentrations range from a few to few tens of nmol N m^{-3} in a selection of remote sites by Jickells et al. (2013), reaching concentrations as high as $\sim 150 \text{ nmol N m}^{-3}$ in the Po Valley (Montero-Martinez et al., 2014) and up to $2 \text{ } \mu\text{mol N m}^{-3}$ at Qingdao (China) (Shi et al., 2011). The above aerosol concentrations determine WSON in rainwater ranging between $5 \text{ } \mu\text{mol L}^{-1}$, in remote regions, and $>100 \text{ } \mu\text{mol L}^{-1}$ as measured in China (Cornell et al., 2011).

The water-soluble fraction of ON contribution to total N in aerosol and rainwater has been investigated in a number of studies, with results ranging from a few percent to more than 40% (Cornell et al., 2011). Jickells et al. (2013) reports a collection of rainwater WSON datasets from around the world from the studies of Cornell et al. (2003; 2011) and Zhang et al. (2012), resulting in an average ON contribution of 24%. Similar contributions, ranging from 19 to 26%, have been observed for aerosols in many other studies (Zhang et al., 2002; Chen et al., 2010; Lesworth et al., 2010; Kunwar and Kawamura, 2014; Miyazaki et al., 2014; Montero-Martinez et al., 2014). Nevertheless, lower WSON/TN contributions have been also reported: 6% in Delaware, USA (Russell et al., 2003), 13% in Crete (Violaki et al., 2010), 16% in the outflow from northeast India over the Bay of Bengal (Srinivas et al., 2011) and 10% in paired urban-rural sites in Georgia (Rastogi et al., 2011). Higher contributions seem, instead, typical of China (Shi et al., 2011; Zhang et al., 2012), likely due to the use of organic manures and urea as fertilizers in agriculture (Jickells et al., 2013).

It is worth highlighting that in one of the few studies in which total ON (and not WSON) was measured, the ON/TN ratio was of the order of 70% (western North Pacific in summer) (Miyazaki et al., 2010), suggesting an important contribution from water-insoluble ON. Russell et al. (2003) also showed an important fraction of aerosol ON in water-insoluble form. Other investigators (Li and Yu, 2004; Duan et al., 2009) which measured the total ON have not confirmed such a result. Finally, the model approach of Kanakidou et al. (2012) estimated ON as 26% of TN deposition, globally.

These numbers provide an insight into the importance of atmospheric ON in the N cycle, even though a full understanding is far from being achieved. In particular, ON can be considered important in the long-range transport of N (e.g., Singh and Hanst, 1981; Gorzelska and Galloway, 1990; Neff et al., 2002; Matsumoto and Uematsu, 2005) because its removal processes tend to be less effective than those for nitrate and ammonium, which are generally deposited closer to their sources (Cornell et al., 2011).

At least a fraction of ON is known to be bioavailable (Timperley et al., 1985; Peierls and Paerl, 1997; Seitzinger and Sanders, 1999) therefore its deposition can provide nutrients for land and marine ecosystems. Nevertheless, the effects of ON on the surface ocean are unclear due to the large uncertainty in the sources and magnitudes of deposition. Even less is known about the potential human and ecosystem toxicity of ON (Paumen et al., 2009).

Recently, atmospheric ON has received attention because of its light-absorbing properties (Desyaterik et al., 2013). Reactions leading to the formation of ON compounds in aerosol particles or evaporating droplets have been indicated as potentially important for the formation of atmospheric brown carbon (Nozriere et al., 2007; 2009; Shapiro et al., 2009; Nguyen et al., 2012; Powelson et al., 2014; Lee et al., 2013).

6.2 Chemical composition

Atmospheric ON is a sub-set of the organic carbon, and in analogy with the latter, is a complex mixture of compounds with different properties and origin (e.g., Saxena and Hildemann, 1996; Jacobson et al., 2000; Neff et al., 2002). Complementary to the total ON (or WSON) determination approach, many studies have focused on measuring the concentration of individual N compounds or groups of compounds in air, aerosols or rainwater. Given the difficulties of measuring total ON, this approach is the usual course in the gas phase.

Although this approach will never account for the whole ON, it can be useful in providing insights to sources and to clarify the contribution of single species to ON. Compounds analysed in individual studies include amines, amino acids, urea, nitrophenols, alkyl amides, N-heterocyclic alkaloids and organic nitrates (Cape et al., 2011; Jickells et al., 2013), none of which resulted dominating the ON composition. This suggests that a large fraction of ON is associated with high molecular weight polymers, constituting the humic-like materials (HULIS) (Chen et al., 2010).

This approach has shown that in certain environments and conditions, some compounds make up a consistent fraction of atmospheric ON. For instance, amino acids have been reported to account for up to 50% in Tasmania (Mace et al., 2003a), while Facchini et al. (2008) reports dimethyl- and diethyl-amines contributing 35% of aerosol ON over the Eastern North-Atlantic Ocean. On the contrary, in other studies these are only minor components (e.g., Mace et al., 2003b; Mace et al., 2003c; Müller et al., 2009; Violaki et al., 2010). Urea was also shown as an important contributor (>20%) by Cornell et al. (2001) and Mace et al. (2003a) in Hawaii and Tasmania, but was reported as a minor ON component in other sites (Mace et al., 2003b; Mace et al., 2003c). Recently, Zhang et al. (2012) showed that urea represents more than 40% of rainwater WSON in China where urea is widely used as a fertilizer.

Recently, ultrahigh resolution mass spectrometry has provided new insights into ON chemical composition in aerosol and rainwater. N-containing molecules have been reported, for instance, by Rincon et al., 2012; Cottrell et al. (2013); O'Brien et al. (2013); Zhao et al. (2013); and Kourtchev et al. (2014), accounting for 40 to more than 50% of the total identified molecules in their samples, for a total of thousands of compounds. These studies suggest that ON is made of both oxidised (organonitrates, nitroxy-organosulfates) and reduced (amines, imines,

imidazoles) N species. Altieri et al. (2009; 2012) found similar results, with more than two-thirds of the detected ON compounds containing reduced nitrogen. Moreover, they observed significant chemical composition differences between marine and continental samples, concluding that, although the concentrations and percent contribution of WSON to total N is fairly consistent across diverse geographic regions, the chemical composition of WSON varies strongly as a function of source region and atmospheric environment. LeClair et al. (2012) reported that approximately 63% of the CHNO and 33% of CHNOS compounds observed in Fresno radiation fog samples exhibited a loss of HNO₃, suggesting that besides organonitrates, there might be other N containing functional groups present, such as amines, imines, and nitro groups.

These techniques detect ON compounds in a wide range of molecular weights with carbon number between 2 and 35 (Zhao et al., 2013; Cottrell et al., 2013). Nevertheless, Chen et al. (2010) has demonstrated that N containing molecules can have masses greater than 1 kDa.

1616

6.3 Organic nitrogen sources

The complexity of ON chemical composition is reflected by its sources. ON source attribution was tentatively achieved in different studies based on size distribution, correlation with source tracers, multivariate analysis and isotopic ratios.


Many investigators report a significant spatial or temporal correlation between ON and inorganic N in aerosol and rainwater samples, with ON constituting roughly a quarter of total N in many environments (Cape et al., 2004; Zhang et al., 2012). Considering that inorganic N emissions are globally dominated by anthropogenic sources, ON has likely an important anthropogenic component (Zhang et al., 2012; Jickells et al., 2013). Many papers highlight important anthropogenic ON sources (e.g., Cornell et al., 2001; Mace et al., 2003c; Chen et al., 2010; Iinuma et al., 2010; Rastogi et al., 2011; Zamora et al., 2011; Zhang et al., 2012; Kourtev et al., 2014; Violaki et al., 2010; Cape et al., 2004; Bencs et al., 2009). More in detail, ON seems to present higher correlation with NH₄⁺ than NO₃⁻ (Zhang et al., 2012; Cape et al., 2004; Srinivas et al., 2011). This points to an atmospheric processing of ON similar to that of reduced nitrogen, or to similar sources (Jickells et al., 2013). Indeed, several papers evidence ON formation processes via interactions between organic matter and reduced N (ammonia, amine compounds or HNCO and related gases) like Nguyen et al. (2012), Lee et al. (2013) and O'Brien et al. (2013). Nevertheless, also ON formation through NO_x or NO₃ radical chemistry is reported in the literature (e.g., Zhao et al., 2013; Fry et al., 2013).

Notwithstanding the very likely global dominance of anthropogenic sources, natural sources of ON have been reported both in the marine (Spokes et al., 2000; Cornell et al., 2011; Mace et al., 2003a; Facchini et al., 2008; Müller et al., 2009; Miyazaki et al. 2010; Kunwar et al., 2014) and continental (Miyazaki et al., 2014; Kieloaho et al., 2014; Laitinen et al., 2014) environments.

As for formation processes, primary ON sources have been reported associated with soil suspension (Chen et al., 2010), sea spray (Miyazaki et al., 2010; Barbaro et al., 2011; Scalabrin et al., 2012) and biomass burning (Desyaterik et al., 2013; Zamora et al., 2011; Mace et al., 2003b; Srinivas et al., 2011; Violaki et al., 2011). Evidence of the importance of secondary ON formation has been presented for a range of environments, involving a number of precursors (De Haan et al., 2011; Nguyen et al., 2012; Rincon et al., 2012; Fry et al., 2013; Lee et al., 2013; Miyazaki et al., 2014; O'Brien et al., 2013; Zhao et al., 2013; Kourtchev et al., 2014). Furthermore, Organic N, amines in particular, is known to play a role in atmospheric new particle formation (Murphy et al., 2007; Kurten et al., 2008; Smith et al., 2010; Kirkby et al., 2011).

Concluding, Jickells et al. (2013) summarises atmospheric ON sources in: 1) Soil dust, including in this source both ON associated with soil organic matter itself and the adsorption of ON onto dust particles; 2) Biomass burning; 3) Marine emission both direct and via emissions of gaseous precursors; 4) Anthropogenic and agricultural sources.

1655



1656

Figure 89: Processes where increased temperatures would be expected to increase atmospheric ON (orange) or decrease ON (green); no colour code implies uncertain effects. Adapted from Jickells et al. (2013).

1659

1660 6.4 Effects of future climate change on ON

The current uncertainty about ON and its sources makes very difficult to estimate how the atmospheric ON contribution to the N cycle will evolve in a climate change scenario. Zamora et al. (2011) predicted that if N pollution emissions aerosols continue to increase as predicted, inorganic N will continue to be the most important source of soluble N to the subtropical North Atlantic, based on observations made at Miami and Barbados. The current interest in atmospheric ON raises hopes about our future understanding, but certainly further efforts are needed in order to improve our predictive capability.

Although predicting the effects of future changes in climate on ON is very uncertain, given the current state of knowledge of the relative contributions of the different sources of ON in the atmosphere, some general points can be made:

1. Observed spatial and temporal patterns of ON concentrations and deposition correlate positively with those of NO_y and NH_x , it seems likely therefore that the projected increases in inorganic N_r will be associated with increases in ON.

2. For material derived from resuspension of soils, periods of prolonged drought and/or increased wind speeds would lead to greater amounts of airborne material than at present;

3. For material derived from biomass burning, future patterns of biomass burning (whether natural, from increased drought, or man-made, from changes in land use) will affect ON emissions;

4. Increases in average temperatures would be expected to lead to increased atmospheric concentrations of volatile and semi-volatile organic compounds, including ON species, and in particular amines and urea, both of which are related to agricultural practices. Increased emissions of volatile organic compounds (VOCs) would provide greater substrate concentrations for reactions which form atmospheric ON. Increased sea-surface temperatures would also lead to increased volatilisation of ON from the sea surface layer.

5. Increases in the oxidised nitrogen (NO_y) content of the atmosphere would lead to faster reaction and conversion of organic matter into N-containing material, probably in the aerosol phase, leading in turn to increased aerosol concentrations of ON. Similarly, increased oxidising capacity would lead to faster conversion of hydrocarbons into oxidised organic matter which would be expected to act as a substrate for subsequent reaction with both reduced and oxidised forms of N.

6. Changes in agricultural practice, for whatever reason, could lead to large changes in ON emissions, e.g. changes in the use of urea as a fertilizer, or changes in the management of animal wastes.

The evidence, while largely qualitative suggests increases in the absolute quantity of ON in the atmosphere due to changes in climate and the amount of N_r fixed by natural and anthropogenic activity. The ON processes which are sensitive to changes in climate and land use are summarised in Fig. 89. However, the knowledge of atmospheric processing and lifetimes of the chemical components preclude quantitative estimates of the changes this century. Many

Formatted: Subscript

more measurements and improvements in understanding are necessary to progress from the current, unsatisfactory position.

6.5 Peroxyacetyl Nitrate (PAN), peroxypropionyl nitrate (PPN) and peroxyethacryloyl nitrate (MPAN).

Acyl peroxy nitrates (APNs) are produced in the boundary layer and lower free troposphere during the photochemical oxidation of volatile organic compounds (VOC) in the presence of NO_x (Cox and Roffey, 1977). The mixing ratios of these compounds are dominated by PAN with measured values commonly in the range 0.1 to 3 ppbV in Europe and North America (McFadyyn and Cape, 1999, Parrish et al., 2004), with PAN contributing 80-90% of the speciated APN measurements (Roberts, 1990). In a study of mixing ratios and fluxes of speciated APNs over a Ponderosa pine forest, PAN values were in the range 200 to 500 pptv while PPN and MPAN values were generally 20 to 50 pptv (Wolfe et al., 2009). A particular interest in APNs has been their role as an atmospheric reservoir of NO_y . APNs are subject to thermal decomposition, being stable in the cool upper troposphere and yet may decompose at lower altitudes and higher temperatures (Parrish et al., 2004). Through these processes APNs can transport NO_y over substantial distances and contribute to ozone formation remote from the NO_x sources.

The temperature sensitivity of PAN through thermal decomposition makes these compounds sensitive to changes in climate as noted in section 5, but important effects also appear at the atmosphere-surface interface over terrestrial ecosystems. Direct measurements of PAN deposition to a grassland were made by Doskey et al., (2004), who showed that thermal decomposition was a greater removal mechanism within the boundary layer than dry deposition to the grassland. Wolfe et al., (2009) reported flux measurements of APNs, above a Ponderosa pine canopy in California, showing daytime deposition fluxes of peroxyacetyl nitrate and deposition velocities peaking at $\sim 5 \text{ mm s}^{-1}$ at mid day and very small deposition velocities at night ($\sim 1 \text{ mm s}^{-1}$). Daytime deposition velocities of PPN were larger peaking at 13 mm s^{-1} , and were similar to those of PAN at night, while deposition velocities of MPAN were similar to those of PAN. In the case of PPN and MPAN the fluxes were an order of magnitude smaller than those of PAN, thus the overall flux was dominated by PAN. The dominant site of uptake within plant canopies for PAN appears to be stomata, from both laboratory and field study. However, night-time deposition values suggest that cuticular uptake contributes to the total deposition flux, and on average may be responsible for between 20 and 30% of the total deposition.

Measurements over Ponderosa pine by Farmer and Cohen (2008), suggest that in canopy chemical production of OH substantially modifies the fluxes of NO_y and VOC species, within and above the canopy. Furthermore it is clear from their work that the exchange of O_3 , NO_x and BVOC is regulated by solar radiation and depends exponentially on temperature, but predictions of the net effect of expected changes in climate have not been quantified. The effects of climate on PAN, PPN and MPAN may change the fate and lifetimes of these species in plant canopies, but since their contribution to nitrogen budgets at regional scales is small, their impact on the wider global nitrogen cycle is limited.

Formatted: Font: Bold

Formatted: Subscript

Formatted: Subscript

7 ~~EFFECTS OF IMPACTS OF~~ GLOBAL CHANGE ON ECOSYSTEM/ATMOSPHERE ~~EXCHANGE OF~~ REACTIVE NITROGEN EXCHANGE PROCESSES

A wide range of atmospheric N_r compounds (reduced N_r including gaseous NH₃, amines and aerosol NH₄⁺; and oxidised N_r including gaseous NO, NO₂, HONO, HNO₃, PAN, PPN and aerosol NO₃⁻) are emitted by, and/or dry-deposited to, the Earth's surface (vegetation, soils, water bodies, built-up areas) (Flechard et al., 2011). The sign and magnitude of their exchange fluxes are governed not only by their chemical properties, but also by meteorological, physical, chemical and biological processes. For many of these species (e.g. NH₃, HONO, NO₂) the exchange can be bi-directional (Flechard et al., 2013; Oswald et al., 2013; Neiryck et al., 2007), with emissions occurring when the surface potential exceeds the atmospheric concentration, or vice-versa.

All transfer processes between the atmosphere and the surface (vertical turbulent transport, ecosystem air column chemistry, surface/vegetation sink or source strength) are potentially affected by global change, not just through altered climate and elevated CO₂ and the knock-on effects on global vegetation and the ocean, but also (i) through changes in the mixing ratios of other pollutants such as O₃ and ~~sulfur dioxide (SO₂)~~ that affect stomatal function and/or surface chemical sinks, (ii) through changes in land use, land cover and agricultural as well as silvicultural practices, and even (iii) through the feed-back of elevated N_r deposition on ecosystem functioning.

7.1 Impacts on processes regulating surface N_r sink/source strength

7.1.1 ~~Vertical atmospheric transport~~

~~Deposition rates are not only governed by the surface properties, but further constrained by turbulence. By contrast, some emission processes are purely governed by the production process, and are thought to occur even when turbulence is low. Other emission processes are governed by the equilibrium between surface pools and the gas phase concentration and are affected by the efficiency with which emitted molecules are dispersed vertically, thus lowering the concentration at the surface and promoting further emission.~~

Compounds whose deposition rates are particularly sensitive to atmospheric turbulence include those for which vegetation is ~~thought to provide~~ a perfect sink, including nitric acid, and those contained in aerosols. Thus, surface wind speed, friction velocity, atmospheric stability and surface roughness control the rates of vertical turbulent transport of N_r trace gases and aerosols through the surface layer and within the canopy. The aerodynamic (R_a) and viscous sub-layer (R_b) resistances to dry deposition are both inversely proportional to the friction velocity (Monteith and Unsworth, 2013). A comparison over the period 1988–2010 of recent linear trends in global surface wind speeds from satellite data, from in situ data and from atmospheric

reanalyses, showed (i) a pattern of positive and negative trend bands across the North Atlantic Ocean and positive trends along the west coast of North America, and (ii) a strengthening of the Southern Ocean winds, consistent with the increasing trend in the Southern Annular Mode and with observed changes in wind stress fields (Fig. 2.38 in IPCC, 2013). The decadal trends in surface winds on land were mostly of the order of 0 to $+0.2 \text{ m s}^{-1} \text{ decade}^{-1}$, with large areas of the Southern Pacific experiencing increases of up to $+0.5 \text{ m s}^{-1} \text{ decade}^{-1}$. Future trends in surface-wind speed are unclear, but it is clear that increases of such magnitudes could enhance atmospheric dry removal rates and shorten pollutant atmospheric lifetimes.

Similarly, changes in land cover and associated surface roughness are likely to affect the atmospheric lifetime of gases and aerosol compounds alike. Large-scale deforestation, for example, would reduce the deposition rate of aerosol significantly, while changes in crop types and tree species would have more subtle, but potentially important effects. For example, Davidson et al. (1982) showed that aerosol deposition rates to different grass species could differ by a factor of 10, in response to the microstructures (e.g. hairs) of the leaves.

7.1.2 Stomatal exchange

The flux of gaseous N_r pollutants into or out of sub-stomatal cavities of vascular plants is controlled primarily by the stoma-atmosphere concentration gradient and by stomatal conductance (G_s) (Baldocchi et al., 1987). Free-air carbon dioxide enrichment (FACE) experiments have suggested that elevated CO_2 concentrations result in a substantial reduction in ecosystem-scale G_s (typically -20% to -30%) (Ainsworth and Rogers, 2007), while the projected elevations in tropospheric O_3 will also reduce G_s by typically 10% to 20% (Wittig et al., 2007). The combined impacts on G_s of elevated CO_2 and O_3 in a future climate are less clear, however, due to nonlinear interactions between plant responses to CO_2 and O_3 . For example, the CO_2 -induced reduction in G_s helps alleviate future O_3 plant damage by mitigating stomatal phytotoxic O_3 uptake (Sitch et al., 2007).

Rising temperatures will on the other hand also impact G_s through a further reduction in stomatal opening under heat waves, or conversely through an increase in G_s in colder climates and an extension of the growing season. Changes in precipitation patterns will however likely affect G_s to a larger extent than temperature if they result in more frequent droughts during the growing season. The N_r species whose dry deposition is most affected by changes in G_s is probably NO_2 , due to its low affinity for non-stomatal sinks (Flechard et al., 2011), but the effect could also be significant for water insoluble organic N compounds such as peroxyacyl nitrates (PANs).

In the specific case of NH_3 , unlike other N_r species, another major control of the stomatal flux is the stomatal compensation point (Meyer, 1973), i.e. the leaf-level NH_3 concentration that reflects the thermodynamic equilibrium with apoplastic NH_4^+ , which itself results from cellular exchange and the balance of cytoplasmic consumption and production (Farquhar et al., 1980; Massad et al., 2010b). The combined temperature-dependent solubility (Henry's law) and dissociation constants result in an effective Q_{10} of 3-4 (Sutton et al., 2013b), which would see the NH_3 compensation point approximately double with a temperature increase of 5 K. This is

the same effect that will increase emissions from agricultural point sources in a future climate (cf Sect. 4.1.1). For vegetation this effect only holds, however, if the emission potential represented by the apoplastic Γ ratio ($\Gamma = [\text{NH}_4^+]/[\text{H}^+]$) remains constant. Ecosystem modelling (e.g. Riedo et al., 2002) suggests that variations in apoplastic $[\text{NH}_4^+]$ might be expected in response to global change, e.g. with rising temperature and CO_2 affecting primary productivity and soil/plant N cycling. Apoplastic pH itself could also be affected by global change; a doubling of CO_2 (from 350 to 700 ppm) can alkalise the apoplast by 0.2 pH units (Felle and Hanstein, 2002); similarly, droughts can induce increased apoplastic alkalinity by a few tenths of a pH unit (Sharp and Davies, 2009; Wilkinson and Davies, 2008). Because nitric oxide (NO) is an important internal signaling compound that is also released in response to ozone exposure (Velikova et al., 2005; Ederli et al., 2006), increased ozone exposure in a future chemical climate might lead to elevated compensation points for NO. This NO source is currently not usually represented in exchange models.

7.1.3 Non-stomatal plant surfaces

Vegetation surfaces other than stomatal apertures (leaf cuticle, stems, bark of tree trunk and branches, also senescent leaves) are generally considered efficient sinks for NH_3 and especially HNO_3 , particularly so if these surfaces are wet from rain or dew. Soluble N_r gases will readily be taken up by surface water films, although their affinity for NH_3 is expected to decrease as pH increases beyond seven (Walker et al., 2013), or if the NH_x accumulated in surface wetness leads to a saturation effect, reducing the sink strength (Jones et al., 2007). The atmospheric SO_2 to NH_3 molar ratio, or the total acids ($2 \cdot \text{SO}_2 + \text{HNO}_3 + \text{HCl}$) to NH_3 ratio, have been used in some inferential or chemical transport models (CTM) to scale non-stomatal resistance to surface NH_3 deposition (Massad et al., 2010a; Simpson et al., 2012).

Chemical composition and size of the wetness pool are thus key to the N_r gas removal efficiency (Flechard et al., 1999). It follows that global changes affecting the frequency and intensity of rain or dew, the subsequent evaporation rate of surface water, and the relative abundances of atmospheric alkaline compounds (NH_x , amines from agriculture; base cations from sea spray and soil erosion) versus acidic species (NO_y , SO_x , HCl from traffic, household and industrial sources) are likely to affect non-stomatal sink strengths for most water-soluble N_r species. Rising atmospheric CO_2 itself will acidify rainfall and any plant or terrestrial surface wetness, as well as freshwater and the ocean. As an illustration, the pH of pure water in equilibrium with ambient CO_2 at 15°C is 5.60 for current (399.5 ppm) CO_2 concentrations; this would drop to 5.59, 5.53, 5.48 and 5.41 for the 2100 CO_2 levels, predicted in the Representative Concentration Pathway scenarios, of 420.9 ppm (RCP2.6), 538.4 ppm (RCP4.5), 669.7 ppm (RCP6.0) and 935.9 (RCP8.5), respectively (IPCC, 2013). In real solutions, buffering effects could mitigate the impact of CO_2 , but global ocean surface pH projections for 2100 do range from 8.05 (RCP2.6) to 7.75 (RCP8.5), versus 8.1 currently, which will mitigate the temperature-induced increase in sea NH_3 emissions.

Global atmospheric emission projections for NO_x and NH_3 for the year 2100 mostly range from around 15 to 75 Tg N yr^{-1} and from around 45 to 65 Tg N yr^{-1} , respectively, compared with similar current emissions levels of around 40 Tg N yr^{-1} for both; those for SO_2 emissions mostly

range from around 15 to 40 Tg S yr⁻¹ in 2100, versus around 55 Tg S yr⁻¹ currently (van Vuuren et al., 2011a, 2011b). If one defines the global emission ratio $(2 \cdot \text{SO}_2 + \text{NO}_x) / \text{NH}_3$ (on a molar basis) as a proxy for global atmospheric acidity/alkalinity, this yields a current global value of around 2.2 mol mol⁻¹, while values based on 2100 emission projections range from 0.4 to 2.4 mol mol⁻¹, with a mean value of 1.2 mol mol⁻¹, i.e. a decrease of the ratio of 45%. If, as suggested by Sutton et al. (2013b), a global temperature rise of 5 K induces an additional – and generally unaccounted for – increase of 42% in global NH₃ emissions (on top of those attributed to increased anthropogenic activities), the reduction in the ratio is 61%. For Europe, where emission reductions are likely to continue for SO₂ and NO_x, by 75-90% and by 65-70%, respectively, by the year 2050, and with more or less constant NH₃ emissions (Engardt and Langner, 2013; Simpson et al., 2014), the ratio would drop by 75% from around 2.3 to 0.6 mol mol⁻¹. The resulting drop in acidity of water films on terrestrial plant surfaces (also reflected in projected reductions in acid deposition – see e.g. Lamarque et al., 2013) is expected to reduce non-stomatal NH₃ uptake significantly, and is a direct consequence of mitigation policies likely being implemented throughout the world for SO_x and NO_x emissions, but not for NH₃ except in very few countries.

This first-order estimate in the acidity ratio ignores nonlinearities caused by a change in the lifetime of individual atmospheric pollutants in response to climate and composition change. Rising temperatures would enhance chemical reaction rates on leaf surfaces as well as in the atmosphere (e.g. SO₂ oxidation to SO₄²⁻), also affecting pH, but perhaps more significantly, a warming would favour the partitioning of dissolved species in water films (NH₃, SO₂) – and of volatile N_r-containing aerosols (e.g. NH₄NO₃, NH₄Cl) – towards the gas phase. The non-stomatal surface resistance to NH₃ deposition has been shown over grassland to be both relative humidity- and temperature- dependent, roughly doubling with every additional 5 K (Flechard et al., 2010), consistent with solubility and dissociation thermodynamics of NH₃(gas)/NH₃·H₂O/NH₄⁺.

Surface warming is thus generally expected to reduce the non-stomatal N_r sink strength, especially for NH₃, with the notable exception of frozen surfaces, over which the effect of warming could be opposite. Surface/atmosphere NH₃ flux measurements over moorland have in fact shown that at sub-zero temperatures the non-stomatal sink is much reduced, but also that the canopy resistance decreases as surface ice or snow melts (Flechard and Fowler, 1998). Warming is expected to be strongest in the mid and especially higher latitudes (IPCC, 2013), such that vast regions in temperate to boreal climates could experience much shorter winters and significantly reduced numbers of frost days, increasing the wintertime N_r sink strength. Further, because ambient NH₃ concentrations should increase globally (higher ground-based emissions, and a decreased volatile aerosol NH₄⁺/total NH_x fraction), predicting the net impact on deposition fluxes is a challenge. Similarly, a reduced aerosol NO₃⁻/total NO_y fraction, and relatively higher HNO₃ concentration, ought to favour overall greater NO_y dry deposition, since HNO₃ deposits much faster than NH₄NO₃ aerosol (Nemitz et al., 2009; Fowler et al., 2009).

7.1.4 Soil surface exchange

Soils and surface leaf litter are both sinks and sources of N_r . The expected impacts of global change on the soil-level source strength for NH_3 , NO and N_2O are described in detail elsewhere in this review (Sect. 4.1.1), ~~and~~ and, in the case of NO have been reviewed by Pilegaard (2013) and by Ludwig et al. (2001). For agricultural soils the changes are essentially controlled by ~~changes in~~ agricultural management and cropping practices (especially fertilizer inputs: form, quantity, technique and timing of application), and by changes in climate that affect soil temperature and moisture, impacting on the turnover of soil organic matter (heterotrophic respiration), fertilizer infiltration, NH_3 volatilisation and the rates of nitrification and denitrification (Butterbach-Bahl and Dannenmann, 2011; Sutton et al., 2013b; Flechard et al., 2013). On the other hand, the N_r sink strength of soils and litter surfaces is governed by the same processes – and should be similarly impacted by changes in meteorological, physical and chemical drivers – as the canopy non-stomatal sink (see above). One essential difference, though, is that soil and decaying plant material in the litter layer are much more buffered media than is leaf surface wetness, such that smaller shifts in pH may be expected in response to the same atmospheric drivers. However, soil acidification may result from increased agricultural intensification in the 21st century, from increased N deposition onto semi-natural systems, and possibly from global hydrological changes impacting on soil oxygen availability and denitrification.

7.1.5 Chemical interactions during the exchange process

Global change may also impact air column chemical processing within and just above vegetation canopies, creating vertical flux divergence and altering the N_r sink (or source) strength. Here, for N, the main chemical interactions are those between NO, O_3 and NO_2 as well as the gas / aerosol partitioning involving volatile ammonium salts, primarily NH_4NO_3 and to a much lesser extent NH_4Cl .

Increasing global tropospheric O_3 concentrations (Sitch et al., 2007) should raise the within-canopy oxidation capacity for soil-emitted NO, thereby transforming more soil NO into NO_2 which can be at least partially recaptured by the overlying canopy, thus reducing total NO_x emission (or increasing net NO_x deposition) (Duyzer et al., 2004). Near and in-canopy chemistry are driven by the sharp gradients in concentrations and meteorological drivers near the ground. Thus they represent subgrid process for typical chemical transport models, where the bottom layer in which chemistry is calculated typically averages over tens of meters. Most models apply empirical formulations of the in-canopy chemical conversion and subsequent canopy reduction of the NO emission (Yienger and Levy, 1995) that do not mechanistically respond to changes in vegetation and chemical climate. Applying a subgrid model within a chemistry-climate model to analyse the impacts of land cover and land use changes on atmospheric chemistry at the global scale by 2050, Ganzeveld et al. (2010) calculated that changes in atmosphere-biosphere fluxes of NO_x would be small, pointing to compensating effects: although global soil NO emissions were expected to increase by $\sim 1.2 \text{ Tg N yr}^{-1}$ (+9%), decreases in soil NO emissions in deforested regions in Africa and elsewhere would be offset by a larger canopy release of NO_x caused by reduced foliage NO_2 uptake. More studies of this type are needed provide a more robust basis for prediction.

Recent advances in instrumentation to measure surface/atmosphere exchange fluxes of individual aerosol chemical components with micrometeorological techniques have led to the revelation that while effective deposition rates of sulfate are of the magnitude predicted by mechanistic aerosol deposition models ($<2 \text{ mm s}^{-1}$ for short vegetation and 1 to 10 mm s^{-1} to forest), measured deposition rates of NO_3^- often reach daytime values in excess of 50 mm s^{-1} (Thomas, 2007; Wolff et al., 2007; Ryder, 2010; Wolff et al., 2011). This observation is due to the fact that some of the aerosol NH_4NO_3 that passes the measurement height dissociates into NH_3 and HNO_3 before interacting with the surface and therefore deposits at an apparent deposition rate that reflects gas-phase deposition rather than physical interaction of particles with vegetation. The volatilisation of NH_4NO_3 is driven by the depletion of NH_3 and HNO_3 near and in canopies, due to their dry deposition, coupled with an increase in temperature which typically peaks at the top of the canopy during daytime.

The impact of near-surface column chemistry on the exchange flux actually depends (i) on the gradients in drivers of disequilibrium (relative mixing ratios of N_r species; gradients in temperature and relative humidity) and (ii) on the comparative time-scales of chemical reactions and turbulent transfer to/from the surface (Nemitz et al., 2000). Global warming will shift the NH_3 - HNO_3 - NH_4NO_3 equilibrium further towards the gas phase, which will reduce the concentrations of NH_4NO_3 . However, as discussed above and in Sect. 4.1.1, NH_3 emissions are likely to increase. NO_x emissions might well decrease, but the oxidation capacity of the atmosphere that governs the conversion of NO_x to HNO_3 is more likely to increase and the change in absolute NH_4NO_3 concentrations is therefore difficult to predict accurately.


The contribution of NH_4NO_3 to European total aerosol concentration is demonstrated in Fig. 249 which summarises campaign-based measurements of submicron aerosol composition across a coordinated network. During the colder seasons in particular, NH_4NO_3 was the single largest contributor to PM_{10} in north-west Europe, often exceeding the importance of organic aerosol and sulfates. Exceptions were sites on Crete (higher temperature), in Finland (few local emissions) and at high elevations sites (long transport time, no local emissions). Even at fairly remote sites such as the Scottish EMEP Supersite ‘Auchencorth Moss’, NH_4NO_3 often accounts for the bulk of the PM_{10} aerosol mass during pollution events (Fig. 104). Thus, the effect of climate change on the evolution of NH_4NO_3 has important consequences for exceedances of PM air quality objectives and for the climate system.

The impact of climate change on the interaction between aerosol volatility and surface exchange is less closely linked to changes in absolute temperature and humidity (these govern the overall atmospheric burden), but to changes in near-surface gradients in temperature, humidity and gas-phase concentrations. Increased solar radiation and reduction in evapotranspiration as a result of decreased stomatal conductance (see above) is likely to increase sensible heat fluxes and associated temperature gradients.

For the NH_3 / HNO_3 / NH_4NO_3 and NH_3 / HCl / NH_4Cl gas-aerosol equilibria, a surface warming and a lowering of relative humidity in a future climate would favor the faster depositing gas phase (NH_3 , HNO_3) over the slower depositing NH_4^+ and NO_3^- aerosol. In addition, reduced NO_x emissions by 2100 (van Vuuren et al., 2011a) may result in lower HNO_3 concentrations


1985 and thus reduce the secondary inorganic aerosol sink for NH₃. The impact of these processes
1986 on the atmospheric lifetimes and travel distances for NH₃ and N_r in general, however, must be
1987 set against the expected (temperature-induced) increase in both non-stomatal resistance and in
1988 stomatal compensation point for NH₃, which would have opposite effects.

1989



1990

1991 **Figure 940:** Relative (non-refractory) submicron aerosol composition measured with an Aerosol Mass
1992 Spectrometer (AMS) network during three pan-European EMEP / EUCAARI campaigns. The sites are arranged
1993 from North to South.



1994

1995 **Figure 104:** Relative PM₁₀ water soluble aerosol composition during an example pollution event observed at a
1996 rural Scottish EMEP site (Auchencorth Moss).

1997

1998 7.2 Regional and global projections for nitrogen deposition

Future trends in total (wet and dry) atmospheric N_r deposition ~~have been~~ ~~can be~~ simulated on the basis of CTM runs forced by climate and emission scenarios (Lamarque et al., 2005, 2013; Engardt and Langner, 2013; Simpson et al., 2014). At the regional scale, European climate/chemistry/deposition studies suggest that with current emission projections the main driver of future N_r deposition changes is the specified future emission change (Engardt and Langner, 2013; Simpson et al., 2014). These two studies both found significant reductions in oxidised N concentrations and deposition over Europe, and much smaller changes (both increases and decreases) in reduced N deposition, with climatic changes in having only moderate impact on total deposition. These two studies also demonstrated that the lack of sulfur and oxidised N in the future atmosphere would result in a much larger fraction of NH_x being present in the form of gaseous NH_3 . Simpson et al. (2014) predicted a large increase in gaseous NH_3 deposition in most of Europe, but with large corresponding decreases in aerosol NH_4^+ . Although not the focus of their study, the change of NH_4^+ to NH_3 , while not greatly reducing the European export, would result in shorter transport distances within Europe with likely important impacts on the protection of sensitive ecosystems.

A separate recent sensitivity study has revealed that the effect of NH_4NO_3 volatilisation near and in plant canopies lowers European surface concentrations of fine NO_3^- by typically 30% at the annual average (Nemitz et al., 2014). At the same time it increases the effective NO_3^- deposition by a factor of four. While some models are now able to account for some of this effect (e.g. the EMEP model; Simpson et al., 2012), it is not included in the majority of models. However, this effect has not yet been projected into the future to quantify the impacts of changes in climate.

Hemispheric N_r deposition projections (Hedegaard et al., 2013) also ~~tend to~~ show that the impact of emission changes dominates and is in some areas (e.g. over Europe) up to an order of magnitude higher than the signal from climate change. Nonetheless, trends in total nitrogen ($NH_x + NO_y$) deposition in parts of the Arctic and at low latitudes *are* dominated by climatic impacts. At the global scale, Lamarque et al. (2013) simulated large regional increases in N_r deposition in Latin America, Africa and parts of Asia (under some of the scenarios considered) by 2100. Increases in South Asia were predicted to be especially large, and were seen in all scenarios, with 2100 values more than double those of 2000 in some scenarios. Region-averaged values under scenarios RCP2.6 and RCP8.5 were typically ~30–50% larger in 2100 than the current values in any region globally.

Most of these studies to date do not account for the full range of global change impacts on surface exchange processes. Surface exchange in most chemistry transport models (CTMs) is generally much simplified in dry deposition (downward-only) modules (Flechard et al., 2011) that cannot respond mechanistically to changes in the physical, chemical and biological drivers of stomatal, non-stomatal and soil sources or sinks. For example, multiplicative algorithms widely used to simulate G_s (Jarvis, 1976) are still the norm in these models, while photosynthesis-based approaches (e.g. Anav et al., 2012) would be needed to quantify the future impacts of rising CO_2 and O_3 on G_s and stomatal pollutant uptake (or release). Similarly, surface chemical interactions and their impact on non-stomatal sinks are not accounted for in CTMs. Indeed, Simpson et al. (2014) noted that modelling for especially NH_x components is

2041 limited by many factors, including process-uncertainties (Massad et al., 2010a; Flechard et al.,
2042 2013), problems of sub-grid heterogeneity (e.g. Loubet et al., 2001, 2009), bi-directional
2043 exchange (Wichink-Kruit et al., 2012, Bash et al., 2013), and lack of necessary and accurate
2044 input data. As one example, it may be argued that such models do not account for a likely
2045 increase in the overall (stomatal and non-stomatal) surface resistance to NH_3 deposition, some
2046 of which may be attributed to feed-backs: higher NH_3 exposure leads to more alkaline surfaces
2047 and higher plant N uptake and a higher NH_3 compensation point, with deposition a self-limiting
2048 process. Improved models which incorporate both better process descriptions and better input-
2049 data, are clearly needed to improve confidence in predictions of future N-deposition.

2050 Two further examples of impacts of climate change can be given, both through new sources
2051 (or forcing) of emissions: the possibility of new shipping routes in the Arctic regions, and
2052 temperature-induced changes in ammonia emission factors. With regard to shipping, the rapid
2053 retreat of the Arctic sea has been one of the most dramatic features of recent decades (Comiso,
2054 2012; Corbett et al., 2010). According to Corbett et al. (2010), ~~the~~ NO_x emissions from Arctic
2055 shipping in high growth scenarios will increase by a factor of ~ 4 by 2050 compared to 2004,
2056 or almost a factor of ~ 14 if high global shipping routes are diverted into Arctic areas. The
2057 impacts of these changes on the phyto-toxic ozone dose, (POD) and N-deposition have been
2058 explored on the regional scale using the EMEP MSC-W model (Simpson et al., 2012) by
2059 Tuovinen et al., (2013). As illustrated in Fig. 112, the impact of shipping emissions is
2060 concentrated along the Norwegian coast. Although the changes are not large, e.g. 50 mg
2061 $(\text{N})/\text{m}^2$, these values are comparable to base-case deposition amounts, and are likely to be
2062 important for the sensitive ecosystems in Arctic Europe. These aspects, and also the results
2063 found for POD, are discussed further in Tuovinen et al. (2013).

2064 Simpson et al. (2014) made a first estimate of the impact of such NH_3 emission increases over
2065 Europe for year 2050 simulations. They explored the impact of both 20 and 30% increases in
2066 NH_3 and calculated the exceedance of critical-levels (CL) for N. Comparison of these runs
2067 against the CL data (Fig. 123) shows that even a 30% increase in NH_3 will not bring
2068 exceedances back to 2000s levels, but such climate-induced increases cause CL exceedances
2069 that are substantially larger than those of the standard 2050 emission scenario. Policy studies
2070 in Europe and elsewhere have been unaware of this hidden potential for increases in NH_3
2071 emissions. As noted by Sutton et al. (2013b), the approaches used to calculate and report NH_3
2072 emissions for both CTM modelling and policy assessments need complete revision to cope this
2073 new paradigm.




Figure 112: Increases in total Nr deposition (mg/m²) due to increased Arctic shipping emissions, including diversion routes in 2030. Results are relative to a 2030 base-case. Calculations from EMEP MSC-W model, redrawn from Tuovinen et al. (2013)




Figure 123: Frequency distribution of exceedances of the Critical Levels for eutrophying Nitrogen in Europe (EU28+). The red line (E05-M00-BC1) represents a year 2000 base-case and the E50-M50-BC3 scenario represents year 2050 with current emission estimates. The E50X20 and E50X30 scenarios illustrate calculations with 20% and 30% extra NH₃ emission due to climate-induced evaporation. See Simpson et al., 2014 for more details.

8 THE EFFECTS OF CLIMATE AND LAND USE CHANGES ON THE WET REMOVAL OF NITROGEN COMPOUNDS FROM THE ATMOSPHERE

The removal of N compounds from the atmosphere and their deposition to land surface by precipitation is known as ‘wet deposition’. Future climate change will cause changes in annual precipitation with some areas of the world being subject to increases in precipitation and others to decreases. Kjellstrom et al. (2011) used an ensemble of 16 regional climate models to show that in the 21st century the precipitation in northern Europe will increase and in the south of Europe, [especially the Mediterranean area](#), it will decrease, with a zone in between where the change is uncertain. Changes in wet deposition of N will be [driven mainly related to future by](#) changes in precipitation. However the degree of increase or decrease in wet deposition [can be](#) [is](#) expected to be smaller than changes to precipitation. The reason for this is that the supply of particulate matter in the atmosphere which can be wet deposited is itself controlled by precipitation. Historically, dryer years have been associated with higher levels of NH_4^+ and NO_3^- particulate concentrations in air. Therefore decreases in deposition due to reduced precipitation are expected to be partially offset by higher concentrations of NH_4^+ and NO_3^- in precipitation.

Certain atmospheric chemical transformations have reaction rates which [are dependent depend](#) on meteorological [vical properties](#) including temperature, humidity and the presence of cloud water. A particular example of this is the equilibrium reaction between ammonia gas and nitric acid vapour to form ammonium nitrate aerosol. The dissociation constant of ammonium nitrate is a strong function of temperature and varies by two orders of magnitude for typical ambient conditions (Seinfeld and Pandis, 1998). Higher temperatures result in a shift towards the gas phase resulting in lower concentrations of ammonium nitrate, a pollutant which is associated with long range transport and contributes to N deposition through wet deposition. Changes in general circulation of air will also result in different patterns in the long range transport of N compounds and the areas in which N is wet deposited. Studies of the long range transport of particulate matter show that the natural inter-annual variation in circulation has a strong influence on N aerosol concentrations (Vieno et al., 2014). Kryza et al. (2012) found that inter-annual variation in annual precipitation could account for changes of 17% and wind direction variation for 14% in total annual N deposition for two European countries. Climate change may therefore lead to a re-balancing in the contributions to N deposition of long range transport and local sources as well as the relative contributions of dry and wet deposition.

An additional wet deposition mechanism by which N can be transferred to the surface is direct cloud droplet deposition. Most types of cloud form in the middle atmosphere and do not come into direct contact with the land surface. However, in mid-latitude regions the formation of orographic clouds in hill areas is a common occurrence due to the forced ascent and condensation of humid air. Such clouds are frequently short lived and the cloud droplets do not grow large enough to form into rain drops. However, where orographic cloud does come into contact with surface vegetation, the cloud water can be deposited to the surface by direct deposition driven by air turbulence ([Fowler et al., 1990](#)). The efficiency of this mechanism depends critically on vegetation type. In grassland areas cloud deposition is generally a much less efficient mechanism for N deposition than wet deposition by precipitation. However in

regions of forested hills, cloud deposition can be the dominant process for the input of nitrogen to sensitive upland ecosystems (Błaś et al., 2008). The impact of future climate change is expected to result in a shift in climatic zones and could cause the migration of forests to higher altitude areas which were previously above the tree line. The consequence of this would be large increases in inputs of nitrogen due to the effect of cloud deposition.

2133

9 EFFECTS OF CLIMATE AND LAND USE CHANGES ON C-N RESPONSES IN TERRESTRIAL ECOSYSTEMS

The close linkage between the terrestrial C and N cycles implies that perturbations of the C cycle, such as the anthropogenic increase in atmospheric CO₂ (and ensuing changes in plant production), man-made climate change (affecting the turnover rates of terrestrial C), or anthropogenic land-use change invariably have repercussions on the terrestrial N cycle (Zaehle, 2013). The level of understanding of these repercussions is generally low, owing to the lack of globally representative empirical studies and sufficiently tested global models (Zaehle and Dalmonech, 2011).

Observational evidence from ecosystem scale CO₂ manipulation experiments consistently shows that the magnitude and persistence of CO₂ fertilization strongly depend on the ability of the vegetation to increase its N acquisition (Finzi et al 2007; Palmroth et al 2006; Norby et al., 2010; Hungate et al., 2013). The sustained increase of vegetation production observed at some experimental sites was associated with increased root exudation and soil organic matter turnover, effectively redistributing N from soils to vegetation (Drake et al., 2011; Hofmockel et al., 2011). Other factors such as increases in N inputs from fixation generally played only a small role in forest ecosystems (Norby et al., 2010; Hofmockel et al., 2007). There is mixed evidence concerning the response of ecosystem N losses to elevated CO₂. The response of gaseous N losses (e.g. as N₂O) to elevated CO₂ ~~is dependent~~ depends on the response of ecosystem N turnover under elevated CO₂ and generally leads to an increase in N₂O emissions in ecosystems where N availability does not strongly limit plant growth (van Groenigen et al., 2011; Butterbach-Bahl and Dannenmenn, 2011). ~~The quantitative understanding of the above responses is limited, because of the difficulty of measuring small changes in the ecosystem's N cycle, in particular against a large spatial and interannual variation of the ecosystem N fluxes.~~

In agreement with the experimental evidence, global modelling studies generally show a strong attenuating effect of the CO₂ fertilization on plant growth and land C storage due to reduced N_r availability (Sokolov et al., 2008; Thornton et al., 2009). Future projections of N cycle models that accounted for varying terrestrial N sources and losses (Xu-Ri and Prentice, 2008; Zaehle et al., 2010a) showed a wide range of responses of the terrestrial N cycle to increasing elevated CO₂ (Fig. 13). This is due to diverging representation of important N cycle processes, in particular those controlling in- and outflows of N from the ecosystem and the coupling of the C and N stoichiometry in plants and soils (Zaehle and Dalmonech, 2011; Zaehle et al., 2014). The increase in terrestrial N by up to 11 Pg N (+10%) during the period 1860-2100 in the LPX model was mostly determined by increasing biological N fixation under elevated CO₂

(Stocker et al., 2013). Over the same time period, the response of the O-CN model was determined by an increase of the vegetation and soil C:N ratios as well as increases in terrestrial N (3 Pg N; +2.5%) due to reduced N losses (Zaehle et al., 2010b). The projections by the CN-TEM model (Sokolov et al., 2008), which assumes that the total terrestrial N store is time-invariant, suggested an increase in terrestrial C between 1860 and 2100 by ~250 Pg C simply due to a prescribed increase in vegetation C:N and redistribution of N from soils to vegetation.

In response to increasing temperature, enhanced decomposition of soil organic matter consistently increases gross and net N mineralisation (Bai et al., 2013; Rustad et al., 2001). Increased mineralisation is generally, but not always, associated with increases in nitrification, and N₂O emissions. There is ambiguous evidence as to the response of N leaching losses, which in some cases increased and in others declined (Bai et al., 2013). Observed growth responses to warming are more diverse, partly owing to difficulties in measuring plant growth and its interannual variability (Rustad et al., 2001). In N-limited ecosystems, increased N mineralisation increases N uptake of vegetation, which causes a long-term fertilization effect in N limited forests (Melillo et al., 2002, 2011). In consequence, despite likely N losses due to warming, the higher C:N ratio in woody vegetation compared to C:N ratio of soil organic matter causes increased ecosystem carbon storage due to the redistribution of N from soil to vegetation (Melillo et al., 2011).

Global models include these mechanisms and consistently suggest an attenuation of the C loss under higher temperatures due to C-N cycle interactions. However, the available climate change projections vary widely in ~~terms of~~ their global N cycle response, partly owing to differences in magnitude and regional patterns of temperature and precipitation changes (Stocker et al., 2013). In general, soil N stocks tend to decline in future projections, due to increased soil N mineralisation and increased ecosystem N losses (Fig. 123). These losses range between 5 and 10 Pg N (roughly 5-10%) between 1860 and 2100, depending on the model and scenario applied (Stocker et al 2013; Zaehle et al., 2010a). Although regionally there are increases in vegetation N associated with the redistribution of N from soils to vegetation, the models project a decline in the global vegetation N store, partly related to declining tropical forest biomass. It is worth noting that the N redistribution effect due to climate warming has important implications for the carbon-cycle - climate interaction, which is generally thought to be positive, i.e. amplifying climate change (Gregory et al., 2009). In two studies, which either assumed a closed N cycle with no losses, or had small positive carbon-cycle climate feedback, the response of vegetation growth was strong enough to turn the carbon-cycle - climate interaction into a small negative feedback (Sokolov et al., 2008; Thornton et al., 2009), whereas in another study that described C-N interactions (Zaehle et al., 2010a), the carbon-climate interaction was reduced but remained positive.

The response of the C and N cycles to land-use changes are diverse, and depend on many details of the conversion process, such that it is difficult to establish generic patterns. Converting the land-use type of an ecosystem causes a pronounced disruption of the N cycle, because typically the vegetation N (and C), and sometime fractions of the litter layer and soil organic material, are removed. This causes a phase of reduced vegetation N uptake and enhanced N losses. Forest regrowth is typically associated with an early phase of vigorous tree growth and associated


high plant N demands, leading to a conservative N cycle with high N accumulation rates compared to pastures and croplands and consequently reduced N losses (e.g., Davidson et al., 2007). Associated with the forest-to-cropland or grassland-forest conversion are typically declines in soil organic matter stocks (Guo and Gifford, 2002). However, the intricate processes of the N cycle can overrule these trends under particular conditions (Kirschbaum et al., 2008). On a decadal to century time-scale, afforestation and reforestation are therefore typically associated with reduced N₂O emission and N as well as C accumulation, whereas the inverse is true for forest to cropland conversions Davidson et al 2007; Kirschbaum et al., 2013).

Not much is known about the large-scale N cycle consequences of land use change per se, partly owing to the simplistic representation of land use and land use change in most global models (Brovkin et al., 2013). Global model simulations suggest (Fig. 123) that the changes in N storage will largely follow the trends in the C cycle (Zaehle, 2013; Stocker et al., 2013; Brovkin et al., 2013), implying that scenarios will lead to a decline in the vegetation N storage because of the removal of the above-ground vegetation (Fig. 134), and vice versa. Using scenarios in which wide-spread increases in agricultural and pasture areas occur at the expense of forests, global soil C stocks decline with land-use change. However, given that croplands are typically extensively fertilized, the C:N ratio of the soil is often lower, given the higher N content of plant matter, such that the soil retains more N after conversion. The LPX model estimates this conversion effect to be in the order of 2 PgN for the RCP2.6 and 8.5 scenarios (Stocker et al., 2013). These estimates should be treated with due caution, given that these models do not account for a lot of the detailed processes, which affect in particular the change of soil N with time, such as the age-structure and age-dependent development of forests, or the effects of cropland management besides fertilizer additions.

Associated with the projected changes in the terrestrial N and C pools (Fig. 134) are large projected changes in the future net ecosystem N and C balance. Of these fluxes, the change in terrestrial N₂O emission is likely the most climatically relevant factor. Figure 145 shows that projections of the effect of increasing atmospheric CO₂ on the N₂O emissions differ more strongly between models than alternative plausible scenarios of atmospheric CO₂. This difference reflects the large impact of alternative hypotheses about the likely changes of biological N fixation with elevated CO₂, which are large in LPX, but insignificant in the O-CN model. In O-CN, this leads to a progressively more conservative N cycle with reduced N losses, as vegetation growth and N sequestration increases due to CO₂ fertilization. Climate change consistently increases N₂O emissions from terrestrial ecosystem. However, the magnitude of this change is both dependent on the model used (with the LPX model having a higher sensitivity to climate change (Ciais et al., 2013), and the particular climate change scenario. An assessment of the effect of diverging model projections of climate change patterns for a given climate change scenario based on the LPX model revealed large uncertainty in the response of the terrestrial N₂O emissions, which is nonetheless smaller than the differences across alternative climate change scenarios (Stocker et al., 2013). Land use change per se has only little influence on the terrestrial N₂O emissions. However, the historical increase in N fertilizer use has led to a significant increase in the terrestrial N source (Zaehle et al., 2011; Stocker et al., 2013). Importantly, there is a strong interaction between the climate response of

2252 terrestrial N₂O emission and N fertilization, as the rate of N₂O production for a given addition
 2253 of fertilizer increases with climate warming (Butterbach-Bahl and Dannenmann, 2011; Stocker
 2254 et al., 2013).

2255



2256
 2257 **Figure 134:** Responses of the terrestrial vegetation and soil C and N pools to projected changes of the atmospheric
 2258 CO₂ burden, climate, and land-use between 1860 and 2100, as simulated by two global terrestrial biosphere models
 2259 (LPX, Stocker et al., 2013; and O-CN, Zaehle et al., 2010a). The LPX simulations are based on the climate change
 2260 projections of HadGEM2-ES model using atmospheric greenhouse gas and land-use forcing for the RCP2.6 and
 2261 8.5 scenarios. The O-CN simulations have been driven by climate change projections of the IPSL-CM4 model
 2262 using the atmospheric greenhouse gas forcing of the SRES-A2 scenario.




Figure 145: Change in terrestrial N₂O emissions from pre-industrial conditions to projected changes of the atmospheric CO₂ burden, climate, and land-use, as simulated by the two global terrestrial biosphere models (LPX and O-CN), as in Fig. 123. Dashed lines in the land-use change panel refer to projected N₂O emission without the change in fertilizer inputs associated with the RCP scenarios (Stocker et al., 2013).

10 DISCUSSION AND POLICY IMPLICATIONS OF THE RESPONSES OF THE NITROGEN CYCLE TO GLOBAL CHANGE

10.1 Emissions and cycling


The changes in fluxes of N within the global cycle discussed in the main sections of this paper are summarised in Fig. 156. Biological fixation of molecular nitrogen (BNF) is expected to increase during the 21st century both in the oceans (120 to 166 Tg N yr⁻¹) and terrestrial environments (128 to 170 Tg N yr⁻¹) due mainly to changes in climate. Anthropogenic emissions of NH₃ are projected to increase substantially, from 60 to 135 Tg N yr⁻¹. The increase has two components, first the effect of climate, in which higher temperature increase terrestrial emissions and second the effect of increases in N_r fixed by anthropogenic activity in part due to increased demand for food, driven by increases in both global population and changes in diet (especially global meat consumption per capita). By contrast, emissions of combustion related NO_x are projected to decline as the widespread use of control technology (catalytic converters on vehicles and SCR on industrial plant) more than compensate for increases in transport and power production.

The changes in emissions are ~~of course~~ spatially very variable, reflecting both the ~~current~~ ~~global~~ ~~current global~~ hotspots of N_r use in Europe, North American and Asia and the expected increases in -South and East Asia, Africa and South America where the largest growth in N_r use is expected.

Not all fluxes are projected to be larger at the end of the century, with smaller emissions of NO_x from anthropogenic sources and reduced emission of NH₃ from the oceans (5.7 declining

to 1.7 Tg N yr⁻¹) due to the effects of ocean acidification more than compensating the effects of higher water temperatures.

The two large cycles of N_r in terrestrial soils and in the oceans both increase substantially, 240 to 328 Tg N yr⁻¹ for soils and 230 to 290 Tg N yr⁻¹ in the oceans (Fig. 15).






Figure 156: Changes in the major fluxes and in the terrestrial, marine and atmospheric processing of reactive nitrogen (N_r) between 2010 and 2100, adapted from Fowler et al., 2013).

10.2 — Effects of changes in atmospheric composition on long range transport of N_r

The removal of sulfur from the atmospheres over Europe and North America has changed the aerosol composition in these regions, with the inorganic aerosol N_r dominated by $(\text{NH}_4)_2\text{SO}_4$ prior to 1990 and by NH_4NO_3 in more recent years, since then. Cool season episodes with high particulate matter (PM) concentrations occur widely in Europe in which NH_4NO_3 is a major contributor (Vieno et al., 2014). Likewise in Beijing, NH_4NO_3 is important in winter PM episodes, contributing on average approximately 30% of the PM_{10} mass (Sun et al., 2014). The change in aerosol composition has changed the atmospheric lifetime, deposition footprint and transport distance of much of the emitted reactive -nitrogen. Aerosols comprising $(\text{NH}_4)_2\text{SO}_4$ are largely non-volatile, once formed the aerosol stays in aerosol form until scavenged from the atmosphere by rain. By contrast, NH_4NO_3 is volatile, and close to terrestrial surfaces the deposition of the gaseous HNO_3 and NH_3 to the surface drive the evaporation of the aerosol, especially in warm daytime conditions. These effects lead to increases in the atmospheric removal rate-rate of removal of NH_4NO_3 relative to $(\text{NH}_4)_2\text{SO}_4$ and a reduction in the lifetime and travel distance of N_r with time during the last 20 years in Europe and North America as the sulfur has been removed from emissions. The trends of increasing importance of NO_3^- aerosols is projected to continue through to the end of this century, with NH_4NO_3 becoming a dominant inorganic component over many regions, despite reductions in NO_x emission due to the increased availability of NH_3 (Hauglustaine et al., 2014). Overall, the changes in atmospheric

composition have increased the importance of nitrogen compounds, as a fraction of the pollutant mixture present and in their role in generating effects on ecosystems, human health and climate. One aspect of the likely changes in the characteristics of Nr nitrogen in the environment is the likely changes in the deposition footprint of reactive nitrogen resulting from changes in climate. To date, the effects of climate change on regional patterns of deposition have been explored using regional chemistry-transport models, as described in section 7. The effect on emission –and deposition footprints at the local scale have not, so far been explored even though the principles have been established, of substantially larger emissions of NH₃ (Sutton et al., 2013b) and increased volatilization of aerosol NH₄NO₃ (Nemitz et al., 2014) close to terrestrial surfaces. These changes raise the importance of control measures for emissions of both ammonia and nitrogen oxides and the need for further modelling and field validation of the interactions between reactive nitrogen in the environment and climate.

10.3 —Costs

Emissions of N_r from farming activities to the atmosphere, soils and freshwaters are large per unit area relative to the fluxes involved in natural ecosystem BNF and are a substantial contributor to the emission and global N_r deposition hot spots, damage to ecosystems and effects on human health. The processing of N_r in soils and vegetation lead to a wide range of mobile gas and solution phase species and leaks to the wider environment. Only a small fraction (between 20% and 30%) of the N_r used in agriculture is consumed by humans in food (Sutton et al., 2013a), most is wasted either in reactive forms or transformed back to N₂. Current societal costs due to these losses of N_r to the environment are very large. Recent cost-benefit analyses of N_r have been attempted for the Chesapeake Bay in the US (Birch et al., 2011), for Europe (Brink et al., 2011) and as a broad overview for the US (Compton et al., 2011). Table 3 shows the ranges of estimated societal costs per N_r component loss and impact, based on the ‘willingness to pay’ method for EU27 (Brink et al., 2011). Based on these costs, the most important component of the N_r cycle is the emission of NO_x, due to the human health impacts of both particulate matters and ozone. Ammonia is also important, but the health effects are less certain. There is a large uncertainty for the cost of N_r enhancement of surface and groundwater. Sutton et al. (2011) Brink et al. (2011) estimated that the agricultural benefits of N_r in Europe are between €1025 and €1030 billion per year, while the total environmental costs based on the values in Table 3 is in the range €2013 and €15065 billion per year, making the point that the costs and benefits are of a similar magnitude, and are approximately half the estimated benefits.

Table 3: Societal costs of nitrogen emissions in ranges based on Brink et al. (2011). Units are euro/kg N_r.

N _r flux	Health	Ecosystem/ coastal systems	crop decline O ₃	Climate	Total
NO _x -N to air	10 - 30	2 - 10	1 - 3		13 - 43
NH ₃ -N to air	1 - 20	2 - 10			3 - 30
N _r to water	0 - 4	5 - 50			5 - 54
N ₂ O-N to air	1 - 3			1 - 15	2 - 18

Formatted: Left

10.4 Policies to reduce the impacts of N_r

The overall mass balance for nitrogen compounds is constrained by mass conservation (what goes up must come down), thus the effect of the deposition rate by itself does not change the amount of N_r deposited globally, but the transport distance of the different compounds and regional [and importantly the country](#) import/export budgets are changed by changes in chemistry and deposition of the N_r forms present. Only changes to the emissions (and to a lesser denitrification losses to N₂ during atmospheric transport) affect the total N_r amount deposited.

In Europe and the US there are examples of successful policies that led to [the reduction of reduced](#) NO_x emissions, through the Air Quality standards for O₃ and NO₂ in the US and through UN-ECE NO_x and Gothenburg protocols in Europe and large combustion plant directives [within of](#) the EU. Successful technologies include the three-way catalysts in vehicle exhausts, the Selective Catalytic Converter systems in industry and energy production. Emissions of NO_x declined by 40% in 2009 relative to 1990 in EU27 (EEA, 2012). [NH₃](#) Policies [to reduce NH₃ emission](#) have been [much](#) less successful. [in Europe and the US. For NH₃, in](#) the US there are no policies while in Europe the Gothenburg protocol (national NO_x and NH₃ emission ceilings) has led to modest (14%) reductions (EEA, 2012). There are, however, two countries that implemented substantial NH₃ abatement measures and reduced emissions by 40% in Denmark and [75](#)0% in the Netherlands. Abatement technologies included: [low](#) emission [poor](#)-housing systems, coverage of manure storage facilities and application of slurry injection technologies. Furthermore, total N inputs in agriculture were reduced by reducing N in feed and by reducing mineral fertilizer application ([Erisman et al., 2005](#)).

Formatted: Subscript

The general options of policies to reduce the cascade effect of N_r are: ([Erisman et al., 2005](#)):

1. Limit N_r production or limit import of N through animal concentrates
2. Increase N_r use efficiency
3. More evenly distribute N production over the country, over the EU or the world
4. Convert N_r to N₂ catalytically or by stimulating de-nitrification.

Substantial reductions in N_r production by fossil fuels may be achieved by use of renewable energy such as solar, wind and water technologies. The use of biomass as an alternative energy

2385 source is not an effective strategy to reduce emissions of N_r (Erisman et al. 2008). Consumer
2386 changes in diet and lifestyle present potentially effective measures to reduce emissions, but
2387 have proved difficult to implement. Reductions in consumption and therewith production of N_r
2388 intensive goods and services offer further valuable control measures that have not been used to
2389 date.

2390 ~~Linked to this option is the i~~Increasing of nitrogen use efficiency (NUE) in agricultural
2391 systems and by closing nutrient cycles on different scales represents an important guiding
2392 principle which has the capacity to deliver both reductions in emission of N_r to the environment
2393 and reduced costs in food production. Furthermore as only between 20% and 30% of the
2394 nitrogen used in agriculture is consumed by humans, the potential gains in NUE are
2395 considerable (MacLeod et al., 2010). ~~These are key options for agricultural management to~~
2396 ~~reduce N_r losses and can generate both environmental and economic benefits for farmers~~
2397 ~~(MacLeod et al., 2010).~~ However, through ~~The~~ concentration and specialisation of intensive
2398 agriculture ~~or industry~~ in certain regions, ~~with import of feed and fertilizer and export of food~~
2399 ~~products,~~ creates N_r hot-spots ~~spots,~~ such as in the Netherlands, the North China Plain, ~~and~~
2400 ~~many other regions and therefore priorities for mitigation~~ (Chen et al., 2014; Shen et al., 2013).
2401 In these regions emissions of N_r are visible not just in local measurements, but increasingly
2402 from space, using satellite remote sensing (Van Damme et al., 2014). ~~If, however,~~ these
2403 agricultural activities were distributed more evenly across the globe, and livestock production
2404 located in places where the nutrients are readily available, the N_r losses would be much
2405 reduced. ~~This is also some form of NUE increase using spatial planning as a policy.~~

2406 Finally whenever the above options do not prove effective, N_r should be converted back into
2407 N_2 by denitrification, to remove N_r from the cascade. Examples of such options include the use
2408 of wetlands and waste water treatment plants.

2409 The most effective measures that were selected based on an evaluation of successful policies
2410 in the Netherlands were:

- 2411 • Increasing nitrogen use efficiency in agriculture
- 2412 • Closing nutrient cycles at different levels
- 2413 • Influencing consumer behaviour towards reduced meat consumption
- 2414 • Using technology to reduce emissions from different compartments
- 2415 • Using spatial planning as a tool to optimise production and environmental protection

2416 In intensive agricultural areas, increasing NUE can be very effective in the short term, whereas
2417 in areas with low N inputs closing the nutrient balances is more effective of major importance.
2418 The same holds for the difference in scale: globally the closing of nutrient balances has highest
2419 priority, whereas on the local scale other priorities might be set. The main target should be to
2420 optimise food production in the world and energy consumption with the lowest possible
2421 environmental consequences.

2422 Policy instruments are needed to increase NUE. Sutton et al. (2013b) proposed increases in
2423 NUE of 20% in agricultural N_r excess areas of the world to reduce the effects of N_r on human
2424 health, climate and ecosystems. This would represent a first step to work towards a global
2425 policy of nutrient management. However, agricultural subsidies and trade restrictions differ
2426 greatly between countries and regions. Such realities distort trade and complicate the
2427 introduction of measures designed to promote environmental protection through increases
2428 nitrogen use efficiency.

2429

2430 Acknowledgements

2431 The authors gratefully acknowledge the support from the following EU projects ACCENT+,
2432 PEGASOS Legume Futures, ECLAIRE, the UK DefraEPRA, and the UK Natural Environment
2433 Research Council (NERC) for the preparation of this paper.

2434

2435 REFERENCES

2436 Abalos, D., Sanchez-Martin, L., Garcia-Torres, L., van Groenigen, J. W., and Vallejo, A.:
2437 Management of irrigation frequency and nitrogen fertilization to mitigate GHG and NO
2438 emissions from drip-fertigated crops, *Sci. Total Environ.*, 490, 880 - 888, doi:
2439 10.1016/j.scitotenv.2014.05.065, 2014.

2440 Abdalla, M., Osborne, B., Lanigan, G., Forristal, D., Williams, M., Smith, P., and Jones, M.
2441 N.: Conservation tillage systems: a review of its consequences for greenhouse gas emissions,
2442 *Soil Use Manage.*, 29, 199-209, doi: 10.1111/sum.12030, 2013.

2443 Ainsworth, E. A. and Rogers, A.: The response of photosynthesis and stomatal conductance to
2444 rising [CO₂]: mechanisms and environmental interactions, *Plant Cell Environ.*, 30, 258–270,
2445 doi: 10.1111/j.1365-3040.2007.01641.x, 2007.

2446 Altieri, K. E., Turpin, B. J., and Seitzinger, S. P.: Composition of dissolved organic nitrogen
2447 in continental precipitation investigated by ultra-high resolution FT-ICR mass spectrometry,
2448 *Environ. Sci. Technol.*, 43, 6950-6955, doi: ~~10.1021/es9007849~~10.1021/es9007849, 2009.

2449 Altieri, K. E., Hastings, M. G., Peters, A. J., and Sigman, D. M.: Molecular characterization of
2450 water soluble organic nitrogen in marine rainwater by ultra-high resolution electrospray
2451 ionization mass spectrometry, *Atmos. Chem. Phys.*, 12, 3557-3571, doi:10.5194/acp-12-3557-
2452 2012, 2012.

2453 Anav, A., Menut, L., Khvorostyanov, D., and Viovy, N.: A comparison of two canopy
2454 conductance parameterizations to quantify the interactions between surface ozone and
2455 vegetation over Europe, *J. Geophys. Res.-Biogeo.*, 117, G3, doi: 10.1029/2012JG001976,
2456 2012.

Field Code Changed

Formatted: No underline

Formatted: No underline

2457 Attard, E., Recous, S., Chabbi, A., De Berranger, C., Guillaumaud, N., Labreuche, J., Philippot,
2458 L., Schmid, B., and Le Roux, X.: Soil environmental conditions rather than denitrifier
2459 abundance and diversity drive potential denitrification after changes in land uses, *Global*
2460 *Change Biol.*, 17, 1975–1989, doi: 10.1111/j.1365-2486.2010.02340.x, 2011.

2461 Baggs, E. M.: A review of stable isotope techniques for N₂O source partitioning in soils: recent
2462 progress, remaining challenges and future considerations. *Rapid Commun. Mass Sp.*, 22, 1664–
2463 1672, doi: 10.1002/rcm.3456, 2008.

2464 Bai, E., Li, S., Xu, W., Li, W., Dai, W., and Jiang, P.: A meta-analysis of experimental warming
2465 effects on terrestrial nitrogen pools and dynamics, *New Phytol.*, 199, 441–451,
2466 doi:10.1111/nph.12252, 2013.

2467 Baldocchi, D. D., Hicks, B. B., and Camara, P.: A canopy stomatal resistance model for
2468 gaseous deposition to vegetated surfaces, *Atmos. Environ.*, 21, 91–101, doi: 10.1016/0004-
2469 6981(87)90274-5, 1987.

2470 Barbaro, E., Zangrando, R., Moret, I., Barbante, C., Cescon, P., and Gambaro, A.: Free amino
2471 acids in atmospheric particulate matter of Venice, Italy, *Atmos. Environ.*, 45, 28, 5050–5057,
2472 doi: 10.1016/j.atmosenv.2011.01.068, 2011.

2473 Barcelos e Ramos, J., Biswas, H., Schulz, K.G., LaRoche, J., Riebesell, U.: Effect of rising
2474 atmospheric carbon dioxide on the marine nitrogen fixer *Trichodesmium*, *Global Biogeochem.*
2475 *Cy.*, 21, GB2028, doi:10.1029/2006GB002898, 2007.

2476 Barton, L., McLay, C. D. A., Schipper, L.A., and Smith C. T., Annual denitrification rates in
2477 agricultural and forest soils: a review, *Aust. J. Soil Res.*, 37, 1073–1093, doi:10.1071/SR99009,
2478 1999.

2479 Bash, J. O., Cooter, E. J., Dennis, R. L., Walker, J. T., and Pleim, J. E.: Evaluation of a regional
2480 air-quality model with bidirectional NH₃ exchange coupled to an agroecosystem model,
2481 *Biogeosciences*, 10, 1635–1645, doi: 10.5194/bg-10-1635-2013, 2013.

2482 Bates, T. S., Cline, J. D., Gammon, R. H., and Kelly-Hansen, S. R.: Regional and seasonal
2483 variations in the flux of oceanic dimethylsulfide to the atmosphere. *J. Geophys. Res.*, 92, 2930–
2484 2938, doi: 10.1029/JC092iC03p02930, 1987.

2485 Bell, T. G., Johnson, M. T., Jickells, T. D., and Liss, P. S.: Ammonia/ammonium dissociation
2486 coefficient in seawater: A significant numerical correction, *Environ. Chem.*, 4, 183–186, doi:
2487 10.1071/ENO7032, 2007.

2488 Bencs, L., Krata, A., Horemans, B., Buczyńska A. J., Dirtu, A. C., Godoi, A. F. L., Godoi, R.
2489 H. M., Potgieter-Vermaak, S., and Van Grieken, R.: Atmospheric nitrogen fluxes at the Belgian
2490 coast: 2004–2006. *Atmos. Environ.*, 43, 3786–3798, doi:10.1016/j.atmosenv.2009.04.002,
2491 2009.

2492 Berman-Frank, I., Cullen, J. T., Shaked, Y., Sherrell, R. M., and Falkowski, P. G.: Iron
 2493 availability, cellular iron quotas, and nitrogen fixation in *Trichodesmium*, *Limnol. Oceanogr.*,
 2494 46, 1249–1260, doi: 10.4319/lo.2001.46.6.1249, 2001.

2495 Birch, M. B. L., Gramig, B. M., Moomaw, W. R., Doering, O. C. III, and Reeling, C. J.: Why
 2496 metrics matter: evaluating policy choices for reactive nitrogen in the Chesapeake Bay
 2497 Watershed, *Environ. Sci. Technol.*, 45, 168–174, doi: 10.1021/es101472z, 2011.

2498 Blackall, T. D., Wilson, L. J., Theobald, M. R., Milford, C., Nemitz, E., Bull, J., Bacon, P. J.,
 2499 Hamer, K. C., Wanless, S., and Sutton, M. A.: Ammonia emissions from seabird colonies,
 2500 *Geophys. Res. Lett.*, 34, L10801, doi: 10.1029/2006GL028928, 2007.

2501 Blais, M., Tremblay, J.-É., Jungblut, A. D., Gagnon, J., Martin, J., Thaler, M., and Lovejoy,
 2502 C.: Nitrogen fixation and identification of potential diazotrophs in the Canadian Arctic, *Global*
 2503 *Biogeochem. Cy.*, 26, GB3022, doi:10.1029/2011GB004096, 2012.

2504 Błaś, M., Sobik, M., and Twarowski, R.: Changes of cloud water chemical composition in the
 2505 Western Sudety Mountains, Poland, *Atmos. Res.*, 87, 224–231, doi:
 2506 10.1016/j.atmosres.2007.11.004, 2008.

2507 Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van Der Hoek, K. W., and
 2508 Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global*
 2509 *Biogeochem. Cy.*, 11, 561–587, doi:10.1029/97GB02266, 1997.

2510 Bouwman, A. F., Beusen, A. H. W., Griffioen, J., Van Groenigen, J. W., Hefting, M. M.,
 2511 Oenema, O., Van Puijenbroek, P. J. T. M., Seitzinger, S., Slomp, C. P., and Stehfest, E.: Global
 2512 trends and uncertainties in terrestrial denitrification and N₂O emissions, *Philos. T. Roy. Soc.*
 2513 *B*, 368, 1621, 20130112, doi:10.1098/rstb.2013.0112, 2013.

2514 Boyd, P. W. and Doney, S. C.: Modelling regional responses by marine pelagic ecosystems to
 2515 global climate change, *Geophys. Res. Lett.*, ~~29~~, 29, 16, 53.1–53.4, doi:
 2516 10.1029/2001GL014130, 2002.

2517 Brink, C., van Grinsven, H., Jacobsen, B. H., Rabl, A., Green, I.-M., Holland, M., Klimont, Z.,
 2518 Hicks, K., Brouwer, R., Dickens, R., Willems, J., Termansen, M., Velthof, G., Alkemade, R.,
 2519 van Oorschoot, M., and Webb, J.: Costs and benefits of nitrogen in the environment, in: *The*
 2520 *European nitrogen assessment. Sources, effects and policy perspectives*, Sutton, M. A.,
 2521 Howard, C. M., Erisman, J. W., Billen, B., Bleeker, A., Grennfelt, P., van Grinsven, H., and
 2522 Grizzetti, B. (Eds.), Cambridge, UK, Cambridge University Press, Chapter 22, pp. 513–540,
 2523 2011.

2524 Brovkin, V., Boysen, L., Arora, V. K., Boiser, J. P., Cadule, P., Chini, L., Claussen, M.,
 2525 Friedlingstein, P., Gayler, V., van den Hurk, B. J. J. M., Hurtt, G. C., Jones, C. D., Kato, E.,
 2526 De Noblet-Ducoudré, N., Pacifico, F., Pongratz, J., and Weiss, M.: Effect of anthropogenic
 2527 land-use and land-cover changes on climate and land carbon storage in CMIP5 projections for
 2528 the twenty-first century, *J. Climate*, 26, 6859–6881, doi:10.1175/JCLI-D-12-00623.1, 2013.

2529 Butterbach-Bahl, K. and Dannenmann, M.: Denitrification and associated soil N₂O emissions
2530 due to agricultural activities in a changing climate, *Curr. Opin. Environ. Sustain.*, 3, 389–395,
2531 doi:10.1016/j.cosust.2011.08.004, 2011.

2532 Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern,
2533 S.: Nitrous oxide emissions from soils: how well do we understand the processes and their
2534 controls?, *Philos. T. Roy. Soc. B*, 368, 1621, 20130122, doi: 10.1098/rstb.2013.0122, 2013.

2535 Canfield, D. E.: Models of oxic respiration, denitrification and sulfate reduction in zones of
2536 coastal upwelling, *Geochim. Cosmochim. Ac.*, 70, 23, 5753–5765,
2537 doi:10.1016/j.gca.2006.07.023, 2006.

2538 Cape, J. N., Kirika, A., Rowland, A. P., Wilson, D. R., Jickells, T. D., and Cornell, S.: Organic
2539 nitrogen in precipitation: real problem or sampling artefact?, *TheScientificWorldJOURNAL*,
2540 1, 230–237, doi:10.1100/tsw.2001.278, 2001.

2541
2542 Cape, J. N., Anderson, M., Rowland, A. P., and Wilson, D.: Organic nitrogen in precipitation
2543 across the United Kingdom, *Water Air Soil Poll.*, 4, 25–35, doi:10.1007/s11267-004-3010-8,
2544 2004.

2545
2546 Cape, J. N., Cornell, S. E., Jickells, T. D., and Nemitz, E.: Organic nitrogen in the atmosphere
2547 — where does it come from? A review of sources and methods, *Atmos. Res.*, 102, 30–48,
2548 doi:10.1016/j.atmosres.2011.07.009, 2011.

2549 Capone, D. G., Zehr, J. P., Paerl, H. W., Bergman, B., and Carpenter, E. J.: Trichodesmium, a
2550 Globally Significant Marine Cyanobacterium, *Science*, 276, 5316, 1221–1229.
2551 doi:10.1126/science.276.5316.1221, 1997.

2552 Carpenter, E. J. and Capone, D. G.: Nitrogen fixation in the marine environment, in: *Nitrogen*
2553 *in the Marine Environment*, 2nd Edition, Capone, D. G., Bronk, D. A., Mulholland, M. R., and
2554 Carpenter, E.J. (Eds.), Academic Press, San Diego, Chapter 4, 141–198, 2008.

2555 Chen, H.-Y., Chen, L.-D., Chiang, Z.-Y., Hung, C.-C., Lin, F.-J., Chou, W.-C., Gong, G.-C.,
2556 and Wen, L.-S.: Size fractionation and molecular composition of water-soluble inorganic and
2557 organic nitrogen in aerosols of a coastal environment, *J. Geophys. Res.*, 115, D22307, doi:
2558 10.1029/2010JD014157, 2010.

2559 Chen, X., Cui, Z., Fan, M., Vitousek, P., Zhao, M., Ma, W., Wang, Z., Zhang, W., Yan, X.,
2560 Yang, J., Deng, X., Gao, Q., Zhang, Q., Guo, S., Ren, J., Li, S., Ye, Y., Wang, Z., Huang, J.,
2561 Tang, Q., Sun, Y., Peng, X., Zhang, J., He, M., Zhu, Y., Xue, J., Wang, G., Wu, L., An, N.,
2562 Wu, L., Ma, L., Zhang, W., and Zhang, F.: Producing more grain with lower environmental
2563 costs, *Nature*, 514, 486–489, doi:10.1038/nature13609, 2014.

2564 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R.,
2565 Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.:
2566 Carbon and other biogeochemical cycles, in: *Climate Change 2013: The physical science basis*.
2567 Working Group I contribution to the to the Fifth Assessment Report of the Intergovernmental

2568 Panel on Climate Change (IPCC), Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M. M. B.,
 2569 Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M. (Eds.), Cambridge
 2570 University Press, Chapter 6, 467-570, 2013.

2571 Cleveland, C. C., Townsend, A. R., Schimel, D. S., Fisher, H., Howarth, R. W., Hedin, L. O.,
 2572 Perakis, S. S., Latty, E. F., von Fischer, J. C., Elseroad, A., and Wasson, M. F.: Global patterns
 2573 of terrestrial biological nitrogen (N₂) fixation in natural ecosystems. *Global Biogeochem. Cy.*,
 2574 13,2, 623-645, doi: 10.1029/1999GB900014, 1999.

2575 Comiso, J.C.: Large decadal decline of the Arctic multiyear ice cover, *J. Climate*, 25, 1176–
 2576 1193, doi: 10.1175/JCLI-D-11-00113.1, 2012.

2577 Compton, J. E., Harrison, J. A. Dennis, R. L., Greaver, T. L., Hill, B. H. Jordan, S. J., Walker,
 2578 H., and Campbell, H. V.: Ecosystem services altered by human changes in the nitrogen cycle:
 2579 a new perspective for US decision making, *Ecol. Lett.*, 14, 8, 804– 815, doi: 10.1111/j.1461-
 2580 0248.2011.01631.x., 2011.

2581 Conrad, R.: Metabolism of nitric oxide in soil and soil microorganisms and regulation of flux
 2582 into the atmosphere, in: *Microbiology of atmospheric trace gases*, Murrell, J. C. and Kelly, D.
 2583 P. (Eds.), NATO ASI Series, vol. 109, Springer, Berlin, pp. 167–203, doi: 10.1007/978-3-642-
 2584 61096-7_11, 1996.

2585 Corbett, J. J., Lack, D. A., Winebrake, J. J., Harder, S., Silberman, J. A., Gold, M.: Arctic
 2586 shipping emissions inventories and future scenarios, *Atmos. Chem. Phys.* 10, 9689–9704,
 2587 doi.org/10.5194/acp-10-9689-2010, 2010.

2588 Cornell, S., Mace, K., Coeppicus, S., Duce, R., Huebert, B., Jickells, T. and Zhuang, L. Z.:
 2589 Organic nitrogen in Hawaiian rain and aerosol, *J. Geophys. Res.*, 106, 7973–7983, doi:
 2590 10.1029/2000JD900655, 2001.

2591
 2592 Cornell, S. E., Jickells, T. D., Cape, J. N., Rowland, A. P., and Duce, R. A.: Organic nitrogen
 2593 deposition on land and coastal environments: a review of methods and data, *Atmos. Env.*, 37,
 2594 16, 2173-2191, doi:10.1016/S1352-2310(03)00133-X, 2003.

2595
 2596 Cornell, S. E.: Atmospheric nitrogen deposition: revising the question of the importance of the
 2597 organic component, *Environ. Pollut.*, 159, 2214–2222, doi:10.1016/j.envpol.2010.11.014,
 2598 2011.

2599
 2600 Cottrell, B. A., Gonsior, M., Isabelle, L. M., Luo, W., Perraud, V., McIntire, T. M., Pankow, J.
 2601 F., Schmitt-Kopplin, P., Cooper, W. J., and Simpson, A. J.: A regional study of the seasonal
 2602 variation in the molecular composition of rainwater, *Atmos. Environ.*, 77, 588-597, doi:
 2603 10.1016/j.atmosenv.2013.05.027, 2013.

2604 Cox, R. A. and Roffey, M. J.: Thermal decomposition of peroxyacetylnitrate in the presence
 2605 of nitric oxide, *Environ. Sci. Technol.*, 11, 9, 900-906, doi: 10.1021/es60132a010, 1977.

Formatted: Font: Not Bold

Davidson, C. I., Miller, J. M., and Pleskow, M.A.: The influence of surface structure on predicted particle dry deposition to natural grass canopies, *Water Air Soil Poll.*, 18, 25-43, doi: 10.1007/BF02419401, 1982.

Davidson, E. A. and Kanter, D.: Inventories and scenarios of nitrous oxide emissions, *Environ. Res. Lett.*, 9, 105012, doi:10.1088/1748-9326/9/10/105012, 2014.

Davidson, C. I., Miller, J. M., and Pleskow, M.A.: The influence of surface structure on predicted particle dry deposition to natural grass canopies, *Water Air Soil Poll.*, 18, 25-43, doi: 10.1007/BF02419401, 1982.

Davidson, E. A., de Carvalho, C. J. R., Figueira, A. M., Ishida, F. Y., Ometto, J. P. H. B., Nardoto, G. B., Sabá, R. T., Hayashi, S. N., Leal, E. C., Vieira, I. C. G., and Martinelli, L.A.: Recuperation of nitrogen cycling in Amazonian forests following agricultural abandonment, *Nature*, 447, 995–998, doi:10.1038/nature05900, 2007.

Davidson, E. A., David, M. B., Galloway, J. N., Goodale, C. L., Haeuber, R., Harrison, J. A., Howarth, R. W., Jaynes, D. B., Lowrance, R. R., Nolan, B. T., Peel, J. L., Pinder, R. W., Porter E., Snyder, C. S., Townsend, A. R., and Ward, M. H.: *Excess nitrogen in the U.S. environment: trends, risks, and solutions, Issues in Ecology, Report Number 15, Ecological Society of America, 1-16, 2012.*

De Haan, D. O., Hawkins, L. N., Kononenko, J. A., Turley, J. J., Corrigan, A. L., Tolbert, M. A., and Jimenez, J. L.: Formation of nitrogen-containing oligomers by methylglyoxal and amines in simulated evaporating cloud droplets, *Environ. Sci. Technol.*, 45, 3, 984-991, doi: 10.1021/es102933x, 2011.

De Vries, W., Du, E., and Butterbach-Bahl, K.: Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems, *Curr. Opin. Environ. Sustain.* 9-10, 90-104, doi: 10.1016/j.cosust.2014.09.001, 2014.

Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings between changes in the climate system and biogeochemistry, in: *Climate Change 2007: The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L. (Eds.), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, Chapter 7, pp. 499-587, 2007.

Dentener, F. J. and Crutzen, P.J.: A three-dimensional model of the global ammonia cycle, *J. Atmos. Chem.*, 19, 331–369, doi:10.1007/BF00694492, 1994.

Derwent, R. G., Collins, W. J., Jenkin, M. E., Johnson, C. E., and Stevenson, D. S.: The global distribution of secondary particulate matter in a 3-D Lagrangian chemistry transport model, *J. Atmos. Chem.*, 44, 57–95, doi:10.1023/A:1022139814102, 2003.

Desyaterik, Y., Sun, Y., Shen, X. H., Lee, T. Y., Wang, X. F., Wang, T., and Collett, J. L.: Speciation of "brown" carbon in cloud water impacted by agricultural biomass burning in eastern China, *J. Geophys. Res.-Atmos.*, 118, 13, 7389–7399, doi: 10.1002/jgrd.50561, 2013.

Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N., and Dunne, J. P.: Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, 445, 163–167, doi: 10.1038/nature05392, 2007.

Doherty, R. M., Wild, O., Shindell, D. T., Zeng, G., MacKenzie, I. A., Collins, W. J., Fiore, A. M., Stevenson, D. S., Dentener, F. J., Schultz, M. G., Hess, P., Derwent, R. G., and Keating, T. J.: Impacts of climate change on surface ozone and intercontinental ozone pollution: A multi-model study, *J. Geophys. Res.-Atmos.*, 118, 3744–3763, doi:10.1002/jgrd.50266, 2013.

Doskey, P. V., Kotamarthi, V. R., Fukui, Y., Cook, D. R., Breitbeil III, F. W., and Wesely, M. L.: Air-surface exchange of peroxyacetyl nitrate at a grassland site, *J. Geophys. Res.-Atmos.*, 109, D10310, doi: 10.1029/2004JD004533, 2004.

Doskey, P. V., V. R. Kotamarthi, Y. Fukui, D. R. Cook, F. W. Breitbeil III, and M. L. Wesely (2004). Air-surface exchange of peroxyacetyl nitrate at a grassland site, *J. Geophys. Res.*, 109, D10310, doi:10.1029/2004JD004

Drake, J. E., Gallet-Budynek, A., Hofmockel, K. S., Bernhardt, E. S., Billings, S. A., Jackson, R. B., Johnson, K. S., Lichter, J., McCarthy, H. R., McCormack, M. L., Moore, D. J. P., Oren, R., Palmroth, S., Phillips, R. P., Pippen, J. S., Pritchard, S. G., Treseder, K. K., Schlesinger, W. H., DeLucia, E. H., and Finzi, A. C.: Increases in the flux of carbon belowground stimulate nitrogen uptake and sustain the long-term enhancement of forest productivity under elevated CO₂, *Ecol. Lett.*, 14, 349–357, doi:10.1111/j.1461-0248.2011.01593.x, 2011.

Duan, F.-K., Liu, X.-D., He, K.-B., and Dong, S.-P.: Measurements and characteristics of nitrogen-containing compounds in atmospheric particulate matter in Beijing, China, *B. Environ. Contam. Tox.*, 82, 3, 332–337, doi: 10.1007/s00128-008-9560-0, 2009.

Duce, R. A., LaRoche, J., Altieri, K., Arrigo, K. R., Baker, A. R., Capone, D. G., Cornell, S., Dentener, F., Galloway, J., Ganeshram, R. S., Geider, R. J., Jickells, T., Kuypers, M. M., Langlois, R., Liss, P. S., Liu, S. M., Middelburg, J. J., Moore, C. M., Nickovic, S., Oschlies, A., Pedersen, T., Prospero, J., Schlitzer, R., Seitzinger, S., Sorensen, L. L., Uematsu, M., Ulloa, O., Voss, M., Ward, B., and Zamora, L.: Impacts of atmospheric anthropogenic nitrogen on the open ocean, *Science*, 320, 893–897, doi:10.1126/science.1150369, 2008.

Dutkiewicz, S., Ward, B. A., Monteiro, F., and Follows, M. J.: Interconnection of nitrogen fixers and iron in the Pacific Ocean: Theory and numerical simulations, *Global Biogeochem. Cy.*, 26, GB1012, doi:10.1029/2011GB004039, 2012.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

2678 Duyzer, J. H., Dorsey, J. R., Gallagher, M. W., Pilegaard, K., and Walton, S.: Oxidized nitrogen
 2679 and ozone interaction with forests. II: multi-layer process-oriented modelling results and a
 2680 sensitivity study for Douglas fir, Q. J. Roy. Meteor. Soc., 130, 1957–1971, doi:
 2681 10.1256/qj.03.125, 2004.

2682 Ederli, L., Morettini, R., Borgogni, A., Wasternack, C., Miersch, O., Reale, L., Ferranti, F.,
 2683 Tosti, N., and Pasqualini, S.: Interaction between nitric oxide and ethylene in the induction of
 2684 alternative oxidase in ozone-treated tobacco plants, Plant Physiol., 142, 595–608, doi:10.1104/
 2685 pp.106.085472, 2006.

2686 EMEP.: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying
 2687 componenets, EMEP Status Report 1/2014 (August 29 2014), 2014.

2688 Engardt, M. and Langner, J.: Simulations of future sulphur and nitrogen deposition over Europe
 2689 using meteorological data from three regional climate projections, Tellus B, 65, 20348,
 2690 doi:10.3402/tellusb.v65i0.20348, 2013.

2691 Erisman, J. W., Domburg, N., de Vries, W., Kros, H., de Haan, Band Sanders, K.: The Dutch
 2692 N-cascade in the European perspective, Science in China Series C: Life Sciences, 48, 2,
 2693 Supplement, 827–842, doi: 10.1007/BF03187122, 2005.

2694 Erisman, J. W., Sutton, M. A., Galloway, J., Klimont, Z., and Winiwarter, W.: How a century
 2695 of ammonia synthesis changed the world, Nat. Geosci., 1, 636–639, doi: 10.1038/ngeo325,
 2696 2008.

2697 Erisman, J. W., Galloway, J. N., Seitzinger, S., Bleeker, A., Dise, N. B., Petrescu, A. M. R.,
 2698 Leach, A. M., and de Vries, W.: Consequences of human modification of the global nitrogen
 2699 cycle, Philos. T. Roy. Soc. B, 368, 1621, 20130116, doi:10.1098/rstb.2013.0116, 2013.

2700 Eugster, O. and Gruber, N.: A probabilistic estimate of global marine N-fixation and
 2701 denitrification, Global Biogeochem. Cy., 26, GB4013, doi:10.1029/2012GB004300, 2012.

2702 Facchini, M. C., Decesari, S., Rinaldi, M., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S.,
 2703 Moretti, F., Tagliavini, E., Ceburnis, D., and O'Dowd, C. D.: Important source of marine
 2704 secondary organic aerosol from biogenic amines, Environ. Sci. Technol., 42, 9116–9121, doi:
 2705 10.1021/es8018385, 2008.

2706 FAO.: Food and Agriculture Organisation's Statistical Database, FAOSTAT, Rome,
 2707 <http://faostat.fao.org/site/291/default.aspx>, 2012.

2708 [Farmer, D. K. and Cohen, R. C.: Observations of HNO₃, ΣAN, ΣPN and NO₂ fluxes: evidence](#)
 2709 [for rapid HO_x chemistry within a pine forest canopy, Atmos. Chem. Phys., 8, 3899–3917, doi:](#)
 2710 [10.5194/acp-8-3899-2008, 2008.](#)

2711 Farquhar, G. D., Firth, P. M., Wetselaar, R., and Weir, B.: On the gaseous exchange of
 2712 ammonia between leaves and the environment: determination of the ammonia compensation
 2713 point, Plant Physiol., 66, 710–714, doi:10.1104/pp.66.4.710, 1980.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

2714 Feick, G. and Hainer, R. M.: On the thermal decomposition of ammonium nitrate. Steady-state
 2715 reaction temperatures and reaction rate, *J. Am. Chem. Soc.*, 76, 22, 5860-5863, doi:
 2716 10.1021/ja01651a096, 1954.

2717 Felle, H. H. and Hanstein, S.: The apoplastic pH of the substomatal cavity of *Vicia faba* leaves
 2718 and its regulation responding to different stress factors, *J. Exp. Bot.*, 53, 73–82, doi:
 2719 10.1093/jexbot/53.366.73, 2002.

2720 Finzi, A. C., Norby, R. J., Calfapietra, C., Gallet-Budynek, A., Gielen, B., Holmes, W. E.,
 2721 Hoosbeek, M. R., Iverson, C. M., Jackson, R. B., Kubiske, M. E., Fedford, J., Liberloo, M.,
 2722 Oren, R., Polle, A., Pritchard, S., Zak, D.R., Schlesinger, W. H., and Ceulemans, R.: Increases
 2723 in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest
 2724 productivity under elevated CO₂, *P. Natl. Acad. Sci. USA*, 104, 35, 14014–14019, doi:
 2725 10.1073/pnas.0706518104, 2007.

2726 Flechard, C. R. and Fowler, D.: Atmospheric ammonia at a moorland site. II: Long-term
 2727 surface-atmosphere micrometeorological flux measurements, *Q. J. Roy. Meteor. Soc.*, 124,
 2728 759–791, doi: 10.1002/qj.49712454706, 1998.

2729 Flechard, C. R., Fowler, D., Sutton, M. A., and Cape, J. N.: A dynamic chemical model of bi-
 2730 directional ammonia exchange between semi-natural vegetation and the atmosphere, *Q. J. Roy.
 2731 Meteor. Soc.*, 125, 2611–2641, doi: 10.1002/qj.49712555914, 1999.

2732 Flechard, C. R., Spirig, C., Neftel, A., and Ammann, C.: The annual ammonia budget of
 2733 fertilized cut grassland – Part 2: Seasonal variations and compensation point modeling,
 2734 *Biogeosciences*, 7, 537–556, doi: 10.5194/bg-7-537-2010, 2010.

2735 Flechard, C. R., Nemitz, E., Smith, R. I., Fowler, D., Vermeulen, A. T., Bleeker, A., Erisman,
 2736 J. W., Simpson, D., Zhang, L., Tang, Y. S., and Sutton, M. A.: Dry deposition of reactive
 2737 nitrogen to European ecosystems: a comparison of inferential models across the NitroEurope
 2738 network, *Atmos. Chem. Phys.*, 11, 2703–2728, doi:10.5194/acp-11-2703-2011, 2011.

2739 Flechard, C. R., Massad, R.-S., Loubet, B., Personne, E., Simpson, D., Bash, J. O., Cooter, E.
 2740 J., Nemitz, E., and Sutton, M. A.: Advances in understanding, models and parameterizations
 2741 of biosphere-atmosphere ammonia exchange, *Biogeosciences*, 10, 5183–5225, doi:
 2742 10.5194/bg-10-5183-2013, 2013.

2743 Fowler, D., Pitcairn, C. E. R., Sutton, M. A., Flechard, C., Loubet, B., Coyle, M., and Munro,
 2744 R.C.: The mass budget of atmospheric ammonia in woodland within 1km of livestock
 2745 buildings, *Environ. Pollut.*, 102, S1, 343-348, doi: 10.1016/S0269-7491(98)80053-5, 1998.

2746

2747 Fowler, D., Pitcairn, C.E.R., Sutton, M.A., Flechard, C., Loubet, B., Coyle, M. and Munro,
 2748 R.C. 1998. The mass budget of atmospheric ammonia in woodland within 1km of livestock
 2749 buildings. *Environ. Pollut.*, 102, S1, 343-348.

Formatted: Font: 12 pt

Formatted: None, Space After: 10 pt, Line spacing: Multiple
1.15 li, No bullets or numbering

2750 ~~Fowler, D., Morse, A. P., Gallagher, M. W., and Choularton, T. W.: 1990. Measurements of~~
 2751 ~~cloud water deposition on vegetation using a lysimeter and a flux gradient technique. Tellus~~
 2752 ~~42(b), 285-293, doi: 10.1034/j.1600-0889.1990.t01-1-00007.x, 1990.~~

2753

2754 ~~Fowler, D., Morse, A. P., Gallagher, M. W. & Choularton, T. W. 1990. Measurements of cloud~~
 2755 ~~water deposition on vegetation using a lysimeter and a flux gradient technique. Tellus 42(b),~~
 2756 ~~285-293.~~

2757

2758 Fowler, D., Pilegaard, K., Sutton, M. A., Ambus, P., Raivonen, M., Duyzer, J., Simpson, D.,
 2759 Fagerli, H., Fuzzi, S., Schjoerring, J. K., Granier, C., Neftel, A., Isaksen, I. S. A., Laj, P.,
 2760 Maione, M., Monks, P. S., Burkhardt, J., Daemmgen, U., Neirynck, J., Personne, E., Wichink-
 2761 Kruit, R., Butterbach-Bahl, K., Flechard, C., Tuovinen, J. P., Coyle, M., Gerosa, G., Loubet,
 2762 B., Altimir, N., Gruenhage, L., Ammann, C., Cieslik, S., Paoletti, E., Mikkelsen, T. N., Ro-
 2763 Poulsen, H., Cellier, P., Cape, J. N., Horváth, L., Loreto, F., Niinemets, Ü., Palmer, P. I.,
 2764 Rinne, J., Misztal, P., Nemitz, E., Nilsson, D., Pryor, S., Gallagher, M. W., Vesala, T., Skiba,
 2765 U., Brüggemann, N., Zechmeister-Boltenstern, S., Williams, J., O'Dowd, C., Facchini, M. C.,
 2766 de Leeuw, G., Flossman, A., Chaumerliac, N., and Erisman, J. W.: Atmospheric composition
 2767 change: Ecosystems–Atmosphere interactions, Atmos. Environ., 43, 5193–5267, doi:
 2768 10.1016/j.atmosenv.2009.07.068, 2009.

2769 Fowler, D., Coyle, M., Skiba, U., Sutton, M. A., Cape, J. N., Reis, S., Sheppard, L. J., Jenkins,
 2770 A., Grizzetti, B., Galloway, J. N., Vitousek, P., Leach, A., Bouwman, A. F., Butterbach-Bahl,
 2771 K., Dentener, F., Stevenson, D., Amann, M., and Voss, M.: The global nitrogen cycle in the
 2772 twenty-first century, Philos. T. Roy. Soc. B, 368, 1621, 20130164, doi:
 2773 10.1098/rstb.2013.0164, 2013.

2774 Fry, J. L., Draper, D. C., Zarzana, K. J., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Brown,
 2775 S. S., Cohen, R. C., Kaser, L., Hansel, A., Cappellin, L., Karl, T., Roux, A. H., Turnipseed, A.,
 2776 Cantrell, C., Lefer, B. L., and Grossberg, N.: Observations of gas- and aerosol-phase organic
 2777 nitrates at BEACHON-RoMBAS 2011, Atmos. Chem. Phys, 13, 8585–8605, doi: 10.5194/acp-
 2778 13-8585-2013, 2013.

2779 Fu, F.-X. and Bell, P. R. F.: Effect of salinity on growth, pigmentation, N₂ fixation and alkaline
 2780 phosphatase activity of cultured *Trichodesmium* sp., Mar. Ecol.-Prog Ser., 257, 69–76, doi:
 2781 10.3354/meps257069, 2003.

2782 Fu, F.-X., Mulholland, M. R., Garcia, N. S., Beck, A., Bernhardt, P. W., Warner, M. E.,
 2783 Sanudo-Wilhelmy, S. A., Hutchins, D. A.: Interactions between changing pCO₂, N₂ fixation,
 2784 and Fe limitation in the marine unicellular cyanobacterium *Crocospaera*, Limnol. Oceanogr.,
 2785 53, 2472–2484, doi: 10.4319/lo.2008.53.6.2472, 2008.

Formatted: Level 1, Space After: 10 pt, Line spacing: Multiple 1.15 li, Tab stops: Not at 16.51 cm

Formatted: Font: Not Italic

Formatted: Font: Not Bold

Formatted: Font: 12 pt

Formatted: None, Space After: 10 pt, Line spacing: Multiple 1.15 li, No bullets or numbering

Formatted: Level 1, Space After: 10 pt, Line spacing: Multiple 1.15 li, Tab stops: -2.01 cm, Left + -1.27 cm, Left + 0 cm, Left + 1.27 cm, Left + 2.06 cm, Left + 3.81 cm, Left + 5.08 cm, Left + 6.35 cm, Left + 7.62 cm, Left + 8.89 cm, Left + 10.16 cm, Left + 11.43 cm, Left + 12.7 cm, Left + 13.97 cm, Left + 15.24 cm, Left + 16.51 cm, Left

Formatted: Level 1, Space After: 10 pt, Line spacing: Multiple 1.15 li

2786 Fu, F., Yu, E., Garcia, N. S., Gale, J., Luo, Y., Webb, E. A., and Hutchins, D. A.: Differing
 2787 responses of marine N₂ fixers to warming and consequences for future diazotroph community
 2788 structure, *Aquat. Microb. Ecol.*, 72, 1, 33–46, doi:10.3354/ame01683, 2014.

2789 Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B.,
 2790 Cosby, J.: The nitrogen cascade, *BioScience*, 53 (4), 341–356, doi: 10.1641/0006-3568, 2003.

2791 Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S.
 2792 P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F.,
 2793 Porter, J. H., Townsend, A. R., and Vörösmarty, C. J.: Nitrogen cycles: past, present, and
 2794 future, *Biogeochemistry*, 70, 153–226. doi:10.1007/s10533-004-0370-0, 2004.

2795 Ganzeveld, L., Bouwman, A., Stehfest, E., van Vuuren, D. P., Eickhout, B., and Lelieveld, J.:
 2796 Impact of future land use and land cover changes on atmospheric chemistry-climate
 2797 interactions, *J. Geophys. Res.*, 115, D23301, doi:10.1029/2010JD014041, 2010.

2798 Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty,
 2799 J., Robinson, S., Thomas, S. M., and Toulmin, C.: Food security: The challenge of feeding 9
 2800 billion people, *Science*, 327, 5967, 812–818, doi:10.1126/science.1185383, 2010.

2801 Gorzelska, K. and Galloway, J. N.: Amine nitrogen in atmospheric environment over ~~the~~
 2802 ~~North~~the North Atlantic ocean, *Global Biogeochem. Cy.*, 4, 309–333, doi:
 2803 10.1029/GB004i003p00309, 1990.

2804 Gregory, J. M., Jones, C. D., Cadule, P., and Friedlingstein, P.: Quantifying carbon cycle
 2805 feedbacks, *J. Climate*, 22, 5232–5250, doi:10.1175/2009JCLI2949.1, 2009.

2806 Groffman, P. M., Altabet, M. A., Böhlke, J.K., Butterbach-Bahl, K., David, M. B., Firestone,
 2807 M. K., Giblin, A.E., Kana, T. M., Nielsen, L. P., and Voytek, M. A.: Methods for measuring
 2808 denitrification: Diverse approaches to a difficult problem, *Ecol. Appl.*, 16, 2091–2122,
 2809 doi:10.1890/1051-0761(2006)016[2091:MFMDDA]2.0.CO;2, 2006.

2810 Groffman, P. M., Butterbach-Bahl, K., Fulweiler, R. W., Gold, A. J., Morse, J. L., Stander, E.
 2811 K., Tague, C., Tonitto, C., and Vidon, P.: Challenges to incorporating spatially and temporally
 2812 explicit phenomena (hotspots and hot moments), *Biogeochemistry*, 93, 49–77, doi:
 2813 10.1007/s10533-008-9277-5, 2009.

2814 Großkopf, T., Mohr, W., Baustian, T., Schunck, H., Gill, D., Kuypers, M. M. M., Lavik, G.,
 2815 Schmitz, R. A., Wallace, D. W. R., and LaRoche, J.: Doubling of marine dinitrogen-fixation
 2816 rates based on direct measurements, *Nature*, 488, 361–364, doi:10.1038/nature11338, 2012.

2817 Guo, L. B. and Gifford, R. M.: Soil carbon stocks and land use change: a meta analysis, ~~Global~~
 2818 Global *Change Biol.*, 8, 345–360, doi: 10.1046/j.1354-1013.2002.00486.x, 2002.

2819 Hauglustaine, D. A., Balkanski, Y., and Schulz, M.: A global model simulation of present and
 2820 future nitrate aerosols and their direct radiative forcing of climate, *Atmos. Chem. Phys.*, 14,
 2821 11031–11063, doi:10.5194/acp-14-11031-2014, 2014.

2822 Hedegaard, G. B., Christensen, J. H., and Brandt, J.: The relative importance of impacts from
 2823 climate change vs. emissions change on air pollution levels in the 21st century, *Atmos. Chem.*
 2824 *Phys.*, 13, 3569–3585, doi:10.5194/acp-13-3569-2013, 2013.

2825 Herridge, D. F., Peoples, M. B., and Boddey, R. M.: Global inputs of biological nitrogen
 2826 fixation in agricultural systems, *Plant Soil*, 311, 1-18, doi: 10.1007/s11104-008-9668-3, 2008.

2827 Hofmockel, K. S. and Schlesinger, W. H.: Carbon dioxide effects on heterotrophic dinitrogen
 2828 fixation in a temperate pine forest, *Soil Sci. Soc. Am. J.*, 71, 140-144,
 2829 doi:10.2136/sssaj2006.110, 2007.

2830 Hofmockel, K. S., Gallet-Budynek, A., McCarthy, H. R., Currie, W. S., Jackson, R. B., and
 2831 ~~Finzi~~and Finzi, A. C.: Sources of increased N uptake in forest trees growing under elevated
 2832 CO₂: results of a large-scale ¹⁵N study, *Global Change Biol.*, 17, 3338–3350,
 2833 doi:10.1111/j.1365-2486.2011.02465.x, 2011.

2834 Houlton, B. Z., Wang, Y.-P., Vitousek, P. M., and Field, C. B.: A unifying framework for
 2835 dinitrogen fixation in the terrestrial biosphere, *Nature*, 454, 327-330, doi:
 2836 10.1038/nature07028, 2008.

2837 Howarth, R.W. and Cole, J.J.: Molybdenum availability, nitrogen limitation, and
 2838 phytoplankton growth in natural waters, *Science*, 229, 653–655, doi:
 2839 10.1126/science.229.4714.653, 1985.

2840 Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C., and
 2841 Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions:
 2842 implementation and space based-constraints, *Atmos. Chem. Phys.*, 12, 7779-7795,
 2843 doi:10.5194/acp-12-7779-2012, 2012.

2844 Hungate, B. A., Dijkstra, P., Wu, Z., Duval, B. D., Day, F. P., Johnson, D. W., Megonigal, J.
 2845 P., Brown, A. L. P., and Garland, J. L.: Cumulative response of ecosystem carbon and nitrogen
 2846 stocks to chronic CO₂ exposure in a subtropical oak woodland, *New Phytol.*, 200, 753–766,
 2847 doi:10.1111/nph.12333, 2013.

2848 Hutchins, D. A., Fu, F.-X., Zhang, Y., Warner, M. E., Feng, Y., Portune, K., Bernhardt, P. W.,
 2849 and Mulholland, M.R.: CO₂ control of *Trichodesmium* N₂ fixation, photosynthesis, growth
 2850 rates, and elemental ratios: Implications for past, present, and future ocean biogeochemistry,
 2851 *Limnol. Oceanogr.*, ~~52~~, 52, 1293–1304, doi: 10.4319/lo.2007.52.4.1293, 2007.

2852 Hutchins, D.A., Mulholland, M.R., Fu, F.-X.: Nutrient Cycles and Marine Microbes in a CO₂-
 2853 Enriched Ocean, *Oceanography*, 22, 4, 128-145, 2009.

2854 Hutchins, D. A., Fu, F.-X., Webb, E. A., Walworth, N., Tagliabue, A.: Taxon-specific response
 2855 of marine nitrogen fixers to elevated carbon dioxide concentrations, *Nature Geosci.*, 6, 790–
 2856 795, doi: 10.1038/ngeo1858, 2013.

2857 Iinuma, Y., Boge, O., Grafe, R., and Herrmann, H.: Methyl-nitrocatechols: Atmospheric tracer
 2858 compounds for biomass burning secondary organic aerosols, *Environ. Sci. Technol.*, 44, 22,
 2859 8453-8459, doi: 10.1021/es102938a, 2010.

2860 IPCC (Intergovernmental Panel on Climate Change): 2006 IPCC Guidelines for National
 2861 Greenhouse Gas Inventories, prepared by the National Greenhouse Gas Inventories
 2862 Programme (Eggleston, H.S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K. (Eds.)), IGES,
 2863 Japan, 2006.

2864 IPCC (Intergovernmental Panel on Climate Change): Climate Change 2013: The physical
 2865 science basis. Working group I contribution to the Fifth Assessment Report of the
 2866 Intergovernmental Panel on Climate Change (IPCC) (Stocker, T. F., Qin, D., Plattner, G.-K.,
 2867 Tignor, M. M. B., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M.
 2868 (Eds.)), Cambridge University Press, 1535 pp., 2013.

2869 Jacobson, M. C., Hansson, H. C., Noone, K. J., and Charlson, R. J.: Organic atmospheric
 2870 aerosols: review and state of the science, *Rev. Geophys.*, 38, 267–294, doi:
 2871 10.1029/1998RG000045, 2000.

2872 Jaeglé, L., Steinberger, L., Martin, R. V., and Chance, K.: Global partitioning of NO_x sources
 2873 using satellite observations: Relative roles of fossil fuel combustion, biomass burning and soil
 2874 emissions, *Faraday Discuss.*, 130, 407–423, doi: 10.1039/b502128f, 2005.

2875 Jarvis, P. G.: The interpretation of the variations in leaf water potential and stomatal
 2876 conductance found in canopies in the field, *Philos. T. Roy. Soc. B*, 273, 927, 593–610, doi:
 2877 10.1098/rstb.1976.0035, 1976.

2878 Jensen, L.S., Schjoerring, J. K., van der Hoek, K. W., Poulsen, H. D., Zevenbergen, J. F.,
 2879 Pallière, C., Lammel, J., Brentrup, F., Jongbloed, A.W., Willems, J., and van Grinsven, H.:
 2880 Benefits of nitrogen for food, fibre and industrial production, in: *The European Nitrogen*
 2881 *Assessment: Sources, effects and policy perspectives*, Sutton, M. A., Howard, C. M., Erisman,
 2882 J. W., Billen, G., Bleeker, A. Grennfelt, P., van Grinsven, H., and Grizzetti, B. (Eds.),
 2883 Cambridge University Press, 32-61, 2011.

2884 Jickells, T., Baker, A. R., Cape, J. N., Cornell, S. E., and Nemitz, E.: The cycling of organic
 2885 nitrogen through the atmosphere, *Philos. T. Roy. Soc. B*, 368, 1621, 20130115, doi:
 2886 10.1098/rstb.2013.0115, 2013.

2887 Johnson, M. T. and Bell, T. G.: Coupling between dimethylsulfide emissions and the ocean–
 2888 atmosphere exchange of ammonia. *Environ. Chem.*, 5, 259–267, doi: 10.1071/EN08030, 2008.

2889 Johnson, M. T., Liss, P. S., Bell, T. G., Lesworth, T. J., Baker, A. R., Hind, A. J., Jickells, T.
 2890 D., Biswas, K. F., Woodward, E. M. S., and Gibb, S. W.: Field observations of the ocean–
 2891 atmosphere exchange of ammonia: Fundamental importance of temperature as revealed by a
 2892 comparison of high and low latitudes, *Global Biogeochem. Cy.* 22, GB1019, doi:
 2893 10.1029/2007GB003039, 2008.

2894 Jones, M. R., Leith, I. D., Fowler, D., Raven, J. A., Sutton, M. A., Nemitz, E., Cape, J. N.,
 2895 Sheppard, L. J., Smith, R. I., and Theobald, M. R.: Concentration-dependent NH₃ deposition
 2896 processes for mixed moorland semi-natural vegetation, *Atmos. Environ.*, 41, 2049–2060, doi:
 2897 10.1016/j.atmosenv.2006.11.003, 2007.

2898 Jungkunst, H. F., Flessa, H., Scherber, C., and Fiedler, S.: Groundwater level controls CO₂,
 2899 N₂O and CH₄ fluxes of three different hydromorphic soil types of a temperate forest ecosystem,
 2900 *Soil Biol. Biochem.*, 40, 2047-2054, doi: 10.1016/j.soilbio.2008.04.015, 2008.

2901 Kammann, C., Müller, C., Grünhage, L., and Jäger, H. J.: Elevated CO₂ stimulates N₂O
 2902 emissions in permanent grassland, *Soil Biol. Biochem.*, 40, 2194-2205,
 2903 doi:10.1016/j.soilbio.2008.04.012, 2008.

2904 Kanakidou, M., Duce, R.A., Prospero, J.M., Baker, A.R., Benitez-Nelson, C., Dentener, F.J.,
 2905 Hunter, K.A., Liss, P.S., Mahowald, N., Okin, G.S., Sarin, M., Tsigaridis, K., Uematsu, M.,
 2906 Zamora, L.M., and Zhu, T.: Atmospheric fluxes of organic N and P to the global ocean, *Glob.*
 2907 *Biogeochem. Cy.*, 26, GB3026, doi: 10.1029/2011GB004277, 2012.

2908 Kicklither, D. W., Cai, Y., Zhuang, Q., Parfenova, E. I., Paltsev, S., Sokolov, A. P., Melillo,
 2909 J. M., Reilly, J. M., Tchekakova, N. M., and Lu, X.: Potential influence of climate-induced
 2910 vegetation shifts on future land use and associated land carbon fluxes in Northern Eurasia.
 2911 *Environ. Res. Lett.*, 9, 035004, doi:10.1088/1748-9326/9/3/035004, 2014.

2912 Kieloaho, A. J., Hellen, H., Hakola, H., Manninen, H. E., Nieminen, T., Kulmala, M., and
 2913 Pihlatie, M.: Gas-phase alkylamines in a boreal Scots pine forest air, *Atmos. Environ.*, 80, 369-
 2914 377, doi: 10.1016/j.atmosenv.2013.08.019, 2013.

2915 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné,
 2916 S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S.,
 2917 Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A.,
 2918 Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W.,
 2919 Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R.,
 2920 Makhmutov, V., Mathot, S., Mikkilä, Minginette, P., Mogo, S., Nieminen, T., Onnela, A.,
 2921 Pereira, P., Petäjä, Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F.,
 2922 Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E.,
 2923 Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.:
 2924 Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation,
 2925 *Nature*, 476, 429-433, doi:10.1038/nature10343, 2011.

2926 Kirschbaum, M. U. F., Guo, L. B., and Gifford, R. M.: Why does rainfall affect the trend in
 2927 soil carbon after converting pastures to forests?: A possible explanation based on nitrogen
 2928 dynamics, *Forest Ecol. Manag.*, 255, 2990–3000, doi:10.1016/j.foreco.2008.02.005, 2008.

2929 Kirschbaum, M. U. F., Saggarr, S., Tate, K. R., Thakur, K. P., and Giltrap, D. L.: Quantifying
 2930 the climate-change consequences of shifting land use between forest and agriculture, *Sci. Total*
 2931 *Environ.*, 465, 314–324, doi:10.1016/j.scitotenv.2013.01.026, 2013.

Field Code Changed

Formatted: No underline

Formatted: No underline

2932 Kjellstrom, E., Nikulin, G., Hansson, U., Strandberg, G., and Ullerstig, A.: 21st century
 2933 changes in the European climate: uncertainties derived from an ensemble of regional climate
 2934 model simulations, *Tellus A*, 63, 24–40, doi:10.1111/j.1600-0870.2010.00475.x, 2011.

2935 Kourtchev, I., O'Connor, I. P., Giorio, C., Fuller, S. J., Kristensen, K., Maenhaut, W., Wenger,
 2936 J. C., Sodeau, J. R., Glasius, M., and Kalberer, M.: Effects of anthropogenic emissions on the
 2937 molecular composition of urban organic aerosols: An ultrahigh resolution mass spectrometry
 2938 study, *Atmos. Environ.*, 89, 525–532, doi: 10.1016/j.atmosenv.2014.02.051, 2014.

2939 Kranz, S., Sültemeyer, D., Richter, K.-U., and Rost, B.: Carbon acquisition in *Trichodesmium*:
 2940 The effect of pCO₂ and diurnal changes, *Limnol. Oceanogr.*, 54, 548–559, doi:
 2941 10.4319/lo.2009.54.2.0548, 2009.

2942 Kryza, M., Werner, M., Dore, A. J., Błaś, M., and Sobik M.: The role of annual circulation and
 2943 precipitation on national scale deposition of atmospheric sulphur and nitrogen compounds, *J.*
 2944 *Environ. Manage.*, 109, 70–79, doi: 10.1016/j.jenvman.2012.04.048, 2012.

2945 Kunwar, B. and Kawamura, K.: One-year observations of carbonaceous and nitrogenous
 2946 components and major ions in the aerosols from subtropical Okinawa Island, an outflow region
 2947 of Asian dusts, *Atmos. Chem. Phys.*, 14, 1819–1836, doi:10.5194/acp-14-1819-2014, 2014.

2948

2949 Kurtén, T., Loukonen, V., Vehkamäki, H., and Kulmala, M.: Amines are likely to enhance
 2950 neutral and ion-induced sulfuric acid-water nucleation in the atmosphere more effectively than
 2951 ammonia, *Atmos. Chem. Phys.*, 8, 4095–4103, doi:10.5194/acp-8-4095-2008, 2008.

2952

2953 Laitinen, T., Junninen, H., Parshintsev, J., Ruiz-Jimenez, J., Petäjä, T., Hautala, S., Hartonen,
 2954 K., Worsnop, D., Kulmala, M., and Riekkola, M. L.: Changes in concentration of nitrogen-
 2955 containing compounds in 10 nm particles of boreal forest atmosphere at snowmelt, *J. Aerosol*
 2956 *Sci.*, 70, 1–10, doi: 10.1016/j.jaerosci.2013.12.009, 2014.

2957 Lamarque, J.-F., Kiehl, J. T., Brasseur, G. P., Butler, T., Cameron-Smith, P., Collins, W. D.,
 2958 Collins, W. J., Granier, C., Hauglustaine, D., Hess, P. G., Holland, E. A., Horowitz, L.,
 2959 Lawrence, M. G., McKenna, D., Merilees, P., Prather, M. J., Rasch, P. J., Rotman, D., Shindell,
 2960 D., and Thornton, P.: Assessing future nitrogen deposition and carbon cycle feedback using a
 2961 multimodel approach: Analysis of nitrogen deposition, *J. Geophys. Res.*, 110, D19303, doi:
 2962 10.1029/2005JD005825, 2005.

2963 Lamarque, J.-F., Kyle, G. P., Meinshausen, M., Riahi, K., Smith, S.J., van Vuuren, D. P.,
 2964 Conley, A. J., and Vitt, F.: Global and regional evolution of short-lived radiatively-active gases
 2965 and aerosols in the Representative Concentration Pathways, *Climatic Change*, 109, 191–212,
 2966 doi: 10.1007/s10584-011-0155-0, 2011.

2967 Lamarque, J.-F., Dentener, F., McConnell, J., Ro, C. U., Shaw, M., Vet, R., Bergmann, D.,
 2968 Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H.,
 2969 MacKenzie, I. A., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S.,
 2970 Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D., and Nolan, M.: Multi-model

mean nitrogen and sulfur deposition from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): evaluation of historical and projected future changes, *Atmos. Chem. Phys.*, 13, 7997–8018, doi:10.5194/acp-13-7997-2013, 2013.

LeClair, J. P., Collett, J. P., and Mazzoleni L. R.: Fragmentation analysis of water-soluble atmospheric organic matter using ultrahigh-resolution FT-ICR mass spectrometry, *Environ. Sci. Technol.*, 46, 4312–4322, doi.org/10.1021/es203509b, 2012.

Ledgard, S. F. and Steele, K. W.: Biological nitrogen-fixation in mixed legume grass pastures, *Plant Soil*, 141, 137–153, doi: 10.1007/BF00011314, 1992.

Lee, A. K. Y., Zhao, R., Li, R., Liggio, J., Li, S.-M., and Abbatt, J. P. D.: Formation of light absorbing organo-nitrogen species from evaporation of droplets containing glyoxal and ammonium sulfate. *Environ. Sci. Technol.*, 47 (22), doi: 10.1021/es402687w, 12819–12826, 2013.

Leemans, R., van Amstel, A., Battjes, C., Kreileman, E., and Toet S.: The land cover and carbon cycle consequences of large scale utilizations of biomass as an energy source, *Glob. Environ. Change*, 6, 335–357, doi: 10.1016/S0959-3780(96)00028-3, 1996.

Lesworth, T., Baker, A. R., and Jickells, T.: Aerosol organic nitrogen over the remote Atlantic Ocean, *Atmos. Environ.*, 44, 1887–1893, doi:10.1016/j.atmosenv.2010.02.021, 2010.

Levitan, O., Rosenberg, G., Setlik, I., Setlikova, E., Grigel, J., Klepetar, J., Prasil, O., and Berman-Frank, I.: Elevated CO₂ enhances nitrogen fixation and growth in the marine cyanobacterium *Trichodesmium*, *Global Change Biol.*, 13, 531–538. doi:10.1111/j.1365-2486.2006.01314.x, 2007.

Li, Q. F. and Yu, J. Z.: Determination of total aerosol nitrogen by thermal evolution, *Aerosol Sci. Tech.*, 38, 382–390, doi: 10.1080/02786820490442626, 2004.

Liao, H., Chen, W. T., and Seinfeld, J. H.: Role of climate change in global predictions of future tropospheric ozone and aerosols, *J. Geophys. Res.*, 111, D12, D12304, doi: 10.1029/2005jd006852, 2006.

Liss, P. S. and Slater, P. G.: Flux of Gases across the Air- Sea Interface, *Nature*, 247, 181–184, doi: 10.1038/247181a0, 1974.

Liu, C., Zheng, X., Zhou, Z., Han, S., Wang, Y., Wang, K., Liang, W., Li, M., Chen, D., and Yang, Z.: Nitrous oxide and nitric oxide emissions from an irrigated cotton field in Northern China, *Plant Soil*, 332, 123–134, 10.1007/s11104-009-0278-5, 2010.

Logan, J. A.: Nitrogen oxides in the troposphere: Global and regional budgets, *J. Geophys. Res.*, 88, C15, 10785–10807, doi: 10.1029/JC088iC15p10785, 1983.

Long, S. P., Ainsworth, E. A., Rogers, A., and Ort, D. R.: Rising atmospheric carbon dioxide: plants FACE the future. *Annu. Rev. Plant Biol.*, 55, 591–628, doi: 10.1146/annurev.arplant.55.031903.141610, 2004.

3007 Loubet, B., Milford, C., Sutton, M. A., and Cellier, P.: Investigation of the interaction between
 3008 sources and sinks of atmospheric ammonia in an upland landscape using a simplified
 3009 dispersion-exchange model, [J. Geophys. Res.](#), 106, 24183-24195, doi:
 3010 10.1029/2001JD900238, 2001.

3011 Loubet, B., Asman, W. A., Theobald, M. R., Hertel, O., Tang, Y. S., Robin, P., Hassouna, M.,
 3012 Dämmgen, U., Genermont, S., Cellier, P., and Sutton, M. A.: Ammonia deposition near hot
 3013 spots: Processes, models and monitoring methods, in: Sutton, M. A., Reis, S., and Baker, S. M.
 3014 (Eds.): Atmospheric ammonia. Detecting emissions changes and environmental impacts,
 3015 Springer, 205-267, doi: 10.1007/978-1-4020-9121-6_15, 2009.

3016 [Ludwig, J., Meixner, F. X., Vogel, B., and Forstner, J.: 2001 Soil-air exchange of nitric oxide: an
 3017 overview of processes, environmental factors and modeling studies, *Biogeochemistry*, 52, 225-
 3018 257, doi:10.1023/A:1006424330555, 2001.](#)

3019

3020 Luo, C., Rodriguez, R. L. M., Johnston, E. R., Wu, L., Cheng, L., Xue, K., Tu, Q., Deng, Y.,
 3021 He, Z., Shi, J. Z., Yuan, M. M., Sherry, R. A., Li, D., Luo, Y., Schuur, E. A. G., Chain, P.,
 3022 Tiedje, J. M., Zhou, J., and Konstantinidis, K. T.: Soil microbial community responses to a
 3023 decade of warming revealed by comparative metagenomics, *Appl. Environ. Microbiol.*, 80,
 3024 1777-1786, doi: 10.1128/AEM.03712-13, 2014.

3025 Luo, Y., Melillo, J., Niu, S., Beier, C., Clark, J. S., Classen, A. T., Davidson, E., Dukes, J. S.,
 3026 Evans, R. D., Field, C. B., Czimczik, C. I., Keller, M., Kimball, B. A., Kueppers, L. M., Norby,
 3027 R. J., Pelini, S. L., Pendall, E., Rastetter, E., Six, J., Smith, M., Tjoelker, M. G., Torn, M. S.:
 3028 Coordinated approaches to quantify long-term ecosystem dynamics in response to global
 3029 change, *Global Change Biol.*, 17, 843-854, doi: 10.1111/j.1365-2486.2010.02265.x, 2011.

3030 Luo, Y.-W., Doney, S. C., Anderson, L. A., Benavides, M., Berman-Frank, I., Bode, A.,
 3031 Bonnet, S., Boström, K. H., Böttjer, D., Capone, D. G., Carpenter, E. J., Chen, Y. L., Church,
 3032 M. J., Dore, J. E., Falcón, L. I., Fernández, A., Foster, R. A., Furuya, K., Gómez, F., Gundersen,
 3033 K., Hynes, A. M., Karl, D. M., Kitajima, S., Langlois, R. J., LaRoche, J., Letelier, R. M.,
 3034 Marañón, E., McGillicuddy Jr., D. J., Moisaner, P. H., Moore, C. M., Mouriño-Carballido,
 3035 B., Mulholland, M. R., Needoba, J. A., Orcutt, K. M., Poulton, A. J., Rahav, E., Raimbault, P.,
 3036 Rees, A. P., Riemann, L., Shiozaki, T., Subramaniam, A., Tyrrell, T., Turk-Kubo, K. A.,
 3037 Varela, M., Villareal, T. A., Webb, E. A., White, A. E., Wu, J., and Zehr, J. P.: Database of
 3038 diazotrophs in global ocean: abundance, biomass and nitrogen fixation rates, *Earth Syst. Sci.*
 3039 *Data* 4, 47–73, doi:10.5194/essd-4-47-2012, 2012.

3040 Luo, Y.-W., Lima, I. D., Karl, D. M., Deutsch, C. A., and Doney, S. C.: Data-based assessment
 3041 of environmental controls on global marine nitrogen fixation, *Biogeosciences*, 11, 691–708,
 3042 doi: 10.5194/bg-11-691-2014, 2014.

3043 Luscher, A., Mueller-Harvey, I., Soussana, J. F., Rees, R. M., and Peyraud, J. L.: Potential of
 3044 legume-based grassland-livestock systems in Europe: a review, *Grass Forage Sci.*, 69, 206-
 3045 228, doi: 10.1111/gfs.12124, 2014.

3046
3047
3048 [G.G. McFadyen, J.N. Cape, Peroxyacetyl nitrate in eastern Scotland Science of the](#)
3049 [Total Environment 337 \(2005\) 213–222](#)
3050
3051 Mace, K. A., Duce, R. A., -and Tindale, N. W.: Organic nitrogen in rain and aerosol at Cape
3052 Grim, Tasmania, Australia, J. Geophys. Res.-Atmos., 108, D11, 4338, [doi:10.1029/2002JD003051](#), 2003a.
3053
3054
3055 Mace, K. A., Artaxo, P., and Duce, R. A.: Water-soluble organic nitrogen in Amazon Basin
3056 aerosols during the dry (biomass burning) and wet seasons, J. Geophys. Res.-Atmos., 108, D16,
3057 doi: [10.1029/2003JD003557](#), 4512, 2003b.
3058
3059 Mace, K. A., Kubilay, N., and Duce, R. A.: Organic nitrogen in rain and aerosol in the eastern
3060 Mediterranean atmosphere: an association with atmospheric dust, J. Geophys. Res.-Atmos.,
3061 108, 4320, D10, [doi:10.1029/2002JD002997](#), 2003c.
3062
3063 MacLeod, M., Moran, D., Eory, V., Rees, R. M., Barnes, A., Topp, C. F. E., Ball, B., Hoad, S.,
3064 Wall, E., McVittie, A., Pajot, G., Matthews, R., Smith, P., and Moxey, A.: Developing
3065 greenhouse gas marginal abatement cost curves for agricultural emissions from crops and soils
in the UK, Agr. Syst., 103, 4, 198-209, doi: 10.1016/j.agry.2010.01.002, 2010.
3066
3067 Mahaffey, C., Michaels, A. F., and Capone, D. G.: The conundrum of marine N₂ fixation, Am.
J. Sci., 305, 546–595, doi:10.2475/ajs.305.6-8.546, 2005.
3068
3069 Marino, R., Howarth, R. W., Chan, F., Cole, J. J., Likens, G. E.: Sulfate inhibition of
3070 molybdenum-dependent nitrogen fixation by planktonic cyanobacteria under sea water
3071 conditions: a non-reversible effect, Hydrobiologia, 500, 277-293, doi:
10.1023/A:1024641904568, 2003.
3072
3073 Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., and Evans, M. J.: Global
3074 inventory of nitrogen oxide emissions constrained by space-based observations of NO₂
columns, J. Geophys. Res., 108, D17, 4537, doi:[10.1029/2003JD003453](#), 2003.
3075
3076 Massad, R.-S., Nemitz, E., and Sutton, M. A.: Review and parameterisation of bi-directional
3077 ammonia exchange between vegetation and the atmosphere, Atmos. Chem. Phys., 10, 10359–
10386, doi: [10.5194/acp-10-10359-2010](#), 2010a.
3078
3079 Massad, R.-S., Tuzet, A., Loubet, B., Perrier, A., and Cellier, P.: Model of stomatal ammonia
3080 compensation point (STAMP) in relation to the plant nitrogen and carbon metabolisms and
3081 environmental conditions, Ecol. Model., 221, 479–494, doi: 10.1016/j.ecolmodel.2009.10.029,
2010b.
3082
3083 Matsumoto, K., and Uematsu, M.: Free amino acids in marine aerosols over the western North
3084 Pacific Ocean, Atmos. Environ., 39, 2163-2170, doi: 10.1016/j.atmosenv.2004.12.022, 2005.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Field Code Changed

3085 ~~G. G. McFadyen, G. G., and Cape, J. N.: Peroxyacetyl nitrate in eastern Scotland, Science~~
3086 ~~of the Total Environment 337, (2005) 213–222, doi:10.1016/j.scitotenv.2004.06.016, 2005.~~

3087

3088 McKee, C. M.: Biogeochemical cycles of ammonia and dimethylsulphide in the marine
3089 environment, Ph.D. thesis, University of East Anglia (UEA), Norwich, UK, 2001.

3090

3091 Medinets, S., Skiba, U., Rennenberg, H., and Butterbach-Bahl, K.: A review on soil NO
3092 transformation: Associated processes and possible physiological significance in organisms,
3093 Soil Biol. Biochem., 80, 92–117, doi: 10.1016/j.soilbio.2014.09.025, 2015.

3094 Melillo, J. M., Steudler, P. A., Aber, J. D., Newkirk, K., Lux, H., Bowles, F. P., Catricala, C.,
3095 Magill, A., Ahrens, T., and Morrisseau, S.: Soil warming and carbon-cycle feedbacks to the
3096 climate system, Science, 298, 2173–2176, doi: 10.1126/science.1074153, 2002.

3097 Melillo, J. M., Butler, S., Johnson, J., Mohan, J., Steudler, P., Lux, H., Burrows, E., Bowles,
3098 F., Smith, R., Scott, L., Vario, C., Hill, T., Burton, A., Zhou, Y.-M., and Tang, J.: Soil warming,
3099 carbon-nitrogen interactions, and forest carbon budgets, P. Natl. Acad. Sci. USA, 108, 9508–
3100 9512, doi:10.1073/pnas.1018189108, 2011.

3101 Meyer, M. W.: Absorption and release of ammonia from and to the atmosphere by plants, Ph.
3102 D. thesis, University of Maryland, College Park, MD, USA, 63 pp., 1973.

3103 Mikkelsen, T. N., Beier, C., Jonasson, S., Holmstrup, M., Schmidt, I. K., Ambus, P., Pilegaard,
3104 K., Michelsen, A., Albert, K., Andresen, L. C., Arndal, M. F., Bruun, N., Christensen, S.,
3105 Danbæk, S., Gundersen, P., Jørgensen, P., Kongstad, J., Maraldo, K., Priemé, A., Riis-Nielsen,
3106 T., Ro-Poulsen, H., Stevnbak, K., Selsted, M. B., Sørensen, P., Larsen, K. S., Carter, M. S.,
3107 Martinussen, T., Miglietta, F., and Sverdrup, H.: Experimental design of multifactor climate
3108 change experiments with elevated CO₂, warming and drought – the CLIMAITE project, Funct.
3109 Ecol., 22, 185–195, doi: 10.1111/j.1365-2435.2007.01362.x, 2008.

3110 Miyazaki, K., Eskes, H.J., and Sudo, K.: Global NO_x emission estimates derived from an
3111 assimilation of OMI tropospheric NO₂ columns, Atmos. Chem. Phys., 12, 2263–2288, doi:
3112 10.5194/acp-12-2263-2012, 2012.

3113 Miyazaki, Y., Kawamura, K., and Sawano, M.: Size distributions of organic nitrogen and
3114 carbon in remote marine aerosols: evidence of marine biological origin based on their isotopic
3115 ratios, Geophys. Res. Lett., 37, L06803, doi: 10.1029/2010GL042483, 2010.

3116

3117 Miyazaki, Y., Fu, P. Q., Ono, K., Tachibana, E., and Kawamura, K.: Seasonal cycles of water-
3118 soluble organic nitrogen aerosols in a deciduous broadleaf forest in northern Japan, J. geophys.
3119 Res.-Atmos., 119, 1440–1454, doi: 10.1002/2013JD020713, 2014.

3120 Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M.,
3121 Baklanov, A., Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K., Cooper, O. R.,
3122 Dentener, F., Fowler, D., Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V.,
3123 Guenther, A., Hansson, H. C., Henne, S., Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen,

3124 I. S. A., Jenkin, M. E., Kaiser, J., Kanakidou, M., Klimont, Z., Kulmala, M., Laj, P., Lawrence,
 3125 M. G., Lee, J. D., Lioussé, C., Maione, M., McFiggans, G., Metzger, A., Mieville, A.,
 3126 Moussiopoulos, N., Orlando, J. J., O'Dowd, C. D., Palmer, P. I., Parrish, D. D., Petzold, A.,
 3127 Platt, U., Poeschl, U., Prevot, A. S. H., Reeves, C. E., Reimann, S., Rudich, Y., Sellegri, K.,
 3128 Steinbrecher, R., Simpson, D., ten Brink, H., Theloke, J., van der Werf, G. R., Vautard, R.,
 3129 Vestreng, V., Vlachokostas, Ch., and von Glasow, R.: Atmospheric composition change –
 3130 global and regional air quality, *Atmos. Env.*, 43, 33, 5268-5350.
 3131 doi:10.1016/j.atmosenv.2009.08.021, 2009.

3132 Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D.,
 3133 Granier, C., Law, K. S., Stevenson, D., Tarasova, O., Thouret, V., von Schneidmeyer, E.,
 3134 Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone and its precursors from the
 3135 urban to the global scale from air quality to short lived climate forcer, *Atmos. Chem. Phys.*
 3136 Discussions (to be submitted), 2014.

3137 Monteith, J. L. and Unsworth, M. H.: Principles of environmental physics, 4th edition, Elsevier,
 3138 Amsterdam, 401 pp., 2013.

3139 Montero-Martinez, G., Rinaldi, M., Gilardoni, S., Giulianelli, L., Paglione, M., Decesari, S.,
 3140 Fuzzi, S., and Facchini, M. C.: On the water-soluble organic nitrogen concentration and mass
 3141 size distribution during the fog season in the Po Valley, Italy, *Sci. Total Environ.*, 485-486,
 3142 103-9, doi: 10.1016/j.scitotenv.2014.03.060, 2014.

3143 Montoya, J. P., Holl, C. M., Zehr, J. P., Hansen, A., Villareal, T. A., and Capone, D. G.: High
 3144 rates of N₂ fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean, *Nature*, 430,
 3145 1027–1032, doi: 10.1038/nature02824, 2004.

3146 Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith,
 3147 E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M.,
 3148 Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito,
 3149 M. A., Thingstad, T. F., Tsuda, A., and Ulloa, O.: Processes and patterns of oceanic nutrient
 3150 limitation, *Nat. Geosci.*, 6, 701–710, doi:10.1038/ngeo1765, 2013.

3151 Moore, J. K. and Doney, S. C.: Iron availability limits the ocean nitrogen inventory stabilizing
 3152 feedbacks between marine denitrification and nitrogen fixation, *Global Biogeochem. Cy.*, 21,
 3153 GB2001, doi:10.1029/2006GB002762, 2007.

3154 Moore, J. K., Doney, S. C., Glover, D. M., and Fung, I. Y.: Iron cycling and nutrient-limitation
 3155 patterns in surface waters of the world ocean, *Deep-Sea Res. Pt. II: Topical Studies in*
 3156 *Oceanography*, 49, 463–507, doi: 10.1016/S0967-0645(01)00109-6, 2002.

3157 Moore, J. K., Doney, S. C., Lindsay, K., Mahowald, N., and Michaels, A. F.: Nitrogen fixation
 3158 amplifies the ocean biogeochemical response to decadal timescale variations in mineral dust
 3159 deposition, *Tellus B*, 58, 560–572, doi:10.1111/j.1600-0889.2006.00209.x, 2006.

3160 Mulholland, M. R., Bernhardt, P. W., Heil, C. A., Bronk, D. A., and O'Neil, J. M.: Nitrogen
 3161 fixation and release of fixed nitrogen by *Trichodesmium* spp. in the Gulf of Mexico. *Limnol.*
 3162 *Oceanogr* 51, 1762–1776, doi: 10.4319/lo.2006.51.4.1762, 2006.

3163 Müller, C., Iinuma, Y., Karstensen, J., van Pinxteren, D., Lehmann, S., Gnauk, T., and
 3164 Herrmann, H.: Seasonal variation of aliphatic amines in marine sub-micrometer particles at the
 3165 Cape Verde islands, *Atmos. Chem. Phys.*, 9, 9587–9597, doi:10.5194/acp-9-9587-2009, 2009.

3166
 3167 Murphy, S. M., Sorooshian, A., Kroll, J. H., Ng, N. L., Chhabra, P., Tong, C., Surratt, J. D.,
 3168 Knipping, E., Flagan, R. C., and Seinfeld, J. H.: Secondary aerosol formation from atmospheric
 3169 reactions of aliphatic amines, *Atmos. Chem. Phys.*, 7, 2313–2337, doi:10.5194/acp-7-2313-
 3170 2007, 2007.

3171
 3172 Neff, J. C., Holland, E. A., Dentener, F. J., McDowell, W. H., and Russell, K. M.: The origin,
 3173 composition and rates of organic nitrogen deposition: a missing piece of the nitrogen cycle?,
 3174 *Biogeochemistry*, 57/58, 99–136, doi: 10.1023/A:1015791622742, 2002.

3175 Neirynck, J., Kowalski, A. S., Carrara, A., Genouw, G., Berghmans, P., and Ceulemans, R.:
 3176 Fluxes of oxidised and reduced nitrogen above a mixed coniferous forest exposed to various
 3177 nitrogen emission sources, *Environ. Pollut.*, 149, 31–43, doi: 10.1016/j.envpol.2006.12.029,
 3178 2007.

3179 Nemitz, E., Sutton, M. A., Wyers, G. P., Otjes, R. P., Schjoerring, J. K., Gallagher, M. W.,
 3180 Parrington, J., Fowler, D., and Choularton, T. W.: Surface/atmosphere exchange and chemical
 3181 interaction of gases and aerosols over oilseed rape. *Agr. Forest Meteorol.*, 105, 427–445, doi:
 3182 10.1016/S0168-1923(00)00207-0, 2000.

3183 Nemitz, E., Dorsey, J. R., Flynn, M. J., Gallagher, M. W., Hensen, A., Erisman, J.-W., Owen,
 3184 S. M., Dämmgen, U., and Sutton, M. A.: Aerosol fluxes and particle growth above managed
 3185 grassland, *Biogeosciences*, 6, 1627–1645, doi: 10.5194/bg-6-1627-2009, 2009.

3186 Nemitz, E., Langford, B., Ryder, J., Vieno, M., Sutton, M.A.: Ammonium nitrate deposition
 3187 modeling – accounting for evaporation during the deposition process, Presentation OP06-6,
 3188 International Aerosol Conference, Busan, South Korea, 28 Aug – 2 Sept, 2014.

3189 Nguyen, T. B., Lee, P. B., Updyke, K. M., Bones, D. L., Laskin, J., Laskin, A., and Nizkorodov,
 3190 S. A.: Formation of nitrogen- and sulfur-containing light-absorbing compounds accelerated by
 3191 evaporation of water from secondary organic aerosols, *J. Geophys. Res.*, 117, D01207, doi:
 3192 10.1029/2011JD016944, 2012.

3193 Noble, A. E., Lamborg, C. H., Ohnemus, D. C., Lam, P. J., Goepfert, T. J., Measures, C. I.,
 3194 Frame, C. H., Casciotti, K. L., Ditullio, G. R., Jennings, J., and Saito, M.A.: Basin-scale inputs
 3195 of cobalt, iron, and manganese from the Benguela-Angola front to the South Atlantic Ocean,
 3196 *Limnol. Oceanogr.*, 57, 989–1010, doi: 10.4319/lo.2012.57.4.0989, 2012.

3197 Norby, R. J., Warren, J. M., Iversen, C. M., Medlyn, B. E., and McMurtrie, R. E.: CO₂
 3198 enhancement of forest productivity constrained by limited nitrogen availability, *P. Natl. Acad.*

3199 Sci. USA, 107, 19368–19373, doi:10.1073/pnas.1006463107 and doi:
3200 10.1073/pnas.1006463107/-DCSupplemental, 2010.

3201 Nozière, B., Dziedzic, P., and Córdova, A.: Formation of secondary light-absorbing “fulvic-
3202 like” oligomers: A common process in aqueous and ionic atmospheric particles? *Geophys. Res.*
3203 *Lett.*, 34, L21812, doi: 10.1029/2007GL031300, 2007.

3204

3205 Nozière, B., Dziedzic, P., and Córdova, A.: Products and kinetics of the liquid-phase reaction
3206 of glyoxal catalyzed by ammonium ions (NH₄⁺), *J. Phys. Chem.*, 113, 1, 231–237, doi:
3207 10.1021/jp8078293, 2009.

3208

3209 O'Brien, R. E., Laskin, A., Laskin, J., Liu, S., Weber, R., Russell, L. M., Goldstein, A. H.:
3210 Molecular characterization of organic aerosol using nanospray desorption/electrospray
3211 ionization mass spectrometry: CalNex 2010 field study, *Atmos. Environ.*, 68, 265–272, doi:
3212 10.1016/j.atmosenv.2012.11.056, 2013.

3213 Oswald, R., Behrendt, T., Ermel, M., Wu, D., Su, H., Cheng, Y., Breuninger, C., Moravek, A.,
3214 Mougin, E., Delon, C., Loubet, B., Pommerening-Röser, A., Sörgel, M., Pöschl, U., Hoffmann,
3215 T., Andreae, M.O., Meixner, F.X., Trebs, I.: HONO emissions from soil bacteria as a major
3216 source of atmospheric reactive nitrogen, *Science*, 341, 1233–1235,
3217 doi:10.1126/science.1242266, 2013.

3218 Paerl, H. W., Crocker, K. M., and Prufert, L. E.: Limitation of N₂ fixation in coastal marine
3219 waters: Relative importance of molybdenum, iron, phosphorus, and organic matter availability,
3220 *Limnol. Oceanogr.*, 32, 525–536, doi: 10.4319/lo.1987.32.3.0525, 1987.

3221 Palmroth, S., Oren, R., McCarthy, H. R., Johnsen, K. H., Finzi, A. C., Butnor, J. R., Ryan, M.
3222 G., and Schlese, U.: Aboveground sink strength in forests controls the allocation of carbon
3223 below ground and its [CO₂]-induced enhancement., *P. Natl. Acad. Sci. USA.*, 103, 19362–
3224 19367, doi:10.1073/pnas.0609492103, 2006.

3225 Parrish, D. D., Dunlea, E. J., Atlas, E. L., Schauffler, S., Donnelly, S., Stroud, V., Goldstein, A.H.,
3226 Millet, D. B., McKay, M., Jaffe, D.A., Price, H. U., Hess, P. G., Flocke, F., and Roberts, J. M.: Changes
3227 in the photochemical environment of the temperate North Pacific troposphere in response to
3228 increased Asian emissions, J. Geophys. Res.-Atmos., 109, D23S18, doi:
3229 10.1029/2004JD004978, 2004.

3230

3231 Parrish, D. D., Dunlea, E. J., Atlas, E. L., Schauffler, S., Donnelly, S., Stroud, V., Goldstein, A. H.,
3232 Millet, D. B., McKay, M., Jaffe, D. A., Price, H. U., Hess, P. G., Flocke, F., and Roberts, J. M.: Changes
3233 in the photochemical environment of the temperate North Pacific troposphere in response to
3234 increased Asian emissions, JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 109,
3235 D23S18, doi:10.1029/2004JD004978, 2004

3236

3237 Park, S., Croteau, P., Boering, K. A., Etheridge, D. M., Ferretti, D., Fraser, P. J., Kim, K.-R.,
3238 Krummel, P. B., Langenfelds, R. L., van Ommen, T. D., Steele, L. P., and Trudinger, C. M.:

Formatted: Left, Space After: 0 pt, Line spacing: single, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

- 3239 Trends and seasonal cycles in the isotopic composition of nitrous oxide since 1940, *Nat.*
3240 *Geosci.*, 5, 261-265, doi:10.1038/ngeo1421, 2012.
- 3241 Paulsen, D. M., Paerl, H. W., and Bishop, P. E.: Evidence that molybdenum-dependent
3242 nitrogen fixation is not limited by high sulfate concentrations in marine environments, *Limnol.*
3243 *Oceanogr.*, 36, 1325–1334, doi:10.4319/lo.1991.36.7.1325, 1991.
- 3244 Paumen, M. L., de Voogt, P., van Gestel, C. A. M., and Kraak, M. H. S.: Comparative chronic
3245 toxicity of homo- and heterocyclic aromatic compounds to benthic and terrestrial invertebrates:
3246 generalizations and exceptions, *Sci. Total Environ.*, 407, 4605-4609, doi:
3247 10.1016/j.scitotenv.2009.04.039, 2009.
- 3248
- 3249 Peierls, B. L. and Paerl, H. W.: Bioavailability of atmospheric organic nitrogen deposition to
3250 coastal phytoplankton, *Limnol. Oceanogr.*, 42, 1819–1823,
3251 doi:10.4319/lo.1997.42.8.1819, 1997.
- 3252 [Phillips, G.J., Pouvesle, N., Thieser, J., Schuster, G., Axinte, R., Fischer, H., Williams, J.,](#)
3253 [Lelieveld, J., and Crowley, J.N.: Peroxyacetyl nitrate \(PAN\) and peroxyacetic acid \(PAA\)](#)
3254 [measurements by iodide chemical ionisation mass spectrometry: first analysis of results in the](#)
3255 [boreal forest and implications for the measurement of PAN fluxes, *Atmos. Chem. Phys.*, 13,](#)
3256 [1129–1139, doi:10.5194/acp-13-1129-2013, 2013.](#)
- 3257
- 3258 [Phillips, G.J., Pouvesle, N., Thieser, J., Schuster, G., Axinte, R., Fischer, H., Williams, J.,](#)
3259 [Lelieveld, J., and Crowley, J.N.: Peroxyacetyl nitrate \(PAN\) and peroxyacetic acid \(PAA\)](#)
3260 [measurements by iodide chemical ionisation mass spectrometry:](#)
3261 [first analysis of results in the boreal forest and implications for the](#)
3262 [measurement of PAN fluxes - *Atmos. Chem. Phys.*, 13, 1129–1139, 2013](#)
3263 [www.atmos-chem-phys.net/13/1129/2013/](#)
3264 [doi:10.5194/acp-13-1129-2013](#)
- 3265 [Pilegaard, K.: Processes regulating nitric oxide emissions from soils, *Philos. T. Roy. Soc. B,*](#)
3266 [368, 1621, 20130112, doi: 10.1098/rstb.2013., 2013.](#)
- 3267 Pinay, G., Gumiero, B., Tabacchi, E., Gimenez, O., Tabacchi-Planty, A. M., Hefting, M. M.,
3268 Burt, T. P., Black, V. A., Nilsson, C., Iordache, V., Bureau, F., Vought, L., Petts, G. E., and
3269 Décamps, H.: Patterns of denitrification rates in European alluvial soils under various
3270 hydrological regimes, *Freshwater Biol.*, 52, 252-266, doi: 10.1111/j.1365-2427.2006.01680.x,
3271 2007.
- 3272 [Porada, P., Weber, B., Elbert, W. Pöschl, U., and Kleidon, A.: Estimating impacts of lichens](#)
3273 [and bryophytes on global biogeochemical cycles, *Global Biogeochem. Cy.*, 28, 71–85, doi:](#)
3274 [10.1002/2013GB004705, 2014.](#)

Commented [MKE3]: Phillips et al reference does not appear in the text - it was in a paragraph since deleted by David. Check with David and if OK delete this reference from the reference list.

3276 Powelson, M. H., Espelien, B. M., Hawkins, L. N., Galloway, M. M., and De Haan, D. O.:
 3277 Brown carbon formation by aqueous-phase carbonyl compound reactions with amines and
 3278 ammonium sulfate, *Environ. Sci. Technol.*, 48, 2, 985-993, doi: 10.1021/es4038325, 2014.

3279 Prather, M. J. and Ehhalt, D.: Atmospheric chemistry and greenhouse gases, in: *Climate*
 3280 *Change 2001: The scientific basis. Contribution of Working Group I to the Third Assessment*
 3281 *Report of the Intergovernmental Panel on Climate Change*, Houghton, J.T., Ding, Y., Griggs,
 3282 D. J., Noguer, M., van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A. (Eds.),
 3283 Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, Cambridge,
 3284 United Kingdom and New York, NY, USA, Chapter 4, pp.239-287, 2001.

3285 Prospero, J. M. and Lamb, P. J.: African droughts and dust transport to the Caribbean: Climate
 3286 change implications, *Science*, 302, 5647, 1024–1027, doi:10.1126/science.1089915, 2003.

3287 Pye, H. O. T., Liao, H., Wu, S., Mickley, L. J., Jacob, D. J., Henze, D. K., and Seinfeld, J.H.:
 3288 Effect of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels
 3289 in the United States, *J. Geophys. Res.*, 114, D01205, doi: 10.1029/2008JD010701, 2009.

3290 Rastogi, N., Zhang, X. L., Edgerton, E. S., Ingall, E., and Weber, R. J.: Filterable water-soluble
 3291 organic nitrogen in fine particles over the southeastern USA during summer, *Atmos. Environ.*,
 3292 45, 33, 6040-6047, doi: 10.1016/j.atmosenv.2011.07.045, 2011.

3293 Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous oxide (N₂O): the dominant
 3294 ozone-depleting substance emitted in the 21st century, *Science*, 326, 5949, 123-12,
 3295 doi:10.1126/science.1176985, 2009.

3296
 3297
 3298 Reeve, N., and Toumi, R.: Lightning activity as an indicator of climate change, *Q. J. Roy.*
 3299 *Meteor. Soc.*, 125, 893-903, doi: 10.1002/qj.49712555507, 1999.

3300 Riddick, S. N., Dragosits, U., Blackall, T. D., Daunt, F., Wanless, S., and Sutton, M. A.: The
 3301 global distribution of ammonia emissions from seabird colonies, *Atmos. Environ.*, 55, 319-
 3302 327, doi:10.1016/j.atmosenv.2012.02.052, 2012.

3303 Riedo, M., Milford, C., Schmid, M., and Sutton, M. A.: Coupling soil-plant-atmosphere
 3304 exchange of ammonia with ecosystem functioning in grasslands, *Ecol. Model.*, 158, 83–110,
 3305 doi: 10.1016/S0304-3800(02)00169-2, 2002.

3306 Rincon, A. G., Calvo, A. I., Dietzel, M., and Kalberer, M.: Seasonal differences of urban
 3307 organic aerosol composition - an ultra-high resolution mass spectrometry study, *Environ.*
 3308 *Chem.*, 9, 3, 298-319, doi: 10.1071/EN12016, 2012.

3309 Roberts, J.M.: The atmospheric chemistry of organic nitrates,. *Atmos. Environ. A-Gen.*, 24, 2,
 3310 243-287, doi: 10.1016/0960-1686(90)90108-Y, 1990.
 3311
 3312

3313 [Roberts, J.M. THE ATMOSPHERIC CHEMISTRY OF ORGANIC NITRATES, *Atmospheric Environment*](#)
 3314 [Vol. 24A, No. 2, pp. 243-287, 1990-Romps, D.M., Seeley, J.T., Vollaro, D., and Molinari, J.](#)
 3315 [Projected increase in lightning strikes in the United States due to global warming, *Science*. 346](#)
 3316 [\(6211\), pp 851-853, doi: 10.1126/science.1259100, 2014.](#)

3317

3318

3319 Rosa, S. M., Kraemer, F. B., Soria, M. A., Guerrero, L. D., Morrás, H. J. M., Figuerola, E. L.
 3320 M., and Erijman, L.: The influence of soil properties on denitrifying bacterial communities and
 3321 denitrification potential in no-till production farms under contrasting management in the
 3322 Argentinean Pampas. *Appl. Soil Ecol.*, 75, 172-180, doi: 10.1016/j.apsoil.2013.11.012, 2014.

3323 Russell, K. M., Keene, W. C., Maben, J. R., Galloway, J.N., and Moody, J. L.: Phase
 3324 partitioning and dry deposition of atmospheric nitrogen at the mid-Atlantic US coast, *J.*
 3325 *Geophys. Res.*, 108, D21, 4656, doi: 10.1029/2003JD003736, 2003.

3326 Rustad, L. E., Campbell, J. L., Marion, G. M., Norby, R. J., Mitchell, M. J., Hartley, A. E.,
 3327 Cornelissen, J. H. C., and Gurevitch, J.: A meta-analysis of the response of soil respiration, net
 3328 nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming,
 3329 *Oecologia* 126, 543–562, doi:10.1007/s004420000544, 2001.

3330 Ryder, J.: Emission, deposition and chemical conversion of atmospheric trace substances in
 3331 and above vegetation canopies, PhD thesis, University of Manchester, Manchester, 2010.

3332 Sarmiento, J. L., Slater, R., Barber, R., Bopp, L., Doney, S. C., Hirst, A. C., Kleypas, J., Matear,
 3333 R., Mikolajewicz, U., Monfray, P., Soldatov, V., Spall, S. A., and Stouffer, R.: Response of
 3334 ocean ecosystems to climate warming, *Global Biogeochem. Cy.*, 18, GB3003,
 3335 doi:10.1029/2003GB002134, 2004.

3336 Saxena, P., and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical
 3337 review of the literature and application of thermodynamics to identify candidate compounds,
 3338 *J. Atmos. Chem.*, 24, 57-109, doi: 10.1007/BF00053823, 1996.

3339

3340 Scalabrin, E., Zangrando, R., Barbaro, E., Kehrwald, N. M., Gabrieli, J., Barbante, C., and
 3341 Gambaro, A.: Amino acids in Arctic aerosols, *Atmos. Chem. Phys.*, 12, 10453-10463, doi:
 3342 10.5194/acp-12-10453-2012, 2012.

3343 Schreiber, F., Wunderlin, P., Udert, K. M., and Wells, G.F.: Nitric oxide and nitrous oxide
 3344 turnover in natural and engineered microbial communities: biological pathways, chemical
 3345 reactions, and novel technologies, *Front. Microbiol.*, 3, 372, doi: 10.3389/fmicb.2012.00372,
 3346 2012.

3347 Schumann, U. And Huntrieser, H.: The global lightning-induced nitrogen oxides source,
 3348 *Atmos. Chem. Phys.*, 7, 3823-3907, doi: 10.5194/acp-7-3823-2007, 2007.

Formatted: Font: 12 pt

Formatted: Left

3349 Seinfeld, J. H. and Pandis, S. N.: Atmospheric physics and chemistry, 1st edition, John Wiley
 3350 & Sons, 1998.

3351 Seitzinger, S. P. and Sanders, R. W.: Atmospheric inputs of dissolved organic nitrogen
 3352 stimulate estuarine bacteria and phytoplankton, *Limnol. Oceanogr.*, 44, 721–730,
 3353 doi:10.4319/lo.1999.44.3.0721, 1999.

3354 Seitzinger, S., Harrison, J. A., Bohlkem J. K., Bouwman, A. F., Lowrance, R., Peterson, B.,
 3355 Tobias, C., and Van Drecht, G., 2006, Denitrification across landscapes and waterscapes: a
 3356 synthesis, *Ecol. Appl.*, 16, 2064–2090, doi:10.1890/1051-
 3357 0761(2006)016[2064:DALAWA]2.0.CO;2, 2006

3358 Shapiro, E. L., Szprengiel, J., Sareen, N., Jen, C. N., Giordano, M. R., and McNeill, V. F.:
 3359 Light-absorbing secondary organic material formed by glyoxal in aqueous aerosol mimics,
 3360 *Atmos. Chem. Phys.*, 9, 2289–2300, doi:10.5194/acp-9-2289-2009, 2009.

3361 Sharp, R. G. and Davies, W. J.: Variability among species in the apoplastic pH signalling
 3362 response to drying soils, *J. Exp. Bot.*, 60, 4363–4370, doi:10.1093/jxb/erp273, 2009.

3363 Shen, J., Cui, Z., Miao, Y., Mi, G., Zhang, H., Fan, M., Zhang, C., Jiang, R., Zhang, W., Li,
 3364 H., Chen, X., Li, X. and Zhang, F.: Transforming agriculture in China: From solely high yield
 3365 to both high yield and high resource use efficiency, *Global Food Security*, 2, 1-8, doi:
 3366 10.1016/j.gfs.2012.12.004, 2013.

3367 Shi, D., Xu, Y., Hopkinson, B. M., Morel, F. M. M.: Effect of ocean acidification on iron
 3368 availability to marine phytoplankton, *Science*, 327, 5966, 676–679,
 3369 doi:10.1126/science.1183517, 2010.

3370 Shi, J. H., Han, J., Fan, D. G., Qi, J. H., and Gao, H. W.: Contribution of water soluble organic
 3371 nitrogen to total nitrogen in atmospheric aerosols in Qingdao, *Huanjing Kexue*, 32, 1, 1-8,
 3372 2011.

3373

3374 Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard,
 3375 C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena,
 3376 V. S., Tsyro, S., Tuovinen, J.-P., Valdebenito, A., and Wind, P.: The EMEP MSC-W chemical
 3377 transport model – technical description, *Atmos. Chem. Phys.*, 12, 7825–7865, doi:10.5194/acp-
 3378 12-7825-2012, 2012.

3379 Simpson, D., Andersson, C., Christensen, J. H., Engardt, M., Geels, C., Nyíri, A., Posch, M.,
 3380 Soares, J., Sofiev, M., Wind, P., and Langner, J.: Impacts of climate and emission changes on
 3381 nitrogen deposition in Europe: a multi-model study, *Atmos. Chem. Phys. Discuss.*, 14, 6663–
 3382 6720, doi: 10.5194/acpd-14-6663-2014, 2014.

3383 Singh, B. K., Bardgett, R. D., Smith, P., and Reay, D. S.: Microorganisms and climate change:
 3384 terrestrial feedbacks and mitigation options, *Nat. Rev. Microbiol.*, 8, 779-790, doi:
 3385 10.1038/nrmicro2439, 2010.

Field Code Changed

3386 Singh, H. B. and Hanst, P. L.: Peroxyacetyl nitrate (PAN) in the unpolluted atmosphere: An
 3387 important reservoir for nitrogen oxides, *Geophys. Res. Lett.*, 8, 941-944, doi:
 3388 10.1029/GL008i008p00941, 1981.

3389 Sitch, S., Cox, P. M., Collins, W. J., and Huntingford, C.: Indirect radiative forcing of climate
 3390 change through ozone effects on the land-carbon sink, *Nature*, 448, 791–795,
 3391 doi:10.1038/nature06059, 2007.

3392 Skiba, U., van Dijk, S., and Ball, B. C: The influence of tillage on NO and N₂O fluxes under
 3393 spring and winter barley, *Soil Use Manage.*, 18, 340-345, doi: 10.1111/j.1475-
 3394 2743.2002.tb00250.x, 2002.

3395 Smil, V.: Nitrogen in crop production: An account of global flows. *Global Biogeochem. Cy.*,
 3396 13, 647-662, doi: 10.1029/1999GB900015, 1999.

3397 Smith, J. N., Barsanti, K. C., Friedli, H. R., Ehn, M., Kulmala, M., Collins, D. R., Scheckman,
 3398 J. H., Williams, B. J., and McMurry, P. H.: Observations of aminium salts in atmospheric
 3399 nanoparticles and possible climatic implications, *P. Natl. Acad. Sci. USA*, 107, 15, 6634–6639,
 3400 doi: 10.1073/pnas.0912127107, 2010.

3401 Sokolov, A. P., Kicklighter, D. W., Melillo, J. M., Felzer, B. S., Schlosser, C. A., and Cronin,
 3402 T. W.: Consequences of considering carbon-nitrogen interactions on the feedbacks between
 3403 climate and the terrestrial carbon cycle. *J. Climate* 21, 3776–3796.
 3404 doi:10.1175/2008JCLI2038.1, 2008.

3405 Spokes, L. J., Yeatman, S. G., Cornell, S. E., and Jickells, T. D.: Nitrogen deposition to the
 3406 eastern Atlantic Ocean. The importance of south-easterly flow, *Tellus B*, 52, 37–49, doi:
 3407 10.1034/j.1600-0889.2000.00062.x, 2000.

3408

3409 Srinivas, B., Sarin, N. M., and Sarma, V. V. S. S.: Atmospheric dry deposition of inorganic
 3410 and organic nitrogen to the Bay of Bengal: Impact of continental outflow, *Mar. Chem.*, 127,
 3411 170–179, doi:10.1016/j.marchem.2011.09.002, 2011.

3412 Staal, M., Meysman, F. J. R., and Stal, L. J.: Temperature excludes N₂-fixing heterocystous
 3413 cyanobacteria in the tropical oceans, *Nature*, 425, 504–507, doi: 10.1038/nature01999, 2003.

3414 Steadman, C. E., Stevenson, D. S., Fowler, D., Heal, M., and Sutton, M.: Ammonia emissions
 3415 from the ocean in the 21st century – the effect of rising temperatures and CO₂ levels on the
 3416 exchange of ammonia between the atmosphere and oceans, 2014 (in preparation).

3417 Stevenson, D. S., Dentener, F. J., Schultz, M. G, Ellingsen, K., van Noije, T. P. C., Wild, O.,
 3418 Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J.,
 3419 Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss,
 3420 M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F.,
 3421 Lawrence, M. G., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S.,
 3422 Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K.,

and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, 111, D8, D08301, doi: 10.1029/2005JD006338, 2006.

Stevenson, F. J., Harrison, R. M., Wetselaar, R., and Leeper, R. A.: Nitrosation of soil organic matter: III. Nature of gases produced by reaction of nitrite with lignins, humic substances, and phenolic constituents under neutral and slightly acidic conditions, *Soil Sci. Soc. Am. J.*, 34, 3, 430-435, doi:10.2136/sssaj1970.03615995003400030024x, 1970.

Stocker, B. D., Roth, R., Joos, F., Spahni, R., Steinacher, M., Zaehle, S., Bouwman, L., Xu-Ri, and Prentice, I. C.: Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios. *Nat. Clim. Change*, 3, 666–672, doi:10.1038/nclimate1864, 2013.

Su, H., Cheng, Y., Oswald, R., Behrendt, T., Trebs, I., Meixner, F.X., Andreae, M.O., Cheng, P., Zhang, Y., and Pöschl, U.: Soil Nitrite as a Source of Atmospheric HONO and OH Radicals, *Science*, 333 (6049), 1616-1618, doi: 10.1126/science.1207687, 2011.

Sun, Y. L., Wang, Z. F., Fu, P. Q., Yang, T., Jiang, Q., Dong, H. B., Li, J., and Jia, J. J.: Aerosol composition, sources and processes during wintertime in Beijing, China, *Atmos. Chem. Phys.*, 13, 4577–4592, doi:10.5194/acp-13-4577-2013, 2013.

Sutton, M. A., Schjørring, J. K., Wyers, G. P., Duyzer, J. H., Ineson, P., and Powlson, D. S.: Plant-atmosphere exchange of ammonia, *Philos. T. Roy. Soc. A*, 351, 1696, 261-275, doi: 10.1098/rsta.1995.0033, 1995.

Sutton, M. A., Burkhardt, J. K., Guerin, D., Nemitz, E., and Fowler, D.: Development of resistance models to describe measurements of bi-directional ammonia surface atmosphere exchange, *Atmos. Environ.*, 32, 3, 473-480, doi: 10.1016/S1352-2310(97)00164-7, 1998.

Sutton, M. A., Nemitz, E., Milford, C., Campbell, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M., Dorsey, J. R., Gallagher, M. W., Horvath, L., Weidinger, T., Meszaros, R., Dämmgen, U., Neftel, A., Herrmann, B., Lehman, B. E., Flechard, C., and Burkhardt, J.: Dynamics of ammonia exchange with cut grassland: synthesis of results and conclusions of the GRAMINAE Integrated Experiment, *Biogeosciences*, 6, 2907–2934, doi:10.5194/bg-6-2907-2009, 2009.

Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., Grizzetti, B. (Eds.): *The European Nitrogen Assessment: Sources, effects and policy perspectives*, Cambridge University Press, 2011.

Sutton, M. A., Bleeker, A., Howard, C. M., Bekunda, M., Grizzetti, B., de Vries, W., van Grinsven, H. J. M., Abrol, Y. P., Adhya, T. K., Billen, G., Davidson, E. A., Datta, A., Diaz, R., Erisman, J. W., Liu, X. J., Oenema, O., Palm, C., Raghuram, N., Reis, S., Scholz, R. W., Sims, T., Westhoek, H., and Zhang F. S.: ~~(2013)~~ Our Nutrient World: The challenge to produce more food and energy with less pollution, *Global Overview of Nutrient Management*, Centre for

Formatted: Font color: Auto

Ecology & Hydrology, Edinburgh on behalf of the Global Partnership on Nutrient Management and the International Nitrogen Initiative, 114 pp., 2013a.

Sutton, M. A., Reis, S., Riddick, S. N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S., Braban, C. F., Vieno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall, T. D., Milford, C., Flechard, C. R., Loubet, B., Massad, R. S., Cellier, P., Personne, E., Coheur, P. F., Clarisse, L., Van Damme, M., Ngadi, Y., Clerbaux, C., Skj  th, C. A., Geels, C., Hertel, O., Wichink Kruit, R. J., Pinder, R. W., Bash, J. O., Walker, J. T., Simpson, D., Horv  th, L., Misselbrook, T. H., Bleeker, A., Dentener, F., and de Vries, W.: Towards a climate-dependent paradigm of ammonia emission and deposition, *Philos. T. Roy. Soc. B*, 368, 1621, 20130166, doi:10.1098/rstb.2013.0166, 2013b.

Syakila, A. and Kroeze, C.: The global nitrous oxide budget revisited, *Greenhouse Gas Measurement & Management*, 1, 1, 17-26, doi: 10.3763/ghgmm.2010.0007, 2011.

Thomas, R.: Measurement of speciated aerosol fluxes, Ph.D thesis, University of Manchester, Manchester, 2007.

Thornton, P. E., Doney, S. C., Lindsay, K., Moore, J. K., Mahowald, N., Randerson, J. T., Fung, I., Lamarque, J.-F., Feddes, J. J., and Lee, Y.-H.: Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model, *Biogeosciences*, 6, 2099–2120, doi:10.5194/bg-6-2099-2009, 2009.

Timperley, M. H., Vigor-Brown, R. J., Kawashima, M., and Ishigami, M.: Organic nitrogen compounds in atmospheric precipitation: ~~their~~ their chemistry and availability to phytoplankton, *Can. J. Fish. Aquat. Sci.*, 42, 1171–1177, doi: 10.1139/f85-145, 1985.

Trost, B., Prochnow, A., Drastig, K., Meyer-Aurich, A., Ellmer, F., and Baumecker, M.: Irrigation, soil organic carbon and N₂O emissions. A review, *Agron. Sustain. Dev.*, 33, 733-749, doi: 10.1007/s13593-013-0134-0, 2013.

Tuomi, R., Haigh, J. D., and Law, K.: A tropospheric ozone-lightning climate feedback, *Geophys. Res. Lett.*, 23, 9, 1037-1040, doi: 10.1029/96GL00944, 1996.

Tuovinen, J.-P., Hakola, H., Karlsson, P. E., and Simpson, D.: Air pollution risks to northern European forests in a changing climate, in: *Air pollution and global Challenges. Understanding and perspectives from forest research*, Matyssek, R., Clarke, N., Cudlin, P., Mikkelsen, T., Tuovinen, J.-P., Wieser, G., and Paoletti, E. (Eds.), Elsevier, Oxford, UK, Chapter 5, 77 – 99, 2013.

Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., and Coheur, P. F.: Global distributions, time series and error characterization of atmospheric ammonia (NH₃) from IASI satellite observations, *Atmos. Chem. Phys.*, 14, 2905-2922, doi:10.5194/acp-14-2905-2014, 2014.

[Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., and Coheur, P. F. 2014 Global distributions, time series and error characterization of atmospheric ammonia \(NH₃\) from IASI satellite observations, *Atmos. Chem. Phys.*, **14**, 2905–2922, doi:10.5194/acp-14-2905-2014](#)

van Groenigen, K. J., Osenberg, C. W., and Hungate, B. A.: Increased soil emissions of potent greenhouse gases under increased atmospheric CO₂, *Nature*, **475**, 214–216, doi:10.1038/nature10176, 2011.

van Vuuren, D. P., Bouwman, A. F., Smith, S. J., and Dentener, F.: Global projections for anthropogenic reactive nitrogen emissions to the atmosphere: an assessment of scenarios in the scientific literature, *Curr. Opin. Environ. Sustain.*, **3**, 359–369, doi: 10.1016/j.cosust.2011.08.014, 2011a.

van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Matsui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Climatic Change*, **109**, 5–31, doi:10.1007/s10584-011-0148-z, 2011b.

Velikova, V., Pinelli, P., Pasqualini, S., Reale, L., Ferranti, F., and Loreto, F.: Isoprene decreases the concentration of nitric oxide in leaves exposed to elevated ozone. *New Phytologist*, **166**, 419–426, doi: 10.1111/j.1469-8137.2005.01409.x, 2005.

Vieno, M., Heal, M. R., Hallsworth, S., Famulari, D., Doherty, R. M., Dore, A. J., Tang, Y. S., Braban, C. F., Leaver, D., Sutton, M. A., and Reis, S.: The role of long-range transport and domestic emissions in determining atmospheric secondary inorganic particle concentrations across the UK, *Atmos. Chem. Phys.*, **14**, 8435–8447, doi:10.5194/acp-14-8435-2014, 2014.

Vile, M. A., Wieder, R. K., Živković, T., Scott, K. D., Vitt, D. H., Hartsock, J. A., Iosue, C. L., Quinn, J. C., Petrix, M., Fillingim, H. M., Popma, J. M. A., Dynarski, K. A., Jackman, T. R., Albright, C. M., and Wyckoff, D. D.: N₂-fixation by methanotrophs sustains carbon and nitrogen accumulation in pristine peatlands, *Biogeochemistry*, **121**, 317–328, doi: 10.1007/s10533-014-0019-6, 2014.

Violaki, K. and Mihalopoulos, N.: Water-soluble organic nitrogen (WSON) in size-segregated atmospheric particles over the Eastern Mediterranean, *Atmos. Environ.*, **44**, 4339–4345, doi:10.1016/j.atmosenv.2010.07.056, 2010.

Violaki, K. and Mihalopoulos, N.: Urea: An important piece of Water Soluble Organic Nitrogen (WSON) over the Eastern Mediterranean, *Sci. Total Environ.*, **409**, 22, 4796–4801, doi: 10.1016/j.scitotenv.2011.07.060, 2011.

Vitousek, P. M., Cassman, K., Cleveland, C., Crews, T., Field, C. B., Grimm, N. B., Howarth, R. W., Marino, R., Martinelli, L., Rastetter, E. B., and Spret, J. I.: Towards an ecological understanding of biological nitrogen fixation, *Biogeochemistry*, **57/58**, 1–45, doi: 10.1023/A:1015798428743, 2002.

3533 Vitousek, P. M., Menge, D. N., Reed, S. C., and Cleveland, C. C.: Biological nitrogen fixation:
 3534 rates, patterns and ecological controls in terrestrial ecosystems, *Philos. T. Roy. Soc. B*, 368,
 3535 1621, 20130119, doi: 10.1098/rstb.2013.0119, 2013.

3536 Voss, M., Bange, H. W., Dippner, J. W., Middelburg, J. J., Montoya, J. P., and Ward, B.: The
 3537 marine nitrogen cycle: recent discoveries, uncertainties and the potential relevance of climate
 3538 change, *Philos. T. Roy. Soc. B*, 368, 1621, 20130121, doi: 10.1098/rstb.2013.0121, 2013.

3539 Walker, J. T., Jones, M. R., Bash, J. O., Myles, L., Meyers, T., Schwede, D., Herrick, J.,
 3540 Nemitz, E., and Robarge, W.: Processes of ammonia air-surface exchange in a fertilized Zea
 3541 mays canopy, *Biogeosciences*, 10, 981–998, doi:10.5194/bg-10-981-2013, 2013.

3542 Wang, Y., Jacob, D. J., and Logan, J. A.: Global simulation of tropospheric O₃-NO_x-
 3543 hydrocarbon chemistry: 1. Model formulation, *J. Geophys. Res.-Atmos.*, 103, D9, 10713–
 3544 10725, doi: 10.1029/98JD00158, 1998.

3545 Westhoek, H., Lesschen, J. P., Rood, T., Wagner, S., De Marco, A., Murphy-Bokern, D., Leip,
 3546 A., van Grinsven, H., Sutton, M. A., and Oenema, O.: Food choices, health and environment:
 3547 Effects of cutting Europe's meat and dairy intake, *Global Environ. Chan.*, 26, 196-205,
 3548 doi:10.1016/j.gloenvcha.2014.02.004, 2014.

3549 Wichink Kruit, R. J., Schaap, M., Sauter, F. J., van Zanten, M. C., and van Pul, W. A. J.:
 3550 Modeling the distribution of ammonia across Europe including bi-directional surface-
 3551 atmosphere exchange, *Biogeosciences*, 9, 5261-5277, doi: 10.5194/bg-9-5261-2012, 2012.

3552 Wilkinson, S. and Davies, W. J.: Manipulation of the apoplastic pH of intact plants mimics
 3553 stomatal and growth responses to water availability and microclimatic variation, *J. Exp. Bot.*,
 3554 59, 3, 619–631, doi: 10.1093/jxb/erm338, 2008.

3555 Williams, E.: Lightning and climate: A review, *Atmos. Res.*, 76, 272-287,
 3556 doi:10.1016/j.atmosres.2004.11.014, 2005.

3557 Williams, J., Crowley, J., Fischer, H., Harder, H., Martinez, M., Petäijä, T., Rinne, J., Bäck,
 3558 J., Boy, M., Dal Maso, M., Hakala, J., Kajos, M., Keronen, P., Rantala, P., Aalto, J., Aaltonen,
 3559 H., Paatero, J., Vesala, T., Hakola, H., Levula, J., Pohja, T., Herrmann, F., Auld, J.,
 3560 Mesarchaki, E., Song, W., Yassaa, N., Nölscher, A., Johnson, A. M., Custer, T., Sinha, V.,
 3561 Thieser, J., Pouvesle, N., Taraborrelli, D., Tang, M. J., Bozem, H., Hosaynali Beygi, Z.,
 3562 Axinte, R., Oswald, R., Novelli, A., Kubistin, D., Hens, K., Javed, U., Trawny, K.,
 3563 Breitenberger, C., Hidalgo, P. J., Ebben, C. J., Geiger, F. M., Corrigan, A. L.,
 3564 Russell, L. M., Ouwersloot, H. G., Vilà-Guerau de Arellano, J., Ganzeveld, L., Vogel, A.,
 3565 Beek, M., Bayerle, A., Kampf, C. J., Bertelmann, M., Köllner, F., Hoffmann, T., Valverde,
 3566 J., González, D., Riekkola, M. L., Kulmala, M., and Lelieveld, J.: The summertime Boreal
 3567 forest field measurement intensive
 3568 (HUMPPA-COPEC-2010): an overview of meteorological and chemical influences, *Atmos.*
 3569 *Chem. Phys.*, 11, 10599–10618, doi:10.5194/acp-11-10599-2011, 2011.

3570 Winiwarter, W., Erisman, J. W., Galloway, J. N., Klimont, Z., Sutton, M. A.: Estimating
 3571 environmentally relevant fixed nitrogen demand in the 21st century. *Climatic Change*, 120,
 3572 889-901, doi: 10.1007/s10584-013-0834-0, 2013.

Formatted: Font: 12 pt

Formatted: Left, Space After: 0 pt, Line spacing: single, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Font: 12 pt

3573 Wittig, V. E., Ainsworth, E. A., and Long, S. P.: To what extent do current and projected
3574 increases in surface ozone affect photosynthesis and stomatal conductance of trees? A meta-
3575 analytic review of the last 3 decades of experiments, *Plant Cell Environ.*, 30, 1150–1162, doi:
3576 10.1111/j.1365-3040.2007.01717.x, 2007.

3577 Wolff, V., Trebs, I., Ammann, C., Spierig, C., Flechard, C., Neftel, A., and Meixner, F.X.:
3578 Concentrations and fluxes of soluble reactive nitrogen compounds over an intensively managed
3579 grassland site, *Geophys. Res. Abstr.*, 9, 02906, 2007.

3580 Wolff, V., Meixner, F.X., and Trebs, I.: Mixing ratios and exchange processes of the ammonia-
3581 nitric acid-ammonium nitrate triad above a spruce forest canopy, in: Proceedings of ‘Earth
3582 observations for land-atmosphere interaction science’, Lacoste-Francid, H. (Ed.), Frascati,
3583 Italy, ESA SP-688, European Space Agency, Noordwijk, 2011.

3584 [Wolfe, G. M., Thornton, J. A., Yatawelli, R. L. N., McKay, M., Goldstein, A. H.,](#)
3585 [LaFranchi, B., Min, K.-E., and Cohen, R. C.: Eddy covariance fluxes of acyl peroxy nitrates](#)
3586 [\(PAN, PPN and MPAN\) above a Ponderosa pine forest, *Atmos. Chem. Phys.*, 9, 615-634,](#)
3587 [doi:10.5194/acp-9-615-2009, 2009.](#)

3588 Xu-Ri and Prentice, I. C.: Terrestrial nitrogen cycle simulation with a dynamic global
3589 vegetation model, *Global Change Biol.*, 14, 1745–1764, doi:10.1111/j.1365-
3590 2486.2008.01625.x, 2008.

3591 Yienger, J. J. and Levy, H. II.: Empirical model of global soil-biogenic NO_x emissions, *J.*
3592 *Geophys. Res.*, 100, D6, 11447-11464, doi: 10.1029/95JD00370, 1995.

3593 Zaehle, S.: Terrestrial nitrogen–carbon cycle interactions at the global scale. *Philos. T. Roy.*
3594 *Soc. B*, 368, 1621, 20130125, doi:10.1098/rstb.2013.0125, 2013.

3595 Zaehle, S. and Dalmonech, D.: Carbon–nitrogen interactions on land at global scales: current
3596 understanding in modelling climate biosphere feedbacks. *Curr. Opin. Environ. Sustain.*, 3,
3597 311–320, doi:10.1016/j.cosust.2011.08.008, 2011.

3598 Zaehle, S., Friend, A. D., Friedlingstein, P., Dentener, F., Peylin, P., and Schulz, M.: Carbon
3599 and nitrogen cycle dynamics in the O-CN land surface model: 2. Role of the nitrogen cycle in
3600 the historical terrestrial carbon balance, *Global Biogeochem. Cy.*, 24, GB1006,
3601 doi:10.1029/2009GB003522, 2010a.

3602 Zaehle, S., Friedlingstein, P., and Friend, A. D.: Terrestrial nitrogen feedbacks may accelerate
3603 future climate change, *Geophys. Res. Lett.* 37, L01401, doi: 10.1029/2009GL041345, 2010b.

3604 Zaehle, S., Ciais, P., Friend, A. D., and Prieur, V.: Carbon benefits of anthropogenic reactive
3605 nitrogen offset by nitrous oxide emissions, *Nature Geosci.*, 4, 601–605, doi:10.1038/ngeo1207,
3606 2011.

3607 Zaehle, S., Medlyn, B. E., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hickler, T., Luo,
3608 Y., Wang, Y.-P., El-Masri, B., Thornton, P., Jain, A., Wang, E., Parton, W., Iverson, C. M.,
3609 Gallet-Budynek, A., McCarthy, H., Finzi, A., Hanson, P. J., Prentice, I. C., Oren, R., and

Formatted: Font: (Default) Times New Roman, 12 pt

3610 Norby, R.J.: Evaluation of 11 terrestrial carbon-nitrogen cycle models against observations
 3611 from two temperate Free-Air CO₂ Enrichment studies, *New Phytol.*, 202, 803–822,
 3612 doi:10.1111/nph.12697, 2014.

3613 Zamora, L. M., Prospero, J. M., and Hansell, D. A.: Organic nitrogen in aerosols and
 3614 precipitation at Barbados and Miami: Implications regarding sources, transport and deposition
 3615 to the western subtropical North Atlantic, *J. Geophys. Res.-Atmos.*, 116, D20309, doi:
 3616 10.1029/2011JD015660, 2011.

3617 Zhang, Q., Anatasio, C., and Jimenez-Cruz, M.: Water-soluble organic nitrogen in atmospheric
 3618 fine particles (PM_{2.5}) from northern California, *J. Geophys. Res.*, 107, D11, doi:
 3619 10.1029/2001JD000870, 2002.

3620 Zhang, Y., Song, L., Liu, X. J., Li, W. Q., Lu, S. H., Zheng, L. X., Bai, Z. C., Cai, G. Y., and
 3621 Zhang, F. S.: Atmospheric organic nitrogen deposition in China, *Atmos. Environ.*, 46, 195–
 3622 204, doi:10.1016/j.atmosenv.2011.09.080, 2012.

3623 Zhao, Y., Hallar, A. G., and Mazzoleni, L. R.: Atmospheric organic matter in clouds: exact
 3624 masses and molecular formula identification using ultrahigh-resolution FT-ICR mass
 3625 spectrometry, *Atmos. Chem. Phys.*, 13, 12343–12362, doi: 10.5194/acp-13-12343-2013, 2013.

3626 Zhuang, Q., Lu, Y., and Chen, M.: An inventory of global N₂O emissions from the soils of
 3627 natural terrestrial ecosystems, *Atmos. Env.*, 47, 66–75, doi:10.1016/j.atmosenv.2011.11.036,
 3628 2012.

3629 Zumft, W. G.: Cell biology and molecular basis of denitrification, *Micobiol. Mol. Biol. R.*, 61,
 3630 4, 533–616, 1997.

3631
 3632
 3633