



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Modeling particle nucleation and growth over northern California during the 2010 CARES campaign

A. Lupascu¹, R. Easter¹, R. Zaveri¹, M. Shrivastava¹, M. Pekour¹, J. Tomlinson¹, Q. Yang¹, H. Matsui^{2,3}, A. Hodzic⁴, Q. Zhang⁵, and J. D. Fast¹

¹Pacific Northwest National Laboratory, Richland, Washington, USA

²Graduate School of Environmental Studies, Nagoya University, Nagoya, Japan

³Department of Environmental Geochemical Cycle Research, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

⁴National Center for Atmospheric Research, Boulder, Colorado, USA

⁵Department of Environmental Toxicology, University of California – Davis, Davis, California, USA

Received: 17 June 2015 – Accepted: 2 July 2015 – Published: 20 July 2015

Correspondence to: J. D. Fast (jerome.fast@pnnl.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Accurate representation of the aerosol lifecycle requires adequate modeling of the particle number concentration and size distribution in addition to their mass, which is often the focus of aerosol modeling studies. This paper compares particle number concentrations and size distributions as predicted by three empirical nucleation parameterizations in the Weather Research and Forecast coupled with chemistry (WRF-Chem) regional model using 20 discrete size bins ranging from 1 nm to 10 μm . Two of the parameterizations are based on H_2SO_4 while one is based on both H_2SO_4 and organic vapors. Budget diagnostic terms for transport, dry deposition, emissions, condensational growth, nucleation, and coagulation of aerosol particles have been added to the model and are used to analyze the differences in how the new particle formation parameterizations influence the evolving aerosol size distribution. The simulations are evaluated using measurements collected at surface sites and from a research aircraft during the Carbonaceous Aerosol and Radiative Effects Study (CARES) conducted in the vicinity of Sacramento, California.

While all three parameterizations captured the temporal variation of the size distribution during observed nucleation events as well as the spatial variability in aerosol number, all overestimated by up to a factor of 2.5 the total particle number concentration for particle diameters greater than 10 nm. Using the budget diagnostic terms, we demonstrate that the combined H_2SO_4 and low-volatility organic vapors parameterization leads to a different diurnal variability of new particle formation and growth to larger sizes compared to the parameterizations based on only H_2SO_4 . At the CARES urban ground site, peak nucleation rates were predicted to occur around 12:00 Pacific (local standard time (PST)) for the H_2SO_4 parameterizations, whereas the highest rates were predicted at 08:00 and 16:00 PST when low-volatility organic gases are included in the parameterization. This can be explained by higher anthropogenic emissions of organic vapors at these times as well as lower boundary layer heights that reduce vertical mixing. The higher nucleation rates in the H_2SO_4 -organic parameterization at these times

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were largely offset by losses due to coagulation. Despite the different budget terms for ultrafine particles, the 10–40 nm diameter particle number concentrations from all three parameterizations increased from 10:00 to 14:00 PST and then decreased later in the afternoon, consistent with changes in the observed size and number distribution. Differences among the three simulations for the 40–100 nm particle diameter range are mostly associated with the timing of the peak total tendencies that shift the morning increase and afternoon decrease in particle number concentration by up to two hours. We found that newly formed particles could explain up to 20–30 % of predicted cloud condensation nuclei at 0.5 % supersaturation, depending on location and the specific nucleation parameterization. A sensitivity simulation using 12 discrete size bins ranging from 1 nm to 10 μm diameter gave a reasonable estimate of particle number and size distribution compared to the 20 size bin simulation, while reducing the associated computational cost by $\sim 36\%$.

1 Introduction

Aerosol particles are ubiquitous in the atmosphere and are important for their potential climate impact and role in atmospheric chemistry. They absorb and scatter solar radiation, act as cloud condensation nuclei (CCN) and ice nuclei (IN), and induce an overall cooling effect (IPCC, 2007, 2013). They are classified depending on their origin and source as primary or secondary particles. Primary particles originate from natural sources, such as dust and sea-salt, as well as anthropogenic sources, such as combustion. Secondary particles are formed through gas-to-particle partitioning processes. The formation of secondary aerosol particles and their subsequent growth has been observed in rural (Place et al., 2010; Ziemba et al., 2006), urban (Betha et al., 2013; Matsui et al., 2011; Jeong et al., 2004), marine (Hoppel et al., 1994; O’Dowd and Hoffmann, 2005), and high altitude (Venzac et al., 2009; Boulon et al., 2011; Cui et al., 2014) environments. Makkonen et al. (2009), Spracklen et al. (2006, 2008, 2010) and many others have shown that secondary particle formation contributes to cloud

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



condensation nuclei (CCN) and subsequently influence cloud droplet number concentrations and other cloud processes.

Sulfuric acid gas (H_2SO_4) plays an important role in the formation of small secondary aerosol particles, due to its very low vapor pressure. Recent studies showed that H_2SO_4 alone cannot explain the abundance of new particles in the troposphere (Merikanto et al., 2007; Kirkby et al., 2011; Schobesberger et al., 2013). Thus, several mechanisms have been proposed to describe the formation of new particles in the troposphere: the binary homogeneous mechanism (BHN) involving H_2SO_4 - H_2O (Wexler et al., 1994; McMurry, 1980; McMurry et al., 2000; Kulmala et al., 1998; Vehkamäki et al., 2002), the ternary nucleation (TN) mechanism involving H_2SO_4 - H_2O -ammonia (NH_3) (Napari et al., 2002; Merikanto et al., 2007), the ion induced nucleation mechanism (IIN) (Turco et al., 1998; Yu and Turco, 2000, 2001; Yu et al., 2008; Kazil et al., 2008), empirical particle formation mechanisms involving H_2SO_4 (Kulmala et al., 2006; Sihto et al., 2006; Riipinen et al., 2007; Kuang et al., 2008), and the combined organic and sulfuric acid (kinetic-type) empirical mechanism (ORG) (Metzger et al., 2010; Paasonen et al., 2010). In the empirical particle formation mechanisms, the nucleation rate is proportional to the H_2SO_4 concentration to the power of 1 to 2; the activation mechanism (ACT) (Kulmala et al., 2006) uses a power of one while the kinetic nucleation mechanism (KIN) (McMurry and Friedlander, 1979; Kuang et al., 2008) uses a power of two. Due to its dependency on temperature, the BHN mechanism produces new particles in the upper and free troposphere (e.g. Spracklen et al., 2005), while the use of state-of-the-science empirical parameterizations like ACT, KIN or ORG lead to better agreement between modeled and observed total number concentration in the boundary layer (i.e., Zhang et al., 2010a; Redington et al., 2011).

In spite of recent advancements, aerosol nucleation and growth processes are not yet completely understood mainly due to limited coincident measurements of aerosol nucleation precursors (such as H_2SO_4 , NH_3 , and organic acids), and the size, composition, and concentration of newly formed nanometer-sized particles. During the Cosmics Leaving OUtdoor Droplets (CLOUD) experiment, Kirby et al. (2011) showed that

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ternary nucleation, with and without ions, did not explain the observed boundary layer nucleation, suggesting that organic vapors participated in atmospheric nucleation. Several studies (i.e., Kuang et al., 2010, 2012; Smith et al., 2008) show that in many environments, H_2SO_4 accounts for less than half of the total growth of the particles. Setyan et al. (2012) suggested that organic vapors and, to a lesser extent, H_2SO_4 play key roles in new particle growth. Still, the role of organic vapors in particle nucleation and subsequent growth processes is very uncertain.

Several studies have been conducted to assess how the choice of nucleation scheme affects simulated particle number concentration (PNC) and their size distribution using global and regional models. For example, Jung et al. (2008) analyzed the impact of different nucleation parameterizations on the modeled size distribution predicted by the Dynamic Model for Aerosol Nucleation (DMAN) and its ability to reproduce the nucleation events and non-events observed during the Pittsburgh Air Quality Study (PAQS) conducted between July 2001 and September 2002. They showed that most of the nucleation schemes included in their study had difficulty reproducing the observed events. In that study, the empirical particle formation mechanism (Sihto et al., 2006) performed well on 70 % of the analyzed days, but still it predicted nucleation events on days when nucleation was not observed. The BHN mechanism of Vehkamäki et al. (2002) did not reproduce any observed nucleation events, and the scaling of the nucleation rate by a correction factor of 10^{10} led to a predicted event on a non-event day. Zhang et al. (2010a) quantified the impact of 11 nucleation schemes on predicted PNC, volume, and surface area using the US Environmental Protection Agency Community Multiscale Air Quality (CMAQ) modeling system version 4.4. They concluded that, although all the parameterizations underpredicted the total PNC (by a factor of 1.3 to 14.5), the ACT empirical formation mechanism (Sihto et al., 2006) led to the best prediction, while the least reliable prediction was given by the Merikanto et al. (2007) TN parameterization. Using the Global Model of Aerosol Processes (GLOMAP), Reddington et al. (2011) showed the use of ACT, KIN and ORG parameterizations led to underpredictions spanning from -53 to -11 % of total PNC in the boundary layer for

continental European monitoring station. In contrast, Jung et al. (2010) found that the PMCAMx-UF model using the TN nucleation parameterization (Merikanto et al., 2007), with the nucleation rate corrected using a nucleation tuner equal to 10^{-5} , overpredicted by 22 % the total number concentration of particles having a diameter larger than 3 nm, while this overprediction was greater than 49 % for particles having a diameter larger than 10 nm.

The aim of this study is to determine the performance of several new particle formation parameterizations implemented in the chemistry version of the Weather Research and Forecasting model (WRF-Chem) and investigate how differences in the parameterization formulations affect simulated spatial and temporal variations in boundary layer particle number concentrations during the Carbonaceous Aerosol and Radiative Effects Study (CARES) (Zaveri et al., 2012). The principal objectives of CARES were to examine the interaction between anthropogenic and biogenic precursors associated with secondary organic aerosol formation, the evolving mixing state of secondary organic and black carbon aerosols, and the impact of aerosols on optical and CCN activation properties. A complete overview of the research objectives and the ground and aircraft measurements collected during the CARES campaign is given by Zaveri et al. (2012). Although the primary objectives of the campaign did not include examining the formation of new particles, measurements of particle number concentration and size distribution showed that new particle formation events frequently occurred in the vicinity of Sacramento and over the western Sierra foothills. This environment is influenced by the SO_2 emissions originating from oil refineries located in the vicinity of the Carquinez Strait and the San Francisco Bay area. Setyan et al. (2014) showed that on days with new particle formation events during CARES, the concentration of organics and sulfate significantly increased in particles in the Aitken mode and that the concentrations of species representative of urban emissions together with the photo-oxidation products of biogenic volatile organic compounds (VOCs) and the biogenically influenced secondary organic aerosols (SOA) were on average 50 % higher during the event days compared to the non-event days. These findings provide means to test

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



several particle formation mechanisms that include the effect of sulfuric acid alone or combined effect of sulfuric acid and organic vapors, in addition to the BHN mechanism currently available in WRF-Chem.

The CARES measurements are described in Sect. 2. Key features and details of the WRF-Chem model and the simulations are described in Sect. 3. Section 4 presents the simulated evolution of the particle number concentration compared with surface and aircraft measurements collected on days in which new particle formation and growth events did and did not occur, and Sect. 5 summarizes the results.

2 Measurements

During the CARES campaign ground measurements were acquired at two sites in northern California: one in Sacramento (38.645° N, 121.34° W, ~ 30 m.a.s.l.) and the other in Cool (38.89° N, 120.97° W, ~ 450 m.a.s.l.), a small town located about 40 km northeast of Sacramento. Following Zaveri et al., (2010), we will refer to the Sacramento and Cool sites as “T0” and “T1”, respectively. Aerosol particle size measurements at both sites were carried out using a Scanning Mobility Particle Sizer (SMPS) to measure the particle the size distribution from 10 to 700 nm. Due to the lack of measurements in the 1–10 nm size range, the formation of new particles was not observed. Therefore, in the forthcoming discussion, following Setyan’s et al. (2014) terminology, instead of “nucleation” and “new particle formation”, we will use the term “new particle formation and growth event” (NPE). While the SMPS does not capture the initiation of NPEs, Zaveri et al. (2010) and Setyan et al. (2014) showed that the growth of small particles frequently occurred during the campaign. The CCN concentration at multiple supersaturations (0.07 to 0.5 %) was measured at both sites using Droplet Measurement Technologies CCN Counters (Model 200-013 and 100-081) (Zaveri et al., 2012). In addition to ground-based measurements, instruments deployed on the ARM Aerial Facility Gulfstream-1 (G-1) research aircraft, such as the condensation particle coun-

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ters CPC-3025 and CPC-3010 (Sem, 2002), measured particle number concentrations for optical diameters greater than 3 and 10 nm, respectively.

3 Model

In this study, the WRF-Chem model version 3.5 was used (Grell et al., 2005; Fast et al., 2006) to simulate new particle formation events and their impact on aerosol size distributions and CCN concentrations. The model domain encompassed California and the surrounding region using a 4 km grid spacing and 65 vertically-stretched layers from the ground up to 50 hPa. The physics options used for this study include the Morrison double-moment microphysics scheme (Morrison et al., 2009), the Kain–Fritsch cumulus parameterization (Kain, 2004), the Rapid Radiative Transfer Model (Iacono et al., 2008) for longwave and shortwave radiation, the Mellor–Yamada–Janjic boundary-layer parameterization (Janjic, 2001), and the Monin–Obukhov scheme for the surface layer.

Gas-phase chemistry is simulated using the SAPRC-99 mechanism (Carter, 2000). Aerosol lifecycle processes are represented by the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) (Zaveri et al., 2008), which has been used in numerous aerosol modeling studies (i.e. Cui et al., 2014; Ritter et al., 2013; Archer-Nicholls et al., 2014). Aerosol species in MOSAIC included sulfate, nitrate, ammonium, sodium, chloride, calcium, carbonate, other inorganics (i.e. dust), methanesulfonate, elemental carbon, primary organic aerosols (POA), and aerosol water (Zaveri et al., 2008). Secondary organic aerosols (SOA) are represented by the simplified volatility basis-set (VBS) approach of Shrivastava et al. (2011), with additional updates for biogenic SOA yields (Shrivastava et al., 2014, in preparation). The VBS approach treats traditional and non-traditional SOA species and precursor gases that partition between particle and gas phases. Traditional SOA precursors derive from oxidation of anthropogenic (e.g., aromatics) and biogenic (e.g., isoprene and terpenes) primary gases, and these are represented with 4 volatility bins having saturation vapor concentrations (C^*) of 0.1, 1, 10 and 100 $\mu\text{g m}^{-3}$ (at 298 K). Non-traditional SOA precursor gases derive from the

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



oxidation of semi- and intermediate volatility primary gases that are emitted during fossil and bio-fuel combustion and biomass burning; these are represented with 2 volatility bins having C^* of 0.001 and $10^6 \mu\text{g m}^{-3}$.

Initial and boundary conditions for the chemical species in our simulations, including the mass and number size distribution of primary aerosol species, are taken from the MOZART-4 global chemistry-transport model (Emmons et al., 2010). We reduced the initial and boundary conditions of aerosol concentrations from MOZART-4 by 50% since Fast et al. (2014) found that long-range transport was likely over-estimated when comparing simulated values to aerosol optical depth and extinction profile measurements.

Biogenic trace gas emissions are calculated online using the MEGAN model (Guenther et al., 2006). Anthropogenic emissions are obtained from the California Air Resources Board (CARB) 2010 project emissions available at <http://www.arb.ca.gov/app/emsmv/fcemssumcat2009.php>. The lognormal size distribution of emitted anthropogenic primary particles uses a geometrical mean diameter of 50 nm and a standard deviation of 2 (Cui et al., 2014). To avoid an artificial increase of 1 nm particles, primary aerosol emissions for particles smaller than 10 nm are not considered. Mineral dust and sea salt emissions are calculated on line (Shaw et al., 2008).

In the public release of WRF-Chem, MOSAIC uses a sectional framework where the aerosol size distribution is divided into 4 or 8 discrete size bins spanning 39 nm to 10 μm diameter. However, these bins do not capture the freshly nucleated particles that have diameters of a few nanometers. Recently, Matsui et al. (2011, 2013) and Cui et al., (2014) added a new sectional framework option for MOSAIC in WRF-Chem that uses 20 bins to represent nucleation and particle growth in the range of 1 nm to 10 μm , and a state-of-the-science empirical cluster activation (ACT) theory parameterization (Kulmala et al., 2006; Sihto et al., 2006). They compared new particle formation (NPF) and particle number concentrations predicted by that approach with land- and aircraft-based measurements. In Matsui et al. (2011), the use of empirical activation type nucleation inside WRF-Chem generally reproduced the number of observed event and

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

non-event days for a measurement site located ~ 50 km south of Beijing; however, the modeled SO_2 concentration was overestimated by a factor of 6 for this location. Matsui et al. (2013) applied the same approach for another case using several stations located in Korea and Japan. In that study, the simulated average number concentration for particles having a diameter greater than 10 nm (CN_{10}) was overestimated within a factor of 2, while the temporal variation of CN_{10} was not well reproduced. Cui et al. (2014) showed that the use of the ACT parameterization combined with an improved treatment of SOA formation from biogenic emissions at a forest site in Colorado led to a more accurate simulation of aerosol particles in the 4–100 nm size range compared with the default WRF-Chem model, including onset times, number concentrations, and number mean diameters.

For this study, we performed simulations using the 20 size bin framework (Matsui et al., 2011, 2013; Cui et al., 2014) with 5 different nucleation parameterizations: two based on classical binary homogeneous nucleation theory (Wexler et al., 1994 and Vehkamäki et al., 2002) and 3 newer empirical parameterizations. The Wexler parameterization is the default nucleation parameterization used with MOSAIC in the public release of WRF-Chem, and is used in our study for that reason. The Wexler parameterization defines a critical H_2SO_4 concentration at which the nucleation rate is approximately $1 \text{ particle cm}^{-3} \text{ s}^{-1}$, based on early classical BHN results. When the ambient gaseous H_2SO_4 concentration exceeds this critical concentration, all gaseous H_2SO_4 in excess of the critical value is used to produce new particles having a diameter equal to the smallest diameter covered by the model's sectional framework (e.g., 1 nm in the 20 size bin framework or 39 nm in the 8 size bin framework).

In the Vehkamäki parameterization, theoretical BHN nucleation rates (derived by Vehkamäki et al., 2002) over a wide range of temperature, relative humidity, and gaseous H_2SO_4 concentration are approximated (fitted) by a rather complicated analytical expression involving exponentials and polynomials of order 1 to 3.

We performed simulations using the Wexler and Vehkamäki parameterizations with the 20 size bin sectional framework, and using the MOSAIC WRF-Chem default con-

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



figuration of the Wexler parameterization with the 8 size bin sectional framework. Due to their poor performance, the results from the 20 size bin Wexler and Vehkamäki simulations are not presented; these simulations were not able to reproduce either the observed new particle formation events or the observed particle number concentrations. In addition, Zhang et al. (2010b) note that the Wexler parameterization often produces unrealistically high nucleation rates. We performed simulations with three of the recent empirical nucleation parameterizations using the 20 size bin sectional framework. These parameterizations are used in the boundary layer, while the Wexler parameterization is applied in the free troposphere. This approach was tested in previous global and regional modeling studies (e.g. Spracklen et al., 2006; Matsui et al., 2011; Reddington et al., 2011). For the activation type nucleation (ACT) and kinetic type nucleation (KIN), the formation rates of 1 nm particles are linear or second-order functions of H_2SO_4 concentration defined as:

$$J = k_{\text{ACT}} \times [\text{H}_2\text{SO}_4], \quad (1)$$

and

$$J = k_{\text{KIN}} \times [\text{H}_2\text{SO}_4]^2, \quad (2)$$

Previous studies showed that k_{ACT} and k_{KIN} coefficients derived from observations in different locations span several orders of magnitude: k_{ACT} ranges from 3.3×10^{-8} to $3.5 \times 10^{-4} \text{ s}^{-1}$ (i.e. Sihto et al., 2006; Riipinen et al., 2007); while k_{KIN} ranges from 2.4×10^{-15} to $1.3 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ (Riipinen et al., 2007; Kuang et al., 2008). In this study we use $k_{\text{ACT}} = 2 \times 10^{-6} \text{ s}^{-1}$ and $k_{\text{KIN}} = 2 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1}$, following Reddington et al. (2011). Cui et al. (2014) also recommend the use of $k_{\text{ACT}} = 2 \times 10^{-6} \text{ s}^{-1}$ based on the measured formation rates of ultrafine particles over a pine forest.

Although the ACT and KIN parameterizations have been used in many studies (i.e. Spracklen et al., 2006; Kuang et al., 2008). Metzger et al. (2010) showed that organic vapors are important for the nucleation processes. We therefore performed an addi-

tional simulation using the combined organic and H₂SO₄ mechanism (ORG):

$$J = k_{\text{ORG}} \times [\text{H}_2\text{SO}_4] \times [\text{OV}], \quad (3)$$

where $k_{\text{ORG}} = 5 \times 10^{-13} \text{ cm}^{-3} \text{ s}^{-1}$ as in Reddington et al. (2011) and OV = organic vapors.

In our WRF-Chem implementation, the OV used in this mechanism consist of the lowest volatility gases of the SOA mechanism (those with $C^* = 0.1$ and $0.001 \mu\text{g m}^{-3}$). In comparison, the OV in Redington et al. (2011) was the first stage oxidation product of monoterpenes. Following Yli-Juunti et al. (2013), an upper limit of $10^8 \text{ molecules cm}^{-3}$ for the OV is used in Eq. (3).

Once formed, the new particles then grow by condensation of inorganic and organic gases, and for the small initial sizes, the Kelvin effect influences the condensation rates. MOSAIC treats the Kelvin effect for the condensation of inorganic gases (although H₂SO₄ is treated as non-volatile), but the SOA gas-particle partitioning treatment was modified to include the Kelvin effect. We assume a surface tension of 50 dyne cm^{-1} , and a molecular weight of 250.0 g mol^{-1} for the condensable OV, so at 298 K, the Kelvin effect increases the effective saturation vapor concentrations over 1 nm particles by a factor of 55.

Budget diagnostics for aerosol nucleation, emission, condensational growth, coagulation, transport, dry deposition and total tendencies (the sum of all the individually computed tendencies) were saved to illustrate when and where new particle formation events occurred and better understand how they influence the evolution of the aerosol size distribution.

We also performed two additional simulations with the ACT parameterization. One used 12 discrete size bins, also ranging from 1 nm to 10 μm diameter, with good resolution in the size range of interest to this study (new particle formation). The other used the MOSAIC default 8 size bins, with diameters rangings from 40 nm to 10 μm , to quantify the performance and computational cost of this size bin structure compared to the 12-bin and 20-bin versions. In order to take into account nuclei growth from 1

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to 40 nm by condensation, cluster scavenging, and self-coagulation in the default 8 bin version, we applied the Kerminen and Kulmala (2002) parameterization. For this purpose, we calculate the growth time from 1 to 40 nm, and from 40 to 63 nm, due to H₂SO₄ condensation only and due to H₂SO₄ + OV condensation for the 20 bin version of the model. Using the linear regression method in which we set the intercept value equal to 0, we estimate that the growth time from 1 to 40 nm is ~ 0.74 h, and tuned the growth rate defined in the Kerminen and Kulmala (2002) parameterization with this factor.

In addition to improving simulations of the aerosol formation and growth processes in MOSAIC, the results of this study will form the basis of incorporating one or more new nucleation schemes into the public version of WRF-Chem.

4 Results

The discussion of the model results focuses on the NPE observed between 7 and 16 June 2010. The prevailing westerly to southwesterly atmospheric flow on 7–9 June and 15–16 June 2010 transported SO₂ from the San Francisco Bay area to the vicinity of Sacramento and produced NPE on these days. Due to changes of synoptic conditions associated with a trough formed over the western US from 10 June to 14 June 2010 (Fast et al., 2012), weak or no NPEs were observed at the T0 (urban) and T1 (rural) sites on these days. The shift to strong northerly winds led to a decrease of Bay Area emissions transported over Sacramento, consequently decreasing SO₂ and the total observed particle number concentration. We first evaluate the overall impact of the nucleation parameterizations on particle number and size distribution at the surface and aloft, using measurements collected at the T0 and T1 sites as well as along G-1 aircraft flight paths during the 7 to 16 June period. We then provide a more detailed analysis of the simulated formation of new particles and growth at ground level for an observed NPE on 8 June 2010 and on 12 June 2010 when no NPE was observed.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Finally, we determine how the differences in particle number and size distribution associated with the nucleation parameterizations affect predicted CCN.

4.1 Temporal variability in particle number concentration

The ability of the model to simulate the observations is evaluated using statistical scores including normalized mean bias (NMB), and the correlation factor between simulated and measured values (R). Due to the minimum cutoff size of the SMPS instrument, we investigated the behavior of CN_{10} (particle number concentration having $D_p \geq 10$ nm), CN_{40} ($D_p \geq 40$ nm) and CN_{100} ($D_p \geq 100$ nm). Tables 1 and 2 present NMB and R for these three parameters from four model configurations at the T0 and T1 sites. Model performance in simulating PM_1 and $PM_{2.5}$ concentrations using the default WRF-Chem 8 size-bin and Wexler nucleation parameterization configuration (WEX-8BIN) has already been presented in Fast et al. (2014); however, we note that small errors in simulated mass due to additional emissions in the 10–40 nm range can lead to relatively large errors in number concentration for small particles.

As shown in Figs. 1 and 2, the overall temporal variability of high particle concentrations before and after the passage of the trough between 10 and 14 June was generally captured by the explicit (20 size bin) nucleation schemes used in our study, although they usually overestimate the observed particle number concentrations.

First, we analyze how the standard configuration of WRF-Chem (8BIN) behaves compared to the measurements. Since the 8BIN simulation does not explicitly resolve nucleation and the geometric diameter range for the smallest size bin is from 39 to 78 nm, only CN_{40} and CN_{100} are examined. As can be seen from Figs. 1 and 2, the WEX-8BIN simulation is able to reproduce the observed diurnal and multi-day variation of CN_{40} ($R = 0.31$ and 0.57 for the T0 and T1 sites, respectively), and largely underestimates the observed number concentration for both the T0 and T1 sites (see Tables 1 and 2). In the case of CN_{100} , the 8BIN simulation also tends to underestimate the observed concentration (up to 65 % at the T1 site) with CN_{100} R values of 0.60 and

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

0.32 calculated between observations and modeling results at the T0 and T1 sites, respectively.

We now discuss how well simulations conducted using the empirical nucleation parameterizations with 20 size bins agree with the measurements. All three empirical nucleation parameterizations reproduce the overall observed daily variability as shown Figs. 1 and 2, although during the NPEs they usually overestimate the PNC in different size ranges. The ACT, KIN and ORG simulations overestimate the observed CN₁₀ (from 178 % for ACT up to 255 % for KIN), although the correlation coefficient suggests reasonable agreement in temporal variability between observations and modeling results ($R > 0.70$). The overestimation is reduced for the CN₄₀ and CN₁₀₀ particles, and the best results are obtained at the rural T1 site that is not influenced by as high anthropogenic emission rates (see Table 1 and 2). Moreover, during the southwesterly flow periods the maximum peaks of CN₁₀ are overpredicted within a factor of 3 by all explicit mechanisms involved in our simulations, while the CN₄₀ and CN₁₀₀ maximum peaks tend to be overpredicted by a factor of 2 (Figs. 1 and 2). The normalized mean bias of CN₁₀ during this period ranges from 187 % for ACT to 347 % for KIN. The overprediction factors for all three empirical parameterizations are reduced during the northerly flow period, when SO₂ transport through the Carquinez Strait to the measurement sites is significantly reduced. During this period the normalized mean bias of CN₁₀ particles is lower and ranges from 129 % for ACT to 278 % for KIN. Based on this analysis of PNC in several size ranges, we conclude that for both measurement sites, the ACT parameterization performs somewhat better than the KIN and ORG parameterizations.

4.2 Analysis of the observed NPE on 8 June 2010

Figures 3 and 4 compare the hourly modeled and observed number size distributions as a function of particle diameter at the T0 and T1 sites, respectively, on 8 June. For this study we have adopted the classification of the observed nucleation events given by Boy et al. (2008). Based on occurrence and clarity of new particle formation events, they define four categories: A-event days, B-event days, undefined, and

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



non-event days. According to their classification an A-event shows a clear nucleation mode particles (the highest number of particles are observed at the lowest measurable SMPS size) and subsequent growth to larger sizes; a B-event is less clearly than an A-event (the highest number of particles are not observed in the first SMPS size bin, but at a greater diameter) and it is followed by the subsequent growth. Non-event days are those with no particle formation, while the days that cannot be classified as event or non-event days are called undefined days. Following this classification, on 8 June 2010 for both the T0 and T1 sites a “class B” event occurred. Thus, the highest concentration of particles are not observed in the first SMPS size bin (~ 12.2 and 8.75 nm diameter at the T0 and T1 sites), but at a greater diameter (~ 13.6 and 16 nm diameter at the T0 and T1 sites on this day), and growth of these particles to Aitken mode was observed (Figs. 3a and 4a).

All of the simulations captured the observed NPE but with different PNC. The ACT, KIN, and ORG schemes behave similarly on this day, leading to a daily NMB for CN_{10} of 234, 346, and 206 % at the T0 site and 143, 198 and 175 % at the T1 site, respectively. Correlation coefficients were between 0.89 and 0.92, indicating a good temporal agreement between observed and simulated CN_{10} .

Observations at both sites show an increase of particles in 10–20 nm range around 09:00 PST (Figs. 3a and 4a). These particles continue to grow in size in the following hours until they reach 80–100 nm. Although the observed onset of nucleation starts around 09:00 PST, in our simulations the onset of nucleation is two hours earlier (Figs. 3b–d and 4b–d). This behavior could be attributed to the break-up of stable nocturnal boundary layer and vertical mixing with cleaner air from the lower troposphere, which can trigger the formation of new particles (Kristensson et al., 2008). On this day, the simulated boundary layer growth at the T0 site was very similar to that derived from radiosonde observations; therefore, vertical mixing associated with the growing convective boundary layer is likely reasonably simulated by the model. From the comparison of simulated CN_{10} aloft with the measurements during the morning G-1 flight on this day, the overestimation factors for CN_{10} aloft (90, 73, and 118 % for ACT, KIN,

and ORG) are much lower than for the surface sites. Therefore, the cleaner simulated free troposphere air could enhance formation of new particles near the surface as the boundary layer grows, as shown in Fig. 5. This figure also shows that for ACT the highest concentration of small particles coming from the nucleation process occurs at $\sim 2/3$ of the boundary layer height. A higher number of small particles exists at this altitude compared to the number predicted at the surface, with concentrations aloft $\sim 30\%$ and $\sim 25\%$ greater at T0 and T1, respectively. These particles formed aloft would be quickly transported to the surface by turbulent vertical mixing. KIN acts in a similar manner as ACT, although it predicts an increased number of small particles. In contrast, ORG predicts that the highest concentration of particles occurs at the surface. We can associate this with the relatively high concentration of OV predicted at the surface compared to the upper layers; therefore, enhanced particle production at this level is not unexpected when the ORG parameterization is used. In addition, the simulated growth of particles corresponds to the increase of fine-particle SOA mass (Fig. 6), which also starts about 2 h before the observed NPE, indicating that SOA might have contributed to the simulated growth of ultrafine particles.

The observed continuous growth of nucleation mode particles through the day at both T0 and T1 sites (Figs. 3a and 4a) suggests that nucleation and growth take places over regional scales. Figure 7a shows the simulated nucleation rate (from ACT) averaged from the surface to ~ 130 m.a.g.l. from 05:00 to 09:00 PST on 8 June 2010. Superimposed on this figure is a back trajectory (at 100 m.a.g.l.) for air arriving at the T0 site at 09:00 PST on this day. The back trajectory indicates the air mass is transported by southwesterly winds from the San Francisco Bay area through the Carquinez Strait into the Central Valley before arriving in Sacramento. The peak nucleation rates occur over industrialized region along northern San Francisco Bay that extends towards the T0 site. Nucleation also occurs over most of the Sacramento Valley, but the rates are about an order of magnitude lower than between the Carquinez Strait and Sacramento. Figure 7b shows the computed PNC in the 1–10 nm (CN_{1-10}), 10–40 nm (CN_{10-40}) and 40–100 nm (CN_{40-100}) ranges along the trajectory. High concentrations

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of CN_{1–10} and CN_{10–40} particles occur along the trajectory prior to their arrival over the T0 site. The sharp increase of CN_{1–10} particles after 05:00 PST 8 June, followed two hours later by the increase of CN_{10–40} particles are associated with the early morning onset of nucleation. The model results indicate that nucleation and growth of aerosol particles were not only a local phenomenon, but also took place upwind of Sacramento before being transported over the T0 site (see Fig. 7a and b).

To obtain a better understanding of the factors that contribute to differences in the simulated number size distribution among the different parameterizations, we use budget diagnostics. These diagnostics are the tendencies associated with dry deposition plus vertical turbulent mixing, other transport (advection and horizontal turbulent mixing), emissions, condensation, nucleation, coagulation, and total tendency that affect PNC in the 1–10, 10–40, and 40–100 nm diameter ranges. As shown in Figs. 8–10 the use of different nucleation parameterizations leads to differences associated with the processes that contribute to aerosol formation and growth.

Even though both the ACT and KIN parameterizations depend only on H₂SO₄ concentration, the differences in their formulation lead to differences in the nucleation rates and consequently CN₁₀ concentration, as shown previously. The KIN nucleation rates are higher than the ACT nucleation rates by as much as 1474 and 930 % at the T0 and T1 sites, respectively. These differences in the nucleation rate could explain the differences in the magnitude of the budget terms associated with condensation (Figs. 8c, 9c, and 10c), coagulation (Figs. 8e, 9e, and 10e) and predicted particle number concentration in the 1 to 10 nm range (Figs. 8g, 9g, and 10g). Moreover, these differences in the nucleation rate could also explain the magnitude of combined nucleation and coagulation processes (Fig. 8h) that suggest for ACT the coagulation loss is more important than nucleation production. Similarly, the higher predicted nucleation rate for KIN (Fig. 8d) could explain the difference in predicted number concentration in 1 to 10 nm range (Fig. 8g). The aforementioned budget terms (nucleation, condensation, and coagulation) that contribute to the formation and growth of particles are strongly correlated with the diurnal variation of H₂SO₄ concentration at both sites on this day

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(less so with the ORG parameterization), although their magnitudes highly differ. Note that the diurnal variations of budget terms for dry deposition plus vertical mixing and for other transport for 10–40 and 40–100 nm particles are similar among the ACT, KIN, and ORG simulations (Figs. 9a, b and 10a, b), but the diurnal variations in these budget terms for 1–10 nm particles are quite different among the three nucleation parameterizations. Figure 8a shows that while the CN_{1-10} in KIN are lost through deposition and vertical mixing processes, a net gain of particles is predicted during mid-day hours and late afternoon using ACT, showing the impact of vertical mixing on predicted number concentration at the surface. Moreover, analysis of the other transport budget term (Fig. 8b) shows an increase of CN_{1-10} at T0 starting with the onset of modeled nucleation for both the ACT and KIN parameterizations, but a weak decrease for the ORG parameterization. These findings confirm that the nucleation and growth of aerosol particles are not just a local phenomenon but take place over a regional scale. A net gain of particles transported from upwind is simulated using ACT in the late afternoon (Fig. 8b), suggesting that the differences associated with the nucleation parameterization formulation and their strength impacts non-local phenomena such as horizontal transport.

We note that the use of the ACT and ORG schemes leads to similar results in terms of total PNC and size distribution during this day, whereas the budget diagnostic terms show several differences. Pierce et al. (2012) found a relatively small difference between ACT and ORG scheme simulations, which they attributed to high correlations of both H_2SO_4 and low-volatility organics with sunlight. In our simulation only H_2SO_4 is highly correlated with downward short wave flux ($R = 0.83$ over 06:00 to 20:00 PST), while the low-volatility OV concentrations are relatively poorly correlated with the downward shortwave radiation flux ($R = 0.38$). As shown in Fig. 11, the daily variability of H_2SO_4 and OV concentrations present a different behavior, with H_2SO_4 concentrations peaking at 1.34×10^8 molecules cm^{-3} close to 13:00 PST and OV concentration peaking at 6.57×10^6 molecules cm^{-3} around 17:00 PST. Although both H_2SO_4 and OV are linked to the formation of OH radicals via photolysis, and hence to the predicted solar

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

radiation that regulates the intensity of photochemical reactions, their different diurnal patterns can be related to lifetime differences. Thus, the longer lifetime of OV compared to H₂SO₄ in association with the shallow boundary layer might explain the timing of predicted peaks for these gases.

5 The nucleation budget term demonstrates the impact of these low-volatility gases on the new particle formation. The ORG budget terms and particle number concentration for 1–10 nm particles have a different diurnal variability compared to ACT (Fig. 8). As can be seen from Fig. 8d and g, the ORG scheme shows two peaks in particle concentration that are associated with the increase of H₂SO₄ concentration in the morning,
10 and the increase of low-volatility OV concentration during the afternoon. As in case of ACT and KIN, the ORG budget terms associated with condensation (Figs. 8c, 9c, and 10c), nucleation (Fig. 8d) and coagulation (Figs. 8e, 9e, and 10e) show the connection of these terms to the way that the nucleation is parameterized. Still, the budget terms and predicted number concentration (Figs. 9g and 10g) in the 10–40 nm and
15 40–100 nm size range present almost the same daily variability for all aforementioned parameterizations. Thus, we can say that even though the treatment of NPF has a great impact on the production rate of freshly nucleated aerosols (Fig. 8), removal processes through condensation, coagulation or dry deposition are efficient and could explain the relatively similar CN₄₀ and CN₁₀₀ predicted by ACT, KIN and ORG schemes. Analyzing
20 the total tendency of particle number we note that particles in 1–10 nm range have a net gain of particles starting around 05:00 PST (Fig. 8f); this trend is kept for larger particles, with a modeled net gain that starts around 06:00 PST for particles in 10–40 nm range (Fig. 9f) and around 08:00 PST for particles in 40–100 nm range (Fig. 10f). This behavior is in accord with new particle formation and growth processes.

25 In general, the simulations also capture the observed simultaneous decrease of particle size and number concentrations during the afternoon, as can be seen in Figs. 1 and 2. The size distribution changes are more clearly seen in Fig. 12, which shows the observed and simulated particle number distributions at 06:00, 14:00, and 19:00 PST on 8 June 2010. The measurements show that the 20–100 nm PNC increases between

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



06:00 and 14:00 PST, presumably due to nucleation and condensational growth. Particle concentrations decrease later in the afternoon as nucleation rates decrease and the wind direction shifts. All nucleation schemes reproduced this variation in 20–100 nm PNC during the day at the T0 site, while at the T1 site both the observed and simulated PNC did not decrease between 14:00 and 19:00 LST. Southwesterly winds during the afternoon of 8 June transported air from the vicinity of Sacramento to the T1 site and kept concentrations of small particles relatively high.

In summary, we show that all the empirical parameterizations used in our simulations were able to qualitatively reproduce characteristics of the observed NPE on 8 June 2010. However, for all parameterizations the onset of the NPE was about 2 h too soon and the simulated PNC was too high. From the analysis of budget diagnostic terms for 1–10 nm particles, we find that at both T0 and T1 sites on this day, the ACT and KIN simulations exhibit almost the same temporal variability and are highly correlated with the diurnal variation of H_2SO_4 concentration. In contrast to the ACT and KIN simulations that show one peak in nucleation rate around noon, the diurnal variability from the ORG simulation is different, showing two peaks for nucleation rates and many other differences in budget diagnostic terms for 1–10 nm particles. Thus, the different nucleation parameterizations lead to different results in both particle number and size distribution.

4.3 Analysis of an observed non-NPE on 12 June 2010

On 12 June 2010, modest increases in CN_{10} were observed at the T0 and T1 sites during the day (Figs. 1 and 2). The observed size distributions in Figs. 13a and 14a show that the number of particles in 10–30 nm size range increased during the day, but this was not followed by particle growth to larger sizes, as was the case on 8 June 2010. Although some nucleation was likely occurring, this behavior does not qualify as a nucleation event, based on the Boy et al. (2008) classification. The daily NMB values show that ACT, KIN, and ORG simulations overestimate CN_{10} by 192, 199, and 307 %, respectively, at the T0 site and by 139, 313 and 118 %, respectively, at the T1 site.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Correlation coefficients between observed and simulated values range between 0.40 and 0.49, indicating that these parameterizations coupled with other model processes did not capture the observed diurnal variability as well as on 8 June.

The observed particle number size distribution at the T0 site shows a shift in the maximum from 10–30 nm range to 20–50 nm range after 19:00 PST (Fig. 13a). All simulations reproduce this feature, although the shift in size is between 20 and 80 nm and the PNC is too high.

A possible explanation for this observed and also modeled feature can be related to the structure of the boundary layer. In contrast to the growth of the convective boundary layer during the day, the simulated boundary layer collapse after 18:00 PST, coinciding with increased stability near the surface. This stability does not allow vertical mixing with the air aloft; therefore, large particles remain constricted to the boundary layer and lead to the simulated increase in the 20–80 nm range. Again, as in the case of 8 June, we speculate that the change in wind direction and the decrease of wind speed between 18:00 and 19:00 PST might also explain the shift in particle size distribution. With lower wind speeds, fewer particles are transported away (Figs. 15b and 16b). At the same time we noticed a net gain of particles in the 40–100 nm size range (Fig. 17c), most likely coming from the condensation of OV, that is within an order of magnitude higher than the predicted H_2SO_4 (Fig. 11).

As in case of 8 June 2010, the analysis of budget terms associated with the formation and growth of particles together with the predicted number concentration shows that the differences in the nucleation rate formulation lead to relatively large daily variability in the 1–10 nm size range (Fig. 15), while these differences are generally much smaller in the 10–40 nm and 40–100 nm size ranges (Figs. 16 and 17). For this particular case, most budget terms for ACT and KIN are similar for all size ranges (Figs. 15–17), although nucleation and coagulation for 1–10 nm particles is larger for KIN, while ORG shows the impact of low-volatility OV on the nucleation rate and subsequent influence on diurnal variation. During this period, the amount of SO_2 transport to the site is small, so there is less H_2SO_4 present compared to the low-volatility OV generated from local

sources. Thus nucleation and subsequent growth of new particles is greater in case of the ORG simulation because of the local sources of OV, leading to the larger overestimation of 1–10 nm (Fig. 15g) and 10–40 nm particles (Fig. 16g) at the T0 site compared to the other mechanisms.

Due to missing measurements between 12:00 and 18:00 PST, the evolving observed particle number size distribution at the T1 site (Fig. 14a) is not complete. Still, as for the T0 site, we can observe at T1 a maximum in size distribution at 11:00 PST in the 10–30 nm range which is shifted towards the 30–50 nm range after 18:00 PST. However, the simulations are not able to reproduce this feature, with the model indicating a shift from the 6–10 to the 10–20 nm range. From an analysis of the budget terms and the OV gas concentration, we can attribute the shift in the T1 size distribution around sunset to the same causes as the T0 shift, namely the boundary layer collapse, variations in wind speed and direction, and (for the ORG parameterization) the condensation of OV.

4.4 Impact of nucleation parameterizations on physical processes associated to the NPEs

Table 3 shows selected budget terms on all NPE days (7–11 June and 14–16 June 2010 at the T0 site, and 7–10 June and 14–16 June at the T1 site) averaged over the period from 04:00 PST through 16:00 PST. On each NPE day, the nucleation budget terms for 1–10 nm particles from the three nucleation parameterizations differ by factors of about 10–50. The ACT nucleation term is always the smallest, and there are large differences between KIN and ORG on many days, with KIN largest on some days and ORG largest on others. Yet, the nucleation and coagulation terms are similar in magnitudes for any given parameterization and day, indicating that coagulation strongly moderates the large differences in nucleation rates. This sustains the findings from the diurnal variability analysis of 8 June 2010 NPE.

The CN_{1-10} condensation tendencies for all three parameterizations are much closer in magnitude to each other, generally within a factor of 2, except for the 7 June 2010 case when the condensation tendencies are higher by factors of 4.6 and 5.6 for KIN and

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ORG compared to ACT. The CN_{1-10} condensation tendency (when multiplied by -1) gives the effective production of particles larger than 10 nm diameter. These effective $CN > 10$ production rates are thus much less sensitive to the nucleation parameterizations, and are also much smaller in magnitude than the nucleation rates, because of the strong modulation of nucleation by coagulation.

The combined condensation and coagulation tendencies for 10–100 nm particles are quite similar to the CN_{1-10} condensation on some days but are lower by factors of about 2–4 on other days, indicating further modulation of the new particle production by coagulation on some days. For the 7 June 2010 case noted above, the combined CN_{10-100} condensation and coagulation terms from the three nucleation parameterizations are somewhat closer (factor of 2.8 range) compared to the CN_{1-10} condensation (factor of 5.6 range) due to this additional modulation. Note that the CN_{10-100} condensation plus coagulation term for KIN is on average about 40 % higher than for ACT, and this is in agreement with the KIN vs. ACT CN_{10} concentration differences presented in Sect. 4.1.

The differences in the CN_{1-10} budget terms for horizontal transport, vertical mixing and dry deposition indicate the impact of nucleation and growth processes over regional spatial scales. As stated previously, the efficiency of removal processes such as condensation, coagulation or dry deposition could explain the relatively small differences among simulations for the predicted particle number concentrations in the 10–40 and 40–100 nm size ranges (see Figs. 1 and 2 and Table 3). Once more, we have shown that although the formulation of the nucleation rate parameterization impacts the prediction of newly formed particles, removal processes strongly modulate the nucleation parameterization differences.

4.5 Comparison with aircraft measurements in the and above boundary layer

As shown in Table 4, all simulations produced average particle number concentrations aloft that were generally too high when compared to the CPC measurements collected during the eight aircraft flights. The statistics for particles with diameters greater than 3 nm (CN_3) show that all parameterizations predicted more particles in the nucleation

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mode than were measured. The best results were obtained from the ACT simulation (NMB ranging from 157 to 274 %) and the worst results were obtained for the KIN simulation (NMB ranging from 162 to 845 %). A similar statistical trend occurred for CN_{10} , except that the differences between model results and observations were greatly reduced compared to CN_3 . For CN_{10} , the KIN simulation had a NMB between 105 and 338 %, while the ACT simulation had a NMB between 69 and 230 %.

There are a few instances, such as the flight during the afternoon on 8 June (flight 08b), in which the statistical analysis shows that all the simulations reproduced the observed spatial variability of CN_3 and CN_{10} reasonably well ($R > 0.67$ and $R > 0.77$), while overestimating the observed concentration (CN_3 concentration has a NMB ranging from 204 % for ORG to 603 % for KIN while CN_{10} concentration has a NMB ranging from 162 % for ACT to 219 % for KIN). There are other cases for which the simulated CN_3 and CN_{10} concentration have similar high biases, but the simulated spatial variabilities do not agree well with the measurements (low correlation), such as on the afternoon of 12 June 2010 (flight 12b).

To summarize the overall performance of the different simulations we also computed statistics for all flights combined (see Table 4). As for individual flights, the best results were obtained when using the ACT parameterization (NMB of 226 and 162 % for CN_3 and CN_{10} , respectively) and the worst results were obtained for the KIN parameterization (NMB of 608 and 239 % for CN_3 and CN_{10} , respectively). The correlation coefficients (between 0.48 and 0.54) show a moderate agreement in capturing the observed spatial variability of both CN_3 and CN_{10} .

We also examined the model performance in simulating CN_3 and CN_{10} as function of aircraft altitude, using intervals from the ground to 500, 500–1000, 1000–1500, and 1500–2000 m a.g.l. Comparisons between the measured and predicted median PNC, together with 25th and 75th percentiles, for the afternoon flights of 8 June and 12 June are shown in Figs. 18 and 19, respectively. Below 1000 m a.g.l., the simulated CN_3 and CN_{10} concentrations are usually 100–200 % higher than the corresponding measurements (see Figs. 18 and 19). Note that the afternoon maximum boundary layer height

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on 8 June was ~ 800 m.a.g.l. (Fig. 3), suggesting that the majority of the measured and simulated CN_3 and CN_{10} particles were located within the boundary layer. Comparison between modeled and observed CN_3 and CN_{10} medians and interquartile ranges above 1000 m.a.g.l. on 8 June 2010 (Fig. 18c, d, g and h) shows that all simulations reasonably represent the measurements. CN_3 concentrations have a NMB ranging from 56 % for ACT to 211 % for KIN, while CN_{10} is overestimated by 70–100 %. This behavior can be associated with lower SO_2 concentrations at these altitudes (simulated but not measured), and consequently lower H_2SO_4 concentrations and less new particle formation.

The strong northerly winds present on 12 June 2010 led to much lower SO_2 concentrations compared to 8 June, and this might explain the relatively homogeneous observed and simulated particle number concentrations at different altitudes (Fig. 19). Yet, the median values of CN_3 show that our simulations still overestimate number concentrations by up to a factor of 6, especially for KIN and ORG, while the overestimation factor reaches 4.5 for CN_{10} .

This comparison of simulated particle concentrations with the aircraft measurements shows that the three nucleation parameterizations were able to reproduce the temporal and spatial variations in particle number for several flights; however, the simulated concentrations in the boundary layer are too high, especially for the smallest particles.

4.6 Impact of nucleation schemes on CCN concentration

CCN concentration depends on both the particle size distribution and aerosol chemical composition. Observational studies have linked nucleation events to CCN production (e.g. Asmi et al., 2011; Wiedensohler et al., 2009). New particles can grow to the size of CCN through condensation and coagulation processes, although only a portion of these particles reach that size. As discussed in Kerminen et al. (2012) and references therein, the connection between the formation of new particles and CCN concentration has recently started to be investigated by using models. They note a non-linear dependence between atmospheric CCN production and the nucleation process due to (1) the

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nucleation rate, (2) subsequent growth of nucleated particles to larger sizes, and (3) the presence of primary aerosol particles, and pointed out the challenge in accurately predicting CCN concentrations given these factors. Several studies (e.g. Merikanto et al., 2009; Westervelt et al., 2014) have shown that boundary layer nucleation plays an important role in determining CCN number concentration. During CARES, CCN number concentrations were measured at both the T0 and T1 sites for several supersaturations (SS = 0.1, 0.2, 0.35, 0.5 %). To assess the impact of the nucleation parameterizations on CCN, we next compare observed CCN with simulated CCN number concentration calculated offline using the simulated chemical composition and the hygroscopicity of particles together with their size-dependent number concentration predicted by WRF-Chem.

As shown in Tables 5 and 6, the calculated CCN number concentrations generally are weakly correlated with the measurements. The calculated CCN are generally lower than the measurements, except for supersaturations of 0.5 %. The underestimation at 0.1 and 0.2 % supersaturations is likely to be closely linked to the large number of ultrafine particles that are not activated at these supersaturations, since the effective diameter of CCN is inversely correlated to the supersaturation (Yue et al., 2011). From the predicted CCN concentration and particle size distribution we estimated the critical diameter for cloud droplet activation. There is no clear distinction in the average simulated critical diameter between the T0 and T1 sites, and it varies from ~ 55 nm at 0.5 % supersaturation to ~ 215 nm at a supersaturation of 0.1 %.

Figures 20 and 21 present the times series of observed and calculated CCN at both the T0 and T1 sites for five supersaturations. Comparing observed CCN and CN₁₀ concentrations shows that several hours after the NPEs take place, an increase in CCN concentration is also observed. This is consistent with observational studies around the world that reported a clear increase in the CCN concentration after the occurrence of nucleation events (e.g., Kuata et al., 2008; Levin et al., 2012). During the northerly wind period, the simulated CCN concentrations better match the observations at the T1 site compared to results at the T0 site. This is likely due to lower PNC biases for all size

ranges at the T1 site during the same period, thus also demonstrating the connection between PNC and CCN.

We next analyze how selection of the nucleation parameterization affects the predicted CCN number concentration for one supersaturation ($SS = 0.5\%$). For this purpose we compare the ACT, KIN, and ORG results with a simulation in which nucleation (and NPF) was turned off. Compared to the simulation with no NPF, CCN levels at $SS = 0.5\%$ from ACT, KIN, and ORG are around 18% higher at the T0 site and 26% higher at the T1 site. These changes indicate that nucleation and subsequent growth moderately influences the CCN at this supersaturation. This result has been found in previous studies that report 5–70% of the CCN could be attributed to nucleation (e.g. Spracklen et al., 2010; Pierce and Adams, 2009; Matsui et al., 2013).

For $SS = 0.5\%$, the average activation ratios (AR), defined as $AR = CCN/CN_{10}$, are around 0.10 at both measurement sites for all parameterizations. As pointed out by Andreae and Rosenfeld (2008), low AR are observed with fresh aerosols, whereas AR approaches 1 for aged aerosols. The relatively small ARs obtained for the ACT, KIN, and ORG parameterizations exhibit the importance of freshly formed secondary aerosols and are relatively close to the observed AR (0.14 and 0.21 for the T0 and T1 sites, respectively).

Thus, from the statistical analysis and the temporal evolution of CCN at different supersaturations, we can conclude that nucleation affects both CCN concentrations and the fraction of particles that are available to act as CCN. However, the ACT, KIN, and ORG parameterizations give fairly similar results, suggesting that the CCN-sized particles are mainly driven by processes other than nucleation.

4.7 Impact of sectional bin number on predicted particle number and mass concentration

Based on the statistics shown in Sect. 4.1, the ACT parameterization performed somewhat better for both surface measurement sites and for most of the G-1 aircraft flights. Therefore, we now focus on comparing the impact of sectional bin number on predicted

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



particle number and mass concentration when the ACT parameterization is used. In addition to the 20-size bin simulation (hereafter referred to ACT-20BIN) presented previously, we performed two additional simulations. Simulation ACT-8BIN employs the default 8-particle-size-bins ranging from 39 nm to 10 μm used by MOSAIC in the publicly released version of WRF-Chem. Note that for this configuration, the Kerminen and Kulmala (2002) parameterization is applied to estimate the coagulation loss of new particles during their growth from 1 to 39 nm. The second simulation, ACT-12BIN, uses 12-particle-size-bins ranging from 1 nm to 10 μm . This bin structure still explicitly resolves new particle formation, while reducing computational costs compared to ACT-20BIN by 36 %. Using ACT-8BIN reduces the computing cost by 45 % compared to ACT-20BIN.

The comparison between standard 8 bin configuration that uses the Wexler parameterization (WEX-8BIN) and 8 bin version that uses ACT (ACT-8BIN) shows a modest improvement in predicted CN_{40} concentration when ACT is used (Tables 7 and 8). For example, at the T0 site the CN_{40} NMB is reduced from -34 % for WEX-8BIN (Table 1) to -8 % for ACT-8BIN (Table 7). However, the statistics for CN_{100} concentration shows that ACT-8BIN has NMBs of -58 and -67 % at the T0 and T1 sites, respectively, compared to WEX-8BIN NMBs of -55 and -65 %. We speculate that CN_{40} and CN_{100} concentrations are more dependent on condensational growth than on coagulation loss.

As can be seen from Tables 7 and 8, the use of ACT with 12 bins produces similar results for CN_{10} particles compared to the ACT-20BIN simulation, while also showing a visible improvement for CN_{40} and CN_{100} particles, reducing the CN_{40} overestimations by more than 40 % and the CN_{100} overestimations by more than 20 %. The correlation coefficients between observed and ACT-12BIN simulated values are similar to the R values between observed and ACT-20BIN.

Applying the same statistics for the particles collected during the G-1 aircraft flights, we note that both ACT-12BIN and ACT-20BIN overestimate CN_3 and CN_{10} particles (Table 9). However, the use of ACT-12BIN tends to decrease the overestimate of ACT-20BIN in predicted CN_3 concentration up to 71 % (10 June, morning flight). The statis-

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the nucleation rate impact the temporal and spatial variations of simulated CN and CCN number concentrations.

By comparing simulation results from the three empirical nucleation parameterizations explicitly treating nucleation using a 20-size bin sectional framework with measurements acquired during the CARES campaign, we showed that the parameterizations generally reproduced the overall observed spatial and temporal variability. However, the simulations with the activation (ACT), kinetic (KIN), and organics + H₂SO₄ (ORG) parameterizations overestimated the observed number concentration for particles with diameters > 10 nm (CN₁₀). In terms of the normalized mean bias (NMB), the best results were obtained using the ACT parameterization (NMB of 178 % over the entire analyzed period at the CARES T0 site), while the least favorable results were obtained for KIN (NMB of 255 % at the T1 site). This overestimation was reduced for larger submicron particles (CN₄₀ and CN₁₀₀). Measurements on a day with a new particle formation and growth event (NPE) indicated a diurnal cycle consisting of the onset of nucleation in the morning, subsequent particle growth, and particle number decrease during the afternoon. The cycles predicted from the ACT, KIN, and ORG simulations were similar to observed; the simulated onset of nucleation, however, was two hours earlier than observed and coincided with the onset of SOA formation. For a non-NPE day, our simulations reproduced the observed particle number concentration and the observed late afternoon shift in size distribution associated with the collapse of the boundary layer.

Using budget diagnostic terms we were able to establish how the nucleation rate parameterization affects the source/sink of particles, and also to explain some simulated features. The different nucleation parameterizations led to differences in the daily variability and magnitude of the budget terms associated with condensation, nucleation and coagulation for particles in the 1–10 nm size range (CN_{1–10}). For example, ACT and KIN exhibit almost the same variability at model locations corresponding to the two surface observation sites on a NPE day, yield one peak in CN_{1–10} particle number concentration around noon local time, and are highly correlated to the diurnal vari-

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ation of H_2SO_4 concentration. In contrast, ORG exhibits a different diurnal variation, showing two peaks (morning and afternoon) for most of the budget diagnostic terms for particles in 1–10 nm range. Thus, the different nucleation parameterizations lead to different results in both particle number and size distribution. However, these differences are greatly reduced for larger particles. We conclude that the loss processes are efficient and could explain the relatively similar biases of CN_{40} and CN_{100} given by the ACT, KIN and ORG parameterizations. Moreover, the budget diagnostic terms help us to understand modeled features such as the late afternoon shift in the observed and simulated size distribution on the non-NPE day.

We also analyzed the impact of the various nucleation parameterizations on CCN concentrations. We found that although the nucleation process could explain a percentage of CCN number concentration (up to 20–30 % at 0.5 % supersaturation), the choice of nucleation parameterization had no impact on the magnitude of the predicted CCN number concentration.

Using an intermediate number of size bins (12) to represent the particle size distribution and to explicitly capture the new particle formation process, we obtain CN_{10} results similar to those obtained using 20 size bins, and improvements (relative to observations) for CN_{40} and CN_{100} . Thus we can state that the use of a 12-size bin sectional framework represents a good compromise between computational time and the physico-chemical processes included in the WRF-Chem model. Simulations using the MOSAIC default 8-size bin (339 nm–10 μm diameter) sectional framework and either the ACT or the default Wexler nucleation parameterization underestimate the observed CN_{40} and CN_{100} concentrations and do not reproduce the observed daily variability as well as the 20 bin simulations.

We also analyzed the impact of size distribution on predicted aerosol mass concentration. We noticed that between 12 and 20 bin frameworks there is little variation over the entire analyzed period. Larger differences are found when we compare 8 and 20 bin frameworks, with up to 6 % more mass (in the case of NH_4) predicted by the 20 bin version.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Our analyses also suggest the ACT parameterization, where the nucleation rate is linearly proportional to the H_2SO_4 concentration, performs reasonably well, although sensitivity studies such as those employing uncertainty quantification (UQ) techniques would be needed to adjust empirical coefficients to better represent the magnitude of total PNC over California. Since the aerosol size distribution is a source of uncertainty in models, additional studies testing the sensitivity of the size distribution to primary particles may also improve the simulated particle number concentration and size distribution. Although the CARES campaign provided data to test and evaluate different new particle formation parameterizations, additional measurements such as H_2SO_4 and organic acid gas concentrations are necessary to better constrain the nucleation rate expressions.

Acknowledgements. We thank the numerous scientists, pilots, and other staff that contributed to the data collection during CARES. CARES was supported by the US Department of Energy's (DOE) Atmospheric Radiation Measurement (ARM) and Atmospheric System Research (ASR) programs. Funding for this research was provided by the US NOAA's Atmospheric Composition and Climate Program (NA11OAR4310160) and utilized resources provided by the Pacific Northwest National Laboratory (PNNL) Institutional Computing program. PNNL is operated for the US DOE by Battelle Memorial Institute.

References

- Andreae, M. O. and Rosenfeld, D.: Aerosol–cloud–precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth-Sci. Rev.*, 89, 13–41, 2008.
- Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., Hodnebrog, Ø., Denier van der Gon, H., and McFiggans, G.: Gaseous chemistry and aerosol mechanism developments for version 3.5.1 of the online regional model, WRF-Chem, *Geosci. Model Dev.*, 7, 2557–2579, doi:10.5194/gmd-7-2557-2014, 2014.
- Asmi, E., Kivekäs, N., Kerminen, V.-M., Komppula, M., Hyvärinen, A.-P., Hatakka, J., Viisanen, Y., and Lihavainen, H.: Secondary new particle formation in Northern Finland Pallas site

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

between the years 2000 and 2010, *Atmos. Chem. Phys.*, 11, 12959–12972, doi:10.5194/acp-11-12959-2011, 2011.

Betha, R., Spracklen, D. V., and Balasubramanian, R.: Observations of new aerosol particle formation in a tropical urban atmosphere, *Atmos. Environ.*, 71, 340–351, doi:10.1016/j.atmosenv.2013.01.049, 2013.

Boulon, J., Sellegri, K., Hervo, M., Picard, D., Pichon, J.-M., Fréville, P., and Laj, P.: Investigation of nucleation events vertical extent: a long term study at two different altitude sites, *Atmos. Chem. Phys.*, 11, 5625–5639, doi:10.5194/acp-11-5625-2011, 2011.

Boy, M., Karl, T., Turnipseed, A., Mauldin, R. L., Kosciuch, E., Greenberg, J., Rathbone, J., Smith, J., Held, A., Barsanti, K., Wehner, B., Bauer, S., Wiedensohler, A., Bonn, B., Kulmala, M., and Guenther, A.: New particle formation in the Front Range of the Colorado Rocky Mountains, *Atmos. Chem. Phys.*, 8, 1577–1590, doi:10.5194/acp-8-1577-2008, 2008.

Carter, W. P. L.: Implementation of the SAPRC-99 chemical mechanism into the models-3 framework, report to the United States Environmental Protection Agency, available at: <http://www.cert.ucr.edu/~carter/absts.htm#s99mod3> (last access: July 2015), 2000.

Cui, Y. Y., Hodzic, A., Smith, J. N., Ortega, J., Brioude, J., Matsui, H., Levin, E. J. T., Turnipseed, A., Winkler, P., and de Foy, B.: Modeling ultrafine particle growth at a pine forest site influenced by anthropogenic pollution during BEACHON-RoMBAS 2011, *Atmos. Chem. Phys.*, 14, 11011–11029, doi:10.5194/acp-14-11011-2014, 2014.

Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010.

Fast, J. D., Gustafson Jr., W. I., Easter Jr., R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology–chemistry–aerosol model, *J. Geophys. Res.-Atmos.*, 111, D21305, doi:10.1029/2005JD006721, 2006.

Fast, J. D., Gustafson Jr., W. I., Berg, L. K., Shaw, W. J., Pekour, M., Shrivastava, M., Barnard, J. C., Ferrare, R. A., Hostetler, C. A., Hair, J. A., Erickson, M., Jobson, B. T., Flowers, B., Dubey, M. K., Springston, S., Pierce, R. B., Dolislager, L., Pederson, J., and Zaveri, R. A.: Transport and mixing patterns over Central California during the carbona-

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ceous aerosol and radiative effects study (CARES), *Atmos. Chem. Phys.*, 12, 1759–1783, doi:10.5194/acp-12-1759-2012, 2012.

Fast, J. D., Allan, J., Bahreini, R., Craven, J., Emmons, L., Ferrare, R., Hayes, P. L., Hodzic, A., Holloway, J., Hostetler, C., Jimenez, J. L., Jonsson, H., Liu, S., Liu, Y., Metcalf, A., Middlebrook, A., Nowak, J., Pekour, M., Perring, A., Russell, L., Sedlacek, A., Seinfeld, J., Setyan, A., Shilling, J., Shrivastava, M., Springston, S., Song, C., Subramanian, R., Taylor, J. W., Vinoj, V., Yang, Q., Zaveri, R. A., and Zhang, Q.: Modeling regional aerosol and aerosol precursor variability over California and its sensitivity to emissions and long-range transport during the 2010 CalNex and CARES campaigns, *Atmos. Chem. Phys.*, 14, 10013–10060, doi:10.5194/acp-14-10013-2014, 2014.

Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled online chemistry within the WRF model, *Atmos. Environ.*, 39, 6957–6975, 2005.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.

Hoppel, W. A., Frick, G. M., Fitzgerald, J. W., and Larson, R. E.: Marine boundary layer measurements of new particle formation and the effects nonprecipitating clouds have on aerosol size distribution, *J. Geophys. Res.*, 99, 14443–14459, 1994.

Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: calculations with the AER radiative transfer models, *J. Geophys. Res.*, 113, D13103, doi:10.1029/2008JD009944, 2008.

IPCC 2007 – Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings between changes in the climate system and biogeochemistry, in: *Climate Change 2007: the Physical Science Basis*, contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK, and New York, NY, USA, 2007.

IPCC 2013 – Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S.,

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Stevens, B., and Zhang, X. Y.: Clouds and aerosols, in: *Climate Change 2013: the Physical Science Basis*, contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK, and New York, NY, USA, 2013.

Janjic, Z. I.: Nonsingular Implementation of the Mellor–Yamada Level 2.5 Scheme in the NCEP Meso model, Technical Report, National Centers for Environmental Prediction, Office Note No. 437, 61 pp., available at: <http://www.emc.ncep.noaa.gov/officenotes/newernotes/on437.pdf> (last access: July 2015), 2001.

Jeong, C. H., Hopke, P. K., Chalupa, D., and Utell, M.: Characteristics of nucleation and growth events of ultrafine particles measured in Rochester, NY, *Environ. Sci. Technol.*, 38, 1933–1940, 2004.

Jung, J. G., Pandis, S. N., Adams, P. J.: Evaluation of Nucleation Theories in a Sulfur-Rich Environment, *Aerosol Sci. Tech.*, 42, 7, 495–504, doi:10.1080/02786820802187085, 2008.

Jung, J., Fountoukis, C., Adams, P. J., and Pandis, S. N.: Simulation of in situ ultrafine particle formation in the eastern United States using PMCAMx-UF, *J. Geophys. Res.*, 115, D03203, doi:10.1029/2009JD012313, 2010.

Kain, J. S.: The Kain–Fritsch convective parameterization: an update, *J. Appl. Meteorol.*, 43, 170–181, doi:10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2, 2004.

Kazil, J., Harrison, R. G., and Lovejoy, E. R.: Tropospheric new particle formation and the role of ions, *Space Sci. Rev.*, 137, 241–255, 2008.

Kerminen, V.-M. and Kulmala, M.: Analytical formulae connecting the “real” and the “apparent” nucleation rate and the nuclei number concentration for atmospheric nucleation events, *J. Aerosol Sci.*, 33, 609–662, 2002.

Kerminen, V.-M., Paramonov, M., Anttila, T., Riipinen, I., Fountoukis, C., Korhonen, H., Asmi, E., Laakso, L., Lihavainen, H., Swietlicki, E., Svenningsson, B., Asmi, A., Pandis, S. N., Kulmala, M., and Petäjä, T.: Cloud condensation nuclei production associated with atmospheric nucleation: a synthesis based on existing literature and new results, *Atmos. Chem. Phys.*, 12, 12037–12059, doi:10.5194/acp-12-12037-2012, 2012.

Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D. Jud, W.,

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–433, doi:10.1038/nature10343, 2011.

Kristensson, A., Dal Maso, M., Swietlicki, E., Hussein, T., Zhou, J., Kerminen, V.-M., and Kulmala, M.: Characterization of new particle formation events at a background site in Southern Sweden: relation to air mass history, *Tellus B*, 60, 330–344, doi:10.1111/j.1600-0889.2008.00345.x, 2008.

Kuang, C., McMurry, P. H., McCormick, A. V., and Eisele, F. L.: Dependence of nucleation rates on sulfuric acid concentration in diverse atmospheric locations, *J. Geophys. Res.*, 113, D10209, doi:10.1029/2007JD009253, 2008.

Kuang, C., Riipinen, I., Sihto, S.-L., Kulmala, M., McCormick, A. V., and McMurry, P. H.: An improved criterion for new particle formation in diverse atmospheric environments, *Atmos. Chem. Phys.*, 10, 8469–8480, doi:10.5194/acp-10-8469-2010, 2010.

Kuang, C., Chen, M., Zhao, J., Smith, J., McMurry, P. H., and Wang, J.: Size and time-resolved growth rate measurements of 1 to 5 nm freshly formed atmospheric nuclei, *Atmos. Chem. Phys.*, 12, 3573–3589, doi:10.5194/acp-12-3573-2012, 2012.

Kulmala, M., Laaksonen, A., and Pirjola, L.: Parameterizations for sulphuric acid/water nucleation rates, *J. Geophys. Res.*, 103, 8301–8307, doi:10.1029/97JD03718, 1998.

Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Cluster activation theory as an explanation of the linear dependence between formation rate of 3nm particles and sulphuric acid concentration, *Atmos. Chem. Phys.*, 6, 787–793, doi:10.5194/acp-6-787-2006, 2006.

Levin, E. J. T., Prenni, A. J., Petters, M. D., Kreidenweis, S. M., Sullivan, R. C., Atwood, S. A., Ortega, J., DeMott, P. J., and Smith, J. N.: An annual cycle of size-resolved aerosol hygroscopicity at a forested site in Colorado, *J. Geophys. Res.*, 117, 06201, doi:10.1029/2011JD016854, 2012.

Makkonen, R., Asmi, A., Korhonen, H., Kokkola, H., Järvenoja, S., Räisänen, P., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., Järvinen, H., Lohmann, U., Bannartz, R., Feichter, J., and Kulmala, M.: Sensitivity of aerosol concentrations and cloud properties to

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nucleation and secondary organic distribution in ECHAM5-HAM global circulation model, Atmos. Chem. Phys., 9, 1747–1766, doi:10.5194/acp-9-1747-2009, 2009.

Matsui, H., Koike, M., Kondo, Y., Takegawa, N., Wiedensohler, A., Fast, J. D., and Zaveri, R. A.: Impact of new particle formation on the concentrations of aerosols and cloud condensation nuclei around Beijing, J. Geophys. Res., 116, D19208, doi:10.1029/2011JD016025, 2011.

Matsui, H., Koike, M., Takegawa, N., Kondo, Y., Takami, A., Takamura, T., Yoon, S., Kim, S.-W., Lim, H.-C., and Fast, J. D.: Spatial and temporal variations of new particle formation in East Asia using an NPF-explicit WRF-chem model: north–south contrast in new particle formation frequency, J. Geophys. Res.-Atmos., 118, 11647–11663, doi:10.1002/jgrd.50821, 2013.

McMurry, P. H.: Photochemical aerosol formation from SO₂: a theoretical analysis of smog chamber data, J. Colloid Interf. Sci., 78, 513–527, doi:10.1016/0021-9797(80)90589-5, 1980.

McMurry, P. H. and Friedlander, S. K.: New particle formation in the presence of an aerosol, Atmos. Environ., 13, 1635–1651, 1979.

McMurry, P. H., Woo, K. S., Weber, R., Chen, D.-R., and Pui, D. Y. H.: Size distributions of 3–10 nm atmospheric particles: implications for nucleation mechanisms, Philos. T. Roy. Soc. A, 358, 2625–2642, doi:10.1098/rsta.2000.0673, 2000.

Merikanto, J., Napari, I., Vehkamäki, H., Anttila, T., and Kulmala, M.: New parameterization of sulfuric acid–ammonia–water ternary nucleation rates at tropospheric conditions, J. Geophys. Res., 112, D15207, doi:10.1029/2006JD007977, 2007.

Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, Atmos. Chem. Phys., 9, 8601–8616, doi:10.5194/acp-9-8601-2009, 2009.

Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the role of organics in aerosol particle formation under atmospheric conditions, P. Natl. Acad. Sci. USA, 107, 6646–6651, doi:10.1073/pnas.0911330107, 2010.

Morrison, H., Thompson, G., and Tatarskii, V.: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: comparison of one- and two-moment schemes, Mon. Weather Rev., 137, 991–1007, doi:10.1175/2008MWR2556.1, 2009.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Napari, I., Noppel, M., Vehkamäki, H., and Kulmala, M.: Parameterization of ternary nucleation rates for $\text{H}_2\text{SO}_4\text{-NH}_3\text{-H}_2\text{O}$ vapors, *J. Geophys. Res.*, 107, 4381, doi:10.1029/2002JD002132, 2002.
- O'Dowd, C. D. and Hoffman, T.: Coastal new particle formation: a review of the current state-of-the-art, *Environ. Chem.*, 2, 245–255, doi:10.1071/EN05077, 2005.
- Paasonen, P., Nieminen, T., Asmi, E., Manninen, H. E., Petäjä, T., Plass-Dülmer, C., Flentje, H., Birmili, W., Wiedensohler, A., Hörrak, U., Metzger, A., Hamed, A., Laaksonen, A., Facchini, M. C., Kerminen, V.-M., and Kulmala, M.: On the roles of sulphuric acid and low-volatility organic vapours in the initial steps of atmospheric new particle formation, *Atmos. Chem. Phys.*, 10, 11223–11242, doi:10.5194/acp-10-11223-2010, 2010.
- Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation and primary emission rates, *Atmos. Chem. Phys.*, 9, 1339–1356, doi:10.5194/acp-9-1339-2009, 2009.
- Pierce, J. R., Leaitch, W. R., Liggio, J., Westervelt, D. M., Wainwright, C. D., Abbatt, J. P. D., Ahlm, L., Al-Basheer, W., Cziczo, D. J., Hayden, K. L., Lee, A. K. Y., Li, S.-M., Russell, L. M., Sjostedt, S. J., Strawbridge, K. B., Travis, M., Vlasenko, A., Wentzell, J. J. B., Wiebe, H. A., Wong, J. P. S., and Macdonald, A. M.: Nucleation and condensational growth to CCN sizes during a sustained pristine biogenic SOA event in a forested mountain valley, *Atmos. Chem. Phys.*, 12, 3147–3163, doi:10.5194/acp-12-3147-2012, 2012.
- Place Jr., P. F., Ziemba, L. D., and Griffin, R. J.: Observations of nucleation-mode particle events and size distributions at a rural New England site, *Atmos. Environ.*, 44, 88–94, 2010.
- Reddington, C. L., Carslaw, K. S., Spracklen, D. V., Frontoso, M. G., Collins, L., Merikanto, J., Minikin, A., Hamburger, T., Coe, H., Kulmala, M., Aalto, P., Flentje, H., Plass-Dülmer, C., Birmili, W., Wiedensohler, A., Wehner, B., Tuch, T., Sonntag, A., O'Dowd, C. D., Jennings, S. G., Dupuy, R., Baltensperger, U., Weingartner, E., Hansson, H.-C., Tunved, P., Laj, P., Selenković, K., Boulon, J., Putaud, J.-P., Gruening, C., Swietlicki, E., Roldin, P., Henzing, J. S., Moerman, M., Mihalopoulos, N., Kouvarakis, G., Ždímal, V., Zíková, N., Marinoni, A., Bonasoni, P., and Duchi, R.: Primary versus secondary contributions to particle number concentrations in the European boundary layer, *Atmos. Chem. Phys.*, 11, 12007–12036, doi:10.5194/acp-11-12007-2011, 2011.
- Riipinen, I., Sihto, S.-L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinilä, K., Kerminen, V.-M., Laaksonen, A., and Lehtinen, K. E. J.: Connections between atmospheric sulphuric acid and new particle formation during QUEST III–IV campaigns in

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heidelberg and Hyytiälä, Atmos. Chem. Phys., 7, 1899–1914, doi:10.5194/acp-7-1899-2007, 2007.

Ritter, M., Müller, M. D., Tsai, M.-Y., and Parlow, E.: Air pollution modeling over very complex terrain: an evaluation of WRF-Chem over Switzerland for two 1 year periods, Atmos. Res., 132–133, 209–222, 2013.

Schobesberger, S., Junninen, H., Bianchi, F., Lönn, G., Ehn, M., Lehtipalo, K., Dommen, J., Ehrhart, S., Ortega, I. K., Franchin, A., Nieminen, T., Riccobono, F., Hutterli, M., Duplissy, J., Almeida, J., Amorim, A., Breitenlechner, M., Downard, A. J., Dunne, E. M., Flanagan, R. C., Kajos, M., Keskinen, H., Kirkby, J., Kupc, A., Kürten, A., Kurtén, T., Laaksonen, A., Mathot, S., Onnela, A., Praplan, A. P., Rondo, L., Santos, F. D., Schallhart, S., Schnitzhofer, R., Sipilä, M., Tomé, A., Tsagkogeorgas, G., Vehkamäki, H., Wimmer, D., Baltensperger, U., Carslaw, K. S., Curtius, J., Hansel, A., Petäjä, T., Kulmala, M., Donahue, N. M., and Worsnop, D. R.: Molecular understanding of atmospheric particle formation from sulfuric acid and large oxidized organic molecules, P. Natl. Acad. Sci. USA, 110, 17223–17228, doi:10.1073/pnas.1306973110, 2013.

Sem, G. J.: Design and performance characteristics of three continuous-flow condensation particle counters: a summary, Atmos. Res., 62, 267–294, doi:10.1016/S0169-8095(02)00014-5, 2002.

Setyan, A., Zhang, Q., Merkel, M., Knighton, W. B., Sun, Y., Song, C., Shilling, J. E., Onasch, T. B., Herndon, S. C., Worsnop, D. R., Fast, J. D., Zaveri, R. A., Berg, L. K., Wiedensohler, A., Flowers, B. A., Dubey, M. K., and Subramanian, R.: Characterization of submicron particles influenced by mixed biogenic and anthropogenic emissions using high-resolution aerosol mass spectrometry: results from CARES, Atmos. Chem. Phys., 12, 8131–8156, doi:10.5194/acp-12-8131-2012, 2012.

Setyan, A., Song, C., Merkel, M., Knighton, W. B., Onasch, T. B., Canagaratna, M. R., Worsnop, D. R., Wiedensohler, A., Shilling, J. E., and Zhang, Q.: Chemistry of new particle growth in mixed urban and biogenic emissions – insights from CARES, Atmos. Chem. Phys., 14, 6477–6494, doi:10.5194/acp-14-6477-2014, 2014.

Shaw, W., Allwine, K. J., Fritz, B. G., Rutz, F. C., Rishel, J. P., and Chapman, E. G.: An evaluation of the wind erosion module in DUSTRAN, Atmos. Environ., 42, 1907–1921, 2008.

Shrivastava, M., Fast, J., Easter, R., Gustafson Jr., W. I., Zaveri, R. A., Jimenez, J. L., Saide, P., and Hodzic, A.: Modeling organic aerosols in a megacity: comparison of simple and complex

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



representations of the volatility basis set approach, *Atmos. Chem. Phys.*, 11, 6639–6662, doi:10.5194/acp-11-6639-2011, 2011.

5 Sihto, S.-L., Kulmala, M., Kerminen, V.-M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A., and Lehtinen, K. E. J.: Atmospheric sulphuric acid and aerosol formation: implications from atmospheric measurements for nucleation and early growth mechanisms, *Atmos. Chem. Phys.*, 6, 4079–4091, doi:10.5194/acp-6-4079-2006, 2006.

10 Smith, J. N., Dunn, M. J., VanReken, T. M., Iida, K., Stolzenburg, M. R., McMurry, P. H., and Huey, L. G.: Chemical composition of atmospheric nanoparticles formed from nucleation in Tecamac, Mexico: evidence for an important role for organic species in nanoparticle growth, *Geophys. Res. Lett.*, 35, L04808, doi:10.1029/2007GL032523, 2008.

15 Spracklen, D. V., Pringle, K. J., Carslaw, K. S., Chipperfield, M. P., and Mann, G. W.: A global off-line model of size-resolved aerosol microphysics: I. Model development and prediction of aerosol properties, *Atmos. Chem. Phys.*, 5, 2227–2252, doi:10.5194/acp-5-2227-2005, 2005.

20 Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Mann, G. W., and Sihto, S.-L.: The contribution of boundary layer nucleation events to total particle concentrations on regional and global scales, *Atmos. Chem. Phys.*, 6, 5631–5648, doi:10.5194/acp-6-5631-2006, 2006.

25 Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I., Merikanto, J., Mann, G. W., Chipperfield, M. P., Wiedensohler, A., Birmili, W., and Lihavainen, H.: Contribution of particle formation to global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 35, L06808, doi:10.1029/2007GL033038, 2008.

30 Spracklen, D. V., Carslaw, K. S., Merikanto, J., Mann, G. W., Reddington, C. L., Pickering, S., Ogren, J. A., Andrews, E., Baltensperger, U., Weingartner, E., Boy, M., Kulmala, M., Laakso, L., Lihavainen, H., Kivekäs, N., Komppula, M., Mihalopoulos, N., Kouvarakis, G., Jennings, S. G., O'Dowd, C., Birmili, W., Wiedensohler, A., Weller, R., Gras, J., Laj, P., Selligri, K., Bonn, B., Krejci, R., Laaksonen, A., Hamed, A., Minikin, A., Harrison, R. M., Talbot, R., and Sun, J.: Explaining global surface aerosol number concentrations in terms of primary emissions and particle formation, *Atmos. Chem. Phys.*, 10, 4775–4793, doi:10.5194/acp-10-4775-2010, 2010.

Turco, R. P., Zhao, J.-X., and Yu, F.: A new source of tropospheric aerosols: Ion-ion recombination, *Geophys. Res. Lett.*, 25, 635–638, doi:10.1029/98GL00253, 1998.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E. J., Timmreck, C., Noppel, M., and Laaksonen, A.: An improved parameterization for sulfuric acid-water nucleation rates for tropospheric and stratospheric conditions, *J. Geophys. Res.*, 107, 4622, doi:10.1029/2002JD002184, 2002.

5 Venzac, H., Sellegri, K., Villani, P., Picard, D., and Laj, P.: Seasonal variation of aerosol size distributions in the free troposphere and residual layer at the puy de Dôme station, France, *Atmos. Chem. Phys.*, 9, 1465–1478, doi:10.5194/acp-9-1465-2009, 2009.

Westervelt, D. M., Pierce, J. R., and Adams, P. J.: Analysis of feedbacks between nucleation rate, survival probability and cloud condensation nuclei formation, *Atmos. Chem. Phys.*, 14, 5577–5597, doi:10.5194/acp-14-5577-2014, 2014.

10 Wexler, A. S., Lurmann, F. W., and Seinfeld, J. H.: Modeling urban and regional aerosols. I. Model development, *Atmos. Environ.*, 28, 531–546, doi:10.1016/1352-2310(94)90129-5, 1994.

15 Wiedensohler, A., Chen, Y. F., Nowak, A., Wehner, B., Achtert, P., Berghof, M., Birmili, W., Wu, Z. J., Hu, M., Zhu, T., Takegawa, N., Kita, K., Kondo, Y., Lou, S. R., Hofzumahaus, A., Holland, F., Wahner, A., Gunthe, S. S., Rose, D., Su, H., and Pöschl, U.: Rapid aerosol particle growth and increase of cloud condensation nucleus activity by secondary aerosol formation and condensation: a case study for regional air pollution in northeastern China, *J. Geophys. Res.*, 114, 00G08, doi:10.1029/2008JD010884, 2009.

20 Yli-Juuti, T., Barsanti, K., Hildebrandt Ruiz, L., Kieloaho, A.-J., Makkonen, U., Petäjä, T., Ruuskanen, T., Kulmala, M., and Riipinen, I.: Model for acid-base chemistry in nanoparticle growth (MABNAG), *Atmos. Chem. Phys.*, 13, 12507–12524, doi:10.5194/acp-13-12507-2013, 2013.

25 Yu, F. and Turco, R. P.: Ultrafine aerosol formation via ion-mediated nucleation, *Geophys. Res. Lett.*, 27, 883–886, 2000.

Yu, F. and Turco, R. P.: From molecular clusters to nanoparticles: role of ambient ionization in tropospheric aerosol formation, *J. Geophys. Res.*, 106, 4797–4814, 2001.

30 Yu, F., Wang, Z., Luo, G., and Turco, R.: Ion-mediated nucleation as an important global source of tropospheric aerosols, *Atmos. Chem. Phys.*, 8, 2537–2554, doi:10.5194/acp-8-2537-2008, 2008.

Yue, D. L., Hu, M., Zhang, R. Y., Wu, Z. J., Su, H., Wang, Z. B., Peng, J. F., He, L. Y., Huang, X. F., Gong, Y. G., and Wiedensohler, A.: Potential contribution of new particle formation to cloud condensation nuclei in Beijing, *Atmos. Environ.*, 45, 6070–6077, 2011.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Zaveri, R. A., Easter Jr., R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res.-Atmos.*, 113, D13204, doi:10.1029/2007JD008782, 2008.
- Zaveri, R. A., Shaw, W. J., Cziczo, D. J., Schmid, B., Ferrare, R. A., Alexander, M. L., Alexandrov, M., Alvarez, R. J., Arnott, W. P., Atkinson, D. B., Baidar, S., Banta, R. M., Barnard, J. C., Beranek, J., Berg, L. K., Brechtel, F., Brewer, W. A., Cahill, J. F., Cairns, B., Cappa, C. D., Chand, D., China, S., Comstock, J. M., Dubey, M. K., Easter, R. C., Erickson, M. H., Fast, J. D., Floerchinger, C., Flowers, B. A., Fortner, E., Gaffney, J. S., Gilles, M. K., Gorkowski, K., Gustafson, W. I., Gyawali, M., Hair, J., Hardesty, R. M., Harworth, J. W., Herndon, S., Hiranuma, N., Hostetler, C., Hubbe, J. M., Jayne, J. T., Jeong, H., Jobson, B. T., Kassianov, E. I., Kleinman, L. I., Kluzek, C., Knighton, B., Kolesar, K. R., Kuang, C., Kubátová, A., Langford, A. O., Laskin, A., Laulainen, N., Marchbanks, R. D., Mazzoleni, C., Mei, F., Moffet, R. C., Nelson, D., Obland, M. D., Oetjen, H., Onasch, T. B., Ortega, I., Ottaviani, M., Pekour, M., Prather, K. A., Radney, J. G., Rogers, R. R., Sandberg, S. P., Sedlacek, A., Senff, C. J., Senum, G., Setyan, A., Shilling, J. E., Shrivastava, M., Song, C., Springston, S. R., Subramanian, R., Suski, K., Tomlinson, J., Volkamer, R., Wallace, H. W., Wang, J., Weickmann, A. M., Worsnop, D. R., Yu, X.-Y., Zelenyuk, A., and Zhang, Q.: Overview of the 2010 Carbonaceous Aerosols and Radiative Effects Study (CARES), *Atmos. Chem. Phys.*, 12, 7647–7687, doi:10.5194/acp-12-7647-2012, 2012.
- Zhang, Y., Liu, P., Liu, X.-H., Jacobson, M. Z., McMurry, P. H., Yu, F., Yu, S., and Schere, K. L.: A comparative study of nucleation parameterizations: 2. Three dimensional model application and evaluation, *J. Geophys. Res.*, 115, D20213, doi:10.1029/2010JD014151, 2010a.
- Zhang, Y., McMurry, P. H., Yu, F., and Jacobson, M. Z.: A comparative study of nucleation parameterizations: 1. examination and evaluation of the formulations, *J. Geophys. Res.*, 115, D20212, doi:10.1029/2010JD014150, 2010b.
- Ziemba, L. D., Griffin, R. J., and Talbot, R. W.: Observations of elevated particle number concentration events at a rural site in northern New England, *J. Geophys. Res.*, 111, D23S24, doi:10.1029/2006JD007607, 2006.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Observed mean and simulation summary statistics for aerosol number concentrations for particle diameters ≥ 10 nm (CN_{10}), ≥ 40 nm (CN_{40}), and ≥ 100 nm (CN_{100}). The normalized mean bias (NMB) and correlation coefficient (R) are calculated between simulated and observed number concentration at the CARES T0 site during the 7–16 June 2010 period.

Analyzed period	Observed mean			Model experiment	NMB (%)			R		
	CN_{10}	CN_{40}	CN_{100}		CN_{10}	CN_{40}	CN_{100}	CN_{10}	CN_{40}	CN_{100}
Entire period	8576	2748	366	ACT	184	66	78	0.71	0.56	0.53
				KIN	255	67	65	0.69	0.56	0.50
				ORG	233	71	68	0.66	0.57	0.51
				8BIN	–	–34	–55	–	0.56	0.32
South-westerly flow	9885	3328	435	ACT	191	60	86	0.71	0.53	0.45
				KIN	278	63	71	0.68	0.52	0.40
				ORG	228	66	75	0.67	0.54	0.40
				8BIN	–	–68	–47	–	0.15	0.53
Northerly flow	6162	1686	202	ACT	168	87	49	0.48	0.17	0.08
				KIN	193	83	47	0.41	0.16	0.07
				ORG	247	88	42	0.44	0.20	0.03
				8BIN	–	–66	–48	–	0.24	0.30

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Table 2. Same as Table 1, except for the CARES T1 site.

Analyzed period	Observed mean			Model experiment	NMB (%)			R		
	CN ₁₀	CN ₄₀	CN ₁₀₀		CN ₁₀	CN ₄₀	CN ₁₀₀	CN ₁₀	CN ₄₀	CN ₁₀₀
Entire period				ACT	178	31	25	0.74	0.70	0.51
				KIN	231	36	13	0.64	0.49	0.55
				ORG	239	36	16	0.67	0.69	0.51
				8BIN	–	–51	–65	–	0.57	0.32
South-westerly flow	7960	4073	691	ACT	187	35	33	0.71	0.66	0.33
				KIN	347	27	6	0.62	0.43	0.41
				ORG	256	41	23	0.59	0.63	0.32
				8BIN	–	–58	–62	–	0.51	0.03
Northerly flow	4429	1878	289	ACT	129	19	–12	0.67	0.11	–0.13
				KIN	278	68	7	0.41	0.06	0.07
				ORG	158	18	–16	0.68	0.10	–0.16
				8BIN	–	–40	–77	–	0.06	–0.16



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Selected budget diagnostic terms ($\text{cm}^{-3}\text{h}^{-1}$) averaged over 04:00–16:00 PST for all NPE days at the T0 and T1 sites. The CN_{1-10} Transport + Deposition term includes advection, turbulent mixing, and dry deposition.

NPE event	Experiment	CN_{1-10} Transport + Deposition	CN_{1-10} Condensation	CN_{1-10} Nucleation	CN_{1-10} Coagulation	CN_{10-100} Condensation + Coagulation
07 Jun 2010	ACT	−467	−6165	64 200	−56 662	3113
	KIN	34 589	−28 112	568 746	−566 077	7851
	ORG	−111 977	−34 309	2 295 385	−2 127 692	8648
08 Jun 2010	ACT	30 028	−33 002	493 769	−498 615	14 378
	KIN	−706 731	−43 362	22 285 077	−21 623 692	11 718
	ORG	−142 762	−19 701	1 323 769	−1 147 923	5311
09 Jun 2010	ACT	13 514	−44 665	136 923	−105 308	37 167
	KIN	−78 154	−85 439	1 812 308	−1 646 923	71 408
	ORG	−163 368	−56 626	1 868 154	−1 646 000	44 537
10 Jun 2010	ACT	38 646	−53 764	45 238	−29 546	50 833
	KIN	7144	−80 066	307 523	−229 331	78 095
	ORG	−194 592	−77 118	630 231	−352 538	68 368
11 Jun 2010	ACT	37 617	−50 877	25 515	−11 409	49 099
	KIN	8629	−58 040	100 538	−51 247	56 856
	ORG	−282 259	−84 154	687 231	−310 454	68 239
14 Jun 2010	ACT	25 291	−31 878	608 846	−594 692	10 094
	KIN	−1 036 169	−50 777	28 470 000	−27 343 923	18 983
	ORG	−72 954	−21 925	900 462	−797 923	8930
15 Jun 2010	ACT	5769	−27 877	267 462	−243 431	11 901
	KIN	−418 362	−25 031	7 323 846	−7 022 969	25 423
	ORG	−77 023	−26 138	1 231 000	−1 129 462	17 232
16 Jun 2010	ACT	54 250	−79 367	84 167	−56 858	74 135
	KIN	−159 425	−49 981	832 727	−615 400	46 482
	ORG	−142 382	−72 017	425 417	−207 458	64 203

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. Observed mean and simulation summary statistics for aerosol number concentrations for particle diameters ≥ 3 nm (CN_3) and CN_{10} . The observed mean, normalized mean bias (NMB) and correlation coefficient (R) are calculated between simulated and observed number concentration collected during the CARES G-1 aircraft flights.

PNC	Flight	Observed mean	NMB (%)			R		
			ACT	KIN	ORG	ACT	KIN	ORG
CN_3	08 Jun, morning (08a)	20 472	157	614	227	0.64	0.67	0.74
	08 Jun, afternoon (08b)	13 990	274	603	204	0.84	0.79	0.67
	10 Jun, morning (10a)	17 392	212	365	212	0.22	0.19	-0.05
	12 Jun, morning (12a)	12 340	198	162	243	0.40	0.29	0.50
	12 Jun, afternoon (12b)	14 459	184	307	359	-0.06	0.10	0.32
	14 Jun, morning (14a)	21 913	259	824	368	0.56	0.57	0.51
	15 Jun, morning (15a)	24 800	250	845	296	0.68	0.69	0.61
	15 Jun, afternoon (15b)	14 800	260	776	330	0.75	0.46	0.38
	All flights	17 564	226	608	282	0.52	0.51	0.48
CN_{10}	08 Jun, morning (08a)	11 588	90	173	118	0.57	0.58	0.58
	08 Jun, afternoon (08b)	10 161	162	219	168	0.80	0.81	0.77
	10 Jun, morning (10a)	10 804	95	138	92	-0.03	0.03	-0.12
	12 Jun, morning (12a)	6 019	69	105	87	0.44	0.49	0.55
	12 Jun, afternoon (12b)	7 940	128	128	105	-0.05	-0.04	-0.04
	14 Jun, morning (14a)	11 999	221	365	273	0.72	0.70	0.60
	15 Jun, morning (15a)	13 601	216	338	250	0.71	0.70	0.61
	15 Jun, afternoon (15b)	12 625	230	296	068	0.48	0.14	0.29
	All flights	10 641	162	239	184	0.54	0.51	0.49

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 5. Summary statistics for CCN number concentration at several super saturations ($SS = 0.1, 0.2, 0.35, 0.5\%$). The observed mean, normalized mean bias (NMB) and correlation coefficient (R) are calculated between simulated and observed number concentration at the T0 site during the 7–16 June 2010 period.

Observed mean				Experiment	NMB (%)				R			
$SS_{0.1}$	$SS_{0.2}$	$SS_{0.35}$	$SS_{0.5}$		$SS_{0.1}$	$SS_{0.2}$	$SS_{0.35}$	$SS_{0.5}$	$SS_{0.1}$	$SS_{0.2}$	$SS_{0.35}$	$SS_{0.5}$
105	329	729	1067	ACT	−49	−34	−5	29	0.39	0.23	0.39	0.30
				KIN	−49	−34	−9	27	0.39	0.22	0.35	0.28
				ORG	−49	−33	−8	26	0.38	0.23	0.36	0.29

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 6. Same as Table 5, except for the T1 site.

Observed mean				Experiment	NMB (%)				<i>R</i>			
SS _{0.1}	SS _{0.2}	SS _{0.35}	SS _{0.5}		SS _{0.1}	SS _{0.2}	SS _{0.35}	SS _{0.5}	SS _{0.1}	SS _{0.2}	SS _{0.35}	SS _{0.5}
109	333	729	1181	ACT	-53	-29	6	26	0.62	0.40	0.43	0.58
				KIN	-53	-32	-1	22	0.64	0.39	0.40	0.57
				ORG	-52	-33	1	23	0.63	0.40	0.41	0.58

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 7. Summary statistics for CN_{10} , CN_{40} and CN_{100} when using the ACT parameterization in a sectional framework with different numbers of particle size bins together with statistics for default configuration of WRF-Chem (WEX-8BIN). It should be noted that WEX-8BIN statistical results are the same as the 8BIN simulation presented in Tables 1 and 2. The observed mean, normalized mean bias (NMB) and correlation coefficient (R) are calculated between simulated and observed number concentration at the T0 site during the 7–16 June 2010 period.

Observed mean			Experiment	NMB (%)			R		
CN_{10}	CN_{40}	CN_{100}		CN_{10}	CN_{40}	CN_{100}	CN_{10}	CN_{40}	CN_{100}
8576	2748	366	WEX-8BIN	–	–34	–55	–	0.56	0.32
			ACT-8BIN	–	–8	–58	–	0.32	0.24
			ACT-12BIN	188	18	49	0.72	0.46	0.48
			ACT-20BIN	184	66	78	0.71	0.56	0.53

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 8. Same as Table 7, except for the T1 site.

Observed mean			Experiment	NMB (%)			<i>R</i>		
CN ₁₀	CN ₄₀	CN ₁₀₀		CN ₁₀	CN ₄₀	CN ₁₀₀	CN ₁₀	CN ₄₀	CN ₁₀₀
6389	3121	518	WEX-8BIN	–	–51	–65	–	0.57	0.32
			ACT-8BIN	–	–17	–67	–	0.43	0.36
			ACT-12BIN	187	–8	4	0.72	0.63	0.48
			ACT-20BIN	178	31	25	0.74	0.70	0.51

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Table 9. Summary statistics for CN₃ and CN₁₀ when using the ACT parameterization in a sectional framework with different numbers of particle size bins. The observed mean, normalized mean bias (NMB) and correlation coefficient (R) are calculated between simulated and observed number concentration collected during the CARES G-1 aircraft flights.

PNC	Flights	Observed mean	NMB (%)		R	
			ACT-12BIN	ACT-20BIN	ACT-12BIN	ACT-20BIN
CN ₃	08 Jun, morning (08a)	20 472	151	157	0.68	0.64
	08 Jun, afternoon (08b)	13 990	223	274	0.75	0.84
	10 Jun, morning (10a)	17 392	141	212	0.21	0.22
	12 Jun, morning (12a)	12 340	112	198	0.35	0.40
	12 Jun, afternoon (12b)	14 459	175	184	0.11	-0.06
	14 Jun, morning (14a)	21 913	213	259	0.61	0.56
	15 Jun, morning (15a)	24 800	242	250	0.64	0.68
	15 Jun, afternoon (15b)	14 800	275	260	0.73	0.75
	All flights	17 564	194	226	0.37	0.52
CN ₁₀	08 Jun, morning (08a)	11 588	19	90	0.62	0.57
	08 Jun, afternoon (08b)	10 161	165	162	0.74	0.80
	10 Jun, morning (10a)	10 804	89	95	-0.11	-0.03
	12 Jun, morning (12a)	6019	61	69	0.39	0.44
	12 Jun, afternoon (12b)	7940	131	128	-0.02	-0.05
	14 Jun, morning (14a)	11 999	219	221	0.72	0.72
	15 Jun, morning (15a)	13 601	230	216	0.68	0.71
	15 Jun, afternoon (15b)	12 625	260	230	0.73	0.48
	All flights	10 641	169	162	0.38	0.54

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 1. Observed and simulated time series of particle number concentration at the T0 site for (a) CN₁₀; (b) CN₄₀; (c) CN₁₀₀. Gray shading indicates nighttime and the arrow denotes the period of northerly synoptic flow associated with the passage of a trough over California. All times are local Pacific Standard Time (PST).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 2. Same as Fig. 1, except for the T1 site.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 3. Number size distribution as a function of geometric diameter and time at the T0 site on 8 June 2010 and from **(a)** observations and the **(b)** ACT, **(c)** KIN, and **(d)** ORG simulations. The vectors along the bottom of each panel represent the observed and calculated winds at 10 m a.g.l. at the T0 site. The maximum wind speed at about 15:00 PST in **(b–d)** is 6 m s^{-1} . The black diamonds represent the observed **(a)** and simulated **(b–d)** boundary layer heights.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 4. Same as Fig. 3, except for the T1 site.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 5. Particle production from the ACT and ORG parameterizations at the T0 and T1 sites on 8 June 2010 together with modeled BL height (circled line). The dashed line shows the altitude of maximum predicted number of small particles.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 6. Simulated aerosol mass concentrations at the T0 site on 8 June 2010 predicted by the ACT parameterization.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 7. (a) Nucleation budget term ($\text{cm}^{-3} \text{h}^{-1}$) averaged over first five model layers from 05:00 to 09:00 PST 8 June 2010 from the ACT simulation, together with a backward trajectory from 08:00 PST on 7 June to 09:00 PST on 8 June 2010; the red dots denote the hourly locations of the air mass prior to its arrival at the T0; (b) Altitude of the air mass (black line) together with particle number concentration in 1–10 nm diameter range (red line), 10–40 nm range (green line), 40–100 nm range (blue line) and nucleation rate (orange circled line); the gray area denotes the terrain height.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 8. Budget diagnostic terms for 1–10 nm particles at the T0 site on 8 June: **(a)** dry deposition; **(b)** transport; **(c)** condensation; **(d)** nucleation; **(e)** coagulation; **(f)** total tendency; **(g)** particle number concentration; and **(h)** tendency of particles coming from nucleation and coagulation processes. The left-hand side (black) vertical axes are for ACT and ORG. The right-hand side (blue) axes are for KIN, which gave larger budget terms.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 9. Budget diagnostic terms for 10–40 nm particles at the T0 site on 8 June: **(a)** deposition; **(b)** transport; **(c)** condensation; **(d)** emissions; **(e)** coagulation; **(f)** total tendency; and **(g)** particle number concentration.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 10. Budget diagnostic terms for 40–100 nm particles at the T0 site on 8 June: **(a)** deposition; **(b)** transport; **(c)** condensation; **(d)** emissions; **(e)** coagulation; **(f)** total tendency; and **(g)** particle number concentration.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.





Figure 11. Modeled H_2SO_4 (red line), and organic gas concentrations (blue line) at the T0 site on 8 June 2010 (circled line) and 12 June 2010 (solid line).

Modeling particle nucleation and growth over northern California

A. Lupascu et al.



6 PST 14 PST 19 PST

Figure 12. Particle number distribution for observations (OBS) and the ACT, KIN, and ORG simulations at the T0 (a, b, c and d) and T1 (e, f, g and h) sites at 10:00 PST (black line), 14:00 PST (blue line) and 19:00 PST (red line). The missing line segments in the simulations correspond to particle number lower than 10 cm^{-3} .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 13. Number size distribution as a function of geometric diameter and time at the T0 site on 12 June 2010 from (a) observations and the (b) ACT, (c) KIN, and (d) ORG simulations. The vectors along the bottom of each panel represent the observed and calculated winds at 10 m a.g.l. at the T0 site. The maximum wind speed at about 12:00 PST in (b–d) is 10.4 m s^{-1} . The black diamonds represent the observed (a) and simulated (b–d) boundary layer heights.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 14. Same as Fig. 13, except for the T1 site.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 15. Budget diagnostic terms for 1–10 nm particles at the T0 site on 12 June: **(a)** deposition; **(b)** transport; **(c)** condensation; **(d)** nucleation; **(e)** coagulation; **(f)** total tendency; **(g)** particle number concentration in 1 to 10 nm range; and **(h)** tendency of particles coming from nucleation and coagulation processes.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 16. Budget diagnostic terms for 10–40 nm particles at the T0 site on 12 June: **(a)** deposition; **(b)** transport; **(c)** condensation; **(d)** emissions; **(e)** coagulation; **(f)** total tendency; and **(g)** particle number concentration.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 17. Budget diagnostic terms for 40–100 nm particles at the T0 site on 12 June: **(a)** deposition; **(b)** transport; **(c)** condensation; **(d)** emissions; **(e)** coagulation; **(f)** total tendency; and **(g)** particle number concentration.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 18. Boxplot for observed and simulated CN_3 and CN_{10} at the surface to 500 m altitude (a and e); 500–1000 m altitude (b and f) 1000–1500 m altitude (c and g); and 1500–2000 m altitude (d and h) for the 8 June 2010 afternoon G-1 flight. Observations are in black, ACT simulation values are in green, KIN in blue, and ORG in red. The line in the middle of each box is the median, while the boxes represent the 25th and 75th percentiles, and “whiskers” the 5th and 95th percentiles.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 19. Same as Fig. 18, except for the 12 June 2010 afternoon G-1 flight.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 20. Observed and simulated time series of CCN number concentration at the T0 site for **(a)** SS = 0.1 %; **(b)** SS = 0.2 %; **(c)** SS = 0.35 % **(e)** SS = 0.5 % together with **(e)** CN₁₀ concentration. Gray shading area indicates nighttime while the black arrow denotes the northerly synoptic flow associated with the passage of troughs over California.

Modeling particle nucleation and growth over northern California

A. Lupascu et al.




Figure 21. Same as Fig. 20, except for the T1 site.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

