

Letter to the Editor

2 Dear Dr. Sullivan,

3 We have striven to take on board as many of the reviewers' comments as possible. The major
4 changes to the article are as follows:

1. Conclusion – this is now bulletized to make the new scientific findings clearer, and we have added a table to summarize the Fennec aircraft publications to minimize the text in the conclusion.
2. Figures – 17 of 18 figures have been altered, many significantly. Keys have been added, lines and text made bolder, titles/colour bar titles added. With a publication of this size and number of authors, consistency across all figures is very challenging but we have done our best to achieve this, and we hope that the figures are now of sufficient clarity. They are attached in a pdf file at the end of the manuscript with changes incorporated (i.e. not this document).
3. Data availability – a new section (2.3) has been added to describe accessing data.
4. Comments about OPCs and size distributions: We have discussed the details in the response to J. Reid. Since articles have already been published which dealt with the issues brought up by the reviewer, perhaps these were glossed over slightly in the present article for conciseness. We have added some extra detail to the article to account for this, in particular emphasizing that our processing accounts for the inherent uncertainties in the OPC instruments, and that two of our instruments (CIP15 and CIP100) use a different (light shadowing, rather than light scattering) technique, which is not susceptible to these sensitivities, yet supports the same conclusions.
5. LIDAR: we have better explained the differences between the two LIDARs used on the different aircraft in Fennec, in the instrumental section in the article, and also in the relevant results sections.

26 We hope that these changes sufficiently address the points brought up by the reviewers. The article
27 with tracked changes can be found below, with figures at the end of the document.

28 Yours sincerely,

29 Claire Ryder

30

31 **Advances in understanding mineral dust and boundary layer**
32 **processes over the Sahara from Fennec aircraft observations**

33 C.L. Ryder^{*1}, J.B. McQuaid^{2,6}, C. Flamant³, R. Washington⁴, H.E. Brindley⁵, E.J. Highwood¹, J.H.
34 Marsham^{2,6}, D.J. Parker², M.C. Todd⁷, J.R. Banks⁵, J.K. Brooke^{2,8}, S. Engelstaedter⁴, V. Estelles^{9,a}, P.
35 Formenti¹⁰, L. Garcia-Carreras^{2,a}, C. Kocha³, F. Marenco⁸, P. Rosenberg², H. Sodemann^{11, b}, C.J.T.
36 Allen⁴, A. Bourdon¹², M. Bart^{2,c}, C. Cavazos-Guerra^{7,d}, S. Chevaillier¹⁰, J. Crosier¹³, E. Darbyshire^{1,13}, A.R.
37 Dean¹⁴, J.R. Dorsey¹³, J. Kent⁸, D. O'Sullivan⁸, K. Schepanski^{2,e}, K. SzpekJ. Trembath¹⁴, A. Woolley¹⁴.

Commented [CLR1]: One of the authors was omitted by mistake in the ACPD submission.

38 [1] Department of Meteorology, University of Reading, RG6 6BB, UK

39 [2] School of Earth and Environment, University of Leeds, LS2 9JT, UK

40 [3] Sorbonne Universités, UPMC, Université Paris 06, CNRS & UVSQ, UMR 8190 LATMOS,
41 Laboratoire Atmosphères Milieux, Observations Spatiales, CNRS Sorbonne Universités UVSQ, Paris,
42 France

43 [4] School of Geography and the Environment, University of Oxford, UK

44 [5] Space and Atmospheric Physics, Department of Physics, Imperial College London, UK

45 [6] National Centre for Atmospheric Science, University of Leeds, LS2 9JT, UK

46 [7] Department of Geography, University of Sussex, Brighton BN1 9QJ, UK

47 [8] Met Office, Exeter, EX1 3PB, UK

48 [9] Dept. Física Fundamental y Experimental, Electrónica y Sistemas, Universidad de La Laguna,
49 Spain

50 [10] LISA, UMR CNRS 7583, Université Paris Est Créteil et Université Paris Diderot, Institut Pierre
51 Simon Laplace, Créteil, France

52 [11] Institute for Atmospheric and Climate Science, ETH Zürich, Switzerland

53 [12] SAFIRE, UMS CNRS-CNES-Météo-France, Francazal, France

54 [13] National Centre for Atmospheric Science, University of Manchester, Manchester, M13 9PL, UK

55 [14] Facility for Airborne Atmospheric Measurements, Cranfield, MK43 0AL, UK

56 [a] Now at Department of Earth Physics and Thermodynamics, Universitat de València, Spain

57 [a] Now at Department of Meteorology and the Bert Bolin Centre for Climate Research, Stockholm
58 University, Sweden

59 [b] Now at Geophysical Institute, University of Bergen, Norway

60 [c] Now at Aeroqual Ltd, 109 Valley Road, Auckland, New Zealand

61 [d] Now at Institute for Advanced Sustainability Studies (IASS), Berliner Straße 130, 14467
62 Potsdam, Germany

63 [ee] Now at Leibniz Institute for Tropospheric - Research - Permoserstr. 15 - 04318 Leipzig,
64 Germany

65 [*] Correspond to c.l.ryder@reading.ac.uk

66 **Abstract**

67 The Fennec climate program aims to improve understanding of the Saharan climate system through
68 a synergy of observations and modelling. We present a description of the Fennec airborne
69 observations during 2011 and 2012 over the remote Sahara (Mauritania and Mali) and the advances
70 in the understanding of mineral dust and boundary layer processes they have provided. Aircraft
71 instrumentation aboard the UK FAAM BAe146 and French SAFIRE Falcon 20 is described, with
72 specific focus on instrumentation specially developed and relevant to Saharan meteorology and
73 dust. Flight locations, aims and associated meteorology are described. Examples and applications of
74 aircraft measurements from the Fennec flights are presented, highlighting new scientific results
75 delivered using a synergy of different instruments and aircraft. These include: (1) the first airborne
76 measurement of dust particles sized up to 300 microns and associated dust fluxes in the Saharan
77 atmospheric boundary layer (SABL), (2) dust uplift from the breakdown of the nocturnal low-level jet
78 before becoming visible in SEVIRI satellite imagery, (3) vertical profiles of the unique vertical
79 structure of turbulent fluxes in the SABL, (4) in-situ observations of processes in SABL clouds showing
80 dust acting as CCN and IN at -15°C, (5) dual-aircraft observations of the SABL dynamics,
81 thermodynamics and composition in the Saharan heat low region (SHL), (6) airborne observations of
82 a dust storm associated with a cold-pool (haboob) issued from deep convection over the Atlas, (7)
83 the first airborne chemical composition measurements of dust in the SHL region with differing
84 composition, sources (determined using Lagrangian backward trajectory calculations) and
85 absorption properties between 2011 and 2012, (8) coincident ozone and dust surface area
86 measurements suggest coarser particles provide a route for ozone depletion, (9) discrepancies
87 between airborne coarse mode size distributions and AERONET sunphotometer retrievals under light
88 dust loadings. These results provide insights into boundary layer and dust processes in the SHL
89 region – a region of substantial global climatic importance.

90 **1 Background and Motivation**

91 The Sahara desert remains one of the most data sparse regions on the planet. During the northern
92 summer a vast low pressure system, the Saharan Heat Low (SHL), exists over the central Sahara
93 caused by the strong solar heating and this drives major dynamical features (e.g. Lavaysse et al.
94 (2009); Chauvin et al. (2010)). Strong sensible surface fluxes generate near-surface temperatures in
95 excess of 40 °C and a deep Saharan Atmospheric Boundary Layer (SABL) that reaches to a height of
96 6000 m, generating what is commonly regarded to be the world's deepest boundary layer (Tompkins
97 et al., 2005; Cuesta et al., 2009) . To the south of the Sahara lies the Sahel and the SHL exerts a
98 significant influence upon this region, in particular the timing of the West African Monsoon (WAM)
99 onset (Lavaysse et al., 2009; Sultan and Janicot, 2003). The prediction of the onset of the WAM has
100 been the topic of a number of recent science programmes, (e.g. the African Monsoon
101 Multidisciplinary Analysis (AMMA), Redelsperger et al. (2006)), as it is critical to the livelihoods of
102 the population in this region: the growing season here is short and the ground must be prepared and
103 planted ahead of the rains arriving.

104 The Sahara is the largest source of mineral dust on the planet, with the highest summer dust
105 loadings co-located with the SHL (Engelstaedter et al., 2006). Mineral dust is an important
106 atmospheric aerosol because of its direct and indirect radiative effects (Forster et al., 2007), its
107 contribution to atmospheric chemistry (de Reus et al., 2005), and its transport and deposition of

108 essential nutrients to the ocean (Jickells et al., 2005). Saharan dust is known to modify hurricane
109 activity by reducing local sea surface temperatures in the Caribbean (Dunion and Velden (2004); Sun
110 et al. (2009); Jenkins et al. (2008)) and in the tropical Atlantic Ocean (Evan et al., 2011; Evan et al.,
111 2009). Saharan dynamics, including haboobs frequently driven by moist convection (Marsham et al.,
112 2013c), low level jets (Washington et al., 2006) and dust devils and convective plumes (Ansmann et
113 al., 2009) ~~mean result that in~~ vast quantities of dust ~~are being~~ lofted on a very regular basis into the
114 atmosphere where they are then susceptible to synoptic-scale atmospheric transport. Thus the
115 Saharan region plays a significant role in the weather and climate in the northern hemisphere
116 (Tompkins et al., 2005; Rodwell and Jung, 2008), influencing regions far beyond its geographical
117 boundaries.

118 There are considerable uncertainties in both climate and numerical weather prediction models for
119 this region (Evan et al., 2014; Marsham et al., 2008b; Messager et al., 2010). Representation of the
120 position and intensity of the SHL in climate models varies considerably. Identifying the cause of such
121 discrepancies and ascertaining which representation most closely matches reality can only be
122 addressed through observational data. The extreme nature of the Saharan climate and also the
123 considerable uncertainties associated with mineral dust aerosols in numerical models all compound
124 the discrepancies between models and reality (e.g. Kim et al. (2014); Huneeus et al. (2011); Evan et
125 al. (2014)). Additionally observations of both dust chemical composition and the full size distribution
126 in this remote region are crucial for accurately representing the radiative effect of dust (Formenti et
127 al. (2014); Mahowald et al. (2014)).

128 In the last decade or so, a number of field programmes have been tasked with improving the
129 observational dataset on meteorological and aerosol conditions in the wider North African sector
130 (Table 1 and Figure 1). With the exception of limited measurements during AMMA (Messager et al.,
131 2010; Cuesta et al., 2008), no previous campaign has focused on this central region of North Africa
132 during the summer dust season. For example, SAMUM1 was based in Morocco, while SAMUM2
133 observations took place at the Cape Verde Islands (Heintzenberg, 2009; Ansmann et al., 2011).
134 Fennec was conceived and designed to fill critical gaps in observations and understanding of the
135 Saharan climate system.

136 The Fennec climate programme aims to improve understanding of and quantify the physical
137 processes controlling the Saharan climate system, through a synergy of observational and modelling
138 approaches in order to evaluate and attribute errors in weather and climate models for this region
139 (Washington et al., 2012). The observational strategy is a large scale, multi-platform approach
140 involving ground-based measurements, airborne observations and Earth observation. Fennec is an
141 international consortium which includes research groups from the United Kingdom, France,
142 Germany, Switzerland and the United States of America working in collaboration with the
143 Meteorological Services of Algeria and Mauritania in North Africa.

144 This paper will focus on the airborne operations that were deployed as part of the Fennec
145 programme and key scientific findings stemming from the airborne programme ~~as well as those that~~
146 ~~were possible through a combination of airborne observations with ground-based, satellite and/or~~
147 ~~modelling approaches. Observations by means of an airborne platform provide an invaluable~~
148 ~~approach, including access to remote, inhostpitable regions of the Sahara, tracking of non-static~~
149 ~~atmospheric features and providing vertical profile observations as well as dust observations above~~

150 the surface layer, which is vital to understanding the capacity for long-range transport of uplifted
151 dust. Many of the fixed target features in the Saharan region, such as albedo gradients and
152 orographic features (e.g. Figure 1), are very remote, whilst the dynamical features, by definition, are
153 non static in nature. Therefore observation by means of an airborne platform provides an invaluable
154 approach. This paper will outline the specific benefits of using an aircraft as an observational
155 platform for tackling the challenges of such a dynamically variable region. Measurements on aircraft
156 platforms provide the ability to link together spatial and temporal features which are simply not
157 accessible through fixed ground sites or satellites or even a combination of both. An aircraft can
158 focus upon specific features such as moving weather systems across a large geographic area and
159 move with it, allowing temporal evolution to be observed. The ability to make vertical profiles
160 through the atmosphere, thus providing vertically resolved measurements, is invaluable with
161 features such as the remote SABL. Airborne platforms can be positioned themselves at appropriate
162 altitudes for dedicated remote sensing surveys such as above/below radiatively active layers of
163 aerosol such as mineral dust. Contrary to ground based measurements, it is possible to make in situ
164 aerosol measurements above the surface dust layer which is vital to understanding the capacity for
165 long range transport of uplifted dust, which is of particular interest in this region. Measurements on
166 aircraft platforms provide the ability to link together spatial and temporal features which are simply
167 not accessible through fixed ground sites or satellites or even a combination of both. Furthermore,
168 specifically in the June 2011 Intensive Observation Period (IOP) two aircraft were operated and their
169 combined power meant that specific events could be followed through staggered missions. This is
170 not achievable with a single aircraft for operational reasons such as aircrew duty periods. Finally, the
171 combination of ground, airborne and satellite observations provide the fullest picture possible of the
172 area of interest.

173 During 2011 and 2012 an extensive dataset was collected as part of the Fennec intensive
174 observation programme. These included the deployment of two airborne platforms: the UK BAe146
175 FAAM and French SAFIRE F-20 aircraft, and also ground based observations via two supersites
176 located on the western and eastern flanks of the central Sahara: Zouerate, Mauritania (Todd et al.,
177 2013) and Bordj Badji Mokhtar, Algeria (Allen et al., 2013; Marsham et al., 2013b). These were
178 supplemented by a network of automated weather stations which were installed in the remote
179 desert (Hobby et al., 2013). An overview of the aircraft deployments are provided in Table 2; more
180 detailed flight information is presented later. As part of the outreach activities of the Fennec project
181 a movie, 'Into the Cauldron: A Meteorological Adventure,' has been also produced (Sternberg,
182 2013).

183 In addition to the Fennec programme, a number of supplementary projects took advantage of the
184 aircraft deployment to the region. The Lagrangian Dust Source Inversion Experiment (LADUNEX)
185 (Sodemann et al., submitted manuscript) used the in-situ and remote sensing observations of
186 mineral dust in order to validate a Lagrangian particle dispersion model FLEXPART and improve its
187 ability to represent dust transport in the atmosphere. RAIN4DUST project exploited the remote
188 sensing data from the French Falcon aircraft to investigate dust sources in relation to sediment
189 supply and surface characteristics in the foothills of the central Saharan mountain ranges
190 (Schepanski et al., 2013). Finally, the Sunphotometer Airborne Validation EXperiment (SAVEX) was
191 designed to take advantage of the use of the island of Fuerteventura as operating base from which
192 to conduct an intercomparison of a number of sunphotometers installed on Tenerife with aircraft
193 observations.

194 The aims of this paper are firstly to document and describe the flights and meteorology during the 3
195 Fennec IOPs in order to provide a reference and context for published and future articles. Secondly,
196 we provide new scientific results that have come about as a result of the Fennec airborne
197 programme, both through airborne observations in isolation over the remote Sahara, and through
198 the integration of data from different platforms – i.e. dual-aircraft observations and ground-based,
199 airborne and satellite platforms. Therefore this paper provides insights into Saharan processes which
200 separate papers cannot. Finally, despite many challenges, the Fennec aircraft campaigns have
201 collected the only comprehensive in-situ data from the Saharan region – a region of substantial
202 global climatic importance. Along with ground-based and satellite measurements, these data
203 provide a much-needed resource with which to develop the science linking dust, dynamics and
204 radiation in the central Sahara, and will be heavily exploited in the coming years. This paper provides
205 a detailed overview of the data and its context, as well as a survey of first results.

206 The paper is structured as follows: in Section 2 we describe the aircraft instrumentation, with a focus
207 on instrumentation specifically developed or installed for Fennec, and also provide information on
208 data provision for the scientific community. Section 3 describes the meteorology during Fennec and
209 provides an overview of the flights performed. Section 4 provides a description of new scientific
210 results, Section 5 concludes the article.

211 **2 Aircraft Instrumentation**

212 Here we describe the instrumentation on both aircraft, the BAe146 and the Falcon F-20, with
213 particular emphasis regarding instrumentation particularly relevant to Fennec measurements.
214 Throughout this article we refer to particle size in diameter.

215 **2.1 FAAM BAe146 Aircraft**

216 The UK's BAe-146-301 Large Atmospheric Research Aircraft operated by the Facility for Airborne
217 Atmospheric Measurements (FAAM) (henceforth the BAe146 aircraft) is available to the ~~UK~~-science
218 community in a number of different configurations. These allow the most efficient use of space and
219 access to inlets (which tend to be in the forward section of the cabin) as well as minimizing the
220 aircraft payload, which in turn maximizes the sortie duration. Due to the remoteness of the areas of
221 interest for Fennec the instrument fit was customized to provide the best balance of observational
222 rigour and range. Table 4 details the instrument fit for the Fennec IOPs; ~~{some instruments were~~
223 ~~only available for some of the deployments, these are indicated in the table}~~. There are a number of
224 excellent descriptions of the standard instrumentation from previous campaigns which have utilised
225 the BAe146 aircraft (e.g. (Renfrew et al. (2008); Highwood et al., 2012; McConnell et al., 2008;
226 Haywood et al., 2011a)); other specific instrumental references are provided in Table 4.
227 Instrumentation specifically developed, installed or configured for Fennec are described in more
228 detail below.

229 **2.1.1 LIDAR**

230 The BAe146 aircraft operates a commercial Leosphere ALS450 aerosol-backscatter LIDAR suitable for
231 aerosol and thin cloud observation (Marenco et al., 2011) ~~suitable for aerosol and thin cloud~~
232 ~~observation~~. A description of the lidar-LIDAR system is provided by Chazette et al. (2012) and
233 technical information is available in Table 1 of Marenco et al. (2014). The nadir-viewing LIDAR
234 provides elastic backscatter at 355 nm and features an uncalibrated qualitative depolarisation

235 channel, fused qualitatively to distinguish depolarising layers). Data are recorded at a vertical
236 resolution of 1.5 m and an integration time of 2 s, giving approximately 200 m horizontal resolution
237 at aircraft speeds. The instrument is lightweight, has a relatively small receiver aperture of 15 cm
238 diameter, has a 12 mJ pulse energy (20 Hz PRF) output and requires a low level of maintenance
239 which makes it ideal for frequent operation aboard the BAe146 aircraft. However, as a consequence
240 the signal to noise ratio is poorer compared to the Falcon LNG LIDAR.

241 Initial quick-look data is provided as range-square corrected signal (arbitrary units) which is
242 proportional to the total backscatter coefficient (from molecules and particles) at a given range, r ,
243 times the two-way transmission of light- from the laser source to the range r (i.e. a function of the
244 atmospheric optical depth), for example as shown in Figure 7 and Figure 17, for which no attempt
245 has been made to correct for attenuation by the aerosol layers). In these cases we use the
246 Leosphere LIDAR data to locate dust layers and clouds, and for which the range-corrected
247 backscattered signal is sufficient, although dust layers lower in the atmosphere may not always be
248 evident with such a representation, due to attenuation at higher altitudes.

249 In a further step, Aaerosol extinction coefficient can be computed from the LIDAR range-square
250 corrected backscatter signal using the method described by Marenco et al. (2013), although this is
251 labour-intensive, since the method is not automated and it requires a profile-by-profile review of
252 assumptions. However Additionally, the signal-to-noise ratio for the dust laden atmosphere in the
253 Fennec region often causes difficulties in inverting the LIDAR backscatter signal to extinction
254 coefficients. This can be overcome by integrating the lidar signals, i.e., For example Sodemann et al.
255 (submitted manuscript) decrease resolution to 300 m in the vertical, and a 60 s integration time,
256 translating to extinction coefficient profiles provided at a ~9 km along-track footprint at a typical
257 ground speed of $\sim 150 \text{ ms}^{-1}$. In the lowest 0-2 km layer the uncertainty in the extinction coefficient is
258 of the order of 100%, but this uncertainty quickly decreases above, the extent of which is dependent
259 on the ambient aerosol conditions (e.g. Marenco et al. (2014)).

260 2.1.2 Low Turbulence Inlet (LTI)

261 A very important consideration when observing aerosol particles is the efficiency of the transmission
262 system which passes external aerosol into the aircraft cabin for collection or in situ analysis. It is
263 highlighted in the difficulty in making accurate and reliable measurements from an aircraft platform,
264 particularly that of coarse mode aerosol (Wendisch et al., 2004). For objectives such as those of the
265 Fennec program, this is of particular importance since a significant fraction of mineral dust is in the
266 coarse mode (Weinzierl et al., 2009). Inlet design can modify aerosol size distribution through either
267 underestimation due to aerosol losses or overestimation due to enhancements.

268 The BAe146 has a specialised Low Turbulence Inlet (LTI) which is designed to provide a characterised
269 community inlet capable of delivering supermicron aerosol into the cabin. This is achieved by
270 reducing turbulent flow within the tip of the inlet, reducing impaction of particles to the walls of the
271 inlet (Wilson et al., 2004). The LTI further maintains isokinetic sampling flow using a feedback
272 controlled pumping system.

273 A Grimm Technik Optical Particle Counter (OPC) was mounted inside the aircraft cabin behind the LTI
274 (LTI-GRIMM), and showed that size distributions behind the LTI compare well with those from the
275 externally mounted aircraft probes. In order to further evaluate inlet efficiency on the BAe146,
276 Grimm OPCs were mounted behind various Rosemount inlets. This allowed evaluation of the size

277 distributions passed by the standard BAe146 Rosemount inlets for the first time, from which many
278 of the internally installed aerosol instruments draw their sample from, such as the nephelometer,
279 particle soot absorption photometer, and aerosol mass spectrometer (Trembath, 2012; Trembath et
280 al., 2012). Significant losses and enhancements of the size distribution have been found to occur at
281 different size ranges.

282 **2.1.3 Double Nephelometer Setup**

283 During Fennec, two TSI 3563 integrating nephelometers measuring scattering at 450, 550 and 700
284 nm were operated inside the aircraft cabin behind a Rosemount Inlet. During Fennec 2011, the
285 nephelometers were run in series with a BGI Very Sharp Cut Cyclone Impactor between them. The
286 impactor has a 50% penetration efficiency at 2.5 μm aerodynamic diameter, or around 1.5 μm
287 geometric diameter, at a flow rate of 16.67 litres per minute (LPM). This therefore allows the 'first'
288 nephelometer to measure scattering due to all particles passing the Rosemount inlet and the
289 pipework (estimated to be particles smaller than 2.5 microns, Trembath (2012)), and the 'second'
290 nephelometer to measure scattering from the fraction of particles smaller than 1.5 microns.
291 However, due to the nephelometers being in series, it was difficult to account for the loss of
292 particles between the two instruments. Therefore during Fennec 2012 the two nephelometers were
293 operated in parallel to avoid this problem. This was possible due to ~~the addition of a newer~~, more
294 powerful pump being used, capable of 50 LPM, even up to altitudes of up to 9000 m. Secondly a
295 volume flow controller was installed to replace the mass flow meter and needle valve.

296 The synergy in the approach of operating a Grimm OPC behind a Rosemount inlet to measure the
297 size distribution, and the use of the impactor to separate the sub-1.5 micron scattering from that
298 measured as standard by the nephelometer is novel, and allows any bias in scattering and
299 absorption due to Rosemount inlet and pipework effects on the BAe146 to be assessed for the first
300 time, which can lead to significant underestimation of dust absorption properties when not
301 accounted for (Ryder et al., 2013b).

302 **2.1.4 Size Distribution Measurements**

303 The BAe146 is well equipped to measure aerosol size distributions (for example, see Haywood et al.
304 (2008); Johnson et al. (2012)). However, the Fennec campaign was unusual amongst aerosol
305 campaigns in the large number of instruments operated to measure particles larger than 3 μm
306 diameter, and in the measurement of 'giant mode' particles – those sized over 30-40 μm .
307 Interestingly, the recent eruption of Eyjafjallajökull in Iceland has reinvigorated the interest in 1-10
308 μm particles since volcanic ash is generally in the same size region as mineral dust and they both
309 have similar challenges such as non-spherical morphology (Ansmann et al, 2012): ~~clearly hence~~ there
310 is considerable benefit to be gained from the concerted efforts surrounding the observation of
311 volcanic ash.

312 Instruments measuring size distribution, and the size ranges measured are shown in Table 4, and
313 also in detail by Ryder et al. (2013b). During Fennec 2011, a total of 6 different instruments
314 successfully measured size distributions between sizes of 0.15 to 300 microns diameter - namely the
315 PCASP (accumulation mode), CDP, LTI-GRIMM, SID2H and CAS (coarse mode), and finally the
316 University of Manchester CIP15 in the giant mode (see Table 4 for explanation of acronyms). All of
317 these are wing mounted except the LTI-GRIMM. Of particular note ~~are is~~ the use of the CIP15 for
318 particles sized 15 microns and above, using image shadowing techniques. This alleviates the need to

319 correct particle size for refractive index, as is required for optical particle probes. Although the CIP15
320 is capable of measuring particles sized up to 930 μm , electrical noise allowed measurements up to
321 300 μm . Both Rosenberg et al. (2012) and Ryder et al. (2013b) show that the CIP15 and CDP/SID2H
322 size distributions agree well in the overlap zone, suggesting accurate measurements of size
323 distributions. Additionally, the PCASP and CDP agree well at their overlap zones. (see Ryder et al.
324 (2013b) and Rosenberg et al. (2012) for full details). Finally, we highlight the regular calibration of
325 the CDP probe during the campaign, which results in better characterised size distributions (see
326 Rosenberg et al., 2012). The combination of these procedures and operation of the various
327 instruments gives good confidence in the measured size distributions, particularly when significant
328 numbers of coarse particles are present (e.g. see Section 4.1.1).

329 During Fennec 2012 a slightly different suite of instruments was operated due to logistical
330 requirements, comprising a PCASP, CDP, 2DC, SID2H, FAAM CIP15 and FAAM CIP100. Unfortunately
331 the CIP15 suffered from electrical noise during the 2012 IOP and the data was not usable. However,
332 the operation of other instruments such as the CDP and 2DC provide alternative measurements for
333 this size range. Additionally the operation of the CIP100 probe extends the measurement range up
334 to 6200 μm .

335 In order to deal with several sources of uncertainty regarding OPC measurements, the size
336 distributions were carefully processed as described in detail by Rosenberg et al. (2012). Briefly, each
337 OPC is considered as an instrument which directly measures particle scattering cross section and is
338 calibrated in terms of this variable. Using the uncertainty in this calibration and Mie theory with an
339 appropriate refractive index for the measured aerosol, we derive a probability density function
340 which gives the probability of a particle of a particular size being counted in a particular OPC bin.
341 Integrating this probability density function allows us to derive the mean diameter and effective
342 width of each bin. This method also permits full uncertainty propagation including ambiguities
343 caused by the nonlinear and non-monotonic Mie theory. Refractive indices spanning 1.53–0.001i to
344 1.53–0.003i have been assumed and errors in diameter and number concentration due to this
345 uncertainty have been propagated. The size distributions have been produced assuming spherical
346 particles rather than non-spherical particles, which has been shown by others to have a negligible
347 impact on the resulting size distributions (Osborne et al., 2011; Veihelmann et al., 2006; Lacis and
348 Mishchenko, 1995; Liu et al., 1992).

349 **2.1.5 Spectrally Resolved Radiation Measurements**

350 In addition to the core pyranometers on the upper and lower of the aircraft fuselage measuring
351 downwelling and upwelling shortwave irradiance respectively, a number of ~~non-core~~^{non-core} specialist
352 radiometers were operated during Fennec which will allow considerably more detailed radiative
353 measurements and radiative closure to be performed. In the shortwave spectrum, the Spectral
354 Hemispheric Irradiance MeasurementS (SHIMS) measured spectrally resolved up and downwelling
355 irradiance from 0.3 to 1.7 μm . The Shortwave Spectrometer (SWS) measures spectrally resolved
356 radiances from 0.3 to 1.7 μm , using an externally mounted scanning telescope designed for viewing
357 at particular angles. In the longwave spectrum, the Airborne Research Interferometer Evaluation
358 System (ARIES) measured spectrally resolved radiances from 3.3 to 18 μm , at either nadir or zenith,
359 as well as several different downward-pointing angles. Further details of SHIMS, SWS and ARIES can
360 be found in Osborne et al. (2011). Operation of these instruments allows detailed radiative closure
361 to be performed (e.g. (Haywood et al. (2011b); Osborne et al., 2011)). Further work will examine the

362 radiative measurements made under extremely high dust loadings when very large particles were
363 present.

364 **2.1.6 Turbulence probe**

365 Due to the scientific objectives of the Fennec program, the ability of the aircraft to make robust
366 observations of atmospheric turbulence was of paramount importance. Three dimensional wind
367 vectors are generated using a 5 port radome mounted turbulence probe at the aircraft nose which
368 provides angle of attack measurements. These are combined with pitot tube measurements of air
369 speed and position information from a GPS inertial navigation unit to generate ground referenced
370 wind vectors at 32 Hz (Petersen & Renfrew 2009). A known linear dependence between the vertical
371 component and aircraft pitch results in additional post-processing. This is likely the result of
372 uncertainties in the calibration of the turbulence or pitot probes. Some of the parameters (static
373 pressure and airspeed required for the processing) are generated through the on-board aircraft
374 computer, this is calibrated in-situ annually as part of the maintenance schedule, using a pressure
375 calibrator. Airspeed is calibrated similarly. The radome transducers are calibrated at a calibration
376 laboratory annually, or as determined by inspection of the data for drifts or other artefacts. The INU
377 alignment is assessed annually by a physical survey for pitch, roll, and heading. Angle of attack
378 (AOA) and angle of sideslip (AOSS) calibrations derive from AOA/AOSS flight manoeuvres that were
379 carried out when the facility was commissioned, as they are physically dependent on the radome
380 mounting. These have been subsequently validated to confirm this. The AOA/AOSS is further
381 corrected using yawing orbits, where further corrections are introduced to these quantities. True
382 airspeed is corrected using reverse-heading manoeuvres, where the correction minimises the
383 difference in derived wind measurement up/down wind.

384 **2.1.7 Cloud Condensation Nuclei Observations**

385 The concentration and properties of Cloud Condensation Nuclei (CCN) were measured using a
386 commercial dual column continuous flow streamwise thermal gradient instrument (Droplet
387 Measurement Technologies, Boulder, Co). The principles of its design are outlined in (Roberts and
388 Nenes (2005); Lance et al. (2006); Rose et al. (2008)). Ambient air is drawn into a pair of temperature
389 controlled columns where it encounters a particle free sheath flow which is humidified to near-
390 saturation. A thermal gradient exists down each of the columns, meaning that supersaturation
391 occurs as the samples flows through the columns. Activated aerosol will form droplets and increase
392 in size dependent upon their hygroscopicity. The instrument is configured to provide a pair of
393 supersaturations at any time and has supersaturation range nominally between 0.07 % and 2 %. The
394 residence time within the humidified zone is sufficient that these activated droplets grow to be
395 diameters larger than 1 μm , all particles with a diameter below this threshold are judged to be
396 unactivated interstitial particles. An optical particle counter at the base of each column estimates
397 the size distribution of the droplets (0.75 - 10 μm across 20 size bins).

398 In order to ensure stable volumetric flow to the CCN instrument, vital for robust measurements
399 across altitude ranges encountered by airborne platforms, it draws air from a reduced pressure
400 buffer volume which is connected to a modified Rosemount 102E inlet (Trembath, 2012). In addition
401 to the CCN, a condensation particle counter, CPC (modified 3786 UCPC, Quant Technologies) also
402 samples from this plenum to allow the total concentration of particles (2.5 nm – 3 μm) to be
403 determined.

404 **2.2 SAFIRE Falcon F-20 Aircraft**

405 The SAFIRE (Service des Avions Français Instrumentés pour la Recherche en Environnement) Falcon
406 20 (F20) performed research flights during the June 2011 IOP. In contrast to the BAe146, it was
407 equipped mostly with instrumentation designed to target the Saharan heat low region remotely
408 from high altitudes (see Table 3 detailing the F20 instrumentation).

409
410 The F20 was equipped with the backscatter LIDAR LEANDRE Nouvelle Génération (LNG, de Villiers et
411 al. (2010)), allowing the measurement of atmospheric reflectivity at three wavelengths (355, 532
412 and 1064 nm) to analyze the structure and radiative characteristics of desert dust plumes with a
413 vertical resolution of 15 m and a horizontal resolution of 2 km (corresponding to a temporal
414 averaging of the data of 10 s - or 200 shots - in order to reach a signal to noise ratio above 100). The
415 LIDAR also has a depolarization capability on the 355 nm channel. During Fennec, the profiles of
416 aerosol extinction coefficient at 532 nm are retrieved with an uncertainty on the order of 15% using
417 a standard lidar-LIDAR inversion technique which is described at length in Banks et al. (2013) and
418 Schepanski et al. (2013). The aerosol lidarLIDAR ratio used for the inversion is considered to be
419 constant with altitude and set to 47 sr. This value is intermediate between the value derived at 532
420 nm from space-borne, airborne, and ground-based LIDAR systems over northern Africa (i.e. 55 sr:
421 Heintzenberg (2009) and Schuster et al. (2012), 50-60 sr: Tesche et al. (2009); Gross et al. (2011)
422 and those derived over Sahelian Africa (i.e. 41 sr: Omar et al. (2009) and Schuster et al. (2012)).

423
424 In addition to the LIDAR, the Falcon 20 was also equipped with a Vaisala AVAPS dropsondes
425 launching system (a total of 136 sondes were launched from the Falcon aircraft during the 2011
426 deployment), radiometers (broadband up- and down-looking Kipp and Zonen pyranometers and
427 pyrgeometers), the radiometer CLIMAT (Legrand et al. (2000)) as well as in situ pressure,
428 temperature, humidity and wind sensors. There was also a nadir pointing visible camera (Basler SCA
429 1400-30FM with a 9 mm lens (Fujion, 2/3")) mounted aboard the Falcon providing high-resolution
430 aerial photographs of the surface (Schepanski et al., 2013).

431 **2.3 Access to Data**

432 UK-Fennec FAAM aircraft data from the BAe146 is available at the British Atmospheric Data Centre
433 (BADC, <http://badc.nerc.ac.uk/home/index.html>) and is freely available subject to registration.
434 Fennec-France aircraft data is available from the Sedoo (Service de données de l'OMP,
435 <http://catalogue.amma-international.org/> and is attached to the AMMA database, subject to free
436 registration, listed under "Fennec" in the project list.

437 **3 Flights and Meteorology**

438 We now provide an overview of the meteorology and dust events during the campaigns, and a
439 description of the flights performed in relation to these. A preliminary mission with the BAe146 was
440 carried out in April 2011, using Ouarzazate, Morocco as the aircraft base, with measurements taken
441 over Mauritania. However, flight restrictions from this base meant that it was logistically more
442 straightforward to operate from Fuerteventura, one of the Canary Islands, Spain, from where
443 subsequent campaigns in June 2011 (both aircraft) and June 2012 (BAe146 only) were based. From
444 Fuerteventura, research flights operated over Mauritania, Mali, Senegal and the Eastern Atlantic
445 Ocean. In following sections, flight numbers prefixed with 'b' refer to BAe146 flights, whereas flight
446 numbers starting with 'F' refer to Falcon flights.

447 **3.1 Meteorology**

448 Here, we consider the synoptic scale structure of the atmosphere in the North African sector during
449 the three Fennec observational phases shown in Table 2. We relate this in general terms to the
450 structure of the SABL and dust conditions observed in the Fennec flight domain of the western
451 Saharan region. In specific relation to the two summertime phases of June 2011 and 201102, we
452 consider the state of the dominant features of the summertime low-level circulation over western
453 North Africa, namely the Azores high pressure system, the SHL and the inter-tropical discontinuity
454 (ITD), as well as the upper level circulation in the adjacent mid-latitudes. The SHL has a pronounced
455 seasonal cycle (Lavaysse et al., 2009) involving a southeast to northwest migration from its position
456 to the south of the Hoggar mountains ($\sim 18^\circ\text{N}, 5^\circ\text{E}$) in May to its most northerly position close to
457 24°N and 0°W during July and August. The climatological mean date of transition between these two
458 states is 20th June.

459 **3.1.1 Fennec Pilot Campaign 2011**

460 The synoptic situation during the short Fennec pilot campaign during April 5th-8th 2011 generated
461 numerous dust emission events characteristic of spring time dust events over the Sahara. On the 1st
462 and 2nd April a high pressure ridge over Algeria-Libya sector drove a strong northeasterly Harmattan
463 surge over the central-eastern Sahara activating multiple dust sources in Algeria, Libya, Niger and
464 Chad created a large dust plume of advected dust southwestward over Northern Mali, Southern
465 Algeria by the 3rd. Further westward transport of this plume into the Fennec aircraft operations
466 zone was prevented by strong northeasterly circulation around an intense cut-off low on the 3rd-4th
467 April (Feature A in Figure 3a). This low tracked northwards from western Algeria to Morocco over
468 this period and was accompanied by strong cyclonic near surface winds with pronounced dust
469 emission along primary and secondary cold fronts penetrating southeastward over southern
470 Morocco and northern Mauritania on the 4th April. Fennec flight b589 was able to observe this dust
471 feature and the accompanying cold surge. Subsequent flights on the 5th-8th (see Table 5) observed
472 the interaction of the cold maritime intrusion with dusty Saharan air, after which the dust was
473 transported towards Portugal (Preissler et al., 2011).

474 **3.1.2 Fennec IOP 2011**

475 During this IOP most of the F20 and BAe146 flights were conducted over northern Mauritania and
476 northern Mali. In terms of the large-scale structure of the atmosphere during June 2011 in this
477 region, a clear distinction can be made between a 'maritime phase' from around the 2nd-12th June
478 and a 'heat-low phase' from around the 13th-30th June (see Todd et al., 2013 for full details). These
479 phases essentially determine conditions across the entire central-western Sahara. These maritime
480 and heat-low phases are broadly congruent with the 'east' and 'west' and phases, respectively, of
481 the intraseasonal SHL mode of variability described by Chauvin et al. (2010). During the maritime
482 ('heat low east') phase the upper level pattern exhibited a trough centered over the Iberian
483 peninsula extending southwards over the northern extremity of North Africa (Feature A in Figure
484 3b). In addition, at low levels the SHL remained relatively stationary in an anomalously eastward
485 location centred $\sim 15^\circ\text{E}$ (Feature B in Figure 3b), similar to the mean state for May, and the Azores
486 high ridged towards the coast of northwest Africa. These conditions combined to drive anomalous
487 westerlies throughout the troposphere over northwest Africa creating a strong northwesterly inflow
488 of maritime air over much of the Fennec flight domain (Feature C in Figure 3b) with the ITD displaced
489 southward (not shown). As such, the Sahara is effectively 'ventilated' by cool advection from the
490 Atlantic sector restricting the heat low to the central/eastern Sahara. Accordingly, Fennec

491 observations at both supersites (not shown) indicate that the SABL during the maritime phase to be
492 anomalously cool and dry with shallow daytime convective boundary layer development (Marsham
493 et al., 2013b; Todd et al., 2013) and generally cloud free conditions. Aerosol loading was low due to
494 the relative absence over the Fennec flight domain of the two dominant dust generating processes,
495 namely ~~the~~ cold pools from moist convective systems, favoured within the southerly monsoon flow
496 (ITD 'bulge') on the eastern flank of the SHL, and ~~the~~ enhanced northeasterly Harmattan winds
497 around the western flank of SHL trough. As a consequence, these two ~~such~~ dust generating activity
498 waswere largely restricted to the central Sahara with the eastward-displaced SHL.

499 Subsequently, during the latter Heat Low (west) phase anomalous positive geopotential heights
500 dominated over Iberia and the extremity of northwest Africa (Feature A in Figure 3c), associated
501 with the passage of three upper level ridges. At lower levels, the SHL exhibited an abrupt westward
502 displacement to ~5-10°W (Feature B in Figure 3c) in two distinct intraseasonal pulses. These
503 conditions combined to drive anomalous mid and upper level easterly flow, with easterlies at lower
504 levels around the SHL, evident over the western Saharan sector (Feature C in Figure 3c) and Fennec
505 flight domain. Fennec ground-based observations indicate the SABL during the Heat Low phase of
506 June 2011 to be substantially hotter with deeper afternoon Convective Boundary Layer (CBL)
507 development and cases of almost 'pure' well-mixed near dry-adiabatic profiles from the surface to
508 the top of the Saharan Residual Layer (SRL) at ~5 km height. Dust aerosol loadings are substantially
509 higher over the western Sahara and Fennec flight domain during the heat low phase associated with
510 enhanced meso-scale convective activity and strong easterlies around the heat low and African
511 Easterly wave troughs. Shallow convective cloud was common developing in the later afternoon in
512 the relatively moist upper SRL.

513 Flight planning to meet Fennec science objectives was largely determined by synoptic meteorology,
514 as well as logistical constraints. As such, the science objectives of specific flights (Table 5, Table 6 and
515 Table 7) are geared to the prevailing meteorology described above. Overall, flights during the
516 maritime phase (Falcon only) were able to sample substantial dust emission events over northern
517 Mauritania (F13, F18). During the heat low phase certain flights were able to measure dust-
518 meteorological processes associated with both northeasterly low level jet -related emission (e.g.
519 b600/601/602, b610, b614) and mesoscale convective system (MCS) cold pool events originating
520 over central Mali (b604) and also the Atlas mountains to the north (b605 and F22/F23). Flights to
521 survey the SABL were able to measure the pronounced evolution in the structure of the PBL over
522 this transition from maritime (e.g. F14-F17) to heat low phases (e.g. b607/b608, F24/F25),
523 representing the intra-seasonal variability and seasonal evolution of the Saharan atmosphere.

524 **3.1.3 Fennec IOP 2012**

525 Unlike the equivalent period of June 2011 Fennec IOP 2012 period 1-17th June there was no clear
526 projection of the circulation onto the east-west heat low mode of Chauvin et al. (2010). As such, the
527 period was characterised by a relatively stationary SHL centred close to the triple point of the
528 Algeria-Niger-Mali border, further west than during the first half of June 2011. However, relatively
529 subtle synoptic-scale variations strongly influenced the circulation over the western Saharan sector
530 and the Fennec flight domain. First, during the early part of June 2012 from 1st-9th a weak upper
531 level trough extended south towards the coast of Morocco (Feature A, Figure 3d) and a heat low
532 extension was established over far western Algeria (Feature B, Figure 3d) driving a strong
533 northwesterly maritime flow over the Fennec domain (Feature C, Figure 3d). As with the maritime

534 phase of IOP 2011 this led to the characteristic maritime conditions of a cool, dry SABL with shallow
535 CBL daytime development and relatively cloud- and aerosol-free conditions over almost all the
536 domain. This maritime flow weakened after the 10th June and a heat low extension west into
537 northwest Mali from 14th-17th June (not shown) established more characteristic heat low SABL
538 conditions over the eastern Fennec flight domain.⁵³⁸ Specifically, a strong northeasterly low level flow
539 around the western flank of the HL trough favourable to dust emission and a northern extension of
540 monsoon flow to the east over Mali developed. MCS activity increased as the maritime flow
541 weakened after the 8th June and substantial cold pool events were observed in the monsoon flow
542 over Southern Mauritania on the 8th (see ITD 'bulge' Feature D in Figure 3d) and over Southern Mali
543 on the 12th and 14th (not shown).

544 Fennec 2012 flights targeted specific features of the evolving Saharan atmosphere, including surveys
545 of the maritime flow in the early period (b699/700), aged dust from MCS cold pools to the south of
546 the flight domain sampled over the ocean (b702-3) and southern Mauritania (b704), boundary layer
547 heat fluxes close to edge of the SHL (b705), the SHL tongue and LLJ dust emission (b706-8) and dust
548 uplift and radiative processes (b708-9).

549 **3.2 Description of Flights**

550 Table 5 to Table 7 list each flight conducted during the various Fennec phases. A brief description of
551 each flight is provided here to link the meteorology described in Section 3.1 to each flight's scientific
552 aims, and to provide information for future reference. Some flights and key scientific results are
553 described further in Section 4.

554 **3.2.1 Flights during the Pilot Campaign 2011**

555 During the Fennec Pilot campaign in April 2011, 7 flights were performed (Table 5, Figure 2a). b589
556 was an initial shake-down flight to test operational logistics, and was conducted at high altitude only,
557 but overflew a dust front which was observed with the LIDAR and dropsondes. b590 (morning) and
558 b591 (afternoon) were the first flights performing in-situ measurements, and sampled maritime
559 inflow over Mauritania, which was overlain by dust layers at higher altitudes. b592 took place 2 days
560 later on 7 April (note b592 was actually two separate flights, one in the morning and one in the
561 afternoon) and sampled the diurnal evolution of the recovering SABL (Saharan boundary layer)
562 following the retreat of marine air. b593 continued the sampling of the recovering SABL, but over a
563 different surface albedo. b594 was a science transit return of the BAe146 to the UK, sampling dust
564 transported northwards by a low pressure system over Morocco.

565 **3.2.2 Flights during Fennec IOP 2011**

566 June 2011 was the main flying period of Fennec, when both the Falcon and the BAe146 were
567 conducting missions over the Sahara. Eleven flights were performed with the F20 during the period
568 2-16 June (Figure 2b, Table 6). The first four flights (F09 – F12) were designed to sample the dust
569 outflow from the continent, over the coastal Atlantic, though almost no dust was sampled during
570 F10. The subsequent seven flights were conducted over the continent, with two flights (F13 and F18)
571 dedicated to the study of the morning dust uplift over alluvial sources of Northern Mauritania in
572 connection with the decay of the low-level jet. The flights were part of the RAIN4DUST project
573 funded by the EUropean Facility for Airborne Research, EUFAR (Schepanski et al., 2013), designed to
574 examine alluvial deposits as a dust source. Four flights were conducted along the exact same track
575 (F14, F16, F17 and F19) to document evolution of the thermodynamics, the dynamics and the

576 composition of the SABL over north central Mauritania in response to an approaching Saharan heat
577 low (SHL), which was migrating westward during that period (see Section 3.1). Flight F15 was
578 conducted to document the SABL over northern Mauritania together with a dust plume transported
579 from Algeria and associated with a Mediterranean wind surge.

580 The first three flights performed by the BAe146 on 17 and 18 June were a set of missions designed
581 to investigate very strong low level winds over northern Mali (b600, b601 and b602). During these
582 flights some of the largest particles encountered during Fennec were measured (see Section 4.1.1),
583 and elevated dust concentrations were seen at altitudes beneath 1km, although vertical mixing
584 played a role in the afternoon. The Falcon also flew on 17 June (F20) with a mission dedicated to the
585 documentation the SABL over northern Mauritania and northern Mali, west of an approaching
586 African easterly wave, as well as the structure the dust plume associated with a Mediterranean wind
587 surge.

588 Flight b603 was a calibration flight performed over the Canary Islands at high altitudes under clear
589 skies for the radiation instruments. Flight b604 was a LADUNEX EUFAR flight sampling dust which
590 had been uplifted more than 24 hours previously by a MCS and associated haboob over Mali, and
591 then transported over Mauritania by prevailing winds (Sodemann et al., submitted manuscript),
592 retaining giant mode dust particles despite large transport distances (Ryder et al., 2013a). The
593 BAe146 crossed the dust front at high and low altitudes for in-situ and remote sensing
594 measurements. F21 consisted of a long rectilinear flight across northern Mauritania and northern
595 Mali to survey the SABL as well as document the dust uplift in the region of the intertropical
596 discontinuity (ITD, i.e. the near surface convergence zone between the monsoon and the harmattan
597 flow) to the south of the SHL, over Mali.

598 On 21 June both the Falcon and BAe146 performed 2 flights (b605, b606, F22 and F23). On the
599 preceding day, convection over the Atlas Mountains had initiated a dust front which had propagated
600 southwards over Mauritania by 21 June, with aged dust overlying it. During the day the layers
601 became mixed together. Both aircraft missions' aimed to sample this dust and diurnal mixing, see
602 Section 4.3.3. On 22 June, again, both aircraft performed missions in the morning and afternoon
603 (b607, b608, F24 and F25). The missions were aimed at sampling the SHL and therefore flight tracks
604 extended well into Mali (Figure 2). LIDAR, dropsondes and radiation instrumentation were used to
605 sample the spatial and diurnal evolution of the SHL (see Engelstaedter et al. (2015)). F26 on 23 June
606 performed a mission dedicated to the study of the morning dust uplift over alluvial sources of
607 Northern Mauritania in connection with the decay of the low-level jet (RAIN4DUST project).

608 From 24 June onwards, dust conditions were generally more well-mixed vertically with less fresh
609 dust being sampled. Flight b609 on 24 June sampled dust and cumulus developing on the top of the
610 dust layers (see Section 4.1.4). Flight b610 sampled the low level jet and dust uplift mechanisms over
611 eastern Mauritania. b611 overflew the Zouerate ground supersite - see Section 4.2.2 for a
612 comparison of in-situ measurements to sunphotometer retrievals. b612 and b613 on 26 June were
613 missions to achieve radiative closure and to measure heat fluxes over the desert. Both were
614 performed under clear sky conditions with a series of stacked runs, under low dust loadings. Flight
615 b614 was a second flight to sample dust uplift and the low level jet early in the morning. Flight b615
616 on 28 June was the return transit to the UK, and included radiation calibration manoeuvres.

617 **3.2.3 Flights during Fennec IOP 2012**

618 Since the initial flying period during Fennec 2012 was initially dominated by Atlantic Inflow, with
619 dust being observed at the confluence of this and Saharan air (see Section 3.1.3), most of the earlier
620 flights aimed to sample this boundary (Table 7). b698 was a science transit from the UK to
621 Fuerteventura, during which calibration manoeuvres for radiation instruments were performed.
622 b699 and b700 were a pair of flights on 6 and 8 June which sampled the gradient of Atlantic Inflow
623 and its eastern boundary at high and low levels over northern Mali and northern Mauritania. b701
624 and b702 were similar flights, but here the edge of the Atlantic inflow was contingent with the ITD,
625 and larger dust loadings were sampled over central and southern Mauritania. Following b702 the
626 BAe146 landed at Dakar, and then returned to Fuerteventura over the Atlantic (b703) sampling
627 continental dust outflow. Flight b704 sampled Atlantic Inflow and the ITD again, this time measuring
628 the highest submicron aerosol optical depths (AODs) of Fennec, 3.4 at 550nm, over southern
629 Mauritania. b705 on 12 June was performed around midday to measure Saharan heat fluxes over a
630 stable pressure gradient.

631 Flights b706 and b707 were a pair of flights examining dust uplift over the Mauritania/Mali border,
632 with exactly the same track, and uplift beginning to happen under stronger winds during b707. b708
633 was designed to measure dust uplift by the LLJ over Mali under clear sky conditions so that the
634 radiative impact of the dust could also be measured. This flight saw the highest scattering
635 measurements on the nephelometer during the campaign (see Section 4.3.4), from dust at very low
636 altitudes. By contrast, b709 on 17 June sampled dust which had been transported into the SHL and
637 was well-mixed vertically up to 6km. This flight aimed to sample the pressure structure of the SHL
638 and also perform radiative closure. b710 overflew the Zouerate ground supersite as part of SAVEX in
639 order to compare AERONET retrievals and aircraft measurements of dust. Finally, b711 was a science
640 transit return to the UK.

641 **4 Key Scientific Results from the Fennec Airborne Programme**

642 Here we present key scientific results from the Fennec airborne programme. They are grouped by
643 dust characterisation (Section 4.1), Cross-platform assessment of dust measurements (Section 4.2),
644 Dust uplift and transport (Section 4.3) and SABL processes, dynamics and interactions with dust
645 (Section 4.4).

646 **4.1 Dust Characterisation**

647 **4.1.1 Size distributions**

648 During Fennec 2011 six different instruments were used to measure size distribution, as described in
649 Section 2, covering the size range 0.1 to 300 μm diameter. Of these, the PCASP, CDP and CIP
650 operated consistently during the whole campaign (see Rosenberg et al. (2012) for details of
651 calibration and errors). Very large particles were measured during Fennec 2011, with effective
652 diameter of the full size distribution ranging from 2.3 to 19.4 μm (Ryder et al., 2013b). Examples of
653 different types of size distribution are shown in Figure 4. The solid lines show measurements from
654 flight b600 at around 700 m above ground level, under aerosol optical depths greater than 3.0 at 550
655 nm when the dust was being actively uplifted by strong winds and was encountered beneath 1 km
656 above ground level. These were some of the largest particles encountered during Fennec 2011, and
657 the size distribution shows a strong coarse and giant mode present with a broad peak in volume

658 concentration from around 10 to 60 μm . Contrastingly, b612 (dashed lines) shows more aged dust
659 (24-48 hours based on satellite imagery) which was well-mixed within the SABL up to 5 km, with
660 optical depths at 550 nm of around 0.6. Here there are fewer particles across all sizes upwards of 0.5
661 μm compared to b600, and the peak volume concentration is now at 10 μm diameter, reflecting a
662 shift to lower number concentrations and fewer coarse particles as dust is mixed vertically through
663 the entire SABL, and larger particles are deposited during transport as well as dispersion decreasing
664 the total number concentration. Ryder et al. (2013a) examine the effects of vertical mixing and
665 transport on dust properties further. Interestingly, at smaller sizes than 0.5 μm there are more
666 particles in the case of b612, which gives the size distribution a flatter shape than b600. This may be
667 due to different dust sources, soil types and uplift wind speeds acting initially.

668 For the first time on the FAAM BAe-146 all size resolved particle measurements were made with
669 high temporal resolution (≥ 10 Hz) allowing correlation with vertical wind speed and measurements
670 of the eddy covariance particle fluxes. This technique has been previously employed to derive heat,
671 momentum and moisture fluxes from FAAM BAe-146 data (Petersen and Renfrew, 2009). During
672 Fennec we are have been able to resolve particle flux both in terms of eddy length scales and
673 particle diameter. During flights b600, b601 and b602 upward particle fluxes were observed
674 associated with synoptic scale wind in Algeria and northern Mali. Upward particle fluxes were also
675 observed during flight b604 again associated with synoptic scale winds in this area. In general it has
676 been found that particles above 10 μm diameter dominate the mass flux and in some cases particles
677 above 100 μm diameter make a significant contribution. Full details are provided in Rosenberg et al.
678 (2014).

679 4.1.2 Chemical Composition

680 To date, information on the mineralogical composition of coarse mineral dust can only be obtained
681 by post-field analysis of filter samples. The mineralogical composition is a fundamental property
682 determining the impacts on mineral dust on climate. It controls the complex refractive index,
683 determining the radiation interactions in the shortwave and longwave spectrum (relevant to the
684 direct radiative effect), the water uptake capability, determining the cloud and ice nuclei activation
685 efficiency (relevant to the indirect radiative effect), the solubility in water, controlling the capability
686 of deposited mineral dust to be assimilated by the marine phytoplankton, and the surface reactivity
687 relevant to interactions with the gas phase (Formenti et al., 2011a; Scheuvens et al., 2013).

688 The mineralogical composition of mineral dust is obtained by X-ray diffraction (XRD) (Caquineau et
689 al., 2002). Nonetheless, this technique is not always applicable to aircraft samples because of limited
690 sampling times yielding light loadings which are incompatible with the detection limits of this
691 analytical technique. Typically, about 800 μg of total dust mass are needed for analysis (Caquineau
692 et al., 1997). As an order of magnitude, this requires at least 1 hour sampling at high volume (~ 50 L
693 min^{-1}) for low to moderate atmospheric concentrations ($< 200 \mu\text{g m}^{-3}$) and at least half an hour for
694 concentrations of the order of $200 \mu\text{g m}^{-3}$ and above.

695 Alternatively, useful indications on the mineralogical composition of mineral dust can be obtained by
696 examining the concentrations of typical trace elements such as Al, Si, Fe, Ti, Ca, K, Mg, Na, which can
697 be obtained by X-ray fluorescence techniques which have typical detection limits of 10 μg or less
698 across a filter sample (Formenti et al., 2011b). In particular, the inter-elemental ratios provide
699 indications of the origin of mineral dust. Typically, Al is used as a unique tracer as alumino-silicates

700 dominate the dust mass. However, the Fe/Ca ratio has also proven useful to trace the origin of the
701 dust plumes (Kandler et al., 2007; Formenti et al., 2011a; Scheuvens et al., 2013; Formenti et al.,
702 2014).

703 Ninety-three samples are available in total from the Fennec 2011 and 2012 campaigns from the
704 BAe146 (fifty-five and thirty-eight for each field phase, respectively). Samples have been collected in
705 the Saharan boundary layer at altitudes ranging between 350 and 2700 m asl. The total dust
706 concentrations, estimated as the sum of oxides of Na, Mg, Al, Si, K, Ca, P, Fe, and Ti, varied between
707 22 to 4012 $\mu\text{g m}^{-3}$.

708 The analysis of Dust Uplift Potential (DUP, section 4.3.1) restricted to the filter sampling legs
709 suggests that the Fennec 2011 was characterized exclusively by emissions from Saharan sources in
710 Algeria, West Sahara and Mauritania, with the exception of samples from b604 where dust had been
711 uplifted by MCS outflow over Mali and transported by a large-scale haboob (Sodemann et al.,
712 submitted manuscript). However, during the Fennec 2012 period additional emissions of Sahelian
713 dust from convective activity in Mali constituted a much larger proportion of the samples. This
714 contrast is a result of the dominant heat low west phase during the latter half of Fennec 2011 driving
715 anomalous northeasterlies over western Algeria (Figure 3c, Section 3.1.2) compared to a northern
716 extension of the monsoon flow over Mali during Fennec 2012 and increased MCS activity (Section
717 3.1.3).

718 The elemental composition is consistent with the DUPs indications for those source regions. This is
719 shown in Figure 5a where the Fe/Ca and the Si/Al ratios obtained for the Fennec 2011 and Fennec
720 2012 samples are compared to those measured during the AMMA, DABEX, DODO and GERBILS
721 campaigns summarized in (Formenti et al., 2014). As a consequence, and with the exception of
722 samples collected during flights b699 and b700 when dust originated from the sources in the Algeria,
723 Western Sahara and Mauritania areas, samples collected during Fennec 2012 had a lower Ca and Mg
724 percent content with respect to Fennec 2011, reflecting the absence of calcium carbonates (calcite
725 and dolomite) in Sahelian soils (Journet et al., 2014).

726 Likewise, there is a clear difference between the measured single scattering albedo (SSA) at 550 nm
727 during Fennec 2011 and Fennec 2012 (Figure 5b). Even when excluding the outlier corresponding to
728 a pollution plume encountered during flight b710 at Zouerate during Fennec 2012, when the single
729 scattering albedo value averaged over the filter collection run was 0.91 (± 0.02), the mean single
730 scattering albedo value for the Fennec 2012 period is lower than that for Fennec 2011 (0.94 ± 0.01
731 and 0.97 ± 0.01 , respectively).

732 Future work will investigate the possible link between the changes in composition and optical
733 properties during the 2011 and 2012 periods. This will also involve taking into account the particle
734 size distribution, as a function of origin and of the age of the sampled air masses.

735 **4.1.3 Column Aerosol Loading from in-situ Measurements**

736 It is possible to use in-situ measurements of scattering and absorption by the nephelometer and
737 PSAP on the BAe146 respectively to calculate extinction profiles, and hence AOD. Measurements are
738 restricted firstly by the altitudes flown by the aircraft, which is usually from-between above the
739 aerosol layer to-and as close to the surface as is safe and permissible. Depending on visibility, this
740 varied between around 50 m to 1 km during Fennec. Secondly the measurements are restricted by

741 the aircraft inlets, which do not sample particles larger than around 2 μm (Ryder et al., 2013b). The
742 former has been accounted for by assuming that the aerosol profile is constant beneath the
743 minimum aircraft altitude to the ground, while the latter is not accounted for and therefore the
744 AODs presented here represent only extinction from the submicron size distribution, and are
745 therefore an underestimate. Scattering and absorption measurements are corrected as described in
746 Ryder et al. (2013b).

747 AODs from Fennec 2011 and 2012 are shown in Figure 6, with circles representing 2011 and
748 diamonds 2012. AODs ranged from 0.2 to 3.6 at 550 nm. Of particular interest were a few heavy dust
749 events which the aircraft sampled, including b600, b601 and b602 on 17 and 18 June 2011 in
750 northern Mali (orange, red and green circles), during which very large dust particles were measured
751 and dust fluxes have been calculated (as described in Section 4.1.1). Secondly, flights b707 (blue and
752 green diamonds on Mali/Mauritania border) and b708 (orange diamonds in northern Mali) in 2012
753 sampled very high dust loadings, the first with very low altitude, fresh dust (see Section 4.3.4), and
754 the second with dust well mixed to above 5 km, both under clear skies (i.e. no cloud). These flights
755 will make excellent radiation closure case studies. Thirdly, we draw the reader's attention to the
756 large number of profiles over the ocean between the land and Fuerteventura. The vertically resolved
757 changes in particle size and optical properties between fresh, aged and oceanic profiles are
758 examined by Ryder et al. (2013a), who find a significant reduction in particle size, number and
759 associated changes in optical properties for dust measured over the ocean.

760 4.1.4 Dust-Cloud Interactions

761 Saharan clouds have the potential to be significantly different to other continental mid-latitude
762 clouds due to the abundance of dust, which can act as ice nuclei (IN) and ~~(giant)~~ cloud condensation
763 nuclei (~~(G)~~CCN), and the fact that the hot dry boundary layer prevents precipitation reaching the
764 surface. Flight b609 on 24th June 2011 investigated a convective system in Northern Mauritania.
765 According to analyses from the Met Office operational Africa Limited Area Model, an overnight
766 monsoon surge associated with an easterly wave brought moist southerlies as far as 24°N at 8°W.
767 Over the course of the day a linear convective feature formed, extending from 18.5°N to link with a
768 system over the Atlas Mountains at 30°N. Dusty cold pool outflows, which affected supersite-2
769 (BBM), were visible in SEVIRI satellite imagery from at least 18 to 23 UTC. ~~The flight b609~~ consisted
770 of an overflight of the system and a series of north-south aligned legs at 8.0°W between 23.8 and
771 25.8°N on the eastern flank of the convective system from 12:42 to 15:36 UTC. The run locations
772 were restricted by operational constraints.

773 Figure 7 ~~shows the flight pattern and measurements. The flight path (thick black lines) consisted of~~
774 ~~An initial high-level leg, (LIDAR data in Figure 7) was~~ followed by a descent to minimum altitude and
775 then three legs beneath the clouds, each increasing in altitude to 4500m, just below cloud base
776 (5400 to 5800 m). Once above cloud base, a series of short legs were performed targeting three
777 cloud cells, with the aircraft finally ascending through the cloud tops at 8000 m ~~(See leg altitudes and~~
778 ~~locations in Figure 7). Cloud droplet concentration is shown in~~ Figure 7 ~~on top of the aircraft track,~~
779 ~~appearing red when the aircraft was in cloud. Range corrected Leosphere LIDAR backscatter signal is~~
780 ~~shown measured during the highest altitude aircraft leg, and is also shown beneath the aircraft~~
781 ~~descent where available, since the signal is strongly attenuated by the clouds along the high-level~~
782 ~~leg. Here we solely use the LIDAR measurements to describe the presence and structure of clouds~~

783 present; not for the vertical distribution of dust, due to the strong attenuation of the LIDAR signal by
784 both the clouds and the thick dust beneath them.

785 The initial LIDAR observations indicated that cloud tops ranged from 6.1 km to above the aircraft
786 altitude of 8.75 km, equivalent to approximately -11 to -28 °C (based on the profile measured during
787 the descent). It was observed visually from the aircraft cockpit that the cloud tops had no
788 observable anvil cirrus outflow. During descent to low level the aircraft passed through one isolated
789 cloud at 24.18 °N. LIDAR observations of this cell 13 minutes prior to the intersection provided a
790 cloud top height of 6.65 km, which is estimated to be at -15.0±0.2 °C. The LIDAR data showed no
791 links to, or particle flow between, any other clouds. Particle images recorded by the CIP showed that
792 this cloud consisted of pristine hexagonal plates. Freezing at this warm temperature is uncommon
793 even for clouds in the vicinity of a source of IN (Kanitz et al., 2011; Ansmann et al., 2008; Sassen et
794 al., 2003; Raymond and Blyth, 1989). An explanation could be the very high dust concentrations
795 acting as IN in the heart of the Sahara.

796 The descent to 500 m provided a measurement of the aerosol input into the cloud. At the surface
797 particle concentrations above 0.13 μm diameter measured by the PCASP and CDP ranged between
798 60 and 80 cm^{-3} south of 25.33 °N. North of this point the concentrations were 200 cm^{-3} . Note that
799 most of this concentration is measured by the PCASP and therefore does not show up on the
800 number concentration scale in Figure 7. As the aircraft climbed to cloud base the aerosol
801 concentration fell to 40 cm^{-3} , although the number of particles above 4 μm diameter rose from 0.05
802 cm^{-3} to 0.15 cm^{-3} .

803 During ascent back towards cloud base sporadic ice precipitation was observed by the CIP probe
804 from altitudes of 4.4 km (4 °C) and graupel was observed impacting the aircraft. Cloud was
805 encountered at 5.75 km (-8 °C) although cloud base could have been slightly lower (minimum of 5.4
806 km or -5 °C). It is of note that cloud base may have been too cold for the Hallett-Mossop ice
807 multiplication process which occurs around -6 °C. No columnar ice crystals typically produced by this
808 process were observed. Near cloud base the cloud was found to be mixed phase with droplet
809 number peaking at 250 cm^{-3} coincident with the peak updraft speed of 10 ms^{-1} . This measured
810 droplet concentration was found to be significantly higher than the aerosol concentration reported
811 by the PCASP and CDP below cloud base: the shortfall in CCN must have been made up of particles
812 smaller than the PCASP detection limit. Twohy et al. (2009) showed that dust with zero
813 hygroscopicity, κ , is entirely activated in cloud by a 10 ms^{-1} updraft and, because of their large size,
814 can dominate the CCN population over other hygroscopic particles when they have a small but non-
815 zero κ (Koehler et al., 2009). It is therefore likely that dust particles were acting as CCN or GCCN in
816 this case. Higher in the cloud there is evidence of liquid water in updraft regions and near cloud top
817 a population of homogeneously nucleated bullet rosettes were observed. No cirrus or precipitating
818 particles were observed above cloud top.

819 These measurements have shown that dust is likely acting as a CCN and is acting as an IN at
820 temperatures of -15 °C. Sampling of further clouds earlier in their evolution would provide further
821 limits on the effectiveness of dust as an IN. The lack of Hallett-Mossop in these clouds makes them a
822 useful case for assessing IN concentrations and the extreme size of the dust particles may provide
823 tests of the impact of GCCN.

824 For a non-precipitating cloud we expect that equivalent potential temperature, θ_e , and total water
825 concentration (condensed plus vapour) are conserved and hence any point in cloud should lie on a
826 mixing line or in a mixing region of these parameters (Paluch, 1979; Blyth et al., 1988). Here the
827 cloud is precipitating meaning that total water concentration is no longer conserved but these
828 variables are still useful in diagnosing the transport and mixing processes (Figure 8).

829 Much of the sampled in-cloud air had higher water content, greater than 5 g kg^{-1} , but similarly high
830 ranges of θ_e compared to boundary layer air (Figure 8). These are inconsistent with clouds being a
831 simple mixture of boundary-layer and entrained air. Out of cloud, above cloud base air had some
832 regions consistent with simple mixing, some in a similar moist warm region to in-cloud air, but also
833 some regions with low moisture content less than 1 g kg^{-1} and high θ_e similar to boundary layer air.
834 Profiles of water vapour mixing ratio (WVMR) (Figure 7) show that in the boundary layer (below
835 2500-5000 m, varying from profile-to-profile) WVMR increases with altitude. Similar behaviour was
836 also seen in the mean WVMR profile at Fennec supersite 1 (BBM) between 15 and 18 UTC, the time
837 of maximum cloudiness (Marsham et al., 2013b). This is again inconsistent with simple mixing of a
838 growing boundary layer. We hypothesise that in this low shear environment precipitation is
839 evaporating in the boundary layer air but is not able to arrest the updraft allowing water to be
840 recycled and concentrated in the cloud. High θ_e air rises in the boundary layer and receives extra
841 water from evaporating precipitation, such that when it enters the cloud base it has more moisture
842 than its environment. Some In cloud air parcels may either precipitate adding to the recycled
843 moisture reservoir before being detrained as dry, high θ_e air, or they do not whereas other parcels
844 may not precipitate and are instead detrained as moist, high θ_e air. We also expect dust and aerosol
845 to be affected by this recycling process. Precipitation accumulates CCN and upon total evaporation
846 releases them as a single aggregate particle. The increase in large dust particle concentration below
847 cloud base is qualitatively consistent with this expectation. This concentrating of moisture and dust
848 in the boundary layer top and the modification of the dust size distribution has implications for long
849 range transport of these atmospheric constituents. This concentration of moisture at mid levels
850 could be extrapolated to dust and aerosol which are also redistributed by precipitation and could
851 cause a peak in dust loading at or near the boundary layer top which could subsequently be
852 adverted over broad regions. This process could be responsible for the increase in large dust
853 particles found just below cloud base. To our knowledge these are the first observations of such a
854 mechanism increasing the moisture content within the SABL mid-levels.

855 4.1.5 Dust-Ozone Interactions

856 Heterogeneous uptake of photochemical species leads to changes in the gas-phase composition of
857 the atmosphere; affecting the global ozone budget (Bauer et al., 2004). Previous campaigns have
858 observed ozone depletion during high dust loadings (de Reus et al., 2000; de Reus et al., 2005).
859 These have also been investigated through modelling studies (Bian and Zender, 2003) and laboratory
860 studies (Chang et al., 2005; Hanisch and Crowley, 2003). ~~Whilst~~ ~~it~~ There is still some debate as to
861 whether the removal of ozone is due to heterogeneous chemistry on the surface of the dust or a
862 feature associated with a change in air mass between high and low dust loadings. The alkalinity of
863 mineral dust has been shown to enhance the uptake of gases on the surface (Grassian, 2002). Bauer
864 et al. (2004) propose that the coarse mode of mineral dust could be important for heterogeneous
865 uptake; whilst Chang et al. (2005) found that there was no mass accommodation limitation to the
866 rate of ozone uptake coefficients, concluding that freshly emitted Saharan dust is potentially a
867 significant route of ozone loss. Hanisch and Crowley (2003) discussed that mineral dust surface sites

868 could be deactivated by the extended presence of ozone. Ultimately the change in the surface of
869 mineral dust may have repercussions for subsequent aerosol-cloud interactions and modify the
870 cloud nucleating properties of the mineral dust. A number of case studies observed during the
871 Fennec campaigns were investigated Brooke (2014).

872 Fennec flight profiles provided the opportunity to sample very recently lofted mineral dust which
873 will not have undergone significant atmospheric 'processing' and thus provide a good opportunity to
874 investigate heterogeneous dust/ozone interactions. These observations of decreased ozone
875 concentrations correspond with increased mineral dust surface area associated with elevated dust
876 concentrations. Figure 9 presents box and whisker diagrams of mineral dust mean surface area
877 correlated with ozone mass mixing ratios observed during b707, where dust uplift was encountered
878 at the far eastern section of the flight track in Northern Mali (orange line in Figure 2d). The red
879 central line of the box and whisker denotes the median, the edges of the box are the 25th and 75th
880 percentiles and the whiskers extend to the most extreme data points. Mean surface areas of 0.15 to
881 $0.35 \mu\text{m}^2\text{cm}^{-3}$ (roughly count median diameters from 0.22 to 0.33 μm) correspond to ozone mass
882 mixing ratio of 49 – 52 ppb. As the mean dust surface area increases to 0.45 to 0.75 $\mu\text{m}^2\text{cm}^{-3}$ (count
883 median diameters from 0.38 to 0.49), the ozone mass mixing ratio decreases to 41 - 44 ppb. The
884 spread in ozone concentrations at mean surface areas of $0.45 \mu\text{m}^2\text{cm}^{-3}$ is associated with crossing
885 into a Harmattan airflow.

886 These in-situ observations suggest that increased mineral dust surface area associated with fresh
887 dust uplift and a large coarse mode contribution to the size distribution act as a route for the
888 reduced ozone concentrations. However, from the analysis presented here it is not possible to
889 unequivocally conclude if the air mass initially contained lower ozone concentrations and mineral
890 dust has subsequently been uplifted, or that mineral dust uplift could have contributed to the
891 reduced ozone concentrations observed. There is scope within the Fennec dataset to further
892 investigate airmass source regions, [potentially with Lagrangian study methods](#).

893 4.2 Cross-Platform Assessment of Dust Measurements

894 4.2.1 Falcon LIDAR and Satellite Validation

895 Aircraft data can play an important role in validating satellite-based retrievals of AOD, covering a
896 more extensive spatial area than that which is viewed from fixed ground-based measurements.
897 Particularly useful in this regard are active remote sensing observations from LIDAR, since they can
898 sample the full depth of the atmosphere below the aircraft instantaneously (i.e. a physical vertical
899 profile by the aircraft is not required) and can provide vertically resolved information.

900 In Figure 10, middle panel, we show an example of the level of agreement seen between three
901 different co-located measures of AOD, one provided at 532 nm by the LIDAR LNG on the F20, one
902 from MODIS Aqua, derived using the Deep-Blue algorithm collection 5.1 (Hsu et al., 2004) and one
903 from the SEVIRI instrument on Meteosat-9 (Brindley and Russell (2009); Banks and Brindley (2013)),
904 both at a wavelength of 550 nm. Here we focus on an afternoon flight (F23, see Figure 2c) made by
905 the Falcon on a track leading across to northern Mali from northern Mauritania on the 21st June
906 2011.

907 The satellite observations are co-located spatially with the LIDAR by averaging the satellite pixels
908 within 25 km of each LIDAR pixel. Temporally, the Aqua satellite overpass time is always within 90

909 minutes of the aircraft observations, with a minimum time difference of 37 minutes. For SEVIRI we
910 take advantage of the improved temporal sampling available from geostationary orbit such that
911 each LIDAR observation is within 30 minutes of the corresponding satellite retrieval. The lower panel
912 in the figure shows the vertical extinction coefficient derived from the LIDAR observations, while the
913 top coloured band illustrates the colouring of the standard 'desert-dust' Red-Green-Blue (RGB)
914 composite (Lensky and Rosenfeld, 2008) extracted from SEVIRI along the flight track.

915 Looking at the middle panel, the longitudinal behaviour of the AOD derived from all three
916 instruments is generally in good agreement although SEVIRI tends to show consistently higher AODs
917 than those derived from the LIDAR and from MODIS. The MODIS retrievals contain more data gaps
918 as a result of various data quality tests: both the LIDAR and SEVIRI retrievals and the RGB composites
919 suggest that these tests may be a touch severe as there is no clear evidence of a break in the aerosol
920 layer or the presence of cloud. The intense pink colour of the composite at the western edge of the
921 track would suggest the largest dust loadings are located here, associated with a thick dust plume at
922 an altitude of \sim 3 km and another distinct layer observable at \sim 5.5 km seen in the LIDAR profile
923 (which may have originated from Mali on the 19th). By the eastern end of the track, the AODs
924 measured by MODIS, SEVIRI and the LIDAR are slightly smaller than the values seen at the western
925 end, the dust is much more uniformly spread throughout the lowest 5 km or so of the atmosphere,
926 and the intensity of the RGB signal is somewhat reduced.

927 Further work has explored co-located aircraft and satellite data in more detail, utilising a more
928 extensive suite of satellite instruments (such as the MISR instrument on Terra and the IASI
929 instrument on the METOP satellites (Banks et al., 2013), and between the BAe146 in-situ
930 measurements and space-borne LIDAR CALIOP (Pappas et al., in prep.). In the former study the
931 differences between retrievals have been investigated, including an evaluation of the sensitivity of
932 the retrievals to variations in dust loading, as well as to atmospheric conditions (such as column
933 water vapour), surface features (such as albedo), and to aerosol height. As diagnosed by Banks et al.
934 (2013), when the dust loadings are high the SEVIRI retrievals appear most capable of retrieving the
935 appropriate AODs, whereas the other retrievals are biased low. On the other hand the SEVIRI
936 retrievals are most sensitive to meteorological conditions, especially column moisture, under high
937 levels of which the SEVIRI retrieved AODs are biased high; meanwhile the MODIS Deep Blue and
938 MISR aerosol retrievals appear to be relatively insensitive to such factors. The aircraft data will be of
939 substantial benefit in interpreting the 'desert-dust' RGB imagery.

940 **4.2.2 Comparison of AERONET and aircraft size distributions**

941 Previous work (Müller et al., 2012; Müller et al., 2010; McConnell et al., 2008) has found relative
942 disagreement between aircraft and AERosol Robotic NETwork (AERONET) size distribution retrievals
943 for dust. Considering the wide application of size distributions from AERONET retrievals such as to
944 aerosol models and climate forcing assessments (e.g. Garcia et al. (2012); Kinne et al. (2003)), it is of
945 importance to repeat comparisons of this type. Moreover, some discrepancies have been found
946 between retrieved size-distributions using the AERONET algorithm (Dubovik and King, 2000; Dubovik
947 et al., 2006) and the same size-distributions derived with the SKYRAD algorithm (Nakajima et al.,
948 1996), as described in Campanelli et al. (2012) and Estellés et al. (2012b). The SAVEX project aims to
949 explore these discrepancies, and has been motivated by studies such as Estellés et al. (2012a) and
950 Estellés et al. (2012b), where differences between different sunphotometer retrieval algorithms are
951 examined.

952 AERONET CIMEL sunphotometers were installed and operated at the two supersites of Zouerate
953 (western Mauritania) and Bordj-Badji Mokhtar (BBM, Algeria) as part of the Fennec programme. As
954 part of the SAVEX project, sunphotometers were also installed and operated at several different
955 sites on Tenerife during June 2012 with the intention of overflying the instruments during dust
956 events. However, overflights were not performed at Tenerife due a lack of dust outflow in this
957 location during the campaign. The aircraft range from Fuerteventura did not permit overflights at
958 BBM. Therefore overflights as close as possible to the Zouerate station under dusty conditions were
959 performed during 2011 (b611, 25 June) and 2012 (b710, 18 June, SAVEX flight).

960 During these flights, profiles and stacked legs were performed to measure in-situ aerosol properties
961 and radiative measurements, to allow radiative closure of the column above the ground site.
962 Radiative flux measurements were also made at the ground site. Here we present some
963 measurements from b611 in 2011. Dust sampled during this flight was around 19 to 43 h old,
964 originating from Algeria (Ryder et al., 2013b), with AERONET AODs at 440nm from 0.8 to 0.94, and
965 was relatively well mixed in the SABL up to around 5.5km, although extinction coefficient
966 measurements from the aircraft approximately doubled beneath 2.5km. Similar measurements are
967 available from flight b710, although for that flight layers of anthropogenic pollution were detected
968 between dust layers, thus making comparisons between platforms more complicated, and are not
969 shown here.

970 Figure 11 shows a comparison of the size distributions measured by the BAe146 compared to
971 AERONET retrievals on 25 June 2011. The in-situ aircraft measurements are taken over a vertical
972 profile close to Zouerate on 25 June 2011 between 8 km to 80 m AGL from 1558 to 1627 UTC. In-situ
973 size distribution measurements shown in Figure 11 are therefore shown as the median, 10th and
974 90th percentiles between 80 m and 5.5 km.

975 Sunphotometer retrievals of size-distribution from almucantar scans are not present during much of
976 the day due to cloud cover over Zouerate. Nevertheless, several retrievals are available during the
977 morning (dark blue), one during the flight (black), and two from 18:06 and 18:30 after the aircraft
978 had left the region (light blue). Size-distribution retrievals shown are those directly available from
979 AERONET (L1.5, V2) and converted to $dV/d\log D$ to match the aircraft measurements, and adjusted to
980 measurements in cm^{-3} assuming the dust layer is distributed evenly over 5.5 km. Further work will
981 examine measurements from aircraft legs at different altitudes, and different ways of representing a
982 column-average measurement from the aircraft measurements, such extinction-weighted averaging.

983 The median aircraft measurements show a peak volume concentration at 12 μm , while the AERONET
984 retrievals show peaks between 3 to 6 μm . This is consistent with previous aircraft-AERONET
985 comparisons finding larger particles measured by aircraft (Reid et al., 2003; Müller et al., 2012;
986 Müller et al., 2010; McConnell et al., 2008). However, one retrieval only shows a peak volume
987 concentration at 13 μm which appears to agree much more closely with the shape of the size
988 distribution from the aircraft measurements. Satellite images show a small convective cloud
989 developing close to, but not over Zouerate around this time. It is possible that small scale
990 downdrafts produced some freshly uplifted dust which may have resulted in different size
991 distribution retrievals. At sizes smaller than 3 μm differences in volume concentration are
992 substantial between AERONET and the aircraft, with AERONET reporting more particles. Further
993 work will explore possible factors causing this difference.

994 Rather few coarse particles were seen during b611 ([towards the end of the campaign](#)) relative to the
995 rest of Fennec, perhaps due to the aged nature of the dust which meant that the largest particles
996 had already been deposited. This is reflected by the absence of particles larger than 16 μm in the
997 median, and the absence of particles larger than 45 μm in the 90th percentile (see the one CIP data
998 point for the 90th percentile), though particles of these sizes *were* measured, but the standard
999 deviation was very large, as shown by the large error bars on the median above sizes of 16 μm .

1000 ~~Flight b611 took place towards the end of the Fennec 2011 campaign, when dust conditions were~~
1001 ~~generally more aged with lower AODs, and the contribution from larger particles as shown in (Ryder~~
1002 ~~et al., 2013b) was rather weak.~~ Unfortunately the flights during Fennec when large particles were
1003 strongly evident did not take place close to AERONET sites, due to the remoteness of the flight
1004 locations. Ryder et al. (2013b) find that particle sizes are larger close to dust sources in remote
1005 locations, and Ryder et al. (2013a) show that giant particles ($d>37.5 \mu\text{m}$) are a feature of freshly
1006 uplifted dust events, and some long-range transported cases. This should act as a caution for using
1007 AERONET retrievals as a basis for dust size distributions over the central Sahara, since they only
1008 extend to 30 μm diameter, the tails of the size distributions are constrained to very small values
1009 (Hashimoto et al., 2012), and encounter large errors (Dubovik and King, 2000). Further studies will
1010 examine aircraft and sun-photometer data from both 25 June 2011 and 18 June 2012, in terms of in-
1011 situ aircraft measurements, airborne and ground-based radiation measurements, and using both the
1012 AERONET and SKYRAD retrievals for the inversion of sun-photometer radiances.

1013 4.3 Dust Uplift and Transport

1014 4.3.1 Dust source areas from Dust Uplift Potential

1015 It is relevant for several areas of dust measurement analysis to identify the sources of dust sampled
1016 during research flights (e.g. Section 4.1.2). Lagrangian backward trajectory calculations with the
1017 FLEXPART model (Stohl et al., 2005) have been initiated in 'tropospheric curtains' run along the track
1018 of each research flight to investigate the sources of the dust sampled. For this a large number of
1019 virtual air parcels (1000) were released at a 30 s interval in a vertical column between the surface
1020 and a pressure of 200 hPa along the flight tracks. Each parcel was tracked for 3 days backward in
1021 time using ECMWF analysis winds at a $1^\circ \times 1^\circ$ horizontal grid spacing. We utilize the metric of Total
1022 dust uplift potential (DUP), defined as $fU^3(1+U_t/U)(1-U_t^2/U^2)$, with f being the desert and bare soil
1023 fraction, the wind velocity U , and the threshold velocity $U_t=6.5 \text{ ms}^{-1}$ (Marsham et al., 2011). Despite
1024 being a simplified representation of likely dust uplift (e.g. variations in soil moisture are neglected,
1025 and dust uplift may not be linear with threshold velocity (Kok et al., 2014), DUP is a useful indicator
1026 of where likely uplift occurred and is relatively easily computed. DUP was then calculated along the
1027 three-day back-trajectories for locations where the tracked air parcels were within the boundary
1028 layer. DUP values were gridded on a $0.25^\circ \times 0.25^\circ$ grid and integrated over time. The DUP thus
1029 calculated for the tropospheric column at the aircraft location characterises the airmass as
1030 measured by the onboard LIDARs when the BAe146 and Falcon were flying at high altitudes. During
1031 lower flight legs this analysis enables interpretation of in-situ dust measurements with respect to
1032 their mobilisation conditions and source regions.

1033 Figure 12 shows the composite of the DUP from (a) all the Fennec 2011 Falcon flights, (b) Fennec
1034 2011 BAe146 flights and (c) Fennec 2012 BAe146 flights. The areas contributing to the sampled air
1035 masses, which experienced strong winds that would be associated with dust uplift for dust-source

1036 regions (i.e. high DUP areas) were mostly located in a NE-SW oriented swath extending from central
1037 Algeria to northern Mali and Mauritania during 2011. This dominant pattern is related to the inflow
1038 into the Saharan heat low, as shown by the 925 hPa winds in Figure 3c over southwest Algeria. DUP
1039 locations from 2012 suggest more southerly dust sources, from southern Mauritania, stretching to
1040 the Mali-Algeria-Niger triple point, and along the Mali-Algeria border towards southern Libya. This is
1041 consistent with additional convective activity in Mali driving emissions which were more Sahelian-
1042 dominated during 2012 (Section 3.1.3).

1043 Individual flights exhibit additional sources and substantial variability (see supplementary material
1044 for DUP maps for individual flights). For example, dust from more southerly sources in Mali and
1045 Mauritania was intercepted during flights b600-602, b604-b606, b608, b611 and b614. Dust from
1046 northern Niger was sampled during flight b607. Note that the connection to dust filter samples to
1047 this figure is not immediate, because only the DUP for the selected legs corresponding to the filter
1048 sampling duration and position are considered in that case (see Section 4.1.2). We note that DUP
1049 from events associated with convective downdrafts such as haboobs may not be accurately
1050 represented due to the ECMWF analyses not fully capturing these events (Marsham et al., 2011). For
1051 example, this is the case for b604, where a large MCS generated a haboob over Mali which
1052 subsequently travelled towards Mauritania (Sodemann et al., submitted manuscript). Therefore in
1053 situations where dust has potential to have been uplifted by events associated with convection, back
1054 trajectories and more generally operational meteorological analysis and forecast data should not be
1055 used in isolation to determine dust source regions. For example, a combination of analysis of SEVIRI
1056 RGB satellite imagery and Lagrangian methods can be used to ensure consistency with observations
1057 (e.g. Ryder et al. (2013b)).

1058 4.3.2 Heavy dust loadings from a low-level jet breakdown over northern Mali

1059 One particularly notable flight was b600 during the morning of 17th June 2011, under which the
1060 highest dust loadings observed during Fennec 2011 and very large particles were measured. This was
1061 followed by flight b601 in the afternoon, and b602 the following morning in the same region. At this
1062 time the SHL was centred on the Mali-Algeria-Niger triple point, producing strong low-level
1063 northeasterlies through Algeria to northern Mali, which were particularly pronounced on the
1064 morning of the 17th (b600, Figure 13a, b, c, d, e). A region of slacker winds in Mauritania was
1065 associated with moisture remaining from the monsoon flow. Flights b600 to b602 were aimed at
1066 sampling these airmasses, travelling out at high-level to descend into the strong winds in northern
1067 Mali and returning northwestwards at low-level into the moister airmass (Figure 13ad, be). In-situ
1068 aircraft profile measurements are shown in Figure 14.

1069 Forecasts showed a pronounced decrease in the strong 925 hPa winds in northern Mali from 06 to
1070 09 UTC, with a corresponding increase in 10 m winds, consistent with the downward mixing of
1071 momentum from the nocturnal LLJ around the SHL, likely deflected around the Hoggar mountains
1072 (Birch et al., 2012). The existence of a LLJ is confirmed by the observation from the b600 descent
1073 into Mali (Figure 14, black) of a wind-maximum of 16.7 ms^{-1} at a pressure height of 1700m (1400m
1074 AGL), located above the growing turbulent moist and dusty CBL found below 1400m AGL. The dust
1075 number and mass concentrations below 1400m were the highest observed during the Fennec 2011
1076 campaign with particularly large particles observed during b600 and b601; the size distribution
1077 during the initial part of the horizontal run in the dusty CBL following the profile descent of b600 can
1078 be seen in Figure 4, with particles present up to nearly 300 μm . The high dust concentrations are

1079 consistent with the very high extinction measurements from the nephelometer and PSAP, of over
1080 1250Mm^{-1} in both profile descents ([Figure 13](#)[d](#)[Figure 14](#)). By the time of the profile descent of b601
1081 at approximately 1700Z the dust had been mixed up into a CBL that reached 3.7 km (Figure 14, red),
1082 with no remaining LLJ. The upwards vertical mixing of the dust resulted in the ‘pinkness’ in the
1083 SEVIRI images ([Figure 13a, b](#)[d, e](#)) becoming more pronounced by the time of the second flight (the
1084 RGB product is sensitive to dust altitude, Brindley et al. (2012)). Flight b601 then travelled back
1085 under the moist convection developing over Mauritania, with some precipitation observed falling
1086 onto the aircraft, but no extensive cold-pool outflows at the aircraft altitude at this time.

1087 To the authors’ knowledge this is the first airborne observation of dust size distributions (including
1088 the presence of coarse and giant particles) measured under uplift conditions caused by the
1089 breakdown of the Saharan nocturnal LLJ. Flights b706, b707 and b708 (Section 4.3.4) from 2012 also
1090 collected in-situ measurements of dust under LLJ breakdown conditions, thus providing scope for
1091 further analysis.

1092 4.3.3 In-situ sampling of an aged Haboob

1093 Recent studies have shown that haboobs (dust fronts occurring at the leading edge of cold pools
1094 emanating from convective storms) are a significant source of dust over the Sahara and Sahel
1095 (Flamant et al., 2007; Knippertz et al., 2007; Schepanski et al., 2009; Tulet et al., 2010). For example,
1096 Marsham et al. (2008b), Marsham et al. (2013b) and Allen et al. (2013) show that haboobs cause
1097 around 50% of dust uplift in the summertime Sahara, contributing to the seasonal cycle in dustiness.
1098 Radiosonde observations show that the transport of cold moist air in haboobs was a major cause of
1099 global model forecast bias at the Fennec BBM supersite in June 2011 (Garcia-Carreras et al., 2013),
1100 [consistent with the role of haboobs diagnosed from convection-permitting simulations](#) (Marsham et
1101 al., 2013a).

1102 On 21 June 2011, aircraft measurements were taken over and through an aged haboob emanating
1103 from convection over the Atlas Mountains in Morocco (Kocha et al., 2013). The cold pool passed
1104 over dust sources and uplifted large quantities of dust. The haboob was observed over the central
1105 Sahara over northern Mauritania and northern Mali in the morning with the LNG LIDAR on the
1106 Falcon 20 during flight F22 (see Figure 2b).

1107 The haboob appears as the layer characterized by large extinction coefficient values at pressure
1108 heights beneath 1.5 km (Figure 15a). The aerosol optical thickness (AOT) derived from the LIDAR
1109 extinction coefficient profiles reached an average of 1 around 0900 UTC. At the same time, the
1110 BAe146 flew through the haboob to directly sample its characteristics during flight b605. In-situ
1111 measurements from the BAe146 show that the dust concentration and observed extinction in the
1112 cold pool air increased by a factor of around three compared to its environment. The number of
1113 large particles of size around $10\text{ }\mu\text{m}$ increased to $0.1\text{ cm}^{-3}\mu\text{m}^{-1}$ (not shown). The properties of the
1114 dust sampled during this event also had a significant impact on the radiative fluxes within the
1115 haboob. For instance, the downward shortwave flux measured by the BAe146 decreased by 100
1116 Wm^{-2} when entering the dusty cold pool (Figure 15b).

1117 In the afternoon, both aircraft sampled the growth of the SABL again (flights F23 and b606) as the
1118 haboob was mixed into the Saharan residual layer above. An unambiguous influence of the haboob
1119 composition and thermodynamics was observed on the development of the SABL (Kocha et al.,

1120 2013). Simulations with and without dust are being used to investigate role of the haboob on the
1121 dynamics/thermodynamics on the development of the SABL over the central Sahara.

1122 **4.3.4 Radiation observations during dust uplift**

1123 Several flights were performed during Fennec to use aircraft in-situ aerosol measurements and
1124 radiative measurements to allow the potential to achieve radiative closure and examine the
1125 radiative properties of dust. Flight b708 on 16 June 2012 aimed to observe freshly uplifted dust at
1126 the time of downwards mixing of strong LLJ winds to the surface which was forecast to uplift dust
1127 over the Mali/Mauritania border. Additionally since clouds were absent, the flight aimed to attain
1128 radiative closure measurements since the dust loadings were high but with very low altitude dust,
1129 with AODs at 550nm of 0.54 and 1.92 measured during the two aircraft profiles by the nephelometer
1130 and the PSAP.

1131 Figure 16b shows information from the aircraft profiles – extinction calculated from corrected
1132 scattering and absorption measurements is shown for the descent (black) and ascent (red) in Mali.
1133 During this flight, the aircraft flew a high level leg at 7.5km for radiative measurements, followed by
1134 a profile down to minimum safe altitude, which was around 100m above ground level (AGL) initially
1135 (see black line in Figure 16a). During the descent the aircraft entered the dust layer at around 900 m.
1136 At this time the dust was not visible in the SEVIRI RGB desert dust imagery, despite an AOD of 0.54,
1137 likely because the RGB imagery is sensitive to dust altitude (Brindley et al., 2012). Absence of a 'pink'
1138 signal in the SEVIRI RGB imagery during active dust uplift such as occurred during this flight would
1139 have major implications for dust source maps that have previously been created based on this
1140 imagery (e.g. Schepanski et al. (2007)). Following the descent, the aircraft flew a low level leg. Figure
1141 16a shows the extinction as a function of longitude. As the aircraft flew eastwards the amount of
1142 dust increased until visibility was so poor that the aircraft had to ascend to 400m AGL. Despite this,
1143 extinction continued to increase to the east, with a maximum of 5500Mm^{-1} , the highest value ever
1144 observed from the FAAM nephelometer and PSAP.

1145 At the end of the low level leg, the aircraft ascended (red line in Figure 16b). The dashed lines in
1146 Figure 16b show potential temperature, which show inversions at the height of the rapid increases in
1147 dust extinction. This is one example of many during Fennec, where the dust was encountered in a
1148 low layer, which was gradually mixed upwards during the day as the SABL grew. The red line in
1149 Figure 16a shows the measured downwelling shortwave irradiance (SWD) during the low level run.
1150 Note that during the legs (around 30 minutes) the solar zenith angle decreased so that SWD would
1151 be expected to increase with increasing longitude. Instead during the western portion of the leg,
1152 SWD decreases with increasing extinction (dust above the aircraft). During the eastern portion of the
1153 leg there is a notable drop in SWD of around 150 Wm^{-2} at around -5.7W at the same time as the
1154 peak in extinction. This flight, as well as b709 in the SHL where dust was well-mixed vertically up to
1155 5km, will be used to examine the radiative effect of dust over the Sahara under different dust
1156 conditions (low level, well-mixed vertically) further, using the spectral radiation instruments SHIMS,
1157 ARIES and SWS on the BAe146 in conjunction with radiative transfer models and satellite
1158 observations.

1159 **4.4 SABL Boundary Layer Processes, Dynamics and Interactions with Dust**

1160 **4.4.1 LIDAR and Dropsonde Observations**

1161 Combining LIDAR observations and dropsonde-derived atmospheric profiles allows for a detailed
1162 analysis of the spatial and vertical structure of the atmosphere as well as the boundary layer
1163 processes that control the emission, vertical mixing and transport of mineral dust. Flights b607 and
1164 b608 were part of an extensive survey of the troposphere in the SHL region with the aim a) to
1165 characterise the spatial variability the SHL, CBL, monsoon inflow and dust distribution in the central
1166 Sahara, b) to analyse how these features change throughout the day, and c) to assess the processes
1167 that control these features and dust dynamics. Both flights followed a straight track crossing from
1168 northern Mauritania into Mali in the morning of 22 June 2011 (see Figure 2c for b607 flight track; the
1169 afternoon flight b608 overlies b607). The aircraft sampled the flight track twice in the morning
1170 (b607) and the afternoon (b608) allowing the evolution of the atmosphere over time to be studied.
1171 Dropsonde measurements were obtained during the out and return flight at fixed locations.
1172 Dropsonde data were interpolated to reference times at each location thereby creating a snapshot
1173 of the state of the atmosphere in the study region at the reference times. Engelstaedter et al. (2015)
1174 analysed the observed SHL characteristics and evaluated the performance of the UK Met Office
1175 limited area model for Africa (Africa-LAM). They identified two moisture transport pathways, one
1176 curving around the SHL core in the north (especially pronounced in a morning near-surface layer),
1177 and the other going towards the northeast within the roughly 2 km deep monsoon surge. The deep
1178 afternoon CBL simulated by the Africa-LAM in the monsoon surge region (more than twice as deep
1179 as observations) suggests a significant model error due to moisture being vertically mixed into
1180 northeasterly flow above about 2 km.

1181 As an example for the combination of observations from different instruments Figure 17 shows
1182 Leosphere LIDAR and dropsonde-derived observations data from the BAe146 flight b607, which was
1183 flying on a straight track crossing from northern Mauritania into Mali in the morning of 22 June 2011
1184 (see Figure 2c for b607 flight track; the afternoon flight b608 overlies b607). This flight was part of
1185 an extensive survey of the troposphere in the SHL region with the aim a) to characterise the spatial
1186 variability the SHL, SABL, monsoon inflow and dust distribution in the central Sahara, b) to analyse
1187 how these features change throughout the day, and c) to assess the processes that control these
1188 features and dust dynamics.

1189 The aircraft sampled the flight track twice in the morning (b607) and the afternoon (b608) allowing
1190 the evolution of the atmosphere over time to be studied. Dropsonde measurements were obtained
1191 during the out and return flight at fixed locations. Dropsonde data were interpolated to reference
1192 times at each location thereby creating a snapshot of the state of the atmosphere in the study
1193 region at the reference times. The range corrected LIDAR backscatter signal (see Section 2.1 for
1194 more detail on the LIDAR measurements) is shown here as coloured blocks and has a vertical
1195 resolution of 45 m and an integration time of 1 min. It is used here as an indicator for the presence
1196 of dust and clouds in the atmosphere but limitations apply. For instance, attenuation of the laser
1197 beam when it passes through an elevated dust layer can limit the LIDAR's ability to detect dust at
1198 lower levels. Dropsonde observations allow for the identification of atmospheric structures such
1199 as the top of the CBL and SRL as well as temperature inversions. The CBL depth was determined by
1200 locating the altitude in the sonde profile (from the surface upwards) where the potential air
1201 temperature (θ) first reaches 0.3°C above the value at 150 m above the surface. In cases where θ

1202 increased monotonically from the surface up to 150 m, the surface θ value was used as a reference.
1203 The top of the SRL was determined manually where possible by identifying a sharp decrease in water
1204 vapour mixing ratio coinciding with a sharp increase in θ . The resulting CBL and SRL tops were linked
1205 by solid lines in Figure 17 in order to illustrate spatio-temporal changes of these features. The depth
1206 of air temperature inversions, defined as an increase in air temperature with altitude, are indicated
1207 as grey boxes along the vertical sonde tracks together with the inversion strength in $^{\circ}\text{C km}^{-1}$ (Figure
1208 17).

1209 Dropsonde-derived near surface winds ranging between 11 and 17 ms^{-1} observed during the b607
1210 outgoing flight (not shown) resulted led to local dust emissions observed by the LIDAR at about
1211 7.3°W (also seen in LIDAR depolarization data, not shown) that were prevented from upward mixing
1212 by a low-level temperature inversion (Figure 17a). At that time in the morning, the CBL was still
1213 relatively shallow (mostly <1 km deep), the top of the SRL varied between about 4.3 and 5.5 km
1214 above MSL, and an aged dust layer of varying intensity could be identified close to the SRL top. Cloud
1215 development was identified west of 11°W in the LIDAR data. In the time that passed between the
1216 outgoing and return flight, surface emissions ceased and the CBL grew up to about 4.5 km above
1217 MSL (B4, Figure 17b) as a result of increasing near surface temperatures. up to about 2.8 km above
1218 MSL (Figure 17b). East of about 7.58°W, the CBL was prevented from growing deep by temperature
1219 inversions and the influence of monsoon flow (not shown). Clouds continued to develop west of
1220 about 10.5°W. CBL growth rates can be calculated for each dropsonde location were calculated
1221 based on the two dropsonde profiles. The SRL top and aged dust layer showed little change
1222 compared to the outward leg apart from at B4 where the SRL was consumed completely by the fast
1223 growing CBL and the low in the SRL top between 8.5 and 7.5°W remained (Figure 17). It should be
1224 noted that SEVIRI imagery did not show any dust presence along the flight tracks on this day
1225 suggesting that the LIDAR dust signal represents background dust levels – some dust is almost
1226 always present over North Africa at this time of year (Israelevich et al., 2003).

1227 As part of this SHL survey, the Falcon 20 took measurements at the same time as the BAe146 but on
1228 a more southern track (flight F24 in Figure 2b). The analysis of the combined aircraft data showed
1229 that the SHL had an elongated shape with a NE-SW orientation. Moisture from the monsoon inflow
1230 was transported around the SHL at low levels (not shown) in the morning. These unique
1231 measurements allow for the first time to challenge climate models in the SHL region and to
1232 understand the processes that control the observed temporal and spatial variability.

1233 4.4.2 First observations of the vertical profile of SABL fluxes and mesoscale circulations 1234 in the SABL

1235 The Saharan atmospheric boundary layer (SABL) is probably the deepest on Earth, commonly
1236 reaching 5-6 km, and is crucial in controlling the vertical redistribution and transport of dust,
1237 moisture, heat and momentum fluxes in the Sahara (Cuesta et al., 2009). Before Fennec, aircraft
1238 observations and radiosondes (Cuesta et al., 2009; Messager et al., 2010; Marsham et al., 2013b)
1239 had shown the persistence of a deep near-neutral Saharan residual layer (SRL) over large areas of
1240 the Sahara throughout the day, with only a very small temperature inversion separating the SRL
1241 from the CBL below. Flamant et al. (2007) and Messager et al. (2010) had shown that the SRL may
1242 have a maximum humidity mixing ratio at its upper levels, and that small errors in model
1243 representation of this humidity can have substantial consequences in terms of relative humidity,
1244 cloud cover and, therefore, radiation. This unusual structure of the SABL means that relatively small

1245 perturbations to CBL temperature (e.g. from a surface albedo anomaly) are expected to have
1246 significant impacts on vertical mixing and perhaps induce circulations that may affect the CBL in
1247 neighbouring regions. There was evidence of such effects in observations from the CBL (Marsham et
1248 al., 2008a) and in modeling studies (Birch et al., 2012; Huang et al., 2010), but observations of
1249 impacts on the SRL were lacking. Observations from Fennec BBM supersite 1 showed that when the
1250 CBL does reach 5 or 6 km this tends to only happen between 15 and 18 UTC (Marsham et al., 2013b).
1251 Fennec flights have provided new insights into the vertical structure of and mixing within the SABL
1252 (see Garcia-Carreras et al. (2015) including schematic (their Figure 14)) and SABL mesoscale
1253 circulations (below).

1254 During Fennec, aircraft LIDAR and in-situ observations were used to better understand the vertical
1255 stratification and transport mechanisms within the SABL, as well as its temporal and spatial
1256 variability (Garcia-Carreras et al., 2015). In order to sample the vertical turbulent structure of the
1257 SABL during Fennec, stacked legs were performed at different heights, determined from inspecting
1258 dropsonde profiles launched at both ends of the leg before descending. Each run was at least 10
1259 times the SABL depth (≥ 60 km) and took place between 13-15LT, when sensible heating was
1260 maximum. Heat fluxes were computed from the stacked legs, as well as the ascents and descents,
1261 taking advantage of the shallow angle of the aircraft profiles. These indicate that entrainment fluxes
1262 are very weak, as a result of detrainment at the CBL top. This is a result of the weak temperature
1263 inversion, and high vertical velocity of overshooting parcels, which are characteristic of the SABL,
1264 and can explain the slow development of the CBL despite the strong surface heating. LIDAR
1265 measurements from high-level runs also showed that the boundary layer depth can vary by up to
1266 100% over distances of a few kilometres due to turbulent processes alone, so that any given
1267 dropsonde profile may not be representative of the whole run.

1268 Figure 18 shows an example from a flight where small variations ~~from-in~~ heating from an albedo
1269 anomaly appear to be generating mesoscale circulations within the SABL. Figure 18 shows the
1270 vertical extinction coefficient at 532 nm retrieved with the LIDAR LNG on 20th June 2011 (1405-1446
1271 UTC, flight F21, see Figure 2b) from the Falcon flying southeastward in Mauritania, with water
1272 vapour mixing ratio (WVMR) and wind profiles from four dropsondes overplotted. The LIDAR
1273 transect highlights a number of BL processes of importance encountered during the Fennec
1274 campaign, showing variability from the turbulent to the synoptic scales, as described below.

1275 At the synoptic scale, there is a temperature and humidity gradient across the transect, with warmer
1276 and drier conditions in the northwest (by ~ 5 K and 7 g/kg), leading to a deeper CBL compared to the
1277 southeast (4 km at 24°N compared with 2 km at 21°N). The monsoon flow at night reached
1278 approximately 20°N along the flight-track, bringing in cool moist air into the southern end of the
1279 transect (from UK Met Office analysis, not shown), which was then redistributed vertically as the CBL
1280 grew during the day. The more spatially homogeneous residual layer, on the other hand, reflects the
1281 conditions from the day before; the monsoon front on the night of the 19th was considerably
1282 further south, leading to a deep CBL throughout the transect. Superimposed on the synoptic
1283 gradient there is substantial variability in the SABL depth and structure. Variability at the turbulent
1284 eddy scale can be observed in the northern end of the transect (24.5 - 25.2°N), with changes in the
1285 depth of the well-mixed aerosol layer (and so the CBL) of ~ 1 km over short horizontal distances (5-
1286 10 km, ~ 0.05 to 0.1°), consistent with idealised simulations and LIDAR measurements described in
1287 Garcia-Carreras et al. (2015).

1288 At the mesoscale, there is a region with cloud and deeper BLs at the boundary between the warm,
1289 dry conditions in the northwest, and the moister conditions in the southeast (21.4-22.5°N), with an
1290 orange plume reaching 6 km at 22.2N. Satellite imagery shows that the clouds observed by the
1291 LIDAR are part of a band of clouds coincident with a ~~dark-negative~~ albedo ~~feature-anomaly of around~~
1292 ~~0.2~~ that is ~~largely just west~~ off the flight-track at 21.6°N (red line, Figure 18). The surface hot-spot
1293 leads to a local increase in the CBL depth, cloud formation and an upward transport of dust. The
1294 impact of ~~another~~ smaller hot-spots can be observed at 22.8, ~~24 and 24.5~~°N. Easterly winds in the
1295 SRL in the southeast lead to the airmass overriding the deeper CBL in the northwest, potentially
1296 contributing to the cloud formation. The 3 gkg⁻¹ contour in Figure 18 has been drawn using the
1297 dropsonde data and the LIDAR-inferred aerosol distribution and suggests that the deeper CBL
1298 around 22°N acts to transport water vapour and dust directly to the top of the SRL, where it spreads
1299 laterally, capping the adjacent CBL and leading to weak maxima in water vapour mixing ratios at the
1300 top of the SRL in the three eastern dropsondes. This supports the hypotheses of Marsham et al.
1301 (2008a) and Messager et al. (2010) of mesoscale variability in the SABL and its role in the transport
1302 of CBL air into the RL, with implications for the long-range transport of dust.

1303 4.4.3 North American wildfire emissions measured over Africa

1304 Approximately 15 pollutant plumes were observed on the BAe146 in the upper troposphere (6 to 8.5
1305 km altitude) above the Sahara desert during the Fennec campaign in June 2011. Using HYSPLIT
1306 trajectory analysis and MODIS satellite fire products, four source regions were identified for these
1307 pollutant plumes: flaring from oil fields in Algeria and biomass burning in the southern USA,
1308 Venezuela and West Africa. The pollutant plumes displayed high concentrations of ozone and sub-
1309 micron particles, with differing characteristics from each source region. Values for the single
1310 scattering albedo ranged from 0.57 to 0.99 and for the Angstrom exponent from -0.85 to 2.44 for
1311 individual plumes. If the HYSPLIT trajectory calculations are robust (uncertain due the substantial
1312 errors identified in the vertical wind fields), it is believed this is the first aircraft measurement of
1313 flaring from oil fields and may require further research attention: this is planned in the forthcoming
1314 DACCIA field campaign in southern West Africa.

1315 5 Conclusions

1316 We have presented a description of the Fennec airborne ~~campaigns fieldwork~~ of 2011 and 2012 over
1317 the western Sahara. ~~We aim firstly to describe and document the aircraft instrumentation of the~~
1318 ~~BAe146 and the F20, flight locations, aims and the associated meteorology~~ in order to provide a
1319 reference and context for published and future articles. ~~Secondly, we have presented~~ new scientific
1320 results which have developed from the airborne measurements. ~~Data has been integrated from~~
1321 ~~different Fennec platforms (i.e. dual aircraft observations, ground based, and satellite based)~~ to
1322 show how the exploitation of aircraft measurements can deepen our understanding of weather,
1323 climate and dust processes over remote regions of the Sahara not otherwise accessible. Finally, the
1324 Fennec airborne data provide the only comprehensive resource of in-situ Saharan observations with
1325 which to develop the science linking dust, dynamics and radiation in the central Sahara. Along with
1326 the ground-based and satellite measurements, these will be heavily exploited in the coming years,
1327 and therefore we have provided a detailed overview of the data and its context.

1328 ~~The research areas and key findings of published articles relating to the Fennec aircraft observations~~
1329 ~~are summarized in Table 8. We emphasize the ~~Notable~~ instrumental developments on the BAe146~~

1330 to include measurement of giant mode dust particles with using Cloud Imaging Probes (up to 300 μ m
1331 during Fennec 2011 and 6200 μ m during Fennec 2012), and the advancement of technologies such
1332 that size distribution measurements across the full size range at 10Hz were possible (Rosenberg et
1333 al., 2012). The former has been used to demonstrate a significant presence of particles sized larger
1334 than 10 μ m over remote parts of the Sahara, with volume distributions peaking between 10 to
1335 60 μ m in many fresh, heavy dust cases while the, with peak volume distribution shifting to 10 to
1336 20 μ m and reduced total concentrations in more aged dust events with a reduction in total
1337 concentrations (Ryder et al., 2013a, b). The measurement of size distributions at 10Hz has allowed
1338 dust fluxes in the SABL to be measured from an aircraft for the first time (Rosenberg et al., 2014).
1339 Measurement of size distributions both on the BAe146 wing probes and behind various inlets has
1340 demonstrated the effects of the BAe146 Rosemount inlets on size distributions and optical
1341 properties for the first time.

1342 The new scientific findings presented in this article are as follows:

1343 The airborne campaign took place in three parts: the pilot study (4–9 April 2011, BAe146 only),
1344 Fennec IOP 2011 (2–28 June 2011, BAe146 and F20), and Fennec IOP 2012 (1–19 June 2012, BAe146
1345 only). During the IOPs both aircraft were based in the Canary Islands and conducted a total of 48
1346 flights over Mauritania, Mali and Senegal. Missions covered various scientific objectives, from
1347 measurement of in situ dust properties, SABL development and processes, haboob structure and
1348 development, and SHL spatial structure and temporal evolution. Aircraft instrumentation was
1349 coordinated so that the F20 made remote sensing measurements, covering large areas at high
1350 altitudes, while the BAe146 made both comprehensive meteorological and aerosol in situ
1351 measurements and remote sensing measurements, geared to study a smaller region of interest
1352 more thoroughly at a range of altitudes. In several instances, the BAe146 and F20 flew coordinated
1353 missions to exploit the strengths of both aircraft.

1354 Meteorological events during the three airborne campaigns have been described. June 2011 was
1355 dominated by a switch from an initial 'maritime phase' to a 'heat low' phase by the end of the
1356 month, congruent with movement of the SHL from a heat low east to west phase. The stronger
1357 easterlies and enhanced MCS activity during the latter phase resulted in more dust events over the
1358 Fennec flight region. June 2012 displayed a more stationary SHL, with strong northeasterlies
1359 favourable to dust emission. Lagrangian backward calculations have been used to determine dust
1360 uplift locations during each campaign and show, for example, that d

1361 • During the second half of June 2011 sources over central Algeria dominated, driven primarily
1362 by stronger easterlies associated with the westward movement of the SHL, in contrast to the
1363 second half of June 2012 when more Sahelian dust sources dominated due to a northern
1364 extension of the monsoon flow and increased MCS and cold pool activity over Mali. This can
1365 be linked to is associated with differences in the chemical composition and optical property
1366 results between campaigns, which show higher dust absorption and lower calcium content
1367 in 2012 compared to 2011, characteristic of dust emitted from Sahelian soils. This change in
1368 composition and associated dust absorption can have significant radiative impacts which can
1369 be driven by dust uplift locations and the dominant meteorology. These first results of dust
1370 chemical composition in the SHL region indicate the importance of large scale meteorology
1371 in affecting dust composition and therefore radiative properties.

1372 • Dust uplift under the breakdown of the nocturnal LLJ has been observed, demonstrating the
1373 presence of coarse and giant particles in these very fresh dust events, which are observed at
1374 low altitudes and often before they become visible in SEVIRI imagery. The comprehensive
1375 aerosol and cloud instrumentation on the BAe146 has been used to explore the interaction
1376 between dust layers and clouds, indicating that dust particles are likely to be acting as CCN
1377 and also as IN at temperatures of -15°C.

1378 • Ozone concentrations have been compared to size distribution measurements of surface
1379 area in an attempt to determine the role of dust on ozone depletion. Results suggest that
1380 coarser, fresher dust particles can provide a route to decrease ozone concentrations, though
1381 in this case a change of air mass during sampling prevented unequivocal attribution.

1382 • Dust uplift under the breakdown of the nocturnal LLJ has been observed, including its impact
1383 on shortwave irradiance and the presence of coarse and giant particles in these very fresh
1384 dust events, which are observed at low altitudes and often before they become visible in
1385 SEVIRI imagery.

1386 • F20 LIDAR measurements have been combined with BAe146 in-situ extinction and vertically
1387 resolved shortwave flux measurements to help describe the influence of a haboob
1388 thermodynamics on the development of the SABL, and subsequent mixing of the haboob
1389 through the SABL.

1390 • Combined LIDAR and dropsonde observations show the spatial and diurnal structure of the
1391 SHL. The CBL develops throughout the day while the influence of the southerly monsoon
1392 flow restricts this growth. Variability in the SABL plays an important role in the transport of
1393 CBL air into the SRL, which has implications for long range transport of dust, with evidence
1394 of surface albedo features driving such variability.

1395 • Vertical profiles of turbulent fluxes have revealed unusual characteristics of entrainment and
1396 detrainment of thermals in the deep, dry SABL, which are a challenge for BL schemes in
1397 global models.

1398 • Unique in-situ observations suggest that precipitation is recycled as it is evaporated into BL
1399 air that feeds clouds (a common feature of the SABL), increasing the total water content of
1400 subsequent clouds and increasing the moisture content at mid-levels in the SABL.
1401 Observations suggest cloud-processing of dust and subsequent evaporation alters the size
1402 distribution of dust.

1403 • In one case, a comparison of aircraft LIDAR data with satellite measurements from SEVIRI
1404 and MODIS show good agreement as to the spatial distribution of dust but disagree as to the
1405 loading, which may be indicative of different sensitivities to varying meteorological
1406 conditions. Further detailed comparisons have taken place (see Table 8), demonstrating the
1407 value of aircraft-satellite validation studies.

1408 • For a flight over the Zouerate supersite, a comparison of column mean size distributions
1409 between AERONET and the BAe146 in-situ measurements shows AERONET retrieved peak
1410 volume size distributions at 3-6 microns, while aircraft measurements measured more
1411 coarse mode, with a peak at 12 microns. This was in a dust event with low concentrations of
1412 coarse and giant particles present – the aircraft frequently encountered cases with a greater
1413 coarse mode present. This provides further evidence that AERONET derived size
1414 distributions should be used with caution when coarse particles are present, and merits
1415 further detailed comparisons under heavy dust loadings.

1416 Notable instrumental developments on the BAe146 include measurement of giant mode dust
1417 particles with Cloud Imaging Probes (up to 300 μm during Fennec 2011 and 6200 μm during Fennec
1418 2012), and size distribution measurements across the full size range at 10Hz. The former has been
1419 used to demonstrate a significant presence of particles sized larger than 10 μm over remote parts of
1420 the Sahara, with volume distributions peaking between 10 to 60 μm in many fresh, heavy dust cases,
1421 with peak volume distribution shifting to 10 to 20 μm and reduced total concentrations in more aged
1422 dust events. Dust uplift under the breakdown of the nocturnal LLJ has been observed, demonstrating
1423 the presence of coarse and giant particles in these very fresh dust events, which are observed at low
1424 altitudes and often before they become visible in SEVIRI imagery. The comprehensive aerosol and
1425 cloud instrumentation on the BAe146 has been used to explore the interaction between dust layers
1426 and clouds, indicating that dust particles are likely to be acting as CCN and also as IN at temperatures
1427 of -15°C. The measurement of size distributions at 10Hz has allowed dust fluxes in the SABL to be
1428 measured from an aircraft for the first time. Measurement of size distributions both on the BAe146
1429 wing probes and behind various inlets has demonstrated the effects of the BAe146 Rosemount inlets
1430 on size distributions and optical properties for the first time. Ozone concentrations have been
1431 compared to size distribution measurements of surface area in an attempt to determine the role of
1432 dust on ozone depletion. Results suggest that coarser, fresher dust particles can provide a route to
1433 decrease ozone concentrations, though in this case a change of air mass during sampling prevented
1434 unequivocal attribution.

1435 In-situ, LIDAR and dropsonde measurements using either one or both aircraft have been used to
1436 study atmospheric processes occurring in the SABL and SHL. Vertical profiles of turbulent fluxes have
1437 revealed the unusual characteristics of entrainment and detrainment of thermals in this deep dry BL,
1438 a challenge for BL schemes in global models. F20 LIDAR measurements have been combined with
1439 BAe146 in-situ extinction and vertically resolved shortwave flux measurements to help describe the
1440 influence of a haboob thermodynamics on the development of the SABL, and subsequent mixing of
1441 the haboob through the SABL. Several flights using LIDAR and dropsondes from both aircraft have
1442 been used to examine the spatial and diurnal structure of the SHL, showing the development of the
1443 CBL throughout the day and the influence of the southerly monsoon flow on restricting this growth.
1444 A case study has exploited LIDAR and water vapor mixing ratios from dropsondes to demonstrate
1445 how variability in the SABL plays an important role in the transport of CBL air into the SRL, which has
1446 implications for long range transport of dust, with evidence of surface albedo features driving such
1447 variability. Finally, clouds are a common feature of the SABL (Stein et al., 2011) and unique in-situ
1448 observations from such clouds suggest that precipitation is recycled as it is evaporated into BL air
1449 that feeds the clouds, increasing the total water content of subsequent clouds and increasing the
1450 moisture content at mid-levels in the SABL.

1451 Ground-based and satellite measurements have been combined with aircraft measurements to
1452 provide insights into dust properties and radiative effects. In one case, a comparison of aircraft
1453 LIDAR data with satellite measurements from SEVIRI and MODIS show good initial agreement and
1454 have allowed further detailed comparisons to take place, demonstrating the value of aircraft-
1455 satellite validation studies. For a flight over the Zouerate supersite, a comparison of column mean
1456 size distributions between AERONET and the BAe146 in-situ measurements shows AERONET
1457 retrieved peak volume size distributions at 3.6 microns, while aircraft measurements measured
1458 more coarse mode, with a peak at 12 microns. This was in a dust event with low concentrations of
1459 coarse and giant particles present – the aircraft frequently encountered cases with a greater coarse

1460 mode present. This provides further evidence that AERONET derived size distributions should be
1461 used with caution when coarse particles are present, and merits further detailed comparisons under
1462 heavy dust loadings. Finally we have presented results showing the impact of dust on shortwave
1463 radiative fluxes concurrent firstly with active dust uplift by a LLJ under a very shallow but thick layer
1464 of dust which was not visible in SEVIRI satellite imagery, and secondly for dust under a haboob
1465 generated by convection over the Atlas mountains transported over Mauritania.

1466 We believe that this paper demonstrates that the Fennec airborne campaign has delivered a novel,
1467 rich dataset through the operation of two aircraft over remote regions of the Sahara. Particular
1468 highlights are the ability of the BAe146 to measure coarse and giant mode dust particles, sized up to
1469 300 μm (Fennec 2011) and 6200 μm (Fennec 2012) for the first time on an aircraft, and the
1470 combined application of LIDAR, dropsonde and in situ measurements to improve understanding of
1471 processes occurring in the SABL – the deepest boundary layer on Earth. The power of these aircraft
1472 measurements will be enhanced via combination with the ground-based measurements available
1473 from the Fennec climate program, providing a unique resource for further in-depth study of the vital
1474 SHL region of the Sahara. These will be further exploited through the Fennec Earth observation and
1475 modelling programs.

1476 **Acknowledgements**

1477 Core project funding for Fennec was from the UK Natural Environmental Research Council (NERC)
1478 under grant NE/G017166/1. In addition, it received support from the NERC National Centre for
1479 Atmospheric Science (NCAS), the Agence Nationale de la Recherche (ANR n°2010 BLAN 606 01), the
1480 Institut National des Sciences de l'Univers (INSU/CNRS) through the LEFE program, the Centre
1481 National d'Etudes Spatiales (CNES) through the TOSCA program and Météo-France. Airborne data
1482 from the BAe146 was obtained using the BAe-146-301 Atmospheric Research Aircraft operated by
1483 Directflight Ltd and managed by FAAM, which is a joint entity of the NERC and the UK Met Office.
1484 Airborne data from the F20 was obtained using the Falcon 20 Environment Research Aircraft
1485 operated and managed by SAFIRE, which is a joint entity of CNRS, Météo-France & CNES. EUFAR
1486 (EUropean Facility for Airborne Research) is acknowledged for its support to the RAIN4DUST Falcon-
1487 20 flights and LADUNEX BAe146 flights. The UK Met Office is acknowledged for funding of flight b710
1488 through SAVEX. SAVEX ground deployment at Tenerife was possible thanks to RIMA/AERONET and
1489 AEMET infrastructure; and support from Juan de la Cierva (JCI-2009-04455), Universidad de La
1490 Laguna (2012/0001624), MICIIN ([CGL2011-24290](#)[CGL2012-33294](#)) and Generalitat Valenciana
1491 (PROMETEO/2010/064) projects. Many other scientists and engineers were involved in the gathering
1492 of this outstanding dataset. Additional partners include: Directflight, AvalonAero, FAAM (Facility for
1493 Airborne Atmospheric Measurements), SAFIRE (Service des Avions Français Instrumentés pour la
1494 Recherche en Environnement), UK Met Office, and DMN Maroc. MODIS data used in this paper were
1495 produced with the LAADS online data system, developed and maintained by NASA Goddard, and we
1496 also acknowledge the MODIS scientists and associated NASA personnel for the production of the
1497 data used in this research effort. Flight forecasting would not have been possible without the model
1498 products made available especially for the Fennec project particularly the UK Met Office, the Météo-
1499 France AROME model team and the DREAM model team.

1500

1501 **References**

1502 Allen, C. J. T., Washington, R., and Engelstaedter, S.: Dust emission and transport mechanisms in
1503 the central Sahara: Fennec ground-based observations from Bordj Badji Mokhtar, June 2011, *J
1504 Geophys Res-Atmos*, 118, 6212-6232, Doi 10.1002/Jgrd.50534, 2013.

1505 Ansmann, A., Tesche, M., Althausen, D., Muller, D., Seifert, P., Freudenthaler, V., Heese, B.,
1506 Wiegner, M., Pisani, G., Knippertz, P., and Dubovik, O.: Influence of Saharan dust on cloud glaciation
1507 in southern Morocco during the Saharan Mineral Dust Experiment, *J Geophys Res-Atmos*, 113, Doi
1508 10.1029/2007jd008785, 2008.

1509 Ansmann, A., Tesche, M., Knippertz, P., Bierwirth, E., Althausen, D., Muller, D., and Schulz, O.:
1510 Vertical profiling of convective dust plumes in southern Morocco during SAMUM, *Tellus B*, 61, 340-
1511 353, DOI 10.1111/j.1600-0889.2008.00384.x, 2009.

1512 Ansmann, A., Petzold, A., Kandler, K., Tegen, I., Wendisch, M., Müller, D., Weinzierl, B., Müller, T.,
1513 and Heintzenberg, J.: Saharan Mineral Dust Experiments SAMUM-1 and SAMUM-2: what have we
1514 learned?, *Tellus B*, 63, 403-429, DOI 10.1111/j.1600-0889.2011.00555.x, 2011.

1515 Banks, J. R., and Brindley, H. E.: Evaluation of MSG-SEVIRI mineral dust retrieval products over
1516 Africa and the Middle East, *Remote Sensing of Environment*, 128, 58-73,
1517 doi:10.1016/j.rse.2012.07.017, 2013.

1518 Banks, J. R., Brindley, H. E., Flamant, C., Garay, M. J., Hsu, N. C., Kalashnikov, O. V., Kluser, L., and
1519 Sayer, A. M.: Intercomparison of satellite dust retrieval products over the west African Sahara during
1520 the Fennec campaign in June 2011, *Remote Sensing of Environment*, 136, 99-116, DOI
1521 10.1016/j.rse.2013.05.003, 2013.

1522 Bauer, S. E., Balkanski, Y., Schulz, M., Hauglustaine, D. A., and Dentener, F.: Global modeling of
1523 heterogeneous chemistry on mineral aerosol surfaces: Influence on tropospheric ozone chemistry
1524 and comparison to observations, *J Geophys Res-Atmos*, 109, doi 10.1029/2003jd003868, 2004.

1525 Baumgardner, D., Jonsson, H., Dawson, W., O'Connor, D., and Newton, R.: The cloud, aerosol and
1526 precipitation spectrometer: a new instrument for cloud investigations, *Atmos Res*, 59, 251-264,
1527 2001.

1528 Bian, H. S., and Zender, C. S.: Mineral dust and global tropospheric chemistry: Relative roles of
1529 photolysis and heterogeneous uptake, *J Geophys Res-Atmos*, 108, Doi 10.1029/2002jd003143, 2003.

1530 Birch, C. E., Parker, D. J., Marsham, J. H., and Devine, G. M.: The effect of orography and surface
1531 albedo on stratification in the summertime Saharan boundary layer: Dynamics and implications for
1532 dust transport, *J Geophys Res-Atmos*, 117, Doi:10.1029/2011jd015965, 2012.

1533 Blyth, A. M., Cooper, W. A., and Jensen, J. B.: A Study of the Source of Entrained Air in Montana
1534 Cumuli, *J Atmos Sci*, 45, 3944-3964, Doi 10.1175/1520-0469(1988)045<3944:Asotso>2.0.Co;2, 1988.

1535 Brindley, H., Knippertz, P., Ryder, C., and Ashpole, I.: A critical evaluation of the ability of the
1536 Spinning Enhanced Visible and Infrared Imager (SEVIRI) thermal infrared red-green-blue rendering to
1537 identify dust events: Theoretical analysis, *J Geophys Res-Atmos*, 117, Doi 10.1029/2011jd017326,
1538 2012.

1539 Brindley, H. E., and Russell, J. E.: An assessment of Saharan dust loading and the corresponding
1540 cloud-free longwave direct radiative effect from geostationary satellite observations, *J Geophys Res-*
1541 *Atmos*, 114, Doi 10.1029/2008jd011635, 2009.

1542 Brooke, J. K.: *Airborne Observations of the Physical and Optical Properties of Atmospheric*
1543 *Aerosol*, PhD, School of Earth and Environment, University of Leeds, 2014.

1544 Campanelli, M., Estelles, V., Smyth, T., Tomasi, C., Martinez-Lozano, M. P., Claxton, B., Muller, P.,
1545 Pappalardo, G., Pietruczuk, A., Shanklin, J., Colwell, S., Wrench, C., Lupi, A., Mazzola, M., Lanconelli,
1546 C., Vitale, V., Congeduti, F., Dionisi, D., Cardillo, F., Cacciani, M., Casasanta, G., and Nakajima, T.:
1547 Monitoring of Eyjafjallajokull volcanic aerosol by the new European Skynet Radiometers (ESR)
1548 network, *Atmos Environ*, 48, 33-45, DOI 10.1016/j.atmosenv.2011.09.070, 2012.

1549 Caquineau, S., Magonthier, M. C., Gaudichet, A., and Gomes, L.: An improved procedure for the
1550 X-ray diffraction analysis of low-mass atmospheric dust samples, *Eur J Mineral*, 9, 157-166, 1997.

1551 Caquineau, S., Gaudichet, A., Gomes, L., and Legrand, M.: Mineralogy of Saharan dust
1552 transported over northwestern tropical Atlantic Ocean in relation to source regions, *J Geophys Res-*
1553 *Atmos*, 107, Doi 10.1029/2000jd000247, 2002.

1554 Chang, R. Y. W., Sullivan, R. C., and Abbatt, J. P. D.: Initial uptake of ozone on Saharan dust at
1555 atmospheric relative humidities, *Geophys Res Lett*, 32, 10.1029/2005gl023317, 2005.

1556 Chauvin, F., Roehrig, R., and Lafore, J. P.: Intraseasonal Variability of the Saharan Heat Low and Its
1557 Link with Midlatitudes, *J Climate*, 23, 2544-2561, Doi 10.1175/2010jcli3093.1, 2010.

1558 Chazette, P., Dabas, A., Sanak, J., Lardier, M., and Royer, P.: French airborne lidar measurements
1559 for Eyjafjallajokull ash plume survey, *Atmos Chem Phys*, 12, 7059-7072, DOI 10.5194/acp-12-7059-
1560 2012, 2012.

1561 Cuesta, J., Edouart, D., Mimouni, M., Flamant, P. H., Loth, C., Gibert, F., Marnas, F., Bouklila, A.,
1562 Kharef, M., Ouchene, B., Kadi, M., and Flamant, C.: Multiplatform observations of the seasonal
1563 evolution of the Saharan atmospheric boundary layer in Tamanrasset, Algeria, in the framework of
1564 the African Monsoon Multidisciplinary Analysis field campaign conducted in 2006, *J Geophys Res-*
1565 *Atmos*, 113, 10.1029/2007jd009417, 2008.

1566 Cuesta, J., Marsham, J. H., Parker, D. J., and Flamant, C.: Dynamical mechanisms controlling the
1567 vertical redistribution of dust and the thermodynamic structure of the West Saharan atmospheric
1568 boundary layer during summer, *Atmos Sci Lett*, 10, 34-42, Doi 10.1002/Asl.207, 2009.

1569 de Reus, M., Dentener, F., Thomas, A., Borrman, S., Strom, J., and Lelieveld, J.: Airborne
1570 observations of dust aerosol over the North Atlantic Ocean during ACE 2: Indications for
1571 heterogeneous ozone destruction, *J Geophys Res-Atmos*, 105, 15263-15275, Doi
1572 10.1029/2000jd900164, 2000.

1573 de Reus, M., Fischer, H., Sander, R., Gros, V., Kormann, R., Salisbury, G., Van Dingenen, R.,
1574 Williams, J., Zollner, M., and Lelieveld, J.: Observations and model calculations of trace gas
1575 scavenging in a dense Saharan dust plume during MINATROC, *Atmos Chem Phys*, 5, 1787-1803,
1576 2005.

1577 de Villiers, R. A., Ancellet, G., Pelon, J., Quennehen, B., Schwarzenboeck, A., Gayet, J. F., and Law,
1578 K. S.: Airborne measurements of aerosol optical properties related to early spring transport of mid-

1579 latitude sources into the Arctic, *Atmos Chem Phys*, 10, 5011-5030, DOI 10.5194/acp-10-5011-2010,
1580 2010.

1581 Dubovik, O., and King, M. D.: A flexible inversion algorithm for retrieval of aerosol optical
1582 properties from Sun and sky radiance measurements, *J Geophys Res-Atmos*, 105, 20673-20696, Doi
1583 10.1029/2000jd900282, 2000.

1584 Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M., Yang, P., Eck, T. F., Volten,
1585 H., Munoz, O., Veihelmann, B., van der Zande, W. J., Leon, J. F., Sorokin, M., and Slutsker, I.:
1586 Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of
1587 desert dust, *J Geophys Res-Atmos*, 111, Doi 10.1029/2005jd006619, 2006.

1588 Dunion, J. P., and Velden, C. S.: The impact of the Saharan air layer on Atlantic tropical cyclone
1589 activity, *B Am Meteorol Soc*, 85, 353-+, Doi 10.1175/Bams-85-3-353, 2004.

1590 Engelstaedter, S., Tegen, I., and Washington, R.: North African dust emissions and transport,
1591 *Earth-Sci Rev*, 79, 73-100, DOI 10.1016/j.earscirev.2006.06.004, 2006.

1592 Engelstaedter, S., Washington, R., Flamant, C., Parker, D. J., Allen, C. J. T., and Todd, M. C.: The
1593 Saharan heat low and moisture transport pathways in the central Sahara – multi-aircraft
1594 observations and Africa-LAM evaluation, *J Geophys Res-Atmos*, 10.1002/2015JD023123, 2015.

1595 Estellés, V., Campanelli, M., Smyth, T. J., Utrillas, M. P., and Martinez-Lozano, J. A.: Evaluation of
1596 the new ESR network software for the retrieval of direct sun products from CIMEL CE318 and PREDE
1597 POM01 sun-sky radiometers, *Atmos. Chem. Phys.*, 12, 11619–11630, doi:10.5194/acp-12-11619-
1598 2012, 2012a.

1599 Estellés, V., Campanelli, M., Utrillas, M. P., Exposito, F., and Martinez-Lozano, J. A.: Comparison of
1600 AERONET and SKYRAD4.2 inversion products retrieved from a Cimel CE318 sunphotometer, *Atmos*
1601 *Meas Tech*, 5, 569-579, DOI 10.5194/amt-5-569-2012, 2012b.

1602 Evan, A. T., Vimont, D. J., Heidinger, A. K., Kossin, J. P., and Bennartz, R.: The Role of Aerosols in
1603 the Evolution of Tropical North Atlantic Ocean Temperature Anomalies, *Science*, 324, 778-781, DOI
1604 10.1126/science.1167404, 2009.

1605 Evan, A. T., Foltz, G. R., Zhang, D. X., and Vimont, D. J.: Influence of African dust on ocean-
1606 atmosphere variability in the tropical Atlantic, *Nat Geosci*, 4, 762-765, Doi 10.1038/Ngeo1276, 2011.

1607 Evan, A. T., Flamant, C., Fiedler, S., and Doherty, S.: An analysis of aeolian dust in climate models,
1608 *Geophys Res Lett*, doi:10.1002/2014GL060545, 2014.

1609 Flamant, C., Chaboureau, J. P., Parker, D. J., Taylor, C. A., Cammas, J. P., Bock, O., Timouk, F., and
1610 Pelon, J.: Airborne observations of the impact of a convective system on the planetary boundary
1611 layer thermodynamics and aerosol distribution in the inter-tropical discontinuity region of the West
1612 African Monsoon, *Q J Roy Meteor Soc*, 133, 1175-1189, Doi 10.1002/qj.97, 2007.

1613 Formenti, P., Rajot, J. L., Desboeufs, K., Said, F., Grand, N., Chevaillier, S., and Schmechtig, C.:
1614 Airborne observations of mineral dust over western Africa in the summer Monsoon season: spatial
1615 and vertical variability of physico-chemical and optical properties, *Atmos Chem Phys*, 11, 6387-6410,
1616 DOI 10.5194/acp-11-6387-2011, 2011a.

1617 Formenti, P., Schütz, L., Balkanski, Y., Desboeufs, K., Ebert, M., Kandler, K., Petzold, A., Scheuvens,
1618 D., Weinbruch, S., and Zhang, D.: Recent progress in understanding physical and chemical properties

1619 of African and Asian mineral dust, *Atmos Chem Phys*, 11, 8231-8256, DOI 10.5194/acp-11-8231-
1620 2011, 2011b.

1621 Formenti, P., Caquineau, S., Desboeufs, K., Klaver, A., Chevaillier, S., Journet, E., and Rajot, J. L.:
1622 Mapping the physico-chemical properties of mineral dust in western Africa: mineralogical
1623 composition, *Atmos. Chem. Phys.*, 14, 10663-10686, doi:10.5194/acp-14-10663-2014, 2014.

1624 Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D. W., Haywood, J. M., Lean,
1625 J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in
1626 Atmospheric Constituents and in Radiative Forcing, in: Climate Change 2007: The Physical Science
1627 Basis, in: Contribution of Working Group I to the Fourth Assessment Report of the
1628 Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z.,
1629 Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK,
1630 2007.

1631 Garcia-Carreras, L., Marsham, J. H., Parker, D. J., Bain, C. L., Milton, S., Saci, A., Salah-Ferroudj, M.,
1632 Ouchene, B., and Washington, R.: The impact of convective cold pool outflows on model biases in
1633 the Sahara, *Geophys Res Lett*, 40, 1647-1652, Doi 10.1002/GrL.50239, 2013.

1634 Garcia-Carreras, L., Parker, D. J., Marsham, J. H., Rosenberg, P. D., Brooks, I. M., Lock, A. P.,
1635 Marenco, F., McQuaid, J. B., and Hobby, M.: The Turbulent Structure and Diurnal Growth of the
1636 Saharan Atmospheric Boundary Layer, *J Atmos Sci*, 72, 693-713, 10.1175/Jas-D-13-0384.1, 2015.

1637 Garcia, O. E., Diaz, J. P., Exposito, F. J., Diaz, A. M., Dubovik, O., Derimian, Y., Dubuisson, P., and
1638 Roger, J. C.: Shortwave radiative forcing and efficiency of key aerosol types using AERONET data,
1639 *Atmos Chem Phys*, 12, 5129-5145, DOI 10.5194/acp-12-5129-2012, 2012.

1640 Grassian, V. H.: Chemical reactions of nitrogen oxides on the surface of oxide, carbonate, soot,
1641 and mineral dust particles: Implications for the chemical balance of the troposphere, *J Phys Chem A*,
1642 106, 860-877, Doi 10.1021/Jp012139h, 2002.

1643 Gross, S., Tesche, M., Freudenthaler, V., Toledano, C., Wiegner, M., Ansmann, A., Althausen, D.,
1644 and Seefeldner, M.: Characterization of Saharan dust, marine aerosols and mixtures of biomass-
1645 burning aerosols and dust by means of multi-wavelength depolarization and Raman lidar
1646 measurements during SAMUM 2, *Tellus B*, 63, 706-724, DOI 10.1111/j.1600-0889.2011.00556.x,
1647 2011.

1648 Hanisch, F., and Crowley, J. N.: Ozone decomposition on Saharan dust: an experimental
1649 investigation, *Atmos Chem Phys*, 3, 119-130, 2003.

1650 Hashimoto, M., Nakajima, T., Dubovik, O., Campanelli, M., Che, H., Khatri, P., Takamura, T., and
1651 Pandithurai, G.: Development of a new data-processing method for SKYNET sky radiometer
1652 observations, *Atmos Meas Tech*, 5, 2723-2737, DOI 10.5194/amt-5-2723-2012, 2012.

1653 Haywood, J. M., Pelon, J., Formenti, P., Bharmal, N., Brooks, M., Capes, G., Chazette, P., Chou, C.,
1654 Christopher, S., Coe, H., Cuesta, J., Derimian, Y., Desboeufs, K., Greed, G., Harrison, M., Heese, B.,
1655 Highwood, E. J., Johnson, B., Mallet, M., Marticorena, B., Marsham, J. H., Milton, S., Myhre, G.,
1656 Osborne, S. R., Parker, D. J., Rajot, J. L., Schulz, M., Slingo, A., Tanre, D., and Tulet, P.: Overview of
1657 the Dust and Biomass-burning Experiment and African Monsoon Multidisciplinary Analysis Special
1658 Observing Period-0, *J Geophys Res-Atmos*, 113, Doi 10.1029/2008jd010077, 2008.

1659 Haywood, J. M., Johnson, B. T., Osborne, S. R., Baran, A. J., Brooks, M., Milton, S. F., Mulcahy, J.,
1660 Walters, D., Allan, R. P., Klaver, A., Formenti, P., Brindley, H. E., Christopher, S., and Gupta, P.:

1661 Motivation, rationale and key results from the GERBILS Saharan dust measurement campaign, Q J
1662 Roy Meteor Soc, 137, 1106-1116, Doi 10.1002/Qj.797, 2011a.

1663 Haywood, J. M., Johnson, B. T., Osborne, S. R., Mulcahy, J., Brooks, M. E., Harrison, M. A. J.,
1664 Milton, S. F., and Brindley, H. E.: Observations and modelling of the solar and terrestrial radiative
1665 effects of Saharan dust: a radiative closure case-study over oceans during the GERBILS campaign, Q J
1666 Roy Meteor Soc, 137, 1211-1226, Doi 10.1002/Qj.770, 2011b.

1667 Heintzenberg, J.: The SAMUM-1 experiment over Southern Morocco: overview and introduction,
1668 Tellus B, 61, 2-11, DOI 10.1111/j.1600-0889.2008.00403.x, 2009.

1669 Highwood, E. J., Northway, M. J., McMeeking, G. R., Morgan, W. T., Liu, D., Osborne, S., Bower, K.,
1670 Coe, H., Ryder, C., and Williams, P.: Aerosol scattering and absorption during the EUCAARI-LONGREX
1671 flights of the Facility for Airborne Atmospheric Measurements (FAAM) BAe-146: can measurements
1672 and models agree?, Atmos Chem Phys, 12, 7251-7267, DOI 10.5194/acp-12-7251-2012, 2012.

1673 Hobby, M., Gascoyne, M., Marsham, J. H., Bart, M., Allen, C., Engelstaedter, S., Fadel, D. M.,
1674 Gandega, A., Lane, R., McQuaid, J. B., Ouchene, B., Ouladichir, A., Parker, D. J., Rosenberg, P.,
1675 Ferroudj, M. S., Saci, A., Seddik, F., Todd, M., Walker, D., and Washington, R.: The Fennec Automatic
1676 Weather Station (AWS) Network: Monitoring the Saharan Climate System, J Atmos Ocean Tech, 30,
1677 709-724, Doi 10.1175/Jtech-D-12-00037.1, 2013.

1678 Hsu, N. C., Tsay, S. C., King, M. D., and Herman, J. R.: Aerosol properties over bright-reflecting
1679 source regions, Ieee T Geosci Remote, 42, 557-569, Doi 10.1109/Tgrs.2004.824067, 2004.

1680 Huang, Q., Marsham, J. H., Parker, D. J., Tian, W. S., and Grams, C. M.: Simulations of the effects
1681 of surface heat flux anomalies on stratification, convective growth, and vertical transport within the
1682 Saharan boundary layer, J Geophys Res-Atmos, 115, Doi 10.1029/2009jd012689, 2010.

1683 Huneeus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher, O.,
1684 Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A., Horowitz, L.,
1685 Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J. J., Myhre, G.,
1686 Penner, J., Perlitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model intercomparison
1687 in AeroCom phase I, Atmos Chem Phys, 11, 7781-7816, DOI 10.5194/acp-11-7781-2011, 2011.

1688 Israelevich, P. L., Ganor, E., Levin, Z., and Joseph, J. H.: Annual variations of physical properties of
1689 desert dust over Israel, J Geophys Res-Atmos, 108, 10.1029/2002jd003163, 2003.

1690 Jenkins, G. S., Pratt, A. S., and Heymsfield, A.: Possible linkages between Saharan dust and
1691 tropical cyclone rain band invigoration in the eastern Atlantic during NAMMA-06, Geophys Res Lett,
1692 35, Doi:10.1029/2008gl034072, 2008.

1693 Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P.
1694 W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N.,
1695 Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert
1696 dust, ocean biogeochemistry, and climate, Science, 308, 67-71, 2005.

1697 Johnson, B., Turnbull, K., Brown, P., Burgess, R., Dorsey, J., Baran, A. J., Webster, H., Haywood, J.,
1698 Cotton, R., Ulanowski, Z., Hesse, E., Woolley, A., and Rosenberg, P.: In situ observations of volcanic
1699 ash clouds from the FAAM aircraft during the eruption of Eyjafjallajokull in 2010, J Geophys Res-
1700 Atmos, 117, Doi 10.1029/2011jd016760, 2012.

1701 Journet, E., Balkanski, Y., and Harrison, S. P.: A new data set of soil mineralogy for dust-cycle
1702 modeling, *Atmos Chem Phys*, 14, 3801-3816, DOI 10.5194/acp-14-3801-2014, 2014.

1703 Kandler, K., Benker, N., Bundke, U., Cuevas, E., Ebert, M., Knippertz, P., Rodriguez, S., Schutz, L.,
1704 and Weinbruch, S.: Chemical composition and complex refractive index of Saharan Mineral Dust at
1705 Izana, Tenerife (Spain) derived by electron microscopy, *Atmos Environ*, 41, 8058-8074, DOI
1706 10.1016/j.atmosenv.2007.06.047, 2007.

1707 Kanitz, T., Seifert, P., Ansmann, A., Engelmann, R., Althausen, D., Casiccia, C., and Rohwer, E. G.:
1708 Contrasting the impact of aerosols at northern and southern midlatitudes on heterogeneous ice
1709 formation, *Geophys Res Lett*, 38, Doi 10.1029/2011gl048532, 2011.

1710 Kim, D., Chin, M., Yu, H. B., Diehl, T., Tan, Q., Kahn, R. A., Tsigaridis, K., Bauer, S. E., Takemura, T.,
1711 Pozzoli, L., Bellouin, N., Schulz, M., Peyridieu, S., Chedin, A., and Koffi, B.: Sources, sinks, and
1712 transatlantic transport of North African dust aerosol: A multimodel analysis and comparison with
1713 remote sensing data, *J Geophys Res-Atmos*, 119, 6259-6277, Doi 10.1002/2013jd021099, 2014.

1714 Kinne, S., Lohmann, U., Feichter, J., Schulz, M., Timmreck, C., Ghan, S., Easter, R., Chin, M.,
1715 Ginoux, P., Takemura, T., Tegen, I., Koch, D., Herzog, M., Penner, J., Pitari, G., Holben, B., Eck, T.,
1716 Smirnov, A., Dubovik, O., Slutsker, I., Tanre, D., Torres, O., Mishchenko, M., Geogdzhayev, I., Chu, D.,
1717 A., and Kaufman, Y.: Monthly averages of aerosol properties: A global comparison among models,
1718 satellite data, and AERONET ground data, *J Geophys Res-Atmos*, 108, Doi 10.1029/2001jd001253,
1719 2003.

1720 Knippertz, P., Deutscher, C., Kandler, K., Muller, T., Schulz, O., and Schutz, L.: Dust mobilization
1721 due to density currents in the Atlas region: Observations from the Saharan Mineral Dust Experiment
1722 2006 field campaign, *J Geophys Res-Atmos*, 112, Doi 10.1029/2007jd008774, 2007.

1723 Kocha, C., Flamant, C., Berckmens, J., Fink, A., Garcia-Carreras, L., Knippertz, P., Lafore, J.-P.,
1724 Marnas, F., Marsham, J. H., Parker, D. J., Rosenberg, P., Ryder, C. L., Tulet, P., and Washington, R.:
1725 How can a dusty cold pool change the diurnal evolution of the Saharan Boundary Layer?, *EGU 2013*,
1726 Vienna, 2013.

1727 Koehler, K. A., Kreidenweis, S. M., DeMott, P. J., Petters, M. D., Prenni, A. J., and Carrico, C. M.:
1728 Hygroscopicity and cloud droplet activation of mineral dust aerosol, *Geophys Res Lett*, 36, Doi
1729 10.1029/2009gl037348, 2009.

1730 Kok, J. F., Mahowald, N. M., Fratini, G., Gillies, J. A., Ishizuka, M., Leys, J. F., Mikami, M., Park, M.,
1731 S., Park, S. U., Van Pelt, R. S., and Zobeck, T. M.: An improved dust emission model - Part 1: Model
1732 description and comparison against measurements, *Atmos Chem Phys*, 14, 13023-13041, DOI
1733 10.5194/acp-14-13023-2014, 2014.

1734 Lacis, A. A., and Mishchenko, M. I.: Climate forcing, climate sensitivity, and climate response: A
1735 radiative modeling perspective on atmospheric aerosols, *Dahl Ws Env*, 17, 11-42, 1995.

1736 Lance, S., Medina, J., Smith, J. N., and Nenes, A.: Mapping the operation of the DMT Continuous
1737 Flow CCN counter, *Aerosol Sci Tech*, 40, 242-254, Doi 10.1080/02786820500543290, 2006.

1738 Lavaysse, C., Flamant, C., Janicot, S., Parker, D. J., Lafore, J. P., Sultan, B., and Pelon, J.: Seasonal
1739 evolution of the West African heat low: a climatological perspective, *Clim Dynam*, 33, 313-330, DOI
1740 10.1007/s00382-009-0553-4, 2009.

1741 Legrand, M., Pietras, C., Brogniez, G., Haeffelin, M., Abuhassan, N. K., and Sicard, M.: A high-
1742 accuracy multiwavelength radiometer for in situ measurements in the thermal infrared. Part I:
1743 Characterization of the instrument, *J Atmos Ocean Tech*, 17, 1203-1214, Doi 10.1175/1520-
1744 0426(2000)017<1203:Ahamrf>2.0.Co;2, 2000.

1745 Lensky, I. M., and Rosenfeld, D.: Clouds-Aerosols-Precipitation Satellite Analysis Tool (CAPSAT),
1746 *Atmos Chem Phys*, 8, 6739-6753, 2008.

1747 Liu, P. S. K., Leaitch, W. R., Strapp, J. W., and Wasey, M. A.: Response of Particle Measuring
1748 Systems Airborne Asasp and Pcap to NaCl and Latex-Particles, *Aerosol Sci Tech*, 16, 83-95, Doi
1749 10.1080/02786829208959539, 1992.

1750 Mahowald, N., Albani, S., Kok, J. F., Engelstaeder, S., Scanza, R., Ward, D. S., and Flanner, M. G.:
1751 The size distribution of desert dust aerosols and its impact on the Earth system, *Aeolian Res*, 15, 53-
1752 71, DOI 10.1016/j.aeolia.2013.09.002, 2014.

1753 Marenco, F., Amiridis, V., Marinou, E., Tsekeli, A., and Pelon, J.: Airborne verification of CALIPSO
1754 products over the Amazon: a case study of daytime observations in a complex atmospheric scene,
1755 *Atmos Chem Phys*, 14, 11871-11881, DOI 10.5194/acp-14-11871-2014, 2014.

1756 Marsham, J. H., Parker, D. J., Grams, C. M., Johnson, B. T., Grey, W. M. F., and Ross, A. N.:
1757 Observations of mesoscale and boundary-layer scale circulations affecting dust transport and uplift
1758 over the Sahara, *Atmos Chem Phys*, 8, 6979-6993, 2008a.

1759 Marsham, J. H., Parker, D. J., Grams, C. M., Taylor, C. M., and Haywood, J. M.: Uplift of Saharan
1760 dust south of the intertropical discontinuity, *J Geophys Res-Atmos*, 113, Doi 10.1029/2008jd009844,
1761 2008b.

1762 Marsham, J. H., Knippertz, P., Dixon, N. S., Parker, D. J., and Lister, G. M. S.: The importance of the
1763 representation of deep convection for modeled dust-generating winds over West Africa during
1764 summer, *Geophys Res Lett*, 38, Doi 10.1029/2011gl048368, 2011.

1765 Marsham, J. H., Dixon, N. S., Garcia-Carreras, L., Lister, G. M. S., Parker, D. J., Knippertz, P., and
1766 Birch, C. E.: The role of moist convection in the West African monsoon system: Insights from
1767 continental-scale convection-permitting simulations, *Geophys Res Lett*, 40, 1843-1849,
1768 10.1002/GrL.50347, 2013a.

1769 Marsham, J. H., Hobby, M., Allen, C., J. B., Bart, M., Brooks, B., Cavazos-Guerra, C., Engelstaeder,
1770 S., Gascoyne, M., McQuaid, J., O'Leary, A., Ouchene, B., Ouladichir, A., Parker, D., Saci, A., Salah-
1771 Ferroudj, M., Todd, M., and Washington, R.: Meteorology and dust in the central Sahara:
1772 Observations from Fennec supersite-1 during the June 2011 Intensive Observation Period, accepted
1773 for publication in *Journal of Geophysical Research-Atmospheres*, 2013b.

1774 Marsham, J. H., Hobby, M., Allen, C. J. T., Banks, J. R., Bart, M., Brooks, B. J., Cavazos-Guerra, C.,
1775 Engelstaeder, S., Gascoyne, M., Lima, A. R., Martins, J. V., McQuaid, J. B., O'Leary, A., Ouchene, B.,
1776 Ouladichir, A., Parker, D. J., Saci, A., Salah-Ferroudj, M., Todd, M. C., and Washington, R.:
1777 Meteorology and dust in the central Sahara: Observations from Fennec supersite-1 during the June
1778 2011 Intensive Observation Period, *J Geophys Res-Atmos*, 118, 4069-4089, Doi 10.1002/Jgrd.50211,
1779 2013c.

1780 McConnell, C. L., Highwood, E. J., Coe, H., Formenti, P., Anderson, B., Osborne, S., Nava, S.,
1781 Desboeufs, K., Chen, G., and Harrison, M. A. J.: Seasonal variations of the physical and optical

1782 characteristics of Saharan dust: Results from the Dust Outflow and Deposition to the Ocean (DODO)
1783 experiment, *J Geophys Res-Atmos*, 113, Doi 10.1029/2007jd009606, 2008.

1784 Messager, C., Parker, D. J., Reitebuch, O., Agusti-Panareda, A., Taylor, C. M., and Cuesta, J.:
1785 Structure and dynamics of the Saharan atmospheric boundary layer during the West African
1786 monsoon onset: observations and analyses from the research flights of 14 and 17 July 2006, *Q J Roy*
1787 *Meteor Soc*, 136, 107-124, doi:10.1002/qj.469, 2010.

1788 Müller, D., Weinzierl, B., Petzold, A., Kandler, K., Ansmann, A., Müller, T., Tesche, M.,
1789 Freudenthaler, V., Esselborn, M., Heese, B., Althausen, D., Schladitz, A., Otto, S., and Knippertz, P.:
1790 Mineral dust observed with AERONET Sun photometer, Raman lidar, and in situ instruments during
1791 SAMUM 2006: Shape-independent particle properties, *J Geophys Res-Atmos*, 115, Doi
1792 10.1029/2009jd012520, 2010.

1793 Müller, D., Lee, K. H., Gasteiger, J., Tesche, M., Weinzierl, B., Kandler, K., Müller, T., Toledano, C.,
1794 Otto, S., Althausen, D., and Ansmann, A.: Comparison of optical and microphysical properties of pure
1795 Saharan mineral dust observed with AERONET Sun photometer, Raman lidar, and in situ instruments
1796 during SAMUM 2006, *J Geophys Res-Atmos*, 117, Doi 10.1029/2011jd016825, 2012.

1797 Nakajima, T., Tonna, G., Rao, R. Z., Boi, P., Kaufman, Y., and Holben, B.: Use of sky brightness
1798 measurements from ground for remote sensing of particulate polydispersions, *Appl Optics*, 35, 2672-
1799 2686, Doi 10.1364/Ao.35.002672, 1996.

1800 Omar, A. H., Winker, D. M., Kittaka, C., Vaughan, M. A., Liu, Z. Y., Hu, Y. X., Trepte, C. R., Rogers, R.
1801 R., Ferrare, R. A., Lee, K. P., Kuehn, R. E., and Hostetler, C. A.: The CALIPSO Automated Aerosol
1802 Classification and Lidar Ratio Selection Algorithm, *J Atmos Ocean Tech*, 26, 1994-2014, Doi
1803 10.1175/2009jtecha1231.1, 2009.

1804 Osborne, S. R., Baran, A. J., Johnson, B. T., Haywood, J. M., Hesse, E., and Newman, S.: Short-wave
1805 and long-wave radiative properties of Saharan dust aerosol, *Q J Roy Meteor Soc*, 137, 1149-1167,
1806 Doi 10.1002/qj.771, 2011.

1807 Paluch, I. R.: Entrainment Mechanism in Colorado Cumuli, *J Atmos Sci*, 36, 2467-2478, Doi
1808 10.1175/1520-0469(1979)036<2467:Temicc>2.0.Co;2, 1979.

1809 Pappas, V., Ryder, C. L., Highwood, E. J., and Young, S.: Comparison of Fennec airborne in-situ
1810 dust measurements with CALIOP spaceborne lidar retrievals of extinction, *Remote Sensing of*
1811 *Environment*, in prep.

1812 Petersen, G. N., and Renfrew, I. A.: Aircraft-based observations of air-sea fluxes over Denmark
1813 Strait and the Irminger Sea during high wind speed conditions, *Q J Roy Meteor Soc*, 135, 2030-2045,
1814 Doi 10.1002/qj.355, 2009.

1815 Preissler, J., Wagner, F., Pereira, S. N., and Guerrero-Rascado, J. L.: Multi-instrumental
1816 observation of an exceptionally strong Saharan dust outbreak over Portugal, *J Geophys Res-Atmos*,
1817 116, Doi:10.1029/2011jd016527, 2011.

1818 Raymond, D. J., and Blyth, A. M.: Precipitation Development in a New-Mexico Thunderstorm, *Q J*
1819 *Roy Meteor Soc*, 115, 1397-1423, DOI 10.1002/qj.49711549011, 1989.

1820 Redelsperger, J. L., Thorncroft, C. D., Diedhiou, A., Lebel, T., Parker, D. J., and Polcher, J.: African
1821 monsoon multidisciplinary analysis - An international research project and field campaign, *B Am*
1822 *Meteorol Soc*, 87, 1739-1746, Doi 10.1175/Bams-87-12-1739, 2006.

1823 Reid, J. S., Jonsson, H. H., Maring, H. B., Smirnov, A., Savoie, D. L., Cliff, S. S., Reid, E. A., Livingston,
1824 J. M., Meier, M. M., Dubovik, O., and Tsay, S. C.: Comparison of size and morphological
1825 measurements of coarse mode dust particles from Africa, *J Geophys Res-Atmos*, 108, Doi
1826 10.1029/2002jd002485, 2003.

1827 Renfrew, I. A., Petersen, Outten, Sproson, Moore, Hay, Ohigashi, Zhang, Kristjansson, Fore,
1828 Olafsson, Gray, Irvine, Bovis, Brown, Swinbank, Haine, Lawrence, Pickart, Shapiro, and Woolley: The
1829 Greenland flow distortion experiment, *B Am Meteorol Soc*, 89, 1307-1324, Doi
1830 10.1175/2008bams2508.1, 2008.

1831 Roberts, G. C., and Nenes, A.: A continuous-flow streamwise thermal-gradient CCN chamber for
1832 atmospheric measurements, *Aerosol Sci Tech*, 39, 206-221, Doi 10.1080/027868290913988, 2005.

1833 Rodwell, M. J., and Jung, T.: Understanding the local and global impacts of model physics
1834 changes: An aerosol example, *Q J Roy Meteor Soc*, 134, 1479-1497, Doi 10.1002/qj.298, 2008.

1835 Rose, D., Gunthe, S. S., Mikhailov, E., Frank, G. P., Dusek, U., Andreae, M. O., and Poschl, U.:
1836 Calibration and measurement uncertainties of a continuous-flow cloud condensation nuclei counter
1837 (DMT-CCNC): CCN activation of ammonium sulfate and sodium chloride aerosol particles in theory
1838 and experiment, *Atmos Chem Phys*, 8, 1153-1179, 2008.

1839 Rosenberg, P., Dean, A., Williams, P., Minikin, A., Pickering, M., and Petzold, A.: Particle sizing
1840 calibration with refractive index correction for light scattering optical particle counters and impacts
1841 upon PCASP and CDP data collected during the Fennec campaign, *Atmospheric Measurement
1842 Technique Discussions*, 5, 97-135, doi:10.5194/amtd-5-97-2012, 2012.

1843 Rosenberg, P. D., Parker, D. J., Ryder, C. L., Marsham, J. H., Garcia-Carreras, L., Dorsey, J. R.,
1844 Brooks, I. M., Dean, A. R., Crosier, J., McQuaid, J. B., and Washington, R.: Quantifying particle size
1845 and turbulent scale dependence of dust flux in the Sahara using aircraft measurements, *J Geophys
1846 Res-Atmos*, 119, 7577-7598, Doi 10.1002/2013jd021255, 2014.

1847 Ryder, C. L., Highwood, E. J., Lai, T. M., Sodemann, H., and Marsham, J. H.: Impact of Atmospheric
1848 Transport on the Evolution of Microphysical and Optical Properties of Saharan Dust, accepted at
1849 *Geophysical Research Letters*, 2013a.

1850 Ryder, C. L., Highwood, E. J., Rosenberg, P., Trembath, J., Brooke, J., Bart, M., Dean, A., Crosier, J.,
1851 Dorsey, J., Brindley, H., Banks, J. R., Marsham, J. H., McQuaid, J. B., Sodemann, H., and Washington,
1852 R.: Optical properties of Saharan dust aerosol and contribution from the coarse mode as measured
1853 during the Fennec 2011 aircraft campaign, *Atmos. Chem. Phys.*, 13, 303-325, doi:10.5194/acp-13-
1854 303-2013, 2013b.

1855 Sassen, K., DeMott, P. J., Prospero, J. M., and Poellot, M. R.: Saharan dust storms and indirect
1856 aerosol effects on clouds: CRYSTAL-FACE results, *Geophys Res Lett*, 30, Doi 10.1029/2003gl017371,
1857 2003.

1858 Schepanski, K., Tegen, I., Laurent, B., Heinold, B., and Macke, A.: A new Saharan dust source
1859 activation frequency map derived from MSG-SEVIRI IR-channels, *Geophys Res Lett*, 34, Doi
1860 10.1029/2007gl030168, 2007.

1861 Schepanski, K., Tegen, I., and Macke, A.: Saharan dust transport and deposition towards the
1862 tropical northern Atlantic, *Atmos Chem Phys*, 9, 1173-1189, 2009.

1863 Schepanski, K., Flamant, C., Chaboureau, J. P., Kocha, C., Banks, J. R., Brindley, H. E., Lavaysse, C.,
 1864 Marnas, F., Pelon, J., and Tulet, P.: Characterization of dust emission from alluvial sources using
 1865 aircraft observations and high-resolution modeling, *J Geophys Res-Atmos*, 118, 7237-7259, Doi
 1866 10.1002/Jgrd.50538, 2013.

1867 Scheuvens, D., Schutz, L., Kandler, K., Ebert, M., and Weinbruch, S.: Bulk composition of northern
 1868 African dust and its source sediments - A compilation, *Earth-Sci Rev*, 116, 170-194, DOI
 1869 10.1016/j.earscirev.2012.08.005, 2013.

1870 Schuster, G. L., Vaughan, M., MacDonnell, D., Su, W., Winker, D., Dubovik, O., Lapyonok, T., and
 1871 Trepte, C.: Comparison of CALIPSO aerosol optical depth retrievals to AERONET measurements, and
 1872 a climatology for the lidar ratio of dust, *Atmos Chem Phys*, 12, 7431-7452, DOI 10.5194/acp-12-
 1873 7431-2012, 2012.

1874 Sodemann, H., Lai, M., Marenco, F., Ryder, C. L., Flamant, C., Knippertz, P., Rosenberg, P., Bart,
 1875 M., and McQuaid, J.: Lagrangian dust model simulations for a case of moist convective dust emission
 1876 and transport in the western Sahara region during Fennec/LADUNEX, *J Geophys Res-Atmos*,
 1877 submitted manuscript.

1878 Stein, T. H. M., Parker, D. J., Delanoe, J., Dixon, N. S., Hogan, R. J., Knippertz, P., Maidment, R. I.,
 1879 and Marsham, J. H.: The vertical cloud structure of the West African monsoon: A 4 year climatology
 1880 using CloudSat and CALIPSO, *J Geophys Res-Atmos*, 116, Doi 10.1029/2011jd016029, 2011.

1881 Sternberg, R.: Into the Cauldron: A Meteorological Adventure,
 1882 <http://fennec.ouce.ox.ac.uk/movie.html>, in: EGU GeoCinema, 2013.

1883 Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian
 1884 particle dispersion model FLEXPART version 6.2, *Atmos Chem Phys*, 5, 2461-2474, 2005.

1885 Sultan, B., and Janicot, S.: The West African monsoon dynamics. Part II: The "preonset" and
 1886 "onset" of the summer monsoon, *J Climate*, 16, 3407-3427, Doi 10.1175/1520-
 1887 0442(2003)016<3407:Twamdp>2.0.Co;2, 2003.

1888 Sun, D. L., Lau, W. K. M., Kafatos, M., Boybeyi, Z., Leptoukh, G., Yang, C. W., and Yang, R. X.:
 1889 Numerical Simulations of the Impacts of the Saharan Air Layer on Atlantic Tropical Cyclone
 1890 Development, *J Climate*, 22, 6230-6250, Doi 10.1175/2009jcli2738.1, 2009.

1891 Tesche, M., Ansmann, A., Muller, D., Althausen, D., Mattis, I., Heese, B., Freudenthaler, V.,
 1892 Wiegner, M., Esselborn, M., Pisani, G., and Knippertz, P.: Vertical profiling of Saharan dust with
 1893 Raman lidars and airborne HSRL in southern Morocco during SAMUM, *Tellus B*, 61, 144-164, DOI
 1894 10.1111/j.1600-0889.2008.00390.x, 2009.

1895 Todd, M. C., Allen, C. J. T., Bart, M., Bechir, M., Bentefouet, J., Brooks, B. J., Cavazos-Guerra, C.,
 1896 Clovis, T., Deyane, S., Dieh, M., Engelstaedter, S., Flamant, C., Garcia-Carreras, L., Gandega, A.,
 1897 Gascoyne, M., Hobby, M., Kocha, C., Lavaysse, C., Marsham, J. H., Martins, J. V., McQuaid, J. B.,
 1898 Ngamini, J. B., Parker, D. J., Podvin, T., Rocha-Lima, A., Traore, S., Wang, Y., and Washington, R.:
 1899 Meteorological and dust aerosol conditions over the western Saharan region observed at Fennec
 1900 Supersite-2 during the intensive observation period in June 2011, *J Geophys Res-Atmos*, 118, 8426-
 1901 8447, Doi 10.1002/Jgrd.50470, 2013.

1902 Tompkins, A. M., Cardinali, C., Morcrette, J. J., and Rodwell, M.: Influence of aerosol climatology
 1903 on forecasts of the African Easterly Jet, *Geophys Res Lett*, 32, Doi:10.1029/2004gl022189, 2005.

1904 Trembath, J.: Airborne CCN Measurements, SAES, University of Manchester, 2012.

1905 Trembath, J., Bart, M., and Brooke, J.: FAAM Technical Note: Efficiencies of modified Rosemount
1906 housings for sampling aerosol on a fast atmospheric research aircraft, Facility for Airborne
1907 Atmospheric Measurements, FAAM, Cranfield, UK,
1908 http://www.faam.ac.uk/index.php/component/docman/cat_view/140-science-instruments, 2012.

1909 Tulet, P., Crahan-Kaku, K., Leriche, M., Aouizerats, B., and Crumeyrolle, S.: Mixing of dust aerosols
1910 into a mesoscale convective system Generation, filtering and possible feedbacks on ice anvils, *Atmos*
1911 *Res*, 96, 302-314, DOI 10.1016/j.atmosres.2009.09.011, 2010.

1912 Twohy, C. H., Kreidenweis, S. M., Eidhammer, T., Browell, E. V., Heymsfield, A. J., Bansemer, A. R.,
1913 Anderson, B. E., Chen, G., Ismail, S., DeMott, P. J., and Van den Heever, S. C.: Saharan dust particles
1914 nucleate droplets in eastern Atlantic clouds, *Geophys Res Lett*, 36, Doi 10.1029/2008gl035846, 2009.

1915 Veihelmann, B., Konert, M., and van der Zande, W. J.: Size distribution of mineral aerosol: using
1916 light-scattering models in laser particle sizing, *Appl Optics*, 45, 6022-6029, Doi
1917 10.1364/Ao.45.006022, 2006.

1918 Washington, R., Todd, M. C., Engelstaedter, S., Mbainayel, S., and Mitchell, F.: Dust and the low-
1919 level circulation over the Bodele Depression, Chad: Observations from BoDEX 2005, *J Geophys Res-*
1920 *Atmos*, 111, 10.1029/2005jd006502, 2006.

1921 Washington, R., Flamant, C., Parker, D. J., Marsham, J. H., McQuaid, J. B., Brindley, H., Todd, M.,
1922 Highwood, E. J., Ryder, C. L., Chaboureau, J.-P., Kocha, C., Bechir, M., and Saci, A.: Fennec - The
1923 Saharan Climate System, *CLIVAR Exchanges*, 17, 31-32, 2012.

1924

1925

1926 **Tables**

Campaign	Date	Reference
JET2000	Summer 2000	Thornicroft et al., 2003
Saharan Dust Experiment (SHADE)	Summer 2000	Haywood et al., 2003
Dust and Biomass Experiment DABEX	Winter 2006	Haywood et al., 2008
Dust Outflow and Deposition to the Ocean (DODO)	Winter/Summer 2006	McConnell et al., 2008
African Monsoon Multidisciplinary Analysis (AMMA)	2006	Redelsperger et al., 2006
NASA AMMA (NAMMA)	Summer 2006	Zipser et al., 2009
Saharan Mineral Dust Experiment 1 (SAMUM1 and 2)	2006/2008	Heintzenberg, 2009
Saharan Mineral Dust Experiment 2 (SAMUM2)	2008	Ansmann et al. (2011)
Geostationary Earth Radiation Budget Intercomparison of Long-wave and Short-wave radiation (GERBILS)	Summer 2007	Haywood et al., 2011

1927 Table 1: Previous aircraft programmes in the region.

1928

1929

IOP	Date	Operating base	Aircraft	Number of Flights	Number of Dropsondes
Pilot study	April 2011	Ouarzazate, Morocco	BAe146	6	42
IOP1	June 2011	Fuerteventura, Canaries	BAe146 FF-20	16 (BAe146) 18 (FF-20)	81 (BAe146) 136 (FF-20)
IOP2	June 2012	Fuerteventura, Canaries	BAe146	14	40

1930

Table 2: Overview of IOPs

1931

Name	Instrument	Measures	Sampling rate	Reference for more detail
LNG LIDAR	Downward facing high spectral resolution LIDAR (CAB)	Atmospheric backscatter coefficients at 532 and 1064 nm. Aerosol extinction coefficients at 532 nm.	20 Hz	Banks et al., 2013; Schepanski et al., 2013
AVAPS II	Airborne Vertical Atmospheric Profiler System & RD94 GPS dropsondes (CAB)	Profiles of position, pressure, temperature, relative humidity, wind speed and direction	2 Hz	
Basler SCA1400-30FM	Downward facing monochrome (black/white) camera (CAB)	Pictures of ground surface with a resolution of 1392 x 1040 pixels. Each photograph covers a horizontal area of 3.3 km x 4.4 km along the track for a nominal aircraft altitude of 11 km asl	1 Hz	Schepanski et al., 2013
Kipp & Zonen CPM22	Precision Spectral Pyranometer (RFM & BLM)	0.2–3.6 μ m up- and downwelling irradiance	0.2 Hz	
Kipp & Zonen CGR4	Precision Infrared Radiometer (RFM & BLM)	4.5–42 μ m up- and downwelling irradiance	0.05 Hz	
CLIMAT CE 332	Downward-facing radiometer (BLM)	Spectrally resolved directional radiance: brightness temperature at 8.7, 10.8 and 12 μ m	1 Hz	Legrand et al., 2000
General Eastern 1011B (RDM)	Hygrometer using the chilled-mirror technique (RDM)	Water vapour (dew point temperature) over range -65 to 50°C	1 Hz	
Aerodata Humicap (RDM)	Humidity capacity sensor (RDM)	Relative humidity (0-100%)	10 Hz	
Rosemount 1201	Pressure sensor (NBM)	Static pressure (250-1035 hPa)	10 Hz	
Rosemount 1221	Pressure sensors (NBM)	Differential incidence and drift pressures (± 70 hPa)	10 Hz	
Rosemount 102 E2AL	Temperature sensor (RDM)	Temperatures (non de-iced), calibrated over range -60° to 40°C; uncertainty $\pm 0.5^\circ$ C	10 Hz	
Rosemount 871	Ice Probe (RDM)	Indication of supercooled water	N/A	
LITTON 90-100	Inertial navigation unit (CAB)	Aircraft position, aircraft velocity components, aircraft attitude (pitch, roll, yaw) and attitude rates, ground speed, wind speed and direction, and drift angle (position and acceleration at 1 Hz)	66 Hz	
TRT AHV 8	Radar altimeter (CAB)	Altitude (0-5000 ft, accuracy $\pm 2\%$)	10 Hz	
Bancom BC635 on Trimble Transducer	Global Positioning System (CAB)	Aircraft position, velocity, and time standard	1 Hz	
Collins ADC 80	Air Data Computer (CAB)	Barometric altitude (-2000 to 7000 ft) and true air speed	10 Hz	

1933
1934 Table 3: Instruments onboard the SAFIRE Falcon 20 during the 2011 IOP. NBM= Nose Boom Mounted; CAB=inside aircraft cabin, RDM=Radome Mounted, BLM=Belly Mounted; RFM=Roof Mounted

Name	Instrument	Property Measured	Sampling rate	Reference	IOP in Use
Aircraft Position and Meteorological Measurements					
GPS	Patch	Aircraft position, velocity, and time standard	1 Hz	FAAM	All
INU	Inertial Navigation Unit	Aircraft velocity components, attitude, attitude rates, ground speed, and drift angle	32 Hz	FAAM	All
RadAlt	Radar Altimeter	Altitude above surface, max 5000 ft AGL (accuracy $\pm 2\%$)	2 Hz	FAAM	All
RVSM	Reduced vertical separation minimum data system	Static and pitot-static pressures, pressure altitude, indicated air speed	32 Hz	FAAM	All
Rosemount Temperature Sensors		Deiced and non-deiced, calibrated over range -60° to 30°C ; ($\pm 0.3^\circ\text{C}$)	32 Hz	FAAM	All
Turbulence probe	Turbulence (see also RVSM)	Air speed and incidence angle; 3-D wind components; measurement uncertainty $\pm 0.2 \text{ m s}^{-1}$	32 Hz	Peterson & Renfrew, 2008	All
AIMMS	Aircraft-Integrated Meteorological Measurement System (Aventech Research, Inc)	General meteorological parameters, generally used as backup to core turbulence probe. WM		FAAM	All
AVAPS	Airborne Vertical Atmospheric Profiler System (Vaisala RD94 GPS dropsondes)	Profiles of position, pressure, temperature, relative humidity, wind speed and direction	2 Hz	FAAM	All
Water Content Measurements					
TWC	Total water content using a Lyman-alpha absorption hygrometer	Water (H_2O) over range $0\text{--}20 \text{ g kg}^{-1}$ and accuracy $\pm 0.15 \text{ g kg}^{-1}$	64 Hz	FAAM	All
General Eastern	Hygrometer (using the chilled-mirror technique)	Water vapor (dewpoint temperature) over 220–320K; instrument response time can be up to 30 s; measurement uncertainty $\pm 0.25\text{K}$ above 273.15 K, $\pm 1\text{K}$ at 210 K	4 Hz	FAAM	All
Johnson Williams	Liquid water content probe	Liquid water concentration in clouds using heated wire resistance bridge over $0\text{--}3 \text{ gm}^{-3}$; uncertainty $\pm 10\%$	4 Hz	FAAM	All
Nevzorov	Liquid and total water content probe	Liquid and total (ice plus liquid) water in clouds using a heated wire over range $0.003\text{--}3 \text{ gm}^{-1}$; accuracy $\pm 10\%$	8 Hz	FAAM	All
Aircraft Inlets					
Rosemount 102E Inlets	Aerosol inlets for cabin instrumentation	Originally designed for PRT Measurements, only accumulation mode particles passed	n/a	Trembath (2012)	All

LTI	Low Turbulence Inlet	Fully characterised inlet, passes coarse mode particles	n/a	Trembath 2012, Wilson et al., 2004	All
Filter sample inlet	Parallel coarse mode samplers	Supplies filter samples for offline analysis	n/a	Formenti et al, 2014	All
In-situ Aerosol Measurements					
PCASP	Passive Cavity Aerosol Spectrometer Probe (PMS canister instrument)	PNC, 0.1–3 µm, OPT, (WM)	1 Hz	Rosenberg et al., 2012, FAAM	All
CDP	Cloud Droplet Probe	PNC, 3-50 µm, OPT, (WM)	1Hz standard, 10Hz during Fennec	Rosenberg et al., 2012, FAAM	All
CIP15	Cloud Imaging Probe	PNC, 15-930 µm, 15 µm resolution, SH, (WM). Provided by U.Manchester in 2011 and by FAAM in 2012. 2012 data suffered from electronic noise.	10Hz	Rosenberg et al., 2012, FAAM	All
CIP100	Cloud Imaging Probe	PNC, 100-6200 µm, 100 µm resolution, SH, (WM)	1Hz	FAAM	2012
GRIMM OPC	Grimm Technik 1.129 Sky Optical Particle Counter	PNC, 0.25-32 µm, placed behind different inlets, CAB, OPT (CAB)	1Hz	Heim et al., 2008	All
2D-C	Two-dimensional cloud particle imaging probe (PMS canister instrument)	5-s-averaged values of PNC, condensed water content, mean volume radius, precipitation rate, and size spectrum (25–800 µm). WM–SH (WM)	1 Hz	FAAM	2012
SID2H	Small Ice Detector	PNC, 2-60 µm, OPT, also non-sphericity, (WM)	1Hz	Cotton et al., 2010	All
CAS	Cloud and Aerosol Spectrometer	PNC, 0.6 – 50 µm, WM, OPT, (WM), part of U.Manchester CAPS probe.	1Hz	(Baumgardner et al., 2001)	2011
University of Manchester CAPS Probe	Cloud, Aerosol and Precipitation Spectrometer (DMT)	Aerosol particle and cloud hydrometeor size (0.51 - 50 µm). Liquid water content from 0.01 to 3 g m ⁻³ . Aerosol probes comprise CAS and CIP15 instruments, (WM)	1Hz	FAAM	2011
CCN	Dual column continuous flow cloud condensation nuclei counter (DMT)	Concentration and properties of cloud condensation nuclei, (CAB)	1 Hz	Trembath (2012)	All
CPC	Modified TSI 3786 condensation particle counter	Aerosol particles (2.5 nm – 3 µm) (CAB)	1 Hz	Trembath (2012)	All
Nephelometer	TSI 3563 Integrating nephelometer	Total scattering and hemispheric backscattering coefficient at 450, 550, and 700 nm, (CAB)	1 Hz	Ryder et al., (2013b), FAAM	All
PSAP	Radiance Research Particle Soot Absorption Photometer	Absorption coefficient at 567 nm, (CAB)	1 Hz	Ryder et al., (2013b), FAAM	All
Radiometric Measurements					

BBR	Broadband shortwave Radiometers (pyranometers)	0.3–3 µm & 0.7–3 µm up and downwelling irradiance	1 Hz	FAAM	All
SHIMS	Spectral Hemispheric Irradiance MeasurementS	Spectrally resolved irradiance, up and downwelling, 0.3-1.7 µm	0.1Hz	Osborne et al., 2011	All
SWS	ShortWave Spectrometer	Spectrally resolved directional radiance, 0.3-1.7 µm	0.1Hz	Osborne et al., 2011	All
ARIES	Airborne Research Interferometer Evaluation System	Spectrally resolved directional radiance, 3.3-18 µm	1Hz	Wilson et al., 1999; Osborne et al., 2011	All
Heimann	Downward-facing radiometer	Downward facing brightness temperature (8–14 µm)	4 Hz	FAAM	All
LIDAR	Downward facing aerosol LIDAR (Leosphere ALS450)	Aerosol and thin cloud retrievals, qualitative depolarisation	2 sec	Marencio et al., (2011; 2013)	All
Video cameras	Up/downward, forward, and rear-view cameras	Digital video recordings		FAAM	All
Chemistry Measurements					
Ozone	TECO 49C UV photometric instrument	Ozone (O ₃); integration time 4 s	1 Hz	FAAM	All
Carbon Monoxide	CO Aerolaser AL5002	Carbon monoxide (CO) by UV fluorescence at 150 nm	1 Hz	FAAM	2012

1936

1937
1938
1939
1940
1941

Table 4: Instrumentation on the BAe146 aircraft relevant to Fennec. WM=Wing Mounted, CAB=inside aircraft cabin, PNC=particle number concentration, OPT=optical scattering measurements, SH=Light Shadowing Measurements. Size ranges shown for optical instruments refer to nominal ranges provided by manufacturers, i.e. not corrected for aerosol type-specific refractive indices. FAAM = refer to FAAM website where full instrumentation details are provided, www.faam.ac.uk/index.php/science-instruments.

1942

1943

Date	Flight Number	Times, UTC	Locations	Purpose
4 April	b589	1551 to 1852	MAU	Overflight of dust front
5 April	b590	0850 to 1328	MAU	Sampling of maritime air underlying dusty continental air
	b591	1505 to 1838	MAU	Sampling of maritime air underlying dusty continental air
7 April	b592 (2 flights)	0652 to 1706	MAU	Sampling of dust in recovering SABL
8 April	b593	0829 to 1341	MAU	Surface albedo impact on recovering SABL
9 April	b594	0913 to 1359	Ouarzazate to UK	Sampling of dust transported northwards towards UK

1944

Table 5: April 2011 pilot campaign flights of BAe146.

1945

Date	Flight Number	Times, UTC	Locations	Purpose
2 June	F09	1527-1858	EAO	Dust outflow over EAO
6 June	F10	1200-1533	EAO	Dust outflow over EAO
10 June	F11	1028-1401	EAO, MAU, SEN	Dust outflow over EAO & PBL over MAU
10 June	F12	161-1940	EAO, MAU, SEN	Dust outflow over EAO & PBL over MAU
11 June	F13	0906-229	N MAU	Dust uplift, RAIN4DUST
11 June	F14	14401-809	N MAU	PBL
13 June	F15	1100-1422	N MAU and N MAL	Survey of N MAU & dust associated with Mediterranean surge
14 June	F16	1437-1809	N MAU	PBL
15 June	F17	1433-1802	N MAU	PBL
16 June	F18	0913-1224	N MAU	Dust uplift, RAIN4DUST
16 June	F19	1442-1812	N MAU	PBL; approaching AEW
17 June	b600	0748-1241	MAL, N MAU	Characterisation of LLJ winds and dust
	F20	1528-1858	N MAL, N MAU	Survey of N MAU and N MAL & dust associated with Mediterranean surge and AEW
	b601	1443-1937	N MAL, N MAU	Characterisation of LLJ winds and dust
18 June	b602	0810-1240	N MAL, N MAU	Characterisation of LLJ winds and dust
	b603	1415-1555	Canary Islands	High altitude radiation instrument calibration
20 June	b604	1247-1751	MAU	Sampling of dust uplifted by MCS, LADUNEX
	F21	1322-1700	N and central MAU	Survey of dust associated with ITD and SHL
21 June	b605	0810-1158	MAU	Sampling of dust uplifted by Atlas Mts density current
	b606	1404-1920	MAU	SABL development and heat fluxes
	F22	0718-1035	N MAU and N MAL	Survey of dust associated with Mediterranean surge and density currents from Atlas Mts
	F23	1313-1630	N MAU and N MAL	Survey of dust associated with Mediterranean surge and density currents from Atlas Mts
22 June	b607	0804-1237	MAU, MAL	Sampling of SHL with LIDAR and dropsondes
	b608	1510-2016	MAU, MAL	Sampling of SHL with LIDAR and dropsondes
	F24	0917-1245	N MAU	Survey SHL; dust associated with Mediterranean surge (N) & ITD (S & E)
	F25	1521-1849	N MAU	Survey of SHL; dust associated with Mediterranean surge (N) & ITD (S & E)
23 June	F26	0833-1200	N MAU	Dust uplift, RAIN4DUST
24 June	b609	1129-1645	MAU	Dust-cloud interactions
25 June	b610	0731-1217	MAU	Dust uplift by LLJ
	b611	1414-1916	MAU	Overflight of Zouerate ground site
26 June	b612	0729-1222	MAU	Dust and radiative fluxes
	b613	1355-1859	MAU	SABL development and heat fluxes
27 June	b614	0634-1139	MAU	Dust uplift by LLJ
28 June	b615	0814-1129	Canary Islands	Radiation instrument calibration

Table 6: June 2011 IoP Flights. Flight numbers with preceding 'b' indicate BAe146 flight, with preceding 'F' indicate Falcon flight. Abbreviations: EAO=Eastern Atlantic Ocean, MAU=Mauritania, MAL=Mali, SEN=Senegal, FUE=Fuerteventura, ZOU=Zouerate supersite.

1946
1947
1948

1949

1950

Date	Flight Number	Times, UTC	Locations	Purpose
1 June	b698	0942 - 1708	UK to FUE	Science transit to FUE with radiation calibrations
6 June	b699	1201 - 1654	N MAL, N MAU	Atlantic Inflow 1
8 June	b700	0756 - 1257	N MAL, N MAU	Atlantic Inflow 2
9 June	b701	0755 - 1308	Central MAU	Dust at ITD 1
10 June	b702	0804 - 1241	Central MAU	Dust at ITD 2 (to Dakar)
	b703	1412 - 1720	EAO	Dust Outflow over EAO
11 June	b704	1214 - 1719	S MAU	Very heavy dust at ITD 3
12 June	b705	1127 - 1707	N MAL	Midday Heat fluxes
14 June	b706	1307 - 1813	N MAL	Dust uplift 1
15 June	b707	0913 - 1433	N MAL	Dust uplift 2
16 June	b708	0756 - 1308	N MAL, W MAU	Dust uplift by LLJ and Radiative Closure
17 June	b709	1214 - 1724	N MAL	Dust in SABL and Radiative Closure
18 June	b710	0751 to 1311	ZOU	SAVEX flight over Zouerate
19 June	b711	0755 to 1039	FUE and EAO	Science transit to Porto

1951 Table 7: June 2012 Fennec IOP flights.

1952

1953

Research Area	Key Findings	Reference
Publications Deriving from Fennec Aircraft Observations		
Size distribution measurements	A new method for correcting OPC data for particle optical properties	Rosenberg et al. (2012)
BAe146 inlets	BAe146 Rosemount inlet significantly excludes particles larger than 3 μm diameter	Trembath (2012)
Size distributions and optical properties of dust	Consistent presence of coarse and giant particles over Sahara; SSA at 550 nm 0.7 to 0.97 strongly related to particle size; inverse relationship between size and dust age.	Ryder et al., (2013b)
Impacts of transport on dust size distribution	d_{eff} decrease of 4.5 μm , and SSA increase from 0.92 to 0.95 between fresh and Atlantic SAL dust.	Ryder et al., (2013a)
Dust-ozone interactions	Increased dust surface area associated with fresh dust uplift and a large coarse mode act as a route for the reduced ozone concentrations.	Brooke (2014)
Dust fluxes	Size resolved dust fluxes follow the power law predicted by the Kok brittle fragmentation theory. Large size cut off is significantly larger than seen in other observations. Large fluxes were correlated with regions of varying topography.	Rosenberg et al. (2014)
Satellite retrievals of dust	Imperial SEVIRI dust AOD products are most effective at high dust loadings, but are sensitive to meteorological conditions; MODIS Deep Blue and MISR AOD products more consistent at lower dust loadings.	Banks et al. (2013)
Lagrangian modelling of dust uplift and transport	Validation of Lagrangian dust transport model with dust mass concentration underlines difficulties to quantify dust emission due to moist convection. Manual inversion approach constrains dust source and flux.	(Sodemann et al., submitted manuscript)
Dust uplift from fluvial sources	Dust emission from alluvial source observed by airborne remote sensing; Nocturnal LLJ drives morning dust uplift; explicit representation of endorheic systems as dust sources required in terms of their role as dust sources.	Schepanski et al.,(2013)
Structure and diurnal growth of the SABL	Turbulent structure, vertical fluxes and diurnal growth of SABL described with radiosondes, aircraft measurements and a LEM. Novel processes found, such as detrainment from the CBL top which acts to slow down CBL growth.	Garcia-Carreras et al. (2015)
Moisture transport pathways in the SHL region	Observation-based SHL characterisation; monsoon surge splits into two moisture transport pathways: a) around the SHL and b) towards northeast; afternoon CBL depth over-estimation by model leads to moisture advection error.	Engelstaedter et al. (2015)
Further information from Fennec		
Introduction to Fennec		Washington et al. (2012)
Ground-based observations	Supersite 1, Bordj Badji Moktar	Marsham et al., (2013)

Supersite 2, Zouerate	Todd et al. (2013)
The Fennec Automatic Weather Station Network	Hobby et al. (2013)

[Table 8: Key publications deriving from Fennec aircraft observations and summarizing other Fennec ground-based observations.](#)

1954

1955

Formatted: Caption

1956

1957 Figure Captions

1958

1959 Figure 1: The Fennec domain and climatology. Figure shows mean (2000–2012) June–September AOD from satellite
1960 MISR data (shaded, contour intervals are 0.4, 0.6, and 0.8) and key mean June–September circulation features derived
1961 from ERA-Interim reanalysis data (1979–2012), specifically the mean position of the Saharan heat low core (1008 hPa
1962 contour of sea level pressure, thick red contour); the mean position of the inter-tropical discontinuity (solid blue line, as
1963 defined by the 10 gkg^{-1} contour of 925 hPa specific humidity). Figure also highlights the location of the two Fennec
1964 supersites (SS1 yellow square, SS2 yellow circle), and approximate aircraft flight zone (green polygon). Also indicated are
1965 surface elevation (dashed cyan contour, 1000, 1500, and 2000m) and the approximate location of recent airborne field
1966 campaigns.

1967

1968 Figure 2: Flight tracks of the BAe146 and Falcon during Fennec: (a) Fennec Pilot, April 2011, BAe146; (b) June 2011,
1969 Falcon, (c) June 2011, BAe146, (d) June 2011, BAe146. Each colour shows a different flight. Note that in (b) and (d), the
1970 tracks of the following flights are the same and therefore not visible: F11, F12 and F26; F13 and F18; F14, F16, F17 and
1971 F19; F22 and F23; F24 and F25; b706 and b707.

1972

1973 Figure 3: Synoptic conditions during the Fennec flight campaigns. (a) 300hPa (m, shaded), ~~and~~ 925hPa geopotential
1974 height (white contours with intervals at 700, 725, 750 and 800m), ~~and~~ 925 hPa winds (ms^{-1}) ~~and~~ 15°C contour of 925 hPa
1975 temperature (blue line) ~~to show cold air advection~~, on 06UTC 4th April 2011. Feature A marks the position of the cut-off
1976 low. (b) Daily mean 200hPa geopotential height (m, shaded), 925hPa winds (ms^{-1}) and mean frequency of the SHL
1977 occurrence (white contours with intervals at 0.25, 0.5 and 0.75, as defined using the method of Lavaysse et al., [2009])
1978 averaged over the period 1-12th June 2011 (the maritime phase). Features A, B and C indicate the approximate
1979 locations of an upper level trough, SHL centre and maritime low level flow, respectively. (c) as (b) except for the period
1980 13-30th June 2011 (heat low phase) and where features A, B and C indicate the approximate locations of an upper level
1981 ridge, SHL centre and enhanced northeasterly 'Harmattan' level flow, respectively. (d) as (b) except for the period 1st-
1982 18th June 2012, and a 10.0 gkg^{-1} 925hPa specific humidity contour (blue line) and where features A, B, C and D indicate
1983 the approximate locations of an upper level trough, SHL extension trough, maritime low level flow and ITD bulge,
1984 respectively.

1985

1986 Figure 4: Example size distributions measured in different dust layers during Fennec 2011. Size distributions were
1987 measured using the PCASP (green), CDP (red) and CIP~~15~~ (purple). Solid lines show measurements from b600, during
1988 active uplift close to the desert surface; dashed lines show measurements from b612 which was dust aged by several
1989 days and well-mixed within a deep SABL. ~~E~~Vertical error bars show one standard deviation of the data combined with
1990 instrumental uncertainty, and only upwards errors are shown for clarity. Horizontal errors show uncertainty in bin
1991 centre diameter.

1992

1993 Figure 5: (a) Scatter plot of the elemental Fe/Ca versus the Si/Al ratios for the Fennec 2011 and 2012 samples compared
1994 to samples collected during the AMMA, DODO and GERBILS campaign (Formenti et al., 2014). Indications of the source
1995 regions according to the values of those tracers are also given. (b) Box plot of SSAs at 550nm measured during Fennec
1996 2011 and 2012 for horizontal runs corresponding to filter samples taken. SSAs are calculated from scattering measured
1997 by the nephelometer and absorption measured by the PSAP on the BAe146 mounted behind Rosemount inlets, and
1998 therefore represent accumulation mode only. Box lines represent the median and interquartile range, whiskers
1999 represent the minimum and maximum values, and square represents the mean.

2000

2001 Figure 6: Aerosol optical depths at 550 nm measured by the nephelometer and PSAP on the BAe146 during profiles,
2002 representing accumulation mode 550nm AOD. AODs are an underestimate since they do not include contribution from
2003 coarse particles. Circles represent 2011 data, diamonds 2012 data.

2004

2005 Figure 7: b609 dropsonde/aircraft moisture profiles and range-corrected LIDAR cross section of the scientific area of
2006 interest (red-blue colour scale, arbitrary logarithmic units) including an aircraft track coloured by the droplet
2007 concentration as measured by the CDP plus PCASP (black to red colour scale). The LIDAR data collected during descent
2008 (thick sloping black line) is plotted instead of the high level data when available. Above this, LIDAR data from the high

2009 level flight leg is shown. Arrows indicate locations of dropsondes. Sondes 1-3 were dropped on entry to the area and

2010 sonde 4 on exit.

2011

2012 Figure 8: Equivalent potential temperature and total water content during the flight. "Environment" points represent
2013 data from the descent out of cloud, "Boundary Layer" points represent data collected during aircraft ascent up to an
2014 altitude of 5000 m, "In cloud" points represent data collected during aircraft ascent above 5000 m where cloud droplet
2015 number was measured greater than 0.5 cm^{-3} and "Out of cloud" points represent data collected during aircraft ascent
2016 above 5000 m where cloud droplet number was less than 0.5 cm^{-3} . Mean in and out of cloud values are plotted shown
2017 with large circles outlined in black.

2018

2019 Figure 9: Box and whisker diagram of mineral dust mean surface area and ozone mass mixing ratio along the b707 (15
2020 June 2012) flight transect. Surface area is calculated from PCASP count median diameter.

2021

2022 Figure 10: Aircraft and satellite observations along the track of the outbound Falcon flight F23 on the 21st June (1352-
2023 1445 UT), across northern Mauritania and ending in northern Mali. Lower panel: LIDAR vertical extinction coefficient
2024 cross-section (at 532 nm); middle panel: co-located SEVIRI, MODIS, and LIDAR (LNG) AOD retrievals along the Falcon
2025 flight-track; upper panel: the along-track SEVIRI RGB 'desert-dust' imagery.

2026

2027 Figure 11: Volume size distributions from BAe146 flight b611 Profile 1 (1558 to 1627 UTC) compared to AERONET
2028 retrievals. Aircraft size distribution measurements are shown by green (PCASP), red (CDP) and purple (CIP 15). Solid lines
2029 show the median volume concentrations over the column up to 5.5km. Error bars show standard deviation over the
2030 column (where lower error bars reach below the plot minimum they have been omitted for clarity). Points with dashed
2031 lines represent the 10th and 90th percentiles across the column. AERONET retrievals from the Zouerate site over the day
2032 are shown in dark blue (morning), black (retrieved during the flight) and light blue (retrieved shortly after the flight).

2033

2034 Figure 12: Composite of the dust uplift potential (DUP, shading, $\text{m}^3 \text{ s}^{-3}$) for the air masses observed by the aircraft LIDARs
2035 during all flights from each campaign (blue lines). Calculations have been performed for tropospheric curtains along the
2036 flight tracks, integrating the dust uplift potential for the 3 days preceding each research flight. DUPs are shown for (a)
2037 Fennec 2011 Falcon flights; (b) Fennec 2011 BAe146 flights; (c) Fennec 2012 BAe146 flights.

2038

2039 Figure 13: (a) and (b) SEVIRI RGB dust imagery for 1000Z and 1700Z, and showing the flight tracks of flight b600 and b601
2040 respectively (BAe146 track in red, F20 track in yellow, black track sections show location of aircraft at satellite image
2041 time. (c), (d) and (e)a, b and e) UK Met Office wind forecasts for 06Z at 925hPa (c), 06Z at 10m (d) and 09Z for 925hPa
2042 (e), all for 17 June 2011 on the morning of the flight. d and e) SEVIRI RGB dust imagery for 1000Z and 1700Z, and
2043 showing the flight tracks of flight b600 and b601 respectively (BAe146 track in red, F20 track in yellow, black track
2044 sections show location of aircraft at satellite image time.

2045

2046 Figure 14: Aircraft measurements from the profile descent of b600 (around 1000Z, black) and b601 (around 1700Z, red)
2047 corresponding to the tracks, imagery and forecasts shown in Figure 13. Figure shows wind speed (u), wind direction,
2048 vertical wind speed (w), corrected extinction coefficient (Ext) calculated from the nephelometer scattering and PSAP

2049 absorption, potential temperature, and water vapour mixing ratio (r). Note that altitude is shown in pressure height,
2050 corresponding to minimum altitudes of 825m and 784m AGL respectively for b600 and b601.

2051

2052 Figure 15: (a) AOD computed from the Falcon 20 LNG LIDAR extinction coefficient profile at around 1000 UTC on 21 June
2053 [2011, flight F22.](#) (ba) Cross section [of](#) the LNG LIDAR extinction coefficient (10^{-6} m^{-1}) derived from the LNG LIDAR flying
2054 [on the Falcon 20 \(bottom\)](#) and the AOT computed from the LIDAR extinction coefficient profile (top) at around 1000 UTC
2055 [on 21 June 2011, flight F22.](#) (bc) Shortwave downwelling irradiance (W m^{-2} , red) and extinction coefficient (10^{-6} m^{-1} ,
2056 black) as a function of the pressure during the ascent of the BAe146 from within the haboob to upper levels, flight b605.
2057 Note that the minimum pressure height of 360 m is equivalent to 105 m above ground level.

2058

2059 Figure 16: a) Measurements made during low level runs in flight b708 on 16 June 2012 sampling uplifted dust by a low
2060 level jet. Black line shows radar altitude (height above ground, left axis), accumulation mode extinction measured by a
2061 nephelometer and PSAP (green line, left axis), and downwelling shortwave irradiance (red line, right axis) measured by a
2062 pyranometer, averaged with a moving window of 20 seconds. Grey shading indicates times when the aircraft was
2063 ascending due to poor visibility. b) Profiles of extinction (solid lines) and potential temperature (dashed lines) measured
2064 during flight b708, for the descent (black) and ascent (red). Potential temperature has been averaged over 5s windows.

2065

2066 Figure 17: LIDAR and dropsonde observations from 22 June 2011 morning flight b607 (yellow line in Figure 2c) plotted
2067 along longitude for a) outgoing and b) return. BAe146 LIDAR measurements (coloured boxes) are shown as the range-
2068 corrected LIDAR backscatter signal 355 nm. White regions identify periods of LIDAR data dropouts. Dotted vertical lines
2069 indicate dropsonde locations. Black solid lines mark top of the CBL ([Convective Boundary Layer](#)) and SRL ([Saharan](#)
2070 [Residual Layer](#)). Grey boxes along dropsonde tracks show depth of temperature inversions (change in $^{\circ}\text{C km}^{-1}$ shown
2071 next to box). [Purple line indicates ground level.](#) Dropsonde location ID and release time are indicated above each
2072 dropsonde track.

2073

2074 Figure 18: LNG LIDAR-derived extinction coefficient at 532 nm on 20 June 2011 during flight F21 from 25.0N, 11.5W to
2075 19.0N, 8.7W. Water vapour mixing ratio (WVMR, g kg^{-1}) and wind profiles from four dropsondes are superimposed (black
2076 lines, dropsonde locations indicated by arrows), with WVMR contours drawn by hand using the LIDAR backscatter; away
2077 from the dropsondes these are by necessity subjective and the 7 and 8 g kg^{-1} contours have not been continued west of
2078 21.2N due to a lack of data. Along-track albedo derived from MODIS satellite data is shown in the upper panel, [and](#)
2079 [albedo 1W of the flight track.](#)

2080

2081