

# Cloud Condensation Nuclei Activity, Droplet Growth Kinetics and Hygroscopicity of Biogenic and Anthropogenic Secondary Organic Aerosol (SOA)

D. F. Zhao<sup>1</sup>, A. Buchholz<sup>1, a</sup>, B. Kortner<sup>1</sup>, P. Schlag<sup>1</sup>, F. Rubach<sup>1, b</sup>, H. Fuchs<sup>1</sup>, A. Kiendler-Scharr<sup>1</sup>, R. Tillmann<sup>1</sup>, A. Wahner<sup>1</sup>, Å. K. Watne<sup>2</sup>, M. Hallquist<sup>2</sup>, J. M. Flores<sup>3</sup>, Y. Rudich<sup>3</sup>, K. Kristensen<sup>4</sup>, A. M. K. Hansen<sup>4</sup>, M. Glasius<sup>4</sup>, I. Kourtchev<sup>5</sup>, M. Kalberer<sup>5</sup>, Th. F. Mentel<sup>1</sup>

<sup>1</sup>Institute for Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich, Jülich, D-52425, Germany

<sup>2</sup>Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, SE-41296, Sweden

<sup>3</sup>Department of Earth and Planetary Sciences, Weizmann Institute of Science, Rehovot, 76100, Israel

<sup>4</sup>Department of Chemistry, Aarhus University, 8000 Aarhus, Denmark

<sup>5</sup>Department of Chemistry, University of Cambridge, Cambridge, CB2 1EW, UK

<sup>a</sup>Now at Department of Applied Physics, University of Eastern Finland, 70211 Kuopio, Finland

<sup>b</sup>Now at Max-Planck-Institute for Chemistry, Mainz, Germany

Correspondence to: Th. F. Mentel ([t.mentel@fz-juelich.de](mailto:t.mentel@fz-juelich.de))

## Abstract

Interaction of biogenic volatile organic compounds (VOC) with anthropogenic VOC affects the physicochemical properties of secondary organic aerosol (SOA). We investigated cloud droplet activation (CCN activity), droplet growth kinetics, and hygroscopicity of mixed anthropogenic and biogenic SOA (ABSOA) compared to pure biogenic SOA (BSOA) and pure anthropogenic SOA (ASOA). Selected monoterpenes and aromatics were used as representative precursors of BSOA and ASOA, respectively.

We found that BSOA, ASOA, and ABSOA had similar CCN activity despite the higher oxygen to carbon ratio (O/C) of ASOA compared to BSOA and ABSOA. For individual

reaction systems, CCN activity increased with the degree of oxidation. Yet, when considering all different types of SOA together, the hygroscopicity parameter,  $\kappa_{\text{CCN}}$ , did not correlate with O/C. Droplet growth kinetics of BSOA, ASOA, and ABSOA was comparable to that of  $(\text{NH}_4)_2\text{SO}_4$ , which indicates that there was no delay in the water uptake for these SOA in supersaturated conditions.

In contrast to CCN activity, the hygroscopicity parameter from hygroscopic tandem differential mobility analyzer (HTDMA) measurement,  $\kappa_{\text{HTDMA}}$ , of ASOA was distinctively higher (0.09-0.10) than that of BSOA (0.03-0.06), which was attributed to the higher degree of oxidation of ASOA. The ASOA components in mixed ABSOA enhanced aerosol hygroscopicity. Changing the ASOA fraction by adding BVOC to ASOA or vice versa AVOC to BSOA changed the hygroscopicity of aerosol, in line with the change in the degree of oxidation of aerosol. However, the hygroscopicity of ABSOA cannot be described by a simple linear combination of pure BSOA and ASOA systems. This indicates that additional processes, possibly oligomerization, affected the hygroscopicity.

Closure analysis of CCN and HTDMA data showed  $\kappa_{\text{HTDMA}}$  was lower than  $\kappa_{\text{CCN}}$  by 30%-70%. Better closure was achieved for ASOA compared to BSOA. This discrepancy can be attributed to several reasons. ASOA seemed to have higher solubility in subsaturated conditions and/or higher surface tension at the activation point than that of BSOA.

## 1 **1 Introduction**

2 Secondary organic aerosol (SOA) is an important class of atmospheric aerosol with impacts on  
3 air quality, human health and climate change (Hallquist et al., 2009; Kanakidou et al., 2005;  
4 Jimenez et al., 2009; Zhang et al., 2011; Verma et al., 2014). Despite substantial improvements  
5 in the understanding of SOA formation mechanisms and properties, considerable uncertainties  
6 remain about the regional and global budget of SOA (e.g. Goldstein and Galbally (2007)).  
7 Models often do not correctly predict the ambient concentrations of organic aerosol (OA) (e.g.,  
8 Spracklen et al. (2011), Heald et al. (2005)), and usually the modeled concentrations  
9 underestimate the observed OA concentrations (Spracklen et al., 2011). Recent studies suggested  
10 that interactions between biogenic volatile organic compounds (VOC) and anthropogenic  
11 emissions can enhance SOA formation and often, ambient OA concentrations correlate with  
12 anthropogenic tracers such as CO or isopropyl nitrate (de Gouw et al., 2008; de Gouw et al.,  
13 2005; Weber et al., 2007; Shilling et al., 2013; Xu et al., 2015). However, <sup>14</sup>C carbon isotopic  
14 analysis showed that the SOA material itself is composed mostly of modern carbon, i.e. arises  
15 from biogenic sources, including biomass burning (Weber et al., 2007; Zotter et al., 2014). These  
16 observations suggest that a consideration of an anthropogenic enhancement can reduce the  
17 discrepancies between models and observations (de Gouw et al., 2005; Goldstein et al., 2009;  
18 Hoyle et al., 2011; Worton et al., 2011; Glasius et al., 2011).

19 Anthropogenic VOCs (AVOC), such as aromatic compounds are possibly important factors that  
20 lead to enhanced SOA formation as their oxidation products can interact with biogenic VOC  
21 (BVOC) oxidation products during SOA formation, as shown by several studies (Hoyle et al.,  
22 2011; Emanuelsson et al., 2013; Flores et al., 2014). In a recent study, Emanuelsson et al. (2013)  
23 found that anthropogenic SOA (ASOA) components reduce the volatility of biogenic SOA  
24 (BSOA) in a non-linear way with respect to the ASOA fraction, possibly by oligomerization or a  
25 phase change such as formation of a glassy state (Emanuelsson et al., 2013; Virtanen et al., 2010;  
26 Koop et al., 2011). The reduced volatility in the mixed SOA (anthropogenic-biogenic SOA,  
27 ABSOA) can enhance SOA persistence and concentrations in the atmosphere. Flores et al.  
28 (2014) investigated the optical properties of BSOA, SOA from simultaneous addition of BVOC  
29 and AVOC and SOA from sequential addition of BVOC and AVOC. They found that both SOA  
30 from mixed AVOC and BVOC show an increase of scattering component of the refraction index

1 with aging (increase of the oxygen to carbon ratio (O/C)) and the increase is greater for SOA  
2 from simultaneous addition of BVOC and AVOC than SOA from sequential addition of VOC.

3 Besides the thermochemical and optical properties, cloud droplet activation (cloud condensation  
4 nuclei (CCN) activity) and hygroscopicity are important physicochemical properties that have  
5 critical implications for the impact of aerosol on climate. It is possible that enhanced  
6 oligomerization which happens in the mixed aerosol particles could modify its CCN activity and  
7 hygroscopicity (Xu et al., 2014). Given that CCN activity and hygroscopicity correlate with the  
8 aerosol O/C in many cases and given that ASOA tends to have a higher O/C (Chhabra et al.,  
9 2011; Emanuelsson et al., 2013), it is expected that ASOA components enhance the CCN  
10 activity and hygroscopicity of mixed SOA. However, to the best of our knowledge, the influence  
11 of the interaction of AVOC with BVOC on CCN activity and hygroscopicity of SOA has not  
12 been studied from the literature.

13 Several field studies found a delay in droplet growth kinetics of the aerosol from anthropogenic  
14 origin when compared with the aerosol from biogenic origin (Shantz et al., 2010; Shantz et al.,  
15 2012). ASOA, as an important anthropogenic aerosol, may contribute to this delay. In addition, a  
16 recent laboratory study suggests limited mixing in SOA formed by sequentially mixing a  
17 biogenic precursor ( $\alpha$ -pinene) with an anthropogenic one (toluene) (Loza et al., 2013). It is  
18 possible that limited mixing could affect water uptake by ABSOA and delay droplet growth  
19 when the diffusion of water within aerosol particle was suppressed. Yet, no laboratory studies on  
20 the droplet growth kinetics of ASOA and ABSOA have been found.

21 In this study, we investigated the effect of the interaction of ASOA and BSOA on CCN activity  
22 and hygroscopicity of aerosol. We also studied the kinetics of droplet growth of ASOA, BSOA  
23 and mixed ABSOA.

## 24 **2 Experimental**

### 25 **2.1 Experiment setup and instrumentation**

26 The experiments were conducted in the atmosphere simulation chamber SAPHIR (Simulation of  
27 Atmospheric PHotochemistry In a large Reaction chamber). SAPHIR is a double-wall Teflon  
28 chamber with a volume of 270 m<sup>3</sup>. Details of the chamber have been previously described  
29 (Rohrer et al., 2005; Bohn et al., 2005). The chamber uses natural sunlight for illumination and is

1 equipped with a louvre system which can be used to simulate dark processes. For the  
2 experiments described here, various instrumentations were used to characterize gas phase and the  
3 particulate phase species.

4 Chamber parameters like temperature, relative humidity, flow rate, and photolysis frequencies  
5 were also recorded. The actinic flux and the corresponding photolysis frequencies were provided  
6 from measurements using a spectral radiometer (Bohn et al., 2005; Bohn and Zilken, 2005).

7 The number concentration and size distributions of aerosol were measured by a scanning  
8 mobility particle sizer (SMPS, DMA model 3081/CPC model 3785, TSI Shoreview, USA) and  
9 separate condensation particle counter (CPC, model 3786, TSI) to allow detection of nucleation  
10 particles down to 3 nm.

11 The chemical composition of aerosol was measured by a High-Resolution Time-of-Flight  
12 Aerosol Mass Spectrometer (HR-ToF-AMS, Aerodyne Research Inc., USA). To characterize the  
13 degree of oxidation of aerosol, the oxygen to carbon ratio (O/C) and  $f_{44}$  (fractional contribution  
14 of m/z 44 to the total organics signal) were obtained from the mass spectra. The O/C (corrected  
15 for the minor influence of gaseous components) was derived by the elemental analysis of mass  
16 spectra obtained in the high mass resolution mode (W-mode) of the mass spectrometer as  
17 described by Aiken et al. (2007) and Aiken et al. (2008). The contributions of gas phase CO<sub>2</sub> and  
18 water vapor to m/z 44 and to m/z 18, respectively, were characterized using a CO<sub>2</sub> and H<sub>2</sub>O  
19 analyzer (Picarro, Santa Clara, USA). The values were subtracted to obtain the particle signals  
20 before the elemental analysis (Allan et al., 2004).

21 Droplet activation and droplet growth were measured using a size scanning CCN method as  
22 described previously (Buchholz, 2010; Zhao et al., 2010). This method, also known as Scanning  
23 Mobility CCN analysis (SCMA, (Moore et al., 2010)), has been successfully used in a number of  
24 previous studies (Asa-Awuku et al., 2008; Padro et al., 2007; Engelhart et al., 2008; Asa-Awuku  
25 et al., 2009; Asa-Awuku et al., 2010; Engelhart et al., 2011). The measurement was done by  
26 coupling a differential mobility analyzer (DMA, model 3081, TSI Shoreview, USA) with a cloud  
27 condensation nuclei counter (CCNC, Droplet Measurement Technique, USA) and condensation  
28 particle counter (CPC3786, TSI). Before entering the instruments, the particles were dried using  
29 a silica gel diffusion drier (gradually drying to ~10% RH) with a residence time of around 3s.  
30 Particles then passed through the DMA and the outgoing air was split into two paths connecting

1 to the CCNC and CPC, which measure the CCN and cloud nuclei (CN) concentrations,  
2 respectively. The flow rate of the CCNC is around 0.5 L/min with a sheath to aerosol flow ratio  
3 of 10. The residence time in the CCN column is around 24 s with the time in the final  
4 supersaturation slightly shorter (Lance et al., 2006). The DMA scanned over a size range  
5 between 10 and 450 nm while the supersaturation (SS) remained constant. And four to five  
6 different supersaturations in the range of 0.1%-1.3% were used depending on the particle sizes.  
7 From the measurement, CCN activation fraction over size and the dry activation diameter (or  
8 critical dry diameter,  $D_{crit}$ ) was obtained using a method as described in Buchholz (2010).  
9 Briefly, for each particle size, the CN and CCN concentrations measured were used to calculate  
10 the activation fraction ( $a_f = CCN/CN$ ). Before  $a_f$  was calculated, the measured CN and CCN  
11 concentrations were corrected for multiple charged particles. To separate the single from the  
12 multiple charged particles, the fraction of multiple charged particles was calculated according to  
13 a Boltzmann charge distribution using the measured size distribution (Wiedensohler, 1988).  
14 Then,  $a_f$  was determined for each charge class separately and fitted with a Gaussian error  
15 function (Rose et al., 2008). The dry activation diameter at the set SS is the turning point of this  
16 function.

17 For each SS at least three full scans were performed and the resulting  $D_{crit}$  were averaged. For the  
18 calibration of SS,  $D_{crit}$  of ammonium sulfate at various SS was measured and compared to  
19 theoretical data in the literature (Rose et al., 2008). The set SS was corrected according to the  
20 theoretical data. From the CCN data, the hygroscopicity parameter  $\kappa_{CCN}$  was calculated  
21 according to the one parameter representation of the Köhler equation proposed by Petters and  
22 Kreidenweis (2007). The error bars of  $\kappa_{CCN}$  were estimated using the standard deviation of  $D_{crit}$   
23 from three duplicate scans. A higher hygroscopicity parameter  $\kappa$  indicates a more hygroscopic  
24 material, i.e. cloud droplet activation at lower SS for particles of a given size or at smaller size  
25 for a given SS.

26 The hygroscopic growth of the aerosol was measured using a home-built hygroscopic tandem  
27 differential mobility analyzer (HTDMA). The details of the HTDMA were described previously  
28 (Buchholz, 2010; Zhao et al., 2010). Particles were selected using the first DMA and then were  
29 exposed to a prescribed relative humidity to measure the growth factor. Hygroscopic growth was  
30 measured at different RH. The sizes of the humidified particles were determined by the second

1 DMA which was operated in a scanning mode in combination with a CPC (model 3022A, TSI).  
2 The size selected aerosol flow and the sheath air flow of the second DMA were humidified at  
3 room temperature (25-30°C depending on the surroundings) to almost the same relative humidity  
4 (RH) with the sheath air at slightly higher RH. The second DMA was kept in a thermo-insulated  
5 box which was cooled to 20 °C. Both aerosol and sheath air flow were cooled down to the same  
6 temperature before entering the second DMA, and thus the RH increased to its final value. The  
7 residence time of particles at the final humidity is approximately 30 s before they entered the  
8 SMPS operated with sheath air of the same RH. The hygroscopic growth factor ( $GF$ ) was  
9 calculated as the ratio of the size of the wet particle ( $D_{wet}$ ) to the selected dry size ( $D_{dry}$ ). The  
10 HTDMA was calibrated using ammonium sulfate aerosol by comparing with the theoretical  
11 growth curve (Rose et al., 2008). From hygroscopic growth factor at 90% RH, the hygroscopicity  
12 parameter,  $\kappa_{HTDMA}$ , was calculated according to Petters and Kreidenweis (2007). The error bars  
13 of  $\kappa_{HTDMA}$  were estimated using the standard deviation of the growth factor at  $(90\pm 1)$  % RH of at  
14 least three duplicate scans.

15 SOA samples were collected on PTFE filters at the end of different experiments to obtain  
16 detailed insight into the chemical composition of the aerosol particles. The details of sample  
17 collection and analysis are described in Emanuelsson et al. (2013) and Kristensen and Glasius  
18 (2011). Before the filters, the air passed through an annular denuder coated with XAD-4 resin to  
19 remove gaseous organic species. The filters were extracted and analyzed using a Dionex  
20 Ultimate 3000 HPLC system coupled through an electrospray (ESI) inlet to a q-TOF mass  
21 spectrometer (micro-TOFq, Bruker Daltonics GmbH, Bremen, Germany), which was operated in  
22 both positive and negative mode. Pinonic acid, cis-pinonic acid, terpenylic acid, diaterpenylic acid  
23 acetate (DTAA) and 3-methyl butane tri-carboxylic acid (MBTCA) were quantified using  
24 authentic standards.

25 For SOA from part of the experiments (experiment #B3, AB4, AB6 as in Table 1), samples were  
26 also collected on quartz fiber filters and analyzed by ultra-high resolution mass spectrometry  
27 (UHRMS). In this analysis, the aerosol samples were extracted as described elsewhere  
28 (Kourchev et al., 2013). The extracts were analyzed using an ultrahigh resolution LTQ Orbitrap  
29 Velos mass spectrometer (Thermo Scientific, Bremen, Germany) equipped with a TriVersa  
30 Nanomate robotic nanoflow chip-based ESI source (Advion Biosciences, Ithaca NY, USA). The

1 Orbitrap MS instrument calibration, settings and mass spectral data interpretation are described  
2 in Kourtchev et al. (2014). The mass accuracy of the instrument was below 1.5 ppm and the  
3 instrument mass resolution was 100,000 at  $m/z$  400. The negative ionization mass spectra were  
4 collected in three replicates and in this study, only ions that appeared in all three analytical  
5 replicates were kept for evaluation.

6 The VOC were measured by a high resolution proton transfer reaction-mass spectrometer (HR-  
7 PTR-MS, Ionicon, Innsbruck, Austria) (Jordan et al., 2009) and gas chromatography coupled to a  
8 mass spectrometer (GC-MS, PerkinElmer, Waltham, USA) (Apel et al., 2008; Kaminiski, 2014).

9 The OH concentration was measured directly using laser-induced fluorescence (LIF) (Fuchs et  
10 al., 2012). The OH radicals inside the chamber are mainly formed by the photolysis of HONO  
11 formed via a photolytic process on the chamber walls, and to a minor fraction by  $O_3$  photolysis  
12 (Rohrer et al., 2005). From the OH concentration, the OH dose was calculated and used as a  
13 common abscissa in order to better compare different experiments. The OH dose is the integral  
14 OH concentration over time that gives the accumulated OH concentrations to which gases and  
15 particles were exposed in the course of experiment. One hour exposure to a typical atmospheric  
16 OH concentration of  $2 \times 10^6$  molecules  $cm^{-3}$  is then equivalent to an OH dose of  $7.2 \times 10^9$   
17 molecules  $cm^{-3}$  s.

## 18 **2.2 Experimental procedure**

19 The experimental procedures have been described elsewhere in details (Emanuelsson et al.,  
20 2013; Flores et al., 2014) and only a short description is given here. The chamber was typically  
21 humidified to 60-70% RH in the beginning of the experiment and relative humidity can vary in  
22 the range of 30-70% due to the ambient temperature change and the dilution by the flow to  
23 compensate the sampling loss. In a typical experiment, VOC was added to the chamber and then  
24 the roof was opened to start the photooxidation. In some experiments,  $O_3$  was added. In all the  
25 experiments, particles formed by homogeneous nucleation and no seed was added. In the BSOA  
26 experiments, a monoterpene mixture of  $\alpha$ -pinene and limonene with a molar ratio of 1:1 was  
27 used as the representative BSOA precursors and its photooxidation induced BSOA formation.  
28 Ozone was added to initialize BVOC oxidation and particle formation. In the ASOA  
29 experiments, toluene or xylene was used as the representative ASOA precursors. In the mixed



1 SOA experiments (ABSOA), AVOC and BVOC were added, either simultaneously or  
2 sequentially to investigate the potential effect of adding order.

3 In total, three BSOA experiments (including one using  $\alpha$ -pinene as precursor), seven ASOA  
4 experiments, and six mixing experiments (ABSOA) with biogenic and anthropogenic precursors  
5 were analyzed in detail and the summary for these experiments is given in Table 1. In two  
6 experiments (#AB1, AB2), AVOC was added 6.3 h before the BVOC. In experiments with  
7 sequential VOC addition, the second VOC was added 1-2 h after the SOA mass concentration  
8 generated from the first addition reached its maximum. Accordingly, the time lag was longer  
9 when AVOC was added first due to its lower reactivity. In two experiments (#AB3, AB4),  
10 BVOC was added 2.5 h and 5 h before the AVOC, respectively. In the other two experiments  
11 (#AB5, AB6), BVOC and AVOC were added simultaneously into the chamber. In the ABSOA  
12 experiments, the mass fraction of ASOA in the total aerosol was estimated using a method based  
13 on the aerosol mass yield and VOC consumed as described by Emanuelsson et al. (2013), where  
14 ideal mixing of ASOA and BSOA components was assumed. Assuming the same density for  
15 ASOA and BSOA, the mass fraction of ASOA is equal to its volume fraction.

### 16 **3 Results and discussion**

#### 17 **3.1 Droplet activation of BSOA, ASOA and ABSOA**

##### 18 **3.1.1 Comparison of droplet activation of BSOA, ASOA and ABSOA**

19 Droplet activation of BSOA, ASOA and ABSOA at various supersaturations (SS) was  
20 parameterized by applying the hygroscopicity parameter  $\kappa_{CCN}$  (Petters and Kreidenweis, 2007).  
21 Generally,  $\kappa_{CCN}$  was found to increase slightly with time in the light periods and remained  
22 relatively constant or slightly decreased in the dark periods (overview in Fig. S1). The increase  
23 of  $\kappa_{CCN}$  with time in the light period is attributed to photochemical aging, which leads to  
24 functionalization, i.e. the formation of oxidized compounds with hydroperoxide, hydroxyl,  
25 carbonyl, and carboxylic acid groups (Kroll and Seinfeld, 2008; Zhao et al., 2015b).  
26 Photochemical aging is thought to mainly happen in the gas phase (Donahue et al., 2012) and  
27 affect the particle composition by partitioning, although heterogeneous oxidation and particle  
28 phase oxidation may also play a role. The formation of oxidized compounds with polar

1 functional groups is also supported by the our filter sample analysis of SOA, showing the  
2 formation of a number of organic acids, carbonyl compounds and multi-generation compounds  
3 such as 3-methyl butane tricarboxylic acid (3-MBTCA), a tracer for photochemical ageing  
4 (Zhang et al. (2010), see also Table 3 of Emanuelsson et al. (2013)). This functionalization  
5 increases the SOA average polarity and thus enhances hygroscopicity (Duplissy et al., 2008;  
6 Topping et al., 2005; Suda et al., 2014). We found that  $\kappa_{\text{CCN}}$  depended on SS. Since different SS  
7 correspond to different critical sizes, this dependence indicates a size dependent chemical  
8 composition of aerosol (Zhao et al., 2015a). When comparing CCN activity of different SOA,  
9  $\kappa_{\text{CCN}}$  at similar supersaturation was examined.

10 The droplet activation of BSOA, ABSOA, and ASOA particles represented by  $\kappa_{\text{CCN}}$  is compared  
11 for two SS (0.51% and 0.34%) in Fig. 1. Different experiments required different ranges of SS  
12 because of the achievable particle size ranges. Usually ASOA particles were smaller and thus  
13 higher SS were needed to get a proper range of activation fraction to derive  $D_{\text{crit}}$ . Therefore, we  
14 choose  $\kappa_{\text{CCN}}$  at two SS in the medium range, 0.34% and 0.51%, to compare between experiments  
15 thus maximizing experiments with overlapping SS. In order to compare CCN activity of different  
16 SOA particles from different experiments,  $\kappa_{\text{CCN}}$  is plotted as a function of OH dose (Fig. 1a), the  
17 integral OH concentration over time to which the particles were exposed. The CCN activity of  
18 the three types of aerosol is generally similar at similar OH dose. In addition, with exception of  
19 BSOA,  $\kappa_{\text{CCN}}$  is largely invariant over a wide range of OH doses, equivalent to 1-4 days of OH  
20 exposure in the atmosphere (assuming average OH concentration  $2 \times 10^6$  molecules  $\text{cm}^{-3}$  and 12 h  
21 sunlight per day).

22 Similarity in CCN activity of ASOA, BSOA and ABSOA was also observed in the ABSOA  
23 experiments with sequential VOC addition, independent of the order of addition of AVOC or  
24 BVOC. When BVOC was added after AVOC to the chamber, besides the reaction with OH,  
25 BVOC also reacted with  $\text{O}_3$  formed previously from photochemical reactions of AVOC. The  
26 reaction products from both ozonolysis and OH oxidation helped to convert ASOA rapidly to  
27 ABSOA reaching significant fractions of BSOA (e.g. 70% within 2.5 h, Fig 2a). However,  $\kappa_{\text{CCN}}$   
28 remained largely unchanged within the experimental uncertainty upon the formation of ABSOA  
29 (green and blue markers). Note that  $\kappa_{\text{CCN}}$  should be compared at similar SS because  $\kappa_{\text{CCN}}$   
30 depended on SS and the SS range changed due to the particle size shift after BVOC addition.

1 Such invariance was also observed when BVOC was added first to form BSOA and then AVOC  
2 was added to form ABSOA (Fig. 2b). Therefore we conclude that BSOA, ASOA and ABSOA  
3 have similar CCN activity.

4 For BSOA,  $\kappa_{\text{CCN}}$  increased with photochemical aging at all SS. For ASOA and ABSOA,  $\kappa_{\text{CCN}}$   
5 increased with photochemical aging only at higher SS while  $\kappa_{\text{CCN}}$  remained relatively stable at  
6 low SS (Fig. S2). The increase of  $\kappa_{\text{CCN}}$  with aging is a result of oxidation to form oxygenated  
7 compounds with higher affinity for water. At the same time, the volatility of BSOA, ASOA, and  
8 ABSOA in these experiments decreased with the OH dose (Emanuelsson et al., 2013), indicating  
9 that these hygroscopic components were at the same time less volatile than the initial aerosol  
10 components.

11 Figure 1b shows  $\kappa_{\text{CCN}}$  as a function of O/C. Although the O/C of ASOA and ABSOA was  
12 generally higher than that of BSOA (Fig. 1b), there is no significant difference between the CCN  
13 activity of ASOA, ABSOA, and that of BSOA. This suggests that although ASOA components  
14 increased the O/C of ABSOA and decreased its volatility (Emanuelsson et al., 2013), it did not  
15 enhance its CCN activity. We conclude that the interaction of AVOC with BVOC oxidation  
16 products, such as potential oligomerization during the photochemical oxidation does not affect  
17 CCN activity of the resulting ABSOA. Decreasing volatility and invariant CCN activity at higher  
18 O/C induced by presence of ASOA components indicates different effects of oligomerization and  
19 photochemistry on volatility and CCN activity of aerosol. While both oligomerization and  
20 photochemical aging reduce the volatility (Emanuelsson et al., 2013), they have  
21 counterbalancing effects on CCN activity. Oligomerization increases the molar volume and  
22 decreases the molar concentration of solutes, which results in an increase of water activity and  
23 thus the saturation ratio. Therefore, oligomerization reduces CCN activity, while photochemical  
24 aging enhances the CCN activity by producing more oxygenated compounds via  
25 functionalization (Zhao et al., 2015b). As shown in Eq. A6,  $\kappa$  is affected by the molar volume of  
26 organics and by the van't Hoff factor (the ratio of the actual concentration of molecules or ions  
27 produced when the substance is dissolved to the concentration of the substance if it does not  
28 dissociate). Functionalization does not significantly change the molar volume of organics, but it  
29 increases the van't Hoff factor due to both its interaction of polar functional groups with water  
30 and additional dissociation effects by carboxylic and hydroxyl group (Petters et al., 2009;

1 Tritscher et al., 2011). In this study we did not observe a significant difference in CCN activity  
2 between ABSOA and other SOA, indicating that the effects of possible oligomerization on the  
3 CCN activity of ABSOA are likely compensated by the effects of photochemical aging.

4 For BSOA and ABSOA,  $\kappa_{\text{CCN}}$  generally increases with O/C at a given SS (Fig. S2). For ASOA,  
5 the relationship between  $\kappa_{\text{CCN}}$  and O/C is not so apparent. The  $\kappa_{\text{CCN}}$  values of BSOA (0.11-0.19)  
6 are consistent with  $\kappa_{\text{CCN}}$  values obtained in previous studies (Massoli et al., 2010; Lambe et al.,  
7 2011; Frosch et al., 2011). An increase of  $\kappa_{\text{CCN}}$  with O/C of SOA has been reported by a number  
8 of previous studies (Massoli et al., 2010; Lambe et al., 2011). Yet, other studies reported that  
9  $\kappa_{\text{CCN}}$  of SOA is largely independent of O/C or  $f_{44}$ , another proxy of the degree of oxidation of  
10 organic aerosol (Frosch et al., 2011; Alfarra et al., 2013). The inconclusive results in the  
11 literature may be attributed to the differences in reaction conditions including the VOC type and  
12 concentration,  $\text{NO}_x$  concentration, photochemical aging (OH dose) and O/C range. In this study,  
13 the observation that  $\kappa_{\text{CCN}}$  is invariant on O/C in ASOA may be associated with the relatively  
14 high and narrow range of observed O/C for ASOA. Emanuelsson et al. (2013) proposed that first  
15 generation products of AVOC (mainly carbonyls) may have higher vapor pressure compared to  
16 BVOC products. Therefore, AVOC need a higher OH dose and more oxidation steps before a  
17 significant amount of material starts to condense on the particles. As a result, once SOA started  
18 to form, ASOA had already reached a relatively high O/C due to aging in the gas phase (Fig. 1b).  
19 As a consequence, a further increase of O/C in ASOA is likely slower and less pronounced.

20 Considering all types of SOA investigated here,  $\kappa_{\text{CCN}}$  did not correlate with O/C here (Fig. 1b),  
21 which is in agreement with the findings of Alfarra et al. (2013) that water uptake is independent  
22 of  $f_{44}$  as a metric of the degree of oxidation for multiple biogenic precursors. This is because O/C  
23 affects the solubility of the solute but other effects such as molecular weight, surface tension and  
24 density are also important. Massoli et al. (2010) also showed, the correlation between  $\kappa_{\text{CCN}}$   
25 derived from CCN and O/C is less significant than in individual systems, when considering all  
26 systems together including  $\alpha$ -pinene, trimethylbenzene, and m-xylene. The relationship between  
27  $\kappa_{\text{CCN}}$  and O/C was further compared to the parameterization in the study by Rickards et al.  
28 (2013) (Fig. S5), which was obtained from their experimental data and a number of literature  
29 data and showed very large variability of  $\kappa$  versus O/C.

1 For ASOA systems, particle formation was studied for different aromatic precursors at low NO<sub>x</sub>  
2 (NO<sub>x</sub><1 ppb) and high NO<sub>x</sub> concentrations (10 ppb of NO added). The CCN activity of ASOA  
3 from different precursors including toluene, xylene, and benzene was found to be similar.  
4 Although NO<sub>x</sub> levels affect the overall SOA particle composition and yields (Ng et al., 2007a;  
5 Hildebrandt et al., 2009; Zhang et al., 2014; Presto et al., 2005; Eddingsaas et al., 2012; Ng et al.,  
6 2007b), they had no significant effect on the CCN activity of ASOA comparing the low NO<sub>x</sub>  
7 (NO<sub>x</sub> <1 ppb) with high NO<sub>x</sub> (10 ppb NO added) condition (Fig. S3). The CCN activity of  
8 ASOA generally agreed with the range obtained from a number of previous studies (Massoli et  
9 al., 2010; Prenni et al., 2007; Lambe et al., 2011). Similarly, NO<sub>x</sub> level has been found to not  
10 influence the CCN activity of SOA from α-pinene ozonolysis and subsequent photochemical  
11 aging (Frosch et al., 2011).

### 12 3.1.2 Droplet growth kinetics

13 Droplet growth kinetics was investigated using the method of ‘threshold droplet growth analysis’  
14 (TDGA), which has been used successfully in many field and laboratory studies (Engelhart et al.,  
15 2008; Asa-Awuku et al., 2009; Asa-Awuku et al., 2010; Bougiatioti et al., 2011). In this method,  
16 the droplet growth kinetics was assessed by comparing the droplet sizes from various SOA with  
17 that from ammonium sulfate, which is highly hygroscopic and rapidly grows under  
18 supersaturated conditions. When two particles are exposed to the same SS, they will grow to  
19 droplets of similar size, if their critical SS *and* the mass transfer of water vapor are similar. In  
20 this study, the TDGA method was applied to the size-resolved CCNC data and droplet size was  
21 compared for activated particles with SS<sub>crit</sub> equal to the instrument SS (i.e. particles with a size of  
22 D<sub>crit</sub>) (Asa-Awuku et al., 2009; Asa-Awuku et al., 2010; Bougiatioti et al., 2011).

23 Droplet size as a function of SS for BSOA, ASOA and ABSOA were compared with that of  
24 ammonium sulfate (Fig. 3). The droplet sizes of BSOA, ABSOA or ASOA are similar to those of  
25 ammonium sulfate. This indicates the absence of a kinetic barrier for the water uptake of these  
26 SOA during droplet activation. Our study is in agreement with several previous studies showing  
27 comparable droplet growth kinetics of SOA from monoterpenes with that of ammonium sulfate  
28 (Engelhart et al., 2008; Frosch et al., 2011). For SOA from toluene or xylene, no report on  
29 droplet growth kinetics was found in the literature. The droplet growth of aerosol from  
30 anthropogenic sources in the field containing both organics and ammonium sulfate has been

1 shown to be slower than that of the pure ammonium sulfate, using a static diffusion cloud  
2 condensation chamber (Shantz et al., 2010; Shantz et al., 2012). Based on our study, ASOA from  
3 common aromatics, does not explain such delay and the observations by Shantz and co-workers  
4 must have been caused by other aerosol components (Shantz et al., 2010; Shantz et al., 2012).

5 A recent laboratory study by Loza et al. (2013) suggests limited mixing of different types of  
6 SOA components in the particles formed from BSOA precursor  $\alpha$ -pinene and ASOA precursor  
7 toluene added sequentially. In contrast, the study of Hildebrandt et al. (2011) supported the  
8 pseudo-ideal mixing of BSOA and ASOA components according to the aerosol mass yield.  
9 Based on the droplet growth kinetics of the ABSOA (Fig. 3) observed in this study, no matter  
10 whether ASOA and BSOA components are well mixed in the particles, mixing issues did not  
11 seem to affect the water uptake of particles in supersaturated conditions. Hence, the kinetics of  
12 ABSOA, ASOA and BSOA activation may be regarded to be similar when cloud activation is  
13 considered.

### 14 **3.2 Hygroscopicity of SOA from HTDMA measurement**

15 Figure 4 shows the hygroscopicity ( $\kappa_{\text{HTDMA}}$ ) of BSOA, ASOA, and ABSOA.  $\kappa_{\text{HTDMA}}$  of BSOA  
16 was between 0.03 and 0.06, and increased slightly with OH dose.  $\kappa_{\text{HTDMA}}$  of ASOA was around  
17 0.09-0.1, significantly higher than that of BSOA. Subsaturated hygroscopic growth of ASOA and  
18 BSOA was obviously different whereas their CCN activity was basically similar. The influence  
19 of the SOA types on the hygroscopic growth is different from their influence on CCN activity.  
20 The comparison between the water uptake in the subsaturated conditions from hygroscopic  
21 growth and that in the supersaturated conditions from CCN activity is discussed in the Sect 3.3.  
22  $\kappa_{\text{HTDMA}}$  of ASOA did not change much with the OH dose and  $\kappa_{\text{HTDMA}}$  of ASOA from different  
23 aromatic precursors (toluene, xylene, and benzene) were similar (#A1, 2, 4, 5 in Fig. 4).

24 The higher  $\kappa_{\text{HTDMA}}$  of ASOA can be related to the chemical composition represented by O/C.  
25 The O/C of BSOA was about 0.3-0.5, distinctively lower than that of ASOA (0.7-0.8). O/C has  
26 been found to correlate with  $\kappa_{\text{HTDMA}}$  for various SOA systems (Jimenez et al., 2009; Massoli et  
27 al., 2010; Duplissy et al., 2008; Duplissy et al., 2011; Lambe et al., 2011). The same arguments  
28 as for CCN activity apply here. ASOA reached much higher O/C at the same OH dose compared  
29 to BSOA (Fig. S4) (Emanuelsson et al., 2013) because first generation products of AVOC have a

1 smaller carbon number and higher vapor pressure compared to BVOC. Thus, first generation  
2 products of AVOC need thus more oxidation steps before starting to condense significantly on  
3 particles as discussed in Sect 3.1. In addition, constituents of aromatic ASOA generally have  
4 lower molecular weights than BSOA molecules here. We can expect that ASOA components  
5 have on average a lower molar volume and thus ASOA has higher  $\kappa_{\text{HTDMA}}$  assuming all other  
6 parameters are the same for ASOA and BSOA (c.f. Appendix A), since  $\kappa$  is by definition the  
7 ratio of the molar volume of water to the average molar volume of the solute.

8 The  $\kappa_{\text{HTDMA}}$  observed for ABSOA formed with AVOC and BVOC added in various orders were  
9 in the range 0.03 - 0.06, close to or slightly higher than the  $\kappa_{\text{HTDMA}}$  of BSOA. The observed O/C  
10 range of ABSOA was slightly higher than the O/C range of BSOA, but partly overlapping. Since  
11 ASOA had much higher  $\kappa_{\text{HTDMA}}$  than BSOA, ASOA enhanced the  $\kappa_{\text{HTDMA}}$  of ABSOA and the  
12 extent of enhancement depended on its fraction.

13 The enhanced  $\kappa_{\text{HTDMA}}$  due to ASOA fraction was also reflected clearly in the ABSOA  
14 experiments when AVOC and BVOC were sequentially added. In the experiment when ASOA  
15 was formed first, the SOA showed higher  $\kappa_{\text{HTDMA}}$ , around 0.09 (Fig. 5a). When BVOC was  
16 added to the system, ASOA was converted to ABSOA with a significant BSOA fraction (e.g.,  
17 70% within 2.5 h) and  $\kappa_{\text{HTDMA}}$  decreased from 0.09 to 0.04 with the formation of the BSOA  
18 components. Meanwhile the degree of oxidation of ABSOA decreased significantly as indicated  
19 by the decrease of  $f_{44}$  (from 0.23 to 0.1). For the experiments when AVOC was added to BSOA  
20 system, an effect was recognizable, however  $\kappa_{\text{HTDMA}}$  only increased slightly (Fig. 5b). This was  
21 because the reaction of aromatics with OH and SOA formation was slow and the fraction of  
22 ASOA did not exceed 10%. Accordingly, only a slight increase of  $f_{44}$  was observed (from 0.10 to  
23 0.12) even with concurrent aging, consistent with the minor effects of the ASOA component on  
24 the chemical composition of ABSOA due to its low fraction.

25 Since ASOA has higher  $\kappa_{\text{HTDMA}}$ , mixing of ASOA with BSOA may directly enhance  $\kappa_{\text{HTDMA}}$  due  
26 to a simple linear mixing. In order to understand the role of ASOA components in enhancing  
27  $\kappa_{\text{HTDMA}}$  of ABSOA,  $\kappa_{\text{HTDMA}}$  was also examined as a function of the ASOA fraction (as shown in  
28 Fig. 6). In the ABSOA experiment, two main factors affect the hygroscopicity: aging of the  
29 BSOA components and increasing fraction of ASOA components. Therefore, the OH dose is  
30 examined to account for the effect of aging. In Fig. 6, the dashed lines connect the  $\kappa_{\text{HTDMA}}$  of

1 pure BSOA and pure ASOA of the same OH dose at a series of OH doses varying from fresh to  
2 aged SOA. Such a graph can help to detect whether the  $\kappa_{\text{HTDMA}}$  of ABSOA can be described by a  
3 simple linear mixing of the  $\kappa_{\text{HTDMA}}$  of BSOA and ASOA components with respect to their  
4 volume fraction or where non-linear response of  $\kappa_{\text{HTDMA}}$  is effective. For each OH dose, a dashed  
5 line connects pure BSOA and pure ASOA at the given OH dose (represented by the size of  
6 marker). This line defines the expected  $\kappa_{\text{HTDMA}}$  range of ABSOA with varying ASOA fraction at  
7 given OH dose. If the  $\kappa_{\text{HTDMA}}$  of ABSOA can be described by a linear combination of the  
8  $\kappa_{\text{HTDMA}}$  of pure ASOA and BSOA components in respect of their volume fraction, the  $\kappa_{\text{HTDMA}}$   
9 data point should be on the line corresponding to the given OH dose of that data point and should  
10 increase with ASOA fraction along the line due to the higher  $\kappa_{\text{HTDMA}}$  of ASOA. If a succession  
11 of points from one experiment cross dashed lines (i.e. points beyond the line corresponding to  
12 the OH dose of those points) would indicate  $\kappa_{\text{HTDMA}}$  cannot be explained by a linear  
13 combination.

14 For ABSOA, several cases with non-linear effects were observed. For the ABSOA in the  
15 experiment #AB1 and #AB2 where AVOC was added first,  $\kappa_{\text{HTDMA}}$  were significantly lower than  
16 the values from the linear combination of pure ASOA and BSOA components (much below the  
17 lines corresponding to the OH doses of the data points). For ABSOA in the experiment #AB5  
18 when AVOC and BVOC were added together,  $\kappa_{\text{HTDMA}}$  did not change significantly in spite of a  
19 significant increase of ASOA fraction. In the beginning,  $\kappa_{\text{HTDMA}}$  of ABSOA was higher than the  
20 value from a linear combination, whereas in the end,  $\kappa_{\text{HTDMA}}$  was lower than the value from a  
21 linear combination of pure systems. These cases indicate that the observed  $\kappa_{\text{HTDMA}}$  of ABSOA  
22 cannot be explained by a simple linear combination of pure ASOA and BSOA systems. There  
23 seems to be some additional effects such as oligomerization, which altered the chemical  
24 composition of ABSOA and thus affected  $\kappa_{\text{HTDMA}}$ . Moreover, for the ABSOA in the experiment  
25 #AB5,  $\kappa_{\text{HTDMA}}$  remained largely unchanged in spite of continuous oxidation and increase of  
26 ASOA fraction, both enhancing hygroscopicity. This further indicates that the possible  
27 oligomerization, which should decrease the Raoult term and thus hygroscopicity, compensates  
28 the effect of photochemical aging which enhances hygroscopicity, consistent with the discussion  
29 in Sect. 3.1.



1 Morphological effects can also play a role. If the ASOA and BSOA components were not well  
2 mixed in the aerosol particles in the experiments with sequential VOC additions, there would be  
3 more BSOA components on SOA particle surface in the experiments #AB1 and #AB2. This  
4 could affect the  $\kappa_{\text{HTDMA}}$  and contributed to the non-linear effect. But this cannot explain the non-  
5 linear effect in the experiment with VOCs added simultaneously. In addition, if ABSOA forms a  
6 glassy state, the lower diffusivity in the particle may hinder water uptake thus decreasing  
7  $\kappa_{\text{HTDMA}}$ . Although similar growth kinetics of SOA to  $(\text{NH}_4)_2\text{SO}_4$  was observed in supersaturated  
8 conditions, in subsaturated conditions the water diffusivity in the particle may be limited thus  
9 limiting water uptake.

10 The ABSOA filter samples from experiment #AB4 and #AB6 were extracted and analyzed for  
11 oligomers. We observed the oligomer formation in these samples (Fig S6). Oligomer in SOA has  
12 been found by a number of studies (Gao et al., 2004; Noziere et al., 2015; Tolocka et al., 2004;  
13 Kalberer et al., 2004; Kourtschev et al., 2015; Kourtschev et al., 2014). Small multi-functional  
14 products from aromatics oxidation (Hamilton et al., 2005; Jenkin et al., 2003; Johnson et al.,  
15 2005) may promote oligomerization between ASOA and BSOA components. But we did not find  
16 indications that ABSOA contained more dimers compared to BSOA. This can be attributed to  
17 the low ASOA fraction  $\leq 5\%$  in experiments #AB4 and #AB6 (estimated using the method as in  
18 Emanuelsson et al. (2013)). The low ASOA fraction was caused by the low OH concentration  
19 and low chemical turnover of the aromatics in these experiments because high concentrations of  
20 VOC were used in order to generate enough particle mass for optical measurement (Flores et al.,  
21 2014). The low fraction of ASOA resulted in little oligomer formation by the interaction between  
22 the ASOA components and BSOA components. In the future experiments, conditions that can  
23 form comparable fractions of both ASOA and BSOA, thus favorable to ASOA and BSOA  
24 interaction such as oligomerization are preferred. Therefore, relatively higher AVOC  
25 concentration and higher OH concentration (as in experiment #AB5) are desirable.

### 26 **3.3 Closure between the hygroscopicity parameter from CCN and HTDMA**

27 The hygroscopicity parameter  $\kappa$  was obtained from CCN and HTDMA measurements in  
28 supersaturated and subsaturated conditions, respectively. For all SOA types studied here, there is  
29 a significant gap between  $\kappa_{\text{HTDMA}}$  and  $\kappa_{\text{CCN}}$ .  $\kappa_{\text{HTDMA}}$  was significantly lower than  $\kappa_{\text{CCN}}$  with

1  $\kappa_{\text{HTDMA}}/\kappa_{\text{CCN}}$  around 0.3-0.7 (Fig. 7). The ratio of  $\kappa_{\text{HTDMA}}/\kappa_{\text{CCN}}$  for BSOA and ABSOA was  
2 lower than that of ASOA, which is closer to 1. This means that there is a smaller gap between  
3  $\kappa_{\text{HTDMA}}$  and  $\kappa_{\text{CCN}}$  for ASOA compared to BSOA and ABSOA.

4 The closure between  $\kappa_{\text{CCN}}$  and  $\kappa_{\text{HTDMA}}$  of SOA has been studied and discussed by a number of  
5 previous studies with varying results (Dusek et al., 2011; Alfarra et al., 2013; Good et al., 2010;  
6 Duplissy et al., 2008; Juranyi et al., 2009; Prenni et al., 2007; Massoli et al., 2010; Hansen et al.,  
7 2015; Wex et al., 2009; Whitehead et al., 2014). The discrepancy between  $\kappa_{\text{CCN}}$  and  $\kappa_{\text{HTDMA}}$   
8 found here can be attributed to several possible reasons as discussed in the literature (Prenni et  
9 al., 2007; Massoli et al., 2010; Frosch et al., 2011; Good et al., 2010; Alfarra et al., 2013; Wex et  
10 al., 2009; Whitehead et al., 2014; Petters et al., 2009; Dusek et al., 2011). An important reason is  
11 the presence of slightly soluble compounds. These compounds only dissolve partly in the  
12 subsaturated condition while they can dissolve completely in the supersaturated conditions due to  
13 more water available. Therefore,  $\kappa$  is underestimated to varied extent in the subsaturated  
14 condition. ASOA components here seemed to have higher solubility compared to BSOA  
15 components, and thus the gap between  $\kappa_{\text{CCN}}$  and  $\kappa_{\text{HTDMA}}$  was smaller than that of BSOA.

16 Surface tension can also play a role in this discrepancy.  $\kappa$  was calculated using the surface  
17 tension of pure water ( $0.073 \text{ N m}^{-1}$ ). If the surface tension of the droplets is lower than that of  
18 water,  $\kappa$  would be overestimated. While  $\kappa_{\text{HTDMA}}$  is not so sensitive to the change of surface  
19 tension,  $\kappa_{\text{CCN}}$  is more sensitive to surface tension at the point of activation according to the  
20 Köhler equation (Petters and Kreidenweis, 2007). The surface tension under the subsaturated  
21 conditions is assumed to be lower than that under the supersaturated conditions due to the more  
22 concentrated organics in the droplets under the subsaturated condition (Prisle et al., 2008). If the  
23 surface tension effect for BSOA would be larger than for ASOA, i.e. lower surface tension in  
24 supersaturated conditions, this would lead to a higher  $\kappa_{\text{CCN}}$  for BSOA. While the surface tension  
25 effect in subsaturated conditions is small, i.e.  $\kappa_{\text{HTDMA}}$  is relatively constant, higher  $\kappa_{\text{CCN}}$  of  
26 BSOA results in a larger discrepancy between  $\kappa_{\text{HTDMA}}$  and  $\kappa_{\text{CCN}}$ . However, surface active  
27 organics can be enriched at the surface to such a high extent that the Raoult term is significantly  
28 diminished (Prisle et al. 2008). This difference can compensate the overestimation by using the  
29 surface tension of water and the compensating effects make using surface tension of water be a  
30 reasonable choice.

1 Furthermore, the  $\kappa$ -Köhler model does not account explicitly for changes in non-ideality of a  
2 solution as a function of saturation ratio i.e. water activity. Instead,  $\kappa$  might be not constant  
3 through the whole range of water activity (Petters and Kreidenweis, 2007). In addition, different  
4 aerosol behaviors such as evaporation and condensation of organics in HTDMA and CCN  
5 instrument and limited diffusivity of water in case of glassy particles can contribute to the  
6 discrepancy (Whitehead et al., 2014; Asa-Awuku et al., 2009; Irwin et al., 2010; Topping and  
7 McFiggans, 2012; Topping et al., 2013; Duplissy et al., 2009).

#### 8 **4 Conclusions and implications**

9 We investigated the droplet activation, droplet growth kinetics and hygroscopicity of the BSOA,  
10 ASOA, and ABSOA formed from monoterpenes and aromatics used as representative BVOC  
11 and AVOC.

12 We found that BSOA, ASOA and ABSOA had similar CCN activity although ASOA had a  
13 higher O/C. Adding BVOC after ASOA formation, or adding AVOC after BSOA formation did  
14 not significantly change the CCN activity of SOA. The similar CCN activity of BSOA, ASOA  
15 and ABSOA suggests that ASOA components and the interaction of ASOA with BSOA did not  
16 significantly change the CCN activity of SOA. This was likely due to compensating effects of  
17 potential oligomerization (reducing effect) and photochemical aging (enhancing effect) on CCN  
18 activity.

19  $\kappa_{\text{CCN}}$  generally increased slightly with photochemical aging, using OH dose as a proxy of  
20 photochemical aging and increased with O/C of aerosol for individual reaction systems. But  
21 when taking all the SOA types into account,  $\kappa_{\text{CCN}}$  did not correlate with O/C.

22 Analysis of the droplet growth kinetics shows that the droplet sizes from BSOA, ASOA, and  
23 ABSOA in supersaturated conditions were similar to those obtained with ammonium sulfate,  
24 indicating that none of these SOA has a kinetic barrier for water uptake. The fast water uptake of  
25 ASOA indicates that ASOA formed by aromatic precursors is not responsible for the droplet  
26 growth delay found in field studies (Shantz et al., 2010; Shantz et al., 2012). This finding also  
27 suggests that potentially limited mixing between BSOA and ASOA reported in the literature  
28 does not hinder the water uptake in supersaturated conditions.

1 In contrast to CCN activity, the hygroscopicity of ASOA was distinctively higher than that of  
2 BSOA. The higher hygroscopicity was related to the higher O/C of ASOA compared to BSOA.  
3 Therefore, the ASOA component in ABSOA enhanced the hygroscopicity of aerosol and the  
4 extent depended on the ASOA fraction. Adding BVOC to ASOA or AVOC to BSOA changed  
5 the aerosol hygroscopicity, which was consistent with co-occurring changes in the ASOA  
6 fraction and the degree of oxidation of the aerosol represented by  $f_{44}$ . However, the  
7 hygroscopicity of ABSOA cannot be described by a linear combination of pure BSOA and  
8 ASOA systems. This indicates that additional processes such as oligomerization suppressed the  
9 hygroscopicity, which is in agreement with the interpretations for CCN activity.

10 Comparing hygroscopicity parameter  $\kappa$  obtained from CCN and HTDMA measurement shows a  
11 discrepancy between  $\kappa$  from the two methods.  $\kappa_{\text{HTDMA}}$  was significantly lower than  $\kappa_{\text{CCN}}$  for all  
12 SOA types studied here, by 30%~70%. This discrepancy could not be resolved but can be  
13 attributed to the presence of slightly soluble materials, possible surface tension effect, or non-  
14 ideality of solutions and different behaviors of aerosol in the instruments. Better closure between  
15 HTDMA and CCN was found for ASOA than BSOA and ABSOA. ASOA seemed to have  
16 higher solubility in the subsaturated condition and/or lower surface tension at the activation point  
17 compared to BSOA.

18 This study has important implications for assessing the impact of SOA formed by the interaction  
19 of biogenic VOC with anthropogenic VOC emissions on the radiative forcing and climate. Since  
20 the interaction of AVOC with BVOC reduces the volatility (Emanuelsson et al., 2013), it  
21 prolongs particle persistence, which further enhances the particle concentration. Yet, based on  
22 this study, the CCN activity is not significantly affected. Therefore, models to assess the climatic  
23 effects of SOA formed through the interaction of biogenic VOC with anthropogenic VOC  
24 emissions could use single series of hygroscopicity parameter  $\kappa_{\text{CCN}}$  for BSOA, ASOA, and  
25 ABSOA to predict CCN concentration. However, significant mixing of ASOA and BSOA  
26 components can change the hygroscopic growth factor of the particles, which further affects the  
27 optical properties of SOA.

28 Comparing emission rates of aromatic compounds and isoprenoids (Lamarque et al., 2010;  
29 Guenther et al., 2012) and considering the turnover rates with OH and O<sub>3</sub> suggest that ABSOA  
30 should be dominated by BSOA components in most cases, as in most of our experiments. Thus

1 globally droplet activation and hygroscopic growth may be determined by BSOA. On the  
 2 regional scale, when an air mass from regions influenced by anthropogenic emissions (e.g. an  
 3 urban region) transports to regions influenced by biogenic emissions (e.g. a forest region), the  
 4 physicochemical properties of the mixed SOA formed will likely shift to be BSOA-dominated  
 5 due to the fast turnover of BVOC, i.e., decreasing hygroscopic growth compared with ASOA. In  
 6 contrast, when an air mass from regions influenced by biogenic emissions transports to regions  
 7 with anthropogenic emissions, the SOA properties will likely remain those of BSOA due to the  
 8 slow turnover of AVOC.

9

## 10 **Appendix A: Equations related to $\kappa$ -Köhler theory**

11 Based on  $\kappa$ -Köhler theory (Petters and Kreidenweis, 2007), the following equations are tenable,

$$12 \quad \frac{1}{a_w} = 1 + \kappa \frac{V_s}{V_w} \quad (A1)$$

$$13 \quad \frac{1}{a_w} = 1 + i \frac{n_s}{n_w} \quad (A2)$$

14 where  $a_w$  is water activity, and  $V_s$  and  $V_w$  are the volume of solute and water, respectively.  $n_s$  and  
 15  $n_w$  are the amount of solute and water in moles, respectively.  $i$  is the van't Hoff factor.  $i$  is the  
 16 ratio of the measured value of a colligative property to the expected value if the solute is a  
 17 nonelectrolyte (Petrucci and Herring, 2007; McDonald, 1953; Low, 1969). It is the ratio of the  
 18 actual concentration of molecules or ions produced when a substance is dissolved to the  
 19 concentration of the substance if it does not dissociate.

20 From Eq. (A1) and Eq. (A2) one can get

$$21 \quad \kappa = i \frac{V_w/n_w}{V_s/n_s} \quad (A3)$$

22 In addition, the amount of water and solute is described by

$$23 \quad n_w = V_w \rho_w / M_w \quad (A4)$$

$$24 \quad n_s = V_s \rho_s / M_s \quad (A5)$$

25  $\rho_s$  and  $\rho_w$  are the density of solute and water, and  $M_s$  and  $M_w$  are the molecular weight of solute  
 26 and water, respectively.

1 Substituting Eq. A4 and A5 into Eq. A3 yields

$$2 \quad \kappa = i \frac{M_w/\rho_w}{M_s/\rho_s} \quad (\text{A6}).$$

3 Assuming  $i$  is constant,  $\kappa$  is the ratio of molar volume of water to the average molar volume of  
4 the solute. Thus, a compound with lower molecular weight at similar density has lower molar  
5 volume and tends to have higher  $\kappa$  provided that other factors are constant.

6

## 7 **Acknowledgements**

8 This study was supported by the EUROCHAMP2 (Integration of European Simulation  
9 Chambers for Investigating Atmospheric Processes) – EC 7th framework. We thank the SAPHIR  
10 team, especially Franz Rohrer, Rolf Häsel, Birger Bohn, Martin Kaminski, Sascha Nehr,  
11 Sebastian Schmitt, Anna Lutz, Eva Emanuelsson, Ismail-Hakki Acir for providing helpful data  
12 and supporting our measurements. Marianne Glasius thanks the funding support from NordForsk  
13 through the Nordic Centre of Excellence Cryosphere-Atmosphere Interactions in a Changing  
14 Arctic Climate (CRAICC) and the VILLUM Foundation. We thank two anonymous reviewers  
15 for the constructive comments.

16

## 1 **References**

- 2 Aiken, A. C., DeCarlo, P. F., and Jimenez, J. L.: Elemental analysis of organic species with  
3 electron ionization high-resolution mass spectrometry, *Anal. Chem.*, 79, 8350-8358,  
4 10.1021/ac071150w, 2007.
- 5 Aiken, A. C., Decarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S.,  
6 Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A.,  
7 Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S.  
8 H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and  
9 OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-  
10 flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42, 4478-4485, 10.1021/es703009q,  
11 2008.
- 12 Alfarra, M. R., Good, N., Wyche, K. P., Hamilton, J. E., Monks, P. S., Lewis, A. C., and  
13 McFiggans, G.: Water uptake is independent of the inferred composition of secondary aerosols  
14 derived from multiple biogenic VOCs, *Atmos. Chem. Phys.*, 13, 11769-11789, 10.5194/acp-13-  
15 11769-2013, 2013.
- 16 Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A.  
17 M., Drewnick, F., Onasch, T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A  
18 generalised method for the extraction of chemically resolved mass spectra from aerodyne aerosol  
19 mass spectrometer data, *J. Aerosol Sci.*, 35, 909-922, 10.1016/j.jaerosci.2004.02.007, 2004.
- 20 Apel, E. C., Brauers, T., Koppmann, R., Bandowe, B., Bossmeyer, J., Holzke, C., Tillmann, R.,  
21 Wahner, A., Wegener, R., Brunner, A., Jocher, M., Ruuskanen, T., Spirig, C., Steigner, D.,  
22 Steinbrecher, R., Alvarez, E. G., Muller, K., Burrows, J. P., Schade, G., Solomon, S. J.,  
23 Ladstatter-Weissenmayer, A., Simmonds, P., Young, D., Hopkins, J. R., Lewis, A. C., Legreid,  
24 G., Reimann, S., Hansel, A., Wisthaler, A., Blake, R. S., Ellis, A. M., Monks, P. S., and Wyche,  
25 K. P.: Intercomparison of oxygenated volatile organic compound measurements at the SAPHIR  
26 atmosphere simulation chamber, *J. Geophys. Res.-Atmos.*, 113, D20307, 10.1029/2008jd009865,  
27 2008.
- 28 Asa-Awuku, A., Sullivan, A. P., Hennigan, C. J., Weber, R. J., and Nenes, A.: Investigation of  
29 molar volume and surfactant characteristics of water-soluble organic compounds in biomass  
30 burning aerosol, *Atmos. Chem. Phys.*, 8, 799-812, 2008.
- 31 Asa-Awuku, A., Engelhart, G. J., Lee, B. H., Pandis, S. N., and Nenes, A.: Relating CCN  
32 activity, volatility, and droplet growth kinetics of beta-caryophyllene secondary organic aerosol,  
33 *Atmos. Chem. Phys.*, 9, 795-812, 2009.
- 34 Asa-Awuku, A., Nenes, A., Gao, S., Flagan, R. C., and Seinfeld, J. H.: Water-soluble SOA from  
35 Alkene ozonolysis: composition and droplet activation kinetics inferences from analysis of CCN  
36 activity, *Atmos. Chem. Phys.*, 10, 1585-1597, 2010.
- 37 Bohn, B., Rohrer, F., Brauers, T., and Wahner, A.: Actinometric measurements of NO<sub>2</sub>  
38 photolysis frequencies in the atmosphere simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 5,  
39 493-503, 2005.
- 40 Bohn, B., and Zilken, H.: Model-aided radiometric determination of photolysis frequencies in a  
41 sunlit atmosphere simulation chamber, *Atmos. Chem. Phys.*, 5, 191-206, 2005.

1 Bougiatioti, A., Nenes, A., Fountoukis, C., Kalivitis, N., Pandis, S. N., and Mihalopoulos, N.:  
2 Size-resolved CCN distributions and activation kinetics of aged continental and marine aerosol,  
3 *Atmos. Chem. Phys.*, 11, 8791-8808, 10.5194/acp-11-8791-2011, 2011.

4 Buchholz, A.: Secondary organic aerosols: Chemical aging, hygroscopicity, and cloud droplet  
5 activation, Ph.D., Department of Chemistry, University of Cologne, Cologne, 2010.

6 Chhabra, P. S., Ng, N. L., Canagaratna, M. R., Corrigan, A. L., Russell, L. M., Worsnop, D. R.,  
7 Flagan, R. C., and Seinfeld, J. H.: Elemental composition and oxidation of chamber organic  
8 aerosol, *Atmos. Chem. Phys.*, 11, 8827-8845, 10.5194/acp-11-8827-2011, 2011.

9 de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M.,  
10 Fehsenfeld, F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C.,  
11 Marchewka, M., Bertman, S. B., and Bates, T. S.: Budget of organic carbon in a polluted  
12 atmosphere: Results from the New England Air Quality Study in 2002, *J. Geophys. Res.-Atmos.*,  
13 110, D16305, 10.1029/2004jd005623, 2005.

14 de Gouw, J. A., Brock, C. A., Atlas, E. L., Bates, T. S., Fehsenfeld, F. C., Goldan, P. D.,  
15 Holloway, J. S., Kuster, W. C., Lerner, B. M., Matthew, B. M., Middlebrook, A. M., Onasch, T.  
16 B., Peltier, R. E., Quinn, P. K., Senff, C. J., Stohl, A., Sullivan, A. P., Trainer, M., Warneke, C.,  
17 Weber, R. J., and Williams, E. J.: Sources of particulate matter in the northeastern United States  
18 in summer: 1. Direct emissions and secondary formation of organic matter in urban plumes, *J.*  
19 *Geophys. Res.-Atmos.*, 113, D08301, 10.1029/2007jd009243, 2008.

20 Donahue, N. M., Henry, K. M., Mentel, T. F., Kiendler-Scharr, A., Spindler, C., Bohn, B.,  
21 Brauers, T., Dorn, H. P., Fuchs, H., Tillmann, R., Wahner, A., Saathoff, H., Naumann, K. H.,  
22 Mohler, O., Leisner, T., Muller, L., Reinnig, M. C., Hoffmann, T., Salo, K., Hallquist, M.,  
23 Frosch, M., Bilde, M., Tritscher, T., Barmet, P., Praplan, A. P., DeCarlo, P. F., Dommen, J.,  
24 Prevot, A. S. H., and Baltensperger, U.: Aging of biogenic secondary organic aerosol via gas-  
25 phase OH radical reactions, *Proc. Nat. Acad. Sci. U.S.A.*, 109, 13503-13508,  
26 10.1073/pnas.1115186109, 2012.

27 Duplissy, J., Gysel, M., Alfarra, M. R., Dommen, J., Metzger, A., Prevot, A. S. H., Weingartner,  
28 E., Laaksonen, A., Raatikainen, T., Good, N., Turner, S. F., McFiggans, G., and Baltensperger,  
29 U.: Cloud forming potential of secondary organic aerosol under near atmospheric conditions,  
30 *Geophys. Res. Lett.*, 35, L03818, 10.1029/2007gl031075, 2008.

31 Duplissy, J., Gysel, M., Sjogren, S., Meyer, N., Good, N., Kammermann, L., Michaud, V.,  
32 Weigel, R., dos Santos, S. M., Gruening, C., Villani, P., Laj, P., Sellegri, K., Metzger, A.,  
33 McFiggans, G. B., Wehrle, G., Richter, R., Dommen, J., Ristovski, Z., Baltensperger, U., and  
34 Weingartner, E.: Intercomparison study of six HTDMAs: results and recommendations,  
35 *Atmospheric Measurement Techniques*, 2, 363-378, 2009.

36 Duplissy, J., DeCarlo, P. F., Dommen, J., Alfarra, M. R., Metzger, A., Barmadimos, I., Prevot,  
37 A. S. H., Weingartner, E., Tritscher, T., Gysel, M., Aiken, A. C., Jimenez, J. L., Canagaratna, M.  
38 R., Worsnop, D. R., Collins, D. R., Tomlinson, J., and Baltensperger, U.: Relating  
39 hygroscopicity and composition of organic aerosol particulate matter, *Atmos. Chem. Phys.*, 11,  
40 1155-1165, 10.5194/acp-11-1155-2011, 2011.

41 Dusek, U., Frank, G. P., Massling, A., Zeromskiene, K., Iinuma, Y., Schmid, O., Helas, G.,  
42 Hennig, T., Wiedensohler, A., and Andreae, M. O.: Water uptake by biomass burning aerosol at



1 sub- and supersaturated conditions: closure studies and implications for the role of organics,  
2 *Atmos. Chem. Phys.*, 11, 9519-9532, 10.5194/acp-11-9519-2011, 2011.

3 Eddingsaas, N. C., Loza, C. L., Yee, L. D., Chan, M., Schilling, K. A., Chhabra, P. S., Seinfeld,  
4 J. H., and Wennberg, P. O.: alpha-pinene photooxidation under controlled chemical conditions -  
5 Part 2: SOA yield and composition in low- and high-NO<sub>x</sub> environments, *Atmos. Chem. Phys.*,  
6 12, 7413-7427, 10.5194/acp-12-7413-2012, 2012.

7 Emanuelsson, E. U., Hallquist, M., Kristensen, K., Glasius, M., Bohn, B., Fuchs, H., Kammer,  
8 B., Kiendler-Scharr, A., Nehr, S., Rubach, F., Tillmann, R., Wahner, A., Wu, H. C., and Mentel,  
9 T. F.: Formation of anthropogenic secondary organic aerosol (SOA) and its influence on  
10 biogenic SOA properties, *Atmos. Chem. Phys.*, 13, 2837-2855, 10.5194/acp-13-2837-2013,  
11 2013.

12 Engelhart, G. J., Asa-Awuku, A., Nenes, A., and Pandis, S. N.: CCN activity and droplet growth  
13 kinetics of fresh and aged monoterpene secondary organic aerosol, *Atmos. Chem. Phys.*, 8,  
14 3937-3949, 2008.

15 Engelhart, G. J., Moore, R. H., Nenes, A., and Pandis, S. N.: Cloud condensation nuclei activity  
16 of isoprene secondary organic aerosol, *J. Geophys. Res.-Atmos.*, 116, D02207,  
17 10.1029/2010jd014706, 2011.

18 Flores, J. M., Zhao, D. F., Segev, L., Schlag, P., Kiendler-Scharr, A., Fuchs, H., Watne, A. K.,  
19 Bluvshstein, N., Mentel, T. F., Hallquist, M., and Rudich, Y.: Evolution of the complex refractive  
20 index in the UV spectral region in ageing secondary organic aerosol, *Atmos. Chem. Phys.*, 14,  
21 5793-5806, 10.5194/acp-14-5793-2014, 2014.

22 Frosch, M., Bilde, M., DeCarlo, P. F., Juranyi, Z., Tritscher, T., Dommen, J., Donahue, N. M.,  
23 Gysel, M., Weingartner, E., and Baltensperger, U.: Relating cloud condensation nuclei activity  
24 and oxidation level of alpha-pinene secondary organic aerosols, *J. Geophys. Res.-Atmos.*, 116,  
25 D22212, 10.1029/2011jd016401, 2011.

26 Fuchs, H., Dorn, H. P., Bachner, M., Bohn, B., Brauers, T., Gomm, S., Hofzumahaus, A.,  
27 Holland, F., Nehr, S., Rohrer, F., Tillmann, R., and Wahner, A.: Comparison of OH  
28 concentration measurements by DOAS and LIF during SAPHIR chamber experiments at high  
29 OH reactivity and low NO concentration, *Atmospheric Measurement Techniques*, 5, 1611-1626,  
30 10.5194/amt-5-1611-2012, 2012.

31 Gao, S., Keywood, M., Ng, N. L., Surratt, J., Varutbangkul, V., Bahreini, R., Flagan, R. C., and  
32 Seinfeld, J. H.: Low-molecular-weight and oligomeric components in secondary organic aerosol  
33 from the ozonolysis of cycloalkenes and alpha-pinene, *J. Phys. Chem. A* 108, 10147-10164,  
34 10.1021/jp047466e, 2004.

35 Glasius, M., la Cour, A., and Lohse, C.: Fossil and nonfossil carbon in fine particulate matter: A  
36 study of five European cities, *Journal of Geophysical Research: Atmospheres*, 116, D11302,  
37 10.1029/2011jd015646, 2011.

38 Goldstein, A. H., and Galbally, I. E.: Known and unexplored organic constituents in the earth's  
39 atmosphere, *Environ. Sci. Technol.*, 41, 1514-1521, 10.1021/es072476p, 2007.

40 Goldstein, A. H., Koven, C. D., Heald, C. L., and Fung, I. Y.: Biogenic carbon and  
41 anthropogenic pollutants combine to form a cooling haze over the southeastern United States,  
42 *Proc. Nat. Acad. Sci. U.S.A.* , 106, 8835-8840, 10.1073/pnas.0904128106, 2009.

1 Good, N., Topping, D. O., Duplissy, J., Gysel, M., Meyer, N. K., Metzger, A., Turner, S. F.,  
2 Baltensperger, U., Ristovski, Z., Weingartner, E., Coe, H., and McFiggans, G.: Widening the gap  
3 between measurement and modelling of secondary organic aerosol properties?, *Atmos. Chem.*  
4 *Phys.*, 10, 2577-2593, 10.5194/acp-10-2577-2010, 2010.

5 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and  
6 Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1  
7 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions,  
8 *Geoscientific Model Development*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.

9 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen,  
10 J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T.,  
11 Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W.,  
12 McFiggans, G., Mentel, T. F., Monod, A., Prevot, A. S. H., Seinfeld, J. H., Surratt, J. D.,  
13 Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic  
14 aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155-5236, 2009.

15 Hamilton, J. F., Webb, P. J., Lewis, A. C., and Reviejo, M. M.: Quantifying small molecules in  
16 secondary organic aerosol formed during the photo-oxidation of toluene with hydroxyl radicals,  
17 *Atmos. Environ.*, 39, 7263-7275, 10.1016/j.atmosenv.2005.09.006, 2005.

18 Hansen, A. M. K., Hong, J., Raatikainen, T., Kristensen, K., Ylisirniö, A., Virtanen, A., Petäjä,  
19 T., Glasius, M., and Prisle, N. L.: Hygroscopic properties and cloud condensation nuclei  
20 activation of limonene-derived organosulfates and their mixtures with ammonium sulfate,  
21 *Atmos. Chem. Phys.*, 15, 14071-14089, 10.5194/acp-15-14071-2015, 2015.

22 Heald, C. L., Jacob, D. J., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H., and  
23 Weber, R. J.: A large organic aerosol source in the free troposphere missing from current  
24 models, *Geophys. Res. Lett.*, 32, L18809, 10.1029/2005gl023831, 2005.

25 Hildebrandt, L., Donahue, N. M., and Pandis, S. N.: High formation of secondary organic aerosol  
26 from the photo-oxidation of toluene, *Atmos. Chem. Phys.*, 9, 2973-2986, 2009.

27 Hildebrandt, L., Henry, K. M., Kroll, J. H., Worsnop, D. R., Pandis, S. N., and Donahue, N. M.:  
28 Evaluating the Mixing of Organic Aerosol Components Using High-Resolution Aerosol Mass  
29 Spectrometry, *Environ. Sci. Technol.*, 45, 6329-6335, 10.1021/es200825g, 2011.

30 Hoyle, C. R., Boy, M., Donahue, N. M., Fry, J. L., Glasius, M., Guenther, A., Hallar, A. G.,  
31 Hartz, K. H., Petters, M. D., Petaja, T., Rosenoern, T., and Sullivan, A. P.: A review of the  
32 anthropogenic influence on biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 11, 321-  
33 343, DOI 10.5194/acp-11-321-2011, 2011.

34 Irwin, M., Good, N., Crosier, J., Choularton, T. W., and McFiggans, G.: Reconciliation of  
35 measurements of hygroscopic growth and critical supersaturation of aerosol particles in central  
36 Germany, *Atmos. Chem. Phys.*, 10, 11737-11752, 10.5194/acp-10-11737-2010, 2010.

37 Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of  
38 the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic  
39 volatile organic compounds, *Atmos. Chem. Phys.*, 3, 181-193, 2003.

40 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H.,  
41 DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M.,  
42 Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin,

1 C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M.,  
2 Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A.,  
3 Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F.,  
4 Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A.,  
5 Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J.  
6 R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C.,  
7 Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic  
8 Aerosols in the Atmosphere, *Science*, 326, 1525-1529, 10.1126/science.1180353, 2009.

9 Johnson, D., Jenkin, M. E., Wirtz, K., and Martin-Reviejo, M.: Simulating the formation of  
10 secondary organic aerosol from the photooxidation of aromatic hydrocarbons, *Environ. Chem.*,  
11 2, 35-48, 10.1071/en04079, 2005.

12 Jordan, A., Haidacher, S., Hanel, G., Hartungen, E., Mark, L., Seehauser, H., Schotchkowsky, R.,  
13 Sulzer, P., and Mark, T. D.: A high resolution and high sensitivity proton-transfer-reaction time-  
14 of-flight mass spectrometer (PTR-TOF-MS), *Int. J. Mass Spectrom.*, 286, 122-128,  
15 10.1016/j.ijms.2009.07.005, 2009.

16 Juranyi, Z., Gysel, M., Duplissy, J., Weingartner, E., Tritscher, T., Dommen, J., Henning, S.,  
17 Ziese, M., Kiselev, A., Stratmann, F., George, I., and Baltensperger, U.: Influence of gas-to-  
18 particle partitioning on the hygroscopic and droplet activation behaviour of alpha-pinene  
19 secondary organic aerosol, *Phys. Chem. Chem. Phys.*, 11, 8091-8097, 10.1039/b904162a, 2009.

20 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R.,  
21 Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as  
22 major components of atmospheric organic aerosols, *Science*, 303, 1659-1662,  
23 10.1126/science.1092185, 2004.

24 Kaminiski, M.: Untersuchung des photochemischen Terpenoidabbaus in der  
25 Atmosphärensimulationskammer SAPHIR, Ph.D. , Institut für Physikalische Chemie, Universität  
26 zu Köln, Jülich, 2014.

27 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van  
28 Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y.,  
29 Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati,  
30 E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review,  
31 *Atmos. Chem. Phys.*, 5, 1053-1123, 2005.

32 Koop, T., Bookhold, J., Shiraiwa, M., and Poeschl, U.: Glass transition and phase state of  
33 organic compounds: dependency on molecular properties and implications for secondary organic  
34 aerosols in the atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19238-19255, 10.1039/c1cp22617g,  
35 2011.

36 Kourtchev, I., Fuller, S., Aalto, J., Ruuskanen, T. M., McLeod, M. W., Maenhaut, W., Jones, R.,  
37 Kulmala, M., and Kalberer, M.: Molecular Composition of Boreal Forest Aerosol from Hyytiälä,  
38 Finland, Using Ultrahigh Resolution Mass Spectrometry, *Environ. Sci. Technol.*, 47, 4069-4079,  
39 10.1021/es3051636, 2013.

40 Kourtchev, I., O'Connor, I. P., Giorio, C., Fuller, S. J., Kristensen, K., Maenhaut, W., Wenger, J.  
41 C., Sodeau, J. R., Glasius, M., and Kalberer, M.: Effects of anthropogenic emissions on the

1 molecular composition of urban organic aerosols: An ultrahigh resolution mass spectrometry  
2 study, *Atmos. Environ.*, 89, 525-532, 10.1016/j.atmosenv.2014.02.051, 2014.

3 Kourtchev, I., Doussin, J. F., Giorio, C., Mahon, B., Wilson, E. M., Maurin, N., Pangui, E.,  
4 Venables, D. S., Wenger, J. C., and Kalberer, M.: Molecular composition of fresh and aged  
5 secondary organic aerosol from a mixture of biogenic volatile compounds: a high-resolution  
6 mass spectrometry study, *Atmos. Chem. Phys.*, 15, 5683-5695, 10.5194/acp-15-5683-2015,  
7 2015.

8 Kristensen, K., and Glasius, M.: Organosulfates and oxidation products from biogenic  
9 hydrocarbons in fine aerosols from a forest in North West Europe during spring, *Atmos.*  
10 *Environ.*, 45, 4546-4556, 10.1016/j.atmosenv.2011.05.063, 2011.

11 Kroll, J. H., and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and  
12 evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593-3624,  
13 10.1016/j.atmosenv.2008.01.003, 2008.

14 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C.,  
15 Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J.,  
16 Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van  
17 Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of  
18 reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017-7039,  
19 10.5194/acp-10-7017-2010, 2010.

20 Lambe, A. T., Onasch, T. B., Massoli, P., Croasdale, D. R., Wright, J. P., Ahern, A. T.,  
21 Williams, L. R., Worsnop, D. R., Brune, W. H., and Davidovits, P.: Laboratory studies of the  
22 chemical composition and cloud condensation nuclei (CCN) activity of secondary organic  
23 aerosol (SOA) and oxidized primary organic aerosol (OPOA), *Atmos. Chem. Phys.*, 11, 8913-  
24 8928, 10.5194/acp-11-8913-2011, 2011.

25 Lance, S., Medina, J., Smith, J. N., and Nenes, A.: Mapping the operation of the DMT  
26 Continuous Flow CCN counter, *Aerosol Sci. Technol.*, 40, 242-254,  
27 10.1080/02786820500543290, 2006.

28 Low, R. D. H.: A Generalized Equation for the Solution Effect in Droplet Growth, *Journal of the*  
29 *Atmospheric Sciences*, 26, 608-611, 10.1175/1520-0469(1969)026<0608:AGEFTS>2.0.CO;2,  
30 1969.

31 Loza, C. L., Coggon, M. M., Nguyen, T. B., Zuend, A., Flagan, R. C., and Seinfeld, J. H.: On the  
32 Mixing and Evaporation of Secondary Organic Aerosol Components, *Environ. Sci. Technol.*, 47,  
33 6173-6180, 10.1021/es400979k, 2013.

34 Massoli, P., Lambe, A. T., Ahern, A. T., Williams, L. R., Ehn, M., Mikkila, J., Canagaratna, M.  
35 R., Brune, W. H., Onasch, T. B., Jayne, J. T., Petaja, T., Kulmala, M., Laaksonen, A., Kolb, C.  
36 E., Davidovits, P., and Worsnop, D. R.: Relationship between aerosol oxidation level and  
37 hygroscopic properties of laboratory generated secondary organic aerosol (SOA) particles,  
38 *Geophys. Res. Lett.*, 37, L24801, 10.1029/2010gl045258, 2010.

39 McDonald, J. E.: Erroneous cloud-physics applications of Raoult's law, *Journal of Meteorology*,  
40 10, 68-70, 10.1175/1520-0469(1953)010<0068:ECPSAO>2.0.CO;2, 1953.

1 Moore, R. H., Nenes, A., and Medina, J.: Scanning Mobility CCN Analysis-A Method for Fast  
2 Measurements of Size-Resolved CCN Distributions and Activation Kinetics, *Aerosol Sci.*  
3 *Technol.*, 44, 861-871, 10.1080/02786826.2010.498715, 2010.

4 Ng, N. L., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Kroll, J. H., Kwan, A. J., McCabe, D.  
5 C., Wennberg, P. O., Sorooshian, A., Murphy, S. M., Dalleska, N. F., Flagan, R. C., and  
6 Seinfeld, J. H.: Effect of NO(x) level on secondary organic aerosol (SOA) formation from the  
7 photooxidation of terpenes, *Atmos. Chem. Phys.*, 7, 5159-5174, 2007a.

8 Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.:  
9 Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem. Phys.*,  
10 7, 3909-3922, 2007b.

11 Noziere, B., Kaberer, M., Claeys, M., Allan, J., D'Anna, B., Decesari, S., Finessi, E., Glasius,  
12 M., Grgic, I., Hamilton, J. F., Hoffmann, T., Iinuma, Y., Jaoui, M., Kahno, A., Kampf, C. J.,  
13 Kourtchev, I., Maenhaut, W., Marsden, N., Saarikoski, S., Schnelle-Kreis, J., Surratt, J. D.,  
14 Szidat, S., Szmigielski, R., and Wisthaler, A.: The Molecular Identification of Organic  
15 Compounds in the Atmosphere: State of the Art and Challenges, *Chem. Rev.*, 115, 3919-3983,  
16 10.1021/cr5003485, 2015.

17 Padro, L. T., Asa-Awuku, A., Morrison, R., and Nenes, A.: Inferring thermodynamic properties  
18 from CCN activation experiments: single-component and binary aerosols, *Atmos. Chem. Phys.*,  
19 7, 5263-5274, 2007.

20 Petrucci, H., and Herring, M.: *General Chemistry, Principles & Modern Applications*, 9th ed.,  
21 Pearson Education, Inc., , Upper Saddle River, NJ, 2007.

22 Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth  
23 and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, 2007.

24 Petters, M. D., Wex, H., Carrico, C. M., Hallbauer, E., Massling, A., McMeeking, G. R.,  
25 Poulain, L., Wu, Z., Kreidenweis, S. M., and Stratmann, F.: Towards closing the gap between  
26 hygroscopic growth and activation for secondary organic aerosol - Part 2: Theoretical  
27 approaches, *Atmos. Chem. Phys.*, 9, 3999-4009, 2009.

28 Prenni, A. J., Petters, M. D., Kreidenweis, S. M., DeMott, P. J., and Ziemann, P. J.: Cloud  
29 droplet activation of secondary organic aerosol, *J. Geophys. Res.-Atmos.*, 112, D10223,  
30 10.1029/2006jd007963, 2007.

31 Presto, A. A., Hartz, K. E. H., and Donahue, N. M.: Secondary organic aerosol production from  
32 terpene ozonolysis. 2. Effect of NOx concentration, *Environ. Sci. Technol.*, 39, 7046-7054,  
33 10.1021/es050400s, 2005.

34 Prisle, N. L., Raatikainen, T., Sorjamaa, R., Svenningsson, B., Laaksonen, A., and Bilde, M.:  
35 Surfactant partitioning in cloud droplet activation: a study of C8, C10, C12 and C14 normal fatty  
36 acid sodium salts, *Tellus Ser. B-Chem. Phys. Meteorol.*, 60, 416-431, 10.1111/j.1600-  
37 0889.2008.00352.x, 2008.

38 Rickards, A. M. J., Miles, R. E. H., Davies, J. F., Marshall, F. H., and Reid, J. P.: Measurements  
39 of the Sensitivity of Aerosol Hygroscopicity and the kappa Parameter to the O/C Ratio, *J. Phys.*  
40 *Chem. A* 117, 14120-14131, 10.1021/jp407991n, 2013.

1 Rohrer, F., Bohn, B., Brauers, T., Bruning, D., Johnen, F. J., Wahner, A., and Kleffmann, J.:  
2 Characterisation of the photolytic HONO-source in the atmosphere simulation chamber  
3 SAPHIR, *Atmos. Chem. Phys.*, 5, 2189-2201, 2005.

4 Rose, D., Gunthe, S. S., Mikhailov, E., Frank, G. P., Dusek, U., Andreae, M. O., and Poschl, U.:  
5 Calibration and measurement uncertainties of a continuous-flow cloud condensation nuclei  
6 counter (DMT-CCNC): CCN activation of ammonium sulfate and sodium chloride aerosol  
7 particles in theory and experiment, *Atmos. Chem. Phys.*, 8, 1153-1179, 2008.

8 Shantz, N. C., Chang, R. Y. W., Slowik, J. G., Vlasenko, A., Abbatt, J. P. D., and Leaitch, W. R.:  
9 Slower CCN growth kinetics of anthropogenic aerosol compared to biogenic aerosol observed at  
10 a rural site, *Atmos. Chem. Phys.*, 10, 299-312, 2010.

11 Shantz, N. C., Pierce, J. R., Chang, R. Y. W., Vlasenko, A., Riipinen, I., Sjostedt, S., Slowik, J.  
12 G., Wiebe, A., Liggi, J., Abbatt, J. P. D., and Leaitch, W. R.: Cloud condensation nuclei droplet  
13 growth kinetics of ultrafine particles during anthropogenic nucleation events, *Atmos. Environ.*,  
14 47, 389-398, 10.1016/j.atmosenv.2011.10.049, 2012.

15 Shilling, J. E., Zaveri, R. A., Fast, J. D., Kleinman, L., Alexander, M. L., Canagaratna, M. R.,  
16 Fortner, E., Hubbe, J. M., Jayne, J. T., Sedlacek, A., Setyan, A., Springston, S., Worsnop, D. R.,  
17 and Zhang, Q.: Enhanced SOA formation from mixed anthropogenic and biogenic emissions  
18 during the CARES campaign, *Atmos. Chem. Phys.*, 13, 2091-2113, 10.5194/acp-13-2091-2013,  
19 2013.

20 Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W.,  
21 Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, A., and Forster, P.:  
22 Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos.*  
23 *Chem. Phys.*, 11, 12109-12136, 10.5194/acp-11-12109-2011, 2011.

24 Suda, S. R., Petters, M. D., Yeh, G. K., Strollo, C., Matsunaga, A., Faulhaber, A., Ziemann, P. J.,  
25 Prenni, A. J., Carrico, C. M., Sullivan, R. C., and Kreidenweis, S. M.: Influence of Functional  
26 Groups on Organic Aerosol Cloud Condensation Nucleus Activity, *Environ. Sci. Technol.*, 48,  
27 10182-10190, 10.1021/es502147y, 2014.

28 Tolocka, M. P., Jang, M., Ginter, J. M., Cox, F. J., Kamens, R. M., and Johnston, M. V.:  
29 Formation of oligomers in secondary organic aerosol, *Environ. Sci. Technol.*, 38, 1428-1434,  
30 10.1021/es035030r, 2004.

31 Topping, D., Connolly, P., and McFiggans, G.: Cloud droplet number enhanced by co-  
32 condensation of organic vapours, *Nat. Geosci.*, 6, 443-446, 10.1038/ngeo1809, 2013.

33 Topping, D. O., McFiggans, G. B., and Coe, H.: A curved multi-component aerosol  
34 hygroscopicity model framework: Part 2 - Including organic compounds, *Atmos. Chem. Phys.*, 5,  
35 1223-1242, 2005.

36 Topping, D. O., and McFiggans, G.: Tight coupling of particle size, number and composition in  
37 atmospheric cloud droplet activation, *Atmos. Chem. Phys.*, 12, 3253-3260, 10.5194/acp-12-  
38 3253-2012, 2012.

39 Tritscher, T., Dommen, J., DeCarlo, P. F., Gysel, M., Barmet, P. B., Praplan, A. P., Weingartner,  
40 E., Prevot, A. S. H., Riipinen, I., Donahue, N. M., and Baltensperger, U.: Volatility and  
41 hygroscopicity of aging secondary organic aerosol in a smog chamber, *Atmos. Chem. Phys.*, 11,  
42 11477-11496, 10.5194/acp-11-11477-2011, 2011.

1 Verma, V., Fang, T., Guo, H., King, L., Bates, J. T., Peltier, R. E., Edgerton, E., Russell, A. G.,  
2 and Weber, R. J.: Reactive oxygen species associated with water-soluble PM<sub>2.5</sub> in the  
3 southeastern United States: spatiotemporal trends and source apportionment, *Atmos. Chem.*  
4 *Phys.*, 14, 12915-12930, 10.5194/acp-14-12915-2014, 2014.

5 Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirila, P., Leskinen, J., Makela, J. M.,  
6 Holopainen, J. K., Poeschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous  
7 solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824-827,  
8 10.1038/nature09455, 2010.

9 Weber, R. J., Sullivan, A. P., Peltier, R. E., Russell, A., Yan, B., Zheng, M., de Gouw, J.,  
10 Warneke, C., Brock, C., Holloway, J. S., Atlas, E. L., and Edgerton, E.: A study of secondary  
11 organic aerosol formation in the anthropogenic-influenced southeastern United States, *J.*  
12 *Geophys. Res.-Atmos.*, 112, D13302, 10.1029/2007jd008408, 2007.

13 Wex, H., Petters, M. D., Carrico, C. M., Hallbauer, E., Massling, A., McMeeking, G. R.,  
14 Poulain, L., Wu, Z., Kreidenweis, S. M., and Stratmann, F.: Towards closing the gap between  
15 hygroscopic growth and activation for secondary organic aerosol: Part 1-Evidence from  
16 measurements, *Atmos. Chem. Phys.*, 9, 3987-3997, 2009.

17 Whitehead, J. D., Irwin, M., Allan, J. D., Good, N., and McFiggans, G.: A meta-analysis of  
18 particle water uptake reconciliation studies, *Atmos. Chem. Phys.*, 14, 11833-11841, 10.5194/acp-  
19 14-11833-2014, 2014.

20 Wiedensohler, A.: An approximation of the bipolar charge-distribution for particles in the sub-  
21 micron size range, *J. Aerosol Sci.*, 19, 387-389, 10.1016/0021-8502(88)90278-9, 1988.

22 Worton, D. R., Goldstein, A. H., Farmer, D. K., Docherty, K. S., Jimenez, J. L., Gilman, J. B.,  
23 Kuster, W. C., de Gouw, J., Williams, B. J., Kreisberg, N. M., Hering, S. V., Bench, G., McKay,  
24 M., Kristensen, K., Glasius, M., Surratt, J. D., and Seinfeld, J. H.: Origins and composition of  
25 fine atmospheric carbonaceous aerosol in the Sierra Nevada Mountains, California, *Atmos.*  
26 *Chem. Phys.*, 11, 10219-10241, 10.5194/acp-11-10219-2011, 2011.

27 Xu, L., Guo, H. Y., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R.,  
28 Isaacman-VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H.,  
29 Hering, S. V., de Gouw, J., Baumann, K., Lee, S. H., Nenes, A., Weber, R. J., and Ng, N. L.:  
30 Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the  
31 southeastern United States, *Proc. Nat. Acad. Sci. U.S.A.* , 112, 37-42, 10.1073/pnas.1417609112,  
32 2015.

33 Xu, W., Guo, S., Gomez-Hernandez, M., Zamora, M. L., Secret, J., Marrero-Ortiz, W., Zhang,  
34 A. L., Collins, D. R., and Zhang, R. Y.: Cloud forming potential of oligomers relevant to  
35 secondary organic aerosols, *Geophys. Res. Lett.* , 41, 6538-6545, 10.1002/2014gl061040, 2014.

36 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and  
37 Sun, Y. L.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass  
38 spectrometry: a review, *Anal. Bioanal. Chem.*, 401, 3045-3067, 10.1007/s00216-011-5355-y,  
39 2011.

40 Zhang, X., Cappa, C. D., Jathar, S. H., McVay, R. C., Ensberg, J. J., Kleeman, M. J., and  
41 Seinfeld, J. H.: Influence of vapor wall loss in laboratory chambers on yields of secondary  
42 organic aerosol, *Proc. Nat. Acad. Sci. U.S.A.* , 111, 5802-5807, 10.1073/pnas.1404727111, 2014.

- 1 Zhang, Y. Y., Mueller, L., Winterhalter, R., Moortgat, G. K., Hoffmann, T., and Poeschl, U.:  
2 Seasonal cycle and temperature dependence of pinene oxidation products, dicarboxylic acids and  
3 nitrophenols in fine and coarse air particulate matter, *Atmos. Chem. Phys.*, 10, 7859-7873,  
4 10.5194/acp-10-7859-2010, 2010.
- 5 Zhao, D. F., Buchholz, A., Mentel, T. F., Muller, K. P., Borchardt, J., Kiendler-Scharr, A.,  
6 Spindler, C., Tillmann, R., Trimborn, A., Zhu, T., and Wahner, A.: Novel method of generation  
7 of  $\text{Ca}(\text{HCO}_3)_2$  and  $\text{CaCO}_3$  aerosols and first determination of hygroscopic and cloud  
8 condensation nuclei activation properties, *Atmos. Chem. Phys.*, 10, 8601-8616, 10.5194/acp-10-  
9 8601-2010, 2010.
- 10 Zhao, D. F., Buchholz, A., Kortner, B., Schlag, P., Rubach, F., Kiendler-Scharr, A., Tillmann,  
11 R., Wahner, A., Watne, Å. K., Hallquist, M., Rudich, Y., Wildt, J., and Mentel, T. F.:  
12 Dependence of cloud condensation nuclei activity of secondary organic aerosol (SOA) on  
13 particle size, in preparation, 2015a.
- 14 Zhao, D. F., Kaminski, M., Schlag, P., Fuchs, H., Acir, I. H., Bohn, B., Häseler, R., Kiendler-  
15 Scharr, A., Rohrer, F., Tillmann, R., Wang, M. J., Wegener, R., Wildt, J., Wahner, A., and  
16 Mentel, T. F.: Secondary organic aerosol formation from hydroxyl radical oxidation and  
17 ozonolysis of monoterpenes, *Atmos. Chem. Phys.*, 15, 991-1012, 10.5194/acp-15-991-2015,  
18 2015b.
- 19 Zotter, P., Ciobanu, V. G., Zhang, Y. L., El-Haddad, I., Macchia, M., Daellenbach, K. R.,  
20 Salazar, G. A., Huang, R. J., Wacker, L., Hueglin, C., Piazzalunga, A., Fermo, P., Schwikowski,  
21 M., Baltensperger, U., Szidat, S., and Prevot, A. S. H.: Radiocarbon analysis of elemental and  
22 organic carbon in Switzerland during winter-smog episodes from 2008 to 2012-Part 1: Source  
23 apportionment and spatial variability, *Atmos. Chem. Phys.*, 14, 13551-13570, 10.5194/acp-14-  
24 13551-2014, 2014.


25



Table 1 Summary of the experiments in this study

Experiment #	Experiment type	Experiment description	BVOC (ppb)	AVOC (ppb)	Ozone added (ppb)
B1	BSOA	BVOC only	$\alpha$ -pinene, limonene (20, 20)	N.A	50
B2	BSOA	BVOC only	$\alpha$ -pinene (40)	N.A	160
B3	BSOA	BVOC only	$\alpha$ -pinene, limonene (48, 48)	N.A	200
A1	ASOA	AVOC only	N.A	toluene (85)	0
A2	ASOA	AVOC only	N.A	toluene (85)	0
A3	ASOA	AVOC only	N.A	p-xylene (30)	0
A4	ASOA	AVOC only	N.A	p-xylene (30)	0
A5	ASOA	AVOC only	N.A	benzene (280)	0
A6	ASOA	AVOC only	N.A	benzene (280)	0
A7	ASOA	AVOC only	N.A	p-xylene-d <sub>10</sub> (200)	200
AB1	ABSOA	AVOC added 6.3 h before BVOC	$\alpha$ -pinene, limonene (20, 20)	p-xylene (30)	0
AB2	ABSOA	AVOC added 6.3 h before BVOC	$\alpha$ -pinene, limonene (20, 20)	toluene (85)	0
AB3	ABSOA	BVOC added 3 h before AVOC	$\alpha$ -pinene, limonene (20, 20)	toluene (85)	50
AB4	ABSOA	BVOC added 5 h before AVOC	$\alpha$ -pinene, limonene (39, 39)	p-xylene-d <sub>10</sub> (51)	200
AB5	ABSOA	AVOC and BVOC added together	$\alpha$ -pinene, limonene (4, 4)	toluene (85)	60
AB6	ABSOA	AVOC and BVOC added together	$\alpha$ -pinene, limonene (42, 42)	p-xylene-d <sub>10</sub> (90)	200


1




2

3 Figure 1. CCN activity of BSOA, ASOA, and ABSOA as a function of OH dose (A) and O/C of  
 4 aerosol (B).  $\kappa_{CCN}$  at two supersaturations (SS) 0.34% (solid markers) and 0.51% (open markers)  
 5 is shown. The green, red and blue colors indicate BSOA, ASOA and ABSOA, respectively.  $\kappa_{CCN}$   
 6 at similar SS is compared since  $\kappa_{CCN}$  depended on SS. BSOA, ASOA, and ABSOA showed no  
 7 significant difference at a given SS. Although ASOA had higher degree of oxidation (O/C) than  
 8 BSOA, ASOA components did not enhance  $\kappa_{CCN}$ .

9




1



2

3 Figure 2. CCN activity of ABSOA from sequential VOC addition at various supersaturations  
 4 (SS). (A) Toluene was added first and monoterpenes were added 6 h afterwards as indicated by  
 5 the vertical arrow. The shaded areas show dark periods. ASOA fraction (red dots, right axis)  
 6 decreased remarkably upon BVOC addition but  $\kappa_{CCN}$  showed no significant change. (B)  
 7 Monoterpenes were added first and toluene was added 2.6 h afterwards as indicated by the  
 8 vertical arrow.  $\kappa_{CCN}$  showed no significant change upon the addition of AVOC to BSOA. Note  
 9 that time series of  $\kappa_{CCN}$  should be followed at similar SS because  $\kappa_{CCN}$  depended on SS and the  
 10 SS range may change due to the particle size change.


11




1

2 Figure 3. Droplet sizes of BSOA, ASOA, ABSOA and ammonium sulfate aerosols at various  
 3 supersaturations (SS). In the CCNC, all SOA particles reached comparable droplet sizes  
 4 compared to ammonium sulfate.

5




1




2

3 Figure 4.  $\kappa_{\text{HTDMA}}$  for BSOA, ASOA and ABSOA as a function of OH dose (A) and O/C of  
 4 aerosol (B). ASOA had a distinctively higher  $\kappa_{\text{HTDMA}}$  and O/C than BSOA.

5




1



2

3 Figure 5.  $\kappa_{\text{HTDMA}}$  of ABSOA from experiments when AVOC and BVOC were added  
 4 sequentially (same experiments as in Fig. 2). (A) Toluene was added first and the monoterpenes  
 5 were added 6 h afterwards as indicated by the vertical arrow. The shaded areas indicate the dark  
 6 periods.  $\kappa_{\text{HTDMA}}$  decreased significantly with the decrease of ASOA fraction due to the addition  
 7 of BVOC. At the same time,  $f_{44}$  decreased distinctively. (B) Monoterpenes were added first and  
 8 toluene was added 2.6 h afterwards as indicated by the vertical arrow.  $\kappa_{\text{HTDMA}}$  increased slightly  
 9 as only a minor fraction (<10%) of ASOA was formed. At the same time,  $f_{44}$  also increased  
 10 slightly.


1



2

3 Figure 6.  $\kappa_{HTDMA}$  of ABSOA as a function of ASOA fraction. The size of the marker denotes the  
4 OH dose. For ASOA,  $\kappa_{HTDMA}$  is largely independent of OH dose and the average value was used,  
5 shown as the same point for different OH dose. The dashed lines connecting the pure BSOA and  
6 ASOA of the same OH doses stand for the linear combination of the  $\kappa_{HTDMA}$  of pure BSOA and  
7 pure ASOA at that given OH dose. A data point beyond the line corresponding to the OH dose of  
8 that point indicates a non-linear effect. The cases of non-linear effect can be found for ABSOA  
9 #AB1, #AB2 and #AB5. The dotted lines with arrow of ABSOA show the time order of data  
10 points.


11




1  
 2 Figure 7. Comparison of  $\kappa_{\text{HTDMA}}$  and  $\kappa_{\text{CCN}}$  from HTDMA and CCN measurements for BSOA,  
 3 ASOA, and ABSOA. The solid bar denotes the  $\kappa_{\text{HTDMA}}$  and the open bar denotes  $\kappa_{\text{CCN}}$  (left axis).  
 4 The diamond denotes the ratio of  $\kappa_{\text{HTDMA}}$  to  $\kappa_{\text{CCN}}$  (right axis). The green, blue and red color  
 5 indicates the BSOA, ABSOA and ASOA, respectively. For all SOA here, there was a  
 6 discrepancy between  $\kappa_{\text{HTDMA}}$  and  $\kappa_{\text{CCN}}$ . For ASOA, the gap between  $\kappa_{\text{HTDMA}}$  and  $\kappa_{\text{CCN}}$  (low  
 7  $\kappa_{\text{HTDMA}}/\kappa_{\text{CCN}}$ ) was smaller compared to BSOA.

8






1  
 2 Figure S1. CCN activity of BSOA (A), ASOA (B) and ABSOA (C) at various supersaturations  
 3 (SS). BSOA was formed by ozonolysis of monoterpene (mixture of  $\alpha$ -pinene and limonene with  
 4 a molar ratio of 1:1) followed by photooxidation. ASOA was formed by photooxidation of  
 5 toluene. ABSOA was formed by the photooxidation of a mixture of toluene and monoterpenes  
 6 ( $\alpha$ -pinene:limonene =1:1). The shaded areas indicate dark periods.




1




2

3 Figure S2. CCN activity of BSOA, ASOA and ABSOA represented by  $\kappa_{CCN}$  at various  
 4 supersaturations (SS) as a function of OH dose (A) and O/C of aerosol (B). The points lining  
 5 vertically for each aerosol type in panel A are from the dark period.



1  
 2 Figure S3. Critical supersaturation as a function of dry particle diameter of ASOA formed from  
 3 toluene, benzene, and xylene photooxidation in the low  $\text{NO}_x$  (<1 ppb) and high  $\text{NO}_x$  condition  
 4 (10 ppb  $\text{NO}$  added). ASOA from different precursors show similar CCN activity. ASOA  
 5 produced at low  $\text{NO}_x$  and high  $\text{NO}_x$  show similar CCN activity.  
 6




1

2 Figure S4. O/C of BSOA, ASOA and ABSOA as a function of OH dose.

3

4


1



2

3 Figure S5. Comparison of  $\kappa$  in this study with the parameterization of the relationship between  $\kappa$   
4 and O/C in the literature (Rickards et al., 2013). The blue markers show  $\kappa_{CCN}$  and the red  
5 markers show  $\kappa_{HTDMA}$ . The lines show the upper and lower limits of the parameterization in  
6 Rickards et al. (2013).

7



1  
2 Figure S6. Mass spectra from nano ESI UHRMS of BSOA (red sticks, from experiment #B3  
3 using  $\alpha$ -pinene + limonene mixture as precursor) and ABSOA (blue sticks, from experiment  
4 #AB6 using  $\alpha$ -pinene+limonene+p-xylene mixture as precursor).