

# Carbonyl sulfide exchange in soils for better estimates of ecosystem carbon uptake

## Response to Reviewers

We thank all of our referees for their contributions to this paper. Below, referee comments are in **bold** and our responses are in plain text.

### 1. Response to Dr. J. Kesselmeier

We thank Dr. Kesselmeier for taking the time to write such helpful feedback. First, we address the effect of COS concentration on fluxes and how it relates to the results from this study. We present new COS net fluxes observed under COS-free sweep air conditions compared to ambient conditions in the Supplementary material. Then we attend to each of the specific comments.

#### **General Comments**

**The goal of this paper is to contribute to a discussion around the role of soils within the exchange of carbonyl sulfide (COS) between a forest ecotype and the atmosphere. The net signal of the COS exchange is a significant matter of discussion for estimating Gross primary Productivity (GPP) independently from the complex practice taking into account net carbon exchange corrected by hetero- and autotrophic respiration. However, in order to exploit the gross uptake of COS, a significant contribution by soils must be excluded. Therefore, the authors investigated soils from 5 different sites and came to the conclusion that soil interaction by uptake and/or emission is not negligible as compared to plant uptake and may interfere with GPP estimation. However, uncertainty of GPP can be regarded to be much larger.**

**This paper deserves publication in ACP after some revision. Data, discussion and conclusion are principally convincing and may help to find a general approach to determine GPP in future. However, the model still deserves some improvements to match observations under natural conditions. There may be missing links. A profound discussion of the lack of matching observed data would be helpful. One important issue for example is not discussed at all in the paper. We know from several studies that uptake fluxes of COS are linearly depending on atmospheric COS concentrations. How will such fluctuations, which may be considerable, be included? Furthermore, the current measurements as described by the authors are performed under “ambient” COS concentrations. But no concentration data are given. It would be of high interest for the reader to get information about concentrations and eventual fluctuations of atmospheric COS concentrations during the measurements. Such data would allow estimating deposition velocities which are easier to compare with other reports. Furthermore, some variabilities, such as in figure 6 might be caused by fluctuations of atmospheric concentrations?**

**Understanding will be improved by presenting data on COS concentrations. This issue should be carefully discussed if it cannot be taken into account.**

This is an excellent point. The relationship between COS concentration and COS uptake has been well established (e.g. Kesselmeier et al. 1999). We have now addressed this in the supplementary material where we present an additional experiment targeting this issue. We now include chamber mixing ratio data. We investigate whether the patterns of Figure 6 could be due to variability of the mixing ratio of COS within the chamber (Supplementary, Figure S1). We believe that the changes in flux due to simulated rainfall overwhelms the influence of chamber mixing ratio on COS uptake.

### **Specific Comments**

**1. Introduction, page 21097, line 2-6: Citation of the first report on the close relationship between COS uptake and GPP as derived from deposition velocity corrected COS/CO<sub>2</sub> uptake by branch level measurements might be nice. Sandoval-Soto et al (2005) were the first to demonstrate that the COS/CO<sub>2</sub> uptake ratios corrected against atmospheric ratios and deposition velocities indicate the potential for GPP estimation. This paper initiated the current COS-related GPP discussions for ecosystems.**

There is some confusion about who initiated the GPP-COS discussion. A 2004 AGU abstract by Steve Montzka, for example, suggests the link. The important contribution by Sandoval-Soto et al. (2005) related COS/CO<sub>2</sub> uptake to NPP, although it is pointed out that it should scale with GPP. This reference is now included in the introduction.

**2. Introduction, page 21097, line 9-13: Enzymatic consumption of COS has not only been demonstrated with the isolated enzyme carbonic anhydrase (CA), but also with Ribulose-1,5-bisphosphate-carboxylase/-oxygenase (RuBisCO) and Phosphoenolpyruvate-Carboxylase (PEP-CO). The cooperation of these three enzymes has already been reported in 1992 (Protoschill-Krebs, G. and Kesselmeier, J., Enzymatic pathways for the consumption of carbonyl sulphide (COS) by higher plants. *Botanica Acta* 105, 206-212, 1992). Furthermore, cooperation of CA with other enzymes was further discussed for soil microorganisms by Wingate et al. (2008). When investigating the function of soils, we should not concentrate only on CA, though it may be the key enzyme, but also mention the potential of other enzymes. In view of the complex “soil organism” our views should not focus too much. Otherwise, we might easily oversee important details.**

We thank Dr. Kesselmeier for pointing this out. While CA is seemingly ubiquitous in soil microorganisms, RuBisCO is also found in soils. The language of the introduction has been changed to reflect this important idea: “Other enzymes such as RuBisCO can also destroy COS (Protoschill-Krebs and Kesselmeier, 1992)”.

**3. Methods, page 21099, lines 26-28: The authors state that by keeping soils whole (unsieved), this reduced the influence of sample processing artifacts on lab-based flux observations. That sounds too optimistic. It helps to hopefully stay a bit closer to natural conditions. However, sieving helps to reach a highly reproducible material to be analyzed and is used in most cases of soil exchange measurements.**

**Skipping this step can lead to unreproducible soil sample mixtures, which may also create problems. Rewriting this statement in order to give both views a right to exist would be appropriate.**

We acknowledge that there are two valid views on the subject of soil-sieving. The language of this statement is changed in the text to reflect this: “By keeping soils whole, we reduced the influence of sample processing artifacts on our lab-based flux observations at the expense of working with non-homogenized and therefore less reproducible samples.”

**4. Methods, page 21100, lines 8 and following: It is always helpful to give units for formula components. This helps the reader to reproduce calculations. Within line 29 and 31, the two different terms “mixing ratio” and “concentration” are used with the same sense. The use should be harmonized.**

Thank you for pointing this out. The terms have been harmonized in the text.

**5. Methods, page 21100, lines 15-16: What do the authors want to describe with “uninterpretable because of variations in ambient CO<sub>2</sub> concentrations, C<sub>i</sub>”? Was the CO<sub>2</sub> concentration in the incoming air highly fluctuating? Were the measurement performed by switching between in- and outlet or were they performed simultaneously? A little bit more information about measurement procedures would be helpful to understand the authors’ concerns.**

The methods section now has more detail to make this point clearer. As described in the supplementary text, we used ambient air for these measurements. While COS varied by +/- 80 ppt, CO<sub>2</sub> occasionally had a larger variation over the course of a 1-hour measurements. If the incoming concentration of CO<sub>2</sub> was inconsistent, trying to calculate a flux is difficult to impossible.

**6. Results, chapter 3.1.: The authors give the amount of water in soils as volumetric water content %. That is fine. But it might be helpful for readers to compare to other units. Why not giving also weight%?**

Table 1 gives the bulk densities of the soils from which gravimetric weight can be calculated from volumetric weight.

**7. Results, page 21109, line 25-27: The authors cite van Diest and Kesselmeier (2008) and Kesselmeier et al. (1999) and mention that all soils measured within this work were sandy soils. A little bit more information was available and given in these papers. Maybe the authors can also add some more information to their table 1 including some characteristics such as maximum soil water content and calculated water filled pore space.**

This is an excellent suggestion. We have now included information about soil texture in Table 1. Water filled pore space can be calculated using the data included in the paper.

**8. Chapter 4.2., page 21110, line 20-21: The authors cite Whelan and Rhew (2015) who autoclaved agricultural soils and found only COS production left. I think this is a questionable prove. Autoclaving cooks the biological material and kills microbes, but what about all the organic stuff, which is still available, may be decomposing now and may be in a very volatile condition. Hence, autoclaving may induce a production and release and may cover any uptake process.**

The main idea of the paragraph in Section 4.2 is that it is possible all soils may be experiencing COS production, but that some soils have large microbial communities that can overcome in situ production and consume COS from the atmosphere, resulting in a net COS sink. It has been established that autoclaved soils both exhibit a net production of COS and have altered organic matter. Autoclaving soils in Whelan and Rhew (2015) was a compromise among many experimental choices. There does not seem to be a method to sterilize soils without altering the organic matter content that we are aware of. Autoclaving should also result in destruction of both cells and extra cellular enzymes. Known abiotic COS uptake processes are dissolution and hydrolysis in water and adsorption onto solid surfaces. At equilibrium with the chamber headspace, these processes should not result in an observable flux. It is unclear what additional uptake process would be covered up by production induced by autoclaving. We still do not have definitive proof, and we used the language “partially supported” in the text. We now include the caveat: “though autoclaved soils have been known to emit COS (Kato et al. 2008).”

**9. Information about calibration, accuracy and precision of the gas analyzers is missing throughout the paper. Within this context, error bars within the figures (5-8) would be of great help.**

We agree with Dr. Kesselmeier: the figures now have error bars, some of which are smaller than the symbols. The discussion in the methods section was expanded to make the measurement procedure clearer.

## **2. Response to Anonymous Referee #2**

We thank Anonymous Referee #2 for his or her insightful comments. We have addressed each one in turn to the betterment of the manuscript.

**The goal of this paper is quantify and understand the soil flux of carbonyl sulfide (COS) so that COS can be used more confidently as a proxy for gross primary productivity (GPP). This paper represents a significant contribution to our understanding of soil fluxes but also presents the uncertainties and suggests future work.**

**General comments:**

**Could the prior history of the soils lead to some of the variability seen in this study?**

We hope that analyzing the soils as soon as possible after collection make them as representative of their in situ state as possible. One would expect that the history of the agricultural fields (fertilizer, pesticides, etc) would affect fluxes. Whatever the history of the agricultural soil, some common management practice is probably at fault. This idea is supported by the soy bean/corn field soil investigated here sharing similarities in flux patterns with an independently managed wheat field 800 km away. The two distinct soil types, crop, and growing strategies both result in exponential COS emissions with temperature.

**Technical details:**

**Pg 21097, line 16: ppt is mole fraction not concentration**

We thank the referee for pointing this out. It has been changed in the text.

**Pg 21100, line 16: This is confusing. Was a constant concentration of COS and CO<sub>2</sub> used for these studies? If lab air was used, what was the variability in the ambient mixing ratio. And if the CO<sub>2</sub> mixing ratios changed to the point of the flux not being useful, how believable were the COS fluxes for the same periods?**

We now include a discussion of soil COS ambient concentration variability in the text, with a larger discussion in the supplemental.

**Pg 21100, line 24: How large was this correction?**

The correction was generally less than 1%. This is now stated in the text.

**Pg 21101, line 16: Awkward phrasing.**

The text now reads: “Data from this study could represent COS exchange from only the top layer of soil. Nonetheless, it would be enlightening to compare controlled experiments to data collected in the field.”

**Pg 21103, line 8: Should litter fluxes included in the ecosystem flux?**

This is an interesting point. We now explicitly include litter: “Ecosystem COS flux, is the sum of leaf COS uptake, and soil COS exchange, including litter.”

**Pg 21103, line 14: What does this mean? GPP greater than 25  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ? And what is the uncertainty in GPP?**

We restricted the dataset to 25  $\mu\text{mol m}^{-2} \text{s}^{-1}$  in order to avoid the complication of nighttime fluxes. We have now added the language, “to include only midday fluxes when photosynthesis was high”.

**Pg 21103, line 23: concept not conceit I assume?**

We have replaced “conceit” with “concept”.

**Pg 21104, line 1: Is there enough previous literature to justify using equation (1) in this way?**

Equation 1 has been used in at least 2 studies to connect GPP to COS uptake: Asaf et al. (2014, Nature), and Billesbach et al. (2014, Agricultural and Forest Meteorology). We have been careful to note that this calculation is an exploration of the possible impacts of soil fluxes.

**Pg 21104, line 8: Not zero but much less than the leaf uptake**

We have changed the phrasing to be more specific: “net COS exchange rates much lower than anticipated leaf COS uptake”

**Pg 21104, line 23: does increased mean more positive?**

We thank the referee for pointing this out. We have rephrased this sentence to be clearer: “COS fluxes tended towards more positive fluxes with hotter temperatures”.

**Pg 21106, line 13: Was were the range of H2O mixing ratios? Could this cause a problem with the spectral COS fit? Could the COS increased flux be an artefact of the nafion? Was the nafion tested for COS hysteresis with stable signals with both increasing and decreasing H2O?**

The Nafion tubing was directed through a container of distilled water sitting in the constant temperature water bath. Both the air that entered the soil chamber and air that bypassed the chamber were directed through the same Nafion tubing. Examining the bypass signal yielded no evidence of a COS hysteresis artifact from the Nafion tubing. That is not to say that Nafion tubing exchanges no COS; however, since both the bypass and the soil chamber air streams underwent the same treatment, using Nafion tubing as a humidifier proved an effective way of ameliorating soil drying during observations. The effect of water in the region of the spectral measurement is generally less than 1%.

**Pg 21107, line 22: Why not use sites where COS has been measured? I know of flux measurements of COS at Harvard Forest and in Finland. There may be more.**

Fortunately, Stunt Ranch and Bondville both have COS measurements which will be published soon. Willow Creek is one of the tall tower sites near where COS measurements are made by NOAA and used by continental scale modelers. The desert and tropical soils were chosen because these two biomes have, to our knowledge, no known soil measurements yet (See Table 3).

**Pg 21114, line 11: agricultural soils?**

We agree that distinguishing the soils as agricultural makes this sentence clearer. The text has been changed.

### 3. Response to Anonymous Referee #3

We thank Anonymous Referee #3 for his or her thorough and perceptive review. We have altered the figures as suggested and changed the language of the manuscript to make the ideas clearer. This has been a great help to improve the paper and we appreciate the effort.

**The topic of this manuscript is on the use of biogenic (preferentially vegetation) COS uptake as a proxy for GPP. The main assumptions behind this approach are that vegetation uptake is the dominant COS flux in land ecosystems and that soil or nonphotosynthetic fluxes are either minor or well characterized (Maseyk et al. 2014). The latter is challenged by this discussion paper, and the potential bias introduced by soil COS exchange is shown. Two GPP estimates were compared: one accounting for COS leaf uptake fluxes alone, the other additionally including soil COS exchange. While soil was generally assumed to predominantly act as sink for COS in earlier studies on a variety of soil types, few recent publications revealed the capacity of agricultural soil being a strong source of COS. Dynamic soil (agricultural, forest, desert, and savannah) incubation chambers were used to assess the COS exchange in a controlled setting in the laboratory. In general the authors confirmed the uptake of COS for most of the soils, at least for the relevant environmental conditions of their habitat. For an agricultural soy field soil type they found strong emissions, which confirm recent field flux studies, but are in contrast to earlier laboratory studies where no emission from agricultural soil was found. The reason for these discrepancies could not be unraveled. The authors used “anticipated” vegetation COS uptake data to estimate the expected bias introduced by including/excluding literature soil COS exchange data, hence showed the potential impact of soil COS exchange on GPP estimates ranging from - 220 to +119 % (Table 3). Furthermore, they used functional dependencies of soil COS emissions on temperature and soil water content, derived specifically from soy field soil, to account for both uptake and emission of COS. Applying their soil COS model to a specific agricultural soy field site (reported CO<sub>2</sub> flux measurements; no in situ COS data for vegetation or soil) the bias was comparably low (-5 to +25 %), due to the fact that the net soil COS exchange was relatively low. The subject fits well into the scope of ACP and publication is justified. The bunch of lab data are not conclusive, but rather disclose the dire need for more information about soil COS exchange. Future studies must disclose the extent of generalization of the model on COS emission from agricultural soils. There are some concerns, mainly on the reported variability of the assimilated database, and the traceability of data used for the different approaches. While solely using soy field soil results for the model is described in the Methods section, it should additionally be mentioned in the main text (beginning of section 3.2, and also mention again that the Bondville site is a soy field).**

The laboratory results are quite diverse with sometimes “no discernible pattern” or contradicting results comparing different soil types. The data on the environmental functional dependencies of soil types other than soy field soil are somewhat distracting from the main idea of the manuscript. More than one environmental parameter is changing at a time, in some cases observed to be accompanied by a change in the sign of the exchange direction. In general soils showed COS uptake at low temperature and being wet, while emission was observed at high temperatures and being dry. The different soils did not show consistent exchange values with respect to, e.g., soil moisture manipulation. However, this work demonstrates the diversity of soil COS exchange behavior. The authors do address this issue, like “the link between soil moisture and COS fluxes for soils collected at other sites is not as clear” or “the pattern of COS fluxes over time after a change in soil water content was not consistent for given changes in soil moisture.” Hence the data base is somewhat vague, and to retrieve a sound model for soil COS exchange characteristics is challenging. For the latter reason (and to avoid misuse of the model) it should specifically be pointed out that the derived model parameters are, if at all, only applicable to agricultural soil types, as the main conclusions do not necessarily hold for (or can be transferred to) other soil types. However, the authors already emphasized: “we present this as a theoretical exercise investigating the possible magnitudes of soil COS exchange on broader scales”. I agree, at least as long as not all ingredients for a GPP estimate are measured simultaneously in situ.

We agree with the referee. We hope that further studies will be able to use the data collected here as a starting point to understand soil COS flux variability. COS fluxes change after simulated precipitation, sometimes appearing like a Birch effect, sometimes having little influence on fluxes. This variability demonstrates the trouble with modeling COS fluxes. The fact that our model is based on the soy field soil has now been re-emphasized in section 3.2.

**Fig. 10: please indicate within the figure (horizontal bars or shaded areas) when wheat was present, when it was senescent, and when it was harvested to better emphasize the involvement of plants. The applied models are not meant to account for the influence of vegetation, although the wheat is assumed to have critical impact on the COS budget. It seems that not even the role of vegetation could adequately be modelled based on current knowledge (as assumed to be a sink) at this specific site, unless the soil source would be able to over-compensate the assumed COS uptake by wheat. May be this critical detail should also be discussed more clearly in the text.**

The fluxes presented in Figure 10 are from a field-based soil chamber containing no wheat. The net exchange eddy flux covariance measurements over the wheat field reported in Maseyk et. al 2014 (PNAS). This idea is re-emphasized in the figure caption: “COS soil fluxes were measured using an automatic soil chamber containing no wheat.”

**Minor corrections/comments:**



**As the COS exchange is dependent on COS mixing ratios: which inlet COS mixing ratios were used for the laboratory dynamic chamber experiments (see respective comment of Referee #1) and could it have had biased the results?**

The influence of COS mixing ratios is an important point, which we've addressed now in the Supplemental. We are grateful to both referees for bringing the issue to light.

**When discussing the “important dimension of soil depth”, the authors might consider a most recent paper by Sun et al. 2015 (A soil diffusion-reaction model for surface COS flux: COSSM v1; Geosci. Model Dev. Discuss., 8, 5139–5182, 2015, doi:10.5194/gmdd-8-5139-2015)**

Thanks for pointing out these new developments. We now refer to the two very recently published models COS soil models in the text: Sun et al. 2015 and Ogee et al. 2015.

**Page 21104, line 23: to prevent misunderstanding, please rephrase “Regardless of sign, COS fluxes increased with temperature”. To my understanding this would mean that increase of temperature on a negative flux (deposition) results in a more negative flux (stronger deposition); this is not what the authors showed in Fig. 4 and 5.**

Thanks for finding this confusing sentence. The sentence now reads: “COS fluxes tended towards more positive fluxes with hotter temperatures.”

**Page 21107, line 16: please remind the reader that the Bondville FLUXNET site, USBo1 is a soy field site.**

We now append the sentence to be clearer: “We used data reported for the Bondville FLUXNET site, US-Bo1, the soy field site in this study.”

**In Table 3, the authors compare the impact of earlier results of (anticipated plant and observed soil) COS exchange from different biomes on GPP. At first view, I would assume that accounting for exclusively soil COS uptake (as in the case of Kuhn et al. 1999) would result in an uncertainty only in one direction, not in both as indicated in the table (+119 to -34%), as “F COS ecosystem” is always higher after including soil uptake. Please correct me if I am wrong. Else: Most of the field measurement references of Table 3 do not appear in the reference list.**

Earlier studies have found both uptake and production from soils. Granted, the number of soil-only observations is small, but we included the full range of whatever data was available for this exercise.

Additionally, we thank the referee for noticing that our bibliography was incomplete. The references have now been amended to include all the citations in the text and tables.

**Please check. Fig. 6: The information on soil moisture is redundant (x-axis title and color code); however, this color code of VWC would help in Fig. 7.**

Great suggestion. We include the color code in Fig 6 to make it consistent with Figure 4 and 5. Figure 7 now retains the same color code for the COS fluxes.

**Fig. 8, legend: to be better comprehensive, I suggest “Estimated fluxes from abiotic (a) and biotic (b) processes of soil COS exchange from soy field soil.”**

We agree with the referee: this makes the figure easier to understand. It has been changed.

# Carbonyl sulfide exchange in soils for better estimates of ecosystem carbon uptake

**M. E. Whelan<sup>1,2</sup>, T. W. Hilton<sup>1</sup>, J. A. Berry<sup>2</sup>, M. Berkelhammer<sup>3</sup>, A. R. Desai<sup>4</sup>, and J. E. Campbell<sup>1</sup>**

<sup>1</sup>University of California, Merced, Merced, CA, USA

<sup>2</sup>Carnegie Institution for Science, Stanford, CA, USA

<sup>3</sup>University of Illinois Chicago, Chicago, IL, USA

<sup>4</sup>University of Wisconsin, Madison, WI, USA

Correspondence to: M. E. Whelan (marywhelan@gmail.com)

## Abstract

Carbonyl sulfide (COS) measurements are one of the emerging tools to better quantify gross primary production (GPP), the largest flux in the global carbon cycle. COS is a gas with a similar structure to CO<sub>2</sub>; COS uptake is thought to be a proxy for GPP. However, soils are a potential source or sink of COS. This study presents a framework for understanding soil-COS interactions. Excluding wetlands, most of the few observations of isolated soils that have been made show small uptake of atmospheric COS. Recently, a series of studies at an agricultural site in the central United States found soil COS production under hot conditions an order of magnitude greater than fluxes at other sites. To investigate the extent of this phenomenon, soils were collected from 5 new sites and incubated in a variety of soil moisture and temperature states. We found that soils from a desert, an oak savannah, a deciduous forest, and a rainforest exhibited small COS fluxes, behavior resembling previous studies. However, soil from an agricultural site in Illinois, > 800 km away from the initial central US study site, demonstrated comparably large soil fluxes under similar conditions. These new data suggest that, for the most part, soil COS interaction is negligible compared to plant uptake of COS. We present a model that anticipates the large agricultural soil fluxes so that they may be taken into account. While COS air-monitoring data are consistent with the dominance of plant uptake, improved interpretation of these data should incorporate the soil flux parameterizations suggested here.

## 1 Introduction

As anthropogenic CO<sub>2</sub> emissions continue increasing, it is necessary to characterize the partitioning of carbon exchange between atmospheric and terrestrial ecosystem reservoirs to predict future CO<sub>2</sub> concentrations in the atmosphere (Wofsy, 2001). Large uncertainties remain in estimates of the amount of carbon removed from the atmosphere by photosynthesis (Beer et al., 2010), called gross primary productivity (GPP). This quantity is essential for describing carbon-climate feedbacks and assessing ecosystem-based CO<sub>2</sub> capture and

storage projects. Using measurements of carbonyl sulfide is one of several emerging approaches to address large uncertainties in GPP estimates (Berry et al., 2013; Campbell et al., 2008; Commane et al., 2013; Montzka et al., 2007; [Sandoval-Soto et al., 2005](#); Seibt et al., 2010; Stimler et al., 2011; Suntharalingam et al., 2008). With a globally averaged tropospheric [concentration-mixing ratio](#) of  $500 \pm 100$  parts-per-trillion (ppt) (Montzka et al., 2007), COS is the most abundant sulfur-containing gas in Earth's atmosphere. Both COS and  $\text{CO}_2$  enter a plant through leaf stomata. Whereas some  $\text{CO}_2$  is released again in back-diffusion or in respiration, COS is irreversibly destroyed by ~~the enzyme~~ carbonic anhydrase (Protoschill-Krebs et al., 1996; Schenk et al., 2004). [Other enzymes such as RuBisCO can also destroy COS \(Protoschill-Krebs and Kesselmeier, 1992\). In soils, algal populations are expected to be smaller than bacterial populations \(Wingate et al., 2009\), and COS uptake is generally attributed to carbonic anhydrase.](#) Soil COS fluxes potentially introduce large uncertainties in estimating the COS leaf uptake flux from atmospheric COS measurements (Maseyk et al., 2014).

To date only three published studies have attempted to use COS concentrations to calculate GPP over individual ecosystems (Asaf et al., 2013; Billesbach et al., 2014; Blonquist et al., 2011). The calculation is performed using this relationship:

$$F_{\text{COS,leaf}} = \text{GPP}[\text{COS}][\text{CO}_2]^{-1}v(p, i, w) \quad (1)$$

$F_{\text{COS,leaf}}$  is the one-way flux of COS into plant leaves in  $\text{pmol m}^{-2} \text{s}^{-1}$ , GPP is the  $\text{CO}_2$  assimilation by plants in  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , [COS] and  $[\text{CO}_2]$  are ambient gas [concentrations mixing ratios](#) in parts-per-trillion (ppt) and parts-per-million (ppm) respectively, and the factor  $v$  is the experimentally determined ratio of deposition velocities for COS and  $\text{CO}_2$ , a function of plant type  $p$ , radiation  $i$ , and water stress  $w$ .

Many of the plant physiological requirements involved in using COS fluxes as a GPP proxy have been empirically investigated. Stimler et al. (2010) confirmed the assumptions about in-leaf processes and COS :  $\text{CO}_2$  exchange that need to be met to use COS as a tracer for GPP, i.e. COS co-diffuses with  $\text{CO}_2$  via the same pathway in plant leaves, COS and  $\text{CO}_2$  do not inhibit one another at reaction sites with carbonic anhydrase, and emis-

sion of COS by leaves is negligible. However, other studies have found species-specific COS emissions by plants (Geng and Mu, 2006; Whelan et al., 2013). For the most part, using COS to predict GPP on the leaf-level was comparable to other methods like  $C^{18}OO$  exchange (Seibt et al., 2010; Stimler et al., 2011).

5 However, a problem arises when the COS : CO<sub>2</sub> scheme is applied to an ecosystem beyond the leaf scale. The uptake ratio is called an ecosystem relative uptake (ERU) when the observation scale encompasses plants and soils (Campbell et al., 2008) or a soil relative uptake (SRU) when soils are observed or modeled apart from plant systems (Berkelhammer et al., 2014). Empirical measurements of ERU deviate from the value of 3 (Sandoval-Soto et al., 2005) when processes other than photosynthesis dominate trace gas exchange over  
10 an ecosystem (Seibt et al., 2010). In these cases, it is assumed that a missing source or sink of COS or CO<sub>2</sub> exchange is present in the system. At continental scales, anthropogenic sources must be taken into account (Campbell et al., 2015). In many natural ecosystems, COS exchange by soils contributes to variations in ERU.

15 Soils in terrestrial biomes usually exhibit low COS exchanges compared to uptake by plants (see review in Whelan et al., 2013). Uncoordinated, individual studies have been undertaken that incidentally quantified soil COS exchange in a limited number of biomes, often with few soil-focused measurements.

The characterization of soil COS exchange should improve the use of COS observations  
20 as a GPP proxy. Here, to better understand soil COS exchange, we collected soil samples from multiple biomes and assessed their COS fluxes in a controlled setting using dynamic incubation chambers. We further develop a framework for interpreting and anticipating soil COS fluxes based on empirical data and gas exchange theory. This model can inform the design of much needed future field experiments.

## 25 **2 Methods**

Soil samples were acquired from agricultural, forest, desert, and savannah sites (Table 1) with a variety of patterns in soil moisture and temperature (Fig. 1). Except for the Peruvian

rainforest sample, soil collection followed the same protocol. First, two 0.0225 m<sup>2</sup> representative sites were selected, one adjacent to the biome's predominant vegetation, the other a meter away. The litter layer was removed and reserved separately. Soil was then excavated from the top 0.05 m of a 0.01 m<sup>2</sup> area, double bagged, and shipped overnight to the Carnegie Institution for Science in Stanford, CA for analysis. The Peruvian rainforest sample was an amalgamation of soils from the top 0.05 m of several sites, collected by auger from the Los Amigos Biological Station in Peru. These soils were air dried, then combined before analysis. Bulk density and soil moisture content for all soils were determined by gravimetric methods. Soil pH was measured with a Corning Pinnacle 530 pH meter (Xylem Inc. White Plains, NY). Locations of sites are shown in Fig. 2.

Sites were selected to capture variability between biomes and address data needs. The Bondville site is an agricultural research station that was rotated between soybean and corn crops; at the time of sampling, soybeans were planted, but soil contained corn litter. The Stunt Ranch Fluxnet site, an oak savannah, and the Boyd Deep Canyon Reserve, to our knowledge the first desert soil investigated for COS exchange, are both located within and managed by the University of California Reserve System. The Willow Creek mature forest, Bondville Fluxnet and Southern Great Plains ARM sites are within the footprints of COS air-monitoring sites that include tall tower and airborne platforms (Montzka et al., 2007). Soil temperature and soil moisture variability for all sites is presented in Fig. 1.

Soil subsamples were placed in individual solid PFA 1 L chambers (Savillex) and weighed. Following Van Diest and Kesselmeier (2008), 75 to 80 g soil samples were used to reduce the presence of concentration gradients in the soil profile during dynamic incubation experiments. One soil subsample from the agricultural site was wet filtered through a 53 µm sieve to remove the sand-sized soil fraction before incubation. Otherwise, soils were not sieved; large pieces of loose litter were already removed when the soils were initially collected. By keeping soils whole, we reduced the influence of sample processing artifacts on our lab-based flux observations [at the expense of working with non-homogenized and therefore less reproducible samples.](#)

## 2.1 Determination of soil COS exchange

Soil fluxes of COS were determined using a dynamic, flow-through chamber approach. A commercially-available Aerodyne quantum cascade laser (QCL, Aerodyne Research, Inc., Billerica, MA, US) was used to quantify COS and CO<sub>2</sub> ~~concentrations~~ mixing ratios in the effluent of a laboratory-based apparatus (Fig. 3). Fluxes were calculated using an equation adapted from de Mello and Hines (1994):

$$F = V(C_f - C_i)m_{\text{soil}}^{-1} \quad (2)$$

$F$  is the COS or CO<sub>2</sub> exchange rate in pmol gas min<sup>-1</sup> g dry soil<sup>-1</sup>.  $C_i$  is the mixing ratio of the compound entering the chamber, determined by analyzing the gas stream bypassing the chamber headspace.  $C_f$  is the ~~concentration~~ mixing ratio of the compound exiting the 1 L PFA chamber headspace.  $V$  represents the sweep rate of the total air through the chamber, measured by the mass flow meter upstream of the QCL and converted to pmol min<sup>-1</sup>. The value  $m_{\text{soil}}$  is the amount of dry soil enclosed inside the chamber in g. The flow of the system was driven by a vacuum pump downstream of the QCL. The instrument also measured H<sub>2</sub>O and applied a ~~correction~~ for water vapor. Some, generally a less than 1% correction. Ambient laboratory air was used as the sweep gas for the incubations performed here. Observing a nitrogen stream between incubations was used to correct for instrument baseline drift. While ambient COS mixing ratios had small variation (510 ppt with 80 ppt standard deviation), some of the CO<sub>2</sub> fluxes were uninterpretable because of variations in ambient CO<sub>2</sub> ~~concentrations~~ mixing ratios,  $C_i$ . CO<sub>2</sub> fluxes that could not be distinguished from 0 are graphically presented at 0.

Each  $F$  quantification is generated from 80 min of 1 Hz air analysis. To promote soil equilibration within a dynamic headspace, air flow was directed through the chamber and the effluent analyzed for 40 min. Before and after each chamber measurement, ambient air and nitrogen gas were each analyzed for 10 min to check for baseline stability. The average COS reported over the last several minutes of chamber flow-through and bypass were corrected for instrument drift using the drift in the nitrogen (COS-free) signal, then used as  $C_f$  and  $C_i$ ,



respectively, in Eq. (2). COS fluxes are reported in pmol COS per gram of dry weight soil per minute ( $\text{pmol COS g}^{-1} \text{min}^{-1}$ ); negative values indicate uptake of COS, when  $C_f < C_i$ .

The temperature of the chamber was manipulated from 10 to 40 °C with a constant temperature water bath. For higher temperature observations of soil fluxes from the soy field soil, the incubation chamber was placed in a container of water on a hotplate. The actual soil temperature was recorded by a small, self-contained temperature data logger with a stainless steel outer casing (iButtons, Maxim Integrated, San Jose, CA, US). In order to prevent the soil from drying out during the analysis, a length of Nafion tubing was placed upstream of the chamber inside a container of distilled water in the same water bath. Even with this precaution, soil samples still dried slightly during the experiment. Samples were weighed daily, and soil moisture content was altered or maintained by adding distilled water. When water content was changed, soil samples were held at 20 °C and COS flux observations continued for at least 12 h.

## 2.2 Scaling laboratory COS measurements to compare to field observations

Performing soil incubation experiments allowed for precise manipulation of environmental variables to reveal underlying patterns in soil COS exchange. Soil in situ has an important dimension not represented by these laboratory experiments: depth. ~~Nonetheless, it would be enlightening to compare controlled experiments to data collected in the field, despite that data~~ (Ogee et al., 2015; Sun et al., 2015). Data from this study could represent COS exchange from only the top layer of soil. Nonetheless, it would be enlightening to compare controlled experiments to data collected in the field.

A further experiment was performed to estimate the relationship between laboratory, per-gram measurements and field, per-area measurements. Soy field soil was gradually added to a 20 °C incubation chamber, starting with 50 g and increasing to 300 g. While the total COS emissions increased with every soil addition, the flux per gram soil increased linearly between 50 and 100 g, then demonstrated saturation behavior with samples greater than 100 g. Thus, all fluxes were scaled up to 100 g and assumed to represent a soil footprint

equal to the area of the incubation chamber base,  $0.00779 \text{ m}^2$ . In short, fluxes were multiplied by a factor of  $(100 \text{ g}) (0.00779 \text{ m}^{-2}) (60 \text{ s min}^{-1})^{-1}$  or  $214 \text{ g min m}^{-2} \text{ s}^{-1}$ .

### 2.3 Modeling patterns in COS soil fluxes

The total net COS flux observed from the soils is thought to be the combination of abiotic and biotic fluxes.

$$F_{\text{COS,soil}} = F_{\text{COS,biotic}} + F_{\text{COS,abiotic}} \quad (3)$$

$F_{\text{COS}}$  is the net flux of COS, whereas  $F_{\text{COS,biotic}}$  and  $F_{\text{COS,abiotic}}$  represent the contribution of biotic and abiotic processes, respectively. The flux units used here were transformed as described in Sect. 2.2 from  $\text{pmol COS min}^{-1} \text{ g dry soil}^{-1}$  to  $\text{pmol COS m}^{-2} \text{ s}^{-1}$ . Two models were fitted to soy field COS soil flux observations to explain  $F_{\text{COS,biotic}}$  and  $F_{\text{COS,abiotic}}$  separately. First, dry agricultural soil COS measurements were described using an exponential equation, as in Maseyk et al. (2014).

$$F_{\text{COS,abiotic}} = \alpha \exp(\beta T_{\text{soil}}) \quad (4)$$

where  $T_{\text{soil}}$  was the temperature of the soil in  $^{\circ}\text{C}$ , and  $\alpha$  and  $\beta$  were parameters determined using the least-squares fitting approach. These driest measurements were assumed to represent the observable fluxes with the least influence from microbial uptake of COS while keeping the soil in tact. The abiotic flux contribution expressed by Eq. (4) was calculated for all soy field soil incubation experiments, then subtracted from their respective  $F_{\text{COS,soil}}$  observations to yield  $F_{\text{COS,biotic}}$ , as in Eq. (3).

To explain  $F_{\text{COS,biotic}}$ , we used a model that was originally developed for soil NO production in Behrendt et al. (2014). Previous work (Van Diest and Kesselmeier, 2008) had used a similar NO soil flux model. The overall form of the equation is the product of a power

function and an exponential function, Eqs. (5) and (6).

$$a = \ln \left( \frac{F_{\text{opt}}}{F_{\theta_g}} \right) \left( \ln \left( \frac{\theta_{\text{opt}}}{\theta_g} \right) + \left( \frac{\theta_g}{\theta_{\text{opt}}} - 1 \right) \right)^{-1} \quad (5)$$

$$F_{\text{COS,biotic}} = F_{\text{opt}} \left( \frac{\theta_i}{\theta_{\text{opt}}} \right)^a \exp \left( -a \left( \frac{\theta_i}{\theta_{\text{opt}}} - 1 \right) \right) \quad (6)$$

Here  $a$  was the curve shape constant,  $F_{\text{opt}}$  and  $F_{\theta_g}$  were the COS fluxes ( $\text{pmol COS m}^{-2} \text{ s}^{-1}$ ) at soil moistures  $\theta_{\text{opt}}$  and  $\theta_g$  (percent volumetric water content, % VWC),  $F_{\text{opt}}$  was the maximum biotic COS uptake, and  $\theta_{\text{opt}} > \theta_g$ .  $F_{\text{COS,biotic}}$  is the COS uptake for a given soil moisture  $\theta_i$  after subtracting  $F_{\text{COS,abiotic}}$  within the specified temperature range. The two models for  $F_{\text{COS,biotic}}$  and  $F_{\text{COS,abiotic}}$  could then be used to predict soil COS fluxes for a given temperature and soil moisture condition.

## 2.4 Assessing the importance of soil COS fluxes to the GPP proxy

Ecosystem COS flux,  $F_{\text{COS,ecosystem}}$ , is the sum of leaf COS uptake,  $F_{\text{COS,leaf}}$  and soil COS exchange  $F_{\text{COS,soil}}$ , [including litter](#). Two approaches were used to explore the error introduced by calculating GPP from ecosystem COS exchange without correcting for  $F_{\text{COS,soil}}$ .

The first method sought to calculate temporal variability in the relative importance of  $F_{\text{COS,soil}}$ . We used GPP estimates for the soy field FLUXNET site (US-Bo1) based on half-hourly  $\text{CO}_2$  eddy flux covariance measurements and a respiration model (Reichstein et al., 2005), restricted to values greater than  $25 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ , [to include only midday fluxes when photosynthesis was high](#).  $F_{\text{COS,leaf}}$  was anticipated from these reported GPP values, using Eq. (1) with relative uptake of 1.8 (Stimler et al., 2011), ambient [concentration-mixing ratio](#) of  $\text{CO}_2$  at 380 ppm and of COS at 500 ppt. The model described in Sect. 2.3 was used to generate  $F_{\text{COS,soil}}$  estimates from field soil moisture and temperature data collected at the site. Estimates of  $F_{\text{COS,leaf}}$  and  $F_{\text{COS,soil}}$  were then added together and used to calculate new GPP estimates with Eq. (1). The difference between the reported GPP estimates and estimates using  $F_{\text{COS,ecosystem}}$  instead of  $F_{\text{COS,leaf}}$  in Eq. (1) was then evaluated.

Secondly, we examined the spatial importance of reported  $F_{\text{COS,soil}}$  from the few values reported in the literature, relying on a similar ~~conceit~~ concept as the global calculation above. Using the biome GPP estimates from Beer et al. (2010), we back calculated anticipated estimates of  $F_{\text{COS,leaf}}$  using Eq. (1). For this purposefully simple calculation, we assume a 100 day growing season with 12 h of light per day to convert between annual estimates of GPP and field measurements calculated in  $\text{s}^{-1}$  units, though this obviously does not represent the diversity of biome carbon assimilation patterns. For each biome where data existed, a range of  $F_{\text{COS,ecosystem}}$  was calculated as the estimated  $F_{\text{COS,leaf}}$  added to the range of reported  $F_{\text{COS,soil}}$  from previous studies. A GPP estimate was then made using Eq. (1) with  $F_{\text{COS,ecosystem}}$  in place of  $F_{\text{COS,leaf}}$ . The percentage difference between the GPP estimate in Beer et al. (2010) and this new GPP estimate was then evaluated.

### 3 Results

With the exception of the soy field sample, soils investigated here exhibited net COS exchange rates ~~constrained near 0~~ much lower than anticipated leaf COS uptake, ranging from  $-8$  to  $+8 \text{ pmol COS m}^{-2} \text{ s}^{-1}$ , compared to leaf uptake rates of  $-27$  to  $-42 \text{ pmol COS m}^{-2} \text{ s}^{-1}$  (Stimler et al., 2011). The overall patterns of COS exchange over temperature and soil moisture gradients are described in Sect. 3.1. The soil samples from the soy field had the highest overall fluxes: the biotic and abiotic components of these fluxes are investigated in Sect. 3.2.

#### 3.1 COS soil flux observations

Overall, desert and rainforest samples had the smallest magnitude net COS exchange rates. The temperate forest samples showed the largest net uptake during the first trials, when the soil sample was at field soil moisture, 41 % VWC. Of the small fluxes presented in Fig. 4, temperate forest soils also had the largest net production when the soil sample was in its hottest and driest state (Fig. 4b,  $38^\circ\text{C}$  and 5 % VWC). Samples from the oak savannah

displayed variable fluxes (Fig. 4c). Observations with the soy field soil generated mostly net production of COS, often 10 times greater than fluxes from other soil samples (Fig. 5).

Regardless of sign, COS fluxes increased with temperature. COS fluxes tended towards more positive fluxes with hotter temperatures (Figs. 4 and 5). Soils incubated at 40 °C exhibited net COS production while incubations at 10 °C yielded net COS consumption in a majority of cases. Except for the desert site, the areas where these soils were collected rarely experienced such high maximum soil temperatures, if at all (Fig. 1).

The temperate forest showed the highest CO<sub>2</sub> fluxes, with increasing fluxes for increasing temperatures and soil moisture (Fig. 4e), contrasted by the small fluxes from the rainforest and desert soils (Fig. 4d). The savannah soils exhibited an optimum temperature for CO<sub>2</sub> fluxes near approximately 30 °C (Fig. 4f).

The soybean agricultural soil incubations yielded net COS emissions for the majority of trials, with a larger range than the other soils investigated: -0.04 to 0.09 pmol COS g<sup>-1</sup> min<sup>-1</sup> when incubated between 10 and 40 °C. When samples of the agricultural soil were heated further, COS net production persisted. To determine the contribution of soil organic matter in the sand-sized fraction (SSF), coarse litter > 53 μm was removed from one subsample and incubated as before. COS net emissions were higher compared to non-sieved samples at similar temperature and water content (Fig. 5).

Soil COS fluxes had a more complicated relationship with soil moisture. When soil samples were waterlogged, net COS exchange shifted towards zero compared to drier trials. For the most part, drier soils have net emissions of COS, except in the case of the varied fluxes from the oak savannah soil (Figs. 4 and 5). In oak savannah soil, increases in soil moisture led to increases in COS uptake. When soil moisture was increased further to near 40% VWC, COS exchange returned to near zero. The savannah site was expected to experience this range of soil moisture (Fig. 1). In contrast, where dry rainforest soil experienced an increase in net COS production, rainforest soil rarely experiences near 0 soil moisture (Fig. 1). Increasing water content to field levels, the rainforest soil COS exchange returned to near zero. This does not take into account the fluctuations in soil moisture and redox

potential experienced in a rainforest in situ. Temperate forest soils appear to experience net COS uptake except under very dry or unusually hot conditions (Fig. 4b).

To observe changes in COS fluxes during changes in soil moisture (i.e. as would happen in situ via precipitation), COS exchange was recorded for at least 12 h after soil moisture was changed during the course of the experiment (Fig. 6). The rainforest and savannah fluxes showed no discernible pattern in fluxes after water additions. For one series of observations with rainforest soil, the Nafion tubing was removed and the soil dried slowly over time, continuing to show little variability. In contrast, the temperate forest and soy field soils (Fig. 6a) responded with a large variability in COS fluxes after soil moisture manipulation, taking several hours to reach a consistent flux value. There was an overall negative relationship between soil moisture and net COS production for the soy field soil samples, but the link between soil moisture and COS fluxes for soils collected at other sites is not as clear.

The pattern of COS fluxes over time after a change in soil water content was not consistent for given changes in soil moisture. However, when water was added to dry soil (< 10 % VWC), many soil subsamples exhibited the pattern in Fig. 7b: CO<sub>2</sub> fluxes remained consistent while COS fluxes increased immediately after water addition, then slowly decreased over many hours. This is contrasted by Fig. 7a, where both COS and CO<sub>2</sub> fluxes demonstrate some variability after changes in water content.

### 3.2 Modeling soil COS production and consumption

Net COS fluxes were a balance of abiotic and biotic processes. If we assume that incubations of air-dried [agricultural](#) soils were representative of an abiotic COS production or desorption (less some physical limitations), we can calculate the relationship between abiotic COS production and temperature for agricultural soil (plotted in Fig. 8a). We fitted Eq. (4) to the data using a least squares approach, much like in Maseyk et al. (2014) (plotted in Fig. 8a). The resulting Eq. (7) had an  $r^2$  value of 0.9.

$$F_{\text{COS,abiotic}} = 0.437 \exp(0.0984T_{\text{soil}}) \quad (7)$$

There were more cold ( $< 15\text{ }^{\circ}\text{C}$ ) incubations performed than hot ( $> 35\text{ }^{\circ}\text{C}$ ) incubations, and some of the coldest incubations were excluded from the fit to give appropriate weight to the hottest incubations.

Subtracting the dry [soy field](#) soil signal component from all other COS incubation results, we found the biotic and physically limited flux component (Fig. 8b). The COS incubation observations were converted to  $\text{pmol m}^{-2} \text{s}^{-1}$  units, binned by incubation temperatures as  $< 20$ ,  $20\text{--}30$ , and  $> 30\text{ }^{\circ}\text{C}$ , fitted to Eq. (4) and plotted in Fig. 8b. The resulting parameters are shown in Table 2. For the purposes of generalizing the equation to any temperature and moisture content pairing,  $\theta_g$  was held constant at 35 % VWC; then the data was binned by different temperature increments to discern how  $F_{\text{opt}}$ ,  $F_{\theta_g}$ , and  $\theta_{\text{opt}}$  in Eqs. (5) and (6) change with temperature. More data needs to be collected to create a robust model; however, we think this is a worthwhile attempt at capturing variability.

$$F_{\text{opt}} = -0.00986T_{\text{soil}}^2 + 0.197T_{\text{soil}} + -9.32 \quad (8)$$

$$\theta_{\text{opt}} = 0.287T_{\text{soil}} + 14.5 \quad (9)$$

$$F_{\theta_g} = -0.0119T_{\text{soil}}^2 + 0.110T_{\text{soil}} + -1.18 \quad (10)$$

The total flux  $F_{\text{COS,soil}}$  can be calculated as the sum of fluxes generated by biotic and abiotic processes.

Using this framework of equations, we estimate the influence of large soil COS fluxes on GPP estimates. We used data reported for the Bondville FLUXNET site, US-Bo1, [the soy field site in this study](#). The model shown in Fig. 8 and described in Eqs. (3)–(10) was based on flux observations from soil collected at this site. There are well known uncertainties associated with reported GPP from flux towers (Desai et al., 2008). However, since we have no in situ measurements of COS from the site, this data is used as a starting point for calculating theoretical error potentials.

Two GPP estimates are presented in Fig. 9a: the first represents GPP estimates with COS leaf uptake fluxes alone, the second was based on theoretical net COS fluxes, including both leaf and soil COS exchange calculated with Eq. (3). The difference between the 1 day moving averages (Fig. 9b) signifies how GPP could have been over- or under-

estimated if net ecosystem COS fluxes were used as leaf uptake fluxes, ranging from  $-5$  to  $+25\%$ .

To explore the possible spatial variation in soil COS exchange influence on the GPP proxy, we perform a similar calculation (described in Sect. 2.4) using in situ soil fluxes from previous studies (Table 3). The potential error in GPP estimates based on these sparse measurements ranges from  $-220$  to  $+119\%$ . More observations and modeling soil COS exchange for different ecosystems could ameliorate this large error.

## 4 Discussion

Generally, non-wetland soils are thought to have a small COS exchange rate compared to uptake by plant leaves. This assumption is based on few chamber measurements, often by severely altering the ecosystem, e.g. extracting plants beforehand (see review in Whelan et al., 2013). During a campaign to measure COS by eddy flux covariance in Oklahoma, Billesbach et al. (2014) noticed that hot soil and particularly hot and dry soil yielded emissions of COS to the atmosphere. This is believed to be a breakdown product from thermal decomposition of soil organic matter (Maseyk et al., 2014; Whelan and Rhew, 2015). This study sought to investigate the ubiquity of this phenomenon by incubating soils from a broad range of ecosystems and under a matrix of controlled conditions. Here we have found that, as assumed previously, most soils have small COS fluxes relative to anticipated plant uptake. However, large emissions like those reported by Billesbach et al. (2014) were generated in incubations of another agricultural soil from a soy field over 800 km away (Figs. 2 and 5).

### 4.1 Mechanisms of soil COS exchange

Multiple mechanisms determined the net COS exchange from soil, which were affected by soil water content and temperature. There are three proposed abiotic processes: COS production from abiotic degradation of soil organic matter (Whelan and Rhew, 2015), the physical limitations of water restricting air exchange between soil pore spaces and the chamber



headspace (Van Diest and Kesselmeier, 2008), and adsorption/desorption of COS onto soil grains. The biotic uptake of COS by soils is theorized to be via enzymes present in the microbial community that are similarly responsible for COS uptake in plants (Kesselmeier et al., 1999; Protoschill-Krebs et al., 1996). There is no known biotic COS production mechanism in soils.

Taking these routes of COS exchange into account, we can explain qualitatively the fluxes observed here. For example, hot, dry soil appeared to produce the highest net COS emissions. Dry soil has a smaller active microbial community (Manzoni et al., 2011), and biotic uptake would be small. Higher temperatures should yield more thermal degradation of organic matter, resulting in higher COS production. In this study, when soy field soils were heated from 40 to 68 °C, COS net emissions continued, suggesting that the trace gas production here had no optimum temperature and was most likely abiotic (Conrad, 1996). Simultaneously, COS within the soil would exchange with the chamber air without the added tortuosity of water-filled pore space. The overall result is more COS produced abiotically, less COS consumed biotically, and the resulting COS excess diffusing quickly out of the soil. After wet up, the temperature response curve shifts towards a COS sink, though often retains a similar shape. When soil moisture is increased further, soil pore spaces are effectively cut off from the chamber headspace. Waterlogged, the soil exhibits COS fluxes nearer to 0 regardless of temperature. This reasoning evidently holds across the temperate forest, savannah, and agricultural soil investigated here.

The desert soil samples, however, demonstrated near zero COS exchange at field moisture and COS uptake when wetted. Since these soils are frequently hot and dry, it could be that there is not sufficient remaining organic material to abiotically degrade into COS, or there are not enough clay or silt surfaces for COS to adsorb/desorb. The behavior of the desert soil resembles the soil COS exchange observed in Van Diest and Kesselmeier (2008) and Kesselmeier et al. (1999), which both investigated exclusively sandy soils.

## 4.2 More COS generated from agricultural soil

For the agricultural soils studied here, it appears that some soil interaction produced much more COS than other soils investigated. Large COS emissions were also observed from a wheat field soil in China (Liu et al., 2010), the previously mentioned wheat field in Oklahoma (Billesbach et al., 2014; Maseyk et al., 2014; Whelan and Rhew, 2015), but not from the sandy arable soil in Germany, Finland, and China (Van Diest and Kesselmeier, 2008) where only net COS uptake was observed. While Melillo and Steudler (1989) found increases in forest soil COS production coincident with nitrogen fertilizer application, the composition of fertilizer used at the sites discussed above is unknown to us. It is unclear what is particular about the agricultural soils in the study by Van Diest and Kesselmeier (2008) that should result in only soil COS net consumption.

Two hypotheses emerge from the theoretical framework detailed above. The first is that all soils experience large COS production from thermal degradation of soil organic matter or desorption from soil surfaces, but most or all COS generated is usually consumed by in situ microbial communities. The agricultural soils collected in Oklahoma and Illinois undergo pesticide/herbicide applications and irrigation during the course of their management that may limit the diversity and size of the microbial community (Griffiths and Philippot, 2013) and the magnitude of the microbial COS sink. This idea is partially supported by Whelan and Rhew (2015), where autoclaved agricultural soils only experienced net COS production: [, though autoclaved soils are known to emit COS \(Kato et al., 2008\).](#)

The second hypothesis suggests that the accessibility of the agricultural soil organic matter allowed more abiotic COS production than in forest or savannah soils. This could also be due to agricultural land management practices, which tend to break down soil aggregates and destabilize soil organic matter (Sollins et al., 1996). Accessibility, rather than litter quality, could explain why we see a similar COS production from agricultural fields with different crop cover, i.e. wheat (Billesbach et al., 2014; Liu et al., 2010) and soy/corn (this study). However, this still does not explain the biotically-driven net COS uptake patterns

found in arable soils by Van Diest and Kesselmeier (2008) and Kesselmeier et al. (1999) which report COS fluxes that resemble more the desert soil fluxes investigated here.

These two hypothesis may both influence COS exchange simultaneously. When the course litter and sand ( $> 53 \mu\text{m}$ ) fraction was removed from a soy field soil sample, COS production increased per gram of incubated sample (Fig. 5). This implies that the origin of the COS emissions resides in the silt and clay-associated fraction of organic matter, which has been shown to consist of plant matter that has undergone some microbial processing (Six et al., 2001, 2002). The combination of microbial activities and increased accessibility of organic matter to degradation may lead to large COS emissions from soils. While these mechanisms may explain differences between managed and non-managed soil COS exchange, we still lack a hypothesis for the difference between the small sinks in European arable soils and the temperature-driven sources in US and Chinese arable soils.

### 4.3 Comparison to field observations

The draw down of COS over North America has been observed from aircraft vertical profiles, appearing to scale with GPP-based uptake of COS by plants (Campbell et al., 2008). Data presented here indicate soil COS emission was maximum during high temperature incubations, coincident with some surface temperatures observed during the North American growing season. We generated a model in Sects. 2.3 and 3.2 to calculate COS fluxes for US agricultural soils, taking these large emissions into account. Relating laboratory measurements to in situ observations has inherent problems, so we present this as a theoretical exercise investigating the possible magnitudes of soil COS exchange on broader scales.

We plotted our equation with one developed by Maseyk et al. (2014) from fluxes (Fig. 10a) and environmental parameters (Fig. 10b) recorded in situ at a wheat field in Oklahoma over the course of that study in 2012. The COS flux model developed by Kesselmeier et al. (1999) is displayed using the same input variables, assuming a constant ambient COS concentration-mixing ratios of 500 ppt and a standard flux of  $75.3 \text{ pmol m}^{-2} \text{ s}^{-1}$  (Fig. 10b). This last equation can only predict COS soil uptake and has been used to model soil COS exchange globally (Kettle et al., 2002).

Key patterns emerged from examining differences between the observations and predictions over the course of the campaign in Maseyk et al. (2014) (Fig. 10), noting first that the model presented by Kesselmeier et al. (1999) and the model presented here were not parameterized using soil from this site. The fact that there are any similarities at all between the model outputs and observations is encouraging for future modeling efforts. None of the three models captured the large emissions observed before day of year (DOY) 130 when wheat was present in the field and higher soil moisture occurred. None of the models captured the large swings from COS source to sink found during large temperature fluctuations between 110 and 115 DOY. After DOY 130, the wheat senesced and was harvested, resulting in hot and dry soils. The simple model from Maseyk et al. (2014) reproduced the COS soil flux variability better under these conditions. The Kesselmeier et al. (1999) model generated some variability, but could not predict any soil COS emissions. This study's model overlapped both the uptake model's variability during wheat senescence and the high emissions predicted by Maseyk et al. (2014) after wheat harvest.

There are several explanations for the discrepancies between models and flux observations. Both this study and the Kesselmeier et al. (1999) model were based on idealized laboratory conditions, not taking into account interactions with soil COS exchange at different depths. No doubt COS is produced or consumed in all layers of soil, not just at the surface, but soil incubations were purposefully designed to avoid these issues. Additionally, there is variability in both soil moisture and temperature even over the area of the soil plot: a heterogeneous soil may experience variations in these parameters on a small scale (Entin et al., 2000). Also, soil temperature was measured at 5 cm, generally cooler than the observed surface temperature for the site (Maseyk et al., 2014). While there was not enough variability in soil moisture and temperature to perform a similar treatment as shown in Fig. 8 for soy field soil incubations, we believe the hybrid model presented here will lead to new investigations that close the gap between lab-based COS observations and COS exchange at larger scales.

#### 4.4 Discussion: implications for uncertainty in COS-based GPP estimates

The main motivation of this work was to make progress towards better estimates of GPP. The draw down of COS over the continents appears to be associated with the uptake of carbon dioxide (Campbell et al., 2008). For some of the biomes explored here, like deserts, soil COS exchange under field conditions may actually be negligible compared to plant uptake. On the other hand, recent work has suggested that soil COS fluxes in agricultural areas might be large and need to be taken into account (Billesbach et al., 2014; Maseyk et al., 2014). The model presented in this study anticipates these agricultural soil COS fluxes using commonly measured variables. With such a correction, applying the COS-GPP tracer will be more feasible to constrain GPP estimates on regional scales.

Taking COS soil fluxes into account when estimating GPP can avoid over- and under-estimations of carbon fluxes presented in Table 3 and Fig. 9. Observations are still scarce: despite a plea for data from desert soils in 2002 by Kettle et al., we were not able to find such a study in the literature over ten years later. Boreal forest soil COS exchange estimates are represented by a single study performed at a single site in Sweden over the course of two months in 1993 (Simmons, 1999). Modeling efforts suggest large COS fluxes in the tropics (Berry et al., 2013; Suntharalingam et al., 2008) and tropical forests and savannas are associated with 60 % of global terrestrial GPP (Beer et al., 2010). However, there remains a dearth of observations in tropical latitudes.

This magnitude of avoidable error suggests that soil fluxes are not negligible; however, the uncertainty of GPP at regional to global scales is much larger. The error introduced by large soil emissions from cropland soils to COS-GPP estimates can be avoided by characterization and correction of COS fluxes. This study's approach deconvolves the production rates seen to dominate the net COS flux in Maseyk et al. (2014) and the small uptake rates observed in sandy soils by Van Diest and Kesselmeier (2008).

## 5 Conclusion

The amount of data in Table 3 suggested a dire need for more information about soil COS exchange. Here we presented a controlled study using soil from multiple ecosystems and cohesive theory for how to interpret observed soil COS fluxes. This study confirms that soil from many biomes exhibited small COS fluxes compared to estimated plant sinks. However, field studies must be conducted to determine the extent of the larger magnitude US agricultural soil COS exchange in order to quantify and correct for soil effects in GPP proxy models. The difference in COS flux behavior between [agricultural](#) soils investigated in the US and Europe also remains an open question.

A final complication arises from water stress: changes in soil moisture can cause the release of pulses of COS to the atmosphere (Fig. 7) while affecting photosynthesis and associated plant COS uptake. Additionally, COS exchange during freeze/thaw events will shed light on conditions that no field or laboratory study has yet determined. If the COS soil sink is indeed overwhelmingly microbial, water stresses will play an important role in their community diversity and function (Schimel et al., 2007), which may control the balance of COS over ecosystems.

*Acknowledgements.* The authors thank D. Chadwick, J. Thom, L. Meredith, J. Chalfant, and W. Sun for sample collection; K. MacFarlane, T. Guilderson, S. Biraud, and K. Maseyk for sampling advice and data sharing; M. Zahniser and A. Kornfeld for QCL technical support; and G. Badgley, K. Caldeira and R. Commane for data analysis suggestions. Equipment was purchased through NSF DBI grant #1 040 106. Funding for the US-WCr AmeriFlux ChEAS cluster core site was provided by the U.S. Department of Energy's Office of Science. The CFSR data was developed by NOAA's National Centers for Environmental Prediction (NCEP). The data for Fig. 1 are from NOAA's National Operational Model Archive and Distribution System (NOMADS) that is maintained at NOAA's National Climatic Data Center (NCDC). This study used the CFSRv2 hourly time series data for soil moisture (soilm1.gdas\*.grb2 files; Soil Moisture Level 1 on T382 Gaussian Grid) and soil temperature (soilt1.gdas\*.grb2 files; Soil Temperature Level 1 on T382 Gaussian Grid). The data used to generate Fig. 10 used eddy covariance data acquired by the FLUXNET community and in particular AmeriFlux (U.S. Department of Energy, Biological and Environmental Research, Terrestrial Carbon Program (DE-FG02-04ER63917 and DE-FG02-04ER63911)). We acknowledge the financial

support to the eddy covariance data harmonization provided by CarboEuropeIP, FAO-GTOS-TCO, iLEAPS, Max Planck Institute for Biogeochemistry, National Science Foundation, University of Tuscia, Université Laval and Environment Canada and US Department of Energy and the database development and technical support from Berkeley Water Center, Lawrence Berkeley National Laboratory, Microsoft Research eScience, Oak Ridge National Laboratory, University of California – Berkeley, and University of Virginia.

## References

- Anderson, R. G. and Goulden, M. L.: Relationships between climate, vegetation, and energy exchange across a montane gradient, *J. Geophys. Res.-Biogeo.*, 116, G01026, doi:10.1029/2010JG001476, 2011.
- Asaf, D., Rotenberg, E., Tatarinov, F., Dicken, U., Montzka, S. A., and Yakir, D.: Ecosystem photosynthesis inferred from measurements of carbonyl sulphide flux, *Nat. Geosci.*, 6, 186–190, doi:10.1038/ngeo1730, 2013.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A., Baldocchi, D., and Bonan, G. B.: Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate, *Science*, 329, 834–838, doi:10.1126/science.1184984, 2010.
- Behrendt, T., Veres, P. R., Ashuri, F., Song, G., Flanz, M., Mamtimin, B., Bruse, M., Williams, J., and Meixner, F. X.: Characterisation of NO production and consumption: new insights by an improved laboratory dynamic chamber technique, *Biogeosciences*, 11, 5463–5492, doi:10.5194/bg-11-5463-2014, 2014.
- Berkelhammer, M., Asaf, D., Still, C., Montzka, S., Noone, D., Gupta, M., Provencal, R., Chen, H., and Yakir, D.: Constraining surface carbon fluxes using in situ measurements of carbonyl sulfide and carbon dioxide, *Global Biogeochem. Cy.*, 28, 161–179, doi:10.1002/2013GB004644, 2014.
- Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., Seibt, U., Stimler, K., Yakir, D., and Zhu, Z.: A coupled model of the global cycles of carbonyl sulfide and CO<sub>2</sub>: a possible new window on the carbon cycle, *J. Geophys. Res.-Biogeo.*, 118, 842–852, doi:10.1002/jgrg.20068, 2013.
- Billesbach, D. P., Berry, J. A., Seibt, U., Maseyk, K., Torn, M. S., Fischer, M. L., Abu-Naser, M., and Campbell, J. E.: Growing season eddy covariance measurements of carbonyl sulfide and CO<sub>2</sub>

- fluxes: COS and CO<sub>2</sub> relationships in Southern Great Plains winter wheat, *Agr. Forest Meteorol.*, 184, 48–55, doi:10.1016/j.agrformet.2013.06.007, 2014.
- Blonquist, J. M., Montzka, S. A., Munger, J. W., Yakir, D., Desai, A. R., Dragoni, D., Griffis, T. J., Monson, R. K., Scott, R. L., and Bowling, D. R.: The potential of carbonyl sulfide as a proxy for gross primary production at flux tower sites, *J. Geophys. Res.-Biogeo.*, 116, 1–18, doi:10.1029/2011JG001723, 2011.
- Campbell, J. E., Carmichael, G. R., Chai, T., Mena-Carrasco, M., Tang, Y., Blake, D. R., Blake, N. J., Vay, S. A., Collatz, G. J., Baker, I., Berry, J. A., Montzka, S. A., Sweeney, C., Schnoor, J. L., and Stanier, C. O.: Photosynthetic control of atmospheric carbonyl sulfide during the growing season, *Science*, 322, 1085–1088, doi:10.1126/science.1164015, 2008.
- Campbell, J. E., Whelan, M. E., Seibt, U., Smith, S. J., Berry, J. A., and Hilton, T. W.: Atmospheric carbonyl sulfide sources from anthropogenic activity: implications for carbon cycle constraints, *Geophys. Res. Lett.*, 42, 3004–3010, doi:10.1002/2015GL063445, 2015.
- [Castro, M. S. and Galloway, J. N.: A comparison of sulfur-free and ambient air enclosure techniques for measuring the exchange of reduced sulfur gases between soils and the atmosphere, \*J. Geophys. Res.\*, 96, 15427–15, doi:10.1029/91JD01399, 1991.](#)
- Commane, R., Herndon, S. C., Zahniser, M. S., Lerner, B. M., McManus, J. B., Munger, J. W., Nelson, D. D., and Wofsy, S. C.: Carbonyl sulfide in the planetary boundary layer: coastal and continental influences, *J. Geophys. Res.-Atmos.*, 118, 8001–8009, doi:10.1002/jgrd.50581, 2013.
- Conrad, R.: Soil microorganisms as controllers of atmospheric trace gases (H<sub>2</sub>, CO, CH<sub>4</sub>, OCS, N<sub>2</sub>O, and NO), *Microbiol. Rev.*, 60, 609–640, 1996.
- Cook, B. D., Davis, K. J., Wang, W., Desai, A., Berger, B. W., Teclaw, R. M., Martin, J. G., Bolstad, P. V., Bakwin, P. S., Yi, C., and Heilman, W.: Carbon exchange and venting anomalies in an upland deciduous forest in northern Wisconsin, USA, *Agr. Forest Meteorol.*, 126, 271–295, doi:10.1016/j.agrformet.2004.06.008, 2004.
- De Mello, W. Z. and Hines, M. E.: Application of static and dynamic enclosures for determining dimethyl sulfide and carbonyl sulfide exchange in Sphagnum peatlands: implications for the magnitude and direction of flux, *J. Geophys. Res.*, 99, 14601–14607, 1994.
- Desai, A. R., Richardson, A. D., Moffat, A. M., Kattge, J., Hollinger, D. Y., Barr, A., Falge, E., Noormets, A., Papale, D., Reichstein, M., and Stauch, V. J.: Cross-site evaluation of eddy covariance GPP and RE decomposition techniques, *Agr. Forest Meteorol.*, 148, 821–838, doi:10.1016/j.agrformet.2007.11.012, 2008.



Entin, J. K., Robock, A., Vinnikov, K. Y., Hollinger, S. E., Liu, S., and Namkhai, A.: Temporal and spatial scales of observed soil moisture variations in the extratropics, *J. Geophys. Res.-Atmos.*, 105, 11865–11877, doi:10.1029/2000JD900051, 2000.

[Fried, A., Klinger, L. F. and Erickson, D. J.: Atmospheric carbonyl sulfide exchange in bog microcosms, \*Geophys. Res. Lett.\*, 20, 129–132, doi:199310.1029/93GL00062, 1993.](#)

Geng, C. and Mu, Y.: Carbonyl sulfide and dimethyl sulfide exchange between trees and the atmosphere, *Atmos. Environ.*, 40, 1373–1383, doi:10.1016/j.atmosenv.2005.10.023, 2006.

Griffiths, B. S. and Philippot, L.: Insights into the resistance and resilience of the soil microbial community, *FEMS Microbiol. Rev.*, 37, 112–129, doi:10.1111/j.1574-6976.2012.00343.x, 2013.

[Kato, H., Saito, M., Nagahata, Y. and Katayama, Y.: Degradation of ambient carbonyl sulfide by \*Mycobacterium\* spp. in soil, \*Microbiology\*, 154, 249–255, doi:10.1099/mic.0.2007/011213-0, 2008.](#)

Kesselmeier, J., Teusch, N., and Kuhn, U.: Controlling variables for the uptake of atmospheric carbonyl sulfide by soil, *J. Geophys. Res.*, 104, 11577–11584, doi:10.1029/1999JD900090, 1999.

Kettle, A. J., Kuhn, U., von Hobe, M., Kesselmeier, J., and Andreae, M. O.: Global budget of atmospheric carbonyl sulfide: temporal and spatial variations of the dominant sources and sinks, *J. Geophys. Res.-Atmos.*, 107, 1–16, doi:10.1029/2002JD002187, 2002.

[Kuhn, U., Ammann, C., Wolf, A., Meixner, F. X., Andreae, M. O. and Kesselmeier, J.: Carbonyl sulfide exchange on an ecosystem scale: soil represents a dominant sink for atmospheric COS, \*Atmos. Environ.\*, 33, 995–1008, 1999.](#)

Liu, J., Geng, C., Mu, Y., Zhang, Y., Xu, Z., and Wu, H.: Exchange of carbonyl sulfide (COS) between the atmosphere and various soils in China, *Biogeosciences*, 7, 753–762, doi:10.5194/bg-7-753-2010, 2010.

Manzoni, S., Schimel, J. P., and Porporato, A.: Responses of soil microbial communities to water stress: results from a meta-analysis, *Ecology*, 93, 930–938, doi:10.1890/11-0026.1, 2011.

Maseyk, K., Berry, J. A., Billesbach, D., Campbell, J. E., Torn, M. S., Zahniser, M., and Seibt, U.: Sources and sinks of carbonyl sulfide in an agricultural field in the Southern Great Plains, *P. Natl. Acad. Sci. USA*, 111, 9064–9069, doi:10.1073/pnas.1319132111, 2014.

Melillo, J. M. and Steudler, P. A.: The effect of nitrogen fertilization on the COS and CS<sub>2</sub> emissions from temperate forest soils, *J. Atmos. Chem.*, 9, 411–417, 1989.

Meyers, T. and Hollinger, S. E.: An assessment of storage terms in the surface energy balance of maize and soybean, *Agr. Forest Meteorol.*, 125, 105–115, doi:10.1016/j.agrformet.2004.03.001, 2004.

Montzka, S. A., Calvert, P., Hall, B. D., Elkins, J. W., Conway, T. J., Tans, P. P., and Sweeney, C.: On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO<sub>2</sub>, *J. Geophys. Res.-Atmos.*, 112, D09302, doi:10.1029/2006JD007665, 2007.

Ogée, J., Sauze, J., Kesselmeier, J., Genty, B., Van Diest, H., Launois, T. and Wingate, L.: [A new mechanistic framework to predict OCS fluxes from soils](#), *Biogeosciences Discuss.* 12, 15687–15736, doi:10.5194/bgd-12-15687-2015, 2015.

Protoschill-Krebs, G. and Kesselmeier, J.: [Enzymatic pathways for the consumption of carbonyl sulphide \(COS\) by higher plants](#), *Bot. Acta.* 105, 206–212, 1992.

Protoschill-Krebs, G., Wilhelm, C., and Kesselmeier, J.: Consumption of carbonyl sulphide (COS) by higher plant carbonic anhydrase (CA), *Atmos. Environ.*, 30, 3151–3156, doi:10.1016/1352-2310(96)00026-X, 1996.

Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., Granier, A., Grünwald, T., Havránková, K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J.-M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., and Valentini, R.: On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm, *Glob. Change Biol.*, 11, 1424–1439, doi:10.1111/j.1365-2486.2005.001002.x, 2005.

Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H.-Y., Juang, H.-M. H., Sela, J., Iredell, M., Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., Van Den Dool, H., Kumar, A., Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J.-K., Ebisuzaki, W., Lin, R., Xie, P., Chen, M., Zhou, S., Higgins, W., Zou, C.-Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G., and Goldberg, M.: The NCEP climate forecast system reanalysis, *B. Am. Meteorol. Soc.*, 91, 1015–1057, doi:10.1175/2010BAMS3001.1, 2010.

Sandoval-Soto, L., Stanimirov, M., von Hobe, M., Schmitt, V., Valdes, J., Wild, A., and Kesselmeier, J.: Global uptake of carbonyl sulfide (COS) by terrestrial vegetation: Estimates corrected by deposition velocities normalized to the uptake of carbon dioxide (CO<sub>2</sub>), *Biogeosciences*, 2, 125–132, doi:10.5194/bg-2-125-2005, 2005.

Schenk, S., Kesselmeier, J., and Anders, E.: How does the exchange of one oxygen atom with sulfur affect the catalytic cycle of carbonic anhydrase?, *Chem.-Eur. J.*, 10, 3091–3105, doi:10.1002/chem.200305754, 2004.

- Schimel, J., Balsler, T. C., and Wallenstein, M.: Microbial stress-response physiology and its implications for ecosystem function, *Ecology*, 88, 1386–1394, doi:10.1890/06-0219, 2007.
- Seibt, U., Kesselmeier, J., Sandoval-Soto, L., Kuhn, U., and Berry, J. A.: A kinetic analysis of leaf uptake of COS and its relation to transpiration, photosynthesis and carbon isotope fractionation, *Biogeosciences*, 7, 333–341, doi:10.5194/bg-7-333-2010, 2010.
- 5 Simmons, J. S.: Consumption of atmospheric carbonyl sulfide by coniferous boreal forest soils, *J. Geophys. Res.*, 104, 11569–11576, doi:10.1029/1999JD900149, 1999.
- Six, J., Guggenberger, G., Paustian, K., Haumaier, L., Elliott, E. T., and Zech, W.: Sources and composition of soil organic matter fractions between and within soil aggregates, *Eur. J. Soil Sci.*, 10 52, 607–618, doi:10.1046/j.1365-2389.2001.00406.x, 2001.
- Six, J., Conant, R. T., Paul, E. A., and Paustian, K.: Stabilization mechanisms of soil organic matter: implications for C-saturation of soils, *Plant Soil*, 241, 155–176, doi:10.1023/A:1016125726789, 2002.
- Sollins, P., Homann, P., and Caldwell, B. A.: Stabilization and destabilization of soil organic matter: 15 mechanisms and controls, *Geoderma*, 74, 65–105, doi:10.1016/S0016-7061(96)00036-5, 1996.
- [Steinbacher, M., Bingemer, H. G. and Schmidt, U.: Measurements of the exchange of carbonyl sulfide \(OCS\) and carbon disulfide \(CS<sub>2</sub>\) between soil and atmosphere in a spruce forest in central Germany, \*Atmos. Environ.\*, 38, 6043–6052, 2004.](#)
- Stimler, K., Montzka, S. A., Berry, J. A., Rudich, Y., and Yakir, D.: Relationships between carbonyl sulfide (COS) and CO<sub>2</sub> during leaf gas exchange, *New Phytol.*, 186, 869–878, doi:10.1111/j.1469- 20 8137.2010.03218.x, 2010.
- Stimler, K., Berry, J. A., Montzka, S. A., and Yakir, D.: Association between Carbonyl Sulfide Uptake and 18Δ during Gas Exchange in C<sub>3</sub> and C<sub>4</sub> Leaves, *Plant Physiol.*, 157, 509–517, doi:10.1104/pp.111.176578, 2011.
- 25 [Sun, W., Maseyk, K., Lett, C. and Seibt, U.: A soil diffusion–reaction model for surface COS flux: COSSM v1, \*Geosci Model Dev\*, 8, 3055–3070, doi:10.5194/gmd-8-3055-2015, 2015](#)
- Suntharalingam, P., Kettle, A. J., Montzka, S. M., and Jacob, D. J.: Global 3-D model analysis of the seasonal cycle of atmospheric carbonyl sulfide: implications for terrestrial vegetation uptake, *Geophys. Res. Lett.*, 35, L19801, doi:10.1029/2008GL034332, 2008.
- 30 Van Diest, H. and Kesselmeier, J.: Soil atmosphere exchange of carbonyl sulfide (COS) regulated by diffusivity depending on water-filled pore space, *Biogeosciences*, 5, 475–483, doi:10.5194/bg-5-475-2008, 2008.

Whelan, M. and Rhew, R.: Carbonyl sulfide produced by abiotic thermal and photo-degradation of soil organic matter from wheat field substrate, *J. Geophys. Res.-Biogeo.*, 120, 54–62, doi:10.1002/2014JG002661, 2015.

5 Whelan, M. E., Min, D.-H., and Rhew, R. C.: Salt marshes as a source of atmospheric carbonyl sulfide, *Atmos. Environ.*, 73, 131–137, doi:10.1016/j.atmosenv.2013.02.048, 2013.

[White, M. L., Zhou, Y., Russo, R. S., Mao, H., Talbot, R., Varner, R. K. and Sive, B. C.: Carbonyl sulfide exchange in a temperate loblolly pine forest grown under ambient and elevated CO<sub>2</sub>, \*Atmos Chem Phys\*, 10, 547–561, 2010.](#)

10 [Wingate, L., Ogée, J., Cuntz, M., Genty, B., Reiter, I., Seibt, U., Yakir, D., Maseyk, K., Pendall, E. G., Barbour, M. M., Mortazavi, B., Burlett, R., Peylin, P., Miller, J., Mencuccini, M., Shim, J. H., Hunt, J. and Grace, J.: The impact of soil microorganisms on the global budget of  \$\delta^{18}\text{O}\$  in atmospheric CO<sub>2</sub>, \*Proc. Natl. Acad. Sci.\*, 106, 22411–22415, doi:10.1073/pnas.0905210106, 2009.](#)

Wofsy, S. C.: Where has all the carbon gone?, *Science*, 292, 2261–2263, doi:10.1126/science.1061077, 2001.

15 [Yi, Z., Wang, X., Sheng, G., Zhang, D., Zhou, G. and Fu, J.: Soil uptake of carbonyl sulfide in subtropical forests with different successional stages in south China, \*J Geophys Res\*, 112, D08302, 2007.](#)

**Table 1.** Site descriptions for soils used in this study and soils from the site used in Billesbach et al. (2014) and Maseyk et al. (2014). Site descriptions for the fluxnet sites can be found in Meyers and Hollinger (2004), Anderson and Goulden (2011) and Cook et al. (2004). The temperature and soil moisture ranges are the maximum and minimum of ten years worth of hourly data from the Climate Forecast System Reanalysis (CFSRv2, Saha et al., 2010).


Site	Description	Bulk Density	pH	Site Temperature Range at 5 cm (C)	Soil Moisture Range
Bondville Fluxnet, US-Bo1 (40.0062° N, 88.2904° W)	Soybean/Corn	1.09	6.1	−14–33	12–46
Stunt Ranch Reserve (34.0939° N, 118.6567° W)	Oak savannah	1.11	7.0	4.2–37	13–45
Boyd Deep Canyon, US-SCd (33.6481° N, 116.3767° W)	Colorado desert	1.46	7.5	−0.23–44	12–38
Willow Creek Fluxnet, US-WCr (45.8060° N, 90.0798° W)	Deciduous forest	0.84	5.8	−22–29	9.5–42
Los Amigos Biological Station, Peru (12.5692° S, 70.1001° W)	Rainforest	0.92	3.9	14–31	15–47
Southern Great Plains ARM site, site of previous studies (36.6050° N, 97.4850° W)	Wheat field	1.14	4.2	−7.8–40	12–46

**Table 2.** Fitting parameters using Eq. (4) for soy field COS fluxes binned by temperature. See Sect. 4.2 for parameter descriptions. Fluxes are in  $\text{pmol COS m}^{-2} \text{s}^{-1}$  and soil moistures are in percentage volumetric water content (%VWC).

Temperature bin ( $^{\circ}\text{C}$ )	$F_{\text{opt}}$	$\theta_{\text{opt}}$	$F_{\theta_g}$	$\theta_g$	$r^2$
10–20	8.38	18.7	1.40	37.2	0.8
21–30	11.6	21.9	9.99	28.6	0.8
31–40	14.8	25.8	8.48	47.6	0.6


**Table 3.** The error introduced to GPP estimates when COS soil fluxes are held negligible. The % uncertainty column describes how much GPP would be overestimated, as a percentage of GPP calculated by Beer et al. (2010), if soil COS uptake determined from chamber measurements was included in the  $F_{\text{COS,leaf}}$  term. Negative values indicate underestimated GPP. Numbers reported for soil COS exchange were often based on a small number of observations, sometimes after forced removal of plants.

Biome	GPP estimated by Beer et al. (2010) in $\text{Pg C yr}^{-1}$	Biome area in $10^9$ ha	$F_{\text{COS,soil}}$ from field studies in $\text{pmol m}^{-2} \text{s}^{-1}$	Anticipated $F_{\text{COS,leaf}}$ in $\text{pmol m}^{-2} \text{s}^{-1}$	% uncertainty in GPP by neglecting soil COS	$F_{\text{COS,soil}}$ field studies
Croplands	14.8	1.35	-18 to 40	-48	+37 to -83	Post-harvest soil exchange estimate from the wheat field from Billesbach et al. (2014).
Temperate grasslands, shrublands	8.5	1.78	-13.3 to -8.8	-21	+119 to -34	Range reported in Kuhn et al. (1999) as an upper limit.
Temperate forests	9.9	1.04	-8 to 1.45	-42	+20 to -3	Range from Castro and Galloway (1991), Steinbacher et al. (2004), White et al. (2010), and Yi et al. (2007).
Boreal forests	8.3	1.37	1.2 to 3.8	-27	-5 to -14	The average and one standard deviation from plots having less than 10% vegetation cover (Simmons, 1999)
Tundra	1.6	0.56	5.27 to 27.6	-13	-42 to -220	The lower production is from De Mello and Hines (1994). The larger production value is an average estimate from Fried et al. (1993)
Deserts	6.4	2.77	No data	-6	No data	No data
Tropical savannas, grasslands	31.3	2.76	No data	-32	No data	No data
Tropical forest	40.8	1.75	No data	-102	No data	No data




**Figure 1.** The normalized concurrence of soil moisture and 5 cm depth temperature at sites where soils were collected for this study and the wheat field where the Maseyk et al. (2014) study was performed, hourly Climate Forecast System Reanalysis (CFSRv2, Saha et al., 2010) data over 2000 through 2009 from the nearest appropriate data point.







**Figure 2.** Locations of soil collection sites. The Southern Great Plains site is referred to in the discussion as the site used in Billesbach et al. (2014) and Maseyk et al. (2014), but was not used in these soil incubation experiments. For site descriptions, see Table 1.



**Figure 3.** The experimental set up for laboratory-based soil incubation experiments. The Nafion tubing was placed in a container of water and used to humidify the incoming gas stream. 3-way valves were used to switch between analyzing a nitrogen stream, the gas stream that flowed through the chamber ( $C_f$ , orientation of valves illustrated above), and the gas stream bypassing the chamber ( $C_i$ ).



**Figure 4.** CO<sub>2</sub> and COS flux observations over a range of temperatures and soil water content. See soil sample descriptions in Table 1.




**Figure 5.** COS net exchange from a soy field soil. For one series of observations, the sand-sized fraction (represented by stars) was removed from a sample by wet sieving, then incubated as before.









**Figure 6.** COS flux observations at 20 °C after soil water content manipulation. A rainforest soil sample in **(b)** was intentionally dried out by removing the Nafion tubing in the experimental set up (see Fig. 3).




**Figure 7.** COS fluxes over time after temperate soil moisture content was changed from **(a)** 10 to 22% VWC and **(b)** 2% VWC (air-dried) to 10% VWC, incubated at 20 °C.



**Figure 8.** Estimated fluxes from abiotic **(a)** and biotic **(b)** processes of soil COS exchange from soy field soil. In **(a)**, COS fluxes from the driest trials (VWC  $\approx$  6%) were related to temperature by Eq. (4). The empirically–derived relationship for soils with soil moisture content less than 20% VWC from Maseyk et al. (2014) is plotted for comparison. In **(b)**, COS fluxes from soy field soil were transformed by subtracting the anticipated driest flux using Eq. (3). A model of COS consumption, Eqs. (5) and (6), was applied to the resulting data, binned into groups of incubations  $<$  21 °C (indigo),  $>$  30 °C (yellow), and the range in between (orange). The parameters of the least squares fit for each temperature bin can be found in Table 2.



**Figure 9.** Comparing theoretical GPP estimates based on gross COS leaf fluxes vs. net ecosystem COS fluxes. **(a)** Theoretical GPP estimates based on leaf COS uptake, GPP estimates based on net ecosystem COS fluxes calculated by Eqs. (1) and (3), and their moving averages for a 24 h window. The yellow shaded region highlights the difference between COS-GPP proxy when no soil correction is included and the reported GPP. **(b)** The percentage difference between the 1 day moving average of reported GPP and the calculated COS flux-GPP estimates with modeled soil COS exchange included.



**Figure 10.** Comparing the model developed here with field observations. **(a)** Soil chamber COS flux observations and the empirically–derived relationship between COS fluxes, soil moisture, and surface temperature from Maseyk et al. (2014) and this study (Eq. 3); the model developed by Kesselmeier et al. (1999) as described in Kettle et al. (2002) adjusted for 10 pmol m<sup>-2</sup> s<sup>-1</sup> as a maximum magnitude uptake. [COS soil fluxes were measured using an automatic soil chamber containing no wheat.](#) **(b)** Environmental variables observed at the Southern Great Plains ARM site in Oklahoma from Maseyk et al. (2014).