Parameterization of oceanic whitecap fraction based on satellite observations

3

M. F. M. A. Albert¹, M. D. Anguelova², A. M. M. Manders¹, M. Schaap¹, and G. de Leeuw^{1,3,4}

6 [1]{TNO, P.O. Box 80015, 3508 TA Utrecht, The Netherlands}

7 [2]{Remote Sensing Division, Naval Research Laboratory, Washington, DC 20375}

8 [3]{Climate Research Unit, Finnish Meteorological Institute, Helsinki, Finland}

9 [4]{Department of Physics, University of Helsinki, Helsinki, Finland}

10 Correspondence to: M. F. M. A. Albert (monique.albert@tno.nl)

11

12 Abstract

In this study the utility of satellite-based whitecap fraction (W) data for the prediction of sea 13 spray aerosol (SSA) emission rates is explored. More specifically, the study aims at 14 15 evaluating how an account for natural variability of whitecaps in the W parameterization 16 would affect SSA mass flux predictions when using a sea spray source function (SSSF) based 17 on the discrete whitecap method. The starting point is a data set containing W data for 2006 18 together with matching wind speed U_{10} and sea surface temperature (SST) T. Whitecap 19 fraction W was estimated from observations of the ocean surface brightness temperature T_B by 20 satellite-borne radiometers at two frequencies (10 and 37 GHz). A global scale assessment of 21 the data set yielded approximately quadratic correlation between W and U_{10} . A new global 22 $W(U_{10})$ parameterization was developed and used to evaluate an intrinsic correlation between 23 W and U_{10} that could have been introduced while estimating W from T_B . A regional scale 24 analysis over different seasons indicated significant differences of the coefficients of regional 25 $W(U_{10})$ relationships. The effect of SST on W is explicitly accounted for in a new $W(U_{10}, T)$ parameterization. The analysis of W values obtained with the new $W(U_{10})$ and $W(U_{10}, T)$ 26 27 parameterizations indicates that the influence of secondary factors on W is for the largest part 28 embedded in the exponent of the wind speed dependence. In addition, the $W(U_{10}, T)$ 29 parameterization is capable to partially model the spread (or variability) of the satellite-based 1 *W* data. The satellite-based parameterization $W(U_{10}, T)$ was applied in an SSSF to estimate the 2 global SSA emission rate. The thus obtained SSA production rate for 2006 of 4.4×10^{12} kg yr⁻¹ 3 is within previously reported estimates, however with distinctly different spatial distribution.

4

5 1 Introduction

6 Whitecaps are the surface phenomenon of bubbles near the ocean surface. They form at wind speeds of around 3 m s⁻¹ and higher, when waves break and entrain air in the water which 7 8 subsequently breaks up into bubbles which rise to the surface (Thorpe, 1982; Monahan and 9 Ó'Muircheartaigh, 1986). The estimated annual global average of whitecap cover, i.e., the 10 fraction of the ocean surface covered with whitecaps W, is 3.4% (Blanchard, 1963). Being 11 visibly distinguishable from the rough sea surface, whitecaps are the most direct way to 12 parameterize the enhancement of many air-sea exchange processes including gas- and heat 13 transfer (Andreas, 1992; Fairall et al., 1994; Woolf, 1997; Wanninkhof et al., 2009), wave energy dissipation (Melville, 1996; Hanson and Phillips, 1999), and the production rate of sea 14 15 spray aerosols (SSA) (e.g., Blanchard, 1963; 1983; Monahan et al., 1983; O'Dowd and de 16 Leeuw, 2007, de Leeuw et al., 2011), because all these processes involve wave breaking and 17 bubbles.

18 Measurements of the whitecap fraction W are usually extracted from photographs and 19 video images collected from ships, towers, and air planes (Monahan, 1971; Asher and Wanninkhof, 1998; Callaghan and White, 2009; Kleiss and Melville, 2011). Whitecap 20 21 fraction is commonly parameterized in terms of wind speed at a reference height of 10 m, U_{10} . 22 Wind speed is the primary driving force for the formation and variability of W (Monahan and 23 Ó'Muircheartaigh, 1986; Salisbury et al., 2013, hereafter SAL13). Whitecap fractions 24 predicted with conventional $W(U_{10})$ parameterizations show a large spread between reported 25 W values (Lewis and Schwartz, 2004; Anguelova and Webster, 2006). Part of these variations 26 is due to differences in methods of extracting W from still and video images. Indeed, the 27 spread of W data has decreased in recently published in situ data sets as image processing 28 improved and data volume increased (de Leeuw et al., 2011). However, an order-of-29 magnitude scatter (spread) of W data remains, suggesting that U_{10} alone cannot fully predict the W variability. Other factors such as atmospheric stability (often expressed in terms of air-30 31 sea temperature difference) and/or sea surface temperature (SST) (Monahan and 32 Ó'Muircheartaigh, 1986), friction velocity (combining wind speed and thermal stability (e.g.,

Wu, 1988; Stramska and Petelski, 2003)), wave field (SAL13), and surfactant activity
(Callaghan et al., 2013) have been indicated to affect *W* with implications for the SSA
production. Thus, parameterizations of *W* that use different, or include additional (secondary),
forcing parameters to better model the spread of *W* data due to natural whitecap variability
have been sought (Monahan and Ó'Muircheartaigh, 1986; Zhao and Toba, 2001; GoddijnMurphy et al., 2011; Norris et al., 2013b; Ovadnevaite et al., 2014; Savelyev et al., 2014).

7 An alternative approach to address the variability of W is to use whitecap fraction 8 estimates from satellite-based observations of the sea state, because such observations provide 9 long-term global data sets which encompass a wide range of meteorological and 10 environmental conditions, as opposed to local measurement campaigns during which a limited variation of conditions is usually encountered. Brightness temperature T_B of the ocean surface 11 12 measured from satellite-based radiometers at microwave frequencies has been successfully 13 used to retrieve geophysical variables, including wind speed (Wentz, 1997; Bettenhausen et al., 2006; Meissner and Wentz, 2012). The feasibility of estimating W from T_B has also been 14 15 demonstrated (Wentz, 1983; Pandey and Kakar, 1982; Anguelova and Webster, 2006).

16 Anguelova et al. (2006; 2009) used WindSat data (Gaiser et al., 2004) to further 17 develop the method of estimating W from T_B , and compiled a database of satellite-based W 18 data accompanied with additional variables (hereafter referred to as whitecap database). An 19 early version of the whitecap database combines whitecap fraction at two frequencies (W_{10} for 20 10 GHz and W_{37} for 37 GHz), with wind speed U_{10} , wind direction U_{dir} , and SST T. Figure 1a 21 shows an example of the global W distribution from WindSat for a randomly chosen day from 22 this whitecap database. An extended version of the whitecap database was compiled later to 23 include three additional environmental variables: air temperature, significant wave height, and 24 peak wave period (Anguelova et al., 2010).

Salisbury et al. (2013) analyzed the extended whitecap database and showed that satellite-based *W* values carry a wealth of information on the variability of *W*. In particular, these authors showed that the global distribution of satellite-based *W* values differs from that obtained using a conventional $W(U_{10})$ parameterization with important implications for modeling SSA production rate in global climate models (GCMs) and chemical transport models (CTMs) (Salisbury et al., 2014). Salisbury et al. (2013) proposed a new $W(U_{10})$ parameterization in power law form using satellite-based *W* data over the entire globe for a 1 full year. They derived wind speed exponents which are approximately quadratic for different2 data sets:

3
$$W_{10} = 4.6 \times 10^{-3} \times U_{10}^{2.26}$$
; $2 < U_{10} \le 20 \text{ m s}^{-1}$,
4 $W_{37} = 3.97 \times 10^{-2} \times U_{10}^{1.59}$; $2 < U_{10} \le 20 \text{ m s}^{-1}$, (1)

where *W* is expressed in %. These exponents are significantly different from the cubic and
higher wind speed dependences proposed by Callaghan et al. (2008, hereafter CAL08):

7
$$W = 3.18 \times 10^{-3} (U_{10} - 3.70)^3$$
; $3.70 < U_{10} \le 11.25 \text{ m s}^{-1}$

8 $W = 4.82 \times 10^{-4} (U_{10} + 1.98)^3;$ 9.25 < $U_{10} \le 23.09 \text{ m s}^{-1}$ (2)

9 and Monahan and O'Muircheartaigh (1980, hereafter MOM80):

10
$$W(U_{10}) = 3.84 \times 10^{-4} U_{10}^{-3.41}$$
 (3).

The MOM80 parameterization was derived on the basis of the data sets of Monahan (1971) and Toba and Chaen (1973). Most of the wind speed values from these two data sets are up to 12 m s^{-1} with only 10% of the data points for winds up to 17 m s⁻¹. The range of SST is from 17 to 31 °C. Monahan and O'Muircheartaigh (1986) emphasized that this is a regionally specific function, but its widespread adoption in global models led to its application at wind speeds and SSTs well beyond its range of validity.

17 In this study we explore the utility of the satellite-based W data from a standpoint of 18 predicting SSA production rate. Whitecaps are used as a proxy for the amount of bubbles at 19 the ocean surface. When these bubbles burst, they generate sea spray droplets which in turn 20 transform to SSA when they equilibrate with the surroundings (Blanchard, 1983). Bursting 21 bubbles produce film and jet droplets, whereas at high wind speeds, exceeding about 9 m s⁻¹, 22 additional sea spray is directly produced as droplets which are blown off the wave crests 23 (Monahan et al., 1983). These spume droplets are larger than the bubble-mediated SSA 24 droplets (Andreas, 1992). In this study we will focus on bubble-mediated production of sea 25 spray.

Sea spray aerosols are important for the climate system because, due to the vast extent of the ocean, SSA particles are amongst the largest aerosol sources globally (de Leeuw et al., 2011). SSA particles contribute to the scattering of short-wave electromagnetic radiation and thus to their direct radiative effect on climate. Also, having high hygroscopicity, SSA

particles are a source for the formation of cloud condensation nuclei (Ghan et al., 1998; 1 2 O'Dowd et al., 1999) and as such influence cloud microphysical properties and thus exert indirect radiative effects on the climate system. While residing in the atmosphere, SSA 3 4 provide surface and volume for a range of multiphase and heterogeneous chemical processes 5 (Andreae and Crutzen, 1997). Through such chemical processes, the SSA contribute to the production of inorganic reactive halogens (Cicerone, 1981; Graedel and Keene, 1996; Keene 6 7 et al., 1999; Saiz-Lopez and von Glasow, 2012), participate in the production or destruction of 8 surface ozone (Keene et al., 1990; Barrie et al., 1988; Koop et al., 2000), and provide a sink in 9 the sulfur atmospheric cycle (Chameides and Stelson, 1992; Luria and Sievering, 1991; 10 Sievering et al., 1992; 1995).

11 The modeling of all these processes in GCMs and CTMs starts with calculation of the 12 production rate of SSA particles (termed also SSA production flux, SSA generation, or SSA 13 emission). Sea spray source function (SSSF) is used to calculate SSA production flux-the 14 number of SSA particles produced per unit of sea surface area per unit time. The most 15 commonly used SSSF, proposed by Monahan et al. (1986, hereafter M86), estimates SSA 16 emission by the indirect, bubble-mediated mechanism. Based on the discrete whitecap 17 method, the SSSF of M86 is formulated in terms of $W(U_{10})$, as defined by MOM80 (Eq. (3)), whitecap decay timescale τ , and the aerosol productivity per unit whitecap dE/dr: 18

19
$$\frac{dF}{dr_{80}} = \frac{W(U_{10})}{\tau} \frac{dE}{dr_{80}} = 1.373 \cdot U_{10}^{3.41} \cdot r_{80}^{-3} (1 + 0.057r_{80}^{1.05}) \times 10^{1.19e^{-B^2}},$$
(4)

In Eq. (4), MOM80 had used a constant value for the timescale $\tau = 3.53$ s, r_{80} is the droplet 20 21 radius at a relative humidity of 80%, and the exponent B is defined as $B = (0.38 - \lg r_{s_0})/0.65$. The term dE/dr, associated with the sea spray size distribution, determines the shape of the 22 23 SSSF (i.e., shape factor); the term W/τ is a scaling (or magnitude) factor as it links 24 predetermined SSA production per unit whitecap area with the amount of whitecapping in different regions at different seasons. Refer to Lewis and Schwartz (2004), de Leeuw et al. 25 (2011), and Callaghan (2013) for clear distinction of the discrete whitecap method from the 26 27 continuous whitecap method.

Estimates of SSA production fluxes using the discrete whitecap method still vary widely (Lewis and Schwartz, 2004; de Leeuw et al., 2011) precluding reliable estimates of the direct and indirect effects by SSA in GCMs, as well as the outcome of heterogeneous chemical reactions taking place in and on SSA particles in CTMs. The wide spread of

predicted SSA emissions is caused by a combination of uncertainties coming from both the 1 2 magnitude and the shape factors of the used SSSFs. The uncertainties associated with the 3 magnitude factor include difficulties of measuring W and τ and their natural variability, which affects the $W(U_{10})$ parameterizations. The assumptions of the discrete whitecap method 4 (detailed in Sect. 2.4) also contribute to the uncertainty. Added to these are the uncertainties 5 associated with the shape factor, such as its natural variability and the model chosen to 6 7 parameterize the SSA size distribution. A source of uncertainty is the difficulty of directly 8 measuring SSA fluxes which are used to develop and/or constrain SSSFs. When 9 measurements of SSA concentrations are used to develop an SSSF, uncertainty comes from 10 the deposition velocity model used to convert the concentrations to fluxes (e.g., Smith et al., 11 1993; Savelyev et al., 2014).

12 Aside from addressing uncertainties due to sea-spray measuring techniques, there are two possible ways to improve the performance of a whitecap-based SSSF as regards the 13 physical processes involved. One way is to address variations and uncertainties in the size-14 15 resolved productivity dE/dr_{80} (i.e., the shape factor in the SSSF), for instance by including the organic matter contribution to SSA at sub-micron sizes (O'Dowd et al., 2004; Albert et al., 16 17 2012) and/or by accounting for its variations with environmental factors instead of keeping it 18 constant for all conditions (de Leeuw et al., 2011, Norris et al., 2013a; Savelyev et al., 2014). 19 Another way is to address the variations and uncertainties in the whitecap fraction W and 20 timescale τ (i.e., the magnitude factor in the SSSF) by steady improvements of the W and τ 21 measurements and by accounting for their natural variability. Both approaches are expected to 22 reduce, or at least to better account for, the variations and uncertainties in parameterizing SSA 23 flux.

24 Here we report on a study investigating the second of these two routes, namely-how 25 using W data, which carry information for secondary factors, would influence the SSA 26 production flux. The objective is to assess how much of the uncertainty in the SSA flux can 27 be explained with the natural variability of W. Using the early version of the whitecap 28 database (consisting of data for W_{10} , W_{37} , U_{10} , U_{dir} and T), we parameterize the W variability in terms of U_{10} and T. Our approach (Sect. 2) involves three steps. We first assess the 29 30 satellite-based whitecap database to evaluate the wind speed dependence of W over as wide a range of U_{10} values as possible (sect. 3.1.1). In assessing the W database, we also evaluate: (i) 31 32 the impact of an intrinsic correlation between W and U_{10} , which could have been introduced

in the process of estimating W from T_B (Sect. 3.1.2); (ii) the influence of the wave field on W 1 2 variability using rising and waning wind speeds as proxy for wave development (Sect. 3.1.3). The $W(U_{10})$ expression resulting from this analysis adjusts the trend of W with U_{10} to the 3 concerted, globally-averaged influence of all secondary factors implicitly. We next apply the 4 5 established $W(U_{10})$ expression to W data on regional scales in order to assess the variability caused by secondary factors in different locations during different seasons (Sect. 3.2). We 6 7 analyze the regional variations of W, remaining after the implicit adjustment with the $W(U_{10})$ 8 expression, and parameterize them explicitly in terms of SST. The new $W(U_{10}, T)$ 9 parameterization is compared to $W(U_{10})$ of MOM80 and SAL13 (Sect. 3.3) in order to assess 10 to what extent SST can account for the W variability. Finally, the new $W(U_{10}, T)$ 11 parameterization is used to estimate SSA emissions and compare results to previous 12 predictions of SSA emissions (Sect. 3.4).

13 2 Methods

To achieve the study objective formulated above, the main task is to develop a parameterization of W that accounts for both the trend and the spread of the W data. Expressions $W(U_{10})$ model (predict) the trend of the whitecap fraction with wind speed. The inclusion of additional variables in $W(U_{10})$ relationships should be able to model (predict) the spread of the W data caused by natural variability. The approach described below aims at deriving an expression $W(U_{10}, T)$ that fulfils these two requirements.

20 **2.1** Approach to derive whitecap fraction parameterization

21 Reasoning on a series of questions shaped our approach to parameterizing W and justified the 22 choices we made for its implementation (Sect. 2.3). We first considered, Why do we need to 23 parameterize W instead of using satellite-based W data directly? A major benefit of using satellite-based W data directly in an SSSF is that these data reflect the amount and persistence 24 25 of whitecaps as they are formed by both primary and secondary forcing factors acting at a given location. This approach limits the uncertainty to that of estimating W from satellite 26 27 measurements and does not add uncertainty from deriving an expression for $W(U_{10})$ or $W(U_{10})$. T, etc.). However, such an approach would limit global predictions of SSA emissions to 28 monthly values because a satellite-based W data set does not provide daily global coverage; 29 i.e., one would need data like those in Fig. 1a for at least two weeks (and more for good 30 estimates of the uncertainties) in order to have full coverage of the globe. 31

1 Alternatively, a parameterization of whitecap fraction derived from satellite-based W 2 data can provide daily estimates of SSA emissions using readily available daily data of wind speed and other variables. Importantly, such a parameterization will be globally applicable 3 4 because the whitecap fraction data cover the full range of meteorological conditions 5 encountered over most of the world oceans. Because the availability of a large number of W data would ensure low error in the derivations of the $W(U_{10})$ or $W(U_{10}, T, \text{etc.})$ expressions, 6 7 we proceed with deriving a parameterization for W using the data in the whitecap database 8 (Sect. 2.2.1).

9 The next question to consider was, How to account for the influence of secondary 10 factors? Generally, to fully account for the variability of whitecap fraction, a parameterization 11 of *W* would involve wind speed and many additional forcings explicitly to derive an 12 expression $W(U_{10}, T, \text{ etc.})$ (MOM80; Monahan and Ó'Muircheartaigh, 1986; Anguelova and 13 Webster, 2006). Using the early version of the whitecap database in this study, we start with 14 parameterization $W(U_{10}, T)$.

15 The question that arises next is, How to combine the different dependences of W? One 16 possibility is to use a single-variable regression to extract the W dependence on each variable 17 separately, e.g., $W(U_{10})$ and W(T). Then, these can be combined to derive an expression for 18 their effects in concert, e.g., $W(U_{10}, T) = W(U_{10})W(T)$. While variables like T, atmospheric 19 stability, surfactants, etc. influence W, they do not cause whitecapping. So a parameterization 20 formulated with dedicated W(T) and other expressions may put undue weight on such 21 influences. This approach can be pursued when we have enough information to judge the relative importance of each influence (e.g., Anguelova et al., 2010, their Fig. 6) and include it 22 23 in a combined expression with a respective weighting factor.

Previous experience points to another possibility to combine causal variables like U_{10} and influential variables like *T* and the likes. The Monahan and O'Muircheartaigh (1986) analysis of five data sets showed that the variability of *W* caused by SST (and the atmospheric stability) affect significantly the coefficients in the wind speed dependence $W(U_{10})$, especially the wind speed exponent. The survey of $W(U_{10})$ parameterizations by Anguelova and Webster (2006, their Tables 1 and 2) also clearly shows that each campaign conducted in different regions and conditions comes up with a specific wind speed exponent. This strongly suggests 1 that the influence of secondary factors is implicitly expressed as a change of the wind speed 2 exponent. On the basis of their principal component analysis, SAL13 also suggested that in 3 describing the *W* variability, it is more effective to combine individual variables with wind 4 speed. On this ground, we proceed to obtain $W(U_{10}, T)$ as a wind speed dependence $W(U_{10})$ 5 whose regression (or parametric) coefficients vary with SST.

6 How to realize this goal knowing that the satellite-based W data carry information for 7 the effect of U_{10} and all other factors? One possible way to proceed is to: (i) express the mean 8 trend in the W data associated with the globally-averaged conditions of U_{10} and all other factors; then (ii) quantify the fluctuations of regional W data around this mean trend as a 9 10 function of a specific secondary factor. Here step (i) implicitly accounts for the effects of all 11 secondary factors on W, while step (ii) quantifies explicitly the effect of a given factor on W. 12 That is, the explicit formulation of the parametric coefficients accounts only partially for the 13 full effect of a given secondary factor; it adds to the implicit account via the mean trend of W 14 with U_{10} . To realize this concept, we first analyze the global W data set to identify a general 15 wind speed dependence $W(U_{10})$ for the mean trend. Then, our analysis of regional W data helps to asses to what extent can SST account for the variations of the regression coefficients 16 17 in a $W(U_{10})$ dependence.

18 The important question now is, What functional form should we use for the general 19 (mean) $W(U_{10})$ dependence? Equations (1)-(3) exemplify the functional forms usually 20 employed to express $W(U_{10})$:

$$21 \qquad W = a U_{10}^n \tag{5a}$$

22
$$W = a(U_{10} + b)^3$$
 (5b).

A general $W(U_{10})$ dependence derived using Eq. (5a) would provide an empirical wind speed exponent *n* determined from available data sets, as MOM80 did using the available data sets at the time (Sect. 1). The wider the range of conditions represented by the data sets is, the closer the resulting $W(U_{10})$ dependence would be to average conditions globally and seasonally.

A general $W(U_{10})$ dependence derived using Eq. (5b) would provide a physicallybased wind speed exponent n = 3 consistent with dimensional (scaling) arguments. Namely, because W is related to the rate at which the wind supplies energy to the sea, W should be proportional to the cube of the friction velocity u_* (Monahan and O'Muircheartaigh, 1986; Wu, 1988). On this basis, Monahan and Lu (1990) related $W^{1/3}$ to U_{10} and derived the cubic power law in Eq. (5b). Subsequently, this relationship was used successfully in whitecap data analyses (e.g., Asher and Wanninkhof, 1998; CAL08). Coefficient *b* in Eq. (5b) is included because it is preferable for a $W(U_{10})$ relationship to involve a finite *y*-intercept (Monahan and O'Muircheartaigh, 1986). A negative *y*-intercept determines *b* from the *x*-intercept and is usually interpreted as the threshold wind speed for whitecap inception.

8 A modified version of Eq. (5) combines the merits of both formulations into the form: 9 $W = a(U_{10} + b)^n$ (6)

where the wind speed exponent is adjustable (i.e., a free parameter) and a finite *y*-intercept is included. A general $W(U_{10})$ dependence derived using Eq. (6) would provide a wind speed exponent as dictated by the whitecap database. Any of the three formulations (Eqs. (5 and 6)) can produce a viable general $W(U_{10})$ dependence, the empirical ones representative of the average conditions of the world oceans and the physical one supported by sound reasoning.

15 **2.2 Data sets**

To implement the approach thus formulated, we use the whitecap database on a global scale 16 17 for the general $W(U_{10})$ dependence, and regional W subsets extracted from the whitecap 18 database for the SST analysis. In describing the data sets used, we start with the whitecap 19 database (Sect. 2.2.1). The considerations given to extract regional data sets from it are 20 described in Sect. 2.2.2. We also introduce the data from the European Centre for Medium 21 range Weather Forecasting (ECMWF) used in this study as an independent source to 22 investigate possible intrinsic correlation among the entries of the whitecap database (Sect. 2.2.3). 23

24 2.2.1 Whitecap database

Anguelova and Webster (2006) describe in detail the general concept of estimating the whitecap fraction *W* from measurements of the brightness temperature T_B of the ocean surface with satellite-borne microwave radiometers. Salisbury et al. (2013) describe the basic points of the algorithm estimating *W* (hereafter referred to as the *W*(T_B) algorithm). Briefly, the algorithm obtains *W* by using measured T_B data for the composite emissivity of the ocean 1 surface and modeled T_B data for the emissivity of the rough sea surface and areas that are 2 covered with foam (Bettenhausen et al., 2006; Anguelova and Gaiser, 2013). An atmospheric 3 model is necessary to evaluate the contribution from the atmosphere to T_B . Minimization of 4 the differences between the measured and modeled T_B data in the $W(T_B)$ algorithm ensures 5 minimal dependence of the *W* estimates on model assumptions and input variables.

6 Wind speed U_{10} is one of the required inputs to the atmospheric, roughness and foam 7 models (Anguelova and Webster, 2006; Salisbury et al., 2013). Wind speed data come from 8 the SeaWinds scatterometer on the QuikSCAT platform or from the Global Data Assimilation 9 System (GDAS), whichever matches up better with the WindSat data in time and space within 60 min and 25 km; hereafter we refer to both QuikSCAT or GDAS wind speed values as U_{10} 10 from QuikSCAT or $U_{100SCAT}$. The use of $U_{100SCAT}$ in the estimates of satellite-based W is 11 12 anticipated to lead to some intrinsic correlation when/if a relationship between W and 13 $U_{10OSCAT}$ is sought.

14 The W data used in this study are obtained from T_B at 10 and 37 GHz, W_{10} and W_{37} ; 15 data for 37 GHz are shown in Fig. 1a. The W_{10} and W_{37} data approximately represent different 16 stages of the whitecaps because of different sensitivity of microwave frequencies to foam 17 thickness (Anguelova and Gaiser, 2011). Data W_{10} are an upper limit for predominantly active 18 wave breaking (stage A whitecaps (Monahan and Woolf, 1989)) partially mixed with 19 decaying (stage B) whitecaps, while W_{37} data quantify both active and decaying whitecaps. 20 Because decaying foam covers a much larger area of the ocean surface than active whitecaps 21 (Monahan and Woolf, 1989), W_{37} data are usually larger than W_{10} data. Comparisons to 22 historic and contemporary in situ W data in Fig. 1b confirm the approximate representations 23 of stage A whitecaps (cyan squares) and A + B whitecaps (blue diamonds) by W_{10} (green) and 24 W_{37} (magenta), respectively. Anguelova et al. (2009) have quantified the differences between 25 satellite-based and in situ W data using both previously published measurements and timespace match-ups of W and discussed possible reasons for the discrepancies. 26

The satellite-based *W* data are gridded into a $0.5^{\circ} \times 0.5^{\circ}$ grid cell together with the variables accompanying each *W* data point, namely $U_{10QSCAT}$, *T* from GDAS, time (average of the times of all samples falling in each grid cell), and statistical data generated during the gridding including the root-mean-square (rms) error, standard deviation (SD), and count (the number of individual samples in a satellite footprint averaged to obtain the daily mean *W* for a grid cell). In this study, we used daily match-ups of *W*, U_{10} , and *T* data for each grid cell for

1 the year 2006. To reiterate, this data set—consisting of W_{10} and W_{37} accompanied with three 2 environmental variables (U_{10} , U_{dir} and T)—is an early version of the whitecap database; the extended W database used by SAL13 (Sect. 1) contains three additional variables suitable to 3 4 quantify explicitly the effects of wave field and atmospheric stability on W. Due to large data 5 gaps in both space and time, the daily W data cannot be interpolated to provide better coverage (Fig. 1a). Therefore, only the available data are used without filling the gaps for 6 7 areas where data are lacking. This global data set was used to assess the globally averaged 8 wind speed dependence of W.

9 2.2.2 Regional data sets

The annual global *W* distributions show regions with valid data points ranging from 100 to 300 samples per grid cell per year when both ascending and descending satellite passes are considered. Thus, different regions were selected using two criteria, namely (i) consider regions with a high number of valid data points, and (ii) obtain a selection representative of different conditions in the northern and southern hemispheres (NH and SH).

15 With these criteria, 12 regions of interest were selected (Fig. 2) and W, U_{10} , and T data for each region were extracted from the whitecap database. The coordinates of the selected 16 17 regions are listed in Table 1, together with the corresponding number of samples (data points) 18 and minimum, maximum, mean, and median values for wind speed and SST for January and 19 July. For 90% of the regional and monthly data used in the study, the percent difference (PD, 20 defined as the difference between two values divided by the average of the two values) 21 between mean and median values of U_{10} and T is less than 4% and 9.5%, respectively. With medians and means approximately the same, the U_{10} and T data have normal distributions; 22 i.e., outliers, though existing, do not affect the mean values significantly. All analyses 23 24 presented here use the mean U_{10} and T values.

Figure 3 shows the seasonal cycles of the mean U_{10} and T values for four of the selected regions (4, 5, 6, and 12) chosen to visualize the full range of regional variations of U_{10} and T data. With the large number of samples, the mean U_{10} and T values plotted in Fig. 3 are determined within 95% confidence interval (CI) of the order of 10^{-2} (Table 1). That is, any uncertainty due to sampling is removed, and Fig. 3 represents well seasonal variations, which we will use in our analyses. Variability of SST within each region is visualized with error bars (\pm one SD) in Fig. 3b. The distinct regional SST variations suggest the effect of SST can be discerned with our data and thus used to parameterize the effect of SST on *W*. The variability of U_{10} within regions is higher (wider error bars are not plotted to avoid clutter), which suggests that the use of the global data set to obtain a generalized wind speed dependence of W (Sect. 2.1) is reasonable.

5 Regions 2-11 are all in the open ocean; region 1 was selected for its landlocked position 6 (Fig. 2). Region 6 in the Pacific Doldrums is used as a reference for the lower limit of U_{10} (Fig. 3a), while region 12 is included to represent the lowest T values (Fig. 3b). Four regions 7 8 (2, 3, 7, and 8) are at latitudes between 0 and 30°S and N (Tropics and Subtropics) 9 representing the Trade winds zone. These are regions with persistent (Easterly) winds blowing over approximately the same fetches (except region 8) in oceans with different 10 11 salinity (Tang et al., 2014) and primary production (Falkowski et al., 1998) (a proxy for surfactant concentrations). Region 4 is in the NH temperate zone representing long-fetched 12 Westerly winds. Region 5 covers the latitudes between 40°S and 50°S known as "The 13 14 Roaring Forties" for the strong Westerly winds there, but is characterized with shorter fetch. 15 Differences in the seasonal cycles of U_{10} and T in regions 4 and 5 (Fig. 3) suggest more 16 uniform conditions and longer fetches in the SH temperate zone. We have chosen regions 8 17 and 9 to represent different zonal conditions and to gauge the effect of narrower range of SST 18 variations (as compared to the SST range in region 5). Chosen at the same latitude, regions 9-19 11 have approximately the same SST, salinity, and surfactants but represent different wind 20 fetches, shortest for region 9 and longest for region 11. Overall, the chosen regions cover the 21 full range of global oceanic conditions and, while representative of diverse regional 22 conditions, each one has distinct regional characteristics.

23 2.2.3 Independent data source

24 Ideally, when deriving a $W(U_{10})$ parameterization, the data for W and U_{10} should come from 25 independent sources. The intrinsic correlation between W and U_{10} that might have arisen from 26 the use of U_{10} from QuikSCAT in the estimates of W from T_B (Sect. 2.2.1), might affect the 27 relationship between W and U_{10} developed here. To evaluate the magnitude of such intrinsic 28 correlation, we used U_{10} from the ECMWF ($U_{10ECMWF}$), which is considered to be a more 29 independent source. Note though that even the ECMWF data are generated by assimilating 30 observational data sets (e.g., from buoys) in a coupled atmosphere-wave model (Goddijn-31 Murphy et al., 2011).

1 To compile this "independent" data set, we made time-space matchups between the 2 gridded W_{10} and W_{37} data and $U_{10\text{ECMWF}}$ from the 3-hourly ECMWF data for 2006. For each $W-U_{100SCAT}$ pair at a time t from the original W database, there is a corresponding W-3 4 $U_{10\text{ECMWF}}$ pair of data within an interval $t \pm 1.5$ h. This matching procedure differs from the $W-U_{10OSCAT}$ matching which was done at the WindSat swath resolution, before gridding the 5 6 variables for the whitecap database. To speed up calculations, and because this already 7 provides a statistically significant amount of data, we used only ascending satellite overpasses. Wind speeds above 35 m s⁻¹ were discarded. Besides ECMWF wind data, for 8 consistency we also extracted ECMWF SST values. 9

Figure 4a shows all ECMWF wind speed data that have been matched in time and space with the available $U_{10QSCAT}$ data for March 2006. The majority of the data is clustered in the range of 5-10 m s⁻¹ (dark red). To characterize the difference between the two wind speed sources, the correlation between U_{10} from ECMWF and U_{10} from QuikSCAT was determined as the best linear fit forced through zero:

15
$$U_{10ECMWF} = 0.953 U_{10QSCAT}$$
 (7)

16 with a coefficient of determination $R^2 = 0.824$. For comparison, the unconstrained fit between 17 $U_{10QSCAT}$ and $U_{10ECMWF}$ is also shown in Fig. 4a (dashed line); both fits are very close (they 18 almost overlap) with identical correlation coefficients ($R^2 = 0.824$ for the unconstrained fit). 19 Similarly, Fig. 4b compares *T* from ECMWF and GDAS showing almost 1:1 correlation. That 20 is, the two data sources provide almost the same values for *T*.

On average, U_{10} from ECMWF is about 5% lower than U_{10} from QuikSCAT. This U_{10} difference can be explained to some extent with the effect of atmospheric stability because QuikSCAT provides equivalent neutral wind which accounts for the stability effects on the wind profile (Kara et al., 2008; Paget et al., 2015), while the ECMWF model gives stability dependent wind speeds (Chelton and Freilich, 2005).

Having the correlation between U_{10} from the whitecap database and U_{10} from the ECMWF quantified (as well as for *T*), one can evaluate differences caused by the use of different data sources. Equation (7) could also be useful when one decides to use ECMWF data because of their availability at 6 or 3 h intervals as compared to the availability of *W*, U_{10} , and *T* match-ups twice a day (Sect. 2.2.1).

1 2.3 Implementation

We aim to develop an expression capable of modeling both the trend of the satellite-based Wdata with U_{10} and their spread (see green and magenta symbols in Fig. 1b).

4 2.3.1 Adjusting the wind speed exponent

5 We first analyze the satellite-based W data to derive a general $W(U_{10})$ expression (i.e., the 6 trend of W with U_{10}). We apply Eq. (6) with coefficients (n, a, b) left as free parameters to 7 global data sets of W_{10} , W_{37} , and both together $(W_{10} \& W_{37})$. Table 2 shows the results for the 8 regression coefficients determined from the fitting procedure within 95% CI. Each set of 9 coefficients (n, a, b) accounts implicitly for U_{10} and all secondary factors.

10 To consistently interpret and explicitly quantify regional and seasonal variations of W data, it is necessary to analyze all W data—global, regional and at different frequencies—with 11 12 the same mean trend given by the $W(U_{10})$ expression. Because Table 2 shows different wind speed exponents, we need to establish a general (unifying) *n* value. With sampling uncertainty 13 removed from the determination of these n values (see the 95% CIs in Table 2), we now 14 15 investigate the variations of the wind speed exponents among data sets. The mean of the free-16 parameter n values in Table 2 is 1.82 with lower and upper limits of the 95% CI of 0.88 and 17 2.77, respectively. A value of n = 2 is within this 95% CI and is thus a reasonable choice for 18 such general (unifying) wind speed exponent. We further verified such a choice by applying 19 two-sample *t*-test for equal means to the *n* values in Table 2 and n = 2. The *t*-test showed that 20 the mean of the wind speed exponents n determined as free parameters is not statistically 21 different from n = 2 (p > 0.05). On this ground, we adjust the free-parameter wind speed 22 exponents to n = 2, a quadratic wind speed dependence of W.

Quadratic wind speed dependence here is not unprecedented. The first reported $W(U_{10})$ relationship of Blanchard (1963) was quadratic. With careful statistical considerations, Bondur and Sharkov (1982) derived a quadratic $W(U_{10})$ relationship for residual W (strip-like structures, in their terminology). Parameterizations of W in waters with different SST have also resulted in wind speed exponents around 2 (see Table 1 in Anguelova and Webster, 2006). Quadratic wind speed dependence is also consistent with the wind speed exponents of SAL13 in Eq. (1). 1 With the adjustment of the free parameter *n* in Eq. (6) to a general (unifying) wind 2 speed exponent n = 2 and for all subsequent analyses we use a functional form for $W(U_{10})$ 3 modified from Eq. (6) to:

4
$$W = a(U_{10} + b)^2$$
 (8a).

5 Following Monahan and Lu (1990), we derive an expression $W(U_{10})$ in the form of Eq. (8a)

6 by plotting $W^{1/2}$ as a function of $U_{10QSCAT}$. Applying linear regression, we find an expression:

7
$$W^{1/2} = mU_{10} + c$$
 (8b)

which is then rearranged and squared to provide coefficients $a = m^2$ and b = c/m in Eq. (8a) 8 (results in Sect. 3.1.1). All linear fits are done on the W data associated with U_{10} from 3 to 20 9 m s⁻¹. The lower limit of 3 m s⁻¹ is chosen as a threshold for observing whitecaps. This 10 restriction is reasonable in light of the SAL13 analysis in which W data with a relative 11 standard deviation $(\sigma_w/W) > 2$ were removed: the discarded W data were about 10% of all W 12 data, mostly in regions with low wind speeds of around 3 m s⁻¹. We exclude the high wind 13 14 speed regime in order to avoid uncertainty due to (i) fewer data points in this regime; and (ii) 15 anticipated larger uncertainty in the W data from the $W(T_B)$ algorithm.

16 **2.3.2 Intrinsic correlation analysis**

For the intrinsic correlation analysis, the $W-U_{10ECMWF}$ data pairs are used in a similar fashion to make $W^{1/2}(U_{10ECMWF})$ linear fits and derive from them a relationship between the satellitebased W data and the ECMWF wind speeds. The two global $W(U_{10})$ parameterizations for the two wind speed sources are then compared to evaluate the magnitude of the intrinsic correlation (results in Sect. 3.1.2).

22 Because Eq. (7) gives the possibility to evaluate discrepancies due to the use of 23 different sources for U_{10} and T, we use U_{10} and T from the whitecap database in all 24 subsequent analyses and results. In this way, with the intrinsic correlation characterized, we restrict the uncertainty in our analyses by using the close matching-up of W, U_{10} , and T data in 25 26 the whitecap database. This decision is reasonable considering that both data sets can be used 27 in practice for different applications. The collocated data in the whitecap database (involving 28 QuikSCAT) are most suitable for analysis (as done in this study). Meanwhile, W data from 29 the whitecap database combined with forcing data from a global model (such as ECMWF or 30 other) are useful for forecasts and climate simulations.

1 2.3.3 Regional analysis

With n = 2 for the general wind speed dependence determined, we then apply Eq. (8b) to the regional monthly sub-sets of W_{10} and W_{37} data. All available data per month were used, ranging from 22 to 31 days of data. Once again, scatter plots of $W^{1/2}(U_{10})$ were generated and the best linear fits were determined providing coefficients *m* and *c* for each region for each month for W_{10} and W_{37} . The regional and seasonal variations of coefficients *a* and *b* are analyzed to inform us how to parameterize them in terms of SST, a(T) and b(T) (results in Sect. 3.2).

9 To quantify how a(T) and b(T) are influenced by different wind speed dependences— 10 our empirically determined (adjusted) wind speed exponent n = 2 (Eq. (8a)) or the physically 11 reasoned cubic wind speed dependence (Eq. (5b))—we also analyzed scatter plots of 12 $W^{1/3}(U_{10})$ and derived a respective set of coefficients a(T) and b(T).

We quantify differences between new and previously published parameterizations with two metrics (results in Sect. 3.3): (i) the PD between *W* values obtained with different parameterizations; and (ii) significance tests (Student *t*-test and ANOVA) of the differences between *W* values obtained with new and previous *W* parameterizations.

17 **2.3.4 Wave field analysis**

Efforts to include wave parameters in *W* parameterizations are well justified because, after wind speed, the most important secondary factor that accounts for variability in *W* is the wave field (SAL13). Lacking wave characteristics, the early version of the whitecap database is not suitable for deriving an explicit expression for the wave field influence on *W*. However, we have investigated the effect of rising and waning winds on the $W(U_{10})$ relationship (results in Sect. 3.1.3); increasing-decreasing winds are considered a proxy for undeveloped-developed seas (Stramska and Petelski, 2003; CAL08).

It is not feasible to determine whether winds are rising or waning from satellite-based wind speed data because of their low temporal resolution: twice a day at a given location. As wind speed provided by ECMWF is available every three hours, $U_{10ECMWF}$ values were used to examine the wind conditions at the satellite overpass time associated with a *W* data point. Wind speed difference between two three-hours intervals ΔU_{10} has been used to detect changing winds. Wind speed differences ΔU_{10} from of 1 to 5 m s⁻¹ in steps of 1 m s⁻¹ were 1 used to examine the sensitivity of the analysis to the choice of ΔU_{10} in identifying rising or 2 waning winds. Higher ΔU_{10} values are associated with the passage of stronger atmospheric 3 low-pressure systems, which come with higher wind speeds and thus stronger wind forcing of 4 waves. The $U_{10QSCAT}$ values were correlated with W using Eq. (8). Only data for 37 GHz from 5 the ascending satellite overpass were used.

6 2.4 Estimation of sea spray aerosol emissions

7 The newly formulated $W(U_{10}, T)$ parameterization is applied to estimate the global annual 8 SSA emission using SSSF of M86 (Eq. (4)). Dividing Eq. (4) by Eq. (3), we modify the M86 9 SSSF to clearly separate the magnitude and shape factors (re-written here as Eq. (4')):

$$10 \qquad \frac{dF}{dr_{80}} = W(U_{10}) \left[\frac{1}{\tau} \frac{dE}{dr_{80}} \right] = W(U_{10}) \cdot \left[3.5755 \times 10^5 \cdot r_{80}^{-3} (1 + 0.057 r_{80}^{1.05}) \times 10^{1.19 e^{-B^2}} \right]$$
(4')

with *B* as defined in Sect. 1. While Eq. (4') shows that the timescale τ is distinct from the shape factor dE/dr, for the calculations the value of τ is included in the numerical coefficient in the brackets. The size range for M86 validity is $r_{80} = 0.8-8$ µm. We calculate the SSA flux for radii r_{80} ranging from 1 to 10 µm. Refer to Anguelova (2016) for using the $W(U_{10})$ parameterization of SAL13 to estimate CO₂ transfer velocity and SSA flux for r_{80} ranging from 0.4 to 250 µm.

17 **2.4.1 Use of discrete whitecap method**

18 The main assumptions of M86 for the SSSF based on the discrete whitecap method-constant 19 values for τ and dE/dr (Sect. 1)—are usually questioned (Lewis and Schwartz, 2004; de 20 Leeuw et al., 2011; Savelyev et al., 2014). It is not expected for either of these assumptions to 21 hold for wave breaking at various scales and under different conditions in different locations. 22 The SSSF proposed by Smith et al. (1993) on the basis of measured size-dependent aerosol 23 concentrations is one of the first formulations to demonstrate that the shape factor cannot be 24 constant. Norris et al. (2013a) also demonstrated that the aerosol flux per unit area whitecap 25 varies with the wind and wave conditions.

Recently, Callaghan (2013) showed that the whitecap timescale is another source of often overlooked variability in SSSF parameterizations based on M86. Because *W* typically includes foam from all stages of whitecap evolution, Callaghan (2013) suggested that the adequate timescale for the aerosol productivity from a discrete whitecap is not just its decay time (as in Eqs. (4) and (4')), but the sum of the whitecap formation and decay timescales τ'.
The value of τ' varies from breaking wave to breaking wave, but an area-weighted mean
whitecap lifetime can be calculated for any given observational period to account for this
natural variability. Analyzing the lifetimes of 552 oceanic whitecaps from a field experiment,
Callaghan (2013) found that the area-weighted mean τ' varies by a factor of 2.7 (from 2.2 to
5.9 s). We refer the reader to Callaghan (2013) for an SSSF that accounts for SSA flux
variability by explicitly incorporating whitecap timescale τ'.

8 Despite these questionable assumptions, the SSSF based on the discrete whitecap 9 method in the form of M86 has been widely used in many models (Textor et al., 2006). 10 Therefore, to those who have worked with M86 until now, a meaningful way to demonstrate 11 how the new satellite-based W data, and W parameterizations based on them, would affect 12 estimates of SSA flux is to hold everything else constant (e.g., the whitecap timescale and 13 productivity in the shape factor) and clearly show differences caused solely by the use of new 14 W expression(s) as a magnitude factor. On these grounds, the choice of the SSSF based on the 15 M86 whitecap method is a suitable baseline for comparisons.

16 **2.4.2 Choice of size distribution**

17 Though the chosen size range of $1-10 \mu m$ for SSA particles is limited, it is well justified for 18 the purposes of this study with the following arguments.

19 Generally, the division of the SSA particles into sizes of small, medium, and large modes (de Leeuw et al., 2011, their §8) is well warranted when one considers the climatic 20 21 effect to be studied (Sect. 1). For example, sub-micron particles are important for scattering 22 by SSA (direct effect) and the formation of cloud condensation nuclei (indirect effect), while 23 super-micron particles are important for heat exchange (via sensible and latent heat fluxes) and heterogeneous chemical reactions (which need surface and volume to proceed 24 25 effectively). However, in this study we do not focus on how the choice of the size distribution 26 will affect the SSA estimates. Nor do we aim to present estimates of specific effect on the 27 climate system. Rather, with a fixed size distribution, we explore how parameterizing W data, 28 which carry information for the influences of many factors, would affect estimates of SSA 29 emission (Sect. 1). In this sense, we can choose to use any published size distribution as a shape factor. 30

1 The chosen size range is the range of medium (super-micron) mode of SSA particles. 2 This is the range for which the size distribution of M86 is valid (Sect. 2.4). The M86 size 3 distribution, in its original or modified form, is widely used in GCMs and CTMs (Textor et 4 al., 2006, their Table 3). The size range of 1–10 μ m is a recurrent part of the various size 5 ranges used in all (or at least most) SSSFs (see Table 2 in Grythe et al. (2014, hereafter G14)).

6 The chemical composition of the SSA particles is another argument favoring the 7 chosen size range. The super-micron particles consist, to a good approximation, solely of sea 8 salt, whereas, in biologically active regions, the sub-micron size range additionally includes 9 organic material, with an increasing contribution as particle size decreases (O'Dowd et al., 10 2004, Facchini et al., 2008; Partanen et al., 2014). Since the organic mass fraction in sub-11 micron SSA particles is still highly uncertain (Albert et al., 2012), we focus on the medium 12 mode SSA emissions.

We evaluate the discrepancy expected due to neglecting particles below 1 μ m using the G14 report of SSA production rate for dry particle diameters D_p = r_{80} obtained with M86 over two different size ranges: 4.51×10^{12} kg yr⁻¹ for the size range of 0.8 μ m < r_{80} < 8 μ m and 5.20×10^{12} kg yr⁻¹ for size range of 0.1 μ m < r_{80} < 10 μ m. The different size ranges bring a difference between the two G14 estimates of about 14%. Neglecting particles with r_{80} < 0.1 μ m would not change significantly the results presented here because they contribute on the order of 1% to the overall mass (Facchini et al., 2008).

20 Because total whitecap fraction, rather than only the active breaking crests, provides 21 bubble-mediated production of SSA, we use W_{37} data to estimate the emission of medium 22 mode SSA. The calculations use a modeling tool (Albert et al., 2010) in which the $W(U_{10})$ parameterization of MOM80, as incorporated in Eq. (4), was replaced with the newly derived 23 $W(U_{10}, T)$ parameterization (Eq. (4')). The resulting size-segregated droplet number emission 24 rate was converted to mass emission rate using the approximation $r_{80} = 2r_d \equiv D_p$, where r_d and 25 $D_{\rm p}$ are the particle dry radius and diameter, respectively (e.g., Lewis and Schwartz, 2004; de 26 Leeuw et al., 2011), and a density of dry sea salt of 2.165 kg m^{-3} . 27

28 **3 Results and Discussion**

The graphs visualizing our results use all W data available for wind speeds from 3 to 35 m s⁻¹. This range of U_{10} is beyond the range $3 \le U_{10} \le 20$ m s⁻¹ used for all fits (Sect. 2.3). In

31 addition, the QuikSCAT instrument, which provided the U_{10} satellite data used in this study,

has a decreased sensitivity for wind speeds over 20 m s⁻¹ (Quilfen et al., 2007). All results regarding higher wind speeds should, therefore, be used with caution.

3 **3.1 Global data sets**

4 Figure 5 shows global W data estimated from WindSat measurements for March 2006 as function of $U_{100SCAT}$ with linear and logarithmic y-axes at 10 GHz (Fig. 5a and c) and 37 GHz 5 6 (Fig. 5b and d). For comparison, the MOM80 relationship (Eq. (3)) is also plotted in each 7 panel (red curves). There are three noteworthy observations in Fig. 5. First, we note the 8 different variability of W_{10} and W_{37} data. The 10 GHz data show far less variability than those 9 at 37 GHz. The W_{37} data at a certain wind speed vary over a much wider range, with the strongest variability for wind speeds of 10-20 m s⁻¹. This observation confirms similar 10 observation reported and analyzed at length by SAL13 in terms of other variables, in addition 11 12 to U_{10} , which influence the whitecap fraction, such as SST, wave field, etc. While SAL13 analyzed this variability, we investigate how well this variability can be parameterized in 13 14 terms of available secondary variables, SST in our case.

15 Another observation in Fig. 5 is noted at low wind speeds. The 10 GHz scatter plots do not show W data for wind speeds lower than about 2 m s^{-1} because at these low wind speeds 16 no active breaking occurs (Sect. 1). In contrast, non-zero W_{37} data are estimated at wind 17 speeds $U_{10} < 2 \text{ m s}^{-1}$. Salisbury et al. (2013) suggested that the presence of foam on the ocean 18 19 surface at these low wind speeds could be due to residual long-lived foam. This residual foam 20 might be stabilized by surfactants, which increases its lifetime (Garrett, 1967; Callaghan et 21 al., 2013). Another explanation could be production of bubbles and foam from biological 22 activity (Medwin, 1977). However, there is not enough information currently to prove any of 23 these conjectures.

The comparison of the MOM80 relationship (Eq. (3)) to W_{10} and W_{37} data clearly reveals the most important feature in Fig. 5—the wind speed dependence of satellite-based *W* data deviates from cubic and cubic-like relationship.

27 **3.1.1 Wind speed dependence**

Following the arguments of our approach (Sect. 2.1) and evaluating the wind speed exponents determined as free parameters (Sect. 2.3.1), we found that a quadratic wind speed exponent (*n* 1 = 2) fits reasonably well both W_{10} and W_{37} data sets. For the same data shown in Fig. 5, Fig.6 2 shows the linear regression of the square root of *W* versus U_{10} :

3
$$W^{1/2} = 0.01U_{10} - 0.011$$
 10 GHz (9a)

4 $W^{1/2} = 0.01U_{10} + 0.019$ 37 GHz (9b)

5 with coefficients of determination R^2 of 0.996 and 0.951, respectively. From Eq. (9), we 6 obtain the following global average wind speed dependence of *W* using U_{10} from QuikSCAT:

7
$$W_{10} = 1 \times 10^{-4} (U_{10} - 1.1)^2$$
 (10)

8
$$W_{37} = 1 \times 10^{-4} (U_{10} + 1.9)^2$$
 (11)

9 where *W* is a fraction (not %).

10 Figure 6c compares $W(U_{10})$ in Eqs. (10-11) to $W(U_{10})$ of SAL13. The trends are close 11 implying that having a different wind speed exponent is largely balanced by corresponding changes to the parametric coefficients. Indeed, the PD between our quadratic $W(U_{10})$ and 12 SAL13 $W(U_{10})$ at 37 GHz ranges from 0.5% to 10% over the wind speed range of 3–20 m s⁻¹. 13 14 ANOVA and Student tests show that these differences are not statistically significant. That is, 15 the global quadratic $W(U_{10})$ parameterization approaches the predictions of the SAL13 parameterization, which has a more specific wind speed exponent (n = 1.59). Note that we do 16 17 not expect our $W(U_{10})$ parameterization to be distinctly different from that of SAL13 because 18 both studies use the same data for W and U_{10} (though from different versions of the whitecap 19 database). Rather, we aim to identify general $W(U_{10})$ trend in order to perform consistent 20 regional analysis.

21 The y-intercept for W_{10} (Eq. (10)) is negative and, following the usual interpretation, yields a threshold wind speed of 1.1 m s⁻¹ for whitecap inception. This is in the range of 22 previously published values from 0.6 (Reising et al., 2002) to 6.33 (Stramska and Petelski, 23 24 2003). Meanwhile, the positive y-intercept b for W_{37} (Eq. (11)) is meaningless at first glance 25 and intriguing upon some pondering. While stabilized residual foam and/or foam from 26 biological sources are possible (Sect. 3.1), it is not known whether such mechanisms are 27 capable of providing a measurable amount of foam patches which produce bubble-mediate 28 sea spray efficiently.

We propose broader interpretation of b in Eqs. (10-11), be it negative or positive. Generally, it is expected that the atmospheric stability (Kara et al., 2008) and fetch (through the wave growth and development) cause inception of the whitecaps at lower or higher wind speed. One can consider the range of values for b mentioned above (0.6 to 6.33) as an expression of such influences. We suppose that b can also incorporate the effect of the seawater properties on the extent of W. The net result of all secondary factors may be either negative or positive b.

8 Specifically, we promote the hypothesis that a positive y-intercept b can be interpreted 9 as a measure of the capacity of seawater with specific characteristics, such as viscosity and 10 surface tension—which are governed by SST, salinity, and surfactant concentration—to affect 11 W. Undoubtedly, none of these secondary factors creates whitecaps per se. Rather, they prolong or shorten the lifetime of the whitecaps via processes governed by the seawater 12 13 properties. For instance, surfactants and salinity influence the persistence of submerged and 14 surface bubbles. This yields variations of bubble rise velocity that replenish the foam on the 15 surface at different rates. Long-lived decaying foam added to foamy areas created by subsequent breaking events would augment W; conversely, conditions that shorten bubble 16 17 lifetimes would reduce *W* (or at least not add to *W*).

A positive *y*-intercept can be thought of as a mathematical expression of this static forcing (as opposed to dynamic forcing from the wind) that given seawater properties can sustain. That is, at any given location, this static forcing acts as though higher wind speed of magnitude $(U_{10} + b)$ is producing more whitecaps than U_{10} alone. By parameterizing coefficients *a* and *b* in terms of different variables, one can evaluate how much the static forcing affects *W* in different geographic regions. By developing parameterizations a(T) and b(T) (Sect. 2.1), here we quantify only one static influence.

25 **3.1.2 Intrinsic correlation**

To quantify the possible intrinsic correlation in the derived $W(U_{10})$ parameterization (Eqs. (10-11)), we derived $W(U_{10})$ using ECMWF wind speeds instead of the QuikSCAT wind speeds (Sect. 2.3.2). Figure 7a shows a scatter plot of $W^{1/2}$ versus $U_{10ECMWF}$ (only data for 37 GHz are shown); dashed and solid lines show unconstrained and zero-forced fits, respectively. The linear regression (given in the figure legend) is used to obtain the global average wind speed dependence using U_{10} from ECMWF as follows:

1
$$W_{37} = 8.1 \times 10^{-5} (U_{10} + 3.33)^2$$

The positive intercept here is interpreted as in Sect. 3.1.1. Using Eq. (12), parameterized Wvalues are plotted as a function of $U_{10\text{ECMWF}}$ in Fig. 7b. Increased scatter of the W data is evident when comparing Figs. 7b and 5d. We use different metrics to detect and evaluate possible intrinsic correlation.

The change of the coefficient of determination R^2 of the $W(U_{10})$ relationship when QuikSCAT winds are substituted with the ECMWF winds is one sign for the presence of intrinsic correlation. Physically, we expect a strong correlation between W and U_{10} , and we see this clearly in Fig. 6b which shows $R^2 = 0.951$ for $W^{1/2}$ and $U_{10QSCAT}$. However, the correlation coefficient might not be as high as in Fig. 6 if U_{10} were from a more independent source. We see this when comparing Figs. 6b and 7a. The $W^{1/2} - U_{10}$ correlation is still strong in Fig. 7a, but the plot shows more scatter and slightly lower correlation with $R^2 = 0.826$.

Figure 8 visualizes the change in the spread of the *W* data with a plot of the residuals (biases) between the *W* data and the derived *W* parameterizations (Eqs. (11) and (12)) as a function of wind speed; Fig. 8a is for $U_{10QSCAT}$ and Fig. 8b is for $U_{10ECMWF}$. Larger biases are evident when $U_{10ECMWF}$ is used. The root-mean-square deviation between *W* data and parameterized *W* values increases from $\Delta W = 0.214\%$ for the data set using $U_{10QSCAT}$ to $\Delta W =$ 0.367% for the data set using $U_{10ECMWF}$.

The slopes in Figs. 6b and 7a differ by about 14%. We evaluate how this translates into differences in W_{37} values as predicted by Eqs. (11) and (12). We found the PD between $W_{37}(U_{10QSCAT})$ and $W_{37}(U_{10ECMWF})$ to be less than \pm 16% for wind speeds of 4–20 m s⁻¹. Specifically, the W_{37} values obtained with $U_{10QSCAT}$ and $U_{10ECMWF}$ are approximately equal for wind speed of 7 m s⁻¹. Below 7 m s⁻¹, $W_{37}(U_{10ECMWF})$ is higher than $W_{37}(U_{10QSCAT})$ by up to 11%. Above 7 m s⁻¹, $W_{37}(U_{10ECMWF})$ is smaller than $W_{37}(U_{10QSCAT})$ by up to 15.7%. The difference goes up to 26% for wind speeds of 3 m s⁻¹.

While different metrics suggest that the intrinsic correlation is present and may contribute to these differences, it is not the only reason for the discrepancies. Different matching procedures (Sect. 2.2.3) and the difference of about 5% between the U_{10} values from the two different sources (Fig. 4a) also contribute to the *W* discrepancies from Eqs. (11) and (12). We, therefore, conclude from the PD values that the effect of the intrinsic correlation alone on *W* is most likely less than about 10% for most frequently encountered wind speeds.

1 **3.1.3 Wave field effect**

Figure 9 shows global $W^{1/2}$ at 37 GHz as function of rising and waning U_{10} from QuikSCAT for March 2006; both rising and waning winds were identified with a wind speed difference $\Delta U_{10} = 1 \text{ m s}^{-1}$ (Sect. 2.3.4). The lines are fits of the data to Eq. (8); solid lines are zero-forced fits. Table 3 shows the slopes *m* and intercepts *c* together with R^2 of the fits for rising and waning winds speeds for ΔU_{10} from 1 to 5 m s⁻¹.

Note the difference between $W^{1/2}$ as a function of rising and waning wind speeds in the 7 range of 10-20 m s⁻¹ (blue colors): more variability is seen at these wind speeds when the 8 wind is waning than in cases when the wind is rising. Larger variability for waning winds 9 lowers their R^2 values compared to R^2 for rising wind speeds for most ΔU_{10} (Table 3). For 10 each, rising and waning wind speeds, R^2 values decrease with ΔU_{10} increasing. The reason for 11 this is that higher ΔU_{10} threshold selects $W^{1/2}$ values associated with more extreme wind 12 conditions (Sect. 2.3.4). Because such conditions are rarer, less $W^{1/2}$ values are selected 13 yielding an increase in the spread of data points. 14

Table 3 shows that slopes *m* for both free and zero-forced fits do not differ substantially for either rising or waning wind speeds for any ΔU_{10} threshold. The intercepts *c* of the free fits increase with the wind threshold for both rising and waning winds. The intercepts are larger for waning winds than for rising. These results yield rising-versuswaning average PD of 10% and 29% for coefficients *a* and *b*, respectively.

20 The rise-wane wind effect, as detected in this study, is not pronounced compared to 21 findings in previous studies that use in situ wind speed data. Goddijn-Murphy et al. (2011) 22 studied wind history and wave development dependencies on in situ W data using wave model 23 (ECMWF), satellite (QuikSCAT), and in situ data for U_{10} . These authors detected significant 24 effects only with in situ U_{10} . The limited wave field effect in our study might be traced back to the method through which U_{10} was determined: wind speeds from satellites are spatial 25 26 averages of scatterometric or radiometric observations that take a snapshot of the surface as it 27 is affected by both wind history and wind local conditions, whereas in situ data for wind 28 speed are single point values averaged over a short time and hence representative for a 29 relatively small area. The effect of the spatial averaging of the satellite data over a much 30 larger area (i.e., the satellite footprint) might be that information on wind history is lost in the 31 process. Limited results on the effect of the wave field obtained with a proxy analysis of the wind history using data paired with a less-than-optimal matching procedure (Sect. 2.2.3) do
not justify further consideration in this study.

3 **3.2** Regional and seasonal data sets

The wind speed exponent in the $W(U_{10})$ relationship derived from the global data set (Eqs. (10-11)) implicitly accounts for the globally-averaged effects of all secondary factors affecting the satellite-based W data. Now we apply Eq. (8) to regional and seasonal sets of satellite-based W data using this wind speed exponent. We analyze the deviations of the parametric coefficients a and b from the globally-averaged trend and parameterize these fluctuations explicitly in terms of SST.

10 **3.2.1** Magnitude of regional and seasonal variations

Table 4 exemplifies the results from Eq. (8b): listed are the slopes m and the intercepts c for 11 $W^{1/2}-U_{10}$ relationships at 10 and 37 GHz in March 2006 in all 12 regions together with 12 coefficients of determination R^2 and 95% CIs from the fitting procedure, as well as mean U_{10} 13 and T values. The results in Table 4 attest that with satellite-based data sets, the sampling 14 15 uncertainty in determining relationships is removed. The remaining geophysical (i.e., regional 16 and seasonal) variations of coefficients a and b, which are obtained from coefficients m and c, are investigated here. Figure 10 shows examples of the $W^{1/2}$ versus $U_{10QSCAT}$ relationships for 17 different regions and seasons. Figures 10a and 10b show scatter plots for the Gulf of Mexico 18 19 (region 1) at both frequencies for January 2006. Statistics are presented at the top of the figures and the fit lines are shown in red. Figures 10c and 10d show the fit lines $W^{1/2}(U_{10})$ for 20 10 and 37 GHz in region 5 for all months, while Figs. 10e and 10f demonstrate variations of 21 the fit lines $W^{1/2}(U_{10})$ for both frequencies over all regions for March 2006. 22

Figure 10 shows that the variations of the $W^{1/2}(U_{10})$ relationships at 10 GHz are 23 24 smaller than those for 37 GHz. Focusing on the results for 37 GHz, we note that geographic differences from region to region for a fixed time period (Fig. 10f) yield more variability in 25 the $W^{1/2}(U_{10})$ relationship than seasonal variations at a fixed location (Fig. 10d). Because the 26 27 37 GHz data provide more information for secondary forcing than the 10 GHz data, the 28 remainder of the data analysis in this study is illustrated with results for W_{37} data. Note that all procedures and analyses described for W_{37} data have also been carried out for the W_{10} data 29 30 and final results are reported (Sect. 3.3).

Figure 10 also shows that variations of $W^{1/2}$ caused by U_{10} from 3 to 20 m s⁻¹ are much 1 larger than the regional and seasonal variations of $W^{1/2}$. While this is expected (because U_{10} is 2 a primary forcing factor), this also points that we need to evaluate whether these regional and 3 4 seasonal variations are statistically significant. For this, we grouped the values of a and b in 5 two ways: (1) by month with the full range of geographical variability (over all 12 regions) for each month; and (2) by region with the full range of seasonal variability (over all 12 6 7 months) for each region. ANOVA test applied to both groups showed that the seasonal 8 variations are not statistically significant, while the regional variations are.

9 We illustrate this in Fig. 11 with values for b; similar graphs for a show the same 10 results. Figure 11a shows the seasonal cycle for the regionally averaged b values with error 11 bars (\pm one SD) representing the regional variability. It is clear that the seasonal variations of 12 the regionally averaged b values lay within the regional variability. That is, variations of b13 from month to month are statistically undistinguishable. Figure 11b illustrates why variations of b from region to region are significantly different. The graph shows the annually averaged 14 15 b values for each region with error bars representing the seasonal variability. It is clear that 16 the geographical variations are not lost in the seasonal variability.

17 **3.2.2 Quantifying SST variations**

The regional differences in Fig. 11b are the variations that we want to quantify with 18 19 coefficients a and b in terms of secondary factors. The deviations of the regional regression coefficients a and b from the regression coefficients $A = 1 \times 10^{-4}$ and B = 1.9 of the general 20 $W(U_{10})$ dependence (Eq. (11)) give a sense for the magnitude of these variations. The PD 21 22 between the annually averaged $\langle a \rangle$ and A is about 5% (average for all regions); the average 23 PD between $\langle b \rangle$ and B is 50%. These regional differences can be caused by any or all other secondary factors. It is not trivial to separate (deconvolve) the effects of different factors 24 25 influencing W data. Because our proxy analysis of the wave field effect produced limited 26 results (Sect. 3.1.3), quantification of the regional differences in terms of wave field with the 27 data we use is not practical. Meanwhile Fig. 3b shows that SST is a distinct characteristic for 28 different regions. This suggests that quantifying the variations of coefficients a and b in terms 29 of SST is a viable possibility. We thus proceed with deriving expressions a(T) and b(T) for 30 the regional variations of the W data; such results are useful to evaluate how well SST can 31 account for the regional variations.

1 We derived a(T) and b(T) for W data at 37 GHz by relating annually averaged a and b 2 values to the annually averaged T for each region (Fig. 12). Figure 12c shows the monthly means of coefficients b for each region and thus demonstrates how the data points in Fig. 12b 3 have been formed; a similar procedure is used for the data points in Fig. 12a. As in Fig. 11b, 4 5 the error bars (± one SD) represent the seasonal variability of SST (horizontal bars) and coefficients a and b (vertical bars). A second order polynomial is fitted to the data points in 6 7 Fig. 12a; a linear fit is applied to the data in Fig. 12b. The correlation coefficients for the derived SST dependences are $R^2 = 0.57$ for a(T) and $R^2 = 0.87$ for b(T). Such R^2 values are 8 9 consistent with the expectation that SST, being a static secondary factor, affects W more via 10 the offset *b* than via the slope *a*.

To evaluate the performance of the quadratic versus cubic wind speed dependence in Eq. (6), we also derived SST dependent coefficients a(T) and b(T) for n = 3 following the same procedure as for the case of n = 2. We applied Eq. (5b) with n = 3 to W_{37} data for all months in regions 4, 5, 6, and 12; we verified that differences due to the use of four instead of twelve regions are not significant. Coefficients *a* and *b* were calculated from the *m* and *c* values and graphs similar to those in Fig. 12 produced. Linear fits for both *a* and *b* were applied to these graphs.

18 **3.3** New parameterization of whitecap fraction

19 New parameterizations for the whitecap fraction $W(U_{10}, T)$ were obtained from 2006 satellite-20 based W data by replacing the fixed coefficients in Eqs. (10-11) with SST-dependent 21 coefficients:

22
$$W = a(T)[U_{10} + b(T)]^2$$
 (13)

23 where

24
$$a(T) = a_0 + a_1 T + a_2 T^2$$
 (14a)

25
$$b(T) = b_0 + b_1 T$$
 (14b)

and the coefficients for data at 10 and 37 GHz are given in Table 5 together with their 95% CIs from the fitting procedure. To evaluate the derived $W(U_{10}, T)$ parameterizations, the whitecap fraction is calculated with Eqs. (13-14) and compared to both parameterized Wvalues and to satellite-based W data.

3.3.1 Comparisons to *W* parameterizations

The $W(U_{10}, T)$ parameterization for 37 GHz is used here. The W values from SAL13 (37 GHz) and MOM80 are used as references for PD calculations and significance tests (Sect. 2.3.3). All parameterizations are run for wind speeds from 3 to 20 m s⁻¹.

Figure 13a compares *W* values from the derived $W(U_{10}, T)$ parameterization at three fixed SST values (T = 2, 12, and 28 °C). Large changes of SST (from 2 to 28 °C) bring relatively small variations between the wind speed trends of *W* at different *T* values. The PDs between the three curves are no more than 15%; indeed, significance tests show that the *W* values at any *T* remain statistically the same. In addition, *W* values at any *T* are not significantly different from the *W* predictions of the global quadratic $W(U_{10})$ parameterization.

11 These results qualitatively illustrate the relative contributions of the implicit and 12 explicit accounts for SST effect in the derived parameterization. Namely, large part of the SST and other influences on W is taken care of implicitly by using quadratic wind speed 13 exponent. Much smaller variations are explicitly expressed with the temperature dependent 14 15 coefficients. Taken together, the set of parametric coefficients—n = 2, a(T), and b(T)— 16 accounts for the: (i) full SST effect (i.e., influence on both the trend and the spread of the W 17 data); and (ii) globally-averaged effects of all other secondary factors (i.e., influences only on 18 the trend of *W* data).

19 We verify the validity of this deduction by comparing in Fig. 13b W values obtained 20 with the quadratic and cubic $W(U_{10}, T)$ parameterizations at T = 20 °C; MOM80 and SAL13 at 37 GHz are shown for reference. The W values from the cubic $W(U_{10}, T)$ parameterization 21 22 are not statistically different from those obtained with either the quadratic $W(U_{10}, T)$ or MOM80 for low winds (< 10 m s⁻¹). Different trends of the W values at higher wind speeds 23 24 suggest that accounting explicitly for SST via a(T) and b(T) in the physically expected cubic wind speed dependence is not sufficient to replicate the satellite-based W data. In other words, 25 26 when the wind speed exponent n is not adjusted to the data but instead follows the physically 27 determined cubic dependence, explicit representation of the SST effect alone via the 28 parametric coefficients a(T) and b(T) cannot account for all observed variations of W. The 29 implication is that when using cubic wind speed exponent, all secondary factors should be 30 introduced explicitly.

1 The PD between the trends of the derived $W(U_{10}, T)$ and MOM80 $W(U_{10})$ is from 5% 2 up to 175% with the largest PDs for wind speeds below 7 m s⁻¹. Figure 13 illustrates this with 3 the different trends of the two parameterizations.

4 **3.3.2** Comparisons to *W* data

5 Here we evaluate how well the derived whitecap fraction parameterizations model the trend 6 and spread of the satellite-based *W* data. The parameterized *W* values are calculated using U_{10} 7 and *T* from the whitecap database (Sect. 2.2.1).

8 Figure 14a compares W values predicted with both new parameterizations, $W(U_{10})$ and 9 $W(U_{10}, T)$, to the same in situ data plotted in Fig. 1b and to independent satellite-based W data 10 for 10 and 37 GHz from 17 March 2007. Comparisons to the in situ W data demonstrate 11 order-of-magnitude consistency of the W values from the new parameterizations. The new 12 global $W(U_{10})$ parameterizations (black symbols in the Fig. 14a) follows reasonably well wind 13 speed trends of the satellite-based W data. The W values predicted with the new $W(U_{10}, T)$ 14 parameterization (red and cyan symbols in Fig. 14a) are spread as the satellite-based W data. The cluster of W values predicted with $W(U_{10}, T)$ are statistically different from the MOM80 15 16 $W(U_{10})$ parameterizations. This is the most important result of this study: we demonstrate that 17 accounting for at least one secondary factor, we are able to model both the trend and the 18 spread of the W values.

19 Note in Fig. 14a that the new $W(U_{10}, T)$ parameterization does not predict the spread 20 of the satellite-based *W* data entirely. This suggests that accounting explicitly for SST in a *W* 21 parameterization is not enough to replicate all natural variability (spread) of *W*. This is 22 consistent with our general understanding of the need to explicitly include many secondary 23 factors in *W* parameterizations, not just SST (Sect. 2.1).

24 Though SST entails small variations in the trend of W with U_{10} (Figs. 13a and 14a), 25 important consequence of the newly derived $W(U_{10}, T)$ parameterization is that it shapes 26 significantly different spatial distribution compared to cubic and higher wind speed 27 dependences like that of the MOM80. Figure 14b shows a difference map between the global 28 annual average W distributions for 2006. MOM80 relationship yields a wider W range with 29 higher values in regions with the highest wind speeds. In particular, this occurs between about 40° and 70° in the Southern ocean and in the North Atlantic. The latitudinal variations from 30 31 the Equator to the poles are more pronounced when using the MOM80 relationship as 1 compared to Eqs. (13-14). The new $W(U_{10}, T)$ parameterization provides a global spatial 2 distribution with similar patterns, but the absolute values are lower at high latitudes and 3 higher at low latitudes. Note that in most studies, as in this study, $W(U_{10})$ of MOM80 is 4 extrapolated beyond the range of the data from which it was derived (Sect. 1). This could be a 5 reason for the large differences between the two parameterizations at higher wind speeds (and 6 especially in cold waters).

7 3.4 Sea spray aerosol production

The newly derived $W(U_{10}, T)$ parameterization (Eqs. (13-14)) was used to estimate the global 8 9 annual average emission of super-micron SSA using M86 SSSF (Eq. (4')). The total (i.e., size integrated) annual SSA mass emission for 2006 is 4359.69 Tg yr⁻¹ (4.4×10^{12} kg yr⁻¹). This is 10 about 50% larger than that calculated with the M86 SSSF using MOM80 (Eq. (4)), 2915 Tg 11 yr⁻¹ (2.9×10¹² kg yr⁻¹). Because we have shown that the new $W(U_{10}, T)$ and MOM80 $W(U_{10})$ 12 are significantly different (Sect. 3.3.2), we infer that the SSA emissions based on SSSFs using 13 each parameterization in combination with the same shape factor (Eq. (4')) also differ 14 15 significantly. The two estimates of SSA emissions are calculated using the same modelling tool (Sect. 2.4) and the same input data (Sect. 2.2.1). Keeping everything the same but the 16 17 magnitude factor guarantees that the 50% difference is due solely to the account for the SST effect on W. The spatial distribution of the mass emission rates obtained with SSSFs using the 18 19 new $W(U_{10}, T)$ is shown in Fig. 15a. The SSA emissions obtained with the new and the 20 MOM80 $W(U_{10})$ parameterizations mimic the patterns of the W distributions. The differences 21 are mapped in Fig. 15b.

Previously modeled total dry SSA mass emissions vary by two orders of magnitude 22 because of a variety of uncertainty sources (Sect. 1): $(2.2-22)\times 10^{12}$ kg yr⁻¹ (Textor et al., 23 2006, their Fig. 1a; de Leeuw et al., 2011, their Table 1); and $(2-74)\times10^{12}$ kg yr⁻¹ for long-24 25 term averages (over 25 years) (G14, their Table 2, excluding 3 outliers). The impact of the 26 modeling method used has to be acknowledged too. Grythe et al. (2014) suggest that the 27 spread in published estimates of global emission based on the same M86 SSSF (Eq. (4)), from 3.3×10^{12} to 11.7×10^{12} kg yr⁻¹ (Lewis and Schwartz, 2004), can be attributed to differences in 28 29 model input data and resolution differences. An example of the same SSSF yielding different 30 results when applied in different models is also seen in the work of de Leeuw et al. (2011, 31 their Table 1).

1 For a meaningful comparison of our results to SSA emissions obtained with other 2 SSSFs, we attempt to remove (or at least minimize) the impact of the modeling method. As in 3 this study, G14 used the same model (i.e., input data and configuration) to evaluate 21 SSSFs, 4 including that of M86, against measurements. We thus can infer a "modelling" factor using 5 our and G14 results obtained with M86 SSSF. We find that the G14 estimate of SSA emission from M86 (4.51×10^{12} kg yr⁻¹) is 1.55 times larger than our estimate of 2.9×10^{12} kg yr⁻¹ from 6 7 M86 and MOM80. We apply this factor of 1.55 to our SSA emission estimated with the new $W(U_{10}, T)$ parameterization and obtain a "model scaled" value of 6.75×10^{12} kg yr⁻¹. Our 8 "model scaled" estimate of the SSA emission is close to the median 5.91×10^{12} kg yr⁻¹ of the 9 10 SSA emissions reported by G14. This shows that an SSSF with a magnitude factor derived 11 from satellite-based W data provides reasonable and realistic predictions of the SSA emission.

12 To narrow down this broad assessment, we now look at the SSSFs evaluated by G14 13 which account for the SST effect on SSA emissions. There are four such SSSFs in the G14 study (see their Table 2): S11T of Sofiev et al. (2011), G03T of Gong (2003), J11T of Jaeglé 14 15 et al. (2011), and G13T of G14. To minimize differences caused by using different size ranges, we focus on S11T and G13T, both applied to dry SSA diameters $D_p = r_{80}$ (Sect. 2.4) 16 17 from 0.01 to 10 µm. The upper limit is the same as in our study, while the lower limit is 18 extended to sub-micron sizes, which, as we have seen (Sect. 2.4.2), introduces a discrepancy 19 of about 14%.

20 The original Sofiev et al. (2011) SSSF is based on the M86 SSSF (Eq. (4)) combined 21 with data from laboratory experiments by Mårtensson et al. (2003) to account for SST and 22 salinity effects and a field experiment by Clarke et al. (2006) to extend the size range. In the G14 study, the salinity weight proposed by Sofiev et al. (2011) is not applied. At a reference 23 salinity of 33 ‰, S11T estimates an SSA emission of 2.59×10¹² kg yr⁻¹. Without the SST 24 effect (the SST factor set to unity), the SSA emission estimated with S11 is 5.87×10^{12} kg yr⁻¹. 25 With everything else the same except for the SST factor in source functions S11 and S11T, 26 27 we evaluate that accounting for the SST effect results in changes by 56%. Correcting for 14% 28 discrepancy due to extended lower size limit, we infer a 42% change when the SST effect is 29 included in the SSSF. This is comparable to the 50% change due to SST in our case. We 30 surmise that parameterizing additional influences on W is a viable way to account and explain 31 some of the uncertainty of SSA emissions.

1 Grythe et al. (2014) used a large data set of ship observations to develop G13T by 2 changing both the magnitude and the shape factors. The authors modified the SSSF of Smith 3 and Harrison (1998) (a sum of two log-normal distributions) to add an extra log-normal mode 4 to cover the accumulation mode. They also added the empirically based SST factor (a third 5 order polynomial) proposed by Jaeglé et al. (2011). With G13T, G14 estimate an SSA emission of 8.91×10^{12} kg yr⁻¹. The functional forms of the magnitude (involving the SST 6 7 effect) and shape (modelling the size distribution) factors of G13T and S11T are very 8 different. This makes it difficult to evaluate the relative contribution of the magnitude and 9 shape factors for variations in SSA emissions. Our results can help.

10 The shape factors of S11T and our SSSF using $W(U_{10}, T)$ have a similar (not identical) 11 functional form (that of M86, original and modified), but the functional forms accounting for SST are different. Our SSA emission estimate is about 62% higher than that of S11T. 12 Allowing for 14% discrepancy due to the lower size limit, we find that different approaches to 13 account for SST lead to about 67% variation in SSA emissions. Compared to G13T, our SSSF 14 15 using $W(U_{10}, T)$ has a different shape factor (that of M86 versus log-normal), and a similar (but not identical) functional form for the SST effect (polynomial). Our SSA emission 16 17 estimate is about 32% lower than that of G13T. Allowing for 14% size discrepancy, we find 18 that different shape factors lead to about 13% variation in SSA emissions.

On the basis of these assessments, we can state that the inclusion of the SST effect in the magnitude factor and/or the choice of the shape factor (size range and model for the size distribution) in the SSSF can explain 13%-67% of the variations in the predictions of SSA emissions. The spread in SSA emission can thus be constrained by more than 100% when improvements of both the magnitude and the shape factor are pursued. Our results on the *W* parameterization (Fig. 14a) suggest that accounting for more secondary forcing in the magnitude factor would explain more fully the spread among SSA emissions.

26 4 Conclusions

The objective of the study presented here is to evaluate how accounting for natural variability of whitecaps in the parameterization of the whitecap fraction W would affect mass flux predictions when using a sea spray source function based on the discrete whitecap method. The study uses satellite-based W data estimated from measurements of the ocean surface brightness temperature T_B by satellite-borne microwave radiometers at frequencies of 10 and 37 GHz, W_{10} and W_{37} . Global and regional data sets comprising W_{10} and W_{37} data, wind speed 1 U_{10} , and sea surface temperature *T* for 2006 were used to derive parameterizations $W(U_{10})$ and 2 $W(U_{10}, T)$. The SSSF of Monahan et al. (1986) combined with the new $W(U_{10}, T)$ was used to 3 estimate sea spray aerosol emission. The conclusions of the study are the following.

4 The global W data set can be parameterized reasonably well with a quadratic 5 correlation between W and U_{10} (Eqs. (10-11) and Sect. 3.1.1). The unconventional positive y-6 intercept for $W_{37}(U_{10})$ could be interpreted as a mathematical expression of the static forcing 7 that given seawater properties (e.g., effects of SST, salinity, and surfactant concentrations) 8 impart on whitecaps. Parameterization $W(U_{10})$ derived with an independent data set (U_{10} from 9 ECMWF instead of QuikSCAT) helps to determine that the intrinsic correlation between W and U_{10} is most likely less than about 10% (Sect. 3.1.2). Proxy analysis of satellite-based W 10 11 data at increasing and decreasing wind speeds (Table 3) yields limited results for the effect of 12 the wave field on W (Sect. 3.1.3). The derived $W(U_{10})$ for both W_{10} and W_{37} replicate the trend 13 of the satellite-based data well (Fig. 14a). That is, the adjusted quadratic wind speed exponent 14 in $W(U_{10})$ accounts implicitly for most of the SST variations. The new quadratic $W(U_{10})$ 15 predicts whitecap fraction significantly different from that obtained with the widely used 16 *W*(*U*₁₀) of MOM80.

17 Applying the global $W(U_{10})$ parameterization on regional scale shows that the seasonal 18 variations of its regression coefficients a and b are not statistically significant, while the 19 regional variations are. On this basis, by relating annually averaged a and b values to the 20 annually averaged T for each region (Fig. 12), the explicit SST dependences a(T) and b(T) for 21 data at 10 and 37 GHz were derived (Sect. 3.3. and Table 5). The new $W(U_{10}, T)$ 22 parameterization (Eqs. (13-14)) is able to model the variability (spread) of the satellite-based 23 W data (Fig. 14a). The capability of the new $W(U_{10}, T)$ parameterization to model both the 24 trend and the spread of the W data sets it apart from all other $W(U_{10})$ parameterizations (e.g., 25 MOM80 and SAL13). Results show that besides SST, one needs to include explicitly other 26 secondary factors in order to model the full spread of the satellite-based W. Including the SST 27 effect via a(T) and b(T) in the physically expected cubic wind speed dependence is not sufficient to replicate the trend of the satellite-based W values. While SAL13 analysis of the 28 29 satellite-based whitecap database demonstrated the influences of secondary factors on whitecap fraction, our study goes a step further in using the satellite-based W data to 30 31 parameterize one of these influences, that of SST.

1 Application of the new $W(U_{10}, T)$ parameterization in the Monahan et al. (1986) SSSF 2 resulted in a total (integrated only over super-micron sizes) SSA mass emission estimate of 4359.69 Tg yr⁻¹ (4.4×10^{12} kg yr⁻¹) for 2006. Scaled for modeling differences (Sect. 3.4), this 3 estimate is 6.75×10^{12} kg yr⁻¹, which is comparable to previously reported estimates. 4 5 Comparing our and previous total SSA emissions, we have been able to assess to what degree 6 accounting for the SST influence on whitecaps can explain the spread of SSA emissions. With 7 or without the SST effect included in the SSSF, SSA emissions obtained with the new $W(U_{10},$ T) parameterization vary by \sim 50%. Different approaches to account for SST effect yield 8 9 ~67% variations. Different models for the size distribution applied to different size ranges 10 lead to 13%-42% variations in SSA emissions. Understanding and constraining the various 11 sources of uncertainty in the SSSF would eventually improve the accuracy of SSSF 12 predictions. Including the natural variability of whitecaps in the SSSF magnitude factor is a 13 viable way toward such accuracy improvement.

14 While the new $W(U_{10}, T)$ parameterization is able to model the trend and the spread of 15 the satellite-based W data, the SST variations are relatively small. To model the full 16 variability of W, future work should focus on the parameterization of the wave field effect. 17 The extended version of the whitecap database contains wave field characteristics and is thus 18 suitable for such quantification. It is recommended that the extended whitecap database 19 includes wind speed data from independent source(s) matched in time and space at WindSat 20 resolution.

21 Data availability

The data analysis and the results reported in this study are available from the corresponding author M.F.M.A. (Monique) Albert (<u>monique.albert@tno.nl</u>).

24 Acknowledgements

25 This study is partly funded by SRON, Netherlands Institute for Space Research, through the

- 26 Dutch Users Support Programme GO-2. MDA was sponsored by the Office of Naval
- 27 Research, NRL Program element 61153N, WU 4500. GdL by was supported by the European
- 28 Space Agency (Support to Science Element: Oceanflux Sea Spray Aerosol, contract No.
- 29 4000104514/11/I-AM), the Centre on Excellence in Atmospheric Science funded by the
- 30 Finnish Academy of Sciences Excellence (project no. 272041), the CRAICC project (part of
- 31 the Top-level Research Initiative).

1 References

- 2 Albert, M. F. M. A., Schaap, M., de Leeuw, G., and Builtjes, P. J. H.: Progress in the
- 3 determination of the sea spray source function using satellite data, Journal of Integrative
- 4 Environmental Sciences, 7, 159-166, 2010.
- 5 Albert, M. F. M. A., Schaap, M., Manders, A. M. M., Scannell, C., O'Dowd, C. D., and de
- 6 Leeuw, G.: Uncertainties in the determination of global sub-micron marine organic matter
- 7 emissions, Atmos. Environ., 57, 289-300, 2012.
- Andreae, M. O. and Crutzen, P. J.: Atmospheric aerosols: biogeochemical sources and role in
 atmospheric chemistry, Science, 276, 1052-1058, 1997.
- Andreas, E. L.: Sea Spray and the turbulent air-sea heat fluxes, J. Geophys. Res., 97, 1142911441, 1992.
- 12 Anguelova, M. D: Assessing the utility of satellite-based whitecap fraction to estimate sea
- 13 spray production and CO2 transfer velocity, IOP Conf. Ser.: Earth Environ. Sci., 35, 012002,
- 14 2016.
- Anguelova, M. D. and Gaiser, P. W.: Skin depth at microwave frequencies of sea foam layers
 with vertical profile of void fraction, J. Geophys. Res., 116, C11002, 2011.
- Anguelova, M. D. and Gaiser, P. W.: Microwave emissivity of sea foam layers with vertically
 inhomogeneous dielectric properties, Remote Sens. Environ., 139, 81-96, 2013.
- Anguelova, M. D. and Webster, F.: Whitecap coverage from satellite measurements: A first
 step toward modeling the variability of oceanic whitecaps. J. Geophys. Res., 111, C03017,
 2006.
- Anguelova, M. D., Bettenhausen, M. H., and Gaiser, P. W.: Passive remote sensing of sea
 foam using physically-based models, in: Proceedings of the IGARSS 2006: IEEE
 International Geoscience and Remote Sensing Symposium, Denver, Colorado, USA, 31 July4 August, 3659-3662, 2006.
- Anguelova, M. D., Bettenhausen, M. H., Johnston, W. F., Gaiser, P. W.: First extensive
 whitecap database and its use to study whitecap fraction variability, in: Proceedings of the
 17th Air-Sea Interaction Conference, AMS, Annapolis, Maryland, USA, 26 30 September,
- 29 2010 (http://ams.confex.com/ams/pdfpapers/174036.pdf).

- 1 Anguelova, M. D., Bobak, J. P., Asher, W. E., Dowgiallo, D. J., Moat, B. I., Pascal, R. W.,
- 2 and Yelland, M. J.: Validation of satellite-based estimates of whitecap coverage: approaches
- 3 and initial results, in: Proceedings of the 16th Air-Sea Interaction conference, AMS, Phoenix,
- 4 Arizona, USA, 10-15 January, 2009, (http://ams.confex.com/ams/pdfpapers/143665.pdf).
- Asher, W. E. and Wanninkhof, R.: The effect of bubble-mediated gas transfer on purposeful
 dual-gaseous tracer experiments, J. Geophys. Res., 103, 10,555-10,560, 1998.
- 7 Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone
- 8 destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere,
- 9 Nature, 334, 138–141, 1988.
- 10 Bettenhausen, M. H., Smith, C. K., Bevilacqua, R. M., Wang, N. -Y., Gaiser, P. W., and
- 11 Cox, S.: A nonlinear optimization algorithm for WindSat wind vector retrievals, IEEE T.
- 12 Geosci. Remote, 44, 597-610, 2006.
- Blanchard, D. C.: The electrification of the atmosphere by particles from bubbles in the sea,
 Prog. Oceanogr., 1, 73-112, 1963.
- 15 Blanchard, D. C.: The production, distribution, and bacterial enrichment of the sea-salt
- aerosol, in: Air-sea exchange of gases and particles, Liss, P. S. and Slinn, W. G. N., D. Reidel
 Publishing Company, Dordrecht, The Netherlands, 407-454, 1983.
- Bondur, V., and Sharkov, E.: Statistical properties of whitecaps on a rough sea, Oceanology,
 22, 274–279, 1982.
- Callaghan, A. H.: An improved whitecap timescale for sea spray aerosol production flux
 modeling using the discrete whitecap method, J. Geophys. Res.-Atmos., 118, 9997-10010,
 2013.
- Callaghan, A. H. and White, M.: Automated processing of sea surface images for the
 determination of whitecap coverage, J. Atmos. Ocean. Tech., 26, 383-394, 2009.
- Callaghan, A. H., de Leeuw, G., Cohen, L., and O'Dowd, C. D.: Relationship of oceanic
 whitecap coverage to wind speed and wind history, Geophys. Res. Lett., 35, L23609, 2008.
- 27 Callaghan, A. H., Deane, G. B., and Stokes, M. D.: Two Regimes of Laboratory Whitecap
- Foam Decay: Bubble-Plume Controlled and Surfactant Stabilized, J. Phys. Oceanogr., 43,
- 29 1114-1126, 2013.

- 1 Chameides, W. L. and Stelson, A. W.: Aqueous-phase chemical processes in deliquescent
- 2 sea-salt aerosols: a mechanism that couples the atmospheric cycles of S and sea salt, J.
- 3 Geophys. Res.- Atmos., 97, 20565-20580, 1992.
- 4 Chelton, D. B. and Freilich, M. H.: Scatterometer-based assessment of 10-m wind analyses
- 5 from the operational ECMWF and NCEP numerical weather prediction models, Mon.
- 6 Weather Rev., 133, 409-429, 2005.
- 7 Cicerone, R. J.: Halogens in the atmosphere, Rev. Geophys. Space Ge., 19 (NO. 1), 123-139,
 8 1981.
- 9 Clarke, A. D., Owens, S. R., and Zhou, J.: An ultrafine sea-salt flux from breaking waves: 10 Implications for cloud condensation nuclei in the remote marine atmosphere, J. Geophys.
- 11 Res., 111, D06202, 2006.
- 12 de Leeuw, G., Andreas, E. L., Anguelova, M. D., Fairall, C. W., Lewis, E. R., O'Dowd, C. D.,
- Schulz, M., and Schwartz, S. E.: Production flux of sea-spray aerosol, Rev. Geophys., 49,
 RG2001, 2011.
- 15 Facchini, M. C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S.,
- 16 Ceburnis, D., Flanagan, R., Nilsson, E. D., de Leeuw, G., Martino, M., Woeltjen, J., and
- 17 O'Dowd, C. D.: Primary submicron marine aerosol dominated by insoluble organic colloids
- 18 and aggregates, Geophys. Res. Lett., 35, L17814, 2008.
- Fairall, C. W., Kepert, J. D., and Holland, G. J.: The effect of sea spray on surface energy
 transports over the ocean, The Global Atmosphere and Ocean System, 2, 121-142, 1994.
- 21 Falkowski, P. G., Barber, R. T., and Smetacek, V.: Biogeochemical controls and feedbacks on
- 22 ocean primary production, Science, 281, 200-206, 1998.
- Ghan, S. J., Guzman, G., and Hayder, A. -R.: Competition between sea salt and Sulfate
 particles as cloud condensation nuclei, J. Atmos. Sci., 55, 3340-3347, 1998.
- 25 Gaiser, P.W., St. Germain, K. M., Twarog, E. M., Poe, G. A., Purdy, W., Richardson, D.,
- 26 Grossman, W., Linwood Jones, W., Spencer D., Golba, G., Cleveland, J., Choy, L.,
- 27 Bevilacqua, R. M., and Chang, P. S.: The WindSat spaceborne polarimetric microwave
- radiometer: sensor description and early orbit performance, IEEE T. Geosci. Remote, 42, NO.
- 29 11, 2347–2361, 2004.

- 1 Garrett, W. D.: Stabilization of air bubbles at the air-sea interface by surface-active material,
- 2 Deep-Sea Res., 14, 661-672, 1967.
- 3 Goddijn-Murphy, L., Woolf, D. K., and Callaghan, A. H.: Parameterizations and algorithms
- 4 for oceanic whitecap coverage, J. Phys. Oceanogr., 41, 742-756, 2011.
- Gong, S. L.: A parameterization of sea-salt aerosol source function for sub- and super-micron
 particles, Global Biogeochem. Cycles, 17, 1097, 2003.
- Graedel, T. E. and Keene, W. C.: The budget and cycle of Earth's natural chlorine, Pure Appl.
 Chem., 68, 1689-1697, 1996.
- 9 Grythe, H., Ström, J., Krejci, R., Quinn, P., and Stohl, A.: A review of sea-spray aerosol
- 10 source functions using a large global set of sea salt aerosol concentration measurements,
- 11 Atmos. Chem. Phys., 14, 1277–1297, 2014.
- 12 Hanson, J. L., and Phillips, O. M.: Wind sea growth and dissipation in the open ocean, J.
- 13 Phys. Oceanogr., 29, 1633-1648, 1999.
- 14 Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J.-T.: Global distribution of sea
- 15 salt aerosols: new constraints from in situ and remote sensing observations, Atmos. Chem.
- 16 Phys., 11, 3137-3157, doi:10.5194/acp-11-3137-2011, 2011.
- Kara, A. B., Wallcraft, A. J., and Bourassa, M. A.: Air-sea stability effects on the 10 m winds
 over the global ocean: Evaluations of air-sea flux algorithms, J. Geophys. Res.-Oceans, 113,
 C04009, 2008.
- 20 Keene, W. C., Pszenny, A. A. P., Jacob, D. J., Duce, R. A., Galloway, J. N., Schultz-Tokos, J.
- J., Sievering, H., and Boatman, J. F.: The geochemical cycling of reactive chlorine through
 the marine troposphere, Global Biogeochem. Cy., 4 (NO. 4), 407-430, 1990.
- 23 Keene, W. C., Khalil, M. A. K., Erickson, D. J., McCulloch, A., Graedel, T. E., Lobert, J. M.,
- 24 Aucott, M. L., Gong, S.-L., Harper, D. B., Kleiman, G., Midgley, P., Moore, R. M., Seuzaret,
- 25 C., Sturges, W. T., Benkovitz, C. M., Koropalov, V., Barrie, L. A., and Li, Y.-F.: Composite
- 26 global emissions of reactive chlorine from anthropogenic and natural sources: reactive
- chlorine emissions inventory, J. Geophys. Res., 104 (NO. D7), 8429-8440, 1999.
- Kleiss, J. M. and Melville, W. K.: The analysis of sea surface imagery for whitecap
 kinematics, J. Atmos. Ocean. Tech, 28, 219-243, 2011.

- 1 Koop, T., Kapilashrami, A., Molina, L.T., and Molina, M. J.: Phase transitions of sea-
- 2 salt/water mixtures at low temperatures: implications for ozone chemistry in the polar marine
- 3 boundary layer, J. Geophys. Res., 105 (NO. D21), 26393-26402, 2000.
- Lewis, E. R. and Schwartz, S. E.: Sea salt aerosol production: mechanisms, methods,
 measurements and models A critical review, Geoph. Monog. Series, 152, American
- 6 Geophysical Union, Washington D. C., 413 pp, 2004.
- 7 Luria, M. and Sievering, H.: Heterogeneous and homogeneous oxidation of SO_2 in the remote
- 8 marine atmosphere, Atmos. Environ., 25A, 1489-1496, 1991.
- 9 Mårtensson, E. M., Nilsson, E. D., de Leeuw, G., Cohen, L. H., and Hansson, H.-C.:
- 10 Laboratory simulations and parameterization of the primary marine aerosol production, J.
- 11 Geophys. Res., 108, 4297, 2003.
- Medwin, H.: In situ acoustic measurements of microbubbles at sea, J. Geophys. Res., 82, 971-976, 1977.
- 14 Meissner, T. and Wentz, F. J.: The emissivity of the ocean surface between 6 and 90 GHz
- over a large range of wind speeds and earth incidence angles, IEEE T. Geosci. Remote, 50,
 3004-3026, 2012.
- Melville, W. K.: The role of surface-wave breaking in air-sea interaction, Annu. Rev. Fluid
 Mech., 28, 279-321, 1996.
- 19 Monahan, E. C.: Oceanic Whitecaps, J. Phys. Oceanogr., 1, 139-144, 1971.
- 20 Monahan, E. C. and O'Muircheartaigh, I.: Optimal power-law description of oceanic
- 21 whitecap coverage dependence on wind speed, J. Phys. Oceanogr., 10, 2094–2099, 1980.
- 22 Monahan, E. C. and O'Muircheartaigh, I.: Whitecaps and the passive remote sensing of the
- 23 ocean surface, Int. J. Remote Sens., 7, 627-642, 1986.
- Monahan, E. C. and Woolf, D. K.: Comments on "Variations of whitecap coverage with wind stress and water temperature, J. Phys. Oceanogr., 19, 706–709, 1989.
- 26 Monahan, E. C., Fairall, C. W., Davidson, K. L., and Boyle, P. J.: Observed inter-relations
- between 10 m winds, ocean whitecaps and marine aerosols, Q. J. Roy. Meteor. Soc., 109,
 379-392, 1983.

- 1 Monahan, E. C., Spiel, D. E., and Davidson, K. L.: A model of marine aerosol generation via
- 2 whitecaps and wave disruption, in: Oceanic whitecaps: and their role in air-sea exchange
- 3 processes, Monahan, E. C., Mac Niocaill, G., D. Reidel Publishing Company, Dordrecht, The
- 4 Netherlands, 167-174, 1986.
- 5 Norris, S. J., Brooks, I. M., Moat, B. I., Yelland, M. J., de Leeuw, G., Pascal, R. W., Brooks,
- 6 B. J.: Near-surface measurements of sea spray aerosol production over whitecaps in the open
- 7 ocean. Ocean Science, 9, 133–145, doi: 10.5194/os-9-133-2013, 2013a.
- 8 Norris, S. J., Brooks, I. M., and Salisbury, D. J.: A wave roughness Reynolds number 9 parameterization of the sea spray source flux, Geophys. Res. Lett., 40, 4415–4419, 2013b.
- O'Dowd, C. D. and de Leeuw, G.: Marine aerosol production: a review of the current
 knowledge, Philos. T. R. Soc. A, 365, 1753-1774, 2007.
- 12 O'Dowd, C. D., Lowe, J. A., Smith, M. H., and Kaye, A. D.: The relative importance of non-
- 13 sea-salt sulphate and sea-salt aerosol to the marine cloud condensation nuclei population: An
- 14 improved multi-component aerosol-cloud droplet parametrization, Q. J. Roy. Meteor. Soc.,
- 15 125, 1295-1313, 1999.
- 16 O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S.,
- Yoon, Y. J., and Putaud, J.-P.: Biogenically driven organic contribution to marine aerosol,
 Nature, 431, 676-680, 2004.
- 19 Ovadnevaite, J., Manders, A., de Leeuw, G., Ceburnis, D., Monahan, C., Partanen, A .- I.,
- 20 Korhonen, H., and O'Dowd, C. D.: A sea spray aerosol flux parameterization encapsulating
- 21 wave state, Atmos. Chem. Phys., 14, 1837-1852, 2014.
- Paget, A. C., Bourassa, M. A., and Anguelova, M. D.: Comparing in situ and satellite-based
 parameterizations of oceanic whitecaps, J. Geophys. Res. Oceans, 120, 2826–2843, 2015.
- Pandey, P. C. and Kakar, R. K.: An empirical microwave emissivity model for a foamcovered sea, IEEE J. Oceanic Eng., 7, 135-140, 1982.
- Partanen, A.-I., Dunne, E. M., Bergman, T., Laakso, A., Kokkola, H., Ovadnevaite, J.,
 Sogacheva, L., Baisnée, D., Sciare, J., Manders, A., O'Dowd, C., de Leeuw, G., and
- 28 Korhonen, H.: Global modelling of direct and indirect effects of sea spray aerosol using a
- source function encapsulating wave state, Atmos. Chem. Phys., 14, 11731-11752, 2014.

- 1 Quilfen, Y., Prigent, C., Chapron, B., Mouche, A. A., and Houti, N.: The potential of
- 2 QuikSCAT and WindSat observations for the estimation of sea surface wind vector under
- 3 severe weather conditions, J. Geophys. Res., 112, C09023, 2007.
- Reising, S., Asher, W., Rose, L., and Aziz, M.: Passive polarimetric remote sensing of the
 ocean surface: The effects of surface roughness and whitecaps, paper presented at the
 International Union of Radio Science, URSI Gen. Assem., Maastrict, Netherlands, 2002.
- 7 Saiz-Lopez, A. and von Glasow, R.: Reactive halogen chemistry in the troposphere, Chem.
- 8 Soc. Rev., 41, 6448-6472, 2012.
- 9 Salisbury, D. J., Anguelova, M. D., and Brooks, I. M.: On the variability of whitecap fraction
- 10 using satellite-based observations, J. Geophys. Res.-Oceans, 118, 6201-6222, 2013.
- 11 Salisbury, D. J., Anguelova, M. D., and Brooks, I. M.: Global distribution and seasonal
- 12 dependence of satellite-based whitecap fraction, Geophys. Res. Lett., 41, 1616–1623, 2014.
- 13 Savelyev, I. B., Anguelova, M. D., Frick, G. M., Dowgiallo, D. J., Hwang, P. A., Caffrey, P.
- 14 F., and Bobak, J. P.: On direct passive microwave remote sensing of sea spray aerosol
- 15 production, Atmos. Chem. Phys., 14, 11611-11631, 2014.
- 16 Sievering, H., Boatman, J., Gorman, E., Kim, Y., Anderson, L., Ennis, G., Luria, M., and
- 17 Pandis, S.: Removal of sulphur from the marine boundary layer by ozone oxidation in sea-salt
- 18 aerosols, Nature, 360, 571-573, 1992.
- 19 Sievering, H., Gorman, E., Ley, T., Pszenny, A., Springer-Young, M., Boatman, J., Kim, Y.,
- 20 Nagamoto, C., and Wellman, D.: Ozone oxidation of sulfur in sea-salt aerosol particles during
- the Azores Marine Aerosol and Gas Exchange experiment, J. Geophys. Res. -Atmos., 100,
 23075-23081, 1995.
- 23 Smith, M. H., Park, P. M., and Consterdine, I. E.: Marine aerosol concentrations and 24 estimated fluxes over the sea, Q. J. Roy. Meteor. Soc., 119, 809–824, 1993.
- Smith, M. H. and Harrison, N. M.: The sea spray generation function, J. Aerosol Sci., 29
 (Suppl. 1), S189-S190, 1998.
- 27 Sofiev, M., Soares, J., Prank, M., de Leeuw, G., and Kukkonen, J.: A regional-to-global
- 28 model of emission and transport of sea salt particles in the atmosphere, J. Geophys. Res., 116,
- 29 D21302, 2011.

- 1 Stramska, M. and Petelski, T.: Observations of oceanic whitecaps in the north polar waters of
- 2 the Atlantic, J. Geophys. Res., 108, NO. C3, 3086, 2003.
- 3 Tang, W., Yueh, S. H., Fore, A. G., and Hayashi A.: Validation of Aquarius sea surface

4 salinity with in situ measurements from Argo floats and moored buoys, J. Geophys. Res.
5 Oceans, 119, 6171–6189, 2014, doi:10.1002/2014JC010101.

- 6 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen,
- 7 T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D.,
- 8 Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I.,
- 9 Iversen, T., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A.,
- 10 Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland,
- 11 Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of
- 12 aerosol life cycles within AeroCom, Atmos. Chem. Phys., 6, 1777-1813, 2006.
- 13 Toba, Y. and Chaen, M.: Quantitative expression of the breaking of wind waves on the sea
- surface, Records of Oceanographic Works in Japan, 12 (NO. 1), 1-11, 1973.
- 15 Thorpe, S. A.: On the clouds of bubbles formed by breaking wind-waves in deep water, and
- 16 their role in air-sea gas transfer, Philos. T. R. Soc. S. -A., 304, 155-210, 1982.
- 17 Wanninkhof, R., Asher, W. E., Ho, D. T., Sweeney, C., and McGillis, W. R.: Advances in
- 18 quantifying air-sea gas exchange and environmental forcing, Annual Review of Marine
- 19 Science, 1, 213-244, 2009.
- Wentz, F. J.: A model function for ocean microwave brightness temperatures, J. Geophys.
 Res., 88, NO. C3, 1892-1908, 1983.
- Wentz, F. J.: A well-calibrated ocean algorithm for special sensor microwave / imager, J.
 Geophys. Res., 102, NO. C4, 8703-8718, 1997.
- Woolf, D. K.: Bubbles and their role in gas exchange, in: The Sea Surface and Global
 Change, Liss, P. S. and Duce, R. A., Cambridge Univ. Press, New York, 173-205, 1997.
- Wu, J.: Variations of whitecap coverage with wind stress and water temperature, J. Phys.
 Oceanogr., 18, 1448-1453, 1988.
- 28 Zhao, D. and Toba, Y.: Dependence of whitecap coverage on wind and wind-wave properties,
- 29 J. Oceanogr., 57, 603-616, 2001.
- 30

1 **Table 1**

Coordinates (longitude and latitude), number of data points are given together with range, mean and median values for wind speed and SST of
all selected regions (a) for January 2006, (b) for July 2006. Confidence intervals (CI) at 95% level are given for regions 4, 5, 6, and 12, whose
seasonal variations are plotted in Fig. 3.

Region Lon		Lon Lat Samples			Wind speed [m s ⁻¹]				SST [°C]		
Region	Lon.	Lat.	Samples	Range	Mean	95% CI	Median	Range	Mean	95% CI	Median
1	86°W–95°W	23°N-28°N	18896	1.3-15.7	7.5		7.6	19.4-26.0	23.8		24.1
2	$1^{\circ}W$ – $15^{\circ}W$	1°S-30°S	169128	0.2-12.9	6.4		6.4	21.4-27.8	24.2		24.1
3	75°E–89°E	1°S-30°S	169056	0.0-13.4	7.0		7.2	23.0-29.4	26.8		27.3
4	11°W-20°W	30°N-44°N	49760	0.2-19.6	8.0	2.7×10 ⁻²	7.6	13.3-20.4	16.4	1.5×10 ⁻²	16.3
5	86°W-100°W	31°S–60°S	200360	0.5-23.0	8.7	1.3×10 ⁻²	8.7	4.8-24.1	12.7	2.2×10 ⁻²	11.7
6	171°W-180°W	15°S–14°N	123328	0.6-15.6	8.2	1.2×10 ⁻²	8.2	26.2-30.4	28.4	0.6×10 ⁻²	28.2
7	31°W-50°W	10°N–29°N	90640	0.3-20.0	8.8		9.0	20.1-27.9	24.9		25.3
8	140°W–160°W	20°S-30°S	50040	0.5-16.3	6.8		6.7	22.2-29.1	26.3		26.6
9	140°W–160°W	40°S-50°S	41840	0.1-20.6	6.9		6.5	9.3-18.2	13.2		13.1
10	0°W-30°W	40°S-50°S	133080	0.5-26.4	9.4		9.3	3.2-16.7	9.6		9.3
11	50°E-70°E	40°S-50°S	50784	0.5-21.6	9.6		9.6	3.2-17.4	9.6		9.5
12	180°E-180°W	60°S-90°S	576576	0.2-20.9	7.0	0.8×10 ⁻²	6.7	-1.9-8.0	1.8	0.5×10 ⁻²	1.4

5 a) For January 2006

2 b) For July 2006

Region Lon		Lat	Samples	Wind speed [m s ⁻¹]				SST [°C]			
11081011	2011.	Lut.	Sumples	Range	Mean	95% CI	Median	Range	Mean	95% CI	Median
1	86°W–95°W	23°N-28°N	13848	0.4-10.0	4.5		4.4	28.7-30.5	29.5		29.4
2	1°W-15°W	1°S-30°S	189600	0.2-14.0	6.6		6.6	17.7–27.1	23.2		23.7
3	75°E–89°E	1°S-30°S	195424	0.6-15.4	8.0		8.1	18.8-30.0	25.4		25.9
4	11°W–20°W	30°N-44°N	43040	0.7-14.0	6.7	2.2×10 ⁻²	6.6	16.9–23.3	20.4	1.3×10 ⁻²	20.5
5	86°W-100°W	31°S–60°S	257496	0.7-22.7	9.8	1.4×10 ⁻²	9.6	2.5-19.1	9.3	1.6×10 ⁻²	8.3
6	171°W-180°W	15°S–14°N	133096	0.1-14.8	6.0	1.1×10 ⁻²	6.0	26.9–29.7	28.8	0.3×10 ⁻²	29.0
7	31°W-50°W	10°N–29°N	88304	0.4-13.6	7.4		7.4	23.6-28.0	26.0		26.1
8	140°W-160°W	20°S-30°S	47504	0.7-24.7	6.9		6.2	18.8-27.0	23.2		23.4
9	140°W-160°W	40°S-50°S	52736	0.5-21.0	10.1		10.3	8.2-14.1	10.9		10.8
10	0°W-30°W	40°S-50°S	160192	0.9–28.9	10.8		10.8	1.8-14.6	8.3		8.3
11	50°E-70°E	40°S-50°S	49344	1.1-28.2	12.9		12.7	2.1–16.1	8.3		7.8
12	180°E-180°W	60°S–90°S	177240	0.8-29.1	11.7	1.9×10 ⁻²	11.9	-1.3–4.3	1.7	0.4×10 ⁻²	1.7

Table 2

Regression coefficients *n*, *a*, and *b* with 95% confidence intervals (CI) derived as free
parameters from fitting of Eq. (6) to different global data sets.

Data set	<i>n</i> ± 95% CI	a ± 95% CI	<i>b</i> ±95% CI		
<i>W</i> ₁₀	2.22 $\pm 3.23 \times 10^{-7}$	$5.23 \times 10^{-5} \pm 5.73 \times 10^{-11}$	$-0.226 \pm 1.54 \times 10^{-6}$		
W ₃₇	$1.46 \pm 6.15 \times 10^{-7}$	$6.17 \times 10^{-4} \pm 1.21 \times 10^{-9}$	$-0.957 \pm 3.58 \times 10^{-6}$		
$W_{10} \& W_{37}$	$1.79 \pm 8.1 \times 10^{-7}$	$2.03 \times 10^{-4} \pm 5.43 \times 10^{-10}$	$-0.409 \pm 4.36 \times 10^{-6}$		

1 **Table 3**

2 Regression coefficients m (slope) and c (intercept) derived by fitting Eq. (8b) to subsets of W

- 3 data associated with rising and waning (increasing and decreasing) wind speeds (a proxy
- 4 analysis for wave field development). Different wind speed differences ΔU_{10} , determined
- 5 from ECMWF wind speed values, were used to select *W* data for rising or waning winds.
- 6 Given is also coefficient m (slope) for a fit forced though zero (intercept c = 0). Coefficients
- 7 of determination R^2 are also given.

8

Wind speed	Slope <i>m</i>		Intercept c		R^2		Slope, <i>m</i>		R^2	
difference								zero intercept		
$\Delta U_{10} [{ m m \ s}^{-1}]$	Rise	Wane	Rise	Wane	Rise	Wane	Rise	Wane	Rise	Wane
1	0.01	0.01	0.021	0.023	0.942	0.947	0.012	0.012	0.887	0.898
2	0.01	0.009	0.024	0.029	0.924	0.915	0.012	0.012	0.854	0.841
3	0.009	0.009	0.027	0.035	0.904	0.863	0.012	0.011	0.819	0.753
4	0.009	0.008	0.028	0.040	0.886	0.794	0.012	0.011	0.789	0.659
5	0.009	0.008	0.028	0.040	0.890	0.755	0.012	0.010	0.798	0.641

2 **Table 4**

Results for slope (coefficient *m*) and intercept (coefficient *c*) with their 95% confidential intervals (CI) from Eq. (8b) applied to satellite-based W data for March 2006 for all 12 regions for a) 10 GHz data; b) 37 GHz data. Mean wind speed U_{10} and sea surface temperature (SST) *T* for each region are also given. Such data were obtained for all months.

Region	Slope $m \times 10^2$	95% CI×10 ⁸	Intercept $c \times 10^2$	95% CI×10 ⁷	R^2	mean U_{10}	mean SST	Samples
1	0.983	6.47	-0.766	5.05	0.995	7.4	23.7	21304
2	0.997	0.84	-0.935	0.56	0.992	6.5	26.5	208560
3	1.006	0.59	-0.967	0.43	0.996	6.8	27.1	211152
4	1.027	1.98	-1.077	1.77	0.996	8.2	15.3	64480
5	1.031	0.55	-1.157	0.57	0.995	9.8	13.3	268320
6	1.004	0.91	-0.946	0.58	0.996	6.1	28.1	140064
7	1.005	1.30	-0.937	0.96	0.995	7.0	23.9	105848
8	1.006	2.82	-0.934	1.95	0.993	6.4	27.5	58112
9	1.014	2.97	-1.055	2.58	0.994	8.04	13.9	52952
10	1.021	0.85	-1.091	0.80	0.995	8.8	10.6	161776
11	1.033	3.12	-1.148	3.07	0.993	9.2	11.5	55200
12	1.028	0.15	-1.145	0.14	0.994	9.3	1.8	1039264

6 a) 10 GHz

2 b) 37 GHz

Slope $m \times 10^2$	95% CI×10 ⁸	Intercept $c \times 10^2$	95% CI×10 ⁷	R^2	mean U_{10}	mean SST	Samples
0.999	22.90	2.270	18.22	0.9574	7.3949	23.7273	18056
1.088	2.67	1.391	1.772	0.9453	6.4370	26.4630	191728
1.032	2.46	1.545	1.812	0.9518	6.6755	27.1823	185224
0.986	7.10	2.623	6.45	0.9604	8.2645	15.3113	55216
1.002	1.68	2.413	1.751	0.9589	9.7181	13.3633	242792
0.985	3.95	1.648	2.49	0.9381	5.9357	28.0589	125632
1.074	3.24	1.886	2.42	0.9784	6.8255	23.8623	96440
0.975	6.59	1.797	4.59	0.9657	6.2512	27.5191	54712
1.008	9.78	2.117	8.67	0.9447	8.0332	13.9375	48888
0.988	2.88	2.474	2.64	0.9521	8.4807	10.6534	150920
0.981	11.21	2.613	10.87	0.9165	9.0372	11.6882	51784
0.963	0.55	2.784	0.53	0.9338	9.0238	1.8538	922080
	Slope <i>m</i> ×10 ² 0.999 1.088 1.032 0.986 1.002 0.985 1.074 0.975 1.008 0.988 0.981 0.963	Slope $m \times 10^2$ 95% CI×10^80.99922.901.0882.671.0322.460.9867.101.0021.680.9853.951.0743.240.9756.591.0089.780.9882.880.98111.210.9630.55	Slope $m \times 10^2$ 95% CI×10^8Intercept $c \times 10^2$ 0.99922.902.2701.0882.671.3911.0322.461.5450.9867.102.6231.0021.682.4130.9853.951.6481.0743.241.8860.9756.591.7971.0089.782.1170.9882.882.4740.98111.212.6130.9630.552.784	Slope $m \times 10^2$ 95% CI×10^8Intercept $c \times 10^2$ 95% CI×10^70.99922.902.27018.221.0882.671.3911.7721.0322.461.5451.8120.9867.102.6236.451.0021.682.4131.7510.9853.951.6482.491.0743.241.8862.420.9756.591.7974.591.0089.782.1178.670.98111.212.61310.870.9630.552.7840.53	Slope $m \times 10^2$ 95% CI×10^8Intercept $c \times 10^2$ 95% CI×10^7 R^2 0.99922.902.27018.220.95741.0882.671.3911.7720.94531.0322.461.5451.8120.95180.9867.102.6236.450.96041.0021.682.4131.7510.95890.9853.951.6482.490.93811.0743.241.8862.420.97840.9756.591.7974.590.96571.0089.782.1178.670.94470.98111.212.61310.870.91650.9630.552.7840.530.9338	Slope $m \times 10^2$ 95% $CI \times 10^8$ Intercept $c \times 10^2$ 95% $CI \times 10^7$ R^2 mean U_{10} 0.99922.902.27018.220.95747.39491.0882.671.3911.7720.94536.43701.0322.461.5451.8120.95186.67550.9867.102.6236.450.96048.26451.0021.682.4131.7510.95899.71810.9853.951.6482.490.93815.93571.0743.241.8862.420.97846.82550.9756.591.7974.590.96576.25121.0089.782.1178.670.94478.03320.98111.212.61310.870.91659.03720.9630.552.7840.530.93389.0238	Slope $m \times 10^2$ 95% CI×10^8Intercept $c \times 10^2$ 95% CI×10^7 R^2 mean U_{10} mean SST0.99922.902.27018.220.95747.394923.72731.0882.671.3911.7720.94536.437026.46301.0322.461.5451.8120.95186.675527.18230.9867.102.6236.450.96048.264515.31131.0021.682.4131.7510.95899.718113.36330.9853.951.6482.490.93815.935728.05891.0743.241.8862.420.97846.825523.86230.9756.591.7974.590.96576.251227.51911.0089.782.1178.670.94478.033213.93750.9882.882.4742.640.95218.480710.65340.9630.552.7840.530.93389.02381.8538

Table 5

Coefficients for the SST dependence of the parametric coefficients *a* and *b* in Eq. (14) with their 95% confidence intervals (CI) from the fitting procedure. The temperature dependent parametric coefficients a(T) and b(T) are used in parameterization $W(U_{10}, T)$ (Eq. (13)) derived from satellite-based *W* data for 10 and 37 GHz for 2006.

Data set	$a_0 \pm 95\%$ CI	<i>a</i> ₁ ± 95% CI	<i>a</i> ₂ ± 95% CI	<i>b</i> ₀ ± 95% CI	<i>b</i> ₁ ± 95% CI
W_{10}	1.08×10 ⁻⁴	-2.45×10 ⁻⁷	-1.45×10 ⁻⁹	-1.203	9.9612×10 ⁻³
	$\pm 1.33 \times 10^{-6}$	$\pm 1.91 \times 10^{-7}$	$\pm 5.78 \times 10^{-9}$	$\pm 1.91 \times 10^{-2}$	$\pm 9.53 \times 10^{-4}$
<i>W</i> ₃₇	8.46×10 ⁻⁵	1.63×10 ⁻⁶	-3.35×10 ⁻⁸	3.354	-6.2×10 ⁻²
	$\pm 3.75 \times 10^{-6}$	$\pm 5.46 \times 10^{-7}$	$\pm 1.65 \times 10^{-8}$	$\pm 9.72 \times 10^{-2}$	$\pm 4.85 \times 10^{-3}$

1 Figure captions

- 2 Figure 1. Satellite estimates of W data at 37 GHz for 11 March 2006. a) Map (0.5°×0.5°) of
- 3 ascending and descending passes for *W* at 37 GHz; b) *W* at 10 and 37 GHz (green and
- 4 magenta symbols, respectively) compared to historical photographic data including total W
- 5 (diamonds) and active whitecap fraction W_A (squares). Parameterization $W(U_{10})$ of Monahan
- 6 and O'Muircheartaigh (1980, MOM80) (purple line) is shown for reference.
- 7 Figure 2. Selected regions to determine regional variations of $W(U_{10})$.
- 8 Figure 3. Seasonal cycle for 2006 in different regions as defined in Fig. 2 and Table 1: a)
- 9 wind speed U_{10} ; b) Sea surface temperature (SST) T. The SST error bars in panel (b) are \pm

10 one standard deviation; the U_{10} error bars are wider and not plotted in panel (a) for clarity.

11 The regions represent: 4-Temperate zone in Northern hemisphere; 5-Temperate zone in

- 12 Southern hemisphere; 6–Doldrums along the Equator; 12–Lowest SST.
- 13 Figure 4. Scatter plot for March 2006 of (a) global $U_{10\text{ECMWF}}$ versus $U_{10\text{QSCAT}}$ and (b) global T
- 14 from ECMWF versus T from GDAS. In both figures the colors indicate the amount of data
- points per hexabin. The black lines are linear fits: the dashed line represents unrestricted fit and the solid line a fit forced through zero. The linear regressions and respective R^2 are listed
- 17 in each panel.
- Figure 5. Global *W* as function of U_{10} from QuikSCAT for March 2006 where *W* is obtained with 10 GHz (a) and 37 GHz (b) measurement frequency. Panels c and d plot the data in panels a and b with logarithmic *y*-axis. The red line indicates the Monahan and O'Muircheartaigh (1980, MOM80) relationship (Eq. (3)). The colors indicate the amount of data points per hexabin.
- Figure 6. Wind speed dependence of whitecap fraction $W(U_{10})$ derived from the global W data 23 set: a) $W^{1/2}$ as function of U_{10} from QuikSCAT for March 2006, where $W^{1/2}$ is obtained with 24 25 10 GHz measurement frequency; b) same as in panel (a) but for 37 GHz. The black line in 26 both panels indicates the best linear fit through the data. The red line in (b) equals the black 27 line in (a). The colors indicate the amount of data points per hexabin. c) Comparison of 28 derived global $W(U_{10})$ at 10 GHz (red line) and 37 GHz (black line) to $W(U_{10})$ parameterizations of Salisbury et al. (2013) in Eq. (1) for 10 GHz (blue line) and 37 GHz 29 30 (magenta). Parameterization $W(U_{10})$ of Monahan and O'Muircheartaigh (1980, MOM80) in 31 Eq. (3) (purple line) is shown for reference.

Figure 7. Scatter plots of *W* data for 37 GHz versus $U_{10\text{ECMWF}}$ for March 2006: a) $W^{1/2}$; b) *W* obtained with Eq. (12). The black lines in panel a are linear fits: the dashed line represents unrestricted fit and the solid line is a fit forced through zero. The linear fits and respective R^2 are listed. The red line in panel b indicates the Monahan and O'Muircheartaigh (1980, MOM80) relationship (Eq. (3)). The colors indicate the amount of data points per hexabin.

6 Figure 8: Scatter plots of residuals ΔW between W data for 37 GHz from the whitecap 7 database and parameterized W values as a function of wind speed from different sources: a) 8 Wind speed values $U_{10QSCAT}$ from the whitecap database used with Eq. (11); b) Wind speed 9 values $U_{10ECMWF}$ from the ECMWF model used with Eq. (12). The rms deviation for each 10 data set is given in each panel.

Figure 9: Global $W^{1/2}$ for data at 37 GHz as a function of rising (a) and waning (b) U_{10} from QuikSCAT for March 2006. The dashed line indicates the best linear fit through the data, whereas the solid line indicates a linear fit, forced through zero.

Figure 10: Linear fits of $W^{1/2}$ versus U_{10} for: region 1 for January 2006 at 10 GHz (a) and 37 GHz (b); region 5 for all months at 10 GHz (c) and 37 GHz (d); regions 1-12 for March 2006 at 10 GHz (e) and 37 GHz (f).

Figure 11: Regional and seasonal variations: a) Regionally averaged b values for each month with error bars (\pm one standard deviation) representing the regional variability; b) Annually averaged b values for each region with error bars representing the seasonal variability.

Figure 12: Sea surface temperature dependences of a) coefficient *a* (slope) and b) coefficient *b* (intercept) in the $W(U_{10})$ dependence. Each point is annual mean for different region. The error bars indicate ± 1 standard deviation for SST (horizontal bars) and coefficients (vertical bars). Panel c) shows the monthly means of coefficients *b* for each region that form one data point in panel b). Regions in Northern hemisphere (NH) are show with squares; regions in Southern hemisphere (SH) are shown with circles. The diamonds are for region 6 at the Equator.

Figure 13: a) Comparison of the new parameterization $W(U_{10}, T)$ (Eqs. 13-14) at three fixed SST values (T = 20 °C, red line; T = 12 °C, green line; T = 2 °C, blue line) to the parameterizations of Salisbury et al. (2013, SAL13) (Eq. (1)) for 37 GHz (magenta line) and Monahan and O'Muircheartaigh (1980, MOM80) (Eq. (3)) (purple line). b) Comparison of the new $W(U_{10}, T)$ parameterizations with quadratic (Eqs. 13-14, blue line) and cubic (green line) wind speed exponents at T = 20 °C to the parameterizations of Salisbury et al. (2013, SAL13) (Eq. (1)) for 37 GHz (magenta line) and Monahan and O'Muircheartaigh (1980, MOM80) (purple line).

5 Figure 14: a) In situ *W* data as in Fig. 1b (gray symbols) and satellite-based *W* data for 17 6 March 2007 at 10 and 37 GHz (green and magenta symbols, respectively) compared to *W* 7 values obtained from $W(U_{10})$ for 10 and 37 GHz (black lines, Eqs. (10-11)) and $W(U_{10}, T)$ for 8 10 (red) and 37 GHz (cyan, Eqs. (13-14)). Wind speed and sea surface temperature from the 9 whitecap database are used for the calculations.

b) Difference map of annual average *W* distribution for 2006 calculated from the Monahan and O'Muircheartaigh (1980, MOM80) $W(U_{10})$ parameterization (Eq. (3)) minus $W(U_{10}, T)$ from Eqs. (13-14). The calculations use wind speed U_{10} from QuikSCAT in the whitecap database.

Figure 15: a) Annual average super-micron mass emission rate for 2006 in μ g m⁻² s⁻¹ calculated from from Eq. (4')). b) Difference map between the annual average super-micron SSA mass emission rate calculated from the Monahan et al. (1986) SSSF and the annual average super-micron SSA mass emission rate calculated from the Monahan et al. (1986) SSSF where *W* is replaced with Eqs. (13-14). The calculations use wind speed U_{10} from QuikSCAT in the whitecap database.

20







3 Figure 1









- 4 Figu







- 14 Figure 5

























6 Figure 9













2 Figure 12



5 Figure 13













