Rebuttal Letter on "Validation of farm-scale methane emissions using nocturnal boundary layer budgets" by J. Stieger et al.

We are thankful to the reviewers for their precise and constructive assessment of our research paper. The reviewes significantly improved our manuscript. We tried to carefully address all mentioned issues by the three reviewers and specified and/or changed our text accordingly in the attached manuscript. Hereafter, we list all mentioned issues from the referees (*in italics*), and provide our responses (normal font, blue) and the new text (in purple, where appropriate) grouped by referee. Where we specify a line number, this line number refers to the numbering in the document that shows the differences between the discussion version and our revised version. We hope that with our proposed changes in the manuscript we were able to address the critique expressed by all three reviewers in such a way that our manuscript can now be accepted for publication in ACP.

Referee 1

(2) I think the numbers of the various livestock emitting CH4 should be given

The numbers of the various livestock emitting CH₄ have been added to section 2.4:

"The CHAI inventory estimates include different animal headcounts for the years 2011 and 2012, respectively. In 2011, 55 diary cattle, seven non-diary cattle, nine young 270 cattle, 18 sheep and 129 swine were located at the chamau farmstead. The inventory estimate for 2012 includes besides 81 diary cattle, two non-diary cattle, 18 young cattle, eight sheep, 47 swine also 39 goats." (lines 275–279)

Referee 2

(1) Advection seems to be a major issue as with many studies. This is discussed fairly detailed in the paper, however I would recommend the following additions:

- When neglecting Fa in section 2.3, it should be mentioned that this will be discussed in more detail in the discussion section. Otherwise the reader wonders how one can characterize the Chamau area as "horizontally sufficiently large and flat terrain with a homogeneous source distribution"
- The fetch of the different profile measurements should be assessed a bit more quantitatively, e.g. by making use of the wind profiles (velocities and directions). Also changing wind direction can cause changes in fetch, and can lead to inclusion or exclusion of emission locations from upwind areas.
- The selection of the wind sector is not very consistent: Section 2.3 mentions the SSE-SW sector, i.e. a wind direction between 202.5 and 225 degrees; the wind direction limits shown in Fig. 3 are at 90 and 270 degrees, and the wind direction observed during the different soundings varies much stronger than the narrow SSE-SW sector. Given the dimension of the farm (buildings cover about a 200 m x 200 m area, estimated using Google Earth), the distance of 150m of the measurement to the nearest farm building seems relatively small. A sketch showing the main farm buildings and the location of the balloon and tower measurements sites would be helpful to the reader to better grasp the geometry.

According to the reviewer's suggestion we additionally informed about the further discussion of neglecting horizontal advection in Section 2.3 :

"The impact of neglected horizontal advective processes on the overall NBL budget estimates will be part of the discussion section." (lines 244–245)

According to the reviewer's concern, that upwind areas can have an impact on the methane budget depending on the wind directions, we re-evaluated the profiles and concluded, that the wind directions were not of major concern for our NBL budget estimates, as the budget calculations were either restricted to specific wind directions and/or to very low wind speeds.

We agreed to the reviewers concern, that the selection of wind direction and its designation within the text put confusion, why we clarified this aspect with the corresponding text in section 2.3 and the removal of the dashed vertical lines in Figure 3(c):

"In addition, only measurements with an average wind direction from the SSE–SW (157.5°–225°) sector were used for the flux calculation, i.e., where the main barn building of the Chamau station and the grazed pastures were located. During very calm weather conditions when mean wind speeds reached zero, measurements were not restricted to wind directions." (lines 240–244)

The vertical dashed lines were removed from Figure 3c. At the same time, we removed the boxes around the legends which were overly prominent in that figure.

(2) It remains unclear whether the limitation of the vertical integration to the level where delta-theta/delta-z approaches zero really avoids influence on the NBL budget from sources further upwind than the Chamau farmstead. This seems especially problematic for the 2012 observations, where profiles show no clear top of the NBL, and where CH4 shows no vertical gradient. This should be explained in more details, as also the various references cited in the manuscript do not really provide this information.

As the top of the NBL was not reached in 2012, we tried to assess the NBL budget flux with selection criteria other than delta-theta/delta-z, i.e. transitions layers in combination with wind direction. In order to clarify this aspect we tried to improve our text in Section 4.2:

"Still, since the upper limit of the NBL height was not reached in 2012 and the wellmixed CH₄ concentration profiles pointed to the presence of advective processes, the emissions from sources further upwind might have had an impact on the resulting NBL budget estimate." (lines 505–507)

3) Usually a budget estimate cannot be given from a single profile as done in Table 1, as a change in the mixing ratio needs to be determined (see Eq. 1). It should be made more clear in the manuscript that this is only possible due to the use of Kriging in time and space and by taking the local derivative.

We added text to be more explicit in the caption to Table 1 (page 24) and in Section 2.5 (lines 298–304). Please note that the \unit{} command in the Copernicus stylesheet prevents latexdiff from correctly interpreting these text elements.

Furthermore, ordinary Kriging algorithms usually also provide an estimate of the uncertainty in the interpolated variable. Have those been used in error propagation to determine the resulting contribution to the uncertainty in estimated NBL budget fluxes?

We are aware of this possibility and thus added further details and expanded the text in our discussion, the real issues are expected in the systematic errors, not in the random errors. We tried to express this in Section 4.1, namely with the new text in lines 451–461:

"Thus, although there are certainly further components that add to random uncertainty in our estimates, the two main issues to overcome in future projects are (i) extending the vertical layer over which high-quality concentration measurements can be made; and (ii) using profiling at slow but steady ascent rates to obtain consistent vertical profiles of all meteorological variables that better allows to detect instationary conditions, and which clearly includes the transition across z_i , which is a critical value in flux estimates. Even with limited payload it may be advisable to have to tethersondes attached to the balloon tether, one which measures the conditions 50– 100 m above the tube inlet (of limited length), and another at the height of the tube inlet. In such a way at least conditions where concentration measurements close to z_i are covered, the upper tethersond would allow to establish that these measurements were actually close to z_i . In most cases a distinct drop in relative humidity is a clear sign of z_i ."

P21771 L10: Are there any tube effects on CH4 when using 220 m tube for tethered balloon measurements?

Tube effects during the tethered balloon measurements were neglected as the used polyethylene inlet tube was of very high chemical and ageing resistance and pre-tests did not show any impact on laboratory concentrations.

But see also our detailed response to Reviewer #3 with a similar question.

P21778 L20, Fig. 2a: It is unclear how the observations from the different soundings listed in Table 1 support the Kriging results shown in Fig. 2a. The text mentions a maximum around 3:00 LT, while the last sounding was made around 00:48-01:47. It would be helpful if the measurement locations (height vs. time) supporting the Kriging results could be shown as thin black lines in Fig. 2a.

We tried to clarify this aspect by adding a statement to the caption of Fig 2(a):

"(a) Kriged time-space interpolation of the CH₄ concentrations obtained from the balloon measurements during 16/17 August 2011, including rejected measurements not used for NBL budget estimates."

P21780 L25: I suggest replacing "gradient" with "difference"

We changed the word 'gradient' by 'difference' (line 388)

Caption Fig. 5: "The circles (A) show the NBL budget flux that was achieved if no interpolation was done beyond the height range" I suggest using "extrapolation" rather than "interpolation".

We changed the word 'interpolation' by 'extrapolation'.

P21782 L25: Also here I suggest using "extrapolation" rather than "interpolation".

We changed the word 'interpolation' by 'extrapolation' (line 436).

P21785 L10-12: This seems like circular reasoning: If the good agreement between NBL budget fluxes with the CHAI estimates is used to validate the experimental approach, the NBL budget fluxes cannot be used to then validate the inventory estimates.

We were very thankful that this miswording was pointed out by the reviewer. We suggested a different wording in Section 4.2:

"Overall, a good agreement of the NBL budget fluxes with the CHAI estimates could be achieved which confirmed the viability of our selection criteria." (lines 511–513)

Referee 3

(1) P21786-Li22 It is very useful to see a comparison of NBL technique with inventory estimates, particularly for the local area "based on actual livestock data" and as the authors mention, few studies have done this. I feel it would add considerable value to this paper and broaden possible comparisons if the authors could give information about the animal numbers for Chamau farmstead for CHAI and for the local area used with SEI estimates and from this add some estimates of emission per head estimated by the NBL method and how this compares to the numbers that have been used in NIR, SEI and CHAI because many animal science and the inventories are based on emission per head (in addition to giving the per area fluxes).

The numbers of the various livestock emitting CH₄ have been added to section 2.4:

"The CHAI inventory estimates include different animal headcounts for the years 2011 and 2012, respectively. In 2011, 55 diary cattles, seven non-diary cattles, nine young 270 cattles, 18 sheeps and 129 swines were located at the chamau farmstead. The inventory estimate for 2012 includes besides 81 diary cattles, two non-diary cattles, 18 young cattles, eight sheeps, 47 swines also 39 goats." (lines 276–279)

(2) I feel that a bit more analysis is warranted given the profiles of 2012 that appear remarkably well mixed with height (in Fig. 3). As mentioned on P21784, the profiles may be influenced by a relatively large area of order hundreds km2 and on P21785. The authors note the "well mixed" nature of the methane and its build through the night. There is useful discussion on P21780 and it appears that with increased wind aloft or possible jet activity may have been sufficient to keep the stable NBL stirred and as mentioned, it appears "that local sources only had a minor influence". However the CHAI would suggest that local sources were larger in 2012 than 2011; alternatives are that these sources have (i) mixed vertically through the lower atmosphere locally which seems unlikely, or (ii) that they have not been captured because too close and do not fall in measurement footprint for winds experienced at the measurement site which is solely "seeing" distant sources contributing to the NBL or (iii) both local and distant sources are well mixed. Can more analysis be done to understand the 2011 / 2012 differences add more interpretation? Figure 4 gives useful additional information for 2011, could a similar figure be looked at for 2012. Alternatively or further it would be instructive to look at the Richardson Number through each of the profiling periods; perhaps a critical number is always exceeded in 2012 to allow turbulent mixing to continue and is perhaps sufficient to mix up local emissions in spite of light wind at the surface (e.g. along lines of Grachev et al 2013) whereas perhaps turbulence is supressed in 2011 case. Whilst this may help explain differences in the profiles, there is also no sign of a capping in the methane profiles and if the NBL flux is only integrated to Zi (50m) then it looks like a substantial component of the flux is missed as it appears that methane above this height is well coupled to the surface layer. There is some discussion of uncertainly introduced by assumptions in Zi etc. This analysis depends to a degree on whether the Chamau farmstead region a relative "hotspot" of methane

emission or is it surrounded over the hundreds of km scale by land of similar or larger mean emission as this affects potential advection flux errors.

After having calculated the gradient Richardson number for each 10-m layer of the vertical profiles, it could be seen, that we did not experience any turbulent conditions, which could have introduced mixing processes. However, as the reviewer correctly stated, the well-mixed CH₄ concentration profiles in 2012 might be influenced by further sources, which could not be excluded although having restricted the NBL budget estimates to specific wind directions and integration heights. We assume that advective effects were causing the well-mixed profiles in different manners (see Section 4.2). We addressed this point with a text change in Section 4.2 :

"Still, since the upper limit of the NBL height was not reached in 2012 and the wellmixed CH_4 concentration profiles pointed to the presence of advective processes, the emissions from sources further upwind might had an impact on the resulting NBL budget estimate." (lines 505–507)

P21768-Line11: the IPCC methodology for NIR could be cited: e.g. IPCC: IPCC Guide- lines for National Greenhouse Gas Inventories. In: Prepared by the National Green- house Gas Inventories Programme, Eggleston, H. S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K. (Eds.), IGES, Japan, 2006.

The IPCC reference was included in the final version (line 53).

P21768-Line16 statement "require detailed knowledge of the transport pathways of tracers and the location of active CH4 sources with which they must be collocated to provide realistic flux estimates" needs clarification. For the individual animal measurements, (Johnston, Deighton) the SF6 tracer source is typically located in the rumen to be collocated with the main source location of enteric methane. It is unclear what if any short-comings the authors are suggesting with this technique.

According to the reviewer's suggestions we changed the mislucked wording in the introduction:

"These tracer techniques have been useful for the quantification of known CH₄ source components and especially of individual emissions per animal, requiring detailed knowledge of the transport pathways of tracers and the location of active CH₄ sources. However, such methods tend to have some limitations as the scale of integration increases to a whole farmstead or even larger spatial scale. For the validation of an emission inventory that is supposed to cover the total of all known and unknown (or neglected source components), mostly volume- integrated budget estimates are used, since such an approach has the potential to also reveal unexpected or unexpectedly large flux components of CH₄."

P21769-Line14: "only few projects, so far, have directly validated livestock CH4 emission estimates via atmospheric concentration measurements without the deployment of chambers". Few projects should be clarified, and only refs to Denmean and Grobler given. There have been a number of studies at a number looking at free grazing animal emissions using a variety of techniques without the deployment of chambers e.g. Judd et al 1999, Lassey et al 2011, 2013, Laubach and Kelliher, 2004, 2005a,b, Laubach et al 2008, Wratt et al 2001 etc. and with comparison with chambers, e.g.: Grainger et al 2007.

In order to avoid a misunderstanding, we changed the wording and introduced further

references in the introductory part of the paper:

"So far, a fair number of studies estimated livestock CH_4 emissions via atmospheric concentration measurement without the deployment of chambers (Judd et al., 1999; Denmead et al., 2000; Laubach et al., 2004; Laubach et al., 2005a; Laubach et al., 2005b; Laubach et al., 2008; Grobler et al., 2014). Unfortunately, only few studies compare their results directly to inventory estimates (Levin et al., 1990, Lowry et al., 2001; Hsu et al., 2010; Hiller et al. 2014b)."

P21771-Line 10 Please give tube diameter as this affects Reynolds number for the tube and how fast and degree of turbulence in gas transported.

We added this information and rearranged the sentences to read:

"A helium-filled blimp (7.2 m³, The Blimp Works Inc., Statesville, NC, USA) with a net lift of 4.35 kg carried a polyethylene inlet tube of 220 m length with inner diameter of 4 mm (Maagtechnic, Dübendorf, Switzerland) connected to a ground measurement station. With this approach it was possible to keep sensitive, heavy, and power consuming instrumentation on the ground at all times. The air sampled was drawn at a flow rate of $\approx 1 \text{ L min}^{-1}$ (Reynolds number 344) to a fast greenhouse gas analyzer (FGGA, Los Gatos Research Inc., Mountain View, CA, USA)."

Please note that a Reynolds number of 344 means laminar flow, which is averaging out short-term fluctuations and increases the quality of the average concentration needed for the NBL approach (in contrast to eddy covariance flux measurements where turbulent flow should be achieved, but this is not the goal with the NBL approach).

Figure 2: what is explanation for the sharp cut-off at 150m in the kriged plot, is this an artifact of the analysis? Further, in a similar way to the 2012 profiles, if Zi is found to be 50m there would appear to be a component of the methane plume that has mixed above this height and do the authors consider possibility of flux underestimation from this by not integrating the whole vertical extent or ignoring the Fent at the top of the NBL?

As the sharp cut-off is an artefact of the applied Kriging method, we suggested to leave Figure 2 as is, but added the time-height trace of the tethered balloon soundings plus white boxes to show availability of near-surface profiles (0–10 m). This information was also added to the figure caption:

"Kriged time-space interpolation of the CH_4 concentrations obtained from the balloon and near-surface (0–10 m) measurements during 16/17 August 2011, including rejected measurements not used for NBL budget estimates. Vertical and horizontal kriging directions were equally weighted, resulting sometimes in sharp vertical concentration changes. The balloon position is shown with a white line, whereas the near-surface concentration measurements are shown with a white box." (Caption to Figure 2).

— end of document —

Manuscript prepared for Atmos. Chem. Phys. with version 2015/04/24 7.83 Copernicus papers of the LATEX class copernicus.cls. Date: 24 November 2015

Validation of farm-scale methane emissions using nocturnal boundary layer budgets

J. Stieger¹, I. Bamberger^{1,2}, N. Buchmann¹, and W. Eugster¹

¹ETH Zurich, Dept. of Environmental Systems Science, Institute of Agricultural Sciences, 8092 Zurich, Switzerland

²Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate Research, 82467 Garmisch-Partenkirchen, Germany

Correspondence to: W. Eugster (eugsterw@ethz.ch)

Abstract. This study provides the first experimental validation of Swiss agricultural methane emission estimates at the farm scale. We measured CH_4 concentrations at a Swiss farmstead during two intensive field campaigns in August 2011 and July 2012 to (1) quantify the source strength of livestock methane emissions using a tethered balloon system, and (2) to validate inventory emis-

- 5 sion estimates via nocturnal boundary layer (NBL) budgets. Field measurements were performed at a distance of 150 m from the nearest farm buildings with a tethered balloon system in combination with gradient measurements at eight heights on a 10 m tower to better resolve the near-surface concentrations. Vertical profiles of air temperature, relative humidity, CH_4 concentration, wind speed and wind direction showed that the NBL was strongly influenced by local transport processes and
- 10 by the valley wind system. Methane concentrations showed a pronounced time course, with highest concentrations in the second half of the night. NBL budget flux estimates were obtained via a time-space kriging approach. Main uncertainties of NBL budget flux estimates were associated with instationary atmospheric conditions and the estimate of the inversion height z_i (top of volume integration). The mean NBL budget fluxes of $1.60 \pm 0.31 \,\mu g \,\mathrm{CH}_4 \,\mathrm{m}^{-2} \,\mathrm{s}^{-1}$ (1.40 ± 0.50 and
- 15 $1.66 \pm 0.20 \,\mu\text{g} \,\text{CH}_4 \,\text{m}^{-2} \,\text{s}^{-1}$ in 2011 and 2012, respectively) were in good agreement with local inventory estimates based on current livestock number and default emission factors, with 1.29 ± 0.47 and $1.74 \pm 0.63 \,\mu\text{g} \,\text{CH}_4 \,\text{m}^{-2} \,\text{s}^{-1}$ for 2011 and 2012, respectively. This indicates that emission factors used for the national inventory reports are adequate, and we conclude that the NBL budget approach is a useful tool to validate emission inventory estimates.

20 1 Introduction

Efforts to mitigate climate change via commitments to reduce national emissions require independent techniques to validate or at least constrain emission estimates reported by each nation. While anthropogenic CO_2 emissions from fossil fuel combustion can be estimated quite accurately from the commercial census data used to determine national tax obligations, the emission rates of CH_4

25 from non-fossil sources are much harder to determine. Here we aim at quantifying the farm-scale CH_4 emissions from a farmstead in Switzerland in order to compare these flux estimates with best estimates used in the national inventory report (NIR) under the Kyoto protocol.

The governmental inventory estimates are based on standard procedures recommended by the IPCC (Lassey, 2008; Nisbet and Weiss, 2010; Heimann, 2011; Wang et al., 2011), e.g., national live-

- 30 stock numbers multiplied with default emission factors (EFs) derived from the IPCC Guidelines and Guidances (Penman et al., 2000). Despite considerable uncertainty of these default values, this estimation methodology has become commonly used in order to ensure conformity and comparability of reported greenhouse gas emissions among different nations (Lassey, 2007). Although an experimental validation of such inventory estimates by independent means would be highly beneficial to
- 35 improve their overall credibility, the assessment of agricultural CH_4 emissions by direct atmospheric measurements is still lacking also for Switzerland. Methane emissions in Switzerland are dominated by the agricultural sector (77.7 %), followed by emissions from waste treatment (16.3 %) and from the energy sector (5.9 %) (FOEN, 2015, data from year 2013).

Because of the high global warming potential of CH₄ and due to its relatively short atmospheric
residence time, already a small reduction of CH₄ emissions will have a high impact on the global CH₄ budget (Dlugokencky et al., 2011; Wang et al., 2011). But in order to formulate feasible mitigation strategies, a deeper understanding and a precise quantification of emission variability and source strength are needed (Lowry et al., 2001; Ulyatt et al., 2002; Dengel et al., 2011).

Yet, due to the lack of direct field experiments and poorly known EFs (Lowry et al., 2001; Lassey,
2007), the NIR estimates are associated with uncertainties of ±18.1 and ±54.4% for enteric fermentation and manure management, respectively (FOEN, 2015). Although recent studies conducted at the animal husbandry level in Switzerland (Staerfl et al., 2012; Zeitz et al., 2012) showed that using country-specific EFs would not substantially alter the total estimate of livestock CH₄ emissions, high uncertainty remains, as differing farming practices can have significant impacts on the EFs (Lowry et al., 2001; Saggar et al., 2004; Christie et al., 2012).

Several methods exist to quantify CH_4 emissions from cattle, e.g., using SF_6 tracer (e.g., Johnson et al., 1994; Deighton et al., 2014) to determine emissions per animal, or via the release of an external N₂O tracer (e.g., Griffith et al., 2008) (e.g., Griffith et al., 2008; IPCC, 2006) to determine the emissions of enteric fermentation from groups of animals. These methods have tracer

55 techniques have been useful for the quantification of known CH_4 source components and especially of individual emissions per animal, requiring detailed knowledge of the transport pathways of tracers and the location of active CH_4 sources. However, such methods tend to have some limitations as the scale of integration increases to a whole farmstead or even larger spatial scale. For the validation of an emission inventory that is supposed to cover the total of all known and unknown (or neglected

60 source components), mostly volume-integrated budget estimates are used, since such an approach has the potential to also reveal unexpected or unexpectedly large flux components of CH_4 .

<u>These methods have</u> been useful for the quantification of known CH_4 source components, but tend to have some limitations as the scale of integration increases to a whole farmstead or even larger spatial scales. Moreover, these tracer techniques require detailed knowledge of the transport pathways

- of tracers and the location of active CH_4 sources with which they must be collocated to provide realistic flux estimates. However, for the validation of an emission inventory that is supposed to cover the total of all known and unknown (or neglected) source components, mostly volume-integrated budget estimates are used, since such an approach has the potential to also reveal unexpected or unexpectedly large flux components of CH_4 .
- Aircraft campaigns measuring vertical CH_4 profiles (Fowler et al., 1996; Denmead et al., 2000; Lassey et al., 2000; Wratt et al., 2001) or horizontal CH_4 transects (Hiller et al., 2014b) have successfully provided regional-scale ($\geq 10^4 \text{ km}^2$) flux estimates. For groups of cattle at the field or paddock scale ($< 0.5 \text{ km}^2$), tracer release (Griffith et al., 2008; Leytem et al., 2011), the integrated horizontal flux method (Laubach and Kelliher, 2004; Griffith et al., 2008), and the point-source dispersion
- 75 modeling and measurement approach (McGinn et al., 2011; McGinn and Beauchemin, 2012) have demonstrated their usefulness. At the farm scale in-between (≈0.5–5 km²), McGinn et al. (2006) have used a combination of tracer release, line-averaged concentration measurements, and inverse modeling of fluxes.

Inventory estimates neither account for the large spatial heterogeneity nor for the temporal variability of CH₄ sources (Wang et al., 2011). While spatially explicit inventories improve the overall spatial representativeness of the NIR, their accuracy still remains unclear due to the lack of small-scale validation data. This underpins the need for detailed local to regional-scale measurements.

Little is known about the temporal variability and spatial heterogeneity of different CH_4 sources, although an increasing number of studies show the importance of assessing CH_4 bud-

- gets via atmospheric measurements (Fowler et al., 1996; Beswick et al., 1998; Ulyatt et al., 2002; Zinchenko et al., 2002; Pattey et al., 2006; Detto et al., 2010; Pendall et al., 2010; Harper et al., 2011; Baldocchi et al., 2012). So far, a fair number of studies estimated livestock
 CH4 emissions via atmospheric concentration measurments without the deployment of chambers (Judd et al., 1999; Denmead et al., 2000; Laubach and Kelliher, 2004, 2005a, b; Laubach et. al., 2008; Grobler et al., 2014).
- 90 Unfortunately, only few studies compare their results to inventory estimates (Levin et al., 1999; Lowry et al., 2001; Hsu et al., 2010; Hiller et al., 2014b), and only few projects, so far, have directly validated livestock emission estimates via atmospheric concentration measurements without the

deployment of chambers (Denmead et al., 2000; Grobler et al., 2014) directly

- 95 (Levin et al., 1999; Lowry et al., 2001; Hsu et al., 2010; Hiller et al., 2014b). Using different budgeting approaches (e.g., convective and nocturnal boundary layer budgets, mass balance method), Denmead et al. (2000) found a good agreement of budget estimates with the respective inventory estimates.
- Therefore, in this study, we focus on the farm scale, by (1) quantifying the CH₄ source strength
 of a typical Swiss farmstead (0.5–5 km²) via nocturnal boundary layer (NBL) CH₄ budget fluxes (average surface fluxes) using vertical CH₄ concentration profiles within the atmospheric boundary layer. In addition, (2) we use experimentally derived NBL budget fluxes for inventory validation at different spatial resolutions, i.e., the NIR and the Swiss spatially explicit inventory (SEI) as well as a direct inventory estimate at the farm scale.

105 2 Materials and methods

2.1 Site description

We performed two intensive measurement campaigns during 16–17 August 2011 and 24–27 July 2012 at the ETH research station Chamau ($47^{\circ}12'37''$ N, $8^{\circ}24'38''$ E at 393 m a.s.l.), which is located in the lower Reuss Valley in central Switzerland. 56.5 % of the Reuss Valley is used for

- 110 agriculture (FSO, 2012), which is the most dominant land use category besides forests (22.5 %). Of the station's total area (62.03 ha), 72.4 % is covered by grassland used for grazing and forage production (Zeeman et al., 2010; Suter, 2011). Cropland used for silage maize production, forests and farm infrastructure cover 11.07 %, 8.28 % and 8.25 %, respectively. The grasslands are intensively managed (cut and fertilized about 5 to 6 times a year; detailed management information for year
- 115 2012 is given by Merbold et al., 2014) and dominated by mixed ryegrass-clover vegetation (Gilgen and Buchmann, 2009). The measurement periods did not include management events (e.g., cutting or fertilization). With respect to CH₄, the grasslands have shown negligible soil CH₄ fluxes on a daily timescale while acting as a small sink on an annual timescale (Imer et al., 2013). Livestock populations at Chamau vary according to the seasonal three-stage farming system. In summer, most cattle
- 120 are moved to higher altitudes, whereas swine, goats and sheep stay stationary at the farm. During our measurement periods, cattle and swine were inside the barns, while goats and sheep were located outside at the meadows.

Local meteorology in the Reuss Valley is dominated by a prominent valley wind regime, where prevailing winds from the NNW towards the Alps during the day are replaced by cold air drainage

125 flows from SSE during the night, promoting the development of a shallow nocturnal boundary layer. Information was available from an earlier project at a nearby location where meteorological tethered balloon soundings were carried out in the lowest 1500 m of the atmospheric boundary layer (Messerli, 1992; Utiger, 1992). These pilot studies showed that the stable nocturnal boundary-layer is rather shallow and restricted to the lowest $\approx 150-200$ m of the atmospheric boundary layer (ABL)

130 during fair weather conditions. Thus, our measurement periods were focusing on fair weather conditions, with a predominant high-pressure system and clear skies during night, when it was expected that probing the lowest 150–200 m of the ABL was sufficient for CH_4 budget calculations via the nocturnal boundary-layer budgeting method.

2.2 Vertical balloon profiles and ground surface measurements

- 135 Tethered balloon measurements were performed at a distance of 150 m from the nearest farm buildings during the night in both years (2011 and 2012) with different temporal coverage (see Table 1 for details). A helium-filled blimp (7.2 m³, The Blimp Works Inc., Statesville, NC, USA) with a net lift of 4.35 kg carried a polyethylene inlet tube of 220 m length with inner diameter of 4 mm (Maagtechnic, Dübendorf, Switzerland) connected to a ground measurement station. With this approach it was
- 140 possible to keep sensitive, heavy, and power consuming instrumentation on the ground at all times. The air sampled was drawn at a flow rate of $\approx 1 \,\mathrm{L}\,\mathrm{min}^{-1}$ (Reynolds number 344) to a fast greenhouse gas analyzer (FGGA, Los Gatos Research Inc., Mountain View, CA, USA). With this approach it was possible to keep sensitive, heavy, and power consuming instrumentation on the ground at all times.
- 145 The precision of the CH₄ concentration measurements, converted to CH₄ dry mole fractions by the FGGA firmware, was $\sigma_{1s} = 1$ ppb and $\sigma_{min} = 0.15$ ppb at 1 s and 200 s integration time, respectively. These values were determined from the Allan variance of a 1 h time series measured in a climate chamber under controlled conditions (1927.7 ± 1.1 ppb, 25 °C, 40 % relative humidity; data not shown).
- Additional measurements of air temperature, atmospheric pressure, relative humidity, wind speed and wind direction were carried out with meteorological tether sondes TS-5A-SEN (Atmospheric Instrument Research, Inc., Boulder, USA), which were attached to the tether line. Each TS-5A-SEN sonde contained a temperature and relative humidity probe in a horizontal radiation shield, with its opening facing the wind direction in order to vent the sensors. A pressure sensor allowed to determine the hydrostatic elevation above ground surface, and a cup anemometer determined
- horizontal wind speed. The wind direction was measured via a built-in compass attached to the tail of the TS-5A-SEN sonde that acted as a wind vane centered at the tether of the balloon.

These meteorological measurements were transmitted via an AIR IS-5A-RCR radio receiver (395–410 MHz) to a Linux computer running an in-house software to collect data. The sondes and

160 the radio receiver correspond to the instruments used in earlier studies (Eugster and Siegrist, 2000; Siegrist, 2001). Pressure (hPa), temperature (°C) and relative humidity (%) were transmitted with a resolution of two decimals, one decimal was transmitted for wind speed (m s⁻¹), while wind direction (°) was transmitted in full numbers. A nearby eddy covariance station provided micrometeorological measurements of air temperature,

- 165 relative humidity, wind speed, wind direction, photosynthetic active radiation (PAR) and turbulence (Zeeman et al., 2010; Merbold et al., 2014), which were used for cross-referencing the meteorological tether sondes at times when the tethered balloon and the sondes were near the ground (ca. 1–2 m height). However, the accuracy of the sensors in the TS-5A-SEN sonde remained clearly below the resolution that the sondes transmitted their data, even after such an intercalibration. The manu-
- 170 facturer did not specify their accuracy, but experience showed that temperature, relative humidity, pressure, wind speed and wind direction may be subject to uncertainties on the order of ± 0.2 °C, $\pm 3-5$ %, ± 0.2 hPa, ± 0.2 m s⁻¹, and $\pm 5^{\circ}$, respectively.

A GPS unit (eflight, SM-Modellbau.de, Germany) provided positioning and height information of the balloon system. In total, 18 soundings were conducted, of which 16 were used for further

- 175 analysis, while two balloon soundings (16/17 August 2011 during 04:48 and 05:56 UTC+1) were rejected due to unfavorable meteorological conditions. Each ascent and descent of the remaining 16 soundings gave a mean vertical profile and a mean NBL budget flux (Table 1). All dates and times are given in Central European Time (UTC+1 h).
- In order to also obtain a good data coverage of the near-surface conditions where barometric height measurements of a tethered balloon tend to be not accurate enough, a guy-wired tower was installed next to the FGGA. Inlet tubes (Maagtechnic, Dübendorf, Switzerland) were installed on the tower at eight different heights (0.2, 0.5, 1, 2, 3, 5, 8 and 10 m a.g.l.), guiding the air into a home-built air inlet selection unit (Zeeman et al., 2008). The tower measurements were performed between balloon soundings, with each height being measured for three minutes.
- Two modes of operation with the tethered balloon were tested: during the 2011 campaign, conventional ascent/descent measurements were made using an ascent and descent velocity on the order of 0.15 m s^{-1} . During the 2012 campaign, an attempt was made to keep the gas inlet at several fixed heights during the ascent to get a better quality of the measured concentration changes over time for these specific layer heights, at the expense of a poorer vertical resolution of the measurements.

190 2.3 Budget flux calculation

The NBL budget technique was applied at night during favorable weather conditions with clear skies and low wind speeds. In principle, nighttime conditions should allow to derive regional-scale average fluxes that can be compared to inventory estimates if the emission flux itself does not show a pronounced diel cycle, so that the diel cycle in concentrations measured in the lowest part of the NBL are primerily a function of (neutureal us, diurnal) atmospheric stability and stratification.

195 NBL are primarily a function of (nocturnal vs. diurnal) atmospheric stability and stratification.

The budget flux F_s , i.e., the average surface flux, can be expressed as follows (Denmead et al., 1996):

$$F_{\rm s} = \int_{0}^{z_i} \frac{\partial \bar{c}}{\partial t} \,\mathrm{d}z \,, \tag{1}$$

where $\partial \bar{c} / \partial t$ is the rate of change in concentration with time, and z_i is the NBL height. Overbars

- 200 indicate averaging over time or space, depending on context. The resulting budget flux describes the accumulation of a scalar within the stable boundary layer at consecutive time steps. In practice, finite differences are used to approximate $\partial \bar{c}/\partial t$, which is expressed as $\Delta \bar{c}/\Delta t$ hereafter. We used time–space kriging to interpolate CH₄ concentrations (see Section 2.5) from the tethered balloon and the 10 m tower.
- The NBL budget method is based on the idea that during nights with ideal weather conditions, a stable nocturnal boundary layer develops, which is capped by a strong temperature inversion at its top that acts as a natural atmospheric chamber, accumulating all emissions of underlying sources (Mathieu et al., 2005; Pattey et al., 2006). Thus, the top of the NBL and hence z_i will be characterized by Δc̄/Δt = 0 ppm s⁻¹ due to a neutral to unstable stratification with Δθ̄/Δz ≤ 0 K m⁻¹. In our case, the budget flux computed with Eq. (1) is an integral measurement of the net flux from all rele-
- vant sources and sinks within the footprint of the balloon measurements. These relevant components can be specified in more detail by:

$$F_{\rm s} = F_{\rm soil} + F_{\rm ent} + F_{\rm HA} + F_{\rm other} , \qquad (2)$$

where F_{soil} is the soil exchange flux, F_{ent} the entrainment flux at the top of the NBL, F_{HA} the horizontal advection, and F_{other} the flux resulting from emissions of other sources in the footprint. Under stationary conditions with a constant NBL height, F_{ent} becomes negligible. At the Chamau site, earlier chamber flux measurements showed that F_{soil} is a small component (-0.0024 µg CH₄ m⁻² s⁻¹ on average, Imer et al., 2013) representing typically less than 1 % of the expected F_s and hence can be neglected in Eq. (2). This minor role of the source and sink strengths of upland grasslands in
the total greenhouse gas budget of a farm has also been found at other sites (e.g., Hartmann et al., 2011, range -0.015 to 0.013 µg CH₄ m⁻² s⁻¹ for single chamber measurements) and in incubation studies (e.g., Wang et al., 1999, maximum uptake of -0.016 µg CH₄ m⁻² s⁻¹ during first 8 h of incu-

bation at 24 °C). Furthermore, the advection term can be neglected if measurements are taken over a horizontally sufficiently large and flat terrain with a homogeneous source distribution (Raupach et al., 1992; Choularton et al., 1995; Fowler et al., 1996; Beswick et al., 1998; Denmead et al., 2000; Cleugh et al., 2004; Mathieu et al., 2005; Pattey et al., 2006). With these simplifications, Eq. (2) can be reduced to:

$$F_{\rm s} \approx F_{\rm other}$$
, (3)

where F_{other} in our case represents CH₄ emissions from the cattle and their products stored at the 230 Chamau farmstead.

Nevertheless, it has been shown elsewhere that even under stable conditions, small advective effects and wind direction play an important role in determining the NBL budget (Lowry et al., 2001; Zinchenko et al., 2002; Hsu et al., 2010). Advected air parcels can include pollutants from sources

lying in the upwind fetch and hence can bias the local CH₄ concentrations, resulting in an overes-

- timation of the NBL budget. In particular, nighttime measurements can include regional information of up to several kilometers upwind (Bamberger et al., 2014). In order to avoid the impact of emissions from upwind sources and to reduce the measurement footprint to an extent, which only includes the barn buildings of the Chamau farmstead, the integration height was set to the height where $\Delta \bar{\theta} / \Delta z \leq 0 \text{ K m}^{-1}$ (Mathieu et al., 2005; Pattey et al., 2006) with θ being the potential
- 240 temperature ($\theta = T (p_0/p)^{1.402}$ with $p_0 = 1000$ hPa and p ambient pressure). In addition, only measurements with a prevailing an average wind direction from the SSE–SW (157.5°–225°) sector were used for the flux calculation, i.e., where the main barn building of the Chamau station and the grazed pastures were located. During very calm weather conditions when mean wind speeds reached zero, measurements were not restricted to wind directions. The impact of neglected horizontal advective
- 245 processes on the overall NBL budget estimates will be part of the discussion section.

2.4 Inventory estimates

Two inventory estimates of CH_4 emissions exist for Switzerland (i.e., NIR and SEI), which both use the same estimation methodology, but differ in their spatial resolution. While the NIR evaluated CH_4 emissions on an annual basis for entire Switzerland (FOEN, 2015), the recently developed

- SEI (Hiller et al., 2014a) distributed the CH_4 emissions, based on the 2007 year stocking census data (FSO, 2009), onto a $500 \text{ m} \times 500 \text{ m}$ grid according to Swiss land use statistics (FSO, 2007), and then scaled the emissions to represent 2011 livestock numbers (see Hiller et al., 2014a). Both inventories estimate the CH_4 emissions from national livestock numbers multiplied with animalspecific emission factors, i.e., the CH_4 conversion rate for enteric fermentation y_m and the CH_4
- 255 conversion factor (MCF) for manure management, which are both given by the IPCC Good Practice Guidance (Soliva, 2006; ART, 2012).

The SEI accounts for the large spatial heterogeneity of CH₄ sources, which the NIR cannot resolve. In the SEI, the Chamau research station was explicitly resolved by several grid cells, with CH₄ emissions ranging from 6 kg CH₄ ha⁻¹ yr⁻¹ to 820 kg CH₄ ha⁻¹ yr⁻¹. This large variation among the individual grid cells arises due to the standardized assignment process of the SEI: about 80 % of the CH₄ emissions are assigned to the grid cell with the main building of a farmstead, while the remaining emissions are evenly distributed to the pasture area of the entire municipality (Hiller et al., 2014a).

In order to compare inventory estimates to budget fluxes, the NIR estimates of enteric fermentation 265 $(119 \pm 21.5 \text{ Gg CH}_4 \text{ yr}^{-1})$ and manure management $(31 \pm 17 \text{ Gg CH}_4 \text{ yr}^{-1})$ for the year 2011 were disaggregated to the area used for pastures (here 5685.8 km²), according to the CORINE land use map (FSO, 1998). For the SEI, the grid cell representing the main building of the research station was used for comparison. To overcome the inconsistencies in spatial resolution and temporal representativeness of both, the

- 270 NIR and the SEI, we specifically estimated CH_4 emissions for the Chamau farmstead (in the following called CHAI as an acronym for **CHA**mau Inventory estimate), using the same EFs as for the SEI, but multiplied with actual stocking densities of the respective measurement periods. This simplified the validation, as livestock represents the main CH_4 source at the Chamau station and hence, the emission of the whole farmstead is included in the CHAI estimates without the need for spatial dis-
- 275 aggregation. The CHAI inventory estimates include different animal headcounts for the years 2011 and 2012, respectively, . In 2011, 55 diary cattle, seven non-diary cattle, nine young cattle, 18 sheep and 129 swine were located at the Chamau farmstead. The inventory estimate for 2012 includes besides 81 diary cattle, two non-diary cattle, 18 young cattle, eight sheep, 47 swine also 39 goats. Since no uncertainty estimates existed for the SEI and CHAI, we assumed the same methodology-
- 280 specific uncertainty as for the NIR. The systematic uncertainty due to the spatial disaggregation of the SEI remains unknown. The standard deviation of all three inventory estimates (i.e., NIR, SEI and CHAI) was calculated as the average uncertainty estimates of enteric fermentation (18.1%) and manure management (54.4%) from the NIR (FOEN, 2015), yielding 36.3%.

2.5 Data processing and general conventions

- 285 Data processing, analysis and flux calculations were done with the statistics software package R, version 2.15.1 (R Core Team, 2014). Nighttime data were defined as $PAR < 20 \,\mu mol \, m^{-2} \, s^{-1}$. Ordinary kriging using the *geoR*-package was performed for spatio-temporal interpolation of CH₄ concentrations and served as a basis for flux calculations.
- Before kriging, the timestamps of the tethered balloon system and the 10 m tower system were synchronized. The grid spacing for the kriging was chosen to be 0.3 h along the time axis in both years. In the vertical direction, the grid spacing was set to 0.2 m between 0 and 10 m a.g.l., where tower profile data were used, and to 5 m between 10 m and 200 m a.g.l., where tethered balloon data were used. In practice, the maximum height that could be reached with the 220 m hose attached to the tethered balloon was roughly 150 m a.g.l. Horizontal drift with the mean wind, and dew formation on the balloon lowering payload capacity were responsible for this height limit.
 - NBL budget flux calculations were restricted to the selection criteria mentioned in Sect. 2.3. Based on the kriged time-height concentration field, the NBL budget flux was calculated separately for each of the balloon soundings using the full duration of each sounding ± 30 min. In practice, the difference between two consecutive time slices was used to approximate the local derivative of concentration
- 300 over time (in ppm s⁻¹), from which the fluxes could be computed by multiplying that value with the thickness of the layer (m) and summing up over the NBL extent covered by measurements and interpolated values. These raw fluxes (in ppm m s⁻¹) were then converted to μ g CH₄ m⁻² s⁻¹ by taking the actual molar density of air into account (computed from ambient pressure and temperature), and the molar mass of CH₄. If not stated otherwise, reported values denote means ±

305 standard deviations. The terms N1–N4 and S1–S16 refer to the different nights and soundings in the respective years (Table 1).

An uncertainty assessment (Sect. 4.1) was performed which also included heights above 200 m up to a z_i of 400 m, which is 2.5× the typical NBL inversion heights observed by Messerli (1992) and Utiger (1992). Since these heights were above the maximum height reached by the tethered balloon

- 310 system, the following approach was employed. A background concentration was determined from the minimum CH_4 concentration observed during each field campaign (diurnal and nocturnal data). Between 200 m (top level of kriged interpolation) and the background at a predefined $z_i > 200$ m, a linear decrease in CH_4 concentrations with height was assumed. This is probably overestimating the true rate of concentration changes at heights > 200 m (which may follow an exponential rather
- than linear decrease with height), and hence our uncertainty estimate should be considered a conservative estimate of potential systematic errors in NBL budget flux estimates in cases where the determination of z_i remains highly uncertain.

3 Results

3.1 Weather conditions and diel course of methane concentrations

- 320 During our measurements in 2011 and 2012, the weather was dominated by a high-pressure system over Central Europe. We experienced warm and dry daytime conditions with temperatures reaching 302.1 K in 2011 and 306.1 K in 2012, and a pronounced surface cooling with clear skies and low wind speeds during all nights (Fig. 1). Radiative cooling (i.e., $\Delta \overline{T} / \Delta t \leq 0$ K h⁻¹) was most pronounced immediately after sunset, providing sufficient atmospheric stability and stationarity for the
- build-up of a NBL. Later during the nights, cooling rates remained negative and enabled the deepening of the stable NBL. The nighttime cooling was most pronounced during N1 in 2011 (286.1 K), whereas air temperatures in N2–N4 in 2012 did not fall below 287.1 K, with the highest nighttime value of 288.7 K being observed during N4. All nights were dominated by low near-surface wind speeds < 1 m s⁻¹, associated with very weak turbulence with friction velocities $u_* < 0.08 \text{ ms}^{-1}$. In
- 330 2012, however, some measurement periods experienced slightly increased wind speeds and friction velocities, probably due to weak advection promoting mixing processes within the stable NBL.

The diel courses of CH_4 concentrations mainly followed the meteorological conditions. Due to convective mixing during daytime, CH_4 concentrations remained relatively low (≈ 1.9 ppm). At night, concentration values increased as the emissions were trapped within the smaller volume of the

335 stable NBL (exemplarily shown for N1, Fig. 2a). Highest mixing ratios, here up to 3.1 ppm, were observed around 03:00 LT at the lowest measurement height close to the surface (i.e., 0.2 m a.g.l.) due to inhibited vertical mixing. For lower measurement heights, the nighttime accumulation of CH₄ started after sunset. Measurements taken from above 70 m a.g.l., however, only showed an increase in CH₄ concentrations during the second half of the night, resulting in a less pronounced diel pattern

compared to lower elevations (Fig. 2b). Besides reaching the top of the NBL at about 70 m a.g.l., variable wind directions affected the nocturnal increase in CH₄ concentrations. The nighttime CH₄ build-up was most pronounced when air masses originated from the wind sector between 180 and 270°, including the cattle barn buildings (in Fig. 2b, at heights below 50 m a.g.l. from 21:00 to 23:00 LT). In contrast, air masses with a northern wind component, here above 100 m a.g.l., showed constant or decreasing CH₄ concentrations during the same measurement period.

3.2 Vertical balloon profiles

Potential temperature gradients of successive vertical balloon profiles indicated stable conditions during all nights (i.e., Δθ/Δz > 0 K m⁻¹) and hence the establishment of a stable NBL (Fig. 3a). However, the height of this NBL, where a transition from stable to neutral or unstable conditions is
expected, strongly varied among the individual profiles. Half of the profiles revealed a temperature inversion at or above 100 m a.g.l. (i.e., S1 and S3 of N1; S7 and S8 of N2; S12 of N3; S14, S15 and S16 of N4). The other half exhibited multiple transition heights, pointing to the development of multiple stable layers within the NBL. Only few soundings (i.e., S5 of N1; S6 of N2; S13 of N4) showed a clear temperature inversion (at 50–60 m a.g.l.), followed by a second stable layer with

- 355 positive temperature gradients with increasing elevation. Other soundings were rather marked by a gradual transition from stable to neutral conditions (e.g., S10 and S11 of N3). However, temperature gradients among individual soundings differed markedly, indicating a temporal evolution (e.g. growth, consolidation or separation) of the different layers within the NBL.
- The CH₄ concentration profiles showed quite different patterns between the two years. In 2011 360 (S1–S5), the profiles exhibited a clear increase in mixing ratios from S1 to S5 within the lowest 100 m, and all profiles reached a background value of ≈ 2 ppm above 100 m a.g.l. (Figure 3b), pointing to the main NBL inversion height. This drop in concentration was associated with a change in wind direction from SSW to N and an increase in wind speeds up to 4 m s⁻¹ (Fig. 3c). In 2012 (S6 to S16), all profiles indicated a well-mixed NBL with small or even absent vertical concentration
- 365 gradients, but with a substantial increase in concentration over time (Fig. 3b). Even though temperature inversions were present, none of the CH_4 concentration profiles reached a background value at or above the inversion height. The same well-mixed conditions were found for the two rejected soundings in 2011. However, no clear changes in wind direction could be observed in the profiles from 2012. Most wind profiles meandered between a WSW sector in the lower part of the NBL and
- an ESE sector in the upper part of the NBL. Only S12 of N3 showed an abrupt change in wind direction from SE to N at 50 m a.g.l. Wind speed in 2012 showed a slightly different picture compared to 2011 as well (data not shown). Except S6 and S8 of N2, all profiles revealed increasing wind speeds with height. Some soundings also showed a peak with maximum wind speeds (i.e., S9 of N2 at 80 m a.g.l.; S11 of N3 at 130 m a.g.l.; S14 and S15 of N4 at 140 and 50 m a.g.l., respectively), indicative
- 375 of an active low-level jet.

3.3 NBL budget fluxes and comparison to inventory estimates

The observed CH_4 concentration and potential temperature gradients showed that the overall NBL at Chamau was not merely influenced by local emissions from the farmstead. The well-mixed CH_4 concentration profiles in 2012 suggested that the NBL was primarily influenced by emissions from

- sources further upwind, which were already well mixed with ambient air, and that local sources only had a minor influence. Restricting the NBL budget either to the height where $\Delta \bar{\theta} / \Delta z \leq 0 \,\mathrm{K \,m^{-1}}$ and/or to wind directions originating from the SSE–SW sector allowed the separation of emissions from the Chamau station vs. emissions from upwind lying sources (see Sect. 2.3). The resulting NBL budget fluxes of the individual profiles varied between 0.79 ± 0.13 and $2.88 \pm 0.61 \,\mathrm{\mu g \, CH_4 \, m^{-2} \, s^{-1}}$
- during the different nights (Table 1), with an overall mean of $1.60\pm0.31 \,\mu\text{g}\,\text{CH}_4 \,\text{m}^{-2} \,\text{s}^{-1}$. All nights showed a clear nocturnal course of NBL budget fluxes, with higher fluxes in the middle of the night and lower fluxes shortly after sunset and before sunrise. This nocturnal course was most pronounced during N2 and N3 in 2012 due to a larger gradient_difference in consecutive temperature and CH₄ profiles.
- Inventory estimates of the Chamau farmstead varied markably markedly among the different inventories (Table 2). Highest values were derived for CHAI with 1.29 ± 0.47 and $1.74 \pm 0.63 \,\mu g \, CH_4 \, m^{-2} \, s^{-1}$ in 2011 and in 2012, respectively. Lower emission fluxes of $0.83 \pm 0.30 \,\mu g \, CH_4 \, m^{-2} \, s^{-1}$ (for all nights in both years) resulted from the spatial disaggregation of the NIR estimates. The SEI emission flux estimate for the grid cell including the main building of the
- 395 farmstead, however, was only 0.42±0.15 µg CH₄ m⁻² s⁻¹, significantly lower than the CHAI estimates. These CHAI estimates, representing the same spatial extent as the main SEI grid cell, were therefore at least three times higher than the SEI, although 80% of total CH₄ emissions from the Chamau farmstead were covered by this main grid cell. Mean NBL budget fluxes for the different nights ranged from 1.40±0.50 to 2.05±0.45 µg CH₄ m⁻² s⁻¹ and were comparable to the CHAI
- 400 estimates (Table 2).

4 Discussion

4.1 Flux uncertainty assessment

405

The determination of the NBL budget flux using Eq. (1) involved the estimation of the inversion height z_i which was considered the top of the relevant layer for the integration of concentration changes to derive F_s . It was however not always easy to determine z_i accurately from available tethered ballon data, namely in cases where the balloon did not clearly rise beyond z_i . Hence, we performed an assessment of the uncertainty of our flux estimates with the NBL budget method. We used the criteria $\Delta \bar{c} / \Delta t = 0$ ppm s⁻¹ under neutral to unstable stratification with $\Delta \bar{\theta} / \Delta z \leq 0$ K m^{-1} for integration of Eq. (1). Therefore, we used relative humidity as an independent variable to 410 assess the uncertainty in our estimate.

During the first part of the night on 16 August 2011 with a growing NBL after 21:30 h (Fig. 4a), the bottom measurement before launching the tethered balloon showed 100 % relative humidity, and hence the marked transition from the saturated near-surface layer to the drier air aloft was clearly seen in both ascents and descents of the balloon. As expected, z_i estimated from ascents did not

- 415 exactly match the values derived from consecutive descents (Fig. 4a), but in contrary to initial expectations, z_i derived from ascents were lower than values derived from descents, with a difference of typically up to 20–30 m. This means that even with a slow ascent $(0.16 \pm 0.03 \text{ m s}^{-1})$, mean \pm standard deviation) and slow descent $(-0.14 \pm 0.03 \text{ m s}^{-1})$ in the example in Fig. 4a, the ascending balloon seemed to perturb the stable atmosphere in a way that the drier air aloft was mixed down to
- 420 the sensor as the balloon rose, whereas during the descent, the sensor first crossed z_i , followed by the balloon, which explained the higher z_i values measured during descents.

Thus, similarly to what was seen with relative humidity, it must be expected that there was some unavoidable mixing of CH_4 in the vertical profiles probed by the tethered balloon that translated to uncertainties in computing the NBL volume budget, both via (a) uncertainties of determining the

- 425 exact height from where a gas sample originated in the vertical profile, and (b) from the associated uncertainty in determining z_i from tethered balloon soundings. As long as the prominent features of increased CH₄ concentrations in the vertical profile mostly remained below z_i as determined from relative humidity (Fig. 4b), the NBL budget fluxes are expected to be rather robust against uncertainties in z_i . The small-scale layer structures evolving in the NBL are in fact emission plumes that
- 430 spread horizontally at a constant buoyancy level (see Fig. 4b), difficult to predict from local measurements alone. The unknown variability of such plumes between two subsequent balloon soundings appeared to be the most problematic source of uncertainty in NBL budget flux estimates.

To assess the uncertainty involved in the NBL budget flux estimates presented in Sect. 3.3 and Table 1, we also calculated the range of fluxes that would result from fixed inversion heights (and

- thus integration heights) z_i of 50, 100, 150, 200, 300 and 400 m, using all tethered balloon sounding data (Fig. 5). In addition to uncertainties in determining z_i , the effect of interpolating extrapolating available data beyond the extent of the maximum height of an actual sounding was assessed (green circles in Fig. 5). For comparison, we repeated all calculations with the subsets of data obtained from ascents and descents only (not shown).
- Using both ascents and descents in flux calculations (Fig. 5) was very comparable to using ascents only, or descents only (not shown). Ascents led to a slightly lower sensitivity of flux calculations to z_i estimates, whereas calculations based only on descents showed a higher sensitivity. In principle, we expected the reverse: when a tethered balloon was lowered into the stable atmosphere below it, the meteorological sensors and gas inlet penetrated the stable layer first, followed by the disturbance
- 445 of the balloon. Hence it was not entirely clear why the ascents appeared to perform better than the

descents, and thus we used both ascents and descents in combination to avoid adding unnecessary bias to our calculations.

In cases of instationary conditions – which were quite frequent during calm nights – the systematic uncertainty was so large that not even the sign of the flux estimate was certain, which led

- 450 to a substantial number of soundings that had to be discarded from the analyses (gray bands in Fig. 5). Thus, although there are certainly further components that add to random uncertainty in our estimates, the two main issues to overcome in future projects are (i) extending the vertical layer over which high-quality concentration measurements can be made; and (ii) using profiling at slow but steady ascent rates to obtain consistent vertical profiles of all meteorological variables that better
- 455 allows to detect instationary conditions, and which clearly includes the transition across z_i , which is a critical value in flux estimates. Even with limited payload it may be advisable to have to tethersondes attached to the balloon tether, one which measures the conditions 50–100 m above the tube inlet (of limited length), and another at the height of the tube inlet. In such a way at least conditions where concentration measurements close to z_i are covered, the upper tethersonde would allow to establish
- 460 that these measurements were actually close to z_i . In most cases a distinct drop in relative humidity is a clear sign of z_i .

4.2 Using the NBL budget approach in complex terrain

In contrast to other studies using the NBL budget approach at horizontally homogeneous and flat sites, our measurements were located in complex terrain with variable NBL structures. Temperature,
CH₄ concentration and wind profiles showed that the NBL in the Reuss Valley underwent a significant temporal evolution, with the establishment of multiple stable layers and with a high dependency on wind direction. With its location north of the Swiss pre-Alps, the NBL was primarily affected by the valley wind system and the resulting thermodynamic processes in the atmosphere. The complex

temporal evolution of our temperature, wind direction and wind speed profiles (Fig. 3) suggested

- 470 different disturbances (e.g., advection of air with higher CH_4 concentration, pulsating down-valley flows, cold air drainage), leading to turbulent mixing events within the lower and upper parts of the NBL. These disturbances finally resulted in well-mixed CH_4 concentration profiles throughout the NBL in 2012 and also in the rejected profiles in 2011, although clear vertical concentration gradients were expected due to the observed temperature inversions. These findings suggested that advection
- 475 and mixing processes played a prominent role during our measurements. In addition, turbulence and wind speed measurements of our eddy-covariance system suggested that intermittent turbulent processes affected the thermodynamic structure of the boundary layer. Although local meteorological conditions favored the build-up of a stable and stratified NBL, turbulent processes within the NBL have also been reported by other studies (Parker and Raman, 1993; Holden et al., 2000; Mahrt
- 480 and Vickers, 2002; Wang et al., 2006). In addition, some of our wind speed profiles indicated the presence of a low-level jet, which besides cold air drainage had a significant impact on vertical

transport within the NBL (Mathieu et al., 2005; Pinto et al., 2006; Sun et al., 2007). Hence, similar to Dorninger et al. (2011), the temporal evolution of the NBL in the Reuss Valley was primarily affected by local-scale mixing processes.

- Besides intermittent turbulence, advection processes challenge the applicability of the NBL budget approach for a feasible estimation of farm-scale CH_4 emissions. Although studies so far have focused on the regional scale with homogeneously distributed sources and hence neglected the advection term in the CH_4 budget, advection has also been reported in other studies as an important factor of the resulting budget flux (Kuck et al., 2000; Pattey et al., 2006). Since we had neglected
- 490 horizontal advection by assuming $F_{\text{HA}} \approx 0$, our NBL budget fluxes represented F_{other} . However, the CH₄ concentration and wind direction profiles suggested that CH₄ concentrations largely depended on wind direction (Fig. 2b). This indicated a heterogeneous distribution of CH₄ sources in the Reuss Valley and hence $F_{\text{HA}} \neq 0$, even though major parts of its relatively broad basin are used for agriculture. In addition, it has been shown that even nighttime observations measured at very low elevations
- 495 above ground (2 m a.g.l.) carry regional information up to 200 km^2 (Bamberger et al., 2014). An application of the NBL budget approach with $F_{\text{HA}} \approx 0$ as often done in other studies will thus result in budgets, which represent an enlarged spatial extent not limited to the Chamau farmstead. Furthermore, the fluxes would include horizontally advected air parcels that carry a CH₄ concentration signal that reflects different rates of agricultural emissions than those from the Chamau farmstead.
- 500 Therefore, in order to estimate emissions at a farm scale, we minimized uncertainties caused by horizontal advection from sources further upwind of the Chamau farmstead by carefully selecting the integration height of the NBL budget flux calculation within specific wind direction and wind speed conditions (see Sect. 2.3). This ensured that the main factor affecting the measurements was kept to the Chamau farmstead and hence reduced the influence of upwind lying sources through advection.
- 505 Still, the since the upper limit of the NBL height was not reached in 2012 and the well-mixed CH_4 concentration profiles pointed to the presence of advective processes, the emissions from sources further upwind might have had an impact on the resulting NBL budget estimate. Therefore, the uncertainty in flux estimates introduced by assumptions necessary with respect to z_i and the time-space interpolation between soundings (Sect. 4.1) revealed some limits of the NBL budget approach that
- 510 will require further attention in future studies.

The Overall, a good agreement of the NBL budget fluxes with the CHAI estimates showed that our emission estimates not only represented the emissions produced at the farm (Table 2), but also could be achieved which confirmed the viability of our selection criteria. Still, not all advective effects and external sources could be accounted for, as the NBL budget fluxes showed both a clear day-to-day

and a nocturnal variability (Tables 1 and 2, Fig. 5), which could not simply be explained by the diel course of livestock CH_4 production. Thus, the NBL budget approach not only integrates over a larger spatial scale, but can also be used to resolve mixing processes on smaller, i.e., hourly time scales.

4.3 Validating inventory estimates

- 520 The validation of inventory emission estimates via atmospheric measurements is not a trivial task. The main challenge lies in matching the different spatial and temporal scales of the inventories and the measurements. The inventory estimates represent annual means, which can deviate substantially from measurements taken at any given time. Seasonally changing farming practices and temporal variability in CH_4 emissions are not resolved within a typical inventory so far and can only be
- 525 covered by direct measurements. Additionally, the coarse resolution of the statistical database in SEI impaired a reliable comparison with atmospheric measurements with a subset of cells in the SEI; the concept of distributing emissions based on statistical considerations is expected to lower emission estimates from grid cells where cattle actually are found, but increases emissions from grid cells where in reality no cattle are found, an artifact that we tried to overcome via our CHAI estimate.
- A differentiation is needed between the validation (1) of livestock CH_4 emission estimates, and (2) of emission factors used for inventory estimates. Most studies address the validation of EFs via direct measurements at the individual animal level (Ulyatt et al., 2002; Zeitz et al., 2012). Our study, on the contrary, focused (a) on the validation of livestock CH_4 emission estimates, by examining the appropriateness of the methodology used in the inventory, and (b) on evaluating whether the
- estimates are consistent with direct atmospheric measurements (Lowry et al., 2001; Lassey, 2007). Using actual livestock data (CHAI), the inventory estimates were of the same order of magnitude as atmospheric measurements. Therefore, statistical assessments and inventories are indeed able to reliably represent CH_4 emissions from livestock, even though the estimates are based on given default values. Beside that, our study showed that the methodology used in the inventory estimates
- 540 is a suitable tool for resolving current livestock CH_4 emissions at farm scale and can at least for the period of data collection – be applied at larger scales as well, e.g., at regional to national scales.

5 Conclusions

Direct inventory estimates based on actual local livestock numbers (CHAI: 1.29 ± 0.47 and $1.74 \pm 0.63 \,\mu\text{g} \,\text{CH}_4 \,\text{m}^{-2} \,\text{s}^{-1}$ in 2011 and 2012, respectively) and mean NBL budget fluxes (ranging between 1.22 ± 0.1 and $2.05 \pm 0.45 \,\mu\text{g} \,\text{CH}_4 \,\text{m}^{-2} \,\text{s}^{-1}$) agreed well, after appropriate selection of avail-

tween 1.22 ± 0.1 and $2.05 \pm 0.45 \,\mu \text{g CH}_4 \,\text{m}^{-2} \,\text{s}^{-1}$) agreed well, after appropriate selection of available sounding data was made (Sect. 2.3). Thus, estimates based on actual livestock data and default emission factors showed that inventory assessments give plausible results, confirming the quality of the methodology used by such inventories.

Nevertheless, further improvements of the NBL budget flux approach are possible and are necessary for future studies. The sensitivity of the NBL approach to uncertain determinations of z_i is one limiting factor, and the completely unrealistic fluxes that can result during instationary conditions – which were quite frequent during our 2011 field campaign – are another limitation. CH₄ accumulations were generally greatest near the surface below the starting height of the tethered balloon system, and hence the combination with a tower profile covering the lowest heights above the surface (i.e., 0-10 m) was an essential component in our NBL budget flux calculations.

The multi-layer structure of the nocturnal atmosphere, with a typical inversion at around 50 m a.g.l. and opposing wind directions above and below, posed the greatest challenge for the NBL budget flux estimates. Hence, it is recommended to rather use the conventional approach with tethered balloon soundings effectuated at constant ascent and descent rates rather than the approach we

- 560 used in summer 2012 with prolonged measurements at fixed heights across the relevant lower part of the NBL. Although this latter approach allowed for a much better assessment of the $\Delta \bar{c}/\Delta t = 0$ ppm s⁻¹ criterion to estimate z_i in the NBL budget flux calculations, this comes at the cost of losing a clear visual control over z_i and potential additional shallower layers below z_i , which may remain undetected if no continuously measured temperature and relative humidity profiles are available.
- 565 In summary, the validation of inventory estimates via the NBL budget flux approach (with appropriate data selection) can be applied at different spatial scales, i.e., from a single farm to larger scales. Within the experimental uncertainty bounds, our study also raises confidence that the Swiss national inventory report reliably represents national livestock CH₄ emissions.

Acknowledgements. Funding for this project was provided by the Swiss National Science Foundation grants
no. 2.77771.10 and CRSII2_136273. We thank Peter Plüss and Thomas Baur for technical support, and Hans-Rudolf Wettstein and his team for their great efforts and support during the measurement campaigns at the ETH research station Chamau. Many thanks to Rebecca Hiller, Zürich, and Mathias Rotach, Innsbruck, for fruitful discussions and valuable comments. The spatially explicit inventory of Meteotest, Bern, was developed within the MAIOLICA project of the ETH Competence Center Environment and Sustainability (CCES).

575 References

590

- ART: Agricultural CH₄ and N₂O emissions in Switzerland, Agroscope Reckenholz Tänikon Research Station (ART), QA/QC Internal Report, Zurich, Switzerland, 61 pp., 2012.
- Baldocchi, D., Detto, M., Sonnentag, O., Verfaillie, J., The, Y., Silver, W., and Kelly, N.: The challenges of measuring methane fluxes and concentrations over a peatland pasture, Agr. Forest Meteorol., 153, 177–187,
- 580 doi:10.1016/j.agrformet.2011.04.013, 2012.
 - Bamberger, I., Stieger, J., Buchmann, N., and Eugster, W.: Spatial variability of methane: Attributing atmospheric concentrations to emissions, Environ. Pollut., 190, 65–74, doi:10.1016/j.envpol.2014.03.028, 2014.
- Beswick, K., Simpson, T., Fowler, D., Choularton, T., Gallagher, M., Hargreaves, K., Sutton, M., and Kaye, A.: Methane emissions on large scales, Atmos. Environ., 32, 3283–3291, doi:10.1016/S1352-2310(98)00080-6, 1998.
 - Choularton, T., Gallagher, M., Bower, K., Fowler, D., Zahniser, M., Kaye, A., Monteith, J., and Harding, R.: Trace gas flux measurements at the landscape scale using boundary-layer budgets, Philos. T. R. Soc. A, 351, 357–369, doi:10.1098/rsta.1995.0039, 1995.
 - Christie, K., Gourley, C., Rawnsley, R., Eckard, R., and Awty, I.: Whole-farm systems analysis of Australian dairy farm greenhouse gas emissions, Anim. Prod. Sci., 62, 998–1011, doi:10.1071/AN12061, 2012.
 - Cleugh, H., Raupach, M., Briggs, P., and Coppin, P.: Regional-scale heat and water vapour fluxes in an agricultural landscape: an evaluation of CBL budget methods at OASIS, Bound.-Lay. Meteorol., 110, 99–137, doi:10.1023/A:1026096711297, 2004.

Deighton, M., Williams, R., Hannah, M., Eckard, R., Boland, T., Wales, W., and Moate, P.: A modified sulphur

- 595 hexafluoride tracer technique enables accurate determination of enteric methane emissions from ruminants, Anim. Feed Sci. Tech., 197, 47–63, doi:10.1016/j.anifeedsci.2014.08.003, 2014.
 - Dengel, S., Levy, P., Grace, J., Jones, S., and Skiba, U.: Methane emissions from sheep pasture, measured with an open-path eddy covariance system, Glob. Change Biol., 17, 3524–3533, doi:10.1111/j.1365-2486.2011.02466.x, 2011.
- 600 Denmead, O., Raupach, M., Dunin, F., Cleugh, H., and Leuning, R.: Boundary layer budgets for regional estimates of scalar fluxes, Glob. Change Biol., 2, 255–264, doi:10.1111/j.1365-2486.1996.tb00077.x, 1996.
 - Denmead, O. T., Leuning, R., Griffith, D. W. T., Jamie, I. M., Esler, M. B., Harper, L. A., and Freney, J. R.: Verifying inventory predictions of animal methane emissions with meteorological measurements, Bound.-Layer Meteorol., 96, 187–209, 2000.
- 605 Detto, M., Baldocchi, D., and Katul, G.: Scaling properties of biologically active scalar concentration fluctuations in the atmospheric surface layer over a managed peatland, Bound.-Lay. Meteorol., 136, 407–430, doi:10.1007/s10546-010-9514-z, 2010.
 - Dlugokencky, E., Nisbet, E., Fisher, R., and Lowry, D.: Global atmospheric methane: budget, changes and dangers, Philos. T. R. Soc. A, 369, 2058–2072, doi:10.1098/rsta.2010.0341, 2011.
- 610 Dorninger, M., Whiteman, C., Bica, B., Eisenbach, S., Pospichal, B., and Steinacker, R.: Meteorological events affecting cold-air pools in a small basin, J. Appl. Meteorol. Clim., 50, 2223–2234, doi:10.1175/2011JAMC2681.1, 2011.
 - Eugster, W. and Siegrist, F. C.: The influence of nocturnal CO₂ advection on CO₂ flux measurements, Basic Appl. Ecol., 1, 177–188, 2000.

- 615 FOEN: Switzerland's Greenhouse Gas Inventory 1990–2013: National Inventory Report 2015, Submission of 15 April 2015 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol, Federal Office for the Environment FOEN, Bern, Switzerland, 596 pp., 2015.
 - Fowler, D., Hargreaves, K. J., Choularton, T. W., Gallagher, M. W., Simpson, T., and Kaye, A.: Measurements of regional CH₄ emissions in the UK using boundary layer budget methods, Energy Conversion and Man-
- 620 agement, 37, 769–775, 1996.

FSO: CORINE Land cover map, Swiss Federal Statistical Office (FSO), Neuchatel, Switzerland, 1998.

FSO: Land use statistics (Arealstatisik) 1992/97, NOAS92, Swiss Federal Statistical Office (FSO), Neuchatel, Switzerland, 2007.

FSO: Data extract of census of agricultural entreprises 2007, Swiss Federal Statistical Office (FSO), Neuchatel,

625 Switzerland, 2009.

- FSO: Land use statistics (Arealstatistik) 2004/09, NOLU04/20120821, Swiss Federal Statistical Office (FSO), Neuchatel, Switzerland, 2012.
- Gilgen, A. and Buchmann, N.: Response of temperate grasslands at different altitudes to simulated summer drought differed but scaled with annual precipitation, Biogeosciences, 6, 2525–2539, doi:10.5194/bg-6-

630 2525-2009, 2009.

640

650

- Griffith, D. W. T., Bryant, G. R., Hsu, D., and Reisinger, A. R.: Methane emissions from free-ranging cattle: Comparison of tracer and integrated horizontal flux techniques, J. Environ. Qual., 37, 582–591, doi:10.2134/jeq2006.0426, 2008.
- Grobler, S., Scholtz, M. M., van Rooyen, H., Mpayipheli, M., and Neser, F. W. C.: Methane production in
- 635 different breeds, grazing different pastures or fed a total mixed ration, as measured by a Laser Methane Detector, S. Afr. J. Anim. Sci., 44, S12–S16, 2014.

Harper, L., Denmead, O., and Flesch, T.: Micrometeorological techniques for measurement of enteric greenhouse gas emissions, Anim. Feed Sci. Tech., 166, 227–239, doi:10.1016/j.anifeedsci.2011.04.013, 2011.

Hartmann, A., Buchmann, N., and Niklaus, P.: A study of soil methane sink regulation in two grasslands exposed to drought and N fertilization, Plant Soil, 342, 265–275, doi:10.1007/s11104-010-0690-x, 2011.

- Heimann, M.: Atmospheric science: Enigma of the recent methane budget, Nature, 476, 157–158, doi:10.1038/476157a, 2011.
 - Hiller, R. V., D.Bretscher, Sontro, T. D., Diem, T., Eugster, W., Henneberger, R., Hobi, S., Hodson, E., Imer, D., Kreuzer, M., Künzle, T., Merbold, L., Niklaus, P. A., Rihm, B., Schellenberger, A., Schroth, M. H., Schu-
- 645 bert, C. J., Siegrist, H., Stieger, J., Buchmann, N., and Brunner, D.: Anthropogenic and natural methane fluxes in Switzerland synthesized within a spatially explicit inventory, Biogeosciences, 11, 1941–1960, doi:10.5194/bg-11-1941-2014, 2014a.
 - Hiller, R. V., Neininger, B., Brunner, D., Gerbig, C., Bretscher, D., Künzle, T., Buchmann, N., and Eugster, W.: Aircraft-based CH₄ flux estimates for validation of emissions from an agriculturally dominated area in
- Holden, J., Derbyshire, S., and Belcher, S.: Tethered balloon observations of the nocturnal stable boundary layer in a valley, Bound.-Lay. Meteorol., 97, 1–24, doi:10.1023/A:1002628924673, 2000.

Switzerland, J. Geophys. Res., 119, 1-14, doi:10.1002/2013JD020918, 2014b.

Hsu, Y., VanCuren, T., Park, S., Jakober, C., Herner, J., FitzGibbon, M., Blake, D., and Parrish, D.: Methane emissions inventory verification in southern California, Atmos. Environ., 44, 1–7, doi:10.1016/j.atmoscamy.2000.10.002.2010

- 655 doi:10.1016/j.atmosenv.2009.10.002, 2010.
 - Imer, D., Merbold, L., Eugster, W., and Buchmann, N.: Temporal and spatial variations of soil CO₂, CH₄ and N₂O fluxes at three differently managed grasslands, Biogeosciences, 10, 5931–5945, doi:10.5194/bg-10-5931-2013, 2013.
- Eggleston, H. S., Buendia, L., Miwa, K., Ngara, T., Tanabe, K. (Eds.): IPCC Guidelines for National Greenhouse
- 660 Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Institute for Global Environmental Strategies, Hayama, Kanagawa, Japan, 2006.
 - Johnson, K., Huyler, M., Westberg, H., Lamb, B., and Zimmerman, P.: Measurement of methane emissions from ruminant livestock using a SF6 tracer technique, Environ. Sci. Technol., 28, 359–362, 1994.
- Judd, M. J., Keillier, F. M., Ulyatt, M. J., Lassey, K. R., Tate, K. R., Shelton, I. D., Harvey, M. J., and
 Walker, C. F.: Net methane emission from grazing sheep, Global Change Biology, 5, 647–657, 1999.
- Kuck, L., Smith Jr, T., Balsley, B., Helmig, D., Conway, T., Tans, P., Davis, K., Jensen, M., Bognar, J., Arrieta, R., and others: Measurements of landscape-scale fluxes of carbon dioxide in the Peruvian Amazon by vertical profiling through the atmospheric boundary layer, J. Geophys. Res., 105, 22137–22146, doi:10.1029/2000JD900105, 2000.
- 670 Lassey, K.: Livestock methane emission: from the individual grazing animal through national inventories to the global methane cycle, Agr. Forest Meteorol., 142, 120–132, doi:10.1016/j.agrformet.2006.03.028, 2007.
 - Lassey, K.: Livestock methane emission and its perspective in the global methane cycle, Anim. Prod. Sci., 48, 114–118, doi:10.1071/EA07220, 2008.
 - Lassey, K. R., Gimson, N. R., Wratt, D. S., Brailsford, G. W., and Bromley, A. M.: Verifying agricultural
- 675 emissions of methane: air sampling from aircraft and mesoscale modelling, in: Non-CO₂ greenhouse gases: Scientific understanding, control and implementation, edited by: Van Ham, J., Baede, A. P. M., Meyer, L. A., and Ybema, R., 107–114, doi:10.1007/978-94-015-9343-4, 2000.
 - Laubach, J. and Kelliher, F. M.: Measuring methane emission rates of a dairy cow herd by two micrometeorological techniques, Agr. Forest Meteorol., 125, 279–303, doi:10.1016/j.agrformet.2004.04.003, 2004.
- 680 Laubach, J. and Kelliher, F. M.: Methane emissions from dairy cows: Comparing open-path laser measurements to profile based techniques, Agr. Forest Meteorol., 135, 340–345, 2005.
 - Laubach, J. and Kelliher, F. M.: Measuring methane emission rates of a diary cow herd (II): results from a backward-Lagrangian stochastic model, Agr. Forest Meteorol., 129, 137–150, 2005.
- Laubach, J., Kelliher, F. M., Knight, T. W., Clark, H., Molano, G., and Cavanagh, A.: Methane emissions from
 beef cattle a comparison of paddock- and animal-scale measurements, Animal Production Science, 48,
 - 132–137, 2008.
 - Levin, I., Glatzel-Mattheier, H., Marik, T., Cuntz, M., and Schmidt, M.: Verification of German methane emission inventories and their recent changes based on atmospheric observations, J. Geophys. Res., 104, 3447–3456, doi:10.1029/1998JD100064, 1999.
- 690 Leytem, A. B., Dungan, R. S., Bjorneberg, D. L., and Koehn, A. C.: Emissions of ammonia, methane, carbon dioxide, and nitrous oxide from dairy cattle housing and manure management systems, J. Environ. Qual., 40, 1383–1394, doi:10.2134/jeq2009.0515, 2011.

Lowry, D., Holmes, C., Rata, N., O'Brian, P., and Nisbet, E.: London methane emissions: Use of diurnal changes in concentration and δ^{13} C to identify urban sources and verify inventories, J. Geophys. Res., 106, 7427–

- 695 7448, doi:10.1029/2000JD900601, 2001.
 - Mahrt, L. and Vickers, D.: Contrasting vertical structures of nocturnal boundary layers, Bound.-Lay. Meteorol., 105, 351–363, doi:10.1023/A:1019964720989, 2002.
 - Mathieu, N., Strachan, I., Leclerc, M., Karipot, A., and Pattey, E.: Role of low-level jets and boundary-layer properties on the NBL budget technique, Agr. Forest Meteorol., 135, 35–43,
- 700 doi:10.1016/j.agrformet.2005.10.001, 2005.
 - McGinn, S. M. and Beauchemin, K. A.: Dairy farm methane emissions using a dispersion model, J. Environ. Qual., 41, 73–79, doi:10.2134/jeq2011.0266, 2012.
 - McGinn, S. M., Flesch, T. K., Harper, L. A., and Beuchemin, K. A.: An approach for measuring methane emissions from whole farms, J. Environ. Qual., 35, 14–20, doi:10.2134/jeq2005.0250, 2006.
- 705 McGinn, S. M., Turner, D., Tomkins, N., Charmley, E., Bishop-Hurley, G., and Chen, D.: Methane emissions from grazing cattle using point-source dispersion, J. Environ. Qual., 40, 22–27, doi:10.2134/jeq2010.0239, 2011.
 - Merbold, L., Eugster, W., Stieger, J., Zahniser, M., Nelson, D., and Buchmann, N.: Greenhouse gas budget (CO₂, CH₄ and N₂O) of intensively managed grassland following restoration, Glob. Change Biol., 20, 1913–

710 1928, doi:10.1111/gcb.12518, 2014.

- Messerli, P.: Die Ballonsondierungen in Hagnau vom 18. bis 20. Mai 1992, Seminararbeit, Institute of Geography, University of Bern, Bern, Switzerland, 71 pp., 1992.
- Nisbet, E. and Weiss, R.: Top-down versus bottom-up, Science, 328, 1241–1243, doi:10.1126/science.1189936, 2010.
- 715 Parker, M. and Raman, S.: A case study of the nocturnal boundary layer over a complex terrain, Bound.-Lay. Meteorol., 66, 303–324, doi:10.1007/BF00705480, 1993.
 - Pattey, E., Strachan, I., Desjardins, R., Edwards, G., Dow, D., and MacPherson, J.: Application of a tunable diode laser to the measurement of CH₄ and N₂O fluxes from field to landscape scale using several micrometeorological techniques, Agr. Forest Meteorol., 136, 222–236, doi:10.1016/j.agrformet.2004.12.009, 2006.
- 720 Pendall, E., Schwendenmann, L., Rahn, T., Millers, J., Tans, P., and White, J.: Land use and season affect fluxes of CO₂, CH₄, CO, N₂O, H₂ and isotopic source signatures in Panama: evidence from nocturnal boundary layer profiles, Glob. Change Biol., 16, 2721–2736, doi:10.1111/j.1365-2486.2010.02199.x, 2010.
 - Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanual, S., Buendia, L., Hoppaus, R., Martinsen, T., Mejer, J., Miwa, K., and Tanabe, K. (Eds.): Good practice guidance and uncertainty management
- 725 in national greenhouse gas inventories, IPCC National Greenhouse Cost Inventories programme, Technical Support Unit, Institute for Global Environmental Strategies, Hayama, Kanagawa, Japan, 2000.
 - Pinto, J., Parsons, D., Brown, W., Cohn, S., Chamberlain, N., and Morley, B.: Coevolution of down-valley flow and the nocturnal boundary layer in complex terrain, J. Appl. Meteorol. Clim., 45, 1429–1449, doi:10.1175/JAM2412.1, 2006.
- 730 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, available at: http://www.R-project.org/, 2014.

- Raupach, M., Denmead, O., and Dunin, F.: Challenges in linking atmospheric CO₂ concentrations to fluxes at local and regional scales, Aust. J. Bot., 40, 697–716, doi:10.1071/BT9920697, 1992.
- Saggar, S., Bolan, N., Bhandral, R., Hedley, C., and Luo, J.: A review of emission of methane, ammonia, and
 nitrous oxide from animal excreta deposition and farm effluent application in grazed pastures, New Zeal. J.
 - Siegrist, F. C.: Determination of energy and trace gas fluxes on a regional scale: Combination of local surface flux measurements and vertical flux profiles throughout the atmospheric boundary layer in complex terrain (Swiss Seeland region), PhD thesis, University of Bern, Institute of Geography, University of Bern, Geo-
- 740 graphical Institute, Hallerstrasse 12, 3012 Bern, Switzerland, vol. G67 of Geographica Bernensia, 113 pp., 2001.
 - Soliva, C.: Report to the attention of IPCC about the data set and calculation method used to estimate methane formation from enteric fermentation of agricultural livestock population and manure management in Swiss agriculture, Institute of Animal Science, ETH Zurich, Switzerland. On behalf of the Federal Office for the Environment (FOEN), Bern, Switzerland, 14 pp., 2006.
- Staerfl, S., Zeitz, J., Kreuzer, M., and Soliva, C.: Methane conversion rate of bulls fattened on grass or maize silage as compared with the IPCC default values, and the long-term methane mitigation efficiency of adding acacia tannin, garlic, maca and lupine, Agric. Ecosyst. Environ., 148, 111–120, doi:10.1016/j.agee.2011.11.003, 2012.
- 750 Sun, J., Burns, S., Delany, A., Oncley, S., Turnipseed, A., Stephens, B., Lenschow, D., LeMone, M., Monson, R., and Anderson, D.: CO₂ transport over complex terrain, Agr. Forest Meteorol., 145, 1–21, doi:10.1016/j.agrformet.2007.02.007, 2007.
 - Suter, F.: Greenhouse gases from managed ecosystems: a farm scale perspective. Insights from a consolidated greenhouse gas budgets of a typical Swiss three-stage grassland farming system, ETH Master thesis, ETH
- 755 Zurich, Zurich, Switzerland, 69 pp., 2011.

Agr. Res., 47, 513-544, 2004.

745

- Ulyatt, M., Lassey, K., Shelton, I., and Walker, C.: Seasonal variation in methane emission from dairy cows and breeding ewes grazing ryegrass/white clover pasture in New Zealand, New Zeal. J. Agr. Res., 45, 217–226, 2002.
 - Utiger, M.: Ballonsondierung Merenschwand vom 16.-18.06.1992, Seminararbeit, Insitute of Geography, Uni-
- 760 versity of Bern, Bern, 98 pp., 1992.
 - van Ham, J., Baede, A. P. M., Meyer, L. A., and Ybema, R. (Eds.): Non-CO₂ greenhouse gases: Scientific understanding, control and implementation, Kluwer Academic, Dordrecht, the Netherlands, 604 pp., 2000.
 - Wang, F. L., Bettany, J. R., and Huang, P. M.: Methane uptake by cultivated Canadian prairie and forest soils, Geoderma, 88, 39–53, doi:10.1016/S0016-7061(98)00076-7, 1999.
- 765 Wang, J., Cardenas, L., Misselbrook, T., and Gilhespy, S.: Development and application of a detailed inventory framework for estimating nitrous oxide and methane emissions from agriculture, Atmos. Environ., 45, 1454– 1463, doi:10.1016/j.atmosenv.2010.12.014, 2011.
 - Wang, S., Ackermann, R., and Stutz, J.: Vertical profiles of O_3 and NO_x chemistry in the polluted nocturnal boundary layer in Phoenix, AZ: I. Field observations by long-path DOAS, Atmos. Chem. Phys., 6, 2671–
- 770 2693, doi:10.5194/acp-6-2671-2006, 2006.



Figure 1. Meteorological conditions at Chamau (30 min averages) during the balloon measurements (gray shaded areas) in 2011 and 2012. Top panel: Air temperature and cooling rates $\Delta \overline{T}/\Delta t$ at 2 m a.g.l. Bottom panel: Wind speed and friction velocity u_* at 2 m a.g.l. Dashed horizontal lines show the cooling rate threshold of 0 K s^{-1} in the top panel and the threshold of low mechanical turbulence $u_* < 0.08 \text{ m s}^{-1}$ (Zeeman et al., 2010) in the lower panel.

Wratt, D. S., Gimson, N. R., Brailsford, G. W., Lassey, K. R., Bromley, A. M., and Bell, M. J.: Estimating regional methane emissions from agriculture using aircraft measurements of concentration profiles, Atmos. Environ., 35, 497–508, doi:10.1016/S1352-2310(00)00336-8, 2001.

Zeeman, M., Werner, R., Eugster, W., Siegwolf, R., Wehrle, G., Mohn, J., and Buchmann, N.: Optimization of

automated gas sample collection and isotope ratio mass spectrometric analysis of δ^{13} C of CO₂ in air, Rapid Commun. Mass Sp., 22, 3883–3892, doi:10.1002/rcm.3772, 2008.

- Zeeman, M., Hiller, R., Gilgen, A., Michna, P., Plüss, P., Buchmann, N., and Eugster, W.: Management and climate impacts on net CO₂ fluxes and carbon budgets of three grasslands along an elevational gradient in Switzerland, Agr. Forest Meteorol., 150, 519–530, doi:10.1016/j.agrformet.2010.01.011, 2010.
- 780 Zeitz, J., Solvia, C., and Kreuzer, M.: Swiss diet types for cattle: how accurately are they reflected by the Intergovernmental Panel on Climate Change default values?, J. Integr. Environ. Sci., 9, 199–216, doi:10.1080/1943815X.2012.709253, 2012.
 - Zinchenko, A., Paramonova, N., Privalov, V., and Reshetnikov, A.: Estimation of methane emissions in the St. Petersburg, Russia, region: an atmospheric nocturnal boundary layer budget approach, J. Geophys. Res., 107,
- 785 1451–1476, doi:10.1029/2001JD001369, 2002.



Figure 2. (a) Kriged time-space interpolation of the CH₄ concentrations obtained from the balloon and near-surface (0–10 m) measurements during 16/17 August 2011. 2011, including rejected measurements not used for NBL budget estimates. Vertical and horizontal kriging directions were equally weighted, resulting sometimes in sharp vertical concentration changes. The balloon position is shown with a white line, whereas the near-surface concentration measurements are shown with a white box. (b) Nocturnal course of CH₄ concentrations at 10, 40, 110 and 130 m a.g.l. The gray shaded time period shows the period when the wind direction changes from easterly (around 20:00) to south-westerly winds (180–270°, down-valley direction) near the surface and anti-wind (in opposite direction) aloft (see legend for details). This clear separation between the down-valley wind and its anti-wind during the August 2011 campaign explains why a relatively low z_i of only 50 m resulted for the integration of the NBL budget fluxes (see Table 1).

Table 1. Overview of the balloon measurements in 2011 and 2012 with the maximum height reached, the integration height in the time-height kriging interpolation, and the resulting averaged NBL budget fluxes (mean \pm SD, where SD is the standard deviation of all 0.3 h time slices in the time-height kriging interpolation included in the calculation of the respective mean flux). The difference between two consecutive interpolated time slices was used to approximate the local derivative of concentration over time as detailed in Section 2.5).

Date	Night	Sounding	Time [UTC+1]	max. Height [m a.g.l.]	Int. Height [m a.g.l.]	NBL budget flux $[\mu g CH_4 m^{-2} s^{-1}]$
16/17 Aug	N1	S1	20:47-21:26	140	50	1.27 ± 0.85
2011		S2	21:26-22:01	150	50	1.85 ± 0.07
		S 3	22:01-22:41	150	50	1.55 ± 0.48
		S4	22:41-23:30	160	70	1.19 ± 1.05
		S5	00:48-01:47	150	50	1.13 ± 0.06
24/25 Jul	N2	S6	20:51-21:24	150	60	1.73 ± 0.14
2012		S 7	23:27-23:58	150	50	2.37 ± 0.72
		S8	01:05-01:37	130	110	2.88 ± 0.61
		S9	05:07-05:37	130	80	1.21 ± 0.34
25/26 Jul	N3	S10	20:56-21:16	170	50	1.18 ± 0.11
2012		S11	23:03-23:23	160	50	1.68 ± 0.05
		S12	01:06-01:22	130	40	0.79 ± 0.13
26/27 Jul	N4	S13	20:53-21:14	190	60	1.93 ± 0.06
2012		S14	23:03-23:23	180	120	1.84 ± 0.09
		S15	01:01-01:25	180	50	1.52 ± 0.01
		S16	03:07-03:29	170	60	1.53 ± 0.12
Overall mean						1.60 ± 0.31

Table 2. Farm-scale CH_4 emission estimates from different inventories (NIR, SEI and CHAI) and the NBL budget approach. The inventory estimates are based on the methodology recommended by the IPCC (see text for details). CHAI estimates use current stocking census data, whereas NIR and SEI estimates are both based on 2011 stocking census data. NBL budget fluxes are denoted as the mean of all budget fluxes from the individual balloon profiles of the respective nights. The fluxes are in $\mu g CH_4 m^{-2} s^{-1}$ (mean \pm SD).

Year	Campaign	NIR	SEI	CHAI	NBL
	16/17 August 2011				1.40 ± 0.50
2011		0.83 ± 0.30	0.42 ± 0.15	1.29 ± 0.47	1.40 ± 0.50
	24/25 July 2012				2.05 ± 0.45
	25/26 July 2012				1.22 ± 0.10
	26/27 July 2012				1.70 ± 0.07
2012		0.83 ± 0.30	0.42 ± 0.15	1.74 ± 0.63	1.66 ± 0.20



Figure 3. Vertical profiles of (**a**) potential temperature gradients, (**b**) CH_4 concentrations and (**c**) wind directions of the individual balloon soundings. The dashed vertical lines indicate neutral stratification in (**a**), and lower and upper wind direction limits used for the NBL budget flux calculation in (**c**).



Figure 4. Inversion height z_i (**a**) approximated via the profile of relative humidity for ascents (bold solid lines) and descents (bold dashed lines) during the first 4 nocturnal tethered balloon profiles, and (**b**) with vertical profiles of CH₄ concentrations superimposed. The balloon height is shown as a time trace (black line). The relative humidity was 100% at the ground surface and the minimum value observed in a profile was used as a reference (thin dashed vertical lines) placed at the time point of the beginning of ascents and descents, respectively. The large and small red circles in (**a**) link the relative humidity showing the top of the saturated surface layer (large circles) with the corresponding height and time point on the black curves (small circle with red horizontal dashed line). The inversion height z_i is derived separately for ascents and descents, yielding the gray band of showing an uncertainty on the order of 20–30 m for z_i .



Figure 5. Sensitivity of NBL budget flux calculations on z_i used as the upper boundary of integration in Eq. (1) using data from all soundings (one ascent and one descent taken as one sounding). The color bars show the range of fluxes resulting if z_i is uncertain between 50 and 400 m (thin orange line), between 100 and 300 m (red medium line), or between 150 and 200 m (bold green line) using interpolated data (see Section 2.5). The circles (**A**) show the NBL budget flux that was achieved if no interpolation extrapolation was done beyond the height range that was actually measured, and the black disks and error bars (**T**) show the values presented in Table 1 following the method described in Section 2.3. Soundings with a gray background were considered instationary and were not used in this study.